1   TITLE:

2   **Stimulus-driven brain rhythms within the alpha band: The attentional-modulation conundrum**

4   ABBREVIATED TITLE: Reversed attentional modulation of alpha and SSRs

AUTHORS:	Affiliation	ORCID	Twitter
Christian Keitel*	1	0000-0003-2597-5499	@KeiCetel
Anne Keitel	1	0000-0003-4498-0146	@anneke_sci
Christopher SY	1	0000-0002-4157-4049	@ChrisSYBenwell
Christoph Daube	1	0000-0002-1763-8508	@christophdaube
Gregor Thut	1	0000-0003-1313-4262	
Joachim Gross	1,2	0000-0002-3994-1006	@Joachim__Gross

AFFILIATIONS:

1 – Institute of Neuroscience and Psychology, University of Glasgow, 58 Hillhead Street, G12 8QB

Glasgow, UK; **2** – Institut für Biomagnetismus und Biosignalanalyse, Westfälische Wilhelms-

Universität, Malmedyweg 15, 48149 Münster, Germany

* – corresponding author, christian.keitel@glasgow.ac.uk

KEYWORDS: alpha rhythm, entrainment, phase synchronisation, spatial attention, steady-state

response (SSR), frequency tagging

ACKNOWLEDGMENTS: Funded by a Wellcome Trust Joint Investigator Grant awarded to GT and JG

(#098433/#098434). Lucy Dewhurst and Jennifer McAllister assisted in data collection. The

experimental stimulation was realized using Cogent Graphics developed by John Romaya at the

Laboratory of Neurobiology, Wellcome Department of Imaging Neuroscience, University College

London (UCL).

0 Table(s), 5 Figure(s), 1 Footnote(s)

ABSTRACT

Two largely independent research lines use rhythmic sensory stimulation to study visual processing. Despite the use of strikingly similar experimental paradigms, they differ crucially in their notion of the stimulus-driven periodic brain responses: One regards them mostly as synchronised (entrained) intrinsic brain rhythms; the other assumes they are predominantly evoked responses (classically termed steady-state responses, or SSRs) that add to the ongoing brain activity. This conceptual difference can produce contradictory predictions about, and interpretations of, experimental outcomes. The effect of spatial attention on brain rhythms in the alpha-band (8 – 13 Hz) is one such instance: alpha-range SSRs have typically been found to *increase* in power when participants focus their spatial attention on laterally presented stimuli, in line with a gain control of the visual evoked response. In nearly identical experiments, retinotopic *decreases* in entrained alpha-band power have been reported, in line with the inhibitory function of intrinsic alpha. Here we reconcile these contradictory findings by showing that they result from a small but far-reaching difference between two common approaches to EEG spectral decomposition. In a new analysis of previously published EEG data, recorded during bilateral rhythmic visual stimulation, we find the typical SSR gain effect when emphasising stimulus-locked neural activity and the typical retinotopic alpha suppression when focusing on ongoing rhythms. These opposite but parallel effects suggest that spatial attention may bias the neural processing of dynamic visual stimulation via two complementary neural mechanisms.

SIGNIFICANCE STATEMENT

Attending to a visual stimulus strengthens its representation in visual cortex and leads to a retinotopic suppression of spontaneous alpha rhythms. To further investigate this process, researchers often attempt to phase-lock, or entrain, alpha through rhythmic visual stimulation under the assumption that this entrained alpha retains the characteristics of spontaneous alpha. Instead, we show that the part of the brain response that is phase-locked to the visual stimulation *increased* with attention (in line with steady-state evoked potentials), while the typical suppression was only present in non-stimulus-locked alpha activity. The opposite signs of these effects suggest that attentional modulation of dynamic visual stimulation relies on two parallel cortical mechanisms – retinotopic alpha suppression and increased temporal tracking.

INTRODUCTION

Cortical visual processing has long been studied using rhythmic sensory stimulation (Adrian and Matthews, 1934; Walter et al., 1946; Regan, 1966). This type of stimulation drives continuous brain responses termed steady-state responses (SSRs) that reflect the temporal periodicities in the stimulation precisely. SSRs allow tracking of individual stimuli in multi-element displays (Vialatte et al., 2010; Norcia et al., 2015). Further, they readily indicate cognitive biases of cortical visual processing, such as the selective allocation of attention (Morgan et al., 1996; Keitel et al., 2013; Stormer et al., 2014).

Although SSRs can be driven using a wide range of frequencies (Herrmann, 2001), stimulation at alpha band frequencies (8 – 13 Hz) has stirred particular interest. Alpha rhythms dominate brain activity in occipital visual cortices (Groppe et al., 2013; Keitel and Gross, 2016) and influence perception (Benwell et al., 2017a; Benwell et al., 2017b; Iemi et al., 2017; Samaha et al., 2017). Researchers have therefore used alpha-rhythmic visual stimulation in attempts to align the phase of – or *entrain* – intrinsic alpha rhythms and consequently provided evidence for visual alpha entrainment (Mathewson et al., 2012; Zauner et al., 2012; Spaak et al., 2014; Gulbinaite et al., 2017b). These findings suggest that at least part of the SSR driven by alpha-band stimulation should be attributed to entrained alpha generators (Notbohm et al., 2016).

Some issues remain with such an account (Capilla et al., 2011; Keitel et al., 2014). For instance, experiments have consistently reported SSR power increases when probing effects of spatial selective attention on SSRs driven by lateralised hemifield stimuli (Müller et al., 1998), also when using alpha-band frequencies (Kim et al., 2007; Kashiwase et al., 2012; Keitel et al., 2013). However, recent studies that used similar paradigms, but treated alpha-frequency SSRs as phase-entrained alpha rhythms in line with an earlier study using rhythmic transcranial magnetic stimulation (Herring et al., 2015), reported the opposite effect (Gulbinaite et al., 2017a; Kizuk and Mathewson, 2017). Oscillatory brain activity showed attentional modulations characteristic of the intrinsic alpha rhythm during stimulation: Alpha power decreased over the hemisphere contralateral to the attended position, an effect known to be part of a retinotopic alpha power lateralisation during selective spatial attention (Worden et al., 2000; Kelly et al., 2006; Thut et al., 2006; Rihs et al., 2007; Capilla et al., 2012). Briefly put, studies analysing SSRs show a power *increase,* whereas studies analysing "entrained alpha" show a power *decrease* with attention*.*

Both neural responses originate from visual cortices contralateral to the hemifield position of the driving stimuli (Keitel et al., 2013; Spaak et al., 2014). Assuming a single underlying neural process, opposite attention effects therefore seemingly contradict each other. However, results in support of

alpha entrainment differed in how exactly responses to the periodic stimulation were quantified.

Effects consistent with SSR modulation resulted from spectral decompositions performed on trial-

averaged EEG waveforms. This approach tunes the resulting power estimate to the part of the neural

response that is sufficiently time-locked to the stimulation (Tallon-Baudry et al., 1996; Delorme and

Makeig, 2004). Effects consistent with alpha entrainment instead typically result from averages of

single-trial spectral transforms, thus emphasising intrinsic non-phase-locked activity (Tallon-Baudry

et al., 1998; Herrmann et al., 2004). Both approaches have been applied before to compare stimulus-

evoked and induced brain rhythms in the gamma frequency range (~40 Hz; Tallon-Baudry et al.,

1998; Picton et al., 2003). Here we focussed on contrasting the attentional modulation of alpha

during- and SSRs driven by an alpha-rhythmic stimulation.

We therefore compared the outcome of both approaches in a new analysis of previously reported

EEG data (Keitel et al., 2017b). Participants viewed two lateralised stimuli, both flickering at alpha

band frequencies (10 and 12 Hz). They were cued to focus on one of the two and perform a target

detection task at the attended position. We quantified spectral power estimates according to both

approaches described above from the same EEG data. Should the outcome depend on the approach

taken, we expected to find the typical alpha power lateralisation (contralateral < ipsilateral) when

averaging single-trial power spectra. In power spectra of trial-averaged EEG instead we expected the

typical SSR power gain modulation in the opposite direction (contralateral > ipsilateral). Crucially,

such an outcome would warrant a re-evaluation of stimulus-driven brain rhythms in the alpha range

and intrinsic alpha as a unitary phenomenon (alpha entrainment).

Figure 1 Stimulus schematics and trial time course. (**a**) shows the exemplary time course of one trial
with a cue displayed for 0.5 sec (here: Attend Right), followed by the bilateral visual stimulation for
3.5 sec. Left (L) stimulus contrast fluctuated with a rate of 10 Hz and Right (R) stimulus contrast at 12
Hz. Targets that participants were instructed to respond to were slightly altered versions of the

stimuli (see inset) that were displayed occasionally for 0.3 sec. (**b**) Rhythmic visual stimulation was
achieved by a frame-by-frame adjustment of global stimulus contrast (through local luminance
changes) as exemplified here in one representative cycle.
114

METHODS

Participants

For the present report, we re-analysed EEG data of 17 volunteers recorded in an earlier study (Keitel

et al., 2017a). Participants (13 women; median age = 22 yrs, range = 19 – 32 yrs) declared normal or

corrected-to-normal vision and no history of neurological diseases or injury. All procedures were

approved by the ethics committee of the College of Science & Engineering at the University of

Glasgow (application no. 300140020) and adhered to the guidelines for the treatment of human

subjects in the Declaration of Helsinki. Volunteers received monetary compensation of £6/h. They

gave informed written consent before participating in the experiment. Note that we excluded five

additional datasets on grounds reported in the original study (four showed excessive eye

movements, one underperformed in the task).

Stimulation

Participants viewed experimental stimuli on a computer screen (refresh rate = 100 frames per sec) at

a distance of 0.8 m that displayed a grey background (luminance = 6.5 cd/m^2). Small concentric

circles in the centre of the screen served as a fixation point (*Figure 1*). Two blurry checkerboard

patches (horizontal/vertical diameter = 4° of visual angle) were positioned at an eccentricity of 4.4°

from central fixation, one each in the lower left and lower right visual quadrants. Both patches

changed contrast rhythmically during trials: Stimulus contrast against the background was modulated

by varying patch peak luminance between 7.5 cd/m^2 (minimum) and 29.1 cd/m^2 (maximum).

On each screen refresh, peak luminance changed incrementally to approach temporally smooth

contrast modulations as opposed to a simple on-off flicker (Andersen and Muller, 2015). Further

details of the stimulation can be found in Keitel et al. (2017a). The contrast modulation followed a

10-Hz periodicity for the left and a 12-Hz periodicity for the right stimulus. Note that the experiment

featured further conditions displaying quasi-rhythmic contrast modulations in different frequency

bands. Corresponding results can be found in the original report and will not be considered in the

present analysis.

Procedure and Task

Participants performed the experiment in an acoustically dampened and electromagnetically

shielded chamber. In total, they were presented with 576 experimental trials, subdivided into 8

blocks with durations of ~5 min each. Between blocks, participants took self-paced breaks. Prior to the experiment, participants practiced the behavioural task (see below) for at least one block. After each block they received feedback regarding their accuracy and response speed. The experiment was comprised of 8 conditions (= 72 trials each) resulting from a manipulation of the two factors attended position (left vs. right patch) and stimulation frequency (one rhythmic and three quasi-rhythmic conditions) in a fully balanced design. Trials of different conditions were presented in pseudo-random order. As stated above, the present study focussed on the two conditions featuring fully rhythmic stimuli. Corresponding trials (N = 144) were thus selected a posteriori from the full design.

Single trials began with cueing participants to attend to the left or right stimulus for 0.5 sec, followed by presentation of the dynamically contrast-modulating patches for 3.5 sec (*Figure* 1). After patch offset, an idle period of 0.7 sec allowed participants to blink before the next trial started.

To control whether participants maintained a focus of spatial attention, they were instructed to respond to occasional brief "flashes" (0.3 sec) of the cued stimulus (= targets) while ignoring similar events in the other stimulus (= distracters). Targets and distracters occurred in one third of all trials and up to 2 times in one trial with a minimum interval of 0.8 sec between subsequent onsets. Detection was reported as speeded responses to flashes (recorded as space bar presses on a standard key board).

Behavioural data recording and analyses

Flash detections were considered a 'hit' when a response occurred from 0.2 to 1 sec after target onset. Delays between target onsets and responses were considered reaction times (RT). Statistical comparisons of mean accuracies (proportion of correct responses to the total number of targets and distracters) and median RTs between experimental conditions were conducted and reported in (2017a). In the present study, we did not consider the behavioural data further. Note that the original statistical analysis found that task performance in Attend-Left and Attend-Right conditions was comparable.

Electrophysiological data recording

EEG was recorded from 128 scalp electrodes and digitally sampled at a rate of 512 Hz using a BioSemi ActiveTwo system (BioSemi, Amsterdam, Netherlands). Scalp electrodes were mounted in an elastic cap and positioned according to an extended 10-20-system (Oostenveld and Praamstra, 2001). Lateral eye movements were monitored with a bipolar outer canthus montage (horizontal electro-

175 oculogram). Vertical eye movements and blinks were monitored with a bipolar montage of

176 electrodes positioned below and above the right eye (vertical electro-oculogram).

Electrophysiological data pre-processing

From continuous data, we extracted epochs of 5 s, starting 1 s before patch onset using the MATLAB

toolbox EEGLAB (Delorme and Makeig, 2004). In further pre-processing, we excluded epochs that

corresponded to trials containing transient targets and distracters (24 per condition) as well as

epochs with horizontal and vertical eye movements exceeding 20 μV (~ 2° of visual angle) or

containing blinks. For treating additional artefacts, such as single noisy electrodes, we applied the

'fully automated statistical thresholding for EEG artefact rejection' (FASTER; Nolan et al., 2010). This

procedure corrected or discarded epochs with residual artefacts based on statistical parameters of

the data. Artefact correction employed a spherical-spline-based channel interpolation. Epochs with

more than 12 artefact-contaminated electrodes were excluded from analysis.

From 48 available epochs per condition, we discarded a median of 14 epochs for the Attend-Left

conditions and 15 epochs for the Attend-Right conditions per participant with a between-subject

range of 6 to 28 (Attend-Left) and 8 to 31 epochs (Attend-Right). Within-subject variation of number

of epochs per condition remained small with a median difference of 3 trials (maximum difference = 9

for one participant).

Subsequent analyses were carried out in Fieldtrip (Oostenveld et al., 2011) in combination with

custom-written routines. We extracted segments of 3 s starting 0.5 s after patch onset from pre-

processed artefact-free epochs (5 s). Data prior to stimulation onset (1 s), only serving to identify eye

movements shortly before and during cue presentation, were omitted. To attenuate the influence of

stimulus-onset evoked activity on EEG spectral decomposition, the initial 0.5 s of stimulation were

excluded. Lastly, because stimulation ceased after 3.5 s, we also discarded the final 0.5 s of original

epochs.

Electrophysiological data analyses – spectral decomposition

Artefact-free 3-sec epochs were converted to scalp current densities (SCDs), a reference-free

measure of brain electrical activity (Ferree, 2006; Kayser and Tenke, 2015), by means of the spherical

spline method (Perrin et al., 1987) as implemented in Fieldtrip (function *ft_scalpcurrentdensity*,

method 'spline', lambda = 10^{-4}). Detrended (i.e. mean and linear trend removed) SCD time series

were then Tukey-tapered and subjected to Fourier transforms while employing zero-padding in order

to achieve a frequency-resolution of 0.25 Hz. Crucially, from resulting complex Fourier spectra we

calculated two sets of aggregate power spectra with slightly different approaches. First, we

calculated power spectra as the average of squared absolute values of complex Fourier spectra (*Z*) as
follows:

$$onPOW(f) = \frac{1}{n} \sum_{i=1}^{n} |Z_i(f)|^2 \qquad [1]$$

where *onPOW* is the classical power estimate for ongoing (intrinsic) oscillatory activity for frequency
f and *n* is the number of trials. Secondly, we additionally calculated the squared absolute value of the
averaged complex Fourier spectra according to:

$$evoPOW(f) = \left| \frac{1}{n} \sum_{i=1}^{n} Z_i(f) \right|^2 \qquad [2]$$

The formula yields *evoPOW*, or evoked power, an estimate that is identical with the frequency-
tagging standard approach of averaging per-trial EEG time series before spectral decomposition. This
step is usually performed to retain only the truly phase-locked response to the stimulus (Tallon-
Baudry et al., 1996). Note that both formulas only differ in the order in which weighted sums and
absolute values are computed. Also note that formula [2] is highly similar to the calculation of inter-
trial phase coherence (ITC), a popular measure of phase locking (Cohen, 2014; Gross, 2014; van
Diepen and Mazaheri, 2018). ITC calculation additionally includes a trial-by-trial amplitude
normalisation. To complement our analysis we thus quantified ITC according to:

$$ITC(f) = \left| \frac{1}{n} \sum_{i=1}^{n} \frac{Z_i(f)}{|Z_i(f)|} \right| \qquad [3]$$

For further analyses, power spectra were normalised by converting them to decibel scale, i.e. taking
the decadic logarithm, then multiplying by 10 (hereafter termed log power spectra). ITC was
converted to ITCz to reduce the bias introduced by differences in trial numbers between conditions
(Bonnefond and Jensen, 2012; Samaha et al., 2015).

Figure 2 EEG spectral decomposition. (**a**) Power spectra collapsed across conditions and all electrode positions below the sagittal midline for single subjects (light grey lines) and group averages (strong black line). Note the characteristic alpha peaks in the frequency range of 8 – 13 Hz. Inset scalp malp shows topographical distribution of alpha power on a dB scale based on scalp current denisities. (**b**) Same as in a but for ‚evoked' power. Distinct peaks are visible at stimulation frequencies 10 & 12 Hz (dashed vertical orange lines across plots). Inset scalp malps shows topographical distribution of SSR power at 10 & 12 on a dB scale. Note the difference in scale between ongoing power in a and evoked power b. (**c**) Same as in a but for inter-trial phase-locking (ITCz). Inset scalp malps shows topographical distribution of SSR ITCz at 10 & 12.

Alpha power – attentional modulation and lateralisation

Spectra of ongoing power (*onPOW*), pooled over both experimental conditions and all electrodes, showed a prominent peak in the alpha frequency range (*Figure 2*). We used mean log ongoing power across the range of 8 – 13 Hz to assess intrinsic alpha power modulations by attention. Analysing Attend-Right and Attend-Left conditions separately, yielded two alpha power topographies for each participant. These were compared by means of cluster-based permutation statistics (Maris and Oostenveld, 2007) using N = 5000 random permutations. We clustered data across channel neighbourhoods with an average size of 7.9 channels that were determined by triangulated sensor proximity (function *ft_prepare_neighbours*, method 'triangulation'). The resulting probabilities (*P*-values) were corrected for two-sided testing. Subtracting left-lateralised (Attend-Left conditions) from right-lateralised (Attend-Right) alpha power topographies, we found a right-hemispheric

positive and a left-hemispheric negative cluster of electrodes that was due to the retinotopic effects of spatial attention on alpha power lateralisation (*Figure 3*), similar to an earlier re-analysis of the other conditions of this experiment (Keitel et al., 2018).

Finally, we tested the difference between Attend-Left and Attend-Right conditions, i.e. attention effects for left- and right-hemispheric clusters separately. To this end we submitted alpha power differences (contralateral hemifield attended minus ignored) to Bayesian one-sample t-tests against zero (Rouder et al., 2009). Attention effects were further compared against each other by means of a Bayesian paired-samples t-test as implemented in JASP (JASP-Team, 2018) with a Cauchy prior scaled to r = 0.5, putting more emphasis on smaller effects (Rouder et al., 2012; Schonbrodt and Wagenmakers, 2017).

This procedure allowed us to quantify the evidence in favour of the null vs the alternative hypothesis (H_0 vs H_1). For each test, the corresponding Bayes factor (called BF_{10}) showed evidence for H_1 (compared to H_0) if it exceeded a value of 3, and no evidence for H_1 if $BF_{10} < 1$, with the intervening range 1 – 3 termed 'anecdotal evidence' by convention (Wagenmakers et al., 2011). Inversing BF_{10}, to yield a quantity termed BF_{01}, served to quantify evidence in favour of H_0 on the same scale. For BF_{10} and BF_{01} values < 1 were taken as inconclusive evidence for either hypothesis. Note that for the sake of brevity we report errors in BF estimates only when exceeding 0.001%.

SSR power – attentional modulation

Spectra of evoked power, pooled over both experimental conditions and all electrodes, revealed periodic responses to the two stimuli at the respective stimulation frequencies, 10 and 12 Hz (*Figure 2*). Therefore, we assessed attention effects for these two spectral SSR representations. Two separate cluster-based permutation tests, one for each stimulation frequency, contrasted evoked power topographies between attended and ignored (= other stimulus attended) conditions. Two-sided tests were performed with the same parameters as for alpha power (see above).

Again, we found one electrode cluster carrying systematic attention effects per frequency. As for alpha, SSR power from these two clusters were subjected to separate Bayesian one-sample t-tests against zero (one-sided, attended > ignored) and compared against each other by means of a Bayesian paired-sample t-test (two-sided).

SSR inter-trial phase coherence – attentional modulation

We also evaluated a pure measure of neural phase-locking to the stimulation, SSR inter-trial phase coherence (ITC), because evoked power can be regarded as a hybrid measure depending on both the amplitude of the underlying rhythmic response and the consistency of its phase across trials. ITC

282 indicates only the latter as SSRs are set to unit amplitude prior to summing across trials (see

283 formula 3). ITC spectra, pooled over both experimental conditions and all electrodes, showed distinct

284 neural phase-locking at the respective driving frequencies, 10 and 12 Hz (*Figure 2*). Cluster-based

285 permutation testing confirmed topographic regions that showed systemic gain effects in ITC.

286 Subsequently, the same Bayesian inference was applied to data from these clusters as for SSR power.

287 Correlation of alpha and SSR attention effects

288 As a consequence of our counter-intuitive finding that SSR attention effects seemed strongest over

289 occipital regions ipsi-lateral to the driving stimulus (see Results section *SSR power & inter-trial phase*

290 *locking – attentional modulation* below) we explored a posteriori whether these effects could be

291 explained by ipsilateral increases in alpha power during focussed attention. We correlated attention

292 effects on alpha and SSR power using Bayesian inference (rank correlation coefficient Kendall's tau-b

293 or τ_b, beta-prior = 0.75) to test for a positive linear relationship. More specifically, we correlated the

294 left-hemispheric alpha power suppression (Ignored minus Attended) with the 10-Hz SSR (evoked)

295 power attention effect (Attended minus Ignored) and the right-hemispheric alpha power suppression

296 with the 12-Hz SSR power attention effect. We opted for these combinations because the

297 corresponding effects overlapped topographically (see Results). Along with the correlation coefficient

298 ρ, we report its 95%-Credible Interval (95%-CrI).

299 We also probed the linear relationship between alpha power and SSR ITC attention effects. Because

300 ITC gains were not clearly lateralised we collapsed gain effects (Attended minus Ignored) across both

301 stimulation frequencies and correlated these with a hemisphere-collapsed alpha suppression index.

302 This index was quantified as the halved sum of left and right-hemispheric suppression effects as

303 retrieved from significant clusters in the topographical analysis of alpha power differences (Attend

304 Left minus Attend Right), shown in *Figure 3*. Again, we expected a positive correlation here if alpha

305 power suppression influenced phase-locking to visual stimulation. For means of comparison, we

306 repeated this analysis with attention effects on SSR power collapsed across frequencies.

Figure 3 Allocation of spatial attention produces retinotopic alpha power modulation. The scalp map
(top, center) depicts alpha power lateralisation (Attend Left – Attend right conditions) on a dB scale.
Black dots indicate left- and right-hemispheric electrode-clusters that showed a consistent difference
in group statistics (two-tailed cluster-based permutation tests). Left and right spectra illustrate alpha
power differences in respective clusters when the contralateral hemifield was attended (Att) versus
ignored (Ign). The bottom grey inset depicts the distribution of individual alpha power suppression
effects (Ignored minus Attended) within left (L) and right (R) hemisphere clusters in the 8 – 13 Hz
band. Boxplots indicate interquartile ranges (boxes) and medians (coloured vertical intersectors).
Dots below show individual effects (1 dot = 1 participant).

RESULTS

Ongoing alpha power – attentional modulation and lateralisation

The power of the ongoing alpha rhythm lateralised with the allocation of spatial attention to left and

right stimuli. A topographic map of the differences in alpha power between Attend-Left and Attend-

Right conditions shows significant left- and right-hemispheric electrode clusters (*Figure 3*). These

clusters signify retinotopic alpha power modulation when participants attended to left vs right

stimulus positions (right cluster: t_{sum} = -21.454, *P* = 0.026; left cluster: t_{sum} = 81.264, *P* = 0.002). The

differences are further illustrated in power spectra pooled over electrodes of each cluster (*Figure 3*).

As predicted, alpha power at each cluster was lower when participants attended to the contralateral

stimulus. Bayesian inference confirmed the alpha power attention effect for the right (*M* = 0.806 dB,

SEM = 0.216; BF_{10} = 21.17) and left cluster (*M* = 0.790 dB, *SEM* = 0.133; BF_{10} = 906.36). Both effects

were of comparable magnitude (BF_{01} = 4.009 ± 0.007).

SSR power & inter-trial phase locking – attentional modulation

Crucially, we found the opposite pattern when looking at SSRs, i.e. the exact same data but with a

slightly different focus on oscillatory brain activity that was time-locked to the stimulation (compare

formulas 1 and 2): SSRs showed increased power when the respective driving stimulus was attended

versus ignored (*Figure 4*). The power of neural responses evoked by our stimuli (SSRs) was at least

one order of magnitude smaller than ongoing alpha power on average (difference > 10dB, i.e.

between 10 – 100 times). Nevertheless, SSRs could be clearly identified as distinct peaks in (evoked)

power and ITC spectra. Consistent with the retinotopic projection to early visual cortices,

topographical distributions of both measures showed a focal maxima contra-lateral to the respective

stimulus positions that were attended (*Figure 2*). Counter-intuitively though, maximum attention

effects on SSR power did not coincide topographically with sites that showed maximum SSR power

overall (compare scalp maps in *Figure 2 & 4*). Also, due to their rather ipsilateral scalp distributions

(with respect to the attended location), SSR attention effects did not match topographies of

attention-related decreases in ongoing alpha power (compare scalp maps in *Figures 3 & 4*). The 10-

Hz SSR driven by the left-hemifield stimulus showed a left-hemispheric power increase when

attended (t_{sum} = 15.837, *P* = 0.059). Similarly, attention increased the power of the 12-Hz SSR driven

by the right-hemifield stimulus in a right-hemispheric cluster (t_{sum} = 53.282, *P* < 0.001). Bayesian

inference confirmed the attention effect on 10-Hz (*M* = 3.727 dB, *SEM* = 0.919; BF_{10} = 37.05) and 12-

Hz SSR power (*M* = 4.473 dB, *SEM* = 0.841; BF_{10} = 329.75) averaged within clusters. Both effects were

of comparable magnitude (BF_{01} = 3.443 ± 0.005).

SSR phase-locking (quantified as ITCz) also increased with attention to the respective stimulus. In

contrast to evoked power, topographical representations of these effects showed greater overlap

with the sites that showed maximum phase-locking in general (*Figure 4*). For both frequencies, ITCz

increased in central occipital clusters (10 Hz: t_{sum} = 41.351, *P* = 0.004; 12 Hz: t_{sum} = 31.116, *P* = 0.012).

Again, Bayesian inference confirmed the attention effect on 10-Hz (*M* = 1.386 au, *SEM* = 0.297;

BF_{10} = 105.71, one-sided) and 12-Hz ITCz (*M* = 1.824 au, *SEM* = 0.451; BF_{10} = 36.11, one-sided).

Evidence for a greater attention effect on 12-Hz than on 10-Hz ITC remained inconclusive

(BF_{10} = 0.473).

Figure 4 Attention effects on SSR evoked power (evoPow) and SSR inter-trial phase coherence. (**a**) SSR evoked power spectra show systematic power differences at the presentation frequency (10 Hz) of the left stimulus when it was attended (dark red) versus ignored (orange). The inset scalp map illustrates the topographical situation of attention effects. Power spectra were averaged across electrodes (black dots in scalp maps) that showed consistent attention effects in group statistics (two-tailed cluster-based permutation tests) for Attended and Ignored conditions separately. (**b**) Same as in a but for the 12-Hz stimulus presented in the right visual hemifield. (**c,d**) Same as in a,b but using ITCz as a measure of SSR inter-trial phase coherence.

Correlation of alpha and SSR attention effects

Lastly, we tested whether the SSR attention gain effects were mere reflections of the topographically coinciding ipsilateral ongoing alpha power increase during focussed attention that co-occurred with the contralateral ongoing alpha-power decrease (*Figure 3*). Speaking against this account, Bayesian inference provided moderate evidence against the expected positive correlations between the left-hemispheric alpha attention effect and the 10-Hz SSR attention effect (τ_b = -0.221, 95%-CrI = [0.002 0.269]; BF_{01} = 5.811) and between the right-hemispheric alpha attention effect and the 12-Hz SSR attention effect (τ_b = -0.088, 95%-CrI = [0.004 0.315]; BF_{01} = 3.904). These relationships are further illustrated by corresponding linear fits in *Figure 5*.

Following this analysis, we further explored the relationship between spatially non-overlapping decreases in alpha-power contralateral to the attended position and the ipsilateral SSR power gain effects. For the lack of a specific hypothesis about the sign of the correlation in this case, we quantified the evidence for any relationship (two-sided test). The results remained inconclusive for a

correlation between the left-hemispheric alpha attention effect and the right-hemispheric 12-Hz SSR

attention effect (τ_b = 0.235, 95%-CrI = [-0.110 0.487]; BF_{01} = 1.280) and between the right-

hemispheric alpha attention effect and the left-hemispheric 10-Hz SSR attention effect (τ_b = 0.103,

95%-CrI = [-0.218 0.383]; BF_{01} = 2.400).

Figure 5 Relationships between attention effects on alpha power and SSRs. (**a**) Individual 10-Hz (left
stimulus) SSR evoked power gain (Attended minus Ignored; z-scored, y-axis) as a function of alpha
suppression (Ignored minus Attended; z-scored, x axis) in overlapping left-hemispheric parieto-
occipital electrode clusters. Grey dots represent participants. Coloured lines depict a straight line fit
and its confidence interval (dashed lines). Goodness of fit of the linear model provided as R^2 along
with corresponding P-Value. As confirmed by additional tests, both attention effects do not show a
positive linear relationship that would be expected if the ipsilateral SSR power gain effect was a
consequence of the ipsilateral alpha suppression. (**b**) Same as in a, but for the 12 Hz SSR driven by
the right stimulus in overlapping right-hemispheric parieto-occipital electrode clusters. (**c,d**) Similar
to a but for attention-related gain effects on SSR ITCz (z-scored, y-axis) in c and gain effects on SSR
evoked power, both collapsed across electrode clusters showing 10- and 12-Hz SSR attention effects.
Alpha suppression was collapsed across left- and right-hemispheric electrode clusters (see Figure 3).
398

Finally, we repeated this analysis for attention effects on inter-trial phase coherence (ITC). Because

SSR ITC attention effects did not show a clear topographical lateralisation (*Figure 4*), they were

collapsed across driving frequencies (10 & 12 Hz). Again, findings were inconclusive when looking

into the correlation between these aggregate SSR ITC gain effects and a hemisphere-collapsed alpha

suppression index (τ_b = -0.059, 95%-CrI = [-0.349 0.251]; BF_{01} = 2.653). Correlating collapsed attention

effects of SSR evoked power with the same pooled alpha suppression index yielded identical results
regarding the rank correlation (also see linear fits in *Figure 5*).

DISCUSSION

We found that two common spectral measures of alpha-band EEG during alpha-rhythmic visual
stimulation reflect effects of spatial attention with opposite signs. In the following we discuss how
this finding supports the notion of two complementary neural mechanisms governing the cortical
processing of dynamic visual input.

Analysis approach determines sign of attentional modulation

When focussing on the spectral representation of EEG ongoing power, we observed the prototypical
broad peak in the alpha frequency range (8 – 13 Hz; *Figure 2*). Moreover, alpha power decreased
over the hemisphere contralateral to the attended stimulus position, indicating a functional
disinhibition of cortical areas representing task-relevant regions of the visual field (Worden et al.,
2000; Kelly et al., 2006; Thut et al., 2006). Concurrently, alpha power increased over the ipsilateral
hemisphere, actively suppressing irrelevant and possibly distracting input (Rihs et al., 2007; Capilla et
al., 2012).

A second approach focussed on the SSRs, i.e. strictly stimulus-locked rhythmic EEG components. As
in classical frequency-tagging studies, we found spectrally distinct SSRs at the stimulation frequencies
(here 10 and 12 Hz). These two concurrent rhythmic brain responses thus precisely reflected the
temporal dynamics of the visual stimulation. Notably, SSR evoked power was between one to two
orders of magnitude (10 – 100 times) lower than ongoing-alpha power. Smaller evoked power also
explained why SSRs remained invisible in spectra of ongoing activity. They were likely masked by the
broad alpha peak (Figure 2; Covic et al., 2017). Note that this is a result of the relatively low-intensity
stimulation used here. Stimulation of higher intensity can evoke SSRs that are readily visible in power
spectra of ongoing activity (Gulbinaite et al. 2017).

Crucially, we examined SSRs for effects of focused spatial attention. Visual cortical regions
contralateral to the respective driving stimuli showed maximum SSR evoked power. We would
expect to observe a decrease in SSR evoked power with attention (Gulbinaite et al., 2017a; Kizuk and
Mathewson, 2017) under the assumption that SSRs are frequency-specific neural signatures of a local
entrainment of intrinsic alpha generators (Spaak et al., 2014; Notbohm et al., 2016) and exhibit
similar functional characteristics. Instead, we found that SSR evoked power increased in line with
earlier reports (Kim et al., 2007; Kashiwase et al., 2012; Keitel et al., 2013).

Note that the attentional gain effects on SSR evoked power did not coincide topographically with
scalp locations of maximum SSR power, or with scalp locations of the attention-related power
decrease in ongoing alpha oscillations. Instead, they were most pronounced over hemispheres
ipsilateral to the position of the respective driving stimuli. A control analysis (*Figure 5*) showed that
this effect was unrelated to the simultaneously occurring ipsilateral alpha power increase (*Figure 3*).
We have described the apparent counter-intuitive lateralisation of this effect before (Keitel et al.,
2017a) when comparing scalp distributions by means of Attended-minus-Unattended contrasts
(Keitel et al., 2017a) Expecting attention effects to emerge at sites of maximum SSR power in that
case entails the implicit assumption that attention only acts as a local response gain mechanism.
Alternatively, neural representations of attended stimuli could access higher order visual processing
(Lithari et al., 2016) and a gain in spatial extent could then produce seemingly ipsilateral effects when
evaluating topographical differences as observed here (*Figure 4*). Considering that SSR inter-trial
phase coherence showed yet another topographical distribution for gain effects (*Figure 4*), and that
EEG only has a limited spatial resolution, this warrants a dedicated analysis of underlying cortical
sources generating these attention effects in further neuroimaging studies.

Opposite but co-occurring attention effects suggest interplay of distinct attention-related
processes

Our analysis compared attention effects between "ongoing" spectral power within the alpha
frequency band and a quantity termed SSR "evoked power" that is commonly used in frequency
tagging research (Colon et al., 2012; Porcu et al., 2013; Stormer et al., 2014; Walter et al., 2016;
Martinovic and Andersen, 2018). This term is somewhat misleading because it conflates a power
estimate with the consistency of the phase of the SSR across trials of the experiment. Inter-trial
phase consistency (ITC) is closely related to evoked power but involves an extra normalisation term
that abolishes (or at least greatly attenuates) the power contribution[1] (Cohen, 2014; Gross, 2014)
and has been used to quantify SSRs before (Ruhnau et al., 2016).

The effects of attention on SSR evoked power and ITC are typically interchangeable (Covic et al.,
2017; Keitel et al., 2017a). In fact, increased ITC, or phase synchronisation, has been considered the
primary effect of attention on stimulus-driven periodic brain responses (Kim et al., 2007; Kranczioch,
2017). Looking at spectral power and ITC separately, as two distinct aspects of rhythmic brain

---

[1] In a noisy, finite signal such as the typical second(s)-long EEG epoch, there will always be a positive
relationship between the power and inter-trial phase consistency at any frequency as is shown by the greater
than zero noise floor in our ITC spectra (*Figure 4*). Also note that ITC only measures SSRs meaningfully if the
neurophysiological signal contains a periodic component at the stimulation frequency.

activity, therefore resolves the attentional modulation conundrum: Seemingly opposing attention-related effects likely index different but parallel influences on cortical processing of rhythmic visual input. To avoid confusion, we therefore suggest opting for ITC (or related measures, e.g. the cosine similarity index (Chou and Hsu, 2018)) instead of "evoked power" to evaluate SSRs.

Incorporating our findings into an account that regards SSRs primarily as stimulus-driven entrainment of intrinsic alpha rhythms would require demonstrating how a decrease in alpha-band power (i.e. the contralateral alpha suppression) can co-occur with increased SSR phase synchronisation. Alternatively, stimulus-locked ("evoked") and intrinsic alpha rhythms could be considered distinct processes (Freunberger et al., 2009; Sauseng, 2012). Consequentially, alpha range SSRs could predominantly reflect an early cortical mechanism for the tracking of fluctuations in stimulus-specific visual input per se (Keitel et al., 2017a) without the need to assume entrainment (Capilla et al., 2011; Keitel et al., 2014).

The underlying neural mechanism might similarly work for a range of rhythmic and quasi-rhythmic stimuli owing to the fact that visual cortex comprises a manifold of different feature detectors that closely mirror changes along the dimensions of colour, luminance, contrast, spatial frequency and more (Buracas et al., 1998; Blaser et al., 2000; Martinovic and Andersen, 2018). Most importantly, for (quasi-)rhythmic sensory input, attention to the driving stimulus may increase neural phase-locking to the stimulus to allow for enhanced tracking of its dynamics, i.e. increased fidelity. This effect has been observed for quasi-rhythmic low-frequency visual speech signals (Crosse et al., 2015; Park et al., 2016; Hauswald et al., 2018) and task-irrelevant visual stimuli at attended vs ignored spatial locations (Keitel et al., 2017a).

Concurrent retinotopic biasing of visual processing through alpha suppression and stronger neural phase-locking to attended stimuli could therefore be regarded as complimentary mechanisms. Both could act to facilitate the processing of behaviourally relevant visual input in parallel. In this context, SSRs would constitute a special case and easy-to-quantify periodic signature of early visual cortices tracking stimulus dynamics over time. Intrinsic alpha suppression instead may gate the access of sensory information to superordinate visual processing stages (Jensen and Mazaheri, 2010; Zumer et al., 2014) and enhanced ipsilateral alpha power may additionally attenuate irrelevant and possibly distracting stimuli at ignored locations (Capilla et al., 2012).

A neuronal implementation may work like this: During rest or inattention, occipital neuronal populations synchronise with a strong internal, thalamo-cortical pacemaker (alpha). During attentive processing of sensory input, retinotopic alpha suppression releases specific neuronal sub-populations from an internal reign and allows them to track the stimulus dynamics at attended locations. A

related mechanism has been observed in the striatum, where local field potentials are dominated by synchronous oscillatory activity across large areas (Courtemanche et al., 2003). However, during task performance focal neuronal populations were found to disengage from this global synchronicity in a consistent and task-specific manner. At the level of EEG/MEG recordings, such a mechanism could lead to task-related decrease of oscillatory power but increase of coherence or ITC, as observed in the current study and previously in the sensorimotor system (Gross et al., 2005; Schoffelen et al., 2005; Schoffelen et al., 2011).

Whereas such an account challenges the occurrence of strictly stimulus-driven alpha entrainment, it may still allow alpha to exert temporally precise top-down influences during predictable and behaviourally relevant rhythmic stimulation – a process that itself could be subject to entrainment (Thut et al., 2011; Nobre et al., 2012; Haegens and Zion Golumbic, 2018; Zoefel et al., 2018).

Conclusion

Our findings reconcile seemingly contradictory findings regarding spatial attention effects on alpha-rhythmic activity, assumed to be entrained by periodic visual stimulation, and SSRs. Focusing on spectral power or phase consistency of the EEG during visual stimulation yielded reversed attention effects in the same dataset. Our findings encourage a careful and consistent choice of measures of ongoing brain dynamics (here power) or measures of stimulus-related activity (here ITC), that should be critically informed by the experimental question, when studying the effects of visuo-spatial selective attention on the cortical processing of dynamic (quasi-) rhythmic visual stimulation. Again, we emphasise that both common data analysis approaches taken here can be equally valid and legitimate, yet they likely represent distinct neural phenomena. These can occur simultaneously, as in our case, and may index distinct cortical processes that work in concert to facilitate the processing of visual stimulation at attended locations.

Competing interests

The authors declare no competing interests.

Author contributions

CK designed research, performed research, analysed data and wrote the article. JG designed research analysed data and wrote the article. AK, CSYB, CD and GT designed research and wrote the article.

528 **Data accessibility**

529 EEG data, pre-processed in Fieldtrip format, that underlie all analyses reported here and a

530 corresponding MATLAB analysis script are available on the Open Science Framework, osf.io/apsyf

531 (Keitel et al., 2017b).

532
533 **References**
534
Adrian ED, Matthews BH (1934) The interpretation of potential waves in the cortex. J Physiol 81:440-

471.

Andersen SK, Muller MM (2015) Driving steady-state visual evoked potentials at arbitrary frequencies

using temporal interpolation of stimulus presentation. BMC Neurosci 16:95.

Benwell CSY, Keitel C, Harvey M, Gross J, Thut G (2017a) Trial-by-trial co-variation of pre-stimulus

EEG alpha power and visuospatial bias reflects a mixture of stochastic and deterministic

effects. Eur J Neurosci.

Benwell CSY, Tagliabue CF, Veniero D, Cecere R, Savazzi S, Thut G (2017b) Prestimulus EEG Power

Predicts Conscious Awareness But Not Objective Visual Performance. eNeuro 4.

Blaser E, Pylyshyn ZW, Holcombe AO (2000) Tracking an object through feature space. Nature

408:196-199.

Bonnefond M, Jensen O (2012) Alpha oscillations serve to protect working memory maintenance

against anticipated distracters. Curr Biol 22:1969-1974.

Buracas GT, Zador AM, DeWeese MR, Albright TD (1998) Efficient discrimination of temporal patterns

by motion-sensitive neurons in primate visual cortex. Neuron 20:959-969.

Capilla A, Pazo-Alvarez P, Darriba A, Campo P, Gross J (2011) Steady-state visual evoked potentials

can be explained by temporal superposition of transient event-related responses. PLoS One

6:e14543.

Capilla A, Schoffelen J-M, Paterson G, Thut G, Gross J (2012) Dissociated α-Band Modulations in the

Dorsal and Ventral Visual Pathways in Visuospatial Attention and Perception. Cerebral

Cortex.

Chou EP, Hsu SM (2018) Cosine similarity as a sample size-free measure to quantify phase clustering

within a single neurophysiological signal. J Neurosci Methods 295:111-120.

Cohen MX (2014) Analyzing neural time series data: theory and practice. Cambridge, Massachusetts:

MIT Press.

Colon E, Nozaradan S, Legrain V, Mouraux A (2012) Steady-state evoked potentials to tag specific

components of nociceptive cortical processing. Neuroimage 60:571-581.

Courtemanche R, Fujii N, Graybiel AM (2003) Synchronous, focally modulated beta-band oscillations

characterize local field potential activity in the striatum of awake behaving monkeys.

Journal of Neuroscience 23:11741-11752.

Covic A, Keitel C, Porcu E, Schroger E, Muller MM (2017) Audio-visual synchrony and spatial attention

enhance processing of dynamic visual stimulation independently and in parallel: A

frequency-tagging study. Neuroimage 161:32-42.

Crosse MJ, Butler JS, Lalor EC (2015) Congruent Visual Speech Enhances Cortical Entrainment to

Continuous Auditory Speech in Noise-Free Conditions. J Neurosci 35:14195-14204.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics

including independent component analysis. J Neurosci Methods 134:9-21.

Ferree TC (2006) Spherical splines and average referencing in scalp electroencephalography. Brain

Topogr 19:43-52.

Freunberger R, Fellinger R, Sauseng P, Gruber W, Klimesch W (2009) Dissociation between phase-

locked and nonphase-locked alpha oscillations in a working memory task. Hum Brain Mapp

30:3417-3425.

Groppe DM, Bickel S, Keller CJ, Jain SK, Hwang ST, Harden C, Mehta AD (2013) Dominant frequencies

of resting human brain activity as measured by the electrocorticogram. Neuroimage

79:223-233.

Gross J (2014) Analytical methods and experimental approaches for electrophysiological studies of

brain oscillations. J Neurosci Methods 228:57-66.

Gross J, Pollok B, Dirks M, Timmermann L, Butz M, Schnitzler A (2005) Task-dependent oscillations

during unimanual and bimanual movements in the human primary motor cortex and SMA

studied with magnetoencephalography. Neuroimage 26:91-98.

Gulbinaite R, Roozendaal D, VanRullen R (2017a) Attention effects on steady-state visual evoked

potentials in response to 3-80 Hz flicker. Journal of Vision 17:977-977.

Gulbinaite R, van Viegen T, Wieling M, Cohen MX, VanRullen R (2017b) Individual Alpha Peak

Frequency Predicts 10 Hz Flicker Effects on Selective Attention. J Neurosci 37:10173-10184.

Haegens S, Zion Golumbic E (2018) Rhythmic facilitation of sensory processing: A critical review.

Neurosci Biobehav Rev 86:150-165.

Hauswald A, Lithari C, Collignon O, Leonardelli E, Weisz N (2018) A Visual Cortical Network for

Deriving Phonological Information from Intelligible Lip Movements. Curr Biol 28:1453-1459

e1453.

Herring JD, Thut G, Jensen O, Bergmann TO (2015) Attention Modulates TMS-Locked Alpha

Oscillations in the Visual Cortex. J Neurosci 35:14435-14447.

Herrmann CS (2001) Human EEG responses to 1-100 Hz flicker: resonance phenomena in visual
cortex and their potential correlation to cognitive phenomena. Exp Brain Res 137:346-353.

Herrmann CS, Munk MH, Engel AK (2004) Cognitive functions of gamma-band activity: memory
match and utilization. Trends Cogn Sci 8:347-355.

Iemi L, Chaumon M, Crouzet SM, Busch NA (2017) Spontaneous Neural Oscillations Bias Perception
by Modulating Baseline Excitability. J Neurosci 37:807-819.

JASP-Team (2018) JASP. In, 0.8.2 Edition.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: gating by
inhibition. Front Hum Neurosci 4:186.

Kashiwase Y, Matsumiya K, Kuriki I, Shioiri S (2012) Time courses of attentional modulation in neural
amplification and synchronization measured with steady-state visual-evoked potentials. J
Cogn Neurosci 24:1779-1793.

Kayser J, Tenke CE (2015) On the benefits of using surface Laplacian (current source density)
methodology in electrophysiology. Int J Psychophysiol 97:171-173.

Keitel A, Gross J (2016) Individual Human Brain Areas Can Be Identified from Their Characteristic
Spectral Activation Fingerprints. PLoS Biology 14:e1002498.

Keitel C, Quigley C, Ruhnau P (2014) Stimulus-Driven Brain Oscillations in the Alpha Range:
Entrainment of Intrinsic Rhythms or Frequency-Following Response? Journal of
Neuroscience 34:10137-10140.

Keitel C, Thut G, Gross J (2017a) Visual cortex responses reflect temporal structure of continuous
quasi-rhythmic sensory stimulation. Neuroimage 146:58-70.

Keitel C, Andersen SK, Quigley C, Muller MM (2013) Independent Effects of Attentional Gain Control
and Competitive Interactions on Visual Stimulus Processing. Cerebral Cortex 23:940-946.

Keitel C, Benwell CSY, Thut G, Gross J (2017b) Alpha during quasi-periodic visual stimulation. In. Open
Science Framework.

Keitel C, Benwell CSY, Thut G, Gross J (2018) No changes in parieto-occipital alpha during neural
phase locking to visual quasi-periodic theta-, alpha-, and beta-band stimulation. Eur J
Neurosci.

Kelly SP, Lalor EC, Reilly RB, Foxe JJ (2006) Increases in alpha oscillatory power reflect an active
retinotopic mechanism for distracter suppression during sustained visuospatial attention. J
Neurophysiol 95:3844-3851.

Kim YJ, Grabowecky M, Paller KA, Muthu K, Suzuki S (2007) Attention induces synchronization-based
response gain in steady-state visual evoked potentials. Nat Neurosci 10:117-125.

Kizuk SA, Mathewson KE (2017) Power and Phase of Alpha Oscillations Reveal an Interaction between
Spatial and Temporal Visual Attention. J Cogn Neurosci 29:480-494.
Kranczioch C (2017) Individual differences in dual-target RSVP task performance relate to
entrainment but not to individual alpha frequency. PLoS One 12:e0178934.
Lithari C, Sanchez-Garcia C, Ruhnau P, Weisz N (2016) Large-scale network-level processes during
entrainment. Brain Res 1635:143-152.
Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J Neurosci
Methods 164:177-190.
Martinovic J, Andersen SK (2018) Cortical summation and attentional modulation of combined
chromatic and luminance signals. Neuroimage 176:390-403.
Mathewson KE, Prudhomme C, Fabiani M, Beck DM, Lleras A, Gratton G (2012) Making waves in the
stream of consciousness: entraining oscillations in EEG alpha and fluctuations in visual
awareness with rhythmic visual stimulation. J Cogn Neurosci 24:2321-2333.
Morgan ST, Hansen JC, Hillyard SA (1996) Selective attention to stimulus location modulates the
steady-state visual evoked potential. Proceedings of the National Academy of Sciences of
the United States of America 93:4770-4774.
Müller MM, Teder-Sälejärvi W, Hillyard SA (1998) The time course of cortical facilitation during cued
shifts of spatial attention. Nat Neurosci 1:631-634.
Nobre AC, Rohenkohl G, Stokes M (2012) Nervous anticipation: top-down biasing across space and
time. In: Cognitive Neuroscience of Attention, 2 Edition (Posner MI, ed), pp 159-186. New
York: Guilford.
Nolan H, Whelan R, Reilly RB (2010) FASTER: Fully Automated Statistical Thresholding for EEG artifact
Rejection. J Neurosci Methods 192:152-162.
Norcia AM, Appelbaum LG, Ales JM, Cottereau BR, Rossion B (2015) The steady-state visual evoked
potential in vision research: A review. J Vis 15:4.
Notbohm A, Kurths J, Herrmann CS (2016) Modification of Brain Oscillations via Rhythmic Light
Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related
Responses. Front Hum Neurosci 10:10.
Oostenveld R, Praamstra P (2001) The five percent electrode system for high-resolution EEG and ERP
measurements. Clin Neurophysiol 112:713-719.
Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for advanced
analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci
2011:156869.

Park H, Kayser C, Thut G, Gross J (2016) Lip movements entrain the observers' low-frequency brain
663   oscillations to facilitate speech intelligibility. Elife 5.

664 Perrin F, Pernier J, Bertrand O, Giard MH, Echallier JF (1987) Mapping of scalp potentials by surface
665   spline interpolation. Electroencephalogr Clin Neurophysiol 66:75-81.

666 Picton TW, John MS, Purcell DW, Plourde G (2003) Human auditory steady-state responses: the
667   effects of recording technique and state of arousal. Anesth Analg 97:1396-1402.

668 Porcu E, Keitel C, Muller MM (2013) Concurrent visual and tactile steady-state evoked potentials
669   index allocation of inter-modal attention: a frequency-tagging study. Neurosci Lett 556:113-
670   117.

671 Regan D (1966) Some characteristics of average steady-state and transient responses evoked by
672   modulated light. Electroencephalogr Clin Neurophysiol 20:238-248.

673 Rihs TA, Michel CM, Thut G (2007) Mechanisms of selective inhibition in visual spatial attention are
674   indexed by alpha-band EEG synchronization. Eur J Neurosci 25:603-610.

675 Rouder JN, Morey RD, Speckman PL, Province JM (2012) Default Bayes factors for ANOVA designs.
676   Journal of Mathematical Psychology 56:356-374.

677 Rouder JN, Speckman PL, Sun DC, Morey RD, Iverson G (2009) Bayesian t tests for accepting and
678   rejecting the null hypothesis. Psychonomic Bulletin & Review 16:225-237.

679 Ruhnau P, Keitel C, Lithari C, Weisz N, Neuling T (2016) Flicker-Driven Responses in Visual Cortex
680   Change during Matched-Frequency Transcranial Alternating Current Stimulation. Front Hum
681   Neurosci 10:184.

682 Samaha J, Iemi L, Postle BR (2017) Prestimulus alpha-band power biases visual discrimination
683   confidence, but not accuracy. Conscious Cogn 54:47-55.

684 Samaha J, Bauer P, Cimaroli S, Postle BR (2015) Top-down control of the phase of alpha-band
685   oscillations as a mechanism for temporal prediction. Proc Natl Acad Sci U S A 112:8439-
686   8444.

687 Sauseng P (2012) Brain oscillations: phase-locked EEG alpha controls perception. Curr Biol 22:R306-
688   308.

689 Schoffelen JM, Oostenveld R, Fries P (2005) Neuronal coherence as a mechanism of effective
690   corticospinal interaction. Science 308:111-113.

691 Schoffelen JM, Poort J, Oostenveld R, Fries P (2011) Selective movement preparation is subserved by
692   selective increases in corticomuscular gamma-band coherence. J Neurosci 31:6750-6758.

693 Schonbrodt FD, Wagenmakers EJ (2017) Bayes factor design analysis: Planning for compelling
694   evidence. Psychon Bull Rev.

Spaak E, de Lange FP, Jensen O (2014) Local entrainment of alpha oscillations by visual stimuli causes

cyclic modulation of perception. J Neurosci 34:3536-3544.

Stormer VS, Alvarez GA, Cavanagh P (2014) Within-hemifield competition in early visual areas limits

the ability to track multiple objects with attention. J Neurosci 34:11526-11533.

Tallon-Baudry C, Bertrand O, Delpuech C, Pernier J (1996) Stimulus specificity of phase-locked and

non-phase-locked 40 Hz visual responses in human. J Neurosci 16:4240-4249.

Tallon-Baudry C, Bertrand O, Peronnet F, Pernier J (1998) Induced gamma-band activity during the

delay of a visual short-term memory task in humans. J Neurosci 18:4244-4254.

Thut G, Schyns PG, Gross J (2011) Entrainment of perceptually relevant brain oscillations by non-

invasive rhythmic stimulation of the human brain. Front Psychol 2:170.

Thut G, Nietzel A, Brandt SA, Pascual-Leone A (2006) Alpha-band electroencephalographic activity

over occipital cortex indexes visuospatial attention bias and predicts visual target detection.

J Neurosci 26:9494-9502.

van Diepen RM, Mazaheri A (2018) The Caveats of observing Inter-Trial Phase-Coherence in Cognitive

Neuroscience. Sci Rep 8:2990.

Vialatte FB, Maurice M, Dauwels J, Cichocki A (2010) Steady-state visually evoked potentials: focus on

essential paradigms and future perspectives. Prog Neurobiol 90:418-438.

Wagenmakers EJ, Wetzels R, Borsboom D, van der Maas HL (2011) Why psychologists must change

the way they analyze their data: the case of psi: comment on Bem (2011). J Pers Soc Psychol

100:426-432.

Walter S, Keitel C, Muller MM (2016) Sustained Splits of Attention within versus across Visual

Hemifields Produce Distinct Spatial Gain Profiles. J Cogn Neurosci 28:111-124.

Walter WG, Dovey VJ, Shipton H (1946) Analysis of the Electrical Response of the Human Cortex to

Photic Stimulation. Nature 158:540-541.

Worden MS, Foxe JJ, Wang N, Simpson GV (2000) Anticipatory biasing of visuospatial attention

indexed by retinotopically specific alpha-band electroencephalography increases over

occipital cortex. J Neurosci 20:RC63.

Zauner A, Fellinger R, Gross J, Hanslmayr S, Shapiro K, Gruber W, Muller S, Klimesch W (2012) Alpha

entrainment is responsible for the attentional blink phenomenon. Neuroimage 63:674-686.

Zoefel B, Ten Oever S, Sack AT (2018) The Involvement of Endogenous Neural Oscillations in the

Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural Responses.

Front Neurosci 12:95.

Zumer JM, Scheeringa R, Schoffelen JM, Norris DG, Jensen O (2014) Occipital alpha activity during

stimulus processing gates the information flow to object-selective cortex. PLoS Biology

12:e1001965.