Broad geographic sampling reveals predictable, pervasive, and strong seasonal adaptation in *Drosophila*

Heather E. Machado[1,2]*‡, Alan O. Bergland[1,3]*‡, Ryan Taylor[1], Susanne Tilk[1], Emily Behrman[4], Kelly Dyer[5], Daniel K. Fabian[6,7], Thomas Flatt[6,8], Josefa González[9], Talia L. Karasov[10], Bernard Kim[1], Iryna Kozeretska[11,12], Brian P. Lazzaro[13], Thomas JS Merritt[14], John E. Pool[15], Katherine O'Brien[4], Subhash Rajpurohit[4], Paula R. Roy[16], Stephen W. Schaeffer[17], Svitlana Serga[11,12], Paul Schmidt[4o‡], Dmitri A. Petrov[1o‡]

* equal contribution
° equal contribution
‡ to whom correspondence should be addressed: heather.machado@sanger.ac.uk, aob2x@virginia.edu, schmidtp@upenn.edu, dpetrov@stanford.edu

[1] Department of Biology, Stanford University, Stanford, CA, United States
[2] Wellcome Sanger Institute, Hinxton CB10 1SA, United Kingdom
[3] Department of Biology, University of Virginia, Charlottesville, VA, United States
[4] Department of Biology, University of Pennsylvania, Philadelphia, PA, United States
[5] Department of Genetics, University of Georgia, Athens, GA, United States
[6] Institute of Population Genetics, Vetmeduni Vienna, A-1210 Vienna, Austria
[7] Centre for Pathogen Evolution, Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, United Kingdom
[8] Department of Biology, University of Fribourg, CH-1700 Fribourg, Switzerland
[9] Institute of Evolutionary Biology, CSIC- Universitat Pompeu Fabra, Barcelona, Spain
[10] Department of Biology, University of Utah, Salt Lake City, UT, United States
[11] Taras Shevchenko National University of Kyiv, Kyiv, Ukraine
[12] National Antarctic Scientific Centre of Ukraine, Taras Shevchenko Blvd., 01601, Kyiv, Ukraine
[13] Department of Entomology, Cornell University, Ithaca, NY, United States
[14] Department of Chemistry & Biochemistry, Laurentian University, Sudbury, ON, Canada
[15] Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI, United States
[16] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, United States
[17] Department of Biology, The Pennsylvania State University, University Park, PA, United States

Abstract

To advance our understanding of adaptation to temporally varying selection pressures, we identified

signatures of seasonal adaptation occurring in parallel among *Drosophila melanogaster* populations.

Specifically, we estimated allele frequencies genome-wide from flies sampled early and late in the

growing season from 20 widely dispersed populations. We identified parallel seasonal allele

frequency shifts across North America and Europe, demonstrating that seasonal adaptation is a

general phenomenon of temperate fly populations. Seasonally fluctuating polymorphisms are enriched

in large chromosomal inversions and we find a broad concordance between seasonal and spatial allele

frequency change. The direction of allele frequency change at seasonally variable polymorphisms can

be predicted by weather conditions in the weeks prior to sampling, linking the environment and the

genomic response to selection. Our results suggest that fluctuating selection is an important

evolutionary force affecting patterns of genetic variation in *Drosophila*.

Introduction

Fluctuations in the environment are an inescapable condition for all organisms. While many of these

fluctuations are entirely unpredictable, some are predictable to a degree, including those that occur on

diurnal and seasonal time scales. The predictability of cyclic fluctuations is reflected in the fact that

many species exhibit plastic physiological and behavioral strategies that enable them to survive the

unfavorable season and exploit the favorable one (Denlinger, 2002; Kostál, 2006); such plastic

responses represent the classical form of seasonal adaptation (Tauber et al., 1986). However,

seasonally varying selection can - in principle - maintain fitness-related genetic variation if some

genotypes have high fitness in one season but not another (Gillespie, 1973; Haldane and Jayakar,

1963). Thus, in organisms that undergo multiple generations per year, a distinct form of seasonal

adaptation occurs when the frequency of alternate genotypes changes in response to seasonal

fluctuations in the environment.

Seasonal adaptation can be seen as a form of local adaptation- local adaptation in time to temporally

varying selection pressures rather than in space. Such adaptation in time has often been considered to

be uncommon and, when present, unlikely to result in long-term balancing selection (Ewing, 1979;

Hedrick, 1976). Classic quantitative genetic theory suggests that an optimal, plastic genotype will

eventually dominate a population that is exposed to periodically changing environments (Bürger and

Gimelfarb, 1999; Korol et al., 1996; Lande, 2008; Scheiner, 1993). This is particularly so when

certain environmental cues are reliable indicators of changes in selection pressure (Levins, 1974; Via

and Lande, 1985). Predictions from traditional population genetic models suggest that periodically

changing environments will lead to the rapid loss of seasonally favored ecotypes as slight changes in

selection pressure from one year to another eventually push allele frequencies at causal alleles to

fixation (Hedrick, 1976).

Recent theoretical models have called these classical predictions into question. For instance, a

quantitative genetic model by Botero and colleagues (Botero et al., 2015) examined whether adaptive

tracking, plasticity, or bet-hedging evolve as a consequence of environmental fluctuations. They

showed that adaptive tracking rather than plasticity is likely to become a predominant feature of

populations living in seasonally fluctuating environments when populations undergo more than three

generations per season. Additionally, a population genetic model by Wittmann and colleagues

(Wittmann et al., 2017) has demonstrated that seasonally varying selection can indeed maintain

fitness related genetic variation at many loci throughout the genome provided that dominance shifts

from season to season in such a way that, on average, the seasonally favored allele remains slightly

dominant (Curtsinger et al., 1994). These recent models, along with others that highlight the

importance of population cycles (Bertram and Masel, 2019), as well as overlapping generations and

age structure (Bertram and Masel, 2019; Ellner, 1996; Ellner and Hairston, 1994; Ellner and Sasaki,

1996), suggest that seasonal adaptation and adaptive tracking (Kain et al., 2015) could be an

important feature of organisms such as *Drosophila* that have multiple generations per year (Behrman

et al., 2015). More generally, it is possible that adaptive tracking of environmental fluctuations on

rapid time scales might be more common than generally acknowledged and has been hidden from us

due to the difficulty of detecting such adaptive tracking reliably (Lynch and Ho, 2020) and from the

lack of finely resolved temporal data (Buffalo and Coop, 2020).

Despite the lack of theoretical consensus on whether and how seasonal adaptation operates, there is

substantial empirical evidence for seasonal adaptation in many organisms including *Drosophila*.

Seasonal adaptation was first observed in *D. pseudoobscura* by Dobzhansky and colleagues (e.g.,

Dobzhansky, 1943) by tracking frequencies of inversion polymorphisms over seasons. Later studies

confirmed and extended these early findings to other species including *D. melanogaster* (Kapun et al.,

2016; Stalker, 1980, 1976) and *D. subobscura* (Rodríguez-Trelles et al., 2013).

In *D. melanogaster*, multiple additional lines of evidence from phenotypic and genetic analyses

demonstrate the presence of seasonal adaptation. When reared in a common laboratory environment,

flies collected in winter or spring (close descendants of flies that successfully overwintered) show

higher stress tolerance (Behrman et al., 2015), greater propensity to enter reproductive dormancy

(Schmidt and Conde, 2006), increased innate immune function (Behrman et al., 2018), and modulated

cuticular hydrocarbon profiles (Rajpurohit et al., 2017) as compared to flies collected in the fall (the

descendants of those flies who prospered during the summer). Rapid adaptation over seasonal time

scales in these and related phenotypes has also been observed in laboratory (Schmidt and Conde,

2006) and field-based mesocosm experiments (Erickson et al., 2020; Grainger et al. 2021; Rajpurohit

et al., 2018, 2017; Rudman et al., 2019). Genome-wide analysis indicated that a large number of

common polymorphisms change in frequency over seasonal time scales in one mid-latitude orchard

(Bergland et al., 2014), and allele frequency change among seasons has been observed using

candidate gene approaches (Cogni et al., 2014). In several cases, these adaptively oscillating

polymorphisms have been linked to seasonally varying phenotypes (Behrman et al., 2018; Erickson et

al., 2020; Paaby et al., 2014).

Despite ample evidence of seasonal adaptation in *D. melanogaster*, many aspects of this system

remain unexplored. First, we do not know whether seasonal adaptation is a general feature of *D.*

melanogaster populations across its range. Previous work (Schmidt and Conde 2006, Bergland et al.

2014, Cogni et al. 2014, Behrman et al. 2015, Behrman et al. 2018) detected seasonal allele frequency

fluctuations in a single locality over the span of one to three years. These results need to be confirmed

at other locations. Second, as many ecological and environmental variables covary with season, it is

unclear which specific factors drive the observed changes in phenotype and genotype in populations

sampled over seasonal time. One straightforward and statistically powerful way to start answering

these questions is to assess the extent of parallel shifts in allele frequencies over seasonal time in

many populations across the *D. melanogaster* range. Should such parallel shifts be detected we can

then infer that seasonal adaptation is widespread and at least partly driven by a common set of

variants. We can also attempt to associate the magnitude and direction of allelic shifts with local

environmental conditions (e.g., temperature) that may vary among populations sampled at equivalent

points in seasonal time. Broad sampling will also allow us to determine the magnitude of the shared

seasonal fluctuations and give us a glimpse into the genetic architecture of seasonal adaptation: for

instance, whether or not the seasonally-responsive variants are found genome-wide or clustered into

linked blocks or even supergenes.

Here we carry out such a study by using allele frequency data from 20 paired seasonal samples from

15 localities in North America and Europe. First, we demonstrate that seasonal adaptation is a general

phenomenon that occurs in multiple populations of *D. melanogaster* on two continents by providing

evidence that at least some of the same polymorphisms cycle between seasons in populations sampled

across vast geographic distances. Seasonal alleles tend to show clinal variation, with the alleles that

increase in frequency through the summer generally being more frequent in lower latitude (more

"summer-like") locations. We also show that seasonal signal is enriched near inversion breakpoints or

within inverted regions on chromosome 2L, 3L, and 3R, although these inverted regions do not

account for all of the signal that we observe. Further, we show that allele frequency change between

seasons is most predictable when taking into account the local temperature prior to sample collections

both in the spring and in the fall, which hints at the complex and nonlinear nature of adaptation with

adaptive tracking occurring at sub-seasonal timescales. Taken together, our work demonstrates that

seasonal adaptation is a general and predictable feature of *D. melanogaster* populations and has

pervasive effects on patterns of allele frequencies genome-wide. More generally we establish that

metazoan populations can exhibit polygenic adaptive tracking of environmental conditions on

extremely short time scales, most likely shorter than a single growing season encompassing ~10

generations.

<u>***Results***</u>

Fly samples and sequence data

We assembled 73 samples of *D. melanogaster* collected from 23 localities in North America and

Europe (Supplemental Table 1, Supplemental Figure 1). For 15 sampling localities, flies were

collected in the spring and fall over the course of one to six years (Figure 1A). Our sampling and

sequencing efforts also complement other genome-wide datasets that investigate genetic variation

within and among *D. melanogaster* populations throughout their range (Campo et al., 2013; Fabian et

al., 2012; Grenier et al., 2015; Kao et al., 2015; Kapun et al., 2020; Kim et al., 2014; Kolaczkowski et

al., 2011; Lack et al., 2016; Mackay et al., 2012; Pool et al., 2012; Reinhardt et al., 2014; Svetec et al.,

2016; Zhao and Begun, 2017).

For our analysis, we divided our samples into two subsets (Figure 1B). The first subset ('Core20') is

composed of 20 pairs of spring and fall samples from the same location. Samples in the Core20 set

are drawn from 15 localities in North America and Europe and we use at most two years of sampling

from any one locality. The second subset of samples ('Clinal') was used to examine patterns of clinal

variation along the East Coast of North America and consists of four populations sampled in the

spring (see Material and Methods and Supplemental Table 1).

To estimate allele frequencies genome-wide we performed pooled DNA sequencing of multiple

individual flies per population (Schlötterer et al., 2014; Zhu et al., 2012). For each sample, we

resequenced pools of ~75 individuals (range 27-164) to an average of 94x coverage (range 22-220,

Supplemental Table 1). Analyses presented here use a total of 43 samples in the Core20 and Clinal

sets described above. Data from the remaining 30 samples, including those that do not constitute

paired spring/fall samples, are not included in our analyses here but nonetheless form a part of our

larger community-based sampling and resequencing effort and are deposited with the rest of the data

(NCBI SRA; BioProject Accession #PRJNA308584; accession numbers for each sample can be found

in Supplemental Table 1).

Figure 1. Sampling times, localities, and basic population structure of samples used in this study. (A) Distribution of collection times during the spring (red) and fall (blue) in relation to latitude. For samples where the collection month, but not day, was recorded, we specified the 15th of the month to calculate Julian Day. (B) Sampling localities for the primary seasonal analysis ('Core20': green) and the latitudinal cline analysis ('Cline': red), distributed across North America and Europe. Numbers represent the number of years of spring/fall comparisons at that locality. (C) Principal components analysis of SNP allele frequencies. Circles are placed around samples from three geographic regions: Europe (EU), California (CA) and Eastern North America (East NA); points are colored by latitude and shapes represent seasons, as defined by collection time.

Pooled resequencing identified ~1.75M SNPs following quality control and filtering for read depth

and average minor allele frequency (minor allele frequency > 0.01). We applied a second filtering and

retained ~775,000 SNPs that have observed polymorphism in each population sampled. This biases us

towards assessing fluctuations at higher frequency polymorphisms (Supplemental Figure 2). Because

this dataset represents approximately half of all detected SNPs, the conclusions drawn from its

analysis are representative of genetic variation in *D. melanogaster*.

To gain insight into basic patterns of differentiation among the sampled populations, we performed a

principal component (PC) analysis across all samples. Samples cluster by locality and geography

(Figure 1C) with PC 1 separating European, North American West Coast and East Coast populations

while PC 2 separates the eastern North America samples by latitude (linear regression $p = 3*10^{-15}$, R^2

= 0.84). No PC is associated with season, following correction for multiple testing, implying that the

spatial differentiation dominates patterns of SNP variation across these samples.

Seasonal adaptation is a general feature of *D. melanogaster* populations

Our first objective was to determine whether or not we find evidence of parallel seasonal cycling of

allele frequencies across multiple populations. To identify a signal of parallel allele frequency change

over time we assessed the per-SNP change in allele frequency between spring and fall among the

Core20 set of populations using three complementary methods that rely on different statistical models.

For all three methods, we contrast the observed signal to permutations in which we shuffled the

seasonal labels within each spring-fall paired sample. With the exception of the seasonal labels within

paired samples, these permutations have the advantage of retaining all of the features of the data

including patterns of genetic covariation and all of the noise in the allele frequency estimation.

We first used a generalized linear model (GLM), regressing allele frequency at each SNP on season

across all of the populations. We find that the observed genome-wide distribution of 'seasonal' *p*-

values from this model is not uniform and there is a pronounced excess of SNPs with low *p*-values

(Figure 2A). The observed data have a stronger seasonal signal than most permutations. For example,

at a *p*-value of ~$4*10^{-3}$, which corresponds to the top 1% (Supplemental Table 2), the observed data

have a greater number of seasonal SNPs than 96.8% of the permutations (Figure 2B). This enrichment

is robust to the choice of quantile (Supplemental Figure 4). Seasonal SNPs identified by Bergland and

colleagues (2014) are slightly enriched among the top 1% of SNPs identified using the GLM, relative

to matched genomic controls, after excluding the overlapping Pennsylvanian populations (log$_2$ odds

ratio $\pm$ SD = 0.59 $\pm$ 0.37, p_{perm} = 0.0512). Our top set of seasonally variable SNPs shift in frequency

by ~4-8% (Fig. 2C). Taken at face value this corresponds to effective selection coefficients of ~10-

30% per season (~1-3% per generation assuming 10 generations per season).

Figure 2. **Signals of seasonal adaptation.** (A) *P*-value distribution of GLM seasonal regression (red line), permutations (solid black lines, 50/500 plotted), and expected values (dashed black line). (B) Comparison of seasonal test statistics (quantile=0.01) for observed (red) and permuted (black) datasets, for the three analyses: i: GLM (-log p-value), ii: Bayenv (Bayes factor; outlier of a Bayes factor of 10 excluded from plotting), iii: RFM (X^2). (C) Average spring/fall allele frequency change for each of the top 1% of seasonally varying SNPs (red) and non-seasonal, matched control SNPs (black), as a function of the folded spring allele frequency. Lines represent a moving average across SNPs with a given spring allele frequency. (D) Enrichment of seasonal SNPs (median of 500 permutations). Enrichment is calculated as the observed number of seasonal SNPs over the number of seasonal SNPs in each permutation at P<0.004. The genome is subset by chromosome and location relative to major cosmopolitan inversions (bkpts.: inversion breakpoints +/- 1Mb, inside: interior to the breakpoints excluding 1Mb buffer, outside: exterior to the breakpoints excluding the 1Mb

buffers). Error bars are 95% confidence intervals and asterisks denote greater seasonal enrichment than more than 95% of permutations. (E/F) Concordance rates of seasonal and clinal polymorphisms by E) geographic region and F) genomic region. The y-axis represents the concordance rate of allele frequency change across time and space, assuming that the winter favored allele is the same as the allele favored in high latitudes. Clinal polymorphisms were identified along the East Coast of North America. Seasonal sites were identified using all populations not used in the clinal analysis (n=18). E) Seasonal sites were also identified using exclusively the California populations (purple: n=3) or the Europe populations (green: n=3). Shaded areas are the 95% confidence intervals of 100 matched control datasets for the Californian (purple) and European (green) analyses. F) Concordance by genomic region. Error bars are 95% confidence intervals (binomial error) and asterisks denote concordance significantly greater than 0.5 (binomial test p<0.05).

Next, we performed a parallel analysis using a Bayesian model, Bayenv, which uses a population

covariance matrix to account for structure among samples (Günther and Coop, 2013). We found an

enrichment of seasonal SNPs at high test statistic values (quantile = 0.01: Bayes factor = 2.6), with a

greater enrichment of seasonal SNPs compared with 94% of permutations (Figure 2B). The set of

seasonal SNPs identified in the Bayenv analysis shows substantial overlap with the GLM set.

Specifically, for the top 1% of SNPs in the GLM analysis, 20% are also in the top 1% of the Bayenv

analysis, compared with a naive expectation of 1% (Supplemental Figure 4).

Finally, we developed a method derived from combining ranked-normalized *p*-values from Fisher's

Exact Tests that compares allele frequencies within each pair of seasonal samples (see Materials and

Methods). We again find a deviation from the expected test statistic distribution indicating an excess

of seasonally varying polymorphisms (Figure 2B) with the observed enrichment of high scores

(quantile = 0.01: X^2 = 32) to be greater than more than 98% of permutations. These three analyses,

using entirely distinct statistical machinery and assumptions, together provide strong evidence of the

genome-wide excess of seasonal SNPs indicating a robust signal of parallel seasonal adaptation across

multiple populations.

We examined whether seasonally variable SNPs are enriched on any particular chromosome. To

perform this test, we calculated the enrichment of seasonal SNPs per chromosome in the observed

data relative to the permutations (Figure 2D). Chromosomes 2L and 3R have the greatest nominal

enrichment of seasonal SNPs relative to permutations, with a mean increase of 30% ($p = 0.06$) and of

18% ($p = 0.08$), respectively. Enrichment is much lower for chromosomes 2R and 3L (-2% and 4%, p

$= 0.54$ and $p = 0.20$, respectively). The variation of the seasonal signal across chromosomes is

statistically significant (ANOVA, $p < 10^{-15}$).

As chromosomal inversions have been observed to fluctuate in frequency over seasonal time

(Dobzhansky, 1948, 1943; Kapun et al., 2016), we extended the analysis to focus on large

cosmopolitan inversions. We find that regions near inversion breakpoints (± 0.5Mb) on chromosomes

2L ($p = 0.03$, inversion *In(2L)t*) and 3L ($p = 0.01$, inversion *In(3L)Payne*) are enriched for seasonal

SNPs. Regions within the three inversions on chromosome 3R (*In(3R)Mo*, *In(3R)K*, *In(3R)Payne*) are

also enriched for seasonal SNPs ($p = 0.05$). We do not find evidence for enrichment of seasonal SNPs

associated with *In(2R)NS* as previously reported by Kapun et al. (2016). These data confirm that

inversions may play a role in the adaptive response to seasonally variable environments, similar to

that of structural variants or large haplotype blocks in other systems (Hager et al., 2021; Jones et al.,

2012; Samuk et al., 2017; Tuttle et al., 2016).

Latitudinal differentiation parallels signatures of seasonal adaptation

Parallel changes in phenotype and genotype across time and space have been observed in *D.*

melanogaster (Bergland et al., 2014; Cogni et al., 2015, 2014; Kapun et al., 2016; Kapun and Flatt,

2019; Paaby et al., 2014; Rajpurohit et al., 2018, 2017). To assess whether we observe similar patterns

in our data, we tested whether SNPs that change in allele frequency along a latitudinal cline and

across seasons do so in a parallel fashion. Parallel changes in allele frequency occur when the sign of

the allele frequency change between spring and fall is the same as between high- and low-latitude

populations. Using four populations along the East Coast latitudinal cline, we identified clinally

varying polymorphisms using single-locus models (GLM, with allele frequency regressed against

latitude) as well as a model that accounts for population structure (Bayenv; Günther and Coop, 2013).

We re-identified seasonally varying SNPs using a GLM (as done previously) using 18 of the Core20

populations to avoid two samples drawn from a shared collection locale with the clinal set (PA_li;

Supplemental Table 1).

We find significant seasonal and latitudinal parallelism, with the magnitude of parallelism increasing

with increasing GLM significance thresholds (Figure 2E). Among the polymorphisms that are in the

top 1% of both seasonal and clinal GLMs (n = 115), 74% have parallel allele frequency change

(greater than 100/100 matched controls; naive expectation: 50%). This pattern of parallelism is also

observed when using a Bayenv model of clinality (Supplemental Figure 5). Parallel changes in allele

frequency between seasons and clines that we observe here is similar to previously published genome-

wide (Bergland et al., 2014) and locus specific results (Cogni et al., 2014; Kapun et al., 2016; Stalker,

1980, 1976, p. 197).

To evaluate whether parallel allele frequency change across time and space are likely to be

independent of shared demographic processes such as local seasonal migration, we compared East

Coast clinal variation to seasonal variation identified in two groups of populations that are

geographically and genetically distant from the East Coast (Campo et al., 2013; and see Figure 1B)

and that are unlikely to be connected via common seasonal migration. We assessed seasonal changes

in allele frequency separately for Californian (n = 3) and European (n = 3) populations and tested for

parallelism with polymorphisms that vary along the East Coast of North America. We found a clear

signal of parallelism between seasonal and latitudinal variation in both the California and the Europe

comparisons: for the top 1% of seasonal and clinal polymorphism, 70% and 63% have parallel allele

frequency change, for California and Europe, respectively (greater than 100/100 and 99/100 matched

controls; Figure 2E).

As SNPs in or near a number of major inversions are enriched for seasonal variation (Figure 2D), we

tested whether there is increased seasonal and latitudinal parallelism in these regions. For the top 5%

of clinal and seasonal SNPs (n = 2132), parallel allele frequency change is highest near inversion

breakpoints on chromosomes 2L and 3R (92%, CI_{95}: 87-96%, and 86%, CI_{95}: 78-93%, respectively)

and inside the 2L inversion (83%, CI_{95}: 80-87%; Figure 2F). While parallelism is greatest overall

around inversion breakpoints (concordance: 81%, CI_{95}: 77-84%) and inside inversions (concordance:

70%, CI$_{95}$: 67-73%), we also observe a statistically significant excess of latitudinal and seasonal

parallelism outside of inversions (concordance: 57%, CI$_{95}$: 53-62%). The elevated signal of

parallelism between spatial and seasonal allele frequency fluctuations across the whole genome, and

in particular outside of inverted regions, suggests that inversions themselves, while important, are not

the sole drivers of allele frequency change between seasons.

Predictability of seasonal adaptation based on local environmental conditions

The signature of seasonal adaptation that we observe indicates that there are consistent changes in

allele frequencies between seasons, broadly defined, among populations sampled across multiple

years, and localities separated by thousands of miles (Figure 1A,B). Next, we asked if seasonal

fluctuations in allele frequencies are predictable within any particular population or year, and if the

extent of predictability is related to aspects of local climate or other factors.

To address this question, we performed a leave-one-out analysis. In this analysis, we sequentially

removed each of the 20 paired spring-fall populations within the Core20 set and re-identified

seasonally variable SNPs among the remaining 19 populations as well as for the dropped, 20th

population. To evaluate predictability, we calculated the proportion of SNPs that have the same

direction of allele frequency change between spring and fall in both the discovery and test models

(i.e., concordant allele frequency change). We estimated concordance across the range of joint

significance thresholds of both models. For each population, we then calculated a 'predictability

score'. This score reflects the change in concordance rate as a function of the joint significance

threshold (see Materials and Methods for details). Note that populations with a positive predictability

score are those in which the concordance rate is greater than 50% over the bulk of the genome or

chromosome.

We find that the majority of populations show a positive predictability score (Figure 3A). For these

populations, the concordance rate reaches 60-70% among SNPs passing a stringent joint significance

threshold (top 0.1% in both models). At more modest joint significance thresholds (e.g., the top 10%

in both sets), the concordance rate is lower but still statistically different from 50% (maximum

nominal *p*-value at the top 5% of sites is 0.008). At the same time, intriguingly, we found that four of

the 20 populations had negative predictability scores (Benton Harbor, Michigan 2014; Lancaster,

Massachusetts 2012; Topeka, Kansas 2014; and Esparto, California 2012). Strong allele frequency

changes in these four populations occur, but the direction of allele frequency change from spring to

fall at many SNPs is opposite that of the remaining populations.

Figure 3. **The thermal limit model.** The thermal limit model argues that weather in the weeks prior to sampling can explain variation in the predictability of allele frequency change. (A) Each line represents the comparison between one population and the remaining 19. The x-axis represents the upper threshold of quantile-ranked *p*-values from the single-population test (Fisher's Exact Test) and the 19-population test (GLM), e.g. the top 1% in both tests. The y-axis is the fraction of SNPs where the sign of allele frequency change in the single population test matches the average sign change among the remaining 19. The color scheme represents the slope of this line and is used as a summary statistic for each population. (B/C). We regressed the summary score from (A) onto a number of characterizations of average temperature (B, first four rows), geography (B, 5th row), technical (B, rows 6-8), and thermal extremes (C), considering weather 14. Diamonds represent the observed R^2 for (B) and observed maximum R^2 across all thermal limits for (C). Violin plots represent the expected distribution of R^2 based on permutations. The red diamond represents the model with nominal *p*-value < 0.01. The empirical *p*-values for these models are listed next to the corresponding red diamond. The 14 day model that uses the counts of hot spring days and cold fall days passes a conservative application of the Bonferroni (see Results & Materials and Methods) at alpha = 0.13. (D) The

distribution of spring maximum (S- Max) daily temperature and fall minimum (F-min) daily temperature in the 2 weeks prior to sampling. Discordant (blue) populations do not cluster in time or space. Populations shown here are those in which we have weather data (15 populations, in total). (E) The stand-out model uses the number of hot spring days and cold fall days. To determine the optimal threshold for what defines hot and cold, we systematically varied the upper and lower thermal limits from 0°C to 40°C and used the count of hot spring ($x1$) and cold fall ($x2$) days as independent, additive variables in a regression model; the genome-wide predictability score was used as the dependent variable (y). The best fit model uses a spring max (S-Max) temp. of ~32°C and a fall min (F-Min) temp of ~5°C and explains ~82% of the variation in the population predictability scores.

We then asked why predictability scores vary among populations. To address this question we built

models that related the genome-wide predictability score (Fig 3A) for each population with a number

of different potential explanatory variables (Fig 3B,C,D). We first considered a number of nuisance

variables such as collection method and substrate, as well as contamination rates by *D. simulans*.

None of these variables are highly correlated with the genome-wide predictability score more than

expected by chance (Fig 3B). Latitude and longitude are also not strong predictors of variation in

genome-wide predictability score (Fig 3B).

We next focused on an environmental model based on aspects of the thermal environment. The

motivation for these characterizations of temperature is based on the hypothesis that populations with

negative genome-wide predictability scores were exposed to particularly warm springs and/or

particularly cool falls prior to our sampling. This hypothesis is plausible considering that short-term

changes in temperature have been linked, either directly or indirectly, to dramatic and partly

predictable changes in allele frequencies in the wild and in the laboratory in multiple Drosophilid

species (Barghi et al., 2019; Bergland et al., 2014; Mallard et al., 2018; Rodríguez-Trelles et al., 2013;

Tobler et al., 2014). We evaluated a number of features of temperature by calculating daily averages,

average minimum and maximums, as well as the cumulative degree days (Figure 3B). Finally, we

calculated the number of days below and above thermal limits, with the thermal limits defined across

a range of possible values (Fig 3C).

We find that the only set of features of the population that is significantly correlated with genome-

wide predictability score is the number of hot spring days and number of cold fall days prior to

sampling. The best fit thermal limit model explains 82.2% of the variation in genome-wide

predictability scores (Figure 3C, 3E), beats >99.5% of permutations, and survives conservative

Bonferroni multiple-test correction at alpha=0.13 (Figure 3C). Specifically, the thermal limits

identified by this best fit model are the number of days in the two weeks prior to spring sampling with

maximum temperature above ~32°C and the number of days in the fall with minimum temperature

below ~5°C. These values roughly correspond to upper and lower thermal limits for flies (Hoffmann,

2010; Kellermann et al., 2012; Overgaard and MacMillan, 2017). Taken together, these results

suggest that exposure to temperature extremes, or to associated environmental variables, acts as a

selective pressure in the wild, causing large fluctuations in allele frequency throughout the genome.

Discussion

In this study we have focused on the identification of genomic signatures of seasonal adaptation in *D.*

melanogaster. Our approach relies on the detection of parallel changes in allele frequency from spring

to fall across 20 populations. We use coarse definitions of spring and fall, with spring as the time of

year close to when *D. melanogaster* first becomes locally abundant and fall as the time close to the

end of the growing season but before populations become too scarce to sample (Behrman et al., 2015;

Ives, 1970). We sampled populations across a wide geographic range, such that the length of the

growing season as well as other environmental conditions varied substantially. Despite this

environmental heterogeneity and the coarseness of the definitions of spring and fall, we detect clear

signals of parallel seasonal allele frequency shifts across these populations (Figure 2A,B).

Specifically, we demonstrate that seasonal adaptation is a general phenomenon of temperate *D.*

melanogaster populations, and not restricted to a single orchard (Linvilla Orchard, Media,

Pennsylvania: Behrman et al., 2015; Bergland et al., 2014; Cogni et al., 2014; Schmidt and Conde,

2006), biogeographic region (East Coast of North America: Behrman et al., 2018), or even a

continent.

By detecting consistent seasonal changes in allele frequencies among multiple populations in North America and Europe we can argue that seasonal adaptation across this whole range is at least partly driven by a consistent set of loci. Note that at the moment we lack power to detect alleles that cycle in a small subset of populations or are private to a single population. We therefore focus only on the polymorphisms that segregate in all populations - and which consequently have a higher population frequency - and only look at the signal of parallel cycling. Thus, our conclusions are limited to the shared genetic basis of seasonal adaptation as revealed by the cycling of ~750,000 common SNPs which represent roughly half of all detected SNPs.

The detected seasonal SNPs can either be causal or, vastly more likely, linked to nearby causal genetic variants, be they SNPs, indels, structural variants, or transposable element insertions. While we cannot determine the precise number and identity of causal sites in the present study, the totality of the evidence is consistent with the polygenic nature of adaptation. Specifically, while we detected stronger seasonal signal near or inside several inversions on chromosomes 2L, 3L, and 3R suggesting that inversions are important drivers of seasonal adaptation (Dobzhansky, 1948, 1943; Kapun et al., 2016; Kapun and Flatt, 2019; Rodríguez-Trelles et al., 2013), we found that by focusing on the subset of seasonal sites that also show clinal patterns of geographic differentiation, we could detect seasonal signal outside inversions as well. Note that cosmopolitan inversions span a substantial portion of the genome and there is also likely to be substantial genetic diversity within these cytologically defined elements. Our study suggests that the strongly polygenic nature of seasonal adaptation will require a much broader seasonal sampling of populations to fine-map the causal variants and that such an endeavor is not futile, given that it is possible to detect a set of loci that show consistent seasonal fluctuations. At the same time, obtaining genomic data capable of identifying structural variation, including inversions, that plausibly underlie much of seasonal adaptation in *Drosophila* will clearly be important.

One key and poorly understood aspect of *Drosophila* seasonal adaptation is how the populations survive annual population collapse during the winter, manage to overwinter, and then manage to

expand and recolonize the available resources during the summer (Ives, 1970; Shpak et al., 2010).

Also poorly understood is the extent to which these patterns of collapse-overwintering-repopulation

involve migration and refugia (Coyne and Milstead, 1987; Ives, 1970). However, it is clear that

seasonal adaptation in *D. melanogaster* cannot involve wholesale population replacements by

migration because the spatial structure of the *D. melanogaster* populations is robust year after year as

evidenced by strong spatial population structure (Figure 1C) and persistent clinal patterns (Machado

et al., 2016; van Heerwaarden and Hoffmann, 2007). While one can conceive of migration from local

refugia playing a role, any such scenario still requires natural selection to operate in order to reset

populations back to the "spring" state every year. The full metapopulation dynamics and the ways that

local *D. melanogaster* survive seasonal crashes and expansions remains to be understood.

Our ability to detect a shared genetic basis of seasonal adaptation across these broadly distributed

populations implies that there are generic features in the environment that exert consistent selective

pressures despite the environmental heterogeneity across sampling times and locations. It is possible

that such generic selective pressures relate to temperature, humidity, population density, resource

availability, or other biotic and abiotic factors (Bogaerts-Márquez et al., 2020). The seasonal and

latitudinal parallelism in allele frequency change that we observe (Figure 2E) suggests that the

environmental factors that drive these patterns must both shift between seasons locally and vary

spatially from south to north.

While the exact nature of the environmental variables that drive seasonal selection remain to be

determined, here we provide evidence that temperature extremes in the weeks prior to sampling are a

strong proximate or even ultimate selective agent (Figure 3). Specifically, we show that the seasonal

behavior of any one population could be statistically associated with temperature extremes in the 2

weeks prior to sampling both in the spring and in the fall. This model accounts for why several

populations showed reversal of direction of movement of seasonal SNPs relative to the bulk of the

populations; these populations had unusually warm springs and unusually cool falls prior to sampling.

We thus hypothesize that, at the time of sampling, some of our spring samples had already

experienced sufficiently strong selection in the generically summer way and that the fall samples had

experienced selective pressures in the generically winter way. This would result in seasonal alleles in

the spring already showing fall-like patterns and the ones in the fall showing spring-like patterns.

These findings also raise an intriguing possibility that natural selection may not push populations in

the same consistent direction across the entire growing season but rather that natural selection and

adaptation may be more heterogeneous through time (Grant and Grant, 2002; Nosil et al., 2020;

Siepielski et al., 2009). In the future, more dense temporal sampling might help map the action of

natural selection to more specific environmental factors and to quantify how fast the selective

pressures change in strength and even direction across a single growing season. Our results thus

illustrate that evolution and ecology do operate on similar timescales in this system (Hendry, 2020;

Schoener, 2011; Thompson, 1998; Yoshida et al., 2003) and that evolutionary adaptation cannot be

generally ignored *a priori* in demographic or ecological investigations even on extremely short

timescales (Rudman et al., 2018)

While the number of detected seasonal loci is likely to be a vast overestimate of the number of causal

seasonal variants, the strength of selection implied by the magnitude of fluctuations at seasonal sites

while already high is likely to be an underestimate. Partially linked loci should show a smaller

magnitude of fluctuation than the true causal sites. Thus the 4-8% seasonal frequency shifts at the top

1% of the seasonal sites and the corresponding effective selection coefficients of ~10-30% per season

(~1-3% per generation) imply even larger magnitudes of selection at the causal sites. Furthermore,

given that we can only detect patterns of seasonal cycling that are at least partially consistent across

all or most sampled populations we are likely to underestimate the impact of linked seasonal selection

due to selection that is occurring in a population-specific manner. Seasonal adaptation therefore likely

constitutes a profound evolutionary force acting on a large number of dispersed loci and affecting

linked variation genome-wide. This seasonal selection force could rival in strength genetic draft due

to selective sweeps and should be much stronger over seasonal timescales than random genetic drift in

populations as large as *D. melanogaster* (Karasov et al., 2010).

Our observation that seasonal adaptation is polygenic and rapid suggests that there exists a substantial

number of common polymorphisms that are subject to strong fluctuating selection that are nonetheless

maintained across the range of *D. melanogaster*. The most straightforward explanation of these results

is that these polymorphisms are subject to some form of balancing selection (Bertram and Masel,

2019; Charlesworth, 2015; Gulisija and Kim, 2015, Levene, 1953; Wittmann et al., 2017). Pervasive

balancing selection is consistent with the recent realization that simple models of mutation-selection

balance are inconsistent with the extent of quantitative genetic variation in a variety of fitness related

traits (Charlesworth, 2015). A full understanding of the genomic consequences of balancing selection

caused by strong fluctuating selection will require the identification of individual causal loci. Doing

so will require substantial additional sampling across seasons as well as from closely and broadly

distributed locales throughout the whole geographic range of *D. melanogaster* in combination with

the genetic mapping of seasonally fluctuating phenotypes (Behrman et al., 2018, 2015; Erickson et al.,

2020; Rajpurohit et al., 2018, 2017; Schmidt and Conde, 2006). Finally, such efforts can be strongly

augmented by manipulative field experiments and functional work to investigate the molecular effects

of putatively causal genetic variants. All of this work will need to be done in the context of other

processes that underlie persistence of populations in the face of environmental heterogeneity such as

phenotypic plasticity and bet hedging (Ayrinhac et al., 2004; Bergland et al., 2008; Kain et al., 2015;

MacMillan et al., 2015; Overgaard et al., 2011; Via and Lande, 1985). Overall, it is becoming clear

that all of the key ecological and evolutionary processes, including demographic changes, mutation,

migration, phenotypic plasticity, and rapid evolution by natural selection, can and do occur on very

short and overlapping timescales and thus must be investigated as part of one complex evolving and

interacting system.

452   *Materials and methods*

Population sampling and sequence data

We assembled 73 samples of *D. melanogaster*, 61 representing newly collected and sequenced

samples and 12 representing previously published samples (Bergland et al., 2014; Kapun et al., 2016).

Locations, collection dates, number of individuals sampled, and depth of sequencing for all samples

are listed in Supplemental Table 1. For each sample, members of the Drosophila Real-Time Evolution

Consortium (DrosRTEC) collected an average of 75 male flies using direct aspiration from substrate,

netting, or trapping at orchards and residential areas. Flies were confirmed to be *D. melanogaster* by

examination of the male genital arch. We extracted DNA by first pooling all individuals from a

sample, grinding the tissue together in extraction buffer, and using a lithium chloride – potassium

acetate extraction protocol (see Bergland et al. 2014 for details on buffers and solutions). We prepared

sequencing libraries using a modified Illumina protocol (Bergland et al., 2014) and Illumina TrueSeq

adapters. Paired-end 125bp libraries were sequenced to an average of 94x coverage either at the

Stanford Sequencing Service Center on an Illumina HiSeq 2000, or at the Stanford Functional

Genomics facility on an Illumina HiSeq 4000.

The following sequence data processing was performed on both the new and the previously published

data. We trimmed low-quality 3' and 5' read ends (sequence quality < 20) using the program cutadapt

v1.8.1 (Martin, 2011). We mapped the raw reads to the *D. melanogaster* genome v5.5 (and for *D.*

simulans genome v2.01, flybase.org) using bwa v0.7.12 mem algorithms, with default parameters (Li

& Durbin, 2009), and used the program SAMtools v1.2 for bam file manipulation (functions index,

sort, and mpileup) (Li et al., 2009). We used the program picard v2.0.1 to remove PCR duplicates

(http://picard.sourceforge.net) and the program GATK v3.2-2 for indel realignment (McKenna et al.,

2010). We called SNPs and indels using the program VarScan v2.3.8 using a *p*-value of 0.05,

minimum variant frequency of 0.005, minimum average quality of 20, and minimum coverage of 10

(Koboldt et al., 2012). We filtered out SNPs within 10bp of an indel (they are more likely to be

spurious), variants in repetitive regions (identified by RepeatMasker and downloaded from the UCSC

Genome browser), SNPs with a median frequency of less than 1% across populations, regions with

low recombination rates (~0 cM/Mb; Comeron et al., 2012), and nucleotides with more than two

alleles. Because we sequenced only male individuals, the X chromosome had lower coverage and was

not used in our analysis. After filtering, we had a total of 1,763,522 autosomal SNPs. This set was

further filtered to include only SNPs found polymorphic in all samples ("common polymorphisms"),

resulting in 774,651 SNPs that represent the core set used in our analyses.

Due to the phenotypic similarity of the species *D. melanogaster* and *D. simulans*, we tested for *D.*

simulans contamination by competitively mapping reads to both genomes. We omitted any of our

pooled samples with greater than 5% of reads mapping preferentially to *D. simulans* (Charlottesville,

VA 2012, fall; Media, PA 2013 fall). For the remaining samples, reads that mapped better to the *D.*

simulans were removed from the analysis. For the populations used in this analysis, the average

proportion of reads mapping preferentially to *D. simulans* was less than 1% (see Supplemental Table

1), and there was no significant difference in the level of *D. simulans* contamination between spring

and fall samples (*t*-test $p = 0.90$).

In silico simulation analysis shows that competitive mapping accurately estimates the fraction of *D.*

simulans contamination. To conduct these simulations, we used a single *D. simulans* genome from an

inbred line derived from an individual caught California (Signor et al., 2017) and a single *D.*

melanogaster genome from a DGRP line (Mackay et al., 2012). We used wgsim to generate simulated

short-reads from each genome and mixed these short reads together in various proportions. We then

mapped the short reads back to the composite genome for competitive mapping, as described above.

We calculated contamination rate as the number of total reads mapping to the *D. simulans* reference

genome divided by the number of reads mapping to both *D. simulans* and *D. melanogaster*. We also

calculated the locus specific residual cross mapping rate by calculating the number of *D. simulans*

reads that still map to the *D. melanogaster* genome even after competitive mapping. These

simulations demonstrate that the estimation of the cross species mapping is precise (Pearson's $r =$

0.9999, $p = 1.6 \times 10^{-24}$, Supplemental Figure 7A), but underestimates the true contamination rate by

~2%. The level of residual *D. simulans* contamination does not correlate with contamination rate and

is roughly 9% (Supplemental Figure 7B).

To assess seasonal variation we analyzed population genomic sequence data from 20 spring and 20

fall samples ('Core20'). These samples represent a subset of the sequenced samples. We used samples

that had both a spring and a fall collection taken from the same locality in the same year. We also

used a maximum of two years of samples for a given locality to prevent the analysis from being

dominated by characteristics of a single population. When there were more than two years of samples

for a given population, we chose to use the two years with the earliest spring collection time. This

decision was made on the assumption that the earlier spring collection would better represent allele

frequencies following overwintering selection. This left 20 paired spring/fall samples, taken from 12

North American localities spread across 6 years and 3 European localities across 2 years

(Supplemental Table 1). The localities and years of sampling are as follows: Esparto, California 2012

and 2013; Tuolumne, California 2013; Lancaster, Massachusetts 2012 and 2014; Media, Pennsylvania

2010 and 2011; Ithaca, New York 2012; Cross Plains, Wisconsin 2012 and 2014; Athens, Georgia

2014; Charlottesville, Virginia 2014 and 2015, State College, Pennsylvania 2014; Topeka, Kansas

2014; Sudbury, Ontario, Canada 2015; Benton Harbor, Michigan 2014, Barcelona, Spain 2012;

Gross-Enzersdorf, Vienna 2012; Odesa, Ukraine 2013. For comparison of seasonal with latitudinal

variation, we used sequence data from four spring samples along the east coast of the United States

(Homestead, Florida 2010; Hahia, Georgia 2008; Eutawville, South Carolina 2010, Linvilla,

Pennsylvania 2009). We performed a principal components analysis of allele frequency per SNP per

sample using the prcomp function in R.

Identification of seasonal sites

We identified seasonal sites using three separate methods: a general linear regression model (GLM), a

Bayesian model (Bayenv; Coop et al., 2010; Günther & Coop, 2013) and a model based on the rank

Fisher's method (RFM). Statistical analyses were performed in R (R Core Team, 2014).

To perform the GLM we used the glm function with binomial error, weighted by the "effective

coverage" (N_c)- a measure of the number of chromosomes sampled, adjusted by the read depth:

537 $N_c = (1/N + 1/R)\text{-}1$

538 where N is the number of chromosomes in the pool and R is the read depth at that site (Kolaczkowski

539 et al., 2011; Feder et al., 2012; Bergland et al., 2014; Machado et al., 2016). This adjusts for the

540 additional error introduced by sampling of the pool at the time of sequencing. The seasonal GLM is a

541 regression of allele frequency by season (e.g., spring versus fall):

542            $y_i = \text{season} + \text{population} + e_i$

543 where y_i is the allele frequency at the ith SNP, population is the unordered categorical variable

544 denoting each of the 20 populations, and ei is the binomial error at the ith SNP. Although the GLM is

545 a powerful test, the corrected binomial sampling variance (N_c) is likely an anti-conservative estimate

546 of the true sampling variance associated with pooled sequencing (Machado et al., 2016; Spitzer et al.,

547 2019). Therefore, we use the results of this test as a seasonal outlier test, rather than an absolute

548 measure of the deviation from genome-wide null expectation of no allele frequency change between

549 seasons. Results are compared to 500 permutations of the spring and fall labels within each population

550 pair.

552 We tested for signals of seasonality using Bayenv. First, we calculated the genetic-covariance matrix

553 between the Core20 set. Next, we calculated Bayes Factor scores for each SNP using the observed

554 spring and fall labels. We used the read counts of reference and alternate based on the corrected

555 binomial sampling variance (N_c). We conducted 100 permutations of the spring and fall labels within

556 each population pair.

558 Finally, we developed a test to identify signals of seasonality which we call the "Rank Fisher's

559 Method" (RFM) that is robust to the inflated sample size estimates that can arise when using pooled

560 allele frequencies (Machado et al., 2016). For this method, we first perform a Fisher's Exact Test for

561 spring-fall allele frequency differentiation within each population pair. We then rank-normalize the

562 resulting *p*-values for any given population by grouping SNPs into categories based on the number of

563 total number of reads and the number of alternate reads, summed between spring and fall. The class-

564 based rank of each SNP becomes its new *p*-value, providing uniform sets of *p*-values across the

565 genome. As the power (minimum *p*-value) is relative to the number of SNPs being ranked per bin,

and as the bin sizes are equivalent across populations, each population has equivalent power.

Additionally, we find that this method is robust to inaccurate estimates of allele frequency precision

(Supplemental Figure 8). We then combine ranked p-values using Fisher's method for each SNP,

taking two times the negative sum of logged p-values across the 20 spring/fall comparisons (each tail

tested separately). The null distribution for this statistic (X^2) is a Chi-squared distribution with

degrees of freedom equal to 40 (two times the number of comparisons). We used the distribution of

these per-SNP X^2 test statistics to test for a seasonal signal above noise and compared to 200

spring/fall permutations.

Inversions

We analyzed seasonal variation associated with 6 major cosmopolitan inversions found in *D.*

melanogaster: *In(2L)t* (2L:2225744-13154180), *In(2R)NS* (2R:11278659-16163839), *In(3L)P*

(3L:3173046-16301941), *In(3R)K* (3R:7576289-21966092), *In(3R)Mo* (3R:17232639-24857019),

In(3R)P (3R:12257931-20569732). As the three inversions on 3R are overlapping, we combined

them. We classified the genome into three categories- inversion breakpoints (500Kb +/- each

inversion breakpoint; 104K SNPs), inside inversions (excluding breakpoint regions; 390K SNPs) and

outside inversions (excluding breakpoint regions; 281K SNPs).

Matched controls

With the assumption that the majority of the genome is not seasonal, we can use matched genomic

controls as a null distribution to test for enrichment of different features of seasonal polymorphisms.

The use of matched control SNP sets was employed for the tests of seasonal and latitudinal

parallelism. We matched each SNP identified as significantly seasonal (at a range p-values) to another

SNP, matched for chromosome, effective coverage, median spring allele frequency, inversion status

either within or outside of the major inversions *In(2L)t*, *In(2R)NS*, *In(3L)P*, *In(3R)K*, *In(3R)Mo*, and

In(3R)P, and recombination rate. We used the same *D. melanogaster* inversion breakpoints used in

Corbett-Detig & Hartl (2012) and the recombination rates from Comeron et al. (2012). We randomly

sampled 100 of the possible matches per SNP (excluding the focal SNP) to produce 100 matched

control sets. Any SNPs with fewer than 10 possible matches were discarded from the matched control

analyses. We defined 95% confidence intervals from the matched controls as the 3rd and 98th ranked

values for the quantity being tested (e.g., percent concordance or proportion of genic classes).

Latitudinal cline concordance

To identify SNPs that changed consistently in allele frequency with latitude (clinal), we first

identified SNPs that varied in allele frequency along a 14.4° latitudinal transect up the east coast of

the United States. We used one spring sample from each of the following populations (identified as

"Region_City_Year"): PA_li_2011 (39.9°N), SC_eu_2010 (33.4°N), GA_ha_2008 (30.1°N), and

FL_ho_2010 (25.5°N).

First, we regressed allele frequency with population latitude in a general linear model (glm: R v3.1.0),

using a binomial error model and weights proportional to the effective coverage (N_c):

$$y_i = \text{latitude} + e_i$$

where y_i is the allele frequency at the ith SNP, and e_i is the binomial error given the ith at the SNP. To

confirm the robustness of the clinal GLM results, clinality also was assessed using a second, more

complex model that accounts for existing covariance in the dataset (Bayenv: Coop et al., 2010;

Günther & Coop, 2013).

We then tested the concordance in allele frequency change by season with the allele frequency change

by latitude. We performed three separate seasonal regressions (see above) for comparison with the

latitudinal regression: spring verses fall for the 18 non-Pennsylvania paired samples, spring versus fall

for the three California paired samples, and spring versus fall for the three Europe paired samples.

With the removal of the Pennsylvania samples, none of these three seasonal regressions contained

samples from any of the four populations used for the latitudinal regression. Taking sets of

increasingly clinal and increasingly seasonal SNPs, we assessed the proportion of sites that both

increase in frequency from spring to fall and increase in frequency from north to south or that

decrease in frequency from spring to fall and decrease in frequency from north to south. We

compared this concordance with the concordance of 100 sets of matched controls.

Predictability analysis

To test the general predictability of seasonal change in our dataset, we performed a leave-one-out

analysis. In this analysis, we performed seasonal regressions for subsets of the data, dropping one

paired sample and comparing it to a seasonal test of the remaining 19. For the single population

model, we used a Fisher's exact test. For the multi-population model, we used a generalized linear

model. For either model, we used the effective number of reads (N_c, defined above). We then

measured the percent concordance of spring/fall allele frequency change, defined as the proportion of

SNPs that agree in the direction of allele frequency change between spring and fall. This was

performed 20 times, once for each paired sample.

To estimate the predictability scores, we calculated the rate of change of the concordance score as a

function of quantile threshold. To estimate this rate, we used the generalized linear model with

binomial error,

$$\text{concordance}_i = \text{quantile}_i + e_i,$$

where 'concordance' is the fraction of SNPs falling below the ith quantile threshold of both the target

(19 population) and test (1 population) models that changed in frequency in a concordant fashion and

e is the binomial error with weights corresponding to the total number of SNPs falling below that

threshold. Thus, the genome-wide (or chromosome specific) predictability score is heavily influenced

by concordance rates of higher quantiles because that is where the bulk of SNPs reside.

We tested whether the variation in genome-wide predictability scores among populations is correlated

with different features of the samples. We evaluated different nuisance parameters, such as collection

method, substrate, and *D. simulans* contamination rate, as well as a number of features related to

temperature. Daily minimum and maximum temperatures were obtained from the Global Historical

Climatology Network-Daily Database (Menne et al., 2012). We matched each locality to a weather

station based on geographic distance. For the Core20 set of populations, four of the collections did not

have precise collection dates for one or both seasonal collections (Athens, Georgia 2014; Media,

Pennsylvania 2010, 2011; Barcelona, Spain 2012), or were not associated with any climate data from

the GHCND database (Odessa, Ukraine 2013). These five populations were omitted from the analysis,

leaving 15 populations for the remainder of the predictability analysis (Figure 3D). Using daily

temperature data, we calculated the cumulative growing degree days following a model developed for

D. suzukii by Wiman et al (2014).

We regressed the predictability score onto each potential explanatory variable, recorded the

proportion of variation explained (R^2), and contrasted the observed R^2 to null distributions generated

via permutation (n = 25,000). For most variables, we performed a univariate model (e.g., latitude or

longitude). In some cases, we used a bivariate model (e.g., latitude + longitude). To test the thermal

limit model, we regressed the observed predictability score on the number of days above and below

the thermal limits. We varied thermal limits from -5°C to 40°C with a step size of 0.1°C.

Data availability

All raw sequence data have been deposited to the NCBI short read archive (SRA; BioProject

Accession #PRJNA308584; accession numbers for each sample can be found in Supplemental Table

1). Code to conduct these analyses, primary results files, and code to reproduce the figures are

available at https://github.com/machadoheather/dmel_seasonal_RTEC. VCF files with the raw allele

frequencies per population and a R-data file of allele frequencies and effective sample sizes (N_c;

compatible with scripts) are available on DataDryad

(https://datadryad.org/stash/dataset/doi:10.5061/dryad.4r7b826).

Acknowledgements

We thank the National Evolutionary Synthesis Center (NESCent) for sponsoring the 2012 Catalysis

meeting that initiated the *Drosophila* Real Time Evolution Consortium. The meeting was attended by

Alan Bergland, Alisa Sedghifar, Brian Helmuth, Brian Lazzarro, Chau-Ti Ting, David Kidd, Dmitri Petrov, Fabian Staubach, Hannah Burrack, Jim Fry, John Lessard, John Coulbourne, John Pool, Josefa Gonzalez, Julien Ayroles, Kelly Dyer, Kim Hughes, Maaria Kankare, Nadia Singh, Paul Schmidt, Regan Early, Stephen Porder, Subhash Rajpurohit, Sui Fai Lee, and Thomas Flatt, and we kindly thank all participants for their participation. We also thank Jamie Blundell and all members of the Schmidt and Petrov labs who provided exceptionally valuable feedback. This work was funded by the NIH grants R01GM100366 to PS and DAP, R35GM118165 to DAP, R01GM100366 to PS, F32GM097837 and R35GM119686 to AOB, European Commission grant H2020-ERC-2014-CoG-647900 to JG, a NSERC Discovery Grant RGPIN-2018-05551 and Canada Research Chair 950-230113 to TJSM.

Author contributions

Heather E. Machado: Designed the study, collected flies, performed analysis, wrote manuscript; Alan O. Bergland: Designed the study, collected flies, performed analysis, wrote manuscript; Ryan Taylor: Performed analysis; Susanne Tilk: Prepared sequencing libraries; Emily Behrman: Collected flies; Kelly Dyer: Collected flies; Daniel K. Fabian: Collected flies; Thomas Flatt: Collected flies; Josefa González: Collected flies; Talia L. Karasov: Collected flies; Iryna Kozeretska: Collected flies; Brian P. Lazzaro: Collected flies; Thomas JS Merritt: Collected flies; John E. Pool: Collected flies; Katherine O'Brien: Collected flies; Subhash Rajpurohit: Collected flies; Paula R. Roy: Collected flies; Stephen W. Schaeffer: Collected flies; Svitlana Serga: Collected flies; Paul Schmidt: Designed the study, collected flies, wrote manuscript; Dmitri Petrov: Designed the study, collected flies, wrote manuscript. All authors edited and approved the manuscript.

References

Ayrinhac A, Debat V, Gibert P, Kister A-G, Legout H, Moreteau B, Vergilino R, David JR. 2004. Cold adaptation in geographical populations of *Drosophila melanogaster*: phenotypic plasticity is more important than genetic variability. *Funct Ecol* **18**:700–706. doi:10.1111/j.0269-8463.2004.00904.x

Barghi N, Tobler R, Nolte V, Jakšić AM, Mallard F, Otte KA, Dolezal M, Taus T, Kofler R, Schlötterer C. 2019. Genetic redundancy fuels polygenic adaptation in *Drosophila*. *PLoS Biol*

17:e3000128. doi:10.1371/journal.pbio.3000128

Behrman EL, Howick VM, Kapun M, Staubach F, Bergland AO, Petrov DA, Lazzaro BP, Schmidt
PS. 2018. Rapid seasonal evolution in innate immunity of wild *Drosophila melanogaster*.
Proc Biol Sci **285**. doi:10.1098/rspb.2017.2599

Behrman EL, Watson SS, O'Brien KR, Heschel MS, Schmidt PS. 2015. Seasonal variation in life
history traits in two *Drosophila* species. *J Evol Biol* **28**:1691–1704. doi:10.1111/jeb.12690

Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. 2014. Genomic evidence of rapid
and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLoS Genet*
10:e1004775. doi:10.1371/journal.pgen.1004775

Bergland AO, Genissel A, Nuzhdin SV, Tatar M. 2008. Quantitative trait loci affecting phenotypic
plasticity and the allometric relationship of ovariole number and thorax length in *Drosophila*
melanogaster. *Genetics* **180**:567–582. doi:10.1534/genetics.108.088906

Bertram J, Masel J. 2019. Different mechanisms drive the maintenance of polymorphism at loci
subject to strong versus weak fluctuating selection. *Evolution* **73**:883–896.
doi:10.1111/evo.13719

Bogaerts-Márquez M, Guirao-Rico S, Gautier M, González J. 2020. Temperature, rainfall and wind
variables underlie environmental adaptation in natural populations of *Drosophila*
melanogaster. *Mol Ecol*. doi:10.1111/mec.15783

Botero CA, Weissing FJ, Wright J, Rubenstein DR. 2015. Evolutionary tipping points in the capacity
to adapt to environmental change. *Proc Natl Acad Sci* **112**:184–189.
doi:10.1073/pnas.1408589111

Buffalo V, Coop G. 2020. Estimating the genome-wide contribution of selection to temporal allele
frequency change. *Proc Natl Acad Sci U S A* **117**:20672–20680.
doi:10.1073/pnas.1919039117

Bürger R, Gimelfarb A. 1999. Genetic variation maintained in multilocus models of additive
quantitative traits under stabilizing selection. *Genetics* **152**:807–820.

Campo D, Lehmann K, Fjeldsted C, Souaiaia T, Kao J, Nuzhdin SV. 2013. Whole-genome
sequencing of two North American *Drosophila melanogaster* populations reveals genetic
differentiation and positive selection. *Mol Ecol* **22**:5084–5097. doi:10.1111/mec.12468

Charlesworth B. 2015. Causes of natural variation in fitness: evidence from studies of *Drosophila*
populations. *Proc Natl Acad Sci U S A* **112**:1662–1669. doi:10.1073/pnas.1423275112

Cogni R, Kuczynski C, Koury S, Lavington E, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF.
2014. The intensity of selection acting on the couch potato gene--spatial-temporal variation in
a diapause cline. *Evol Int J Org Evol* **68**:538–548. doi:10.1111/evo.12291

Cogni R, Kuczynski K, Lavington E, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF.
2015. Variation in *Drosophila melanogaster* central metabolic genes appears driven by
natural selection both within and between populations. *Proc Biol Sci* **282**:20142688.
doi:10.1098/rspb.2014.2688

Comeron JM, Ratnappan R, Bailin S. 2012. The Many Landscapes of Recombination in *Drosophila*
melanogaster. *PLOS Genet* **8**:e1002905. doi:10.1371/journal.pgen.1002905

Coop G, Witonsky D, Rienzo AD, Pritchard JK. 2010. Using Environmental Correlations to Identify
Loci Underlying Local Adaptation. *Genetics* **185**:1411–1423.
doi:10.1534/genetics.110.114819

Corbett-Detig RB, Hartl DL. 2012. Population Genomics of Inversion Polymorphisms in *Drosophila*
melanogaster. *PLOS Genet* **8**:e1003056. doi:10.1371/journal.pgen.1003056

Coyne JA, Milstead B. 1987. Long-Distance Migration of *Drosophila*. 3. Dispersal of *D.
melanogaster* Alleles from a Maryland Orchard. *Am Nat* **130**:70–82. doi:10.1086/284698

Curtsinger JW, Service PM, Prout T. 1994. Antagonistic Pleiotropy, Reversal of Dominance, and
Genetic Polymorphism. *Am Nat* **144**:210–228. doi:10.1086/285671

Denlinger DL. 2002. Regulation of Diapause. *Annu Rev Entomol* **47**:93–122.
doi:10.1146/annurev.ento.47.091201.145137

Dobzhansky T. 1948. Genetics of natural populations; altitudinal and seasonal changes produced by
natural selection in certain populations of *Drosophila persimilis*. *Genetics* **33**:158–176.

Dobzhansky T. 1943. Genetics of Natural Populations IX. Temporal Changes in the Composition of
Populations of *Drosophila pseudoobscura*. *Genetics* **28**:162–186.

Ellner S. 1996. Environmental fluctuations and the maintenance of genetic diversity in age or stage-
structured populations. *Bull Math Biol* **58**:103–127. doi:10.1007/BF02458284
Ellner S, Hairston, NG. 1994. Role of Overlapping Generations in Maintaining Genetic Variation in a
Fluctuating Environment. *Am Nat* **143**:403–417. doi:10.1086/285610
Ellner S, Sasaki A. 1996. Patterns of Genetic Polymorphism Maintained by Fluctuating Selection with
Overlapping Generations. *Theor Popul Biol* **50**:31–65. doi:10.1006/tpbi.1996.0022
Erickson PA, Weller CA, Song DY, Bangerter AS, Schmidt P, Bergland AO. 2020. Unique genetic
signatures of local adaptation over space and time for diapause, an ecologically relevant
complex trait, in *Drosophila melanogaster*. *PLoS Genet* **16**:e1009110.
doi:10.1371/journal.pgen.1009110
Ewing EP. 1979. Genetic Variation in a Heterogeneous Environment VII. Temporal and Spatial
Heterogeneity in Infinite Populations. *Am Nat* **114**:197–212. doi:10.1086/283468
Fabian DK, Kapun M, Nolte V, Kofler R, Schmidt PS, Schlötterer C, Flatt T. 2012. Genome-wide
patterns of latitudinal differentiation among populations of *Drosophila melanogaster* from
North America. *Mol Ecol* **21**:4748–4769. doi:https://doi.org/10.1111/j.1365-
294X.2012.05731.x
Feder AF, Petrov DA, Bergland AO. 2012. LDx: Estimation of Linkage Disequilibrium from High-
Throughput Pooled Resequencing Data. *PLoS ONE* **7**:e48588.
doi:10.1371/journal.pone.0048588
Grant PR, Grant BR. 2002. Unpredictable evolution in a 30-year study of Darwin's finches. *Science*
296:707–711. doi:10.1126/science.1070315
Grenier JK, Arguello JR, Moreira MC, Gottipati S, Mohammed J, Hackett SR, Boughton R,
Greenberg AJ, Clark AG. 2015. Global diversity lines - a five-continent reference panel of
sequenced *Drosophila melanogaster* strains. *G3 Bethesda Md* **5**:593–603.
doi:10.1534/g3.114.015883
Günther T, Coop G. 2013a. Robust identification of local adaptation from allele frequencies. *Genetics*
195:205–220. doi:10.1534/genetics.113.152462
Günther T, Coop G. 2013b. Robust Identification of Local Adaptation from Allele Frequencies.
Genetics **195**:205–220. doi:10.1534/genetics.113.152462
Hager ER, Harringmeyer OS, Wooldridge TB, Theingi S, Gable JT, McFadden S, Neugeboren B,
Turner KM, Hoekstra HE. 2021. A chromosomal inversion drives evolution of multiple
adaptive traits in deer mice. *bioRxiv* 2021.01.21.427490. doi:10.1101/2021.01.21.427490
Hedrick PW. 1976. Genetic variation in a heterogeneous environment. II. Temporal heterogeneity and
directional selection. *Genetics* **84**:145–157.
Hendry AP. 2020. Eco-Evolutionary Dynamics. Princeton; Jackson: Princeton University Press Two
Rivers Distribution.
Hoffmann AA. 2010. Physiological climatic limits in *Drosophila*: patterns and implications. *J Exp*
Biol **213**:870–880. doi:10.1242/jeb.037630
Ives PT. 1970. Further genetic studies of the South Amherst population of *Drosophila melanogaster*.
Evol Int J Org Evol **24**:507–518. doi:10.1111/j.1558-5646.1970.tb01785.x
Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, Swofford R, Pirun M, Zody MC,
White S, Birney E, Searle S, Schmutz J, Grimwood J, Dickson MC, Myers RM, Miller CT,
Summers BR, Knecht AK, Brady SD, Zhang H, Pollen AA, Howes T, Amemiya C, Lander
ES, Palma FD, Lindblad-Toh K, Kingsley DM. 2012. The genomic basis of adaptive
evolution in threespine sticklebacks. *Nature* **484**:55–61. doi:10.1038/nature10944
Kain JS, Zhang S, Akhund-Zade J, Samuel ADT, Klein M, Bivort BL. 2015. Variability in thermal
and phototactic preferences in *Drosophila* may reflect an adaptive bet-hedging strategy.
Evolution **69**:3171–3185. doi:10.1111/evo.12813
Kao JY, Zubair A, Salomon MP, Nuzhdin SV, Campo D. 2015. Population genomic analysis
uncovers African and European admixture in *Drosophila melanogaster* populations from the
south-eastern United States and Caribbean Islands. *Mol Ecol* **24**:1499–1509.
doi:10.1111/mec.13137
Kapun M, Barrón MG, Staubach F, Obbard DJ, Wiberg RAW, Vieira J, Goubert C, Rota-Stabelli O,
Kankare M, Bogaerts-Márquez M, Haudry A, Waidele L, Kozeretska I, Pasyukova EG,
Loeschcke V, Pascual M, Vieira CP, Serga S, Montchamp-Moreau C, Abbott J, Gibert P,

Porcelli D, Posnien N, Sánchez-Gracia A, Grath S, Sucena É, Bergland AO, Guerreiro MPG,
Onder BS, Argyridou E, Guio L, Schou MF, Deplancke B, Vieira C, Ritchie MG, Zwaan BJ,
Tauber E, Orengo DJ, Puerma E, Aguadé M, Schmidt P, Parsch J, Betancourt AJ, Flatt T,
González J. 2020. Genomic Analysis of European *Drosophila melanogaster* Populations
Reveals Longitudinal Structure, Continent-Wide Selection, and Previously Unknown DNA
Viruses. *Mol Biol Evol* **37**:2661–2678. doi:10.1093/molbev/msaa120
Kapun M, Fabian DK, Goudet J, Flatt T. 2016. Genomic Evidence for Adaptive Inversion Clines in
Drosophila melanogaster. *Mol Biol Evol* **33**:1317–1336. doi:10.1093/molbev/msw016
Kapun M, Flatt T. 2019. The adaptive significance of chromosomal inversion polymorphisms in
Drosophila melanogaster. *Mol Ecol* **28**:1263–1282. doi:10.1111/mec.14871
Karasov T, Messer PW, Petrov DA. 2010. Evidence that adaptation in *Drosophila* is not limited by
mutation at single sites. *PLoS Genet* **6**:e1000924. doi:10.1371/journal.pgen.1000924
Kellermann V, Overgaard J, Hoffmann AA, Fløjgaard C, Svenning J-C, Loeschcke V. 2012. Upper
thermal limits of *Drosophila* are linked to species distributions and strongly constrained
phylogenetically. *Proc Natl Acad Sci U S A* **109**:16228–16233. doi:10.1073/pnas.1207553109
Kim YB, Oh JH, McIver LJ, Rashkovetsky E, Michalak K, Garner HR, Kang L, Nevo E, Korol AB,
Michalak P. 2014. Divergence of *Drosophila melanogaster* repeatomes in response to a sharp
microclimate contrast in Evolution Canyon, Israel. *Proc Natl Acad Sci U S A* **111**:10630–
10635. doi:10.1073/pnas.1410372111
Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L,
Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in
cancer by exome sequencing. *Genome Res* **22**:568–576. doi:10.1101/gr.129684.111
Kolaczkowski B, Kern AD, Holloway AK, Begun DJ. 2011. Genomic differentiation between
temperate and tropical Australian populations of *Drosophila melanogaster*. *Genetics*
187:245–260. doi:10.1534/genetics.110.123059
Korol AB, Kirzhner VM, Ronin YI, Nevo E. 1996. Cyclical environmental changes as a factor
maintaining genetic polymorphism. 2. Diploid selection for an additive trait. *Evol Int J Org*
Evol **50**:1432–1441. doi:10.1111/j.1558-5646.1996.tb03917.x
Kostál V. 2006. Eco-physiological phases of insect diapause. *J Insect Physiol* **52**:113–127.
doi:10.1016/j.jinsphys.2005.09.008
Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016. A Thousand Fly Genomes: An
Expanded *Drosophila* Genome Nexus. *Mol Biol Evol* **33**:3308–3313.
doi:10.1093/molbev/msw195
Lande R. 2008. Adaptive topography of fluctuating selection in a Mendelian population. *J Evol Biol*
21:1096–1105. doi:10.1111/j.1420-9101.2008.01533.x
Levins RA. 1974. Evolution in changing environments ; some theoretical explorations, 3. printing. ed,
Monographs in population biology. Princeton, NJ: Princeton Univ. Press.
Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinforma Oxf Engl **25**:1754–1760. doi:10.1093/bioinformatics/btp324
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000
Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinforma Oxf Engl* **25**:2078–2079. doi:10.1093/bioinformatics/btp352
Lynch M, Ho W-C. 2020. The Limits to Estimating Population-Genetic Parameters with Temporal
Data. *Genome Biol Evol* **12**:443–455. doi:10.1093/gbe/evaa056
Machado HE, Bergland AO, O'Brien KR, Behrman EL, Schmidt PS, Petrov DA. 2016. Comparative
population genomics of latitudinal variation in *Drosophila simulans* and *Drosophila*
melanogaster. *Mol Ecol* **25**:723–740. doi:https://doi.org/10.1111/mec.13446
Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire
MM, Cridland JM, Richardson MF, Anholt RRH, Barrón M, Bess C, Blankenburg KP,
Carbone MA, Castellano D, Chaboub L, Duncan L, Harris Z, Javaid M, Jayaseelan JC,
Jhangiani SN, Jordan KW, Lara F, Lawrence F, Lee SL, Librado P, Linheiro RS, Lyman RF,
Mackey AJ, Munidasa M, Muzny DM, Nazareth L, Newsham I, Perales L, Pu L-L, Qu C,
Ràmia M, Reid JG, Rollmann SM, Rozas J, Saada N, Turlapati L, Worley KC, Wu Y-Q,
Yamamoto A, Zhu Y, Bergman CM, Thornton KR, Mittelman D, Gibbs RA. 2012. The
Drosophila melanogaster Genetic Reference Panel. *Nature* **482**:173–178.

doi:10.1038/nature10811

MacMillan HA, Ferguson LV, Nicolai A, Donini A, Staples JF, Sinclair BJ. 2015. Parallel
ionoregulatory adjustments underlie phenotypic plasticity and evolution of *Drosophila* cold
tolerance. *J Exp Biol* **218**:423–432. doi:10.1242/jeb.115790

Mallard F, Nolte V, Tobler R, Kapun M, Schlötterer C. 2018. A simple genetic basis of adaptation to
a novel thermal environment results in complex metabolic rewiring in *Drosophila*. *Genome*
Biol **19**:119. doi:10.1186/s13059-018-1503-4

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**:10–12. doi:10.14806/ej.17.1.200

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**:1297–1303.
doi:10.1101/gr.107524.110

Menne MJ, Durre I, Vose RS, Gleason BE, Houston TG. 2012. An Overview of the Global Historical
Climatology Network-Daily Database. *J Atmospheric Ocean Technol* **29**:897–910.
doi:10.1175/JTECH-D-11-00103.1

Nosil P, Villoutreix R, de Carvalho CF, Feder JL, Parchman TL, Gompert Z. 2020. Ecology shapes
epistasis in a genotype-phenotype-fitness map for stick insect colour. *Nat Ecol Evol* **4**:1673–
1684. doi:10.1038/s41559-020-01305-y

Overgaard J, Kristensen TN, Mitchell KA, Hoffmann AA. 2011. Thermal tolerance in widespread and
tropical *Drosophila* species: does phenotypic plasticity increase with latitude? *Am Nat* **178**
Suppl 1:S80-96. doi:10.1086/661780

Overgaard J, MacMillan HA. 2017. The Integrative Physiology of Insect Chill Tolerance. *Annu Rev*
Physiol **79**:187–208. doi:10.1146/annurev-physiol-022516-034142

Paaby AB, Bergland AO, Behrman EL, Schmidt PS. 2014. A highly pleiotropic amino acid
polymorphism in the *Drosophila* insulin receptor contributes to life-history adaptation. *Evol*
Int J Org Evol **68**:3395–3409. doi:10.1111/evo.12546

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, Duchen P, Emerson
JJ, Saelao P, Begun DJ, Langley CH. 2012. Population Genomics of sub-saharan *Drosophila*
melanogaster: African diversity and non-African admixture. *PLoS Genet* **8**:e1003080.
doi:10.1371/journal.pgen.1003080

Rajpurohit S, Gefen E, Bergland AO, Petrov DA, Gibbs AG, Schmidt PS. 2018. Spatiotemporal
dynamics and genome-wide association analysis of desiccation tolerance in *Drosophila*
melanogaster. *Mol Ecol* **27**:3525–3540. doi:10.1111/mec.14814

Rajpurohit S, Hanus R, Vrkoslav V, Behrman EL, Bergland AO, Petrov D, Cvačka J, Schmidt PS.
2017. Adaptive dynamics of cuticular hydrocarbons in *Drosophila*. *J Evol Biol* **30**:66–80.
doi:10.1111/jeb.12988

Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD. 2014. Parallel geographic variation
in *Drosophila melanogaster*. *Genetics* **197**:361–373. doi:10.1534/genetics.114.161463

Rodríguez-Trelles F, Tarrío R, Santos M. 2013. Genome-wide evolutionary response to a heat wave
in *Drosophila*. *Biol Lett* **9**:20130228. doi:10.1098/rsbl.2013.0228

Rudman SM, Barbour MA, Csilléry K, Gienapp P, Guillaume F, Hairston NG, Hendry AP, Lasky JR,
Rafajlović M, Räsänen K, Schmidt PS, Seehausen O, Therkildsen NO, Turcotte MM, Levine
JM. 2018. What genomic data can reveal about eco-evolutionary dynamics. *Nat Ecol Evol*
2:9–15. doi:10.1038/s41559-017-0385-2

Rudman SM, Greenblum S, Hughes RC, Rajpurohit S, Kiratli O, Lowder DB, Lemmon SG, Petrov
DA, Chaston JM, Schmidt P. 2019. Microbiome composition shapes rapid genomic
adaptation of *Drosophila melanogaster*. *Proc Natl Acad Sci* **116**:20025–20032.
doi:10.1073/pnas.1907787116

Samuk K, Owens GL, Delmore KE, Miller SE, Rennison DJ, Schluter D. 2017. Gene flow and
selection interact to promote adaptive divergence in regions of low recombination. *Mol Ecol*
26:4378–4390. doi:https://doi.org/10.1111/mec.14226

Scheiner SM. 1993. Genetics and Evolution of Phenotypic Plasticity. *Annu Rev Ecol Syst* **24**:35–68.
doi:10.1146/annurev.es.24.110193.000343

Schlötterer C, Tobler R, Kofler R, Nolte V. 2014. Sequencing pools of individuals - mining genome-

wide polymorphism data without big funding. *Nat Rev Genet* **15**:749–763. doi:10.1038/nrg3803

Schmidt PS, Conde DR. 2006. Environmental heterogeneity and the maintenance of genetic variation for reproductive diapause in *Drosophila melanogaster*. *Evol Int J Org Evol* **60**:1602–1611.

Schoener TW. 2011. The Newest Synthesis: Understanding the Interplay of Evolutionary and Ecological Dynamics. *Science* **331**:426–429. doi:10.1126/science.1193954

Shpak M, Wakeley J, Garrigan D, Lewontin RC. 2010. A structured coalescent process for seasonally fluctuating populations. *Evol Int J Org Evol* **64**:1395–1409. doi:10.1111/j.1558-5646.2009.00891.x

Siepielski AM, DiBattista JD, Carlson SM. 2009. It's about time: the temporal dynamics of phenotypic selection in the wild. *Ecol Lett* **12**:1261–1276. doi:https://doi.org/10.1111/j.1461-0248.2009.01381.x

Signor S. 2017. Population genomics of Wolbachia and mtDNA in *Drosophila simulans* from California. *Sci Rep* **7**:13369. doi:10.1038/s41598-017-13901-3

Stalker HD. 1980. Chromosome Studies in Wild Populations of *Drosophila melanogaster*. II. Relationship of Inversion Frequencies to Latitude, Season, Wing-Loading and Flight Activity. *Genetics* **95**:211–223.

Stalker HD. 1976. Chromosome studies in wild populations of *D. melanogaster*. *Genetics* **82**:323–347.

Svetec N, Cridland JM, Zhao L, Begun DJ. 2016. The Adaptive Significance of Natural Genetic Variation in the DNA Damage Response of *Drosophila melanogaster*. *PLoS Genet* **12**:e1005869. doi:10.1371/journal.pgen.1005869

Tauber MJ, Tauber CA, Masaki S. 1986. Seasonal adaptations of insects. New York: Oxford University Press.

Thompson JN. 1998. Rapid evolution as an ecological process. *Trends Ecol Evol* **13**:329–332. doi:10.1016/S0169-5347(98)01378-0

Tobler R, Franssen SU, Kofler R, Orozco-Terwengel P, Nolte V, Hermisson J, Schlötterer C. 2014. Massive habitat-specific genomic response in *D. melanogaster* populations during experimental evolution in hot and cold environments. *Mol Biol Evol* **31**:364–375. doi:10.1093/molbev/mst205

Tuttle EM, Bergland AO, Korody ML, Brewer MS, Newhouse DJ, Minx P, Stager M, Betuel A, Cheviron ZA, Warren WC, Gonser RA, Balakrishnan CN. 2016. Divergence and Functional Degradation of a Sex Chromosome-like Supergene. *Curr Biol* **26**:344–350. doi:10.1016/j.cub.2015.11.069

van Heerwaarden B, Hoffmann AA. 2007. Global Warming: Fly Populations Are Responding Rapidly to Climate Change. *Curr Biol* **17**:R16–R18. doi:10.1016/j.cub.2006.11.035

Via S, Lande R. 1985. Genotype-Environment Interaction and the Evolution of Phenotypic Plasticity. *Evolution* **39**:505. doi:10.2307/2408649

Wiman NG, Walton VM, Dalton DT, Anfora G, Burrack HJ, Chiu JC, Daane KM, Grassi A, Miller B, Tochen S, Wang X, Ioriatti C. 2014. Integrating Temperature-Dependent Life Table Data into a Matrix Projection Model for *Drosophila* suzukii Population Estimation. *PLoS ONE* **9**:e106909. doi:10.1371/journal.pone.0106909

Wittmann MJ, Bergland AO, Feldman MW, Schmidt PS, Petrov DA. 2017. Seasonally fluctuating selection can maintain polymorphism at many loci via segregation lift. *Proc Natl Acad Sci* **114**:E9932–E9941. doi:10.1073/pnas.1702994114

Yoshida T, Jones LE, Ellner SP, Fussmann GF, Hairston NG. 2003. Rapid evolution drives ecological dynamics in a predator–prey system. *Nature* **424**:303–306. doi:10.1038/nature01767

Zhao L, Begun DJ. 2017. Genomics of parallel adaptation at two timescales in *Drosophila*. *PLoS Genet* **13**:e1007016. doi:10.1371/journal.pgen.1007016

Zhu Y, Bergland AO, González J, Petrov DA. 2012. Empirical validation of pooled whole genome population re-sequencing in *Drosophila melanogaster*. *PloS One* **7**:e41901. doi:10.1371/journal.pone.0041901

Supplemental Information

Supplemental Table 1: Population information with sampling meta-data

Supplemental Table 2: Top 1% of seasonal SNPs (regression beta and p-values) based on GLM model. Coordinates are in *D. melanogaster* reference v5.5.

Supplemental Figure 1. Extended sampling figure.

Supplemental Figure 2. Site frequency spectrum.

Supplemental Figure 3. Chromosome level enrichment tests.

Supplemental Figure 4. Bayenv seasonal analysis.

Supplemental Figure 5. Comparison of the GLM and the Bayenv model for assessing clinal and seasonal concordance.

Supplemental Figure 6. Inversion frequencies.

Supplemental Figure 7. Simulation of competitive mapping to evaluation *D. simulans* contamination.

Supplemental Figure 8. The effect of artificial increases of sample size on model output.

Supplemental Figure 1

Supplemental Figure 1. Sampling localities in A) North America and B) Europe for all samples collected, regardless of use in the present analysis. Sizes of circles reflect number of samples per locality.

Supplemental Figure 2

Supplemental Figure 2. Distributions of the mean spring (A/B) and mean fall (C/D) allele frequencies for the dataset before (A/C) and after (B/D) filtering SNP found to be monomorphic in any population.

983
984
985 **Supplemental Figure 3**

986
987

Supplemental Figure 3. Enrichment of seasonal SNPs (median of 500 permutations) by chromosome and significance threshold. Enrichment is calculated as the observed number of seasonal SNPs over the number of seasonal SNPs in each permutation. Error bars are 95% confidence intervals and asterisks denote greater seasonal enrichment than more than 95% of permutations.

990 **Supplemental Figure 4**

991
992
993

Supplemental Figure 4. Bayenv seasonal analysis. A) Proportion of SNPs found highly seasonal in both the GLM and the bayenv seasonal analyses. B) Seasonal and latitudinal concordance for both the GLM and the bayenv analyses.

995 **Supplemental Figure 5**

Supplemental Figure 5. Comparison of the GLM and the Bayenv model for assessing clinal and seasonal concordance. A) The "joint quantile threshold" is the quantile threshold for both the seasonal and the clinal analysis, and includes all SNPs at that quantile or less. B) The "joint quantile bin" includes only those SNPs at that specific quantile.

998   **Supplemental Figure 6**

999
1000

Supplemental Figure 6. Inversion frequencies. Each datapoint is a separate population. Numbers are the fraction of populations where spring is greater than fall.

1001

1002 **Supplemental Figure 7**
1003

1004
1005

Supplemental Figure 7. Competitive mapping effectively reduces cross species mapping between *D. simulans* and *D. melanogaster*. (A) The estimated rate of *D. simulans* contamination can be precisely estimated by calculating the number of reads mapping preferentially to the *D. simulans* genome. Estimated contamination rates are ~2% lower than the expected rate. (B) The residual cross species mapping rate, i.e. the fraction of *D. simulans* reads mapping to the *D. melanogaster* genome is roughly consistent at ~9%.

1006
1007

1008 **Supplemental Figure 8**

Supplemental Figure 8. The effect of artificial increases of sample size on the Core20 seasonal analysis using A,C) the RFM and B,D) the GLM analyses. Top row: sample size inflated by a fixed amount. Bottom row: sample size inflated by a Poisson sampled amount. Shown is the proportion of SNPs in each bin (solid lines) compared with the null expectation (dashed line). Black: original dataset. Dark grey: 10 reads added to each sample (same allele frequencies). Light grey: 100 reads added to each sample (same allele frequencies).