1
Broad geographic sampling reveals predictable, pervasive, and strong seasonal adaptation in Drosophila

Heather E. Machado[1,2]*‡, Alan O. Bergland[1,3]*‡, Ryan Taylor[1], Susanne Tilk[1], Emily Behrman[4], Kelly Dyer[5], Daniel K. Fabian[6,7], Thomas Flatt[6,8], Josefa González[9], Talia L. Karasov[10], Iryna Kozeretska[11,12], Brian P. Lazzaro[13], Thomas JS Merritt[14], John E. Pool[15], Katherine O'Brien[4], Subhash Rajpurohit[4], Paula R. Roy[16], Stephen W. Schaeffer[17], Svitlana Serga[9], Paul Schmidt[4o‡], Dmitri A. Petrov[1o‡]

* equal contribution
° equal contribution
‡ to whom correspondence should be addressed: heather.machado@sanger.ac.uk, aob2x@virginia.edu, schmidtp@sas.upenn.edu, dpetrov@stanford.edu

[1] Department of Biology, Stanford University, Stanford, CA
[2] Wellcome Trust Sanger Institute, Hinxton CB10 1SA, UK
[3] Department of Biology, University of Virginia, Charlottesville, VA
[4] Department of Biology, University of Pennsylvania, Philadelphia, PA
[5] Department of Genetics, University of Georgia, Athens, GA 30602
[6] Institute of Population Genetics, Vetmeduni Vienna, A-1210 Vienna, Austria
[7] Centre for Pathogen Evolution, Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, United Kingdom
[8] Department of Biology, University of Fribourg, CH-1700 Fribourg, Switzerland
[9] Institute of Evolutionary Biology, CSIC- Universitat Pompeu Fabra, Barcelona, Spain
[10] Department of Ecology and Evolution, University of Chicago, Chicago, Illinois 60637, USA
[11] Taras Shevchenko National University of Kyiv, Kyiv, Ukraine
[12] State Institution National Antarctic Center of Ministry of Education and Science of Ukraine, Taras Shevchenko Blvd., 01601, Kyiv, Ukraine
[13] Department of Entomology, Cornell University, Ithaca, NY, 14853
[14] Department of Chemistry & Biochemistry, Laurentian University, Sudbury, ON, Canada
[15] Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI
[16] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence KS 66045
[17] Department of Biology, The Pennsylvania State University, University Park, PA

Abstract

To advance our understanding of adaptation to temporally varying selection pressures, we identified signatures of seasonal adaptation occurring in parallel among *Drosophila melanogaster* populations. To study these evolutionary dynamics, we estimated allele frequencies genome-wide from flies sampled early and late in the growing season from 20 widely dispersed populations. We identify parallel seasonal allele frequency shifts across North America and Europe, demonstrating that seasonal adaptation is a general phenomenon of temperate fly populations. The direction of allele frequency change at seasonally variable polymorphisms can be predicted by weather conditions in the weeks prior to sampling, linking the environment and the genomic response to selection. The extent of allele frequency

47 fluctuations implies that seasonal evolution drives substantial (5-10%) allele frequency
48 fluctuations at >1% of common polymorphisms across the genome. Our results suggest that
49 fluctuating selection is an important evolutionary force affecting the extent and stability of linked
50 and functional variation.
51
52   *Introduction*
Fluctuations in the environment are an inescapable condition for all organisms. While many of
these fluctuations are unpredictable, some are predictable to a degree, including those that occur
on diurnal and seasonal time scales. The predictability of cyclic fluctuations is reflected in the
fact that many species exhibit plastic physiological and behavioral strategies that enable them to
survive the unfavorable season and exploit the favorable one (Denlinger 2003; Kostál 2006);
such plastic responses represent the classical form of seasonal adaptation (Tauber *et al.* 1986).
However, seasonally varying selection can - in principle - maintain fitness related genetic
variation if some genotypes have high fitness in one season but not another (Gillespie 1973).
Thus, in the organisms that undergo multiple generations per year (multivoltine), a distinct form
of seasonal adaptation occurs when the frequency of alternate genotypes changes in response to
seasonal fluctuations in the environment.
64
Seasonal adaptation can be seen as a form of local adaptation but local adaptation in time –
adaptation to temporally varying selection pressures - rather than in space. However, such
adaptation in time has been considered by some to be uncommon and, when present, unlikely to
result in long-term balancing selection (Ewing 1979; Hedrick 2006). Classic quantitative genetic
theory suggests that an optimal, plastic genotype will eventually dominate a population that is
exposed to periodically changing environments (Scheiner 1993). This is particularly so when
certain environmental cues are reliable indicators of changes in selection pressure (Levins 1968;
Via & Lande 1985). Predictions from traditional population genetic models suggested that
periodically changing environments will lead to the rapid loss of seasonally favored ecotypes as
slight changes in selection pressure from one year to another eventually push allele frequencies
at causal alleles to fixation (Hedrick 1976).
76
Recent theoretical models have called these classical predictions into question. For instance, a
quantitative genetic model by Botero et al. (Botero *et al.* 2015) examined whether adaptive
tracking, plasticity, or bet-hedging evolve as a consequence of environmental fluctuations. They
showed that adaptive tracking rather that plasticity is likely to become a predominant feature of
populations living in seasonally fluctuating environments when populations undergo more than
three generations per season. Additionally, a population genetic model by Wittmann et al
(Wittmann *et al.* 2017) has demonstrated that seasonally varying selection can maintain fitness
related genetic variation at many loci throughout the genome provided that dominance shifts
from season to season in such a way that, on average, the seasonally favored allele remains even
slightly dominant (Curtsinger et al 1994). These recent models, along with others that highlight
the importance of population cycles (Bertram & Masel 2019), as well as overlapping generations
and age structure (Ellner 1996; Ellner & Sasaki 1996; Ellner & Hairston 2015; Bertram & Masel
2019), suggest that seasonal adaptation and adaptive tracking (Kain *et al.* 2015) could be an
important feature of multivoltine organisms such as Drosophila (Behrman *et al.* 2015). More
generally, it is possible that adaptive tracking of environmental fluctuations on other time scales
might be more common than generally acknowledged.

Despite the lack of theoretical agreement on whether and how seasonal adaptation operates in multivoltine organisms, there is substantial empirical evidence for seasonal adaptation in many organisms including *Drosophila*. Seasonal adaptation was first observed in *D. pseudoobscura* by Dobzhansky and colleagues (e.g., Dobzhansky 1948) by tracking allele frequencies of inversions over seasons. Later studies confirmed and extended these early findings to other species including *D. melanogaster* (Stalker 1976; Stalker 1980; Kapun *et al.* 2016) and *D. subobscura* (Rodríguez-Trelles *et al.* 2013).

In *D. melanogaster*, multiple additional lines of evidence from phenotypic and genetic analyses demonstrate the presence of seasonal adaptation. When reared in a common laboratory environment, flies collected in the spring show higher stress tolerance (Behrman *et al.* 2015), greater propensity to enter reproductive dormancy (Schmidt & Conde 2006), increased innate immune function (Behrman *et al.* 2018), and modulated cuticular hydrocarbon profiles (Rajpurohit *et al.* 2017) as compared to flies collected in the fall. Rapid adaptation over seasonal time scales in these and related phenotypes has also been observed in laboratory (Schmidt & Conde 2006) and field-based mesocosm experiments (Rajpurohit *et al.* 2017; 2018). Genome-wide analysis indicated that a large number of common polymorphisms change in frequency over seasonal time scales in one mid-latitude orchard (Bergland *et al.* 2014), and allele frequency change among seasons has been observed using candidate gene approaches (Cogni *et al.* 2014). In several cases, these adaptively oscillating polymorphisms have been linked to seasonally varying phenotypes (Paaby *et al.* 2014; Behrman *et al.* 2018).

Despite ample evidence of seasonal adaptation in *D. melanogaster*, many aspects of this system remain unexplored. First, we do not know whether seasonal adaptation is a general feature of *D. melanogaster* populations across its range. Previous work (Schmidt and Conde 2006, Bergland *et al.* 2014, Cogni *et al.* 2014, Behrman et al. 2015, Behrman *et al.* 2018) detected seasonal fluctuations in a single locality over the span of one to three years. However, we do not know whether these results of fluctuating selection can be extrapolated to other localities. Second, it is unclear how responsive such adaptation is to specific local environmental conditions. Do seasons generate entirely idiosyncratic selective pressures that are not shared across populations, or does Drosophila experience partly generic selective pressures that are common to temperate populations? A related question is whether seasonal adaptation across the range involves a partially shared set of seasonal polymorphisms. Finally, we do not yet have estimates of the magnitude of effect and the fraction of polymorphisms genome-wide that are affected by seasonal adaptation through direct or linked selection, both in terms of the number of causal variants and the overall stochastic effect on linked variants.

The most straightforward and statistically powerful way to start answering these questions is to assess the extent of parallel shifts in allele frequencies over seasonal time across the *D. melanogaster* range. Should such parallel shifts be detected we can then infer that seasonal adaptation is widespread and at least partly driven by a common set of variants. It will also help us determine the magnitude of the shifts and give us a glimpse into the genetic architecture of seasonal adaptation, for instance whether or not the variants are found genome-wide or clustered into linked blocks or even supergenes.

Examining patterns of allele frequency change on seasonal timescales across a species' range
faces two problems. The first problem is logistical: the power of any analysis of parallel seasonal
adaptation across dispersed populations requires a large sample size. Such sampling can become
prohibitively expensive and logistically difficult for any single lab to carry out, and these
difficulties are further exacerbated if heavy constraints are placed on the exact timing of the
sampling. The second, and related problem, is the determination of the timing of sampling. We
have an intuitive sense of what seasons are; however, it is not clear how to match seasons across
locations. While it is possible, albeit logistically difficult, to sample flies according to some
matching of seasons using the calendar window (e.g. first week of June for the "spring" and the
first week of November for the "fall") or using a pre-determined physiological time (e.g.,
"spring" is the time after 21 cumulative growing degree days), it is not at all clear that any such
matching would be meaningful in terms of the ecology, physiology, and evolution of local
populations.
152
We have elected to solve these two related problems by (i) working as a consortium of
Drosophila biologists (DrosRTEC or the Drosophila Real Time Evolution Consortium) and (ii)
sampling in a somewhat haphazard manner. A similar effort is being carried out primarily in
Europe by the parallel *Drosophila* Population Genomics Consortium DrosEU (Kapun *et al.*
2018). Both consortia began as a result of the 2012 Catalysis Meeting at the National
Evolutionary Synthesis Center (NESCent) in Durham, NC, USA. These consortia organize a
collective effort of Drosophila biologists who sample at their own locations which then in turn
allows for the sampling to be carried out range-wide. We are continuing this effort and
expanding the consortium going forward ("*Drosophila Evolution in Space and Time*", DEST) to
meet this ambitious goal.
163
We chose to carry out haphazard sampling for logistical and inferential reasons. In this study, the
"spring" sample was generally taken close to the time when the populations of *D. melanogaster*
become locally abundant. The "fall" is either simply later in the summer or early fall, generally
prior to the time that populations crash at the onset of winter. Beyond the impracticality of
requiring that everyone collect flies at a given point in time we simply do not know when,
exactly, would be the most appropriate time to collect flies or what the fine-scaled temporal or
spatial factors matter for driving changes in allele frequency. Given this lack of knowledge,
choosing a specific time point, collection method, or substrate limits the covariance of collection
attributes with allele frequencies thereby preventing us from testing for specific gene-
environment associations *a posteriori*. Thus, a haphazard approach might in fact have the
greatest inferential power provided that the signal of seasonal adaptation can be detected in such
a complex dataset.
176
Using allele frequency data from 20 paired seasonal samples from 15 localities in North America
and Europe, we find evidence of parallel, pervasive, and strong seasonal adaptation in *D.*
melanogaster. First, we demonstrate that seasonal adaptation is a general phenomenon that
occurs in multiple populations of *D. melanogaster* on two continents. We provide evidence
suggesting that at least some of the same polymorphisms cycle between seasons in populations
sampled across vast geographic distances. Further, we show that allele frequency change
between seasons is predictable to some extent when taking into account weather in the weeks
immediately prior to sample collection, which hints at the complex and nonlinear nature of

seasonal adaptation. Seasonal alleles tend to show clinal variation, with the alleles that increase
in frequency through the summer generally being more frequent in lower latitude locations.
Finally, using a simulation approach, we show that the detected seasonal signal is consistent with
substantial (5-10%) seasonal allele frequency fluctuations at more than 1% of all common
polymorphisms. Taken together, our work demonstrates that seasonal adaptation is a general and
predictable feature of *D. melanogaster* populations and has pervasive effects on patterns of allele
frequencies genome-wide. More generally we establish that metazoan populations can exhibit
adaptive tracking of environmental conditions on extremely short time scales, most likely shorter
than a single growing season encompassing ~10 generations, suggesting that such rapid and
cyclic evolutionary shifts cannot be discounted *a priori*.
195
196   *Results*
Fly samples and sequence data. We assembled 72 samples of *D. melanogaster* collected from 23
localities in North America and Europe (Supplemental Table 1, Supplemental Figure 1). For 15
sampling localities, flies were collected in the spring and fall over the course of one to six years
(Figure 1A). Our sampling and sequencing efforts also complement other genome-wide datasets
that investigate genetic variation within and among *D. melanogaster* populations throughout
their range (Kolaczkowski *et al.* 2011; Mackay *et al.* 2012; Pool *et al.* 2012; Campo *et al.* 2013;
Reinhardt *et al.* 2014; Kim *et al.* 2014; Grenier *et al.* 2015; Svetec *et al.* 2016; Lack *et al.* 2016;
Zhao & Begun 2017; Kapun *et al*. 2018).
205
For our analysis, we divided our samples into three subsets (Figure 1B). The first subset
(hereafter 'Core20') is composed of 20 populations consisting of one spring and one fall sample.
Populations in the Core20 set are drawn from 15 localities in North America and Europe and we
use at most two years of sampling from any one locality. Samples in the Core20 set are used in
our basic analyses of seasonal adaptation. The second subset (hereafter 'ValidationSet') is
composed of spring and fall samples from four populations. Populations in the ValidationSet are
drawn from the two localities where there are more than two years of sampling. The
ValidationSet is used as a part of our leave-one-out analysis. The third subset (hereafter the
'clinal' set) of samples was used to examine patterns of clinal variation along the East Coast of
North America and consists of four populations sampled in the spring (see Material and Methods
and Supplemental Table 1).
217
To estimate allele frequencies genome-wide we performed pooled DNA sequencing of multiple
individual flies per population (Zhu *et al.* 2012; Schlötterer *et al.* 2014). For each sample, we
resequenced pools of ~75 individuals (range 27-164) to an average of 94x coverage (range 22-
220, Supplemental Table 1). Analyses presented here use a total of 51 samples. Data from the
remaining samples, including those that do not constitute paired spring/fall samples, are not
included in our analyses here but nonetheless form a part of our larger community-based
sampling and resequencing effort and are deposited with the rest of the data. All raw sequence
data have been deposited at the NCBI Short Read Archive (SRA; BioProject Accession
#PRJNA308584; individual accession numbers for each sample can be found in Supplemental
Table 1), a VCF file with allele frequencies from all populations, scripts of the analyses
presented here, and genome-wide SNP statistics for seasonality and clinality are available on
DataDryad (doi:10.5061/dryad.4r7b826).

230
Pooled resequencing identified ~1.75M SNPs following quality control and filtering for read
depth and average minor allele frequency (minor allele frequency > 0.01). We applied a second
filtering to identify ~775,000 SNPs that have observed polymorphism in each population
sampled. Unless otherwise noted, we use the smaller set of more common SNPs. Whether this
SNP selection process generates bias in our estimates of the strength and magnitude of
seasonally variable selection remains to be determined but note that the smaller set still
represents approximately half of all detected SNPs.

To gain insight into basic patterns of differentiation among the sampled populations, we
performed a principal component (PC) analysis across all samples. Samples cluster by locality
and geography (Figure 1C) with PC 1 separating European, North American West Coast and
East Coast populations while PC 2 separates the eastern North America samples by latitude
(linear regression $p = 3*10^{-15}$, $R^2 = 0.84$). No PC is associated with season, as defined by the time
of collection, following correction for multiple testing.

Figure 1:

247
248

Figure 1. Sampling times, localities, and basic population structure of samples used in this
study. A) Distribution of collection times during the spring (red) and fall (blue) in relation to
latitude. For samples where the collection month, but not day, was recorded, we specified the
15th of the month to calculate Julian Day. B) Sampling localities for the primary seasonal
analysis ('Core20': green), the cross-validation analysis ('ValidationSet': blue) and the
latitudinal cline analysis ('Cline': red), distributed across North America and Europe. Numbers

represent the number of years of spring/fall comparisons at that locality. C) Principal
components analysis of SNP allele frequencies. Circles are placed around samples from three
geographic regions: Europe (EU), California (CA) and Eastern North America (East NA); points
are colored by latitude and shapes represent seasons, as defined by collection time.

Seasonal adaptation is a general feature of fly populations. Our first objective was to determine
whether or not we find evidence of seasonally cycling genetic variation. To identify such a signal
of allele frequency change over time we assessed the per-SNP change in allele frequency
between spring and fall among the Core20 set of populations. We first used a generalized linear
model (GLM), regressing allele frequency at each SNP on season across all of the populations
(see Materials and Methods for details). Indeed, we find that the observed genome-wide
distribution of 'seasonal' p-values from this model is not uniform and there is a pronounced
excess of SNPs with low p-values (Figure 2A; compare the red line for the observed data with
the grey lines for the shuffled data). We assessed the statistical significance of this result by
utilizing the paired spring/fall nature of our dataset. Specifically, we performed a permutation
analysis whereby we shuffled the seasonal labels within populations to create 100 permuted
datasets and then refit the seasonal GLM model to each such permutation. These permutations
have the advantage of retaining all of the features of the data except for the seasonal labels
within populations. We do find an excess of low p-values in the observed distribution compared
to the permutations (Figure 2A), with a greater proportion of SNPs with $p<0.001$ in the
unpermuted dataset than in 98% of the permutations. Our permutation analysis thus suggests that
the genome-wide excess of SNPs with low p-values contains a signal of parallel seasonal
adaptation across multiple populations.

To further verify the observed signal of seasonal adaptation, we used a second statistical method
for measuring seasonal variation. For this we developed a semi-parametric test, the "rank
Fisher's method" (RFM), which is calculated by combining the per-population changes in allele
frequency (represented as p-values from a Fisher's exact test) between the two seasons
normalized by the genome-wide distribution of SNPs with similar attributes (see Materials and
Methods for details). This model was motivated by the observation that the GLM model can be
affected by an inflated precision of allele frequency estimates (Supplemental Figure 2) or other
statistical complexities (Machado *et al* 2016; Wiberg *et al* 2017) and that the observed
distribution of the p-values in the GLM is shifted to small values even in the shuffled datasets
(Figure 2A). The limitations of the GLM are particularly relevant, as our data are from pooled
sequencing, for which we can only estimate an effective sample size (Machado *et al.* 2016). The
RFM, by contrast, is relatively insensitive to the sample size estimate (Supplemental Figure 2).
Using the RFM we again find a deviation from the expected test statistic distribution indicating
an excess of seasonally varying polymorphisms (Figure 2B; compare the red line for the
observed data with the grey lines for the shuffled data). Performing a permutation analysis
analogous to that used for the GLM, we found the observed enrichment to be greater than in
98.2% of permutations. Together, our GLM and RFM results provide robust evidence that
parallel seasonal adaptation is a general feature of these fly populations.

Predictability of seasonal adaptation among populations

The genome-wide signature of seasonal adaptation that we observe indicates that there are
consistent changes in allele frequencies between seasons, broadly defined, among populations
sampled across multiple years, and localities separated by thousands of miles (Figure 1A,B).
However, it is not yet clear whether all populations show signatures of seasonal adaptation at a
common set of SNPs and if not, whether such populations are associated with specific
environmental effects, such as unusually short summer or an unusually mild winter. To
investigate these questions, we performed a leave-one-out analysis. In this analysis, we
sequentially removed each of the 20 paired spring-fall populations within the Core20 set and re-
identified seasonally variable SNPs among the remaining 19 populations (hereafter, 'discovery
set') as well as for the dropped, 20[th] population ('test set'). We then tested (1) if there is an
enrichment of SNPs showing signals of seasonal evolution in both the discovery and test sets,
and (2) if allele frequencies changed in the same direction (i.e., concordant allele frequency
change) for SNPs that show a significant allele frequency change in both the discovery and test
sets (see Materials and Methods for more details). For each population, we then calculated a
'*predictability score*'. This score reflects the genome-wide rate of concordance change as a
function of the joint significance threshold. Note that populations with a positive predictability
score are those in which the concordance rate is greater than 50% over the bulk of the genome or
chromosome.
317
Our predictability analysis yielded several basic results. First, we find that there is an enrichment of
polymorphisms identified as strongly seasonal in both the test and discovery sets for a limited number of
populations. For instance, at a joint significance quantile of 1%, 4 out of the 20 test sets show a
significant enrichment of seasonal SNPs compared to their conjugate discovery set (nominal *p*-value <
0.05 based on genome-wide Fisher's exact test; Supplemental Figure 3). The populations that show
significant enrichment of strongly seasonal sites in the discovery and test sets include: Esparto, CA,
USA, 2012 (CA_es); Gross-Enzersdorf, Austria 2012 (AT_gr); Sudbury, Ontario, Canada 2015
(ON_su); and, Charlottesville, VA 2015 (VA_ch). For these populations, enrichment was modest – at
most on the order of 25-50%. Although only four populations show nominally significant enrichment at
a joint significance quantile of 1%, 14 out of 20 populations show a positive enrichment, and no
population shows a significant negative enrichment at a joint significance threshold of 1%. It is unclear
whether the extent of this modest, yet sometimes statistically significant, enrichment is due to statistical
power, sampling effort, ecology, or the idiosyncratic nature of the evolutionary process (Gould 1989).
331
Second, we find that the majority of populations show a positive predictability score (Figure 2C).
For populations that show the strongest positive predictability score, the concordance rate
reaches ~65% (range 52-69%; all populations significantly different from 50% at a nominal *p*-
value < 0.05) among the ~65 (range 36-199) SNPs with the most stringent joint significance
threshold. At more modest joint significance thresholds (e.g., the top 10% in both sets), the
concordance rate is lower but still statistically different from 50% (maximum nominal *p*-value at
the top 5% of sites is 0.008). Intriguingly, we found that four of the 20 populations had
significantly negative predictability scores (Benton Harbor, Michigan 2014; Lancaster,
Massachusetts 2012; Topeka, Kansas 2014; and Esparto, California 2012). For these populations,
the strength of negative predictability was, in general, as strong as those populations with
positive predictability. The predictability score per population does not vary in a systematic way
with enrichment (e.g., at a joint significance threshold of 1%, the correlation between enrichment
and predictability score is -0.25, *p* = 0.41) and, notably, populations with negative predictability

scores have some of the strongest enrichment in the leave-one-out analysis. The sign and
magnitude of predictability scores varies among chromosomes (Supplemental Figure 4),
although the genome-wide score is correlated with the per-chromosome scores, particularly for
chromosome arms 2L and 2R (Supplemental Figure 5).
349
350 **Figure 2.**
351

352
353
Figure 2. Generality and predictability of seasonal adaptation. A) GLM *p*-value distribution
of per-SNP seasonal regressions (red = original dataset in which all seasonal labels are taken at
face value; blue = flipped dataset that was generated after detecting that some populations exhibit
the opposite pattern of fluctuations compared to the rest of the populations – see the text and
Figs. 2CDEF) and permutations (grey)). The y-axis represents the proportion of SNPs within *p*-
value bins between 0 and 1 with bin size 0.001. Note, the expected value of a uniform
distribution is 0.001, as represented by the dashed line. B) RFM X^2 distribution of per-SNP

We sought to explain why some populations display negative predictability by constructing a
weather based, thermal-limit model. We hypothesized that populations with negative genome-
wide predictability scores were exposed to warm springs or cool falls prior to our sampling. To
test this hypothesis, we calculated the number of days in the three weeks prior to spring sampling
when maximum temperatures fell above a specified upper thermal limit and the number of cool
days in the three weeks prior to fall sampling when minimum temperatures fell below a lower
limit. We refer to these counts of days as 'critical days.' The use of a three-week window to
assess critical days is somewhat arbitrary but we note that this duration corresponds to one to two
generations and the general period from egg to peak adult reproduction (McMillan *et al.* 1970;
Schmidt *et al.* 2005; Bergland *et al.* 2012; Klepsatel *et al.* 2013). Thus, a three-week window is a
biologically relevant time window. We tabulated the critical days for each population across a
range of spring and fall thermal limits (0°C - 40°C). For each combination of spring and fall
thermal limits, we regressed predictability scores on the number of spring and fall critical days
and assessed model fit in contrast with a null distribution obtained via permutation. We found
that the number of days prior to sampling in the spring above ca. 35°C (32-37°C) and number of
days in the fall below ca. 5°C (0-10°C) is significantly correlated with genome-wide
predictability scores of the 20 populations (Fig. 2D,E; $R^2_{max} = 75.7$, $p_{perm} = 0.0025$ for the most
significant set of spring and fall thermal limits). Inferred model parameters for the best fit models
(those with $p_{perm} < 0.05$) and critical days across the range of thermal limits for each population
can be found in Supplemental Tables 2 and 3, respectively.

These best fit models indicate that populations experiencing warm temperatures prior to the
spring sampling or cool temperatures prior to the fall sampling will have negative genome-wide
predictability scores (Figure 2E). This situation would naturally arise if these particular spring
and fall fly samples were collected at a tail of the season. However, genome-wide predictability
scores are not correlated with spring or fall collection date ($p = 0.6$ and 0.9, respectively),
cumulative growing degrees ($p = 0.5, 0.86$), collection locale (residential vs. orchard, $p = 0.6$,
0.4), collection method (aspirator vs. net vs. trap, $p = 0.9, 0.9$), collection substrate (contrasting
various types of fruit, compost, $p = 0.8, 0.9$), percent *D. simulans* contamination (0.8, 0.8),
latitude ($p = 0.9$), or the difference in cumulative growing degree days between spring and fall (p

407   = 0.46). The genome-wide predictability score is more related to the specific temperature
408   patterns observed in the spring and the fall than any other factors tested. Although we cannot rule
409   out other explanations such as inadvertent sample swapping, our analysis is generally consistent
410   with a model which suggests that changes in selection pressure within and among seasons may
411   lead to dramatic changes in allele frequencies.
412
We tested the predictive power of our thermal limit model by applying it to the four populations
in the ValidationSet (Figure 1B). We find that populations in the ValidationSet show a similar
variability of predictability scores as those estimated in the leave-one-out analysis of the Core20
set and that two populations have negative predictability scores (Supplemental Figure 6B;
Media, Pennsylvania 2012; Cross Plains, Wisconsin 2013). Next, we used the parameters from
the best fit thermal models derived from the Core20 set (i.e., those where $p_{perm} < 0.05$; Figure
2D,E; Supplemental Table 2,3) to model the genome-wide predictability scores of the
ValidationSet populations for which collection dates are known. We find that the modeled
genome-wide predictability scores are positively correlated with the observed genome-wide
predictability score of the ValidationSet populations (Fig 2F; median Pearson's $r = 0.37$ across
the range of best-fit models, minimum Pearson's $r = 0.09$, maximum $r = 0.60$; see Material and
Methods). While this correlation of three points is not significantly different from zero ($p = 0.77$,
based on the best fit thermal model), we nonetheless find this correlation is greater than 76% of
the best-fit permuted models (range 56 - 89% across all models with $p_{perm} < 0.05$). Taken
together with the leave-one-out analysis of the Core20 set, we conclude that coarse-grained
temperature data of the weeks prior to sampling is sufficient to predict some of the seasonal
changes in allele frequencies genome-wide. Our conclusion is consistent with work in other
drosophilids which demonstrate that short term changes in temperature, either directly or
indirectly, elicits dramatic changes in allele frequencies in the wild and in the laboratory
(Rodríguez-Trelles *et al.* 2013; Tobler *et al.* 2014; Mallard *et al.* 2018; Barghi et al 2019).
433
The flipped model. In the analysis presented above, we used the time of year that flies were
collected to designate season. However, our leave-one-out analysis and thermal limit model
suggest that the designation of season based on time of year may not completely reflect the
recent selective history of the populations that we sampled. Notably, our analysis suggests that
some populations experienced a particularly warm spring or a cold fall prior to sampling, and
that such temperature extremes elicit dramatic changes in allele frequencies that are opposite
from what would be expected from the calendar time season designation. We therefore reasoned
that our ability to detect seasonally variable polymorphisms, along with our ability to assess
general signals of seasonal adaptation, would improve if we flip the season label of the
populations that show negative genome-wide predictability scores.
444
To evaluate this idea, we flipped the season label for the four populations with negative genome-
wide predictability (Figure 2C) and calculated seasonal p-values using a GLM (hereafter, the
'flipped model'). As expected, the genome-wide distribution of p-values from the flipped model
is more heavily enriched for low p-values than the original model. The flipped model out-
performs the original model in the permutation analysis, where there are more SNPs with
$p<0.001$ than 100% of 100 permutations for the flipped model (compared to 98% of the
permutations for the original model; Figure 2A). We also re-calculated the RFM statistic after
flipping the season labels on these four populations and again find a stronger signal than for the

original model and greater than 100% of the permutations (Figure 2B). Of course, the stronger
signal of seasonal allele frequency change in the flipped model is a foregone conclusion given
the particular way we are changing the seasonal labels.

To provide an orthogonal assessment of whether flipping the seasonal label for these four
populations improves signals of seasonal adaptation we re-ran our predictability analysis. In this
re-analysis, we calculated the predictability score for each ValidationSet population using the
flipped Core20 model as our discovery set. The observed predictability scores estimated using
the flipped Core20 model is more strongly correlated with the model-based estimates from our
thermal limit model: the median correlation between model-estimated and observed
predictability scores of $r = 0.51$ (original model: $r = 0.37$). This correlation coefficient is greater
than 83% of the best-fit permuted models (compared with 68.8% for the original model;
minimum 0.25, maximum 0.73; Supplemental Figure 6ABC). Thus, the use of the flipped model
as a discovery set increased the predictive power of our thermal limit model in an independent
dataset.

Another way to assess the increased predictive power of the flipped model is to examine the
overlap between seasonal SNPs identified here with those identified by Bergland *et al.* (2014).
Two populations are shared between those included in the Core20 and those analyzed in
Bergland *et al.* (2014) and these shared populations could drive the signal of enrichment that we
potentially observe. Accordingly, we identified the top 1% most seasonally variable SNPs from a
model fit with 18 of the Core20 populations that do not overlap with those used in the earlier
Bergland *et al.* (2014) study. These top SNPs identified here with 18 populations are marginally
enriched for those identified by Bergland *et al.* (2014) relative to control SNPs matched for
chromosome, heterozygosity, and average recombination rate ($\log_2$ odds ratio $\pm$ SD = 0.59 $\pm$
0.37, p_{perm} = 0.0512). Applying the flipped season labels for this reduced Core20 set substantially
improves the overlap ($\log_2$OR $\pm$ SD = 1.02 $\pm$ 0.39; p_{perm} = 6e-4). Note that the Pennsylvanian
populations studied by Bergland *et al.* (2014) were not among those populations with season
labels altered in our flipped model.

Taken together, we conclude that there is support for the idea that the flipped model generates a
higher confidence set of seasonally varying SNPs. For the remainder of our analyses, presented
below, we used seasonal SNPs identified by both the original and flipped models. In the results
that follow we find that the use of the flipped model generally increases the strength of the
signals that we observe, suggesting improved biological relevance of the flipped model.

Latitudinal differentiation parallels signatures of seasonal adaptation. Phenotypic and genetic
analyses have demonstrated local adaptation among populations of *D. melanogaster* sampled
across latitudinal clines (see Adrion *et al.* 2015 and references therein). Signals of local
adaptation across latitudinal gradients have even been suggested to result from clines in
seasonality (Rhomberg & Singh 1989): higher latitudes are more stressful as a consequence of
severe winters and shorter growing seasons whereas more benign conditions are found in tropical
and sub-tropical areas. Thus, spatial heterogeneity in selection pressures associated with
temperate environments may be paralleled by seasonal fluctuations in similar selection pressures.

 Figure 3.

500
501
Figure 3. Concordance rates of seasonal and clinal polymorphisms. A) Seasonal sites were
identified using 18 of the Core20 populations to avoid a shared collection locale (Linvilla,
Pennsylvania) with populations used for clinal analysis. The y-axis represents the concordance
rate of allele frequency change across time and space, assuming that the winter favored allele is
the same as the allele favored in high latitudes. Seasonal sites were also identified in B)
California populations (n=3) and C) Europe populations (n=3) and contrasted with clinal
polymorphisms identified along the East Coast of North America. Shaded areas are the 96%
confidence intervals of 100 matched control datasets. Stars represent significant differences
between the original and flipped concordance rates based on a Fisher's exact test (nominal $p <$
0.05).

512
Parallel changes in phenotype and genotype across time and space have been observed in *D.*
melanogaster (Cogni *et al.* 2014; Bergland *et al.* 2014; Paaby *et al.* 2014; Cogni *et al.* 2015;
Kapun *et al.* 2016; Rajpurohit *et al.* 2017; 2018; Kapun & Flatt 2019). To assess whether we
observe similar general patterns in our data, we tested whether SNPs that are clinal and seasonal
below a range of significance thresholds change in allele frequency in a parallel fashion. Parallel
changes in allele frequency across time and space occur when the sign of allele frequency change
between spring and fall is the same as between high- and low-latitude populations. We re-
identified seasonally varying SNPs using 18 of the Core20 populations to avoid a shared
collection locale with the clinal set (see Supplemental Table 1). Seasonally varying SNPs were
identified among these 18 populations using both the original and flipped seasonal labels.
Clinally varying polymorphisms were identified using single-locus models (GLM, with allele
frequency regressed against latitude) as well as models that account for population structure
(BayEnv2; Günther & Coop 2013).

526
We find that the rate of parallelism increases with increasingly stringent seasonal and clinal
significance thresholds (Figure 3A) using either model of clinality (Supplemental Figure 7).
Parallel changes in allele frequency between seasons and clines is similar to previously published
genome-wide (Bergland *et al.* 2014) and locus specific results (Stalker 1976; Stalker 1980;
Cogni *et al.* 2014; Kapun *et al* 2016). Parallelism rates are higher when assessing seasonal
variation using the flipped model as compared to when we use the original model (Figure 3A).

This increase in parallelism rates using the flipped model can be taken as yet another strong, and
orthogonal, evidence that the flipped model generates a higher confidence set of seasonal SNPs.
535
Based on these analyses, there is clear evidence that parallel changes in allele frequency occur
across time and space. Are these parallel changes driven by large scale seasonal migration? To
address this question, we identified seasonal SNPs using populations that are geographically and
genetically isolated from the East Coast (Campo *et al.* 2013; and see Figure 1A). We assessed
seasonal changes in allele frequency separately for Californian (n=3) and European (n=3)
populations and tested for parallelism with polymorphisms that vary along the East Coast of
North America. One of the Californian populations contained a 'flipped' sample and so we also
re-assessed seasonal changes in allele frequency for those populations using a flipped model. We
found a strong signal of parallelism between seasonal and latitudinal variation in both the
California and the Europe comparisons (Figure 3B,C). For the Californian populations,
parallelism was greater for the flipped model than for the original model (Figure 3B).
547
These results suggest that parallel changes in allele frequency between seasons and across the
latitudinal cline is not exclusively driven by seasonal migration in the spring from neighboring
southern populations. Our conclusion is consistent with patterns of enrichment of jointly
seasonal and clinal polymorphisms (Supplemental Figure 8). Notably, for the flipped seasonal
model, there is weak evidence of an enrichment of highly seasonal and clinal polymorphisms, a
pattern otherwise expected if seasonal migration was a strong determinant of seasonal evolution.
Indeed, strongly seasonal sites identified from the flipped model are enriched for moderately
clinal polymorphism, and *vice versa*, a pattern ostensibly consistent with independent seasonal
evolution among populations arrayed along a spatial gradient. In contrast, seasonal sites
identified with the original seasonal model, in which seasons are defined by calendar time and
not temperature, are significantly enriched for strongly clinal polymorphisms suggesting that
seasonal migration may contribute to certain aspects of seasonal evolution. We note, however,
that the observed enrichment of seasonal and clinal polymorphisms only represents a small
fraction of all seasonal sites: for instance, using the original model at a joint quantile of 1%, we
expect ~84 (95% CI based on bootstrap resampling of matched control polymorphism: 63-104)
of the top 1% (n=~7500) seasonally varying SNPs to be among the top 1% of clinal sites;
however, we observe ~115 to be clinal. Thus, while we observe a significant overlap of seasonal
and clinal polymorphisms the vast majority of strongly seasonal polymorphisms are not strongly
clinal, again suggesting that clinal and seasonal evolution in *D. melanogaster* are largely
decoupled. Additionally, our analysis presented here assumes that patterns of clinal variation
along the West Coast of North America and Europe are distinct from patterns of clinal variation
along the East Coast; whether or not this assumption is valid remains to an open question (but
see Oakeshott *et al.* 1982).
571
572   *The strength and genomic extent of seasonal adaptation*. Our results suggest that seasonal
573   adaptation is driven, at least in part, by changes in allele frequency at a shared set of
574   polymorphisms spread throughout the genome. Here, we investigate the magnitude of change
575   and the number of sites that vary seasonally.
576

Figure 4. **The strength and magnitude of seasonal adaptation.** A) Average spring/fall allele
frequency change for each of the top 1% of seasonally varying SNPs (red/blue) and non-
seasonal, matched control SNPs (black), as a function of the folded spring allele frequency.
Lines represent a moving average across SNPs with a given spring allele frequency. BC) ABC
estimates for the proportion of sites affected by seasonal selection and the associated *s* for the
original (A) and flipped model (B), showing the sum of squared distances (SSD) between the
observed and simulation RFM X^2 test statistic distributions (see Materials and Methods for
details**)**, over varying strengths of selection and proportion of sites under selection. Yellow
points represent the top 1% of simulated parameter value combinations, reflecting an ABC
acceptance rate of 0.01).

To gain basic insight into the magnitude of seasonally variable changes in allele frequency, we
first examined the difference in allele frequencies between seasons for the most significantly
seasonal polymorphisms. The top 1% most seasonally variable SNPs identified by the GLM
change in frequency by a difference of 4 and 8%, on average (Figure 4A). This change is not
different between the original and flipped models (Figure 4A), but is substantially smaller than
estimates of ~20% reported by Bergland *et al.* (Bergland *et al.* 2014).

To robustly estimate the strength and extent of seasonally varying selection, we used our semi-
parametric RFM test in an Approximate Bayesian Computation (ABC) analysis. This analysis
allows us to estimate the fraction of the genome affected by seasonal adaptation and the average
strength of seasonal selection by comparing the observed distributions of the test statistic from
both the flipped and original models to simulations where a fraction of polymorphisms was
randomly selected to change in frequency between seasons with a given selection coefficient (see
Materials and Methods for more details). Using the sum of the square distances between the
observed and simulated test statistic as our metric, we identified which parameter estimates were
most likely to produce the observed genome-wide distribution of the test statistic. The model we
employ for this analysis is based on one generation per season and assumes that every site in the
genome changes in frequency independently and is subject to an equal strength of selection. Note
that this model is appropriate here as we aim to estimate the proportion of both causal and linked
SNPs that change in frequency due to seasonal selection and not to assess the number of causal
sites. We note that our analysis is potentially biased because we are conditioning on
polymorphisms that are present in all populations tested, including North American and
European ones with different long-term dynamics and histories of colonization, expansion, and
admixture (Keller 2007; Bergland *et al.* 2016; Kapopoulou *et al.* 2018). The filtering requirement
that we impose may therefore bias our SNP set to those which are subject to balancing selection,

causing an upward bias in our estimates. We note, however, there are no qualitative differences
found as a function of filtering for common SNPs (compare Figure 4B to Supplemental Figure
9).
618
We find a ridge of best fit parameters, with a trade-off between the proportion of sites under
selection and strength of selection (Figure 4B,C). For the original model, the data are consistent
with a minimum of ~1% of sites subject to an effective cumulative seasonal selection coefficient
("*s*", for simplicity*)* of ~0.3, or a maximum of ~10% of sites at a *s* of ~0.05. For the flipped
model, the data are consistent with a minimum of ~2.5% at an *s* of ~0.3 and a maximum of ~5%
at an *s* of ~0.2. As expected, the estimated strength of selection and the fraction of the genome
affected by seasonally varying selection is greater for the flipped model than the original model
(Figure 4B vs. 4C). In order to assess whether and to what extent our estimates are confounded
by clustering of seasonally varying polymorphisms, we performed the same analyses using
datasets of subsampled polymorphisms every 1Kb and 5Kb (Supplemental Figure 10). We found
no substantial shift in our parameter estimates across subsampled datasets.
630
631   *The distribution of seasonally variable SNPs throughout the genome*. The analyses presented
632   above suggest that seasonal adaptation affects patterns of allele frequency genome-wide. We
633   therefore aimed to characterize several aspects of the genomic distribution and functional
634   enrichment of seasonally varying polymorphisms.
635
First, we examined seasonal changes in the frequency of the major, cosmopolitan inversions that
segregate in wild *D. melanogaster* populations. We estimated inversion frequency from pooled
allele frequency data by calculating the average frequency of SNPs previously identified as being
closely linked to these inversions (Kapun *et al.* 2014). With the exception of *In(2L)t*, we find that
most inversions are rare in the populations that we examined, segregating at less than 15%
frequency in most populations (Supplemental Figure 11). Using a simple sign test, we find that
no inversion consistently changes in frequency using season labels as defined in either the
original or the flipped model. We note that *In(2R)Ns* is at a higher frequency in the fall compared
to the spring in a quarter of the Core20 populations ($p_{binomial-test}$ = 0.048; Bonferroni adjusted p =
0.28), consistent with previously reported signals of seasonal change (Stalker 1976; Stalker
1980; Kapun *et al.* 2016; Kapun & Flatt 2019). However, *In(2R)Ns* segregates at a lower
frequency (<15% across populations surveyed) than many of the most strongly seasonal SNPs on
chromosome 2R (over 85% of the top seasonally varying SNPs have minor allele frequency
greater than 15%), and seasonal changes of *In(2R)Ns* are relatively mild compared to other
seasonally varying polymorphisms (cf. Figure 4A Supplemental Figure 11). We conclude that
inversions are not the exclusive drivers of seasonal adaptation in *D. melanogaster*, although we
cannot rule out their role in contributing to seasonal adaptation.
653
To determine if seasonally variable SNPs are non-randomly distributed throughout the genome,
we examined the abundance of the top 1% of seasonally variable SNPs (GLM model; n=7,748)
among bins of 100 consecutive SNPs. We found significant over-dispersion in the distribution of
seasonal SNPs per bin compared to both the theoretical and the matched control distributions
(Figure 5A; Kolmogorov-Smirnov test $p=10^{-16}$), indicating that seasonal SNPs are more
clustered than expected, consistent with an elevated seasonal F_{st} surrounding the most strongly

seasonal sites (Figure 5B). We cannot determine to what extent this clustering is due to linkage
to causal SNPs or to the clustering of causal seasonal variants. The distribution of control
polymorphisms was indistinguishable from the binomial expectation in our analysis
(Kolmogorov-Smirnov test p=1).
664
Although seasonally variable SNPs are non-randomly distributed throughout the genome, they
are nonetheless broadly distributed. To further quantify the distribution of these sites, we
calculated the probability of observing at least one of the top 1% most seasonally variable SNPs
among 1000 randomly selected genomic windows ranging in size from 100bp to 100Kb. We find
that seasonally variable SNPs are distributed across all autosomes but are enriched on
chromosome 2L (Figure 5C,D). Across the window sizes tested, the probability of observing at
least one top seasonal SNP in any given window is less than expected by chance given a Poisson
distribution, consistent with clustering at both large and small genomic scales (Figure 5C).
Despite significant clustering, seasonally variable SNPs show broad genomic distribution, as
evidenced by a >75% chance of identifying a seasonally variable SNP in any randomly selected
50Kb window (Figure 5C). The general signal of over-dispersion is similar for both the flipped
and original models (Figure 5A,C). The broad genomic distribution and large number of
seasonally variable SNPs suggests that seasonal adaptation is highly polygenic. A polygenic
model is consistent with the view that many of the traits underlying seasonal adaptation (Schmidt
& Conde 2006; Behrman *et al.* 2015; Betini *et al.* 2017; Rajpurohit *et al.* 2017; Noh *et al.* 2017;
Behrman *et al.* 2018; Rajpurohit *et al.* 2018), and adaptation to temperate environments in
general (Rand *et al.* 2010; Levine *et al.* 2011; Chen *et al.* 2012; Rajpurohit *et al.* 2013; Cockerell
et al. 2014; Svetec *et al.* 2015; Chakraborty & Fry 2016; Juneja *et al.* 2016; Bozicevic *et al.*
2016; Svetec *et al.* 2016), are themselves quantitative traits (Mackay & Huang 2018).
684
Next, we tested whether the seasonally varying polymorphisms that we identify here using either
the original or flipped model are enriched among different functional annotation classes
(Supplemental Figure 12). The top 1% most seasonally varying polymorphisms are not
significantly enriched for any functional class that we tested. The top 5% of seasonally varying
polymorphisms are weakly enriched for non-synonymous sites as compared to matched control
polymorphisms (enrichment = 8%, 95% CI = 2-14%, nominal p=0.01). The ratio of non-
synonymous to synonymous polymorphisms among the top 5% of seasonal SNPs is also
marginally higher than that for matched controls (enrichment = 7%, 95% CI = 1 − 15%, p=0.03).
We found no significant enrichment or under-enrichment for intergenic regions, introns, UTR's,
or coding regions.
695
Figure 5.

Figure 5. Genomic distribution of seasonal SNPs. A) Distribution of seasonal sites (quantile:
0.01) per 100 SNP window. The observed distribution (red and blue) is over-disbursed compared
to both the theoretical distribution (dark grey) and that of matched controls (light grey). Error
bars are 2 standard deviations. B) Spring/fall genetic differentiation (F_{st}) at highly seasonal SNPs
(distance = 0) and for SNPs with increasing distance from a seasonal SNP. Also plotted is the
mean (black) and 96% confidence interval (grey area) of 100 matched controls. C) Proportion of
windows with at least one seasonal SNP, with increasing window size (Kb). The observed
proportions (solid lines) and expected theoretical proportion (dashed line) is plotted for each
chromosome arm. D) Sliding window analysis of regions with high densities of seasonal SNPs.
Open circles denote regions of the genome with an excess of seasonally variable SNPs following
multiple testing correction. Red lines, above the x-axis, represent the strength of enrichment of

We performed a sliding-window analysis to identify regions of the genome with an excess of
seasonally varying polymorphisms. We found 25 regions of the genome with a significant excess
of seasonally varying polymorphisms for either model. The locations of these regions change
substantially between the 'original' and 'flipped' models (Figure 5D). For simplicity, we focus
the remainder of our analysis here on the regions identified using the 'flipped' model, however
the coordinates of all identified regions of both the 'original' and 'flipped' are reported in
Supplemental Table 4. Regions of the genome enriched for seasonally variable SNPs are found
on both arms of chromosome 2 and deficient along chromosome 3 ($\chi^2 = 22.9$, $p = 4e\text{-}5$).
Intriguingly, one of 40 most seasonally variable regions identified with the 'flipped model'
contains the genes *Tep2* and *Tep3,* which we have previously associated with seasonal variation
in immune function in flies (Behrman *et al.* 2018). The *Tep2/3* region contains 11 of the top
seasonally variable SNPs, ranking this region among the top 0.1% genomic windows. The
genomic window surrounding *couch-potato* (*cpo*), a seasonally varying gene (Cogni *et al.* 2014)
found to be associated with reproductive dormancy (Schmidt *et al.* 2008), has seven of the top
seasonally variable SNPs ranking it among the top 1% of windows genome-wide. Note that the
enrichment of seasonal SNPs surrounding *cpo* is not significant after conservative, multiple
testing correction ($p_{adjusted} = 1$) which would be appropriate for *de novo* discovery of such
regions. If we treat *cpo* as the true seasonal locus, this result implies that we currently lack power
to discover many causal genes involved in seasonal adaptation. There was no excess of seasonal
SNPs surrounding the insulin-like receptor gene (*InR*), wherein seasonally variable indels and
SNPs have been previously reported and linked to variation in life-history traits (Paaby *et al.*
2014).

Gene-ontology analysis (Huang *et al.* 2009a; b) of the 198 genes within the 25 regions we
identified show significant enrichment for different biological processes, cellular components,
and molecular function following multiple testing correction (Supplemental Table 5). For
instance, we find that genes associated with several molecular function categories including
inositol metabolism and phosphorylation are enriched following multiple testing correction
($p_{adjusted} < 0.013$). This result is *prima facie* consistent with a role of inositol related compounds
in drosophilid overwintering and cold-tolerance (Vesala *et al.* 2012). However, extreme caution
should be taken when interpreting this result, as in any GO analysis (Pavlidis *et al.* 2012), as it is
driven by a set of three, closely linked genes (CG17026, CG17028, CG17029).

Discussion
In this study we identify genomic signatures of seasonal adaptation in *D. melanogaster*. Our
approach relies on the detection of parallel changes in allele frequency from spring to fall across
20 populations. We use coarse definitions of spring and fall, with spring as the time of year close
to when *D. melanogaster* first becomes locally abundant and fall as the time close to the end of
the growing season but before populations become too scarce to sample (Ives 1970; Behrman et
al. 2015). We sampled populations across a wide geographic range, such that the length of
growing season as well as other environmental conditions vary substantially. Despite this

environmental heterogeneity and the coarseness of the definitions of spring and fall, we detect a
clear signal of parallel seasonal allele frequency shifts across these populations.
756
Our analysis revealing pervasive seasonal adaptation would not be possible without accurate
allele frequency estimates (Zhu *et al.* 2012), genome-wide, from paired spring-fall samples
collected across *D. melanogaster*'s range. Our approach used less expensive pooled sample
sequencing which were collected by the DrosRTEC consortium, to facilitate seasonal sampling
across broad spatial scales, which is mirrored in Europe by the DrosEU consortium (Kapun *et al.*
2018).
763
By detecting consistent seasonal changes in allele frequencies among multiple populations in
North America and Europe, we demonstrate that seasonal adaptation is a general phenomenon of
temperate *D. melanogaster* populations, and not restricted to a single orchard (Linvilla Orchard,
Media, Pennsylvania; Schmidt & Conde 2006; Cogni *et al.* 2014; Bergland *et al.* 2014; Behrman
et al 2015), biogeographic region (East Coast of North America; Behrman *et al* 2018), or even a
continent. Furthermore, the parallel frequency shifts across multiple populations that we describe
here suggests that seasonal adaptation across the whole range is at least partly driven by a
consistent set of loci. Note that we generally lack power to detect alleles that cycle in a small
subset of populations or are private to a single population. Therefore, we focus only on
polymorphisms that segregate in all populations – and which therefore have an overall high
population frequency - and only look at the signal of parallel cycling. Moreover, we did not
attempt to investigate cycling of structural variants such as indels and transposable elements that
are known to generate adaptive mutations at an appreciable rate (reviewed in Barrón *et al.* 2014).
Thus, our conclusions are limited to the shared genetic basis of seasonal adaptation as revealed
by the cycling of common SNPs.
779
Our ability to detect a shared genetic basis of seasonal adaptation across these populations
implies that our admittedly crude definitions of spring and fall nonetheless carry biological
meaning. It also implies that there are generic features in the environment that exert consistent
selective pressures despite the environmental heterogeneity across sampling times and locations.
It is possible that such generic selective pressures relate to temperature, humidity, population
density, resource availability, or other biotic and abiotic factors. However, what these selective
pressures are specifically and how they act across seasons remains to be determined. Note,
however, that our results here (Figure 2) and elsewhere (Bergland *et al.* 2014) suggest that
temperature extremes may be a strong proximate or ultimate selective agent.
789
Although we identified consistent patterns of evolution between spring and fall, not all
populations fit the general pattern, with some showing strong signals of reversed allele frequency
change (Figure 2C). These 'flipped' populations are geographically dispersed (California,
Kansas, Michigan, and Massachusetts) with samples collected over two non-consecutive years
(2012, 2014). We demonstrated that this flipping is associated with particularly warm springs
and particularly cool falls prior to sampling. Indeed, the ecological model we built using
information about local temperature extremes allowed us to predict the sign and magnitude of
allele frequency changes genome-wide for additional populations not used to build the model
(Figure 2F, Supplemental Figure 6). We thus hypothesize that, at the time of sampling, some of
our spring samples had already experienced sufficiently strong selection in the generically

summer way and that the fall samples had experienced selective pressures in the generically
winter way. This would result in seasonal alleles in the spring already showing fall-like patterns
and the ones in the fall showing spring-like patterns. More generally, it is likely that natural
selection does not push populations in the same consistent direction across the growing season
and that there are phases within the growing season with distinct ecological and selection
pressures. In the future, more dense temporal sampling across the growing season might help
map the action of natural selection to more specific environmental factors. Taken at face value
the implication is that metazoan populations are capable of adaptive tracking on extremely short
temporal scales on the order of less than 10 generations. These data thus further argue that
evolution and ecology do often operate on similar timescales (Hendry 2009) and evolutionary
adaptation cannot be generally ignored *a priori* in demographic or ecological investigations
(Rudman *et al*. 2017).

Our analysis allowed us to estimate the proportion of the genome affected by seasonal cycling
and the magnitude of seasonal fluctuations. Our best guess is that the frequency of ~4% of
common polymorphisms are affected by seasonally variable selection coefficients, cumulatively
on the order of ~20% per season selection. However, there is a large range of plausible values
that tradeoff the number of sites (1-10% of all sites) with the strength of selection (10-30% per
season). Note that we do not claim that these 1-10% of common polymorphisms are causal nor
that they cycle independently. In fact, we speculate that the vast majority of the seasonal sites are
only linked to causal loci and many cycle together by fluctuating linked selection. Indeed, the
number of seasonal loci is implausibly large for all of them to be individually causal (Bergland *et*
al. 2014) under any of these scenarios. The lack of strong enrichment for specifically functional
sites such as non-synonymous compared to synonymous loci (Supplemental Figure 12) is also
consistent with the vast majority not being causal. At the same time, our model of unlinked loci
captures the breadth and magnitude of such linked selection and demonstrates that seasonal
adaptation is affecting, either directly or indirectly, a large number of variants across a range of
populations. Given that we can only detect consistent pattern of seasonal cycling we are likely to
underestimate the impact of linked seasonal selection. Seasonal adaptation therefore constitutes a
profound stochastic evolutionary force acting on linked variation that is likely to rival in strength
genetic draft due to selective sweeps and should be much stronger over seasonal timescales than
random genetic drift in populations as large as *D. melanogaster* (Karasov *et al.* 2010).

While we cannot determine the precise number and location of causal sites in the present study,
the the broad distribution of the seasonal SNPs across the genome implies that seasonal
adaptation is highly polygenic. Indeed, there is greater than a 75% chance that a random 50Kb
region of the genome will have at least one strongly seasonal site, reminiscent of the data on the
density of GWAS hits in human populations (Boyle *et al.* 2017). Therefore, the cycling we
observe cannot be due to a single causal region such as an inversion or a region of particularly
low recombination. At the same time, seasonal SNPs are significantly clustered, suggesting that
some genomic regions are more likely to contain one or more causal sites. We conducted a
screen for regions of the genome with an excess of highly seasonal SNPs (Figure 5D). We found
some outlier regions contain genes previously linked to seasonally variable phenotypes (
immunity and *Tep2*/*Tep3* - Behrman *et al.* 2018; reproductive dormancy - *cpo* (Cogni *et al.*
2014; Schmidt *et al.* 2008). Additional sampling may reduce the false positive rate and should
help us map individual seasonally causal loci.

846
Our observation that seasonal adaptation is polygenic and rapid suggests that there exists a
substantial number of common polymorphisms that are subject to strong fluctuating selection
and which are nonetheless maintained across the range of *D. melanogaster*. The most
straightforward explanation of these results is that these polymorphisms are subject to some form
of balancing selection (Levene 1953; Wittmann *et al.* 2017; Bertram & Masel 2019). Pervasive
balancing selection is consistent with the recent realization that simple models of mutation-
selection balance are inconsistent with the extent of quantitative genetic variation in a variety of
fitness related traits (Charlesworth 2015). A full understanding of the genomic consequences of
balancing selection caused by strong fluctuating selection will require the identification of
individual causal loci. Doing so will require substantial additional sampling across seasons as
well as from closely and broadly distributed locales throughout the whole geographic range of *D.*
melanogaster in combination with the genetic mapping of seasonally fluctuating phenotypes
(Schmidt & Conde 2006; Behrman *et al.* 2015; Rajpurohit *et al.* 2017; Behrman *et al.* 2018;
Rajpurohit *et al.* 2018). Finally, such efforts can be strongly augmented by manipulative field
experiments and functional work to investigate the molecular effects of putatively causal genetic
variants. All of this work will need to be done in the context of other processes that underlie
persistence of populations in the face of environmental heterogeneity such as phenotypic
plasticity and bet hedging (Via & Lande 1985; David *et al.* 1997; Ayrinhac *et al.* 2004; Bergland
et al. 2008; MacMillan *et al.* 2015; Overgaard *et al.* 2015; Kain *et al.* 2015) as it is becoming
clear that all of these processes are occurring contemporaneously with seasonal adaptation.
867
Materials and methods
Population sampling and sequence data. We assembled 72 samples of *D. melanogaster*, 60
representing newly collected and sequenced samples and 26 representing previously published
samples (Bergland *et al.* 2014; Kapun *et al.* 2016). Locations, collection dates, number of
individuals sampled, and depth of sequencing for all samples are listed in Supplemental Table 1.
For each sample, members of the Drosophila Real-Time Evolution Consortium collected an
average of 75 male flies using direct aspiration from substrate, netting, or trapping at orchards
and residential areas. Flies were confirmed to be *D. melanogaster* by examination of the male
genital arch. We extracted DNA by first pooling all individuals from a sample, grinding the
tissue together in extraction buffer, and using a lithium chloride – potassium acetate extraction
protocol (see Bergland *et al.* 2014 for details on buffers and solutions). We prepared sequencing
libraries using a modified Illumina protocol (Bergland *et al.* 2014) and Illumina TrueSeq
adapters. Paired-end 125bp libraries were sequenced to an average of 94x coverage either at the
Stanford Sequencing Service Center on an Illumina HiSeq 2000, or at the Stanford Functional
Genomics facility on an Illumina HiSeq 4000.
883
The following sequence data processing was performed on both the new and the previously
published data. We trimmed low-quality 3' and 5' read ends (sequence quality < 20) using the
program *cutadapt* v1.8.1 (Martin 2011). We mapped the raw reads to the *D. melanogaster*
genome v5.5 (and for *D. simulans* genome v2.01, flybase.org) using *bwa* v0.7.12 *mem*
algorithms, with default parameters (Li & Durbin 2009), and used the program *SAMtools* v1.2
for bam file manipulation (functions *index*, *sort*, and *mpileup*) (Li *et al.* 2009). We used the
program *picard* v2.0.1 to remove PCR duplicates (http://picard.sourceforge.net) and the program

GATK v3.2-2 for indel realignment (McKenna *et al.* 2010). We called SNPs and indels using the
program *VarScan* v2.3.8 using a *p*-value of 0.05, minimum variant frequency of 0.005, minimum
average quality of 20, and minimum coverage of 10 (Koboldt *et al.* 2012). We filtered out SNPs
within 10bp of an indel (they are more likely to be spurious), variants in repetitive regions
(identified by *RepeatMasker* and downloaded from the UCSC Genome browser), SNPs with a
median frequency of less than 1% across populations, regions with low recombination rates (~0
cM/Mb; Comeron *et al.* 2012), and nucleotides with more than two alleles. Because we
sequenced only male individuals, the X chromosome had lower coverage and was not used in our
analysis. After filtering, we had a total of 1,763,522 autosomal SNPs. This set was further
filtered to include only SNPs found polymorphic in all samples ("common polymorphisms"),
resulting in 774,651 SNPs that represent the core set used in our analyses.

Due to the phenotypic similarity of the species *D. melanogaster* and *D. simulans*, we tested for
D. simulans contamination by competitively mapping reads to both genomes. We omitted any of
our pooled samples with greater than 5% of reads mapping preferentially to *D. simulans*
(Charlottesville, VA 2012, fall; Media, PA 2013 fall). For the remaining samples reads that
mapped better to the *D. simulans* were removed from the analysis. For the populations used in
this analysis, the average proportion of reads mapping preferentially to *D. simulans* was less than
1% (see Supplemental Table 1), and there was no significant difference in the level of *D.*
simulans contamination between spring and fall samples (t-test *p*=0.90).

In silico simulation analysis shows that competitive mapping accurately estimates the fraction of
D. simulans contamination. To conduct these simulations, we used a single *D. simulans* genome
from an inbred line derived from an individual caught California (Signor *et al.* 2017) and a single
D. melanogaster genome from a DGRP line (Mackay *et al.* 2012). We used *wgsim* to generate
simulated short-reads from each genome and mixed these short reads together in various
proportions. We then mapped the short reads back to the composite genome for competitive
mapping, as described above. We calculated contamination rate as the number of total reads
mapping to the *D. simulans* reference genome divided by the number of reads mapping to both
D. simulans and *D. melanogaster*. We also calculated the locus specific residual cross mapping
rate by calculating the number of *D. simulans* reads still mapping to the *D. melanogaster* genome
even after competitive mapping. These simulations demonstrate that the estimation of the cross-
species mapping is precise (Pearson's $r = 0.9999$, $p = 1.6 \times 10^{-24}$, Supplemental Figure 13A), but
underestimates the true contamination rate by ~2%. The level of residual *D. simulans*
contamination does not correlate with contamination rate and is roughly 9% (Supplemental
Figure 13B). Additionally, the proportion of seasonal sites (top 1%) is not correlated with the
rate of residual mapping of *D. simulans* reads to the *D. melanogaster* genome as inferred through
our simulation ($r = -0.0026$, $p=0.93$).

To assess seasonal variation we analyzed population genomic sequence data from 20 spring and
20 fall samples ('Core20'). These samples represent a subset of the sequenced samples. We used
samples that had both a spring and a fall samples taken from the same locality in the same year.
We also used a maximum of two years of samples for a given locality to prevent the analysis
from being dominated by characteristics of a single population. When there was more than two
years of samples for a given population, we chose to use the two years with the earliest spring

collection time. This decision was made on the assumption that the earlier spring collection
would better represent allele frequencies following overwintering selection. This left 20 paired
spring/fall samples, taken from 12 North American localities spread across 6 years and 3
European localities across 2 years (Supplemental Table 1). The localities and years of sampling
are as follows: Esparto, California 2012 and 2013; Tuolumne, California 2013; Lancaster,
Massachusetts 2012 and 2014; Media, Pennsylvania 2010 and 2011; Ithaca, New York 2012;
Cross Plains, Wisconsin 2012 and 2014; Athens, Georgia 2014; Charlottesville, Virginia 2014
and 2015, State College, Pennsylvania 2014; Topeka, Kansas 2014; Sudbury, Ontario, Canada
2015; Benton Harbor, Michigan 2014, Barcelona, Spain 2012; Gross-Enzersdorf, Vienna 2012;
Odesa, Ukraine 2013. The four sets of paired spring/fall samples that were not included in the
Core20 set were used for cross-validation ('ValidationSet': Media, Pennsylvania 2009, 2012,
2014, Cross Plains, Wisconsin 2013). For comparison of seasonal with latitudinal variation, we
used sequence data from four spring samples along the east coast of the United States
(Homestead, Florida 2010; Hahia, Georgia 2008; Eutawville, South Carolina 2010, Linvilla,
Pennsylvania 2009). We performed a principal components analysis of allele frequency per SNP
per sample using the *prcomp* function in *R* (frequency data scaled by SNP).

GLM identification of seasonal sites. We identified seasonal sites using two separate methods, a
general linear regression model (GLM) and a rank Fisher's method (RFM). All statistical
analyses were performed in *R* (R Core Team 2014). To perform the GLM we used the *glm*
function with binomial error, weighted by the "effective coverage" (N_c)- a measure of the
number of chromosomes sampled, adjusted by the read depth:

$$N_c = (1/N + 1/R)^{-1}$$

where N is the number of chromosomes in the pool and R is the read depth at that site
(Kolaczkowski *et al.* 2011; Feder *et al.* 2012; Bergland *et al.* 2014; Machado *et al.* 2016). This
adjusts for the additional error introduced by sampling of the pool at the time of sequencing. The
seasonal GLM is a regression of allele frequency by season (e.g., spring versus fall):

$$y_i = \text{season} + \text{population} + e_i$$

where y_i is the allele frequency at the i[th] SNP, and e_i is the binomial error at the i[th] SNP.
Although the GLM is a powerful test, the corrected binomial sampling variance (N_c) is likely an
anti-conservative estimate of the true sampling variance associated with pooled sequencing
(Machado *et al.* 2016; Spitzer et al 2019). Therefore, we use the results of this test as a seasonal
outlier test, rather than an absolute measure of the deviation from genome-wide null expectation
of no allele frequency change between seasons.

RFM identification of seasonal sites.
As a secondary measure of enrichment of seasonally varying sites above neutrality, as well as to
estimate the proportion and average selection coefficient of seasonally varying sites, we used a
method in which calculated combined *p*-values based on class-based, ranked *p*-values from a
Fisher's exact test estimated for each population. In this "rank Fisher's method" (RFM), we
calculated a per-SNP *p*-value for a single spring/fall population comparison using a Fisher's

exact test. Then, *p*-values for either a spring to fall increase in allele frequency or spring to fall
decrease in allele frequency (i.e., each tail tested separately) were ranked among all other SNPs
with the same total number of reads and same total number of alternate reads. The class-based
rank of each SNP becomes its new *p*-value, providing uniform sets of *p*-values across the
genome. These rank *p*-value calculations were made for each of the 20 CoreSet populations,
separately. As the power (minimum *p*-value) is relative to the number of SNPs being ranked per
bin, and as the bin sizes are equivalent across populations, each population has equivalent power.
We then combine ranked *p*-values using Fisher's method for each SNP, taking two times the
negative sum of logged *p*-values across the 20 spring/fall comparisons (each tail tested
separately). The null distribution for this statistic (X^2) is a Chi-squared distribution with degrees
of freedom equal to 40 (two times the number of comparisons). We used the distribution of these
per-SNP X^2 test statistics to 1) test for a seasonal signal above noise and 2) perform an ABC
estimation of selection parameters (see below, *The strength …*).

Matched controls. With the assumption that the majority of the genome is not seasonal, we can
use matched genomic controls as a null distribution to test for enrichment of different features of
seasonal polymorphisms. The use of matched control SNP sets was employed for tests of
enrichment of clinal polymorphism, enrichment with seasonal SNPs identified by Bergland *et al.*
(Bergland *et al.* 2014), and in tests of genomic clustering. We matched each SNP identified as
significantly seasonal (at a range *p*-values) to another SNP, matched for chromosome, effective
coverage, median spring allele frequency, inversion status either within or outside of the major
inversions *In(2L)t, In(2R)NS, In(3L)P, In(3R)K, In(3R)Mo*, and *In(3R)P*, and recombination rate.
We used the same *D. melanogaster* inversion breakpoints used in Corbett-Detig & Hartl (2012)
and the recombination rates from Comeron *et al.* (2012). We randomly sampled 100 of the
possible matches per SNP (excluding the focal SNP) to produce 100 matched control sets. Any
SNPs with fewer than 10 possible matches were discarded from the matched control analyses.
We defined 95% confidence intervals from the matched controls as the 3rd and 98th ranked values
for the quantity being tested (e.g., percent concordance or proportion of genic classes).

1009

Leave-one-out analysis. To test the general predictability of seasonal change in our dataset, we
performed a leave-one-out analysis. In this analysis, we performed seasonal regressions for
subsets of the data, dropping one paired sample and comparing it to a seasonal test of the
remaining 19. We then measured the percent concordance of spring/fall allele frequency change,
defined as the proportion of SNPs that agree in the direction of allele frequency change and sign
of regression coefficient. This was performed 20 times, once for each paired sample.

1016

1017   To estimate genome-wide (or chromosome specific) predictability scores, we calculated the rate
1018   of change of the concordance score as a function of quantile threshold. To estimate this rate, we
1019   used the generalized linear model with binomial error,

1020 $concordance_i = quantile_i + e,$

where 'concordance' is the fraction of SNPs falling below the i^{th} quantile threshold of both the
target (19 population) and test (1 population) models that changed in frequency in a concordance
fashion and *e* is the binomial error with weights corresponding to the total number of SNPs
falling below that threshold. Thus, the genome-wide (or chromosome specific) predictability

score is heavily influenced by concordance rates of higher quantiles because that is where the
bulk of SNPs reside.

1027

The thermal limit model. To test whether heterogeneity in genome-wide predictability scores can
be explained by aspects of weather, we obtained climate data (daily minimum and maximum
temperatures) from the Global Historical Climatology Network-Daily Database (Menne *et al*
2012). We matched each locality to a weather station based on geographic distance. For the
Core20 set of populations, three of the collections did not have precise collection dates for one or
both seasonal collections (Athens, Georgia 2014; Media, Pennsylvania 2010, 2011), or were not
associated with any climate data from the GHCND database (Odesa, Ukraine); for the
ValidationSet populations, one of the populations did not have precise collection dates for one of
both seasonal collections (Media, Pennsylvania 2009). These populations were removed from
our weather model analysis.

1038

1039 To fit the thermal limit model for the Core20 and ValidationSet populations, we regressed the
1040 observed predictability score on the number of days above and below the thermal limits using a
1041 simple linear regression of the form,

1042

1043 $$\text{PredictabilityScore} = \text{CriticalDays}_{\text{Spring}} + \text{CriticalDays}_{\text{Fall}}$$

1044

and recorded model fit (R^2). We calculated the null distribution of R^2 values by permuting the
genome-wide predictability scores across among populations (n=10,000). For each permutation,
we recorded the maximum R^2 across the full range of thermal limits and used the distribution of
R^2 values as a null distribution. Using this null distribution, we then identified which thermal
limits are more strongly correlated with genome-wide predictability scores than expected by
chance (p_{perm}).

1051

1052 Predicted predictability scores were calculated using the best fit thermal limit models (p_{perm} <
1053 0.05). We calculated the median predicted score across the range of models. Confidence intervals
1054 were calculated using one standard deviation across the range of best fit models.

1055

1056 *Latitudinal cline concordance.* To identify SNPs that changed consistently in allele frequency
1057 with latitude (clinal), we first identified SNPs that varied in allele frequency along a 14.4°
1058 latitudinal transect up the east coast of the United States. We used one spring sample from each
1059 of the following populations (identified as "Region_City_Year"): PA_li_2011 (39.9°N),
1060 SC_eu_2010 (33.4°N), GA_ha_2008 (30.1°N), and FL_ho_2010 (25.5°N).

1061

1062 First, we regressed allele frequency with population latitude in a general linear model (*glm*: *R*
1063 v3.1.0), using a binomial error model and weights proportional to the effective coverage (N_c):

1064

1065 $$y_i = \text{latitude} + e_i$$

1066

where y_i is the allele frequency at the i^{th} SNP, and e^i is the binomial error given the i^{th} at the SNP.
To confirm the robustness of the clinal GLM results, clinality also was assessed using a second,
more complex model, that accounts for existing covariance in the dataset (bayenv2: Coop *et al.*
2010; Günther & Coop 2013). Results of the seasonal and clinal concordance analysis using the
bayenv2 model were comparable to that of the GLM model (Supplementary Figure 11) and are
not presented in the main text.
1073
We then tested the concordance in allele frequency change by season with the allele frequency
change by latitude. We performed three separate seasonal regressions (see above) for comparison
with the latitudinal regression: spring verses fall for the 18 non-Pennsylvania paired samples,
spring versus fall for the three California paired samples, and spring versus fall for the three
Europe paired samples. With the removal of the Pennsylvania samples, none of these three
seasonal regressions contained samples from any of the four populations used for the latitudinal
regression. Taking sets of increasingly clinal and increasingly seasonal SNPs, we assessed the
proportion of sites that both increase in frequency from spring to fall and increase in frequency
from north to south or that decrease in frequency from spring to fall and decrease in frequency
from north to south. We compared this concordance with the concordance of 100 sets of matched
controls.
1085
The strength and genomic extent of seasonal adaptation. We estimated the proportion of sites
under selection and the selection strength using the RFM in an ABC analysis (R package *abc*;
Csillery *et al.* 2012) using simulated datasets of neutral and selected sites. To simulate neutral
SNPs we used the observed spring allele frequency for a given SNP in a given paired sample as
the "true" allele frequency and performed a binomial draw of the size equal to the spring and of
the fall samples. This was done across all SNPs for each paired sample, producing a simulated,
null dataset the same size as the observed dataset, with the same allele frequency distributions as
the observed data. This simulated dataset will not necessarily reflect the same sampling (i.e.,
drift) process as observed in nature from spring to fall, and as neither the local census nor
variance effective population size are known, such processes cannot be accurately modeled.
However, as our rank *p*-value Fisher's method relies only on the rank within a paired sample,
and the consistency of the rank across paired samples, there is not likely to be an artificially
inflated seasonal signal due to incorrectly estimated sampling error (as there can be for a
regression *p*-value; Supplemental Figure 2).
1100
1101   Selected sites were drawn from the same spring allele frequencies and read depths as the neutral
1102   sites, with the exception that the fall allele frequency was drawn from a new allele frequency,
1103   following a specified shift from the observed spring frequency by a factor of *s*:
1104
$$AF_{fall} = AF_{spring} + AF_{spring} * (1\text{-}AF_{spring}) * s$$

where *AF* is the allele frequency and *s* is the cumulative "seasonal selection coefficient" from the
spring to the fall sample. We use this cumulative *s* since the true number of generations between
the spring and fall samples is not known. We simulated datasets with seasonal selection
coefficients ranging from 0 to 0.5 (in 0.01 increments), and proportions of sites under selection
ranging from 0 to 0.1 (in 0.001 increments). Each simulated dataset was then analyzed in the
same manner as the observed data, producing per-dataset distributions of X^2 test statistics. For

each X^2 bin of each dataset, we calculated the proportion of sites greater than expected. To avoid
double counting, we excluded the bottom half of the distribution ($X^2<20$), which consists of
negative observed minus expected values and is largely symmetric with the top half. We
calculated the square difference between the simulation and the observed proportion of sites per
bin, resulting in 28 summary statistics. We also calculated the sum of squared differences across
bins, which we refer to as the SSD. ABC parameter estimates were robust to using either all 28
per-bin summary statistics or the SSD, therefore we reduced our analysis to the single SSD
summary statistic. We used the rejection method in the *abc* function, choosing the tolerance level
of 0.01 based on lowest error in a cross-validation (*cv4abc* function; Supplemental Figure 14).
1122
1123   *Seasonal changes in inversion frequencies.* To test whether large cosmopolitan inversions
1124   change in frequency between spring and fall we calculated the average frequency of SNPs that
1125   are closely linked to inversion karyotype (Kapun *et al.* 2014).
1126
Genomic characteristics of seasonal SNPs. We assessed the uniformity of the genomic
distribution of highly seasonal SNPs by comparing the observed dataset to that of matched
controls and theoretical null distributions. For this analysis we examined the genomic
distribution of the top 1% of seasonally varying polymorphisms (n=7,748). We first measured
the number of seasonal SNPs per bin 100 SNP bin (average of 1 SNP per bin) and compared this
distribution to that of the mean of 100 sets of matched controls to the expectation from a
binomial distribution. A distribution that is more strongly peaked around 1 than expected is an
under-dispersed genomic distribution of seasonal SNPs (more evenly distributed than expected)
whereas a distribution less strongly peaked around 1 is over-dispersed and indicates clustering of
seasonal SNPs. We assessed the similarity of the observed and expected distributions using a
Kolmogorov-Smirnov test. We also assessed this distribution across genomic length scales. We
calculated the probability of observing at least one highly seasonal SNP among 1000 randomly
selected genomic windows ranging in size from 100bp to 100Kb. We compared this to the
expected probability given a Poisson distribution.
1141
1142   We calculated a "seasonal F_{st}" for each SNP by taking the median of the spring/fall F_{st} (Weir &
1143   Cockerham 1984, equations 1:4) values across paired samples.
1144
1145   Data availability. All raw sequence data have been deposited to SRA (BioProject Accession
1146   #PRJNA308584; individual accession numbers for each sample can be found in Supplemental
1147   Table 1), a VCF file with allele frequencies from all populations, scripts to conduct analyses
1148   presented here, and genome-wide SNP statistics for seasonality and clinality are available on
1149   DataDryad (doi:10.5061/dryad.4r7b826).
1150
1151
1152

1153   *Acknowledgements*
We thank the National Evolutionary Synthesis Center (NESCent) for sponsoring the 2012
Catalysis meeting that initiated the Drosophila Real Time Evolution Consortium. The meeting
was attended by Alan Bergland, Alisa Sedghifar, Brian Helmuth, Brian Lazzarro, Chau-Ti Ting,
David Kidd, Dmitri Petrov, Fabian Staubach, Hannah Burrack, Jim Fry, John Lessard, John
Coulbourne, John Pool, Josefa Gonzalez, Julien Ayroles, Kelly Dyer, Kim Hughes, Maaria

Kankare, Nadia Singh, Paul Schmidt, Regan Early, Stephen Porder, Subhash Rajpurohit, Sui Fai
Lee, and Thomas Flatt and we kindly thank all participants for their participation. We also thank
Jamie Blundell and all members of the Schmidt and Petrov labs who provided exceptionally
valuable feedback. This work was funded by the NIH grants R01GM100366 to PS and DAP,
R35GM118165 to DAP, R01GM100366 to PS, F32GM097837 and R35GM119686 to AOB,
European Commission grant H2020-ERC-2014-CoG-647900 to JG, a NSERC Discovery Grant
RGPIN-2018-05551 and Canada Research Chair 950-230113 to TJSM.

1166

Author contributions

Heather E. Machado: Designed the study, performed analysis, wrote manuscript, edited
manuscript, collected flies; Alan O. Bergland: Designed the study, performed analysis, wrote
manuscript, edited manuscript, collected flies; Ryan Taylor: Performed analysis, edited
manuscript; Susanne Tilk: Prepared sequencing libraries; Emily Behrman: Collected flies; Kelly
Dyer: Collected flies; Daniel K. Fabian: Collected flies; Thomas Flatt: Collected flies, edited
manuscript; Josefa González: Collected flies, edited manuscript; Talia L. Karasov: Collected
flies; Iryna Kozeretska: Collected flies; Brian P. Lazzaro: Collected flies, edited manuscript;
Thomas JS Merritt: Collected flies; John E. Pool: Collected flies, edited manuscript; Katherine
O'Brien: Collected flies; Subhash Rajpurohit: Collected flies; Paula R. Roy: Collected flies;
Stephen W. Schaeffer: Collected flies, edited manuscript; Svitlana Serga: Collected flies; Paul
Schmidt: Designed the study, wrote manuscript, edited manuscript, collected flies; Dmitri
Petrov: Designed the study, wrote manuscript, edited manuscript.

1180
1181

Supplemental Tables

1183
1184   Supplemental Table 1: Detailed information for each sample, including time and location of
1185   sample, collector, and SRA accession number.

1186
1187   Supplemental Table 2: Model coefficients for best-fit thermal limit models ($p_{perm} < 0.05$).

1188
1189   Supplemental Table 3: Critical days across a range of thermal limits for each population.

1190
1191   Supplemental Table 4: Regions of the genome with elevated density of seasonally variable SNPs
1192   for the original and flipped models

1193
1194   Supplemental Table 5: Results from Gene Ontology analysis of regions harboring an excess of
1195   seasonal SNPs using the flipped model.

1205
1206
1207

1208
1209   *References*
Ayrinhac A, Debat V, Gibert P *et al.* (2004) Cold adaptation in geographical populations of
Drosophila melanogaster: phenotypic plasticity is more important than genetic variability.
Functional Ecology, **18**, 700–706.
Adrion JR, Hahn MW, Cooper BS (2015) Revisiting classic clines in Drosophila melanogaster in
the age of genomics. *Trends in Genetics*, **31**, 434-444.
Barghi N, Tobler R, Nolte V, Jakšić AM, Mallard F, Otte KA, et al. (2019) Genetic redundancy
Fuels polygenic adaptation in *Drosophila*. *PloS Biology*, **17**, e3000128.
Barron MG, Fiston-Lavier AS, Petrov DA, Gonzalez J. (2014) Population genomics of
transposable elements in Drosophila. *Annu Rev Genet*, **48** , 561-81.
Behrman EL, Howick VM, Kapun M, Staubach F, Bergland AO, Petrov DA, Lazzaro BP,
Schmidt PS (2018) Rapid seasonal evolution in innate immunity of wild Drosophila
melanogaster. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20172599.
Behrman EL, Watson SS, O'Brien KR, Heschel MS, Schmidt PS (2015) Seasonal variation in
life history traits in two Drosophila species. *Journal of Evolutionary Biology*, **28**, 1691–
1704.
Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA (2014) Genomic evidence of
rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLoS
Genetics*, **10**, e1004775.
Bergland AO, Chae H-S, Kim Y-J, Tatar M (2012) Fine-scale mapping of natural variation in fly
fecundity identifies neuronal domain of expression and function of an aquaporin. *PLoS
Genetics*, **8**, e1002631.
Bergland AO, Genissel A, Nuzhdin SV, Tatar M (2008) Quantitative trait loci affecting
phenotypic plasticity and the allometric relationship of ovariole number and thorax length in
Drosophila melanogaster. *Genetics*, **180**, 567–582.
Bergland AO, Tobler R, González J, Schmidt P, Petrov D (2016) Secondary contact and local
adaptation contribute to genome-wide patterns of clinal variation in *Drosophila
melanogaster*. *Molecular Ecology*, **25**, 1157-1174.
Bertram J, Masel J (2019) Different mechanisms drive the maintenance of polymorphism at loci
subject to strong versus weak fluctuating selection. *Evolution,* **73***,* 883-896.
Betini GS, McAdam AG, Griswold CK, Norris DR (2017) A fitness trade-off between seasons
causes multigenerational cycles in phenotype and population size. *eLife*, **6**, e18770.
Botero CA, Weissing FJ, Wright J, Rubenstein DR (2015) Evolutionary tipping points in the
capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences
of the United States of America*, **112**, 184–189.
Boyle EA, Li YI, Pritchard JK. (2017) An expanded view of complex traits: from polygenic to
omnigenic. *Cell*, **169**, 1177–1186.
Bozicevic V, Hutter S, Stephan W, Wollstein A (2016) Population genetic evidence for cold
adaptation in European *Drosophila melanogaster* populations. *Molecular Ecology*, **25**,
1175–1191.
Campo D, Lehmann K, Fjeldsted C *et al.* (2013) Whole-genome sequencing of two North
American *Drosophila melanogaster* populations reveals genetic differentiation and positive
selection. *Molecular Ecology*, **22**, 5084–5097.
Carson HL, Stalker HD (1948) Reproductive Diapause in *Drosophila robusta*. *Proceedings of
the National Academy of Sciences of the United States of America*, **34**, 124–129.

Chakraborty M, Fry JD (2016) Evidence that Environmental Heterogeneity Maintains a Detoxifying Enzyme Polymorphism in *Drosophila melanogaster*. *Current Biology*, **26**, 219–223.

Charlesworth B. (2015) Causes of natural variation in fitness: Evidence from studies of Drosophila populations. *PNAS*, **112** ,1662-1669

Chen Y, Lee SF, Blanc E *et al.* (2012) Genome-Wide Transcription Analysis of Clinal Genetic Variation in Drosophila. *PloS one*, **7**, e34620.

Cockerell FE, Sgró CM, McKechnie SW (2014) Latitudinal clines in heat tolerance, protein synthesis rate and transcript level of a candidate gene in *Drosophila melanogaster*. *Journal of Insect Physiology*, **60**, 136–144.

Cogni R, Kuczynski C, Koury S *et al.* (2014) The intensity of selection acting on the *couch potato* gene -- spatial–temporal variation in a diapause cline. *Evolution*, **68**, 538–548.

Cogni R, Kuczynski K, Lavington E *et al.* (2015) Variation in *Drosophila melanogaster* central metabolic genes appears driven by natural selection both within and between populations. *Proceedings. Biological sciences / The Royal Society*, **282**, 20142688–20142688.

Comeron JM, Ratnappan R, Bailin S (2012) The many landscapes of recombination in *Drosophila melanogaster*. *PLoS Genetics*, **8**, e1002905.

Coop G, Witonsky D, Di Rienzo A, Pritchard J.K. (2010). Using Environmental Correlations to Identify Loci Underlying Local Adaptation. *Genetics*, **185**, 1411-1423.

Corbett-Detig RB, Hartl DL (2013) Correction: Population Genomics of Inversion Polymorphisms in *Drosophila melanogaster*. PLOS Genetics, 9, 10.1371

Csillery K, Francois O, Blum MGB. (2012), abc: an R package for approximate Bayesian computation (ABC). *Methods in Ecology and Evolution,* **3**, 475-479.

Curtsinger JW, Service PM, Prout T. (1994) Antagonistic Pleiotropy, Reversal of Dominance, and Genetic Polymorphism. *The American Naturalist,* **144**, 210-228.

David JR, Gibert P, Gravot E *et al.* (1997) Phenotypic plasticity and developmental temperature in Drosophila: Analysis and significance of reaction norms of morphometrical traits. *Journal of Thermal Biology*, **22**, 441–451.

Denlinger DL (2003) Regulation of diapause. *Annual Review of Entomology*, **47**, 93–122.

Dobzhansky T (1948) Genetics of natural populations; altitudinal and seasonal changes produced by natural selection in certain populations of *Drosophila persimilis*. *Genetics*, **33**, 158–176.

Ellner S (1996) Environmental fluctuations and the maintenance of genetic diversity in age or stage-structured populations. *Bulletin of Mathematical Biology*, **58**, 103–127.

Ellner S, Nelson G Hairston J (2015) Role of Overlapping Generations in Maintaining Genetic Variation in a Fluctuating Environment. *The American Naturalist*, **143**, 403–417.

Ellner S, Sasaki A (1996) Patterns of genetic polymorphism maintained by fluctuating selection with overlapping generations. *Theoretical Population Biology*, **50**, 31–65.

Ewing EP (1979) Genetic variation in a heterogeneous environment VII. Temporal and spatial heterogeneity in infinite populations. *American Naturalist*, **114**, 197–212.

Feder AF, Petrov DA, Bergland AO (2012) LDx: Estimation of Linkage Disequilibrium from High-Throughput Pooled Resequencing Data (R Wu, Ed,). *PloS one*, **7**, e48588.

Gould, SJ. (1989). *Wonderful life: the Burgess Shale and the nature of history.* W.W. Norton & Company, Inc. New York, NY.

Gillespie J (1973) Polymorphism in random environments. *Theoretical Population Biology*, **4**, 193–195.

Grenier JK, Arguello JR, Moreira MC *et al.* (2015) Global diversity lines - a five-continent
reference panel of sequenced *Drosophila melanogaster* strains. *G3:*
Genes|Genomes|Genetics, **5**, 593–603.
Günther T, Coop G. (2013) Robust identification of local adaptation from allele frequencies.
Genetics, **195**: 205-220.
Hedrick PW (1976) Genetic variation in a heterogeneous environment. II. Temporal
heterogeneity and directional selection. *Genetics*, **84**, 145–157.
Hedrick PW (2006) Genetic polymorphism in heterogeneous environments: The age of
genomics. *Annual Review of Ecology, Evolution, and Systematics*, **37**, 67–93.
Hendry AP. (2017) *Eco-evolutionary dyanmics*. Princeton University Press, Princeton, NJ.
Huang DW, Sherman BT, Lempicki RA (2009a) Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nature protocols*, **4**, 44–57.
Huang DW, Sherman BT, Lempicki RA (2009b) Bioinformatics enrichment tools: paths toward
the comprehensive functional analysis of large gene lists. *Nucleic Acids Research*, **37**, 1–13.
Ives PT (1970) Further genetic studies of the south Amherst population of *Drosophila*
melanogaster. *Evolution*, **24**, 507–518.
Juneja P, Quinn A, Jiggins FM (2016) Latitudinal clines in gene expression and cis-regulatory
element variation in *Drosophila melanogaster*. *BMC Genomics*, **17**, 981.
Kain JS, Zhang S, Zade JA *et al.* (2015) Variability in thermal and phototactic preferences in
Drosophila may reflect an adaptive bet-hedging strategy. *Evolution*, **69**, 3171–3185.
Kapopoulou A, Pfeifer SP, Jensen JD, Laurent S. (2018) The demographic history of African
Drosophila melanogaster. *Genome Biology and Evolution,* **10**, 2338-2342.
Kapun M, Flatt T (2019) The adaptive significance of chromosomal inversion polymorphisms in
Drosophila melanogaster. *Molecular Ecology*, **28**, 1263-1282.
Kapun M, Aduriz MGB, Staubach F *et al.* (2018) Genomic analysis of European *Drosophila*
melanogaster populations on a dense spatial scale reveals longitudinal population structure
and continent-wide selection. *bioRxiv*, 313759.
Kapun M, Fabian DK, Goudet J, Flatt T (2016) Genomic Evidence for Adaptive Inversion Clines
in *Drosophila melanogaster*. *Molecular Biology and Evolution*, **33**, 1317–1336.
Kapun M, van Schalkwyk H, McAllister B, Flatt T, Schlötterer C (2014) Inference of
chromosomal inversion dynamics from Pool-Seq data in natural and laboratory populations
of *Drosophila melanogaster*. *Molecular Ecology*, **23**, 1813–1827.
Karasov T, Messer P, Petrov DA (2010) Evidence that Adaptation in *Drosophila* Is Not Limited
by Mutation at Single Sites. *PLoS Genetics*, **6**, e1000924.
Keller A. (2007) *Drosophila melanogaster*'s history as a human commensal. *Current Biology*,
17, R77-81.
Kim YB, Oh JH, McIver LJ *et al.* (2014) Divergence of *Drosophila melanogaster* repeatomes in
response to a sharp microclimate contrast in Evolution Canyon, Israel. *Proceedings of the*
National Academy of Sciences of the United States of America, **111**, 201410372–10635.
Klepsatel P, Gáliková M, De Maio N *et al.* (2013) Reproductive and post-reproductive life
history of wild-caught *Drosophila melanogaster* under laboratory conditions. *Journal of*
Evolutionary Biology, **26**, 1508–1520.
Koboldt DC, Zhang Q, Larson DE *et al.* (2012) VarScan 2: somatic mutation and copy number
alteration discovery in cancer by exome sequencing. *Genome Research*, **22**, 568–576.

Kolaczkowski B, Kern AD, Holloway AK, Begun DJ (2011) Genomic differentiation between temperate and tropical Australian populations of *Drosophila melanogaster*. *Genetics*, **187**, 245–260.

Kostál V (2006) Eco-physiological phases of insect diapause. *Journal of Insect Physiology*, **52**, 113–127.

Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE (2016) A Thousand Fly Genomes: An Expanded Drosophila Genome Nexus. *Molecular Biology and Evolution*, **33**, 3308–3313.

Levene H (1953) Genetic equilibrium when more than one ecological niche is available. *The American Naturalist*, **87**, 331–333.

Levine MT, Eckert ML, Begun DJ (2011) Whole-Genome Expression Plasticity across Tropical and Temperate *Drosophila melanogaster* Populations from Eastern Australia. *Molecular Biology and Evolution*, **28**, 249–256.

Levins R (1968) *Evolution in changing environments: some theoretical explorations*. Princeton University Press, Princeton, NJ.

Lynch M, Bost D, Wilson S, Maruki T, Harrision S (2014) Population-genetic inference from pooled-sequencing data. *Genome Biology and Evolution,* **6**, 1210-1218.

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, **25**, 1754–1760.

Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

Machado HE, Bergland AO, O'Brien KR *et al.* (2016) Comparative population genomics of latitudinal variation in *Drosophila simulans* and *Drosophila melanogaster*. *Molecular Ecology*, **25**, 723–740.

Mackay TFC, Huang W (2018) Charting the genotype-phenotype map: lessons from the *Drosophila melanogaster* Genetic Reference Panel. *WIREs Dev Biol,* **7**, e289.

Mackay TFC, Richards S, Stone EA *et al.* (2012) The *Drosophila melanogaster* Genetic Reference Panel. *Nature*, **482**, 173–178.

MacMillan HA, Ferguson LV, Nicolai A *et al.* (2015) Parallel ionoregulatory adjustments underlie phenotypic plasticity and evolution of Drosophila cold tolerance. *Journal of Experimental Biology*, **218**, 423–432.

Mallard F, Nolte V, Tobler R, Kapun M, Schlötterer C (2018) A simple genetic basis of adaptation to a novel thermal environment results in complex metabolic rewiring in Drosophila. *Genome Biology*, **19**, 119.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, **17**, pp. 10–12.

McKenna A, Hanna M, Banks E *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, **20**, 1297–1303.

McMillan I, Fitz-Earle M, Robson DS (1970) Quantitative genetics of fertility. II. Lifetime egg production of *Drosophila melanogaster*--experimental. *Genetics*, **65**, 355–369.

Noh S, Everman ER, Berger CM, Morgan TJ (2017) Seasonal variation in basal and plastic cold tolerance: Adaptation is influenced by both long- and short-term phenotypic plasticity. *Ecology and Evolution*, **7**, 5248–5257.

Oakeshott JG, Gibson JB, Anderson PR, Knibb WR, Anderson DG, Chambers GK. (1982).
Alcohol Dehydrogenase and Glycerol-3-Phosphate Dehydrogenase Clines in *Drosophila*
melanogaster on Different Continents. *Evolution*, **36**, 86-96
Overgaard J, Kristensen TN, Mitchell KA, Hoffmann AA (2015) Thermal Tolerance in
Widespread and Tropical Drosophila Species: Does Phenotypic Plasticity Increase with
Latitude? *The American Naturalist*, **178**, S80–S96.
Paaby AB, Bergland AO, Behrman EL, Schmidt PS (2014) A highly pleiotropic amino acid
polymorphism in the Drosophila insulin receptor contributes to life-history adaptation.
Evolution, **68**, 3395–3409.
Pavlidis P, Jensen JD, Stephan W, Stamatakis A (2012) A Critical Assessment of Storytelling:
Gene Ontology Categories and the Importance of Validating Genomic Scans. *Molecular
Biology and Evolution*, **29**, 3237–3248.
Pool JE, Corbett-Detig RB, Sugino RP *et al.* (2012) Population genomics of Sub-Saharan
Drosophila melanogaster: African diversity and Non-African admixture. *PLoS Genetics*, **8**,
e1003080.
R Core Team. (2019) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Rajpurohit S, Gefen E, Bergland AO *et al.* (2018) Spatiotemporal dynamics and genome-wide
association analysis of desiccation tolerance in *Drosophila melanogaster*. *Molecular
Ecology*, **27**, 3525-3540.
Rajpurohit S, Hanus R, Vrkoslav V *et al.* (2017) Adaptive dynamics of cuticular hydrocarbons in
Drosophila. *Journal of Evolutionary Biology*, **30**, 66–80.
Rajpurohit S, Nedved O, Gibbs AG (2013) Meta-analysis of geographical clines in desiccation
tolerance of Indian drosophilids. *Comparative Biochemistry and Physiology a-Molecular &
Integrative Physiology*, **164**, 391–398.
Rand DM, Weinreich DM, Lerman D, Folk D, Gilchrist GW (2010) Three Selections Are Better
Than One: Clinal Variation of Thermal QTL From Independent Selection Experiments in
Drosophila. *Evolution*, **64**, 2921–2934.
Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD (2014) Parallel geographic
variation in *Drosophila melanogaster*. *Genetics*, **197**, 361–373.
Rhomberg, L.R. and R.S. Singh. (1989) Evidence for a link between local and seasonal cycles in
gene frequencies and latitudinal clines in a cyclic parthenogen. *Genetica*, **78**, 73-79.
Rodríguez-Trelles F, Tarrío R, Santos M (2013) Genome-wide evolutionary response to a heat
wave in Drosophila. *Biology Letters*, **9**, 20130228.
Rudman SM, Barbour M, Csillery K, Gienapp P, Guillaume F, Hairston N, Hendry A, Lasky JR,
Rafajlović M, Räsänen K, Schmidt PS, Seehausen O, Therkildsen NO, Turcotte MM, Levine
J. (2017). What genomic data can reveal about eco-evolutionary dynamics. *Nature Ecology
& Evolution, **2**, 9-15.
Scheiner SM (1993) Genetics and Evolution of Phenotypic Plasticity. *Annual Review of Ecology
and Systematics, **24**, 35-68.
Schlötterer C, Tobler R, Kofler R, Nolte V (2014) Sequencing pools of individuals — mining
genome-wide polymorphism data without big funding. *Nature Reviews Genetics*, **15**, 749–
763.
Schmidt PS, Conde DR (2006) Environmental heterogeneity and the maintenance of genetic
variation for reproductive diapause in *Drosophila melanogaster*. *Evolution*, **60**, 1602-1611.

Schmidt PS, Matzkin L, Ippolito M, Eanes WF (2005) Geographic variation in diapause incidence, life-history traits, and climatic adaptation in *Drosophila melanogaster. Evolution*, **59**, 1721–1732.

Schmidt PS, Zhu C-T, Das J *et al.* (2008) An amino acid polymorphism in the couch potato gene forms the basis for climatic adaptation in *Drosophila melanogaster. Proceedings of the National Academy of Sciences of the United States of America*, **105**, 16207–16211.

Signor SA, New FN, Nuzhdin S. A large panel of *Drosophila simulans* reveals an abundance of common variants. *Genome Biology and Evolution*, **10**, 189–206.

Spitzer K, Pelizzola M, Futschik A. (2019). Modifying the Chi-square and the CMH test for population genetic inference: adapting to over-dispersion. http://arxiv.org/abs/1902.08127.

Stalker HD. (1976). Chromosome studies in wild populations of *Drosophila melanogaster. Genetics,* **82**: 323-347.

Stalker HD (1980). Chromosome Studies in Wild Populations of *Drosophila melanogaster*. II. Relationship of Inversion Frequencies to Latitude, Season, Wing-Loading and Flight Activity. *Genetics*, **95**, 211-223.

Svetec N, Cridland JM, Zhao L, Begun DJ (2016) The Adaptive Significance of Natural Genetic Variation in the DNA Damage Response of *Drosophila melanogaster. PLoS Genetics*, **12**, e1005869.

Svetec N, Zhao L, Saelao P, Chiu JC, Begun DJ (2015) Evidence that natural selection maintains genetic variation for sleep in *Drosophila melanogaster. BMC evolutionary biology*, **15**, 41.

Tauber MJ, Tauber CA, Masaki S (1986) *Seasonal Adaptations of Insects*. Oxford University Press, Oxford, UK.

Tobler R, Franssen SU, Kofler R *et al.* (2014) Massive habitat-specific genomic response in *D. melanogaster* populations during experimental evolution in hot and cold environments. *Molecular Biology and Evolution*, **31**, 364–375.

Vesala L, Salminen TS, Kostál V, Zahradnickova H, Hoikkala A (2012) Myo-inositol as a main metabolite in overwintering flies: seasonal metabolomic profiles and cold stress tolerance in a northern drosophilid fly. *Journal of Experimental Biology*, **215**, 2891–2897.

Via S, Lande R (1985) Genotype-environment interaction and the evolution of phenotypic plasticity. *Evolution*, **39**, 505–522.

Weir BS, Cockerham CC. (1984) Estimating F-Statistics for the Analysis of Population Structure. *Evolution*, **38**, 1358-1370

Wiberg RAW, Gaggiotti OE, Morrissey MB, Ritchie MG (2017) Identifying consistent allele frequency differences in studies of stratified populations. (L Johnson, Ed,). *Methods in Ecology and Evolution*, **8**, 1899–1909.

Wittmann MJ, Bergland AO, Feldman MW, Schmidt PS, Petrov DA (2017) Seasonally fluctuating selection can maintain polymorphism at many loci via segregation lift. *Proceedings of the National Academy of Sciences of the United States of America*, **114**, E9932–E9941.

Zhao L, Begun DJ (2017) Genomics of parallel adaptation at two timescales in Drosophila. *PLoS Genetics*, **13**, e1007016.

Zhu Y, Bergland AO, González J, Petrov DA (2012) Empirical Validation of Pooled Whole Genome Population Re-Sequencing in Drosophila melanogaster. *PloS one*, **7**.

1476 Supplemental Figures
1477

1478
1479
1480 **Supplemental Figure 1**. Sampling localities in A) North America and B) Europe for all samples
1481 collected, regardless of use in the present analysis. Sizes of circles reflect number of samples per
1482 locality.

1495
Supplemental Figure 2. The effect of artificial increases of sample size on the Core20 seasonal
analysis using A,C) the rank Fisher's method and B,D) the GLM analyses. Top row: sample size
inflated by a fixed amount. Bottom row: sample size inflated by a Poisson sampled amount.
Shown is the proportion of SNPs in each bin (solid lines) compared with the null expectation
(dashed line). Black: original dataset. Dark grey: 10 reads added to each sample (same allele
frequencies). Light grey: 100 reads added to each sample (same allele frequencies).

1508
1509 **Supplemental Figure 3**. The enrichment of shared seasonal sites from the leave-one-out
1510 analysis of the Core20 populations. Each line represents one population, colored by the inferred
1511 genome-wide predictability score. (A) Enrichment scores for all populations; (B) Enrichment
1512 scores for populations with significant enrichment (nominal $p < 0.05$) at a joint significance
1513 threshold less than 1% (-2).
1514
1515
1516
1517

1518
1519 **Supplemental Figure 4**. Predictability of seasonal adaptation, by chromosome. The fraction of
1520 concordant allele frequency changes from the leave-one-out cross validation analysis of the
1521 Core20 populations. Each line represents one population, colored by the inferred genome-wide
1522 predictability score.
1523

1524
1525
1526   Supplemental Figure 5. Correlation between genome-wide predictability scores and per-
1527   chromosome predictability scores.
1528

Supplemental Figure 6. Additional plots related to the predictability analysis. A) Pearson's correlation coefficient between the ValidationSet's observed genome-wide concordance score and their predicted scores for the best-fit permuted models (null) versus observed scores generated from the original Core20 model versus the flipped Core20 model. B) Genome-wide concordance scores from the ValidationSet of populations in relation to the flipped Core20 model. C) The best thermal limit models from the Core20 analysis predict genome-wide concordance scores of the ValidationSet (diamonds).

1542

1543
1544 **Supplemental Figure 7**. Comparison of the general linear model (GLM: main text) and the
1545 bayenv model for assessing clinal and seasonal concordance. The joint quantile is the quantile
1546 threshold for both the seasonal and the clinal analysis (plot is cumulative).

1557
Supplemental Figure 8. The enrichment of SNPs found to be both seasonal and latitudinal,
regardless of concordance, as a function of *p*-value quantile thresholds. Darker hues indicate a
greater proportion of concordant SNPs. The percent enrichment is calculated as the percent
increase in SNPs at a joint quantile threshold over matched controls. A,C) Original dataset. B,D)
Flipped dataset. A,B) All values, regardless of a significant enrichment. C,D) Only values greater
than 95% of controls (otherwise, colored white).

1574

1575

1576 **Supplemental Figure 9.** ABC estimates for the proportion of sites affected by seasonal selection
1577 and the associated *s* for the original model *without* filtering sites found to be monomorphic in
1578 one or more populations. Plotted is the sum of squared distances (SSD) between the observed
1579 and simulation RFM X^2 test statistic distributions, for varying strengths of selection and
1580 proportion of sites under selection.
1581

1582

Supplemental Figure 10. Effect of subsampling on ABC parameter estimates. Top row: original dataset. Bottom row: flipped dataset. A,D) No subsampling. B,E) Sampling one SNP per 1Kb. C,F) Sampling one SNP per 5Kb.

1601

1602 **Supplemental Figure 11**. Inversion frequencies. Each datapoint is a separate population.

1603 Numbers are the fraction of populations where spring is greater than fall. A) Original dataset. B)

1604 Flipped dataset.

1605

1609 **Supplemental Figure 12**. Enrichment of genic classes for A) the top 1% and B) top 5% of
1610 seasonal sites for the original analysis, and C) the top 1% and D) top 5% for the flipped analysis.
1611 Points represent means and error bars represent 95% CI across100 bootstrapped sets of matched
1612 control SNPs.
1613

1614
1615
1616

1617
1618
Supplemental Figure 13. Competitive mapping effectively reduces cross species mapping
between *D. simulans* and *D. melanogaster*. (A) The estimated rate of *D. simulans* contamination
can be precisely estimated by calculating the number of reads mapping preferentially to the *D.*
simulans genome. Estimated contamination rates are ~2% lower than the expected rate. (B) The
residual cross species mapping rate, i.e. the fraction of *D. simulans* reads mapping to the *D.*
melanogaster genome is roughly consistent at ~9%.
1625

1626

1627
1628 **Supplemental Figure 14**. Distribution of the summary statistic SSD used in the ABC parameter
1629 estimation. Red dashed line indicates the cutoff for the top 1% of simulations (corresponds to the
1630 ABC tolerance level). A) Original dataset. B) Flipped dataset.