Dynamic decoupling of biomass and lipid biosynthesis by

autonomously regulated switch

short title: Dynamic decoupling of biomass and lipid synthesis

Suvi Santala[1]*, Elena Efimova[1], Ville Santala[1]

[1]Laboratory of Chemistry and Bioengineering, Tampere University of Technology,

Korkeakoulunkatu 8, FI-33720, Tampere, Finland

*Corresponding author

Email addresses:

SS: suvi.santala@tut.fi

EE: elena.efimova@tut.fi

VS: ville.santala@tut.fi

Abstract

For improving the microbial production of fuels and chemicals, gene knock-outs and overexpression are routinely applied to intensify the carbon flow from substrate to product. However, their possibilities in dynamic control of the flux between the biomass and product synthesis are limited, whereas dynamic metabolic switches can be used for optimizing the distribution of carbon and resources. The production of single cell oils is especially challenging, as the synthesis is strongly regulated, competes directly with biomass, and requires defined conditions, such as nitrogen limitation. Here, we engineered a metabolic switch for redirecting carbon flow from biomass to wax ester production in *Acinetobacter baylyi* ADP1 using acetate as a carbon source. Isocitrate lyase, an essential enzyme for growth on acetate, was expressed under an arabinose inducible promoter. The autonomous downregulation of the expression is based on the gradual oxidation of the arabinose inducer by a glucose dehydrogenase *gcd*. The depletion of the inducer, occurring simultaneously to acetate consumption, switches the cells from a biomass mode to a lipid synthesis mode, enabling the efficient channelling of carbon to wax esters in a simple batch culture. In the engineered strain, the yield and titer of wax esters were improved by 3.8 and 3.1 folds, respectively, over the control strain. In addition, the engineered strain accumulated wax esters 19% of cell dry weight, being the highest reported among microbes. The study provides important insights into the dynamic engineering of the biomass-dependent synthesis pathways for the improved production of biocompounds from low-cost, sustainable substrates.

Keywords

Lipid biosynthesis, wax esters, acetate, dynamic control, decoupling, autonomous

circuit

Abbreviations

WE – wax esters, AceA – isocitrate lyase

1. Background

Metabolic engineering and synthetic biology provide powerful means for the bio-based production of a variety of chemicals and other commodities by engineered microbes. Compared to conventional chemical synthesis, the superiority of biological production systems lies in the possibility to synthesize both the catalyst (i.e. the cell factory) and the product itself from very simple chemical compounds, such as sugars or organic acids. However, the challenge is to develop a system, which optimally distributes the resources and carbon flux between building up the catalyst and operating the catalyst for the actual production; extensive cell growth takes resources from the product synthesis, whereas too excessive flux towards product synthesis may result in reduced growth, insufficient cofactor regeneration, low enzyme expression levels, and eventually poor titers. To address the challenges related to the optimal distribution of cellular resources, a number of dynamic circuit designs targeting the central pathway nodes have been recently developed (1). While growth-associated genes responsible for central carbon metabolism cannot be directly deleted, various strategies for decoupling growth and product synthesis have been introduced; Soma et al. constructed a metabolic toggle switch for conditional knockout of citrate

synthase *gltA,* an enzyme required for functional tricarboxylic acid (TCA) cycle (2). The

switch allowed an induced shift of carbon flow from TCA cycle to synthetic isopropanol

pathway. More recently, the system was further improved by introducing a sensor-

regulator system responsive to a defined cell density (3). Solomon et al. introduced a

dynamic approach to controlling the glycolytic flux; antisense RNA technology and an

inverting gene circuit were employed for inhibiting the activity of glucokinase (Glk),

resulting in a controlled growth rate and a reduced production of acetate (4).

Brockman and Prather introduced another example of a dynamic regulation system,

where they developed a circuit for dynamic knockdown of phosphofructokinase-1

(Pfk-1), the enzyme responsible for the key step in the glycolytic pathway regulating

glucose-6-phosphate flux. By the temporal control of Pfk-1 degradation, glucose-6-

phosphate could be efficiently directed to heterologous *myo*-inositol synthesis

pathway instead of biomass production (5). In a previous work, Doong et al. further

improved the system with a *myo*-inositol responsive dynamic sensor that regulated

the downstream enzymes of the pathway in converting *myo*-inositol to glucarate. The

introduced systems represent highly elegant examples of advanced metabolic control,

but the complex circuit designs can be prone to destabilization in prolonged

cultivations and function unexpectedly in scaled-up processes (6). Thus, reliable

metabolic control systems with simple and robust operation would be also desired.

Microbial storage compounds, such as polyhydroxyalkanoates (PHA), triacylglycerols

(TAG) and wax esters (WE), are industrially relevant and desirable molecules due to

their vegetable oil like properties and broad applicability in e.g. fuel, nutritional,

cosmetic, and pharmaceutical industries. However, the production of the long carbon

chain products derived from fatty acyl-CoA requires significant energy investments

from the cell. In addition, the synthesis of storage compounds directly competes with

biomass production and is strongly regulated, growth-phase dependent, and requires

high amounts of cofactors and excess carbon along with limitation on other nutrients,

such as nitrogen (7, 8). While strategies for dynamically regulated production of free

fatty acids (FFA) and FA derived products have been introduced (9, 10), means for

overcoming the challenges of storage lipid synthesis regulation are still lacking.

Therefore, the production processes are conventionally improved by non-specific

means, such as bioprocess optimization or in conditions with a defined carbon-

nitrogen ratio (11). As an example of a more advanced approach, Xu et al. established

a semi-continuous culture system with model-aided bioprocess optimization and cell

recycling, which allowed efficient TAG production and high productivities even with

dilute acetate feed (12). While also metabolic engineering strategies have been

employed for improving TAG (13-16) and WE synthesis (17, 18) in microbes, efficient

overproduction of acetyl-CoA coupled with dynamic resource distribution between

biomass and storage lipid synthesis remains a challenge.

When microbes are cultivated on non-glycolytic substrates, such as acetate or fatty

acids, complete TCA cycle yielding carbon dioxide cannot be utilized for anabolic

processes. In those circumstances, the cells have to rely on an alternative route,

namely glyoxylate cycle, which bypasses the two decarboxylation steps of TCA cycle.

The key enzyme in the glyoxylate cycle is isocitrate lyase (AceA), responsible for

converting isocitrate to glyoxylate and succinate. It was previously demonstrated that

a knock-out of the isocitrate lyase in *Pseudomonas putida* improved the production of

PHAs when grown on gluconate, apparently for providing surplus acetyl-CoA for the

PHA synthesis (19). Some bacteria also exhibit an alternative pathway for glycolysis,

such as the modified Entner-Doudoroff pathway of certain *Acinetobacter* strains (20,

21). An interesting feature of the glycolysis of *Acinetobacter* is the oxidation of glucose

to gluconate prior the transport to cells. In the absence of glucose, the first enzyme of

the pathway, glucose dehydrogenase Gcd, can unselectively oxidize other sugar

compounds present in the medium, such as pentoses, without the capability to utilize

them as a carbon source (22-24). Importantly, this feature does not interfere with the

utilization of non-glycolytic carbon sources, such as organic acids, and can be

considered being 'orthogonal' to the substrate utilization. Thus, this feature could be

exploited in the regulation of metabolic pathways in the cells.

Here, we construct a dynamic switch for autonomous shifting of cells from the

biomass mode to the storage lipid synthesis mode by introducing a circuit for a

conditional elimination of the glyoxylate cycle, which is the essential bypass for cells

growing on acetate, and the key control node in lipid biosynthesis pathway. We

demonstrate the functionality of the switch by the improved (both yield and titer)

production of wax esters using *A. baylyi* ADP1 as a host.

2. Material and Methods

Strains and molecular work

A. baylyi ADP1 (DSM 24193, Deutsche Sammlung von Microorganismen und Zellkulturen, Germany) was used in the study. The single gene knock-out strain of *A. baylyi* ADP1Δ*aceA*::*tdk*/*Kan*[r] (ACIAD1084 deleted) was kindly provided by Veronique de Berardinis (Genoscope, France). In the single gene knock-out mutant, the gene in question is replaced with a gene cassette containing a kanamycin resistance gene (Kan[r]) (25). For cloning and plasmid amplification, *Escherichia coli* XL1-Blue (Stratagene, USA) was used. The reagents and primers for molecular work were purchased from ThermoFisher Scientific (USA), unless stated otherwise. The primers used in the study are presented in Table 1. The complementation of *aceA* in the knock-out strain was carried out by constructing a gene cassette with the *aceA* gene under an arabinose inducible promoter AraC-pBAD; a previously described gene cassette (16, 26) was used as the scaffold for the construction. The arabinose promoter and AraC repressor, designated as ara, were amplified from pBAV1C-ara-LuxCDE plasmid (18) using primers VS10_09 and VS10_10 and inserted to the plasmid iluxAB/pIX (26) using restriction sites MfeI and NdeI. The gene *aceA* (ACIAD1084) was amplified from the genome of ADP1 wt with primers SS17_08 and SS17_09 and cloned to the ara-iluxAB/pIX using restriction sites NdeI and XhoI. Transformations of ADP1 were carried out as described previously (16). The transformed cells were selected on LA plates containing 25 µg/ml chloramphenicol. ADP1 strain with a *poxB* deletion, *A. baylyi* ADP1Δ*poxB*::*Cm*[r], was used as the reference strain designated as ADP1 wt. All genetic modifications were confirmed with PCR and sequencing.

Cultivations

The components for culture mediums were purchased from Sigma (USA). The strains were cultivated in modified MA/9 minimal salts medium (17) at 25 °C and 300 rpm unless stated otherwise. The medium was supplemented with 25-100 mM Na-acetate or 250 mM glucose, 0.1 % casein amino acids (w/v), and 0-1.0 % L(+)arabinose when appropriate.

The cultivations for optimizing the arabinose concentration for ADP1-ara-aceA growth were carried out using Tecan Spark® (Tecan, Switzerland) microplate reader in 25 °C for 21 hours with two replicate wells for each strain and arabinose concentration. The medium was supplemented with 25 mM Na-acetate and 0.1 % casein amino acids, and arabinose (concentrations 0; 0.05; 0.1; 0.2; 0.5 or 1.0 %) was used for the induction of *aceA*. Mediums without acetate supplementation was used as the control medium to determine the growth on casein amino acids. The strain ADP1 wt was used as the positive control, whereas the knock-out strains ADP1 Δ*aceA*::*Tdk/kan*[r] and *A. baylyi* ADP1Δ*aceA*::*tdk/Kan*[r] Δ*poxB*::*Cm*[r] were used as negative controls for growth on acetate. For semi-quantitative determination of the WE production of ADP1-ara-aceA, the cells were cultivated in two parallel 5 ml cultures in the same medium except with 50 mM Na-acetate in 5 ml tubes for 30 hours.

For the quantitative determination of the WEs with NMR, the strain ADP1 wt and ADP1-ara-aceA were cultured in total 600 ml of medium supplemented with 50 mM Na-acetate and 0.5 % (~30 mM) arabinose distributed in 12 Erlenmeyer flasks. HPLC samples were taken every 1-5 hours and two parallel 40 ml samples were taken in five

different time-points to quantitatively determine WE (by NMR) and biomass

production (as cell dry weight).

Lipid and end-metabolite analyses

The amount of total lipids and WEs were estimated by TLC or quantified by NMR. For

TLC, equal volumes of samples (3 ml) from different cultures were taken and the lipids

were extracted using 'miniscale' chloroform-methanol extraction as described

previously (17). Thirty μl of the chloroform phase was applied on 20 × 10 cm Silica Gel

60 F_{254} HPTLC glass plates with 2.5 × 10 cm concentrating zone (Merck, USA). Mobile

phase used was n-hexane: diethyl ether: acetic acid 90: 15: 1 and iodine was used for

visualization. Jojoba oil was used as the standard for WEs. For comparative evaluation

of the intensities of the WE bands on TLC, the Gel analysis method of ImageJ software

(rsb.info.nih.gov/ij/index.html) was applied as described in the ImageJ

documentation.

For NMR analyses, the 40-ml biomass samples were freeze-dried and the cell dry

weight (CDW) was determined gravimetrically. The lipid extraction and the

quantitative [1]H NMR analysis of WEs was carried out as described earlier (26). The

amount of total lipids was determined gravimetrically. The areas of the peaks in the

NMR spectrum are directly proportional to the molar concentration of each functional

group, yielding specific concentration for WEs in total biomass. The concentration of

WEs was calculated from the integrated signal at 4.05 ppm which is characteristic for

protons of α-alkoxy-methylene group of esters (–CH2-COO-CH2–). For the calculation

of the WE titer in grams per liter, an average molar mass of 506 g/mol was used (17).

The glucose, acetate, and arabinose concentrations were determined by LC-20AC

prominence liquid chromatograph (Shimadzu, USA) equipped with RID-10A refractive

index detector, DGU-20A5 prominence degasser, CBM-20A prominence

communications bus module, SIL-20AC prominence autosampler, and Shodex SUGAR

SH1011 (Showa Denko KK, Japan) as described previously (16).

204　3. Results

In order to investigate the effect of the knock-out of isocitrate lyase AceA on the

growth and WE production in *A. baylyi* ADP1, we employed a knock-out mutant strain

A. baylyi ADP1 Δ*aceA*::Tdk/kanr (25) for preliminary test cultivations. We observed

that when grown on glucose, the cells grow more slowly, but produce WE titers

comparable to those of the wild type (wt) strain; after 48 hours of cultivation, the wild

type had produced 470±150 mg/l WEs compared to 460±40 mg/l WEs produced by

the knock-out strain. In opposite to the wt strain, however, the mutant strain did not

exhibit growth on minimal medium supplied with acetate as the sole carbon source.

This is due to the lack of route for acetyl-CoA to be directed in biosynthetic pathways

via malate. Thus, as acetyl-CoA represents the key precursor in both the biomass

production through the glyoxylate shunt and the wax ester biosynthesis, we

hypothesized that by dynamically regulating the isocitrate lyase, the state of the cells

could be switched between biomass and lipid synthesis modes (Fig 1). In order to make

the shift dynamic, we introduced an approach for autonomous regulation of the

isocitrate lyase AceA; by expressing the enzyme under an arabinose-inducible

promoter AraC-pBAD, the induction is gradually repressed due to the depletion of

arabinose by the glucose dehydrogenase activity of ADP1. The arabinose inducible

promoter has been previously shown to function in *A. baylyi* ADP1 (18, 27). In order

to establish a system with maximal linearity and controllability, we constructed a gene

cassette for genomic expression of *aceA* (Fig 1c, d). Exploiting the natural

transformation machinery of ADP1, the gene cassette was integrated in the genome

to replace a gene *poxB* (ACIAD3381), which has been previously shown to be a neutral

target site in terms of growth and wax ester production (26, 28). The resulting strain

A. baylyi ADP1Δ*aceA*::*tdk*/*Kan*[r] Δ*poxB*::*araC*-pBAD-*aceA*-*Cm*[r] was designated as ADP1-

ara-aceA. The strain *A. baylyi* ADP1Δ*poxB*::*Cm*[r] was used as the reference strain, from

now on designated as the ADP1 wt.

We investigated the functionality of the complementation in the strain ADP1-ara-aceA

in minimal medium supplemented with 50 mM acetate (Figure 2). Arabinose (at

concentrations 0; 0.1 and 1.0 %) was added to the cultures, and the cultivations were

continued for 68 hours. Cells did not grow or consume acetate in the absence of

arabinose, indicating sufficiently tight regulation of the arabinose promoter. In the

cultures with small amount of arabinose (0.1 %), the cells stopped growing after

reaching an optical density (OD) of 0.3 and consumed only 5 mM acetate, whereas

with 1.0 % arabinose the cells reached OD ~3.5 along with complete consumption of

acetate. The ADP1 wt grew to slightly lower biomass (OD ~2.4) and consumed all the

acetate; arabinose supplementation had no effect on the ADP1 wt growth. After 68

hours of cultivation, approximately 85 % of the 1% arabinose had been oxidized. The

control knockout strains ADP1Δ*aceA*::*tdk*/*Kan*[r] and ADP1Δ*aceA*::*tdk*/*Kan*[r] Δ*poxB*::*Cm*[r]

did not exhibit growth nor acetate consumption with or without the presence of

arabinose (OD 0 at 0-68 h). We also confirmed, that the AceA expression is repressed

due to arabinose oxidation, i.e. the conversion of arabinose to non-inducive form,

arabino-lactone and further to arabonate (Figure S1).

In order to find the optimal arabinose concentration in terms of both biomass and wax

ester production, the strain ADP1-ara-aceA was cultivated in several different

arabinose concentrations in minimal salts medium supplemented with acetate (Figure

3) for 21 hours. ADP1 wt was cultured as the reference strain. Casein amino acids

(0.1%) were added to the culture in order to promote the growth and to prevent

nitrogen limitation. As indicated by the previous growth experiment, we found that

1% arabinose was sufficient to allow the engineered strain to reach the same biomass

as ADP1 wt, albeit the cells grew slower. Within the concentration range 0 – 0.2%, only

small differences in growth pattern or biomass production were observed. The slight

increase in OD of uninduced cells is due to the utilization of the casein amino acids

present in the growth medium; a same amount of biomass is achieved without acetate

supplementation   with   the   wild   type   strain   and   the   knock-out   strain

ADP1ΔaceA::tdk/Kan$^r\Delta$poxB::Cm^r with   both   acetate   and   casein   amino   acid

supplementation (data not shown). For ADP1 wt, all the growth curves were similar

regardless of the arabinose concentration used (data not shown).

Next, we determined which initial arabinose concentration most optimally distributes

the carbon between the biomass and the WE production. The strain ADP1-ara-aceA

was cultivated in 5 ml minimal salts medium supplemented with 0.1 % casamino acids

and 50 mM acetate with different arabinose concentrations (0; 0.1; 0.2; 0.5% and

1.0%) for 30 hours, after which biomass and WE production were determined. A clear

correlation between the arabinose concentration and biomass production (OD600)

was detected (Figure 4). Without arabinose supplementation, the cells grew to an OD

of approximately 0.8, which is due to the utilization of casamino acids (the same OD

was   obtained   with   ADP1*ΔaceA::tdk/Kan^r ΔpoxB::Cm^r*).   A   semi-quantitative   lipid

analyses based on thin layer chromatography (TLC) was carried out to compare the

amount of WEs produced (Figure 5B). For all the cultures, the same sample volume

was taken for the analysis, thus representing the titer of WEs produced. Based on

image analysis, the intensity of the band representing the wax ester titer increases

along with the biomass concentration. However, only a slight difference was observed

between the bands of the 0.5% and 1.0% cultures, suggesting that in the culture which

contains saturating amount of arabinose the growth corresponds to that of the wild-

type strain. When the intensities were divided with the optical densities, the highest

amounts of WEs (per cell) were produced in the cultures with 0.2% and 0.5%

arabinose. Thus, considering both the volumetric titer and the yield of WEs per

biomass, the arabinose concentration of 0.5% was found to be optimal in terms of

distributing the carbon and cellular resources between biomass and WEs.

Batch cultivations for ADP1 wt and ADP1-ara-aceA were carried out in minimal

medium supplemented with 0.1 % casein amino acids, 50 mM Na-acetate and 0.5 %

arabinose. Acetate and arabinose concentrations as well as the WE production (at 5,

7, 9 and 10 h time-points) were determined (Figure 5). ADP1 wt consumed all the

acetate in approximately 9-10 hours; the amount of biomass increased until the 10-

hour time-point, the CDW being 1.3 g/l. The highest amount of WEs was measured at

the 9-hour time-point, being 47 mg/g CDW and 60 mg/l. In ADP1 wt, the WEs

accounted for 44 % of total lipids. The WE yield was found to be 0.02 g WE/g consumed

acetate. The strain ADP1-ara-aceA utilized acetate more steadily and produced less

biomass compared to ADP1 wt; the growth ceased after 16 hours along with the

arabinose depletion: the biomass remained at 0.6-0.7 g/l CDW between the 16-24

hours of cultivation. Thereafter, the WE content of the cells strongly increased, being

highest at 38 h time-point, which also increased the amount of total biomass to 1.0

g/l. The WE titer was found to be 184 mg/l representing 19 % of CDW, which was 3.8-

fold higher compared to the ADP1 wt. In addition, the WEs accounted for 80 % of all

cellular lipids in ADP1-ara-aceA. The WE yield was 0.08 g WE/g consumed acetate, also

being 4-fold higher over the ADP1 wt.

4. Discussion

Sugars, mainly glucose, have been the major carbon source for the heterotrophic

microbial production of fatty acid derived compounds, such as TAGs and WEs, which

can be used for the production of biofuels, biochemicals, and other biocommodities

(29, 30). However, in order to increase the feasibility and sustainability of the

processes, the possibility to utilize alternative carbon sources is of high interest.

Organic acids, such as acetate, serves as a low-cost, abundant carbon source for

microbial lipid synthesis. Acetate can be readily derived from the hemicellulose

fraction of plant biomass or waste streams, or produced from syngases by microbial

fermentation (31). For example, we have previously demonstrated the conversion of

carbon dioxide to WEs via acetate intermediate by combining microbial

electrosynthesis with aerobic lipid synthesis in a two-stage process (32). Many of the

potential acetate streams, however, may have dilute acetate concentrations, thus

sustaining conditions that are preferable for biomass production rather than for

efficient storage carbon synthesis. On the other hand, highly concentrated acetate

feeds can inhibit cell growth (33).

Nitrogen starvation is a commonly used and efficient means to trigger storage lipid

accumulation in microbes (34-36). However, in such conditions the cell biomass

typically remains low (37), which can result in lower overall product titres. Thus,

genetic reprogramming would serve as a means to bypass the natural regulation for

storage lipid synthesis. Mechanisms behind the regulation in microbes are not well

understood, and therefore it has been challenging to genetically drive cells to

overproduce the storage compounds or other fatty acid derived products. Previous

strategies include for example the manipulation of the conserved carbon storage

regulator CsrA through the CsrA-CsrB ribonucleoprotein complex, by which alterations

in the central carbon metabolism and fatty acid synthesis regulation have led to

favourable changes in both native and non-native product synthesis pathways in *E.*

coli (38). This approach, however, is rather unspecific and potentially difficult to

combine with other (targeted) engineering strategies.

In this study, we developed an autonomously regulated circuit for programmable

synthesis of WEs in a native production host, *A. baylyi* ADP1. The circuit allows the

cells to shift from the biomass mode to the WE synthesis mode independent from the

carbon/nitrogen ratio or the growth phase of the culture. In practice, we replaced the

native isocitrate lyase aceA with an arabinose inducible system, which allows a

conditional and timed knockdown of the expression of *aceA*. This enzyme is essential

for the biomass production when the cells grow on acetate. The timed repression of

aceA expression is achieved by gradually eliminating the inducer, namely arabinose;

the native enzyme activity of glucose dehydrogenase Gcd of *A. baylyi* oxidizes

arabinose to arabino-lactone and further to arabonate, which in turn cannot serve as

inducers. Importantly, and in contrast to other auto-induction-based systems,

arabinose oxidation does not interfere with the utilization of the carbon source, here

acetate, and can be thus considered as an orthogonal system. By adjusting the

arabinose concentration, a predefined and optimal amount of biomass can be

produced.　When the inducer concentration is oxidized below the 'threshold'

concentration, the cells shift from the biomass producing mode to the synthesis mode,

efficiently directing carbon to product synthesis.

First, we confirmed that the engineered strain ADP1-ara-aceA with complemented

isocitrate lyase was able to grow on acetate as the sole carbon source. We observed

that without the presence of arabinose, the cells did not exhibit growth, showing

phenotype similar to the knockout strain *A. baylyi* ADP1Δ*aceA*::*tdk*/*Kan*[r]Δ*poxB*::*Cm*[r].

We also observed that arabinose concentration 1% is sufficient to allow the growth of

ADP1-ara-aceA to reach at least the same biomass as the ADP1 wt.

Initial arabinose concentrations were in correlation with the obtained biomasses

between concentrations 0.2% and 1.0%. The concentration 0.2% was found to be the

'threshold' for sufficient biomass production; below this concentration, the cells

produced only slightly more biomass compared to the uninduced cells, indicating that

arabinose concentrations >0.2% are required for sufficient growth in the studied

conditions. This finding was also supported by the reporter induction test; when the

cells containing the bacterial luciferase *luxAB* under arabinose promoter were induced

with the supernatant from different cultivation time-points (thus having different

arabinose concentrations), clear induction of *luxAB* determined as luminescence

production was only observed with the sample that contained >0.2% arabinose.

According to the semi-quantitative WE analyses, the cultures supplemented with 0.2%

or 0.5% arabinose produced the highest WE yields (per biomass). The cultures that

were induced with 1% arabinose produced nearly two times more biomass compared

to that of the 0.5% culture but had lower WE yield, indicating that a significant

proportion of the carbon was directed to the biomass when 1% arabinose was used.

Interestingly, the cultures with little (0.1%) or no arabinose produced the lowest WE

yield, suggesting that at least subtle levels of *aceA* expression are required, not only

for biomass production but also to support WE synthesis. The WE titres (WEs/volume)

of 0.5% and 1.0% cultures were estimated to be very close to each other in the two

cultures, whereas 0.2% culture had clearly lower volumetric WE production due to low

biomass production. Thus, we considered 0.5% arabinose as the most effective

inducer concentration in terms of optimal distribution of carbon between biomass and

products.

For validation of the system, batch cultures for the ADP1 wt and ADP1-ara-aceA were

carried out. It was shown that the engineered strain ADP1-ara-aceA efficiently

accumulated WEs in simple batch conditions supplied with relatively low substrate

concentration (50 mM acetate) and in non-optimal carbon-nitrogen ratio. The strain

ADP1-ara-aceA produced 187 mg/l WEs representing 19 % of the CDW and a yield of

0.08 g WE/g consumed acetate. As expected, the strain grew more slowly and

produced less biomass than ADP1 wt, but the yield of WEs per biomass and per

consumed acetate were 3.8 and 4 folds higher compared to ADP1 wt, respectively. In

addition, the WE titer was found to be 3.1 folds higher compared to that of the ADP1

wt. Thus, the dynamic regulation not only improved the yield of WEs per biomass and

per used carbon, but clearly excelled the volumetric titer of that of the wild type strain.

For comparison, in a previous study (32), *A. baylyi* ADP1 produced WEs from acetate

with a titer of approximately 90 mg/l (from higher initial acetate concentration, 100

mM), with an average yield of 4 % (carbon/carbon), being equal to 0.02 g WE/g

consumed acetate. The highest WE titer reported so far has been 450 mg/l, which was

obtained when the key enzyme of the pathway (fatty acyl-CoA reductase) was

overexpressed and 5 % glucose was used as the substrate (17). However, the yields

(0.04g WE/g glucose and 12.5% WEs of CDW) were lower compared to this study. With

external alkane supplementation, up to 17% WEs of CDW has been obtained (39).

Notably, the amount of WEs per cell was nearly constant in ADP1 wt at the analysed

time points, being 3.9-4.3% of the CDW. In ADP1-ara-aceA, by contrast, the amount of

WEs per cell strongly increased along with the arabinose depletion; the percentage of

WEs per cell increased from 5.2% to 19% between the sampling points. The highest

increase in the WE content (from 7.5 to 19%) was achieved after the arabinose

concentration reached the 'threshold' 0.2% (equivalent to 15 mM arabinose).

Although the shift from biomass mode to lipid mode was rather gradual, the arabinose

concentration 0.2% seems to be the key turning point in the cellular state. In the batch

culture, initial concentrations 50 mM acetate and 0.5% arabinose were used. By the

end of the culture, the arabinose was completely oxidized, and only a small amount

(5 mM) of acetate remained unutilized. Thus, by adjusting the substrate and inducer

concentrations, the system is potentially scalable to a wide range of substrate

concentrations. Moreover, coupling this system with other engineering strategies,

such as introducing additional knock-outs (16) and/or overexpression of key enzymes

of the pathway (17) could further improve the WE production.

Considering not only the efficient redirection of carbon to product, but also the

downstream processing, the purity of the product is important. In ADP1-ara-aceA,

WEs constituted 80% of total lipids, indicating high purity of the desired product. In

ADP1 wt, only 44% of the total lipids were WEs.

The results from different experiments indicate that at least low levels of isocitrate

lyase are required to maintain WE production from acetate, potentially due to the

requirements for cells to generate e.g. NADPH for the synthesis. This hypothesis was

also supported by a further observation in an additional experimental set-up where

the WE content of the *aceA* knockout strain did not increase after a transfer from a

glucose medium to an acetate medium (data not shown). While arabinose

concentration <0.2% is not sufficient to promote biomass production, it allows the

cells to synthetize and maintain the required cofactor balance, and to efficiently

produce WEs.

Rapid advancements in the CRISPR/Cas9 technologies have broaden the tools

available for targeted genome engineering, and especially the employment of the

deactivated Cas9 (dCas9) has recently gained interest in the context of targeted gene

silencing (40, 41). While the dCas9 –based tools have been shown to be functional and

applicable in a wide range of (microbial) hosts, challenges related to unpredictability,

cellular burden and off-targeting may limit its use (42, 43). In addition, at least two

different constructs are typically required for the fine-tuned expression of dCas9 and

the RNA elements, and the system cannot be easily operated 'hands-free' without a

timed addition of an inducer. In this context, the system described here provides a

straight-forward, readily controllable, and reliable set-up for conditional and timed

gene knock-down. In addition, other interesting synthetic biology hosts such as *P.*

putida (44, 45) exhibit the same glucose dehydrogenase activity and could thus find

this strategy applicable. However, the transferability of this system to others hosts

such as *E. coli* and *S. cerevisiae* remains to be investigated in the future.

Our system serves as a simple and scalable method for dynamic, 'hands-free'

regulation of growth-essential reactions in the cell, which allows targeted and

adjusted biomass and product synthesis. Here, the dynamic regulation system was

exploited in the conversion of acetate to carbon-rich storage compounds, namely wax

esters, that are otherwise not efficiently accumulated in cells without optimized

conditions and high carbon-nitrogen ratio. Moreover, the system could be potentially

utilized and generalized for a broad range of synthesis pathways that are dependent

on acetyl-CoA (17, 46). In addition, other sustainable carbon sources, such as lignin-

derived compounds (47-49) would be compatible with this system.

5. Conclusions

We showed that an autonomously regulated genetic switch allowed the dynamic

decoupling of biomass and wax ester production in engineered *A. baylyi* ADP1, which

resulted in 3-4 fold improvements in the wax ester yield and titer compared to the

wild type strain. Shifting the cells from a biomass mode to a product synthesis mode

was achieved by gradually repressing the growth essential gene *aceA* by a simple and

robust set-up. The engineered strain produced 19% WEs of its cell dry weight, being

the highest reported among microbes. The study demonstrates the possibility to

bypass the challenges related to highly regulated storage lipid synthesis, and

strengthens the status of *A. baylyi* ADP1 as a convenient host for metabolic

engineering and high-value lipid production from sustainable substrates.

Acknowledgements

472 Funding: This work was supported by the Academy of Finland (grant numbers 286450,

473 310135, 310188, and 311986).

References

1. Tan SZ & Prather KL (2017) Dynamic pathway regulation: recent advances and

methods of construction. *Curr Opin Chem Biol* 41:28-35.

2. Soma Y, Tsuruno K, Wada M, Yokota A, & Hanai T (2014) Metabolic flux

redirection from a central metabolic pathway toward a synthetic pathway using

a metabolic toggle switch. *Metabolic engineering* 23:175-184.

3. Soma Y & Hanai T (2015) Self-induced metabolic state switching by a tunable

cell density sensor for microbial isopropanol production. *Metabolic engineering*

30:7-15.

4. Solomon KV, Sanders TM, & Prather KL (2012) A dynamic metabolite valve for

the control of central carbon metabolism. *Metabolic engineering* 14(6):661-671.

5.   Brockman IM & Prather KL (2015) Dynamic knockdown of E. coli central

metabolism for redirecting fluxes of primary metabolites. *Metabolic engineering*

28:104-113.

6.   Moser F*, et al.* (2012) Genetic circuit performance under conditions relevant for

industrial bioreactors. *ACS Synthetic Biology* 1(11):555-564.

7.   Wältermann M*, et al.* (2005) Mechanism of lipid-body formation in prokaryotes:

how bacteria fatten up. *Molecular microbiology* 55(3):750-763.

8.   Wältermann M & Steinbüchel A (2005) Neutral lipid bodies in prokaryotes:

recent insights into structure, formation, and relationship to eukaryotic lipid

depots. *Journal of bacteriology* 187(11):3607-3619.

9.   Xu P, Li L, Zhang F, Stephanopoulos G, & Koffas M (2014) Improving fatty

acids production by engineering dynamic pathway regulation and metabolic

control. *Proceedings of the National Academy of Sciences of the United States*

of America 111(31):11299-11304.

10.   Teixeira PG, Ferreira R, Zhou YJ, Siewers V, & Nielsen J (2017) Dynamic

regulation of fatty acid pools for improved production of fatty alcohols in

Saccharomyces cerevisiae. *Microbial Cell Factories* 16(1):45.

11.   Kurosawa K, Boccazzi P, de Almeida NM, & Sinskey AJ (2010) High-cell-
513   density batch fermentation of Rhodococcus opacus PD630 using a high
514   glucose concentration for triacylglycerol production. *J Biotechnol* 147(3-4):212-
515   218.

517   12.   Xu J, Liu N, Qiao K, Vogg S, & Stephanopoulos G (2017) Application of
518   metabolic controls for the maximization of lipid production in semicontinuous
519   fermentation. *Proceedings of the National Academy of Sciences of the United*
520   *States of America* 114(27):E5308-E5316.

522   13.   Runguphan W & Keasling JD (2014) Metabolic engineering of Saccharomyces
523   cerevisiae for production of fatty acid-derived biofuels and chemicals.
524   *Metabolic engineering* 21:103-113.

526   14.   Plassmeier J, Li Y, Rueckert C, & Sinskey AJ (2016) Metabolic engineering
527   Corynebacterium glutamicum to produce triacylglycerols. *Metabolic*
528   *engineering* 33:86-97.

530   15.   Tai M & Stephanopoulos G (2013) Engineering the push and pull of lipid
531   biosynthesis in oleaginous yeast Yarrowia lipolytica for biofuel production.
532   *Metabolic engineering* 15:1-9.

534   16.   Santala S*, et al.* (2011) Improved triacylglycerol production in *Acinetobacter*
535   *baylyi* ADP1 by metabolic engineering. *Microb Cell Fact* 10:36.

17. Lehtinen T, Efimova E, Santala S, & Santala V (2018) Improved fatty aldehyde and wax ester production by overexpression of fatty acyl-CoA reductases. *Microbial Cell Factories* 17(1):19.

18. Santala S, Efimova E, Koskinen P, Karp MT, & Santala V (2014) Rewiring the wax ester production pathway of *Acinetobacter baylyi* ADP1. *ACS Synth Biol* 3(3):145-151.

19. Klinke S, Dauner M, Scott G, Kessler B, & Witholt B (2000) Inactivation of isocitrate lyase leads to increased production of medium-chain-length poly(3-hydroxyalkanoates) in Pseudomonas putida. *Applied and environmental microbiology* 66(3):909-913.

20. Young DM, Parke D, & Ornston LN (2005) Opportunities for genetic investigation afforded by Acinetobacter baylyi, a nutritionally versatile bacterial species that is highly competent for natural transformation. *Annual review of microbiology* 59:519-551.

21. Kannisto M, Aho T, Karp M, & Santala V (2014) Metabolic engineering of Acinetobacter baylyi ADP1 for improved growth on gluconate and glucose. *Applied and environmental microbiology* 80(22):7021-7027.

22.   Kannisto MS*, et al.* (2015) Metabolic engineering of Acinetobacter baylyi ADP1

for removal of Clostridium butyricum growth inhibitors produced from

lignocellulosic hydrolysates. *Biotechnol Biofuels* 8(1):1-10.

23.   Carr EL, Kampfer P, Patel BK, Gurtler V, & Seviour RJ (2003) Seven novel

species of Acinetobacter isolated from activated sludge. *International journal of*

systematic and evolutionary microbiology 53(Pt 4):953-963.

24.   Barbe V*, et al.* (2004) Unique features revealed by the genome sequence of

Acinetobacter sp. ADP1, a versatile and naturally transformation competent

bacterium. *Nucleic acids research* 32(19):5766-5779.

25.   de Berardinis V*, et al.* (2008) A complete collection of single-gene deletion

mutants of *Acinetobacter baylyi* ADP1. *Molecular systems biology* 4:174.

26.   Santala S, Efimova E, Karp M, & Santala V (2011) Real-time monitoring of

intracellular wax ester metabolism. *Microb Cell Fact* 10:75.

27.   Murin CD, Segal K, Bryksin A, & Matsumura I (2012) Expression vectors for

Acinetobacter baylyi ADP1. *Applied and environmental microbiology*

78(1):280-283.

28. Santala V, Karp M, & Santala S (2016) Bioluminescence based system for
rapid detection of natural transformation. *FEMS microbiology letters*.

29. Lennen RM & Pfleger BF (2013) Microbial production of fatty acid-derived fuels
and chemicals. *Current opinion in biotechnology* 24(6):1044-1053.

30. Marella ER, Holkenbrink C, Siewers V, & Borodina I (2018) Engineering
microbial fatty acid metabolism for biofuels and biochemicals. *Current opinion*
in biotechnology 50:39-46.

31. Bengelsdorf FR, Straub M, & Durre P (2013) Bacterial synthesis gas (syngas)
fermentation. *Environ Technol* 34(13-16):1639-1651.

32. Lehtinen T*, et al.* (2017) Production of long chain alkyl esters from carbon
dioxide and electricity by a two-stage bacterial process. *Bioresource*
Technology 243:30-36.

33. Trček J, Mira NP, & Jarboe LR (2015) Adaptation and tolerance of bacteria
against acetic acid. *Applied microbiology and biotechnology* 99(15):6215-6229.

34. Breuer G, Lamers PP, Martens DE, Draaisma RB, & Wijffels RH (2012) The
impact of nitrogen starvation on the dynamics of triacylglycerol accumulation in
nine microalgae strains. *Bioresource Technology* 124:217-226.

35. Alvarez HM & Steinbüchel A (2002) Triacylglycerols in prokaryotic
microorganisms. *Applied microbiology and biotechnology* 60(4):367-376.

36. Ishige T, Tani A, Sakai Y, & Kato N (2003) Wax ester production by bacteria.
Current opinion in microbiology 6(3):244-250.

37. Amara S*, et al.* (2016) Characterization of key triacylglycerol biosynthesis
processes in rhodococci. *Sci Rep* 6:24985.

38. McKee AE*, et al.* (2012) Manipulation of the carbon storage regulator system
for metabolite remodeling and biofuel production in Escherichia coli. *Microbial*
Cell Factories 11:79.

39. Ishige T*, et al.* (2002) Wax ester production from n-alkanes by Acinetobacter
sp. strain M-1: ultrastructure of cellular inclusions and role of acyl coenzyme A
reductase. *Applied and environmental microbiology* 68(3):1192-1195.

40. Larson MH*, et al.* (2013) CRISPR interference (CRISPRi) for sequence-specific
control of gene expression. *Nat Protoc* 8(11):2180-2196.

41. Peters JM*, et al.* (2016) A Comprehensive, CRISPR-based Functional Analysis
of Essential Genes in Bacteria. *Cell* 165(6):1493-1506.

42. Nielsen AA & Voigt CA (2014) Multi-input CRISPR/Cas genetic circuits that interface host regulatory networks. *Molecular systems biology* 10:763.

43. Cui L*, et al.* (2018) A CRISPRi screen in E. coli reveals sequence-specific toxicity of dCas9. *Nat Commun* 9(1):1912.

44. Nikel PI & de Lorenzo V (2018) Pseudomonas putida as a functional chassis for industrial biocatalysis: From native biochemistry to trans-metabolism. *Metabolic engineering*.

45. Nikel PI, Chavarria M, Danchin A, & de Lorenzo V (2016) From dirt to industrial applications: Pseudomonas putida as a Synthetic Biology chassis for hosting harsh biochemical reactions. *Curr Opin Chem Biol* 34:20-29.

46. Lehtinen T, Santala V, & Santala S (2017) Twin-layer biosensor for real-time monitoring of alkane metabolism. *FEMS microbiology letters* 364(6).

47. Linger JG*, et al.* (2014) Lignin valorization through integrated biological funneling and chemical catalysis. *Proceedings of the National Academy of Sciences of the United States of America* 111(33):12013-12018.

649   48.   Salvachúa D, Karp EM, Nimlos CT, Vardon DR, & Beckham GT (2015)

650   Towards lignin consolidated bioprocessing: simultaneous lignin

651   depolymerization and product generation by bacteria. *Green Chemistry* DOI:

652   10.1039/c5gc01165e.

654   49.   Salmela MM*, et al.* (2018) Molecular tools for selective recovery and detection

655   of lignin-derived molecules. *Green Chemistry*.

Figure legends

Figure 1. An autonomously regulated switch for shifting the cells from a biomass to a product synthesis mode. A) Wax ester synthesis strongly competes with biomass production; In the wild type cells, efficient wax ester synthesis is triggered only in defined conditions (i.e. in conditions with nitrogen starvation and excess carbon). In normal growth conditions, most of the carbon is directed to biomass production and cell maintenance. B) The ADP1 wild type strain utilizes acetate for biomass production through a glyoxylate shunt in citric acid cycle. Isocitrate lyase (AceA) is the key enzyme in the glyoxylate shunt and thus essential for growth on acetate. Strain lacking this enzyme is unable to grow on acetate as the sole carbon source. C) The genetic construct for dynamic regulation of *aceA* expression and growth. In the construct, *aceA* is placed under the arabinose-inducible promoter AraC-pBAD. When cultured on acetate, the arabinose used as the inducer is slowly oxidized by the native glucose dehydrogenase Gcd of ADP1, gradually repressing the expression of AceA. Arabinose oxidation does not serve as the carbon source or interfere with the acetate utilization, thus being orthogonal to the circuit function. D) Along with the inducer depletion (arabinose oxidation), the cells gradually shift from the biomass mode to the lipid synthesis mode; the less there is arabinose left in the culture, a higher proportion of the carbon flux is directed to the product. The amount of biomass can be simply regulated by adjusting the initial arabinose concentration.

Figure 2. Growth of ADP1 wt and the engineered strain ADP1-ara-aceA in minimal

medium. The cells were cultured for 68 hours at 25 °C in MSM supplemented with 50

mM Na-acetate as the sole carbon source. Arabinose concentrations of 0; 0.1 and 1.0%

(for ADP1 wt 0 and 1%) were used for the induction. Optical densities representing

biomasses are presented as an average of two individual replicates.

Figure 3. Growth of ADP1-ara-aceA and ADP1 wt with different arabinose

concentrations. The cells were cultured in MA/9 medium supplemented with 25 mM

acetate, 0.1% cas.amino acids, and arabinose (0-1%) at 25 °C in micro well-plates for

21 hours. The optical densities are presented as the average of three different

replicate wells. For ADP1 wt, only the growth curve of the culture containing 1%

arabinose is shown.

Figure 4. Biomass and wax ester production by ADP1-ara-aceA induced with different

arabinose concentrations. ADP1-ara-aceA was cultured in MA/9 supplemented with

50 mM acetate, 0.1% casam, and arabinose (0, 0.1, 0.2, 0.5, or 1.0%) for 30 hours. A)

The amount of the produced biomass (at the end-point) was determined by optical

density (600 nm) measurement, presented as an average of two individual replica

cultures. B) Wax ester production in the cultures with different arabinose

concentrations were determined by thin layer chromatography analysis, for which 3

ml samples for taken from each culture. Jojoba oil was used as the standard.

Figure 5. Acetate utilization, arabinose oxidation, and the accumulation of wax esters

(WE) in the batch cultures of A) ADP1 wt and B) ADP1-ara-aceA. The cells were

cultivated in MA/9 medium supplemented with 50 mM acetate, 0.1% cas. amino acids,

and 0.5% arabinose for 12 h (ADP1 wt) or 38 h (ADP1-ara-aceA). For the arabinose and

acetate concentrations determined by HPLC, the average and standard deviation for

samples from two individual cultures are presented. Similarly, the WEs were

quantitatively analyzed by NMR from two individual cultures. The WE production at

different time-points is presented as mg/g CDW to demonstrate the different

accumulation patterns of the strains.

713
714

Tables

Table 1. List of primers used in the study.

Name	Sequence (5'-> 3')
VS10_9	CAATGAATTCCGATAAAAGCGGATTCCTGAC
VS10_10	ATCCCATATGTAATTCCTCCTGTTAG
SS17_08	AATACATATGACATATCAATCAGCTCTTGAGC
SS17_09	TAATACAATTGCGAACAGGCTTATGTCAAGACGTC

727 Figure 1.

730   Figure 2.

733   Figure 3.

737   Figure 4
738

742　　　Figure 5A.

745
746
747