Inhibition of thrombocyte activation restores protective immunity to

mycobacterial infection

Elinor Hortle[1], Khelsey E. Johnson[2], Matt D. Johansen[1], Tuong Nguyen[1], Jordan A. Shavit[3],

Warwick J. Britton[1,4], David M. Tobin[2], Stefan H. Oehlers[1,4] *

[1] Tuberculosis Research Program, Centenary Institute, Camperdown, NSW 2050, Australia

[2] Department of Molecular Genetics and Microbiology, Duke University School of Medicine,

Durham, NC 27710, USA

[3] Department of Pediatrics and Cellular and Molecular Biology Program, University of

Michigan, Ann Arbor, MI 48107, USA

[4] Central Clinical School and Marie Bashir Institute, The University of Sydney,

Camperdown, NSW 2050, Australia

* Corresponding author email: s.oehlers@centenary.org.au, Twitter: @oehlerslab

Summary

Infection-induced thrombocytosis is a clinically important complication of tuberculosis

infection. Recent studies have highlighted the utility of aspirin as a host-directed therapy

modulating the inflammatory response to infection, but have not investigated the possibility

of an anti-platelet mode of action. Here we utilise the zebrafish-*Mycobacterium*

marinum model to show mycobacteria drive host haemostasis through the formation of

granulomas. Treatment of infected zebrafish with aspirin markedly reduced mycobacterial

burden. This effect is phenocopied by treatment with platelet-specific glycoprotein IIb/IIIa

inhibitors demonstrating a detrimental role for infection-induced thrombocyte activation. We

find that the reduction in mycobacterial burden is dependent on macrophages and granuloma

formation providing the first *in vivo* experimental evidence that infection-induced platelet

activation compromises protective host immunity to mycobacterial infection. Our study

illuminates platelet activation as an efficacious target of aspirin, a widely available and

affordable host-directed therapy candidate for tuberculosis.

Keywords

Mycobacterial infection, haemostasis, innate immunity, clotting

Introduction

Mycobacterium tuberculosis is the world's most lethal pathogen, causing nearly 2 million

deaths each year (World Health Organization. and Global Tuberculosis Programme.). The

increasing incidence of both multi- and extremely- drug resistant tuberculosis (TB) urgently

require the development of therapeutics that overcome the shortcomings of conventional

antibiotics. Pathogenic mycobacteria co-opt numerous host pathways to establish persistent

infection, and subversion of these interactions with host-directed therapies (HDTs) has been

shown to reduce the severity of infection in animal models. For example, we have recently

shown that mycobacteria induce host angiogenesis and increase host vascular permeability;

blockade of either of these processes reduced both the growth and spread of bacteria (Oehlers

et al., 2017; Oehlers et al., 2015). Therefore, host processes co-opted by the bacteria provide

attractive targets for novel TB treatments. One such pathway may be haemostasis.

Thrombocytosis has long been recognised as a biomarker for advanced TB, and infection is

often accompanied by the induction of a hyper-coagulable state, resulting in increased risk of

deep vein thrombosis and stroke (Kutiyal et al., 2017; Robson et al., 1996). Recent evidence

hints that mycobacteria may drive this process, and that it may aid their growth. For example,

cell wall components from *M. tuberculosis* can induce expression of tissue factor - an

important activator of coagulation - in macrophages (Kothari et al., 2012). In mice and

humans markers of platelet activation are upregulated during *M. tuberculosis* infection (Fox

et al., 2018), and it has been shown *in vitro* that interaction with activated platelets increases

the conversion of infected macrophages into cells permissive for bacterial growth (Feng et

al., 2014; Fox et al., 2018). To date the pathogenic roles of haemostasis have not been studied

in an intact *in vivo* model of mycobacterial infection.

Here we used the zebrafish-*M. marinum* model to investigate the role of host thrombocytes in

mycobacterial infection. We show evidence that coagulation and thrombocyte activation are

both bacterially driven, and that infection-induced activation of thrombocytes compromises

protective immunity. However, clot formation itself has no effect on infection suggesting

direct communication between the mycobacterial infection milieu and thrombocytes.

Results

Mycobacterium marinum **induces coagulation in zebrafish.**

To determine if *M. marinum* induces coagulation in zebrafish, we infected *Tg(fabp10a:fgb-*

EGFP) transgenic embryos expressing EGFP tagged fibrinogen beta (FGB) with *M.*

marinum-tdtomato, and imaged the developing infection every 15 minutes from 3 days post

infection (DPI), until 6 DPI (Supplementary Video 1). We observed that clots formed only at

areas of bacterial growth, and that the size of the clots increased as the number of bacteria

increased over the course of the infection (Figure 1A). When we infected fish with ΔESX1

mutant *M. marinum* - which lacks the ability to export key virulence proteins, and does not

form granulomas – significantly less clot formation was observed (Figures 1B and 1C).

Together, this data demonstrates that coagulation is a conserved consequence of infection

driven by pathogenic mycobacteria across host species. Treatment with the anti-coagulant

warfarin to prevent clot formation during infection (Figures 1D-E) did not affect bacterial

burden, suggesting coagulation does not affect bacterial growth within the host (Figure 1F).

***Mycobacterium marinum* induces thrombosis in zebrafish.**

To determine if *M. marinum* induces thrombosis in zebrafish, we infected *Tg(cd41:GFP)*

embryos, where thrombocytes are marked by GFP expression, with red fluorescent

M. marinum. Infected embryos had significantly increased density of thrombocytes around

the tail venous plexus where granulomas preferentially form (Figures S1A-B). Amongst

infected embryos, there was a strong positive correlation between thrombocyte density and

mycobacterial burden (Figure S1C). Both increased thrombocyte density, and a positive

correlation between density and burden, were also observed in ΔESX1 mutant *M. marinum*

infection (Figures S1C-D), suggesting that bacterially induced thrombosis is ESX1

independent.

Because the *cd41* promoter is active in non-motile thrombocyte precursors within their

caudal haematopoietic tissue, we could not conclusively determine if these thrombocytes had

actively migrated to and been retained at the site of infection (Lin et al., 2005). To determine

if zebrafish thrombocytes are recruited to sites of mycobacterial infection, we performed

trunk injection of *M. marinum* in *Tg(cd41:GFP)* at 3 days post fertilization (DPF), a time

point after which mature thrombocytes are in the circulation. Embryos were then imaged at 2,

3, and 4 DPI. Thrombocytes were co-localised with sites of infection in 87% of fish. Rather

than forming a stable and growing clot over a period of days (as we observed for FGB),

thrombocytes appeared to form transient associations with sites of infection (Figure S1D). In

some instances, new thrombocytes seemed to be retained at sites of infection in different

locations each day (Figure 2A I-IV), and in others, thrombocytes appeared to stay stationary

for multiple days (Figure 2A V-VIII). The size of infection-associated thrombocytes varied

greatly, suggesting activation and progressive degranulation or death of thrombocytes at sites

of infection. Thrombocytes were most often observed on the edges of granulomas consistent

with the location of granuloma-defining macrophages (Cronan et al., 2016).

Anti-platelet drugs reduce mycobacterial burden in zebrafish.

To determine if the previously reported infection-controlling effect of aspirin is conserved

across vertebrate species (Mai et al., 2018), we treated *M. marinum*-infected fish with aspirin

by immersion. Mycobacterial burden was reduced by approximately 50% in aspirin treated

embryos (Figure 2A).

Although aspirin is a widely used platelet inhibitor, it is a broadly acting nonsteroidal anti-

inflammatory drug that is known to modulate infection-relevant prostaglandin metabolism.

Thus, to determine if the anti-platelet effects of aspirin treatment contribute to the reduced

mycobacterial burden, we treated *M. marinum*-infected fish with the platelet-specific small

molecule Glycoprotein IIb/IIIa inhibitor tirofiban or eptifibatide. Treatment with either

Glycoprotein IIb/IIIa inhibitor phenocopied aspirin by reducing bacterial burden providing

the first direct evidence of a detrimental role for platelet activation in the immune response to

mycobacterial infection (Figure 2B-C).

Surprisingly, tirofiban treatment resulted in significantly higher number of visible

thrombocytes at sites of infection compared to controls (Figures 2D-E). This suggests that, in

contrast to our observations in the tail wound thrombosis model, tirofiban does not inhibit

thrombocyte migration to areas of bacterial infection but may act by preventing activation

and subsequent degranulation or death of zebrafish thrombocytes in response to

mycobacterial infection.

We next examined the cellular target of anti-platelet drugs in our infection system. We

performed antibacterial testing of the anti-platelet drugs in axenic cultures of *M. marinum*

and did not observe any effect on bacterial growth *in vitro* demonstrating host-directed

activity (Figure 2F).

The specificity of anti-platelet drugs against mycobacterial infection was tested in acute

Pseudomonas aeruginosa infection. Treatment of *P. aeruginosa*-infected embryos with either

class of anti-platelet drug did not affect survival demonstrating specificity against chronic

mycobacterial infection (Figure 2G).

The anti-bacterial effect of anti-platelet drugs is thrombocyte dependent.

We next confirmed that anti-platelet drugs inhibit thrombocyte activation in zebrafish using a

tail wound thrombosis assay (Figure S4A). Anti-platelet drugs reduced the number of

thrombocytes recruited to tail wound clots demonstrating conservation of cellular target in

zebrafish embryos (Figure S4B-C). Of the two Glycoprotein IIb/IIIa inhibitors tirofiban and

eptifibatide, tirofiban is the more stable small molecule and therefore we chose to use tirofiban

for inhibition of thrombocytes in all further experiments.

To determine if zebrafish thrombocytes are the conserved target for tirofiban in the zebrafish-

M. marinum infection model, we inhibited thrombopoiesis by injection with a morpholino

against the thrombopoietin receptor *cmpl* (Lin et al., 2005). Surprisingly, inhibition of

thrombopoiesis did not affect the outcome of infection. However, tirofiban treatment failed to

reduce bacterial burden in thrombocyte-depleted embryos, demonstrating thrombocytes are

the cellular target of this drug in the zebrafish infection model (Figure 3A).

To confirm that tirofiban was acting specifically through inhibition of Glycoprotein IIb/IIIa, we

performed infection experiments in Glycoprotein IIb/IIIa knock-out (KO) transgenic embryos

(*itga2b* mutants). The *itga2b^{sa10134}* allele caused a dose-dependent reduction in thrombocytes

recruited to tail wound clots (Figure S4A). As with *cmpl* knockdown, complete KO of *itga2b*

had no effect on bacterial burden (Figure 3F), and thrombocyte accumulation at the site of

infection was also unaffected (Figures S4B-C). However, heterozygous embryos phenocopied

tirofiban treatment, showing a trend to reduced bacterial burden and increased thrombocyte

accumulation (Figures 3B and S4B-C). This suggests that treatment with tirofiban may only

partially inhibit Glycoprotein IIb/IIIa in our model.

Thrombocyte activation and coagulation are independently induced by mycobacterial

infection.

We hypothesised that infection-induced coagulation and the resulting fibrin deposition in clots

would activate thrombocytes through fibrin cross-linking. To investigate this hypothesis, we

again used warfarin, which prevented clot formation during infection and did not affect bacterial

burden (Figure 1E). Surprisingly, when warfarin was added to our tirofiban treatment model, it

had no effect on the ability of tirofiban to reduce bacterial burden (Figure 3C).

To discount the possibility that warfarin and tirofiban co-treatment resulted in non-specific

toxicity, we utilised a *fibrinogen alpha chain* mutant zebrafish line that does not readily form

clots. Tirofiban treatment of FGA mutant zebrafish still significantly reduced bacterial burden

further demonstrating that infection-induced activation of coagulation is not upstream of

thrombocyte activation by mycobacterial infection (Figure 3D).

Conversely, we next asked if thrombocyte activation contributes to infection-induced

coagulation. Analysis of FGB-GFP clots in tirofiban-treated embryos revealed reduced total clot

formation (Figure 3E). However, correction for relative bacterial burden revealed that the

reduced clot formation was a burden-dependent and thrombus formation was not additionally

impacted by tirofiban treatment (Figure 3F).

Thrombocyte activation compromises immunity through granuloma-associated

macrophages.

To further delineate the mechanism by which thrombocyte activation compromises innate

immune control of mycobacterial infection but not the response to Pseudomonas infection, we

asked if granuloma formation and maturation is essential to pathological thrombocyte activation.

To examine the requirement of granuloma formation and necrosis for thrombocyte-inhibiting

drug efficacy, we infected embryos with ΔESX1 *M. marinum* that are unable to drive granuloma

maturation or necrosis. Tirofiban treatment had no effect on mutant bacterial burden at 5 DPI

(Figure 4A). Similarly, in embryos infected with *M. marinum* at a dose too low to form necrotic

granulomas by 5 DPI, continuous tirofiban treatment had no effect on bacterial burden at 5 DPI

(Figure S6A).

We next took advantage of the stereotypical progression of innate immune granulomas in

zebrafish embryos to investigate the temporal activity of thrombocyte inhibition by tirofiban.

We found that at 3 DPI, a time-point with nascent granuloma formation but prior to significant

granuloma organisation and necrosis, tirofiban had no effect on bacterial burden (Figure 4B).

Conversely, treatment of infection from 4 to 5 DPI, a time-point when granulomas are organised

and start to become necrotic, tirofiban significantly reduced bacterial burden within 24 hours

(Figure 4C).

Macrophages are key cells in the innate granuloma, therefore to determine if thrombocyte

inhibition exerts a protective effect through boosting macrophage-dependent immunity, we to

depleted macrophages. We injected clodronate liposomes to deplete macrophages early during

granuloma formation at 3 DPI (Figure S6B-C). Anti-platelet drug treatment was started

immediately after liposome injection. In macrophage-depleted fish tirofiban did not reduce

bacterial burden (Figure 4D).

It has previously been shown that in the presence of mycobacteria *in vitro*, platelets

accelerate the conversion of macrophages to foam cells, which are permissive for

mycobacterial growth, suggesting a mechanism for infection-induced thrombocytosis to

compromise innate immunity to mycobacterial infection (Feng et al., 2014). We therefore

hypothesised that thrombocyte inhibition would reduce the conversion of macrophages into

foam cells. We investigated this by performing Oil-red O staining to measure lipid

accumulation within individual granulomas (Figure 4E). Because tirofiban treatment reduced

bacterial burden, Oil-red O density was only compared between size-matched granulomas in

treatment group. Tirofiban-treated embryos had significantly less Oil-red O accumulation in

their granulomas when compared to DMSO control, even after correction for reduced

bacterial burden (Figure 4F). Together, these data demonstrate an *in vivo* effect of

thrombocyte activation inhibiting an effective immune response by converting macrophages

into foam cells in mycobacterial granuloma.

Given that foam cell formation is closely associated with necrosis in tuberculosis (Russell et

al., 2009) we hypothesised that tirofiban treatment would reduce cell death within the

granuloma. We therefore used TUNEL staining to detect the fragmented DNA of dying cells

in *M. marinum* infected embryos. At 5 DPI tirofiban-treated embryos showed significantly

less TUNEL staining, indicating significantly reduced cell death within the granuloma

(Figures 4G-H). Together these results suggest that infection-induced platelet activation

aggravates a basal rate of macrophage cell death in the granuloma.

Discussion

Here we have used the zebrafish-*M. marinum* model to identify infection-induced

haemostasis as a detrimental host response that is co-opted by pathogenic mycobacteria. Our

data builds on previous studies that have shown coagulation, thrombocytosis and

thrombocyte activation are associated with mycobacterial infection, and provide the first *in*

vivo study to demonstrate a direct role for thrombocyte activation in promoting mycobacterial

growth. We have shown that infection-induced haemostasis is conserved in the zebrafish-*M.*

marinum infection model, and that the platelet inhibiting drugs aspirin, tirofiban, and

eptifibatide are able to reduce bacterial burden, independently of effects on coagulation,

through host-mediated effects (Graphical abstract).

A number of studies have investigated aspirin as a possible adjunctive treatment for TB in

both mice and humans (reviewed in (Hawn et al., 2013; Kalantri and Kalantri, 2018; Kroesen

et al., 2017; Kroesen et al., 2018)). The results of these studies have been far from

conclusive, with some finding beneficial effects, others finding no effect, and still others

finding aspirin may reduce the effectiveness of anti-tubercular drugs. This lack of consensus

may be due to the fact that NSAIDs affect many cell types and processes important for the

host response to mycobacterial infection, not just coagulation and platelet activation. Our

study expands this literature by demonstrating a significant reduction in bacterial burden

following treatment with two specific anti-platelet drugs – tirofiban and eptifibatide, in addition

to aspirin.

Our study found that coagulation and thrombocytes are independently activated by infection

and have distinct roles during the pathogenesis of mycobacterial infection of zebrafish.

Inhibiting coagulation alone did not significantly reduce bacterial burden, and therefore we

considered anti-platelet treatment as a more attractive HDT than anti-coagulant treatment.

Although we found lower total clot formation in tirofiban-treated embryos, this was only

proportional to bacterial load suggesting infection-induced coagulation could be independent of

infection-induced thrombocyte activation. It must be noted that we only measure a simple single

end-point in our zebrafish embryo experiments (bacterial load) at a relatively early time point for

a chronic infection. In more complex animals, where stroke and DVT are important secondary

complications of mycobacterial infection, reducing coagulation may yet prove to be efficacious

as a HDT to reduce morbidity. Conversely, data from the more complex mouse model of TB

suggests tissue factor-induced fibrin is necessary to contain mycobacteria within granulomas

(Venkatasubramanian et al., 2016).

Our study provides evidence that infection-induced haemostasis is a conserved function of

the core mycobacterial pathogenicity program centred around granuloma formation. Our

experiments with ΔESX1 mutant *M. marinum*, which cannot secrete key virulence proteins

that drive granuloma formation, demonstrated ESX1-dependent haemostasis and

responsiveness to growth restriction by platelet inhibiting drugs. These data fit well with our

observations that stationary thrombocytes were only observed around well-developed

mycobacterial granulomas, and platelet inhibition was only effective at reducing bacterial

burden after the development of significant granuloma pathology. Together these results point

to a bidirectional relationship between thrombocyte activation and granuloma maturation

(Figure 6).

Interestingly we observed that tirofiban treatment increased the number of thrombocytes

accumulated at sites of infection. This may represent increased migration, increased

accumulation through coagulation or decreased activation and subsequent

degranulation/death. Given Tirofiban blocks the glycoprotein IIb/IIIa integrin - on which

platelet migration is dependent (Gaertner et al., 2017) - and we observed that tirofiban

reduced migration of thrombocytes in our tail wound model, we do not think that increased

migration is likely. Similarly, an increase in coagulation 'trapping' thrombocytes at the site of

infection is not supported by our experiments in *Tg(fabp10a:fgb-EGFP)* transgenic embryos.

We therefore propose that this 'increase' actually represents a decrease in thrombocyte

activation. Previous research has shown that during infection activated platelets can undergo

phagocytosis and/or apoptosis (Feng et al., 2014; Gaertner et al., 2017; Hottz et al., 2014);

both of these processes would result in the loss of visible GFP+ thrombocytes from our

system.

Recent research has highlighted the important role of platelets as innate immune cells; they

are able to release anti-microbial peptides, pick up and 'bundle' bacteria, and initiate the

recruitment of other innate immune cells to sites of infection (Gaertner et al., 2017; Li et al.,

2017; Morrell et al., 2014). The low frequency at which direct thrombocyte-mycobacterial

interactions were observed in our study argues against thrombocytes having a significant role in

directly mediating immunity to mycobacterial infection. Our experiments demonstrating

macrophage-dependence for anti-platelet drug mediated control of infection and reduced lipid

accumulation in the granuloma, suggests infection-induced thrombocyte activation modulates

macrophage containment of infection.

Our study joins several *in vitro* studies that have shown platelets can induce macrophages to

have differential responses to bacteria or other pathologic stimuli. In the presence of

S. aureus, thrombin activated platelets induce macrophages to increase phagocytosis and

slow the growth of bacteria (Ali et al., 2017). Conversely, platelets can supress the

macrophage response to LPS (Ando et al., 2016), and induce macrophages to produce less

TNFa, more IL-10, and more IL-1ß in response to both BCG and *M. tuberculosis* (Feng et al.,

2014; Fox et al., 2018). Crucially it has been shown that platelets are necessary for the

formation of foam cells in the context of mycobacterial infection and atheroma (Feng et al.,

2014). Our experiments with both Oil Red O and TUNEL demonstrate that thrombocytes

induce similar pathways *in vivo* in response to *M. marinum*; directing macrophages towards

foam-cell formation and necrosis.

Here we used the zebrafish-*M.marinum* model to show that mycobacteria drive coagulation

and thrombosis through the formation of granulomas. We found that inhibition of

thrombocyte activation was able to reduce foam-cell formation and cell death within the

granuloma, leading to dramatically reduced bacterial burden. This is the first *in*

vivo experimental evidence that infection-induced platelet activation contributes to the

pathogenesis of mycobacterial infection. Our study identifies platelet activation as a potential

target for tuberculosis host-directed therapy.

Acknowledgements

Dr Kristina Jahn and Sydney Cytometry for assistance with imaging equipment; Garvan

Biological Testing Facility staff Ms Jennifer Brand, Mr Michael Pickering, Ms Rola Bazzi,

Dr Lucie Nedved and Dr Stephanie Allison at the Garvan Institute of Medical Research for

maintenance of zebrafish breeding stock; Professors Lalita Ramakrishnan, Shaun Jackson and

Georges Grau, Associate Professor Carl Feng and Dr Jorn Coers for helpful discussion of

results.

This work was supported by the Australian National Health and Medical Research Council

APP1099912 and APP1053407; University of Sydney Fellowship; NSW Ministry of Health

under the NSW Health Early-Mid Career Fellowships Scheme; and the Kenyon Family

Inflammation Award (S.H.O.), Duke Summer Research Opportunities Program (K.J.), NIH

Director's New Innovator Award 1DP2-OD008614 (D.M.T), NIH R01-HL124232 and R01-

HL125774 (J.A.S.).

Author contributions

E.H., D.M.T. and S.H.O designed the experiments. E.H., K.E.J., M.D.J., T.N. and S.H.O

performed the experiments. J.A.S. generated transgenic and mutant zebrafish lines. E.H. and

S.O. wrote the paper. W.J.B., D.M.T. and S.H.O. supervised the project.

Declaration of Interests

The authors declare no competing interests.

References

Ali, R.A., Wuescher, L.M., Dona, K.R., and Worth, R.G. (2017). Platelets Mediate Host

Defense against Staphylococcus aureus through Direct Bactericidal Activity and by

Enhancing Macrophage Activities. J Immunol *198*, 344-351.

Ando, Y., Oku, T., and Tsuji, T. (2016). Platelets attenuate production of cytokines and nitric

oxide by macrophages in response to bacterial endotoxin. Platelets *27*, 344-350.

Cronan, M.R., Beerman, R.W., Rosenberg, A.F., Saelens, J.W., Johnson, M.G., Oehlers,

S.H., Sisk, D.M., Jurcic Smith, K.L., Medvitz, N.A., Miller, S.E.*, et al.* (2016). Macrophage

Epithelial Reprogramming Underlies Mycobacterial Granuloma Formation and Promotes

Infection. Immunity *45*, 861-876.

Feng, Y., Dorhoi, A., Mollenkopf, H.J., Yin, H., Dong, Z., Mao, L., Zhou, J., Bi, A., Weber,

S., Maertzdorf, J.*, et al.* (2014). Platelets direct monocyte differentiation into epithelioid-like

multinucleated giant foam cells with suppressive capacity upon mycobacterial stimulation. J

Infect Dis *210*, 1700-1710.

Fox, K.A., Kirwan, D.E., Whittington, A.M., Krishnan, N., Robertson, B.D., Gilman, R.H.,

Lopez, J.W., Singh, S., Porter, J.C., and Friedland, J.S. (2018). Platelets Regulate Pulmonary

Inflammation and Tissue Destruction in Tuberculosis. Am J Respir Crit Care Med.

Gaertner, F., Ahmad, Z., Rosenberger, G., Fan, S., Nicolai, L., Busch, B., Yavuz, G.,

Luckner, M., Ishikawa-Ankerhold, H., Hennel, R._, et al._ (2017). Migrating Platelets Are

Mechano-scavengers that Collect and Bundle Bacteria. Cell *171*, 1368-1382 e1323.

Hawn, T.R., Matheson, A.I., Maley, S.N., and Vandal, O. (2013). Host-directed therapeutics

for tuberculosis: can we harness the host? Microbiol Mol Biol Rev *77*, 608-627.

Hottz, E.D., Medeiros-de-Moraes, I.M., Vieira-de-Abreu, A., de Assis, E.F., Vals-de-Souza,

R., Castro-Faria-Neto, H.C., Weyrich, A.S., Zimmerman, G.A., Bozza, F.A., and Bozza, P.T.

(2014). Platelet activation and apoptosis modulate monocyte inflammatory responses in

dengue. J Immunol *193*, 1864-1872.

Johansen, M.D., Kasparian, J.A., Hortle, E., Britton, W.J., Purdie, A.C., and Oehlers, S.H.

(2018). Mycobacterium marinum infection drives foam cell differentiation in zebrafish

infection models. bioRxiv.

Kalantri, A., and Kalantri, S. (2018). Can aspirin help? Elife *7*.

Kothari, H., Rao, L.V., Vankayalapati, R., and Pendurthi, U.R. (2012). Mycobacterium

tuberculosis infection and tissue factor expression in macrophages. PLoS ONE *7*, e45700.

Kroesen, V.M., Groschel, M.I., Martinson, N., Zumla, A., Maeurer, M., van der Werf, T.S.,

and Vilaplana, C. (2017). Non-Steroidal Anti-inflammatory Drugs As Host-Directed Therapy

for Tuberculosis: A Systematic Review. Front Immunol *8*, 772.

Kroesen, V.M., Rodriguez-Martinez, P., Garcia, E., Rosales, Y., Diaz, J., Martin-Cespedes,

M., Tapia, G., Sarrias, M.R., Cardona, P.J., and Vilaplana, C. (2018). A Beneficial Effect of

Low-Dose Aspirin in a Murine Model of Active Tuberculosis. Front Immunol *9*, 798.

Kutiyal, A.S., Gupta, N., Garg, S., and Hira, H.S. (2017). A Study of Haematological and

Haemostasis Parameters and Hypercoagulable State in Tuberculosis Patients in Northern

India and the Outcome with Anti-Tubercular Therapy. Journal of clinical and diagnostic

research : JCDR *11*, OC09-OC13.

Li, J.L., Zarbock, A., and Hidalgo, A. (2017). Platelets as autonomous drones for hemostatic

and immune surveillance. J Exp Med.

Lin, H.F., Traver, D., Zhu, H., Dooley, K., Paw, B.H., Zon, L.I., and Handin, R.I. (2005).

Analysis of thrombocyte development in CD41-GFP transgenic zebrafish. Blood *106*, 3803-

3810.

Mai, N.T., Dobbs, N., Phu, N.H., Colas, R.A., Thao, L.T., Thuong, N.T., Nghia, H.D., Hanh,

N.H., Hang, N.T., Heemskerk, A.D.*, et al.* (2018). A randomised double blind placebo

controlled phase 2 trial of adjunctive aspirin for tuberculous meningitis in HIV-uninfected

adults. eLife *7*.

Matty, M.A., Oehlers, S.H., and Tobin, D.M. (2016). Live Imaging of Host-Pathogen

Interactions in Zebrafish Larvae. Methods Mol Biol *1451*, 207-223.

Morrell, C.N., Aggrey, A.A., Chapman, L.M., and Modjeski, K.L. (2014). Emerging roles for

platelets as immune and inflammatory cells. Blood *123*, 2759-2767.

Oehlers, S.H., Cronan, M.R., Beerman, R.W., Johnson, M.G., Huang, J., Kontos, C.D., Stout,

J.E., and Tobin, D.M. (2017). Infection-induced vascular permeability aids mycobacterial

growth. J Infect Dis *215*, 813-817.

Oehlers, S.H., Cronan, M.R., Scott, N.R., Thomas, M.I., Okuda, K.S., Walton, E.M.,

Beerman, R.W., Crosier, P.S., and Tobin, D.M. (2015). Interception of host angiogenic

signalling limits mycobacterial growth. Nature *517*, 612-615.

Passeri, M.J., Cinaroglu, A., Gao, C., and Sadler, K.C. (2009). Hepatic steatosis in response

to acute alcohol exposure in zebrafish requires sterol regulatory element binding protein

activation. Hepatology *49*, 443-452.

Robson, S.C., White, N.W., Aronson, I., Woollgar, R., Goodman, H., and Jacobs, P. (1996).

Acute-phase response and the hypercoagulable state in pulmonary tuberculosis. Br J

Haematol *93*, 943-949.

Russell, D.G., Cardona, P.J., Kim, M.J., Allain, S., and Altare, F. (2009). Foamy

macrophages and the progression of the human tuberculosis granuloma. Nat Immunol *10*,

943-948.

Venkatasubramanian, S., Tripathi, D., Tucker, T., Paidipally, P., Cheekatla, S., Welch, E.,

Raghunath, A., Jeffers, A., Tvinnereim, A.R., Schechter, M.E.*, et al.* (2016). Tissue factor

expression by myeloid cells contributes to protective immune response against

Mycobacterium tuberculosis infection. Eur J Immunol *46*, 464-479.

Vo, A.H., Swaroop, A., Liu, Y., Norris, Z.G., and Shavit, J.A. (2013). Loss of fibrinogen in

zebrafish results in symptoms consistent with human hypofibrinogenemia. PLoS ONE *8*,

e74682.

Walton, E.M., Cronan, M.R., Beerman, R.W., and Tobin, D.M. (2015). The Macrophage-

Specific Promoter mfap4 Allows Live, Long-Term Analysis of Macrophage Behavior during

Mycobacterial Infection in Zebrafish. PLoS ONE *10*, e0138949.

World Health Organization., and Global Tuberculosis Programme. Global tuberculosis

control : WHO report. (Geneva, Global Tuberculosis Programme), p. 15 volumes.

Figure legends

Figure 1: *Mycobacterium marinum* induces coagulation and thrombosis around sites of

infection in zebrafish

A) Representative images of a *Tg(fabp10a:fgb-EGFP)* embryo infected with *M.marinum*-tdTomato by caudal vein injection, showing clot formation (green) at sites of infection (red) at 3, 4, and 5 days post infection. B) Representative images of 5 DPI *Tg(fabp10a:fgb-EGFP)* embryos infected with *M.marinum*-tdTomato WT and ΔESX1 *M. marinum*-td-tomato (ΔESX1) showing clot formation (green) at sites of infection (red). Arrows indicate clotting. C) Quantification of clot formation in burden-matched ΔESX1 mutant-infected *Tg(fabp10a:fgb-EGFP)* embryos normalised to WT *M. marinum* control. Error lines represent mean ± SEM, statistical analysis by T test. Data show combined results from two independent experiments. D) Representative images of *Tg(fabp10a:fgb-EGFP)* embryos, where clot formation can be visualised by GFP fluorescence, infected with *M.marinum*-tdTomato (red) and treated with either DMSO or warfarin. E) Quantification of clotting in warfarin-treated *Tg(fabp10a:fgb-EGFP)* embryos. F) Quantification of bacterial burden in embryos treated with warfarin. Data are combined results of two independent experiments. F) Representative 2DPI, 3DPI and 4DPI images of *Tg(cd41:GFP)* fish infected with *M.marinum*-tdTomato at 3DPF. White arrow heads show thrombocyte (green) association with areas of bacterial growth (red). Thrombocytes not indicated with an arrow head were circulating and not considered to be associated with bacteria (more detail available in Figure S2C).

Figure 2: Anti-platelet drugs reduce bacterial burden in *M. marinum* infection

A) Quantification of bacterial burden in embryos treated with aspirin normalised to DMSO control. Data are combined results of two independent experiments. B) Quantification of bacterial burden in embryos treated with tirofiban normalised to DMSO control. Data are combined results of two independent experiments. C) Quantification of bacterial burden in embryos treated with eptifibatide or DMSO control. D) Representative images of

Tg(cd41:gfp) embryos treated with tirofiban or DMSO control. E) Quantification of total

thrombocyte *Tg*(*cd41:GFP*) area within the tail of infected larvae treated with tirofiban. F)

Quantification of bacterial growth by relative fluorescence in 7H9 broth culture

supplemented with drugs as indicated. G) Survival analysis of *P. aerogenosa* PA01 infected

embryos treated with anti-platelet drugs as indicated.

All graphs show Mean ± SEM. Statistical analyses performed by T tests and Log-Rank tests

where appropriate.

Figure 3: The anti-bacterial effect of anti-platelet drugs is thrombocyte dependent

A) Total fluorescence area of *M. marinum* bacteria in larvae injected with either control or

cmpl morpholino (MO) to deplete thrombocytes, and then treated with tirofiban (Tiro).

Values are normalised to DMSO treated control MO larvae. Graphs show the combined

results of 2 independent experiments. B) Quantification of bacterial burden in in *itga2b*

mutant embryos normalised to WT control. Data are combined results of four independent

experiments. C) Quantification of bacterial burden in embryos treated with tirofiban,

warfarin, or tirofiban and warfarin, normalised to DMSO control. Data are combined results

of two independent experiments. D) Quantification of bacterial burden in *fga*[-/-] embryos

treated with tirofiban. E) Representative images of 5 DPI *Tg(fabp10a:fgb-EGFP)* embryos

infected with *M.marinum*-tdTomato and treated with either DMSO or tirofiban. F)

Quantification of clotting relative to bacterial burden in embryos treated with tirofiban

normalised to DMSO control. Data are combined results of two independent experiments.

All graphs show Mean ± SEM. Statistical analyses performed by T tests or ANOVA where

appropriate.

Figure 4: Thrombocyte activation compromises immunity through granuloma-associated

macrophages.

A) Quantification of bacterial burden in embryos infected with ΔESX1 *M.marinum*-tdTomato

and treated with tirofiban. B) Quantification of bacterial burden at 3 DPI after continuous

tirofiban treatment from 0 DPI. C) Quantification of bacterial burden at 5 DPI after overnight

drug treatment initiated at 4 DPI. D) Quantification of bacterial burden in embryos infected

with *M.marinum*-tdTomato, injected with clodronate liposomes and treated with tirofiban

from at 3 DPI. E) Representative images of bacterial granulomas chosen for analysis

(bacteria are white in greyscale images), and corresponding Oil Red O (ORO) staining (red-

brown in color images). F) Quantification of ORO pixel density relative to granuloma

bacterial area, in embryos treated with tirofiban, normalised to DMSO control. G)

Representative images of bacterial granulomas showing bacteria in red and TUNEL staining

in green. F) Quantification of TUNEL positive area within the largest granuloma of

individual embryos.

All graphs show Mean ± SEM, statistical tests by T-tests. Data are combined results of two

independent experiments, except D) and H) which represent single experiments.

Graphical abstract.

M. marinum indicues two independent haemostatic processes in the zebrafish: ESX1

dependent coagulation (1), and ESX1 independent thrombocyte recruitment (2). Bacteria then

induce ESX1 dependent thrombocyte activation - either directly or via the macrophage - (3),

which leads to foam cell formation and macrophage necrosis, promoting bacterial growth (4).

Inhibition of thrombocyte activation with anti-platelet drugs like Tirofiban, stops this

progression and thereby limits the growth of bacteria. Drugs which reduce coagulation (1)

have no effect on bacterial growth.

Supplementary Figure Legends

Figure S1

Related to Figure 1. Quantification of clot formation in warfarin-treated embryos infected

with *M. marinum*. Error lines represent mean ± SEM, statistical analysis by T test.

Figure S2

Related to Figure 1. A) Representative images of 5 DPI *Tg(cd41:GFP)* embryos infected

with *M.marinum*-tdTomato, showing thrombocyte accumulation (green) at sites of bacterial

infection (red). B) Quantification of total thrombocyte *Tg(cd41:GFP)* area within the tail of

uninfected or WT *M.marinum*-tdTomato infected embryos. C) Correlation between *M.*

marinum bacterial burden and total thrombocyte *Tg(cd41:GFP)* area within the tail of

infected embryos. D) Quantification of total thrombocyte *Tg(cd41:GFP)* area within the tail

of uninfected or ΔESX1 *M.marinum*-tdTomato infected larvae at 5 DPI. E) Correlation

between ΔESX1 *M. marinum* bacterial burden and total thrombocyte *Tg (cd41:GFP)* area

within the tail of infected embryos at 5 DPI.

Figure S3

Related to Figure 1. Images of 2 DPI, 3 DPI and 4 DPI *Tg(cd41:GFP)* embryos infected with

M.marinum-tdTomato at 3 DPF. At each time-point, 2 images were taken ~10 seconds apart.

Thrombocytes located in the same position in both images were considered stationary (non-

circulating), and labelled with a white arrow head.

Graphs show Mean ± SEM, statistical testing by T-test or linear regression.

Figure S4

Related to Figure 2. A) Representative images of thrombocyte accumulation at tail wound in

4 DPF *Tg(cd41:GFP)* embryos , 4 hours post injury (hpi). B) Quantification of thrombocytes

within 100 µm of the cut site 4 hpi in aspirin-treated embryos. C) Quantification of

thrombocytes within 100 µm of the cut site 4 hpi in tirofiban-treated embryos.

Graphs show Mean ± SEM, statistical testing by T-test (B and C) or ANOVA (D).

Figure S5

Related to Figure 3. A) Quantification of thrombocytes within 100 µm of the cut site 4 hours

after injury in *itga2b* WT, heterozygous and knock-out embryos. B) Representative images of

Tg(cd41:gfp) itga2b mutant embryos infected with *M. marinum*-tdTomato. C) Quantification

of total thrombocyte *Tg*(*cd41:GFP)* area within the tail of infected *itga2b* mutant embryos.

Figure S6

Related to Figure 4. A) Quantification of bacterial burden in embryos infected with a low

dose of 10-50 fluorescent *M. marinum* and treated with tirofiban. **B)** Representative images

of caudal haematopoietic tissue in 6 DPF *Tg(mfap4:tdTomato)* embryos, where macrophages

are fluorescently labelled, injected with clodronate or PBS liposomes at 4 DPF. **C)**

Quantification of macrophage number by fluorescent pixel count at 5 DPI in embryos

injected with clodronate or PBS liposomes at 3 DPI, and treated with tirofiban. Graphs show

Mean ± SEM, statistical tests by T-tests (A and B) or ANOVA (D).

Methods

Zebrafish husbandry

Adult zebrafish were housed at the Garvan Institute of Medical Research Biological Testing

Facility (St Vincent's Hospital AWC Approval 1511) and embryos were produced by natural

spawning for infection experiments at the Centenary Institute (Sydney Local Health District

AWC Approval 2016-022). Zebrafish embryos were obtained by natural spawning and

embryos were raised at 28°C in E3 media.

Zebrafish lines

Wild type zebrafish are the TAB background. Transgenic lines are: *Tg(fabp10a:fgb-*

EGFP)^{mi4001} referred to as *Tg(fabp10a:fgb-EGFP)(Vo et al., 2013)*, *Tg(-6.0itga2b:eGFP)^{la2}*

referred to as *Tg(cd41:EGFP)(Lin et al., 2005)*, *Tg(mfap4:tdTomato)^{xt12}* referred to as

Tg(mfap4:tdTomato)(Walton et al., 2015), Mutant allele *fga^{mi}* contains a 26 bp insertion in

the *fibrinogen alpha chain* gene (manuscript in preparation).

Infection of zebrafish embryos

Aliquots of single cell suspensions of midlog-phase *Mycobacterium marinum* M strain,

ΔESX1 *M. marinum* and *Pseudomonas aeruginosa* PA01 were frozen at -80°C for use in

infection experiments. Bacterial aliquots were thawed and diluted with phenol red dye (0.5%

w/v). 10-15 nL was injected into the caudal vein or trunk of M-222 (tricaine)-anaesthetised

30-48 hpf embryos resulting in a standard infectious dose ~400 fluorescent *M. marinum*.

Embryos were recovered into E3 supplemented with 0.036 g/L PTU, housed at 28 °C and

imaged on day 5 of infection unless otherwise stated.

Drug treatments

Embryos were treated with vehicle control (DMSO or water as appropriate), 10 µg/ml

aspirin, 20 µg/ml tirofiban, 10 µM eptifibatide, or 5 µM warfarin. Drugs and E3 were

replaced on days 0, 2, and 4 days post infection (DPI) unless otherwise stated.

Tail wound thrombosis assay

3 day post fertilisation (DPF) embryos were treated over-night with anti-platelet drugs. They

were anaesthetised, and then a small amount of their tail was removed with a scalpel.

Embryos were imaged 4 hours post wounding and the number of GFP positive cells within

100 µm of the cut site was counted.

Imaging

Live zebrafish embryos were anaesthetized in M-222 (Tricaine) and mounted in 3%

methylcellulose for imaging on a Leica M205FA or DM6000B fluorescence

stereomicroscope. Image analysis was carried out with Image J Software Version 1.51j using

fluorescent pixel counts and intensity measurements as previously described (Matty et al.,

2016).

Axenic culture

A midlog culture of fluorescent *M. marinum* was diluted 1:100 and aliquoted into 96 well

plates for drug treatment. Cultures were maintained at 28°C in a static incubator and bacterial

fluorescence was measured in a plate reader.

Morpholinos:

Embryos were injected at the single cell stage with 1 pmol

cMPL (5′-CAGAACTCTCACCCTTCAATTATAT-3′),

or control morpholino (5′-CCTCTTACCTCAGTTACAATTTATA-3′).

Clodronate liposome injections:

Larvae were injected at 3 DPI (4 DPF) with 10 nl of 5 mg/ml clodronate liposomes or 5

mg/ml PBS vehicle liposomes by caudal vein injection.

Oil-red O:

Oil Red O lipid staining on whole mount embryos was performed and analysed as previously

described (Johansen et al., 2018; Passeri et al., 2009). Briefly, embryos were individually

imaged for bacterial distribution by fluorescent microscopy, fixed, and stained in Oil Red O

(0.5% w/v in propylene glycol). Oil Red O density was calculated by using the 'measure'

function in Image J, and subtracting the mean brightness of a representative region within

each granuloma from the mean brightness of a representative adjacent 'background' region.

Statistics

All t-tests were unpaired t-tests with Welch's correction. All ANOVA were ordinary one-way

ANOVA, comparing the mean of each group with the mean of every other group, using

Turkey's multiple comparisons test with a single pooled variance. In cases where data was

pooled from multiple experiments, data from each was normalised to its own

within-experiment control (usually 'DMSO') before pooling. Outliers were removed using

ROUT, with Q=1%.

A

3DPI   4DPI   5DPI

M. marinum FGB

B

WT

Bacteria FGB

ΔESX1

C

0.0200

Normalised FGB-GFP Area

M.marinum WT   *M. marinum* ΔESX1

D

DMSO

Warfarin

M. marinum FGB

E

0.3129

Normalised Total Bacteria Area

DMSO   Warfarin

F

M. marinum cd41:gfp

2 DPI   3 DPI   5 DPI

Thrombocytes

②

M. marinum

Macrophage

Tirofiban

ESX1

Activated thrombocytes

ESX1

③

①

Fibrinogen clot

④

Necrotic, foamy macrophage

Fish

1 2 3 4 5 6 7 8 9 10 11 12 13 14

2DPI

3DPI

4DPI

A

B