Thrombocyte inhibition restores protective immunity to mycobacterial

infection in zebrafish

Short title:

Thrombocytes compromise immunity to mycobacteria

Elinor Hortle[1], Khelsey E. Johnson[2], Matt D. Johansen[1], Tuong Nguyen[1], Jordan A. Shavit[3],

Warwick J. Britton[1,4], David M. Tobin[2], Stefan H. Oehlers[1,4] *

[1] Tuberculosis Research Program, Centenary Institute, University of Sydney, Camperdown,

NSW 2050, Australia

[2] Department of Molecular Genetics and Microbiology, Duke University School of Medicine,

Durham, NC 27710, USA

[3] Department of Pediatrics and Cellular and Molecular Biology Program, University of

Michigan, Ann Arbor, MI 48107, USA

[4] The University of Sydney, Central Clinical School and Marie Bashir Institute,

Camperdown, NSW 2050, Australia

* Corresponding author email: stefan.oehlers@sydney.edu.au, Twitter: @oehlerslab

Key Points

1) Inhibition of thrombocyte activation improves control of mycobacterial infection.

2) Inhibition of thrombocyte activation reduces thrombocyte-macrophage interactions and improves indices of macrophage immune function against mycobacterial infection.

Abstract

Infection-induced thrombocytosis is a clinically important complication of tuberculosis (TB). Recent studies have separately highlighted a correlation of platelet activation with TB severity and utility of aspirin as a host-directed therapy for TB that modulates the inflammatory response. Here we investigate the possibility that the beneficial effects of aspirin are related to an anti-platelet mode of action. We utilize the zebrafish-*Mycobacterium marinum* model to show mycobacteria drive host hemostasis through the formation of granulomas. Treatment of infected zebrafish with aspirin or platelet-specific glycoprotein IIb/IIIa inhibitors reduced mycobacterial burden demonstrating a detrimental role for infection-induced thrombocyte activation. We found platelet inhibition reduced thrombocyte-macrophage interactions and restored indices of macrophage-mediated immunity to mycobacterial infection. Pathological thrombocyte activation and granuloma formation were found to be intrinsically linked illustrating a bidirectional relationship between host hemostasis and TB pathogenesis. Our study illuminates platelet activation as an efficacious target of anti-platelets drugs including aspirin, a widely available and affordable host-directed therapy candidate for tuberculosis.

Keywords

Mycobacterial infection, haemostasis, innate immunity, clotting

Introduction

Mycobacterium tuberculosis is the world's most lethal pathogen, causing nearly 2 million

deaths each year [1]. The increasing incidence of both multi- and extremely-drug resistant

tuberculosis (TB) urgently require the development of therapeutics that overcome the

shortcomings of conventional antibiotics. Pathogenic mycobacteria co-opt numerous host

pathways to establish persistent infection, and subversion of these interactions with host-

directed therapies (HDTs) has been shown to reduce the severity of infection in animal

models. For example, we have recently shown that mycobacteria induce host angiogenesis

and increase host vascular permeability; blockade of either of these processes reduced both

the growth and spread of bacteria [2,3]. Therefore, host processes co-opted by the bacteria

provide attractive targets for novel TB treatments. One such pathway may be hemostasis.

Thrombocytosis has long been recognized as a biomarker for advanced TB, and infection is

often accompanied by the induction of a hyper-coagulable state, resulting in increased risk of

deep vein thrombosis and stroke [4,5]. Recent evidence hints that mycobacteria may drive this

process, and that it may aid their growth. For example, cell wall components from *M.*

tuberculosis can induce expression of tissue factor - an important activator of coagulation - in

macrophages [6]. In mice and humans markers of platelet activation are upregulated during *M.*

tuberculosis infection [7,8], and it has been shown *in vitro* that interaction with activated

platelets increases the conversion of infected macrophages into cells permissive for bacterial

growth [7,9]. To date the pathogenic roles of hemostasis have not been studied in an intact *in*

vivo model of mycobacterial infection.

Here we used the zebrafish-*M. marinum* model to investigate the role of host thrombocytes in

mycobacterial infection. We present evidence that while coagulation and thrombocyte

activation are both driven by mycobacteria, it is only infection-induced activation of

thrombocytes that specifically compromises protective immunity through direct thrombocyte-

macrophage interactions.

Methods

Zebrafish husbandry

Adult zebrafish were housed at the Garvan Institute of Medical Research Biological Testing

Facility (St Vincent's Hospital AWC Approval 1511) and embryos were produced by natural

spawning for infection experiments at the Centenary Institute (Sydney Local Health District

AWC Approval 2016-022). Zebrafish embryos were obtained by natural spawning and

embryos were raised at 28°C in E3 media.

Zebrafish lines

Wild type zebrafish are the TAB background. Transgenic lines are: *Tg(fabp10a:fgb-*

EGFP)^{mi4001} referred to as *Tg(fabp10a:fgb-EGFP)*[10], *Tg(-6.0itga2b:eGFP)^{la2}* referred to as

Tg(cd41:EGFP)[11], *Tg(mfap4:tdTomato)^{xt12}* referred to as *Tg(mfap4:tdTomato)*[12], Mutant

allele *fga^{mi}* contains a 26 bp insertion in the *fibrinogen alpha chain* gene (manuscript in

preparation).

Infection of zebrafish embryos

Aliquots of single cell suspensions of midlog-phase *Mycobacterium marinum* M strain and

ΔESX1 *M. marinum* were frozen at -80°C for use in infection experiments. Bacterial aliquots

were thawed and diluted with phenol red dye (0.5% w/v). 10-15 nL was injected into the

caudal vein or trunk of M-222 (tricaine)-anaesthetized 30-48 hpf embryos resulting in a

standard infectious dose ~400 fluorescent *M. marinum*. Embryos were recovered into E3

supplemented with 0.036 g/L PTU, housed at 28 °C and imaged on day 5 of infection unless

otherwise stated.

Drug treatments

Embryos were treated with vehicle control (DMSO or water as appropriate), 10 µg/ml

aspirin, 20 µg/ml tirofiban, 10 µM eptifibatide, or 5 µM warfarin. Drugs and E3 were

replaced on days 0, 2, and 4 days post infection (DPI) unless otherwise stated.

Tail wound thrombosis assay

Three day post fertilization (DPF) embryos were treated over-night with anti-platelet drugs.

They were anaesthetized, and then a small amount of their tail was removed with a scalpel.

Embryos were imaged 4 hours post wounding and the number of GFP positive cells within

100 µm of the cut site was counted.

Imaging

Live zebrafish embryos were anaesthetized in M-222 (Tricaine) and mounted in 3%

methylcellulose for static imaging on a Leica M205FA or DM6000B fluorescence

stereomicroscope. Image analysis was carried out with Image J Software Version 1.51j using

fluorescent pixel counts and intensity measurements as previously described [13].

Video and timelapse imaging was carried out on anaesthetized embryos mounted in 0.75%

low melting point agarose on a Leica M205FA or Deltavision Elite fluorescence microscope.

Video editing was carried out with Image J Software Version 1.51j and iMovie.

Axenic culture

A midlog culture of fluorescent *M. marinum* was diluted 1:100 and aliquoted into 96 well

plates for drug treatment. Cultures were maintained at 28°C in a static incubator and bacterial

fluorescence was measured in a plate reader.

Morpholinos:

Embryos   were   injected   at   the   single   cell   stage   with   1   pmol

cMPL (5′-CAGAACTCTCACCCTTCAATTATAT-3′),

or control morpholino (5′-CCTCTTACCTCAGTTACAATTTATA-3′).

Clodronate liposome injections:

Larvae were injected at 3 DPI (4 DPF) with 10 nl of 5 mg/ml clodronate liposomes or 5

mg/ml PBS vehicle liposomes by caudal vein injection.

Oil-red O:

Oil Red O lipid staining on whole mount embryos was performed and analyzed as previously

described [14,15]. Briefly, embryos were individually imaged for bacterial distribution by

fluorescent microscopy, fixed, and stained in Oil Red O (0.5% w/v in propylene glycol). Oil

Red O density was calculated by using the 'measure' function in Image J, and subtracting the

mean brightness of a representative region within each granuloma from the mean brightness

of a representative adjacent 'background' region.

Statistics

All t-tests were unpaired t-tests with Welch's correction. All ANOVA were ordinary one-way

ANOVA, comparing the mean of each group with the mean of every other group, using

Turkey's multiple comparisons test with a single pooled variance. In cases where data was

pooled from multiple experiments, data from each was normalized to its own

within-experiment control (usually 'DMSO') before pooling. Outliers were removed using

ROUT, with Q=1%.

Results

***Mycobacterium marinum* infection induces thrombocytosis in zebrafish.**

To determine if *M. marinum* induces thrombocytosis in zebrafish, we infected *Tg(cd41:GFP)*

embryos, where thrombocytes are marked by GFP expression, with red fluorescent

M. marinum. Infected embryos had significantly increased density of thrombocytes around

the tail venous plexus where granulomas preferentially form (Figures 1A-B). Amongst

infected embryos, there was a strong positive correlation between thrombocyte density and

mycobacterial burden (Figure 1C).

Because the *cd41* promoter is active in non-motile thrombocyte precursors within their

caudal hematopoietic tissue, we could not conclusively determine if these thrombocytes had

actively migrated to and been retained at the site of infection [11]. To determine if zebrafish

thrombocytes are recruited to sites of mycobacterial infection, we performed trunk injection

of *M. marinum* in *Tg(cd41:GFP)* at 3 days post fertilization (DPF), a time point after which

mature thrombocytes are in the circulation. Embryos were then imaged at 2, 3, and 4 DPI.

Rather than forming a stable and growing clot over a period of days, thrombocytes appeared

to form transient associations with sites of infection, and new thrombocytes seemed to be

retained at sites of infection in different locations each day (Figure 1D). We therefore

recorded videos of *Tg(cd41:GFP)* embryos infected with *M. marinum*-tomato using a long

pass GFP filter to capture bacterial and thrombocyte fluorescence simultaneously.

Thrombocytes were most often observed on the edges of granulomas consistent with the

location of granuloma-defining macrophages [16]. They also formed short associations with

sites of infection, sometimes lasting only 5-10 minutes (Supplementary Video 1). Therefore

thrombocyte-granuloma interactions appear to be a conserved feature of mycobacterial

infection across species.

Anti-platelet drugs reduce mycobacterial burden in zebrafish.

It has previously been reported that aspirin has a host-protective effect during TB infection [17-

20]. Most of these studies have focused on the fact that aspirin is a broadly acting nonsteroidal

anti-inflammatory drug (NSAID) that is known to modulate infection-relevant prostaglandin

metabolism [21]. However, aspirin is also a widely-used platelet inhibitor, and we theorized this

capacity may also play a role in the drug's effectiveness against TB.

To test this hypothesis we first confirmed that aspirin's protective effect was seen across

species, by treating *M. marinum*-infected fish with aspirin by immersion. Mycobacterial

burden was reduced by approximately 50% in aspirin-treated embryos (Figure 2A).

To determine if the anti-platelet effects of aspirin treatment contribute to the reduced

mycobacterial burden, we treated *M. marinum*-infected fish with the platelet-specific, small

molecule glycoprotein IIb/IIIa inhibitors, tirofiban or eptifibatide. These drugs do not inhibit

platelet activation and de-granulation, but rather inhibit activated platelets from binding to

one-another, and to monocytes, via fibrinogen. Treatment with either glycoprotein IIb/IIIa

inhibitor phenocopied aspirin by reducing bacterial burden providing direct evidence of a

pathological role for thrombocyte activation in the immune response to mycobacterial

infection (Figure 2B-C).

We next examined the cellular target of anti-platelet drugs in our infection system. We

performed antibacterial testing of the anti-platelet drugs in axenic cultures of *M. marinum*

and did not observe any effect on bacterial growth *in vitro* demonstrating host-dependent

activity (Figure 2D).

The anti-bacterial effect of anti-platelet drugs is thrombocyte dependent.

We next confirmed that anti-platelet drugs inhibit thrombocytes in zebrafish using a tail wound

thrombosis assay (Figure S1A). Anti-platelet drugs reduced the number of thrombocytes

recruited to tail wound clots demonstrating conservation of their cellular target in zebrafish

embryos (Figure S1B-C).

To determine if zebrafish thrombocytes are the conserved target for anti-platelet drugs in the

zebrafish-*M. marinum* infection model, we inhibited thrombopoiesis by injection with a

morpholino against the thrombopoietin receptor *cmpl* [11]. Inhibition of thrombopoiesis did not

affect the outcome of infection. However, both aspirin and tirofiban treatment failed to

reduce bacterial burden in thrombocyte-depleted embryos, demonstrating thrombocytes are

the cellular target of this drug in the zebrafish infection model (Figure 3A-B).

Eptifibatide and tirofiban were designed as specific inhibitors to prevent binding between

Glycoprotein IIb/IIIa and fibrinogen in mammals [22], but nothing is known about their potential

off-target effects in the zebrafish. Therefore, to confirm that disruption of Glycoprotein IIb/IIIa

binding alone can reduce bacterial burden, we performed infection experiments in

Glycoprotein IIb/IIIa knock-out (KO) transgenic embryos (*itga2b* mutants). The *itga2b*^{sa10134}

allele caused a dose-dependent reduction in thrombocytes recruited to tail wound clots (Figure

S1D), and KO of *itga2b* significantly reduced bacterial burden (Figure 3C).

Similarly, when we when we addressed the same question using a *fibrinogen alpha chain* (fga)

mutant zebrafish line that does not produce mature fibrinogen, we saw that KO fish had

significantly reduced bacterial burden (Figure 3D). Furthermore, while tirofiban was able to

reduce bacterial burden in fga sufficient mutants, it did not reduce bacterial burden in fga KO.

Together these data suggest that tirofiban is reducing bacterial burden by inhibiting binding

between glycoprotein IIb/IIIa and fibrinogen.

Mycobacterium marinum induces coagulation in zebrafish but inhibition of coagulation

does not affect the outcome of infection.

Seeing that there was a marginal decrease in bacterial burden in *fga-/-* embryos compared to

WT and heterozygous clutchmates, we next sought to determine if *M. marinum* induces

coagulation in zebrafish and the role of coagulation in *M. marinum* infection. We infected

Tg(fabp10a:fgb-EGFP) transgenic embryos expressing EGFP-tagged fibrinogen beta (FGB)

with *M. marinum*-tdtomato, and imaged the developing infection every 15 minutes from 3

days post infection (DPI), until 6 DPI (Supplementary Video 2). We observed that clots

formed only at areas of bacterial growth, and that the size of the clots increased as the

number of bacteria increased over the course of the infection (Figure 4A). When we infected

fish with ΔESX1 mutant *M. marinum*, which lacks the ability to export key virulence proteins

and does not form granulomas, we observed significantly reduced clot formation (Figures

4B).

These data suggested clotting could be driven by mycobacterial and may thus contribute to

mycobacterial pathogenesis. To test this hypothesis, we treated infected embryos with the

anti-coagulant warfarin to prevent clot formation. We did not observe significant changes in

bacterial burden, suggesting coagulation itself does not affect bacterial growth within the host

(Figures 4C-D). Together, these data demonstrate that while coagulation is a consequence of

infection driven by a conserved mycobacterial pathogenicity program across host species, the

effects of coagulation on mycobacterial pathogenesis vary between host species [6].

Thrombocytes increase mycobacterial burden independently of coagulation.

To assess the contribution of thrombocytes to infection-induced coagulation, we analyzed the

formation of FGB-GFP clots in tirofiban-treated *Tg(fabp10a:fgb-EGFP)* embryos. Tirofiban

visibly reduced total clot formation (Figure 5A). However, correction for relative bacterial

burden suggested that the reduced clot formation was burden-dependent and thrombus

formation was not additionally impacted by tirofiban treatment (Figure 5B). Therefore, we

hypothesized that tirofiban was reducing bacterial burden independently of infection-induced

coagulation. To investigate this hypothesis, we again used warfarin, which prevented clot

formation during infection and did not affect bacterial burden (Figure 4C-D and 5C). As

expected, the addition of warfarin to our tirofiban treatment model had no effect on the ability of

tirofiban to reduce bacterial burden (Figure 5D), indicating that tirofiban acts through an

independent process. This suggests that the protective effect of tirofiban occurs independently of

fibrin clot formation, but requires the presence of soluble fibrinogen.

Thrombocytes compromise immunity through physical interactions with granuloma macrophages.

To investigate the effect of glycoprotein IIb/IIIa inhibitors on thrombocyte-granuloma interactions, we first measured the effect of tirofiban on the infection-induced thrombocytosis phenotype. Surprisingly, tirofiban treatment increased thrombocytosis compared to the untreated infected group (Figure 6A). Increased thrombocyte density was also observed in infections with ΔESX1 mutant *M. marinum*, that are unable to drive granuloma maturation or necrosis, (Figure 6B), yet tirofiban treatment did not reduce ΔESX1 *M. marinum* burden (Figure 6C), further suggesting that the infection-induced thrombocytosis phenotype is independent of granuloma immunity.

Given that glycoprotein IIb/IIIa-mediated platelet-monocyte binding can occur via fibrinogen [23,24] and patients with pulmonary tuberculosis have been shown to have significantly increased platelet-monocyte aggregation [25], we next aimed to determine whether anti-platelet drugs disrupted the interaction of thrombocytes and macrophages *in situ* within our model by live imaging. We infected double transgenic *Tg(cd41:GFP; mfap4:tdtomato)* with *M. marinum*-cerulean and performed timelapse imaging of 5 DPI embryos. Thrombocytes formed transient associations with *M. marinum*-infected macrophages (Supplementary Video 3), the frequency of which were increased upon eptifibatide treatment consistent with the increased thrombocytosis seen in Figure 6A (Supplementary Video 4), with an average dwell time of 30 minutes in untreated controls, and 80 minutes in eptifibatide-treated embryos (Figure 6D). We also observed a significant increase in the distance travelled by thrombocytes associated with *M. marinum*-infected macrophages in eptifibatide-treated embryos (Figure 6E). Thrombocyte

tracks in control embryos were most often short and straight (Figure 6F I-II, Supplementary

Video 5), maintaining a consistent distance from the center of the macrophage (Figure 6G III,

Supplementary Video 6). In contrast, thrombocyte tracks in eptifibatide-treated embryos were

long and circuitous, and varied in their distance from the center of the macrophage (Figure 6G

IV-VI, Supplementary Videos 7 and 8).

To demonstrate that thrombocyte inhibition exerts a protective effect through boosting

macrophage-dependent immunity, we depleted macrophages by injecting clodronate liposomes

to deplete macrophages early during granuloma formation at 3 DPI (Figure 6C). Macrophage-

depleted fish were unresponsive to tirofiban treatment demonstrating that pathological

thrombocyte activation promotes bacterial growth via interactions with macrophages (Figure

6D). Together these results demonstrate glycoprotein IIb/IIIa inhibitor treatment disrupt

pathological thrombocyte-macrophage attachments in mycobacterial granulomas.

Granuloma maturation and pathological thrombocyte activation have a bidirectional

relationship.

Our observation that infection with ΔESX1 mutant *M. marinum* did not result in pathological

thrombocyte activation (Figures 6C-D) suggested the existence of a bidirectional relationship

between granuloma maturation, which is deficient in ΔESX1 mutant infections, and pathological

thrombocyte activation. To further delineate this relationship we next took advantage of the

stereotypical progression of innate immune granulomas in zebrafish embryos and used the

burden reducing effect of tirofiban as a small molecule probe for thrombocyte activation. We

found that at 3 DPI, a time-point with nascent granuloma formation but prior to significant

granuloma organization and necrosis, tirofiban had no effect on bacterial burden (Figure 7A).

Conversely, treatment of established infections from 4 to 5 DPI, a time-point when granulomas

become organized and necrotic, tirofiban significantly reduced bacterial burden within 24 hours

(Figure 7B). Together, these data demonstrate the existence of a switch point in granuloma

maturity when thrombocytes are either activated or the activation of thrombocytes becomes

pathological.

Co-incident with the appearance of granuloma necrosis at 4 to 5 DPI, we have demonstrated the

appearance of foam cells in zebrafish embryo granulomas at this stage of infection [15]. It has

previously been shown that platelets accelerate the conversion of macrophages to foam cells

in the presence of mycobacteria *in vitro* [9]. Foam cells are permissive for mycobacterial

growth, suggesting a mechanism for infection-induced thrombocyte activation to compromise

innate immunity. We therefore hypothesized that thrombocyte inhibition would reduce the

conversion of macrophages into foam cells. We investigated this by performing Oil-red O

staining to measure lipid accumulation within size-matched granulomas (Figure 7C).

Tirofiban-treated embryos had significantly less Oil-red O accumulation in their granulomas

when compared to DMSO control, even after correction for reduced bacterial burden (Figure

7D). Together, these data demonstrate an *in vivo* effect of thrombocyte activation inhibiting

an effective immune response by converting macrophages into foam cells in the maturing

mycobacterial granuloma.

Given that foam cell formation is closely associated with necrosis in tuberculosis [26] we

hypothesized that tirofiban treatment would reduce cell death within the granuloma. We

therefore used TUNEL staining to detect the fragmented DNA of dying cells in *M. marinum*

infected embryos. At 5 DPI, tirofiban-treated embryos showed significantly less TUNEL

staining, indicating significantly reduced cell death within the granuloma (Figures 7E-F).

Together these results indicate that infection-induced thrombocyte activation aggravates

pathological markers of granuloma maturation and compromise immune control of

mycobacterial infection.

Discussion

Here we have used the zebrafish-*M. marinum* model to identify thrombocyte activation as a

detrimental host response that is co-opted by pathogenic mycobacteria. Our data builds on

previous studies that have shown coagulation, thrombocytosis and thrombocyte activation are

associated with mycobacterial infection, and provides *in vivo* evidence of a direct role for

thrombocyte activation in promoting mycobacterial growth. We have shown that infection-

induced hemostasis is conserved in the zebrafish-*M. marinum* infection model, and that the

platelet inhibiting drugs, aspirin, tirofiban, and eptifibatide, are able to reduce bacterial

burden through host-mediated effects, independently of effects on coagulation.

A number of studies have investigated aspirin as a possible adjunctive treatment for TB in a

range of animal models and human trials [17-21,27,28]. The results of these studies have been far

from conclusive, while most found beneficial effects [17-19,21,27], one human trial observed no

effect [20], and a mouse study identified an antagonistic relationship between aspirin and the

frontline anti-tubercular drug isoniazid [28]. This lack of consensus may be due to the fact that

the NSAID effect of aspirin will affect many cell types and processes important in the

heterogeneous host response to mycobacterial infection. Our study expands this literature by

delineating a role for glycoprotein IIb/IIIa in compromising the host response to infection.

Our study found that coagulation, thrombocytosis, and thrombocyte activation have distinct

roles during the pathogenesis of mycobacterial infection of zebrafish. Inhibiting coagulation

alone did not significantly reduce bacterial burden, and therefore we considered anti-platelet

treatment as a more attractive HDT than anti-coagulant treatment. Although we found lower

total clot formation in tirofiban-treated embryos, this was only proportional to bacterial load,

suggesting infection-induced coagulation could be independent of infection-induced

thrombocyte aggregation. It must be noted that we only measure a simple single end-point in our

zebrafish embryo experiments (bacterial load) at a relatively early time point for a chronic

infection. In more complex animals, where stroke and DVT are important secondary

complications of mycobacterial infection, reducing coagulation may yet prove to be efficacious

as a HDT during TB therapy to reduce morbidity. Conversely, data from the mouse model of TB

suggests tissue factor-induced fibrin is necessary to contain mycobacteria within granulomas [29].

Our study provides evidence that while infection-induced thrombocytosis is a conserved

function of infection, pathological thrombocyte-macrophage interactions are only driven in

pathogenic mycobacteria during granuloma formation. Our experiments with ΔESX1 mutant

M. marinum, which cannot secrete key virulence proteins that drive granuloma formation,

demonstrated ESX1-dependent responsiveness to growth restriction by platelet inhibiting

drugs. These data fit well with our observations that stationary thrombocytes were only

observed around well-developed mycobacterial granulomas, and platelet inhibition was only

effective at reducing bacterial burden after the development of significant granuloma

pathology.

Our imaging and experimental data adds evidence that as for mammalian platelets, activated

thrombocytes can form complexes with leukocytes through fibrinogen binding to glycoprotein

IIb/IIIa, and that this alters immune cell function [25,30-32]. Recent research has highlighted the

important role of platelets as innate immune cells; they are able to release anti-microbial

peptides, pick up and 'bundle' bacteria, and initiate the recruitment of other innate immune

cells to sites of infection [33-35]. However, the low frequency at which direct thrombocyte-

mycobacterial interactions were observed in our study argues against thrombocytes having a

significant role in directly mediating immunity to mycobacterial infection.

Platelets can induce macrophages to produce less pro-inflammatory TNF and IL-1ß, and

more anti-inflammatory IL-10 in response to both BCG and *M. tuberculosis* [7,9,25]. Crucially it

has been shown that platelets are necessary for the formation of foam cells in the context of

mycobacterial infection and atheroma [9]. Our experiments demonstrating anti-platelet drug

mediated control of infection, and reduced lipid accumulation and cell death in the granuloma

suggest that infection-induced thrombocyte activation can be therapeutically modulated in

mycobacterial infection.

Our findings that inhibition of thrombocyte activation reduces foam cell formation and cell

death within the granuloma, leading to reduced bacterial burden provide important *in*

vivo experimental evidence that infection-induced thrombocyte activation is a potential target

for TB host-directed therapy. Specifically, our experimental demonstration of a thrombocyte-

mediated effect of aspirin identifies a novel application for this well-tolerated and low cost

drug in a disease of global importance.

Acknowledgements

We thank Dr Kristina Jahn and Sydney Cytometry for assistance with imaging equipment;

Garvan Biological Testing Facility staff Ms Jennifer Brand, Mr Michael Pickering, Ms Rola

Bazzi, Dr Lucie Nedved and Dr Stephanie Allison at the Garvan Institute of Medical

Research for maintenance of zebrafish breeding stock; and Professors Lalita Ramakrishnan,

Shaun Jackson and Georges Grau, Associate Professor Carl Feng, and Drs Jorn Coers and

Imala Alwis for helpful discussion of results.

This work was supported by the Australian National Health and Medical Research Council

APP1099912 and APP1053407; University of Sydney Fellowship; NSW Ministry of Health

under the NSW Health Early-Mid Career Fellowships Scheme; and the Kenyon Family

Inflammation Award (S.H.O.), Duke Summer Research Opportunities Program (K.J.), NIH

Director's New Innovator Award 1DP2-OD008614 (D.M.T), NIH R01-HL124232 and R01-

HL125774 (J.A.S.).

Author contributions

Declaration of Interests

The authors declare no competing interests.

References

1.   World Health Organization., Global Tuberculosis Programme. Global tuberculosis

control : WHO report. Geneva: Global Tuberculosis Programme:15 volumes.

2.   Oehlers SH, Cronan MR, Scott NR, et al. Interception of host angiogenic signalling

limits mycobacterial growth. *Nature*. 2015;517(7536):612-615.

3.   Oehlers SH, Cronan MR, Beerman RW, et al. Infection-induced vascular permeability

aids mycobacterial growth. *J Infect Dis*. 2017;215(5):813-817.

4.   Robson SC, White NW, Aronson I, Woollgar R, Goodman H, Jacobs P. Acute-phase

response and the hypercoagulable state in pulmonary tuberculosis. *Br J Haematol*.

1996;93(4):943-949.

5.   Kutiyal AS, Gupta N, Garg S, Hira HS. A Study of Haematological and Haemostasis

Parameters and Hypercoagulable State in Tuberculosis Patients in Northern India and the

Outcome with Anti-Tubercular Therapy. *J Clin Diagn Res*. 2017;11(2):OC09-OC13.

6.   Kothari H, Rao LV, Vankayalapati R, Pendurthi UR. Mycobacterium tuberculosis

infection and tissue factor expression in macrophages. *PLoS One*. 2012;7(9):e45700.

7.   Fox KA, Kirwan DE, Whittington AM, et al. Platelets Regulate Pulmonary

Inflammation and Tissue Destruction in Tuberculosis. *Am J Respir Crit Care Med*. 2018.

8.   Dong Z, Shi J, Dorhoi A, et al. Hemostasis and Lipoprotein Indices Signify

Exacerbated Lung Injury in TB With Diabetes Comorbidity. *Chest*. 2018;153(5):1187-1200.

9.   Feng Y, Dorhoi A, Mollenkopf HJ, et al. Platelets direct monocyte differentiation into

epithelioid-like multinucleated giant foam cells with suppressive capacity upon

mycobacterial stimulation. *J Infect Dis*. 2014;210(11):1700-1710.

10.   Vo AH, Swaroop A, Liu Y, Norris ZG, Shavit JA. Loss of fibrinogen in zebrafish

results in symptoms consistent with human hypofibrinogenemia. *PLoS One*.

2013;8(9):e74682.

11.   Lin HF, Traver D, Zhu H, et al. Analysis of thrombocyte development in CD41-GFP

transgenic zebrafish. *Blood*. 2005;106(12):3803-3810.

12.   Walton EM, Cronan MR, Beerman RW, Tobin DM. The Macrophage-Specific

Promoter mfap4 Allows Live, Long-Term Analysis of Macrophage Behavior during

Mycobacterial Infection in Zebrafish. *PLoS One*. 2015;10(10):e0138949.

13.   Matty MA, Oehlers SH, Tobin DM. Live Imaging of Host-Pathogen Interactions in

Zebrafish Larvae. *Methods Mol Biol*. 2016;1451:207-223.

14.   Passeri MJ, Cinaroglu A, Gao C, Sadler KC. Hepatic steatosis in response to acute

alcohol exposure in zebrafish requires sterol regulatory element binding protein activation.

Hepatology. 2009;49(2):443-452.

15.   Johansen MD, Kasparian JA, Hortle E, Britton WJ, Purdie AC, Oehlers SH.

Mycobacterium marinum infection drives foam cell differentiation in zebrafish infection

models. *Dev Comp Immunol*. 2018;88:169-172.

16.   Cronan MR, Beerman RW, Rosenberg AF, et al. Macrophage Epithelial

Reprogramming Underlies Mycobacterial Granuloma Formation and Promotes Infection.

Immunity. 2016;45(4):861-876.

17.   Mai NT, Dobbs N, Phu NH, et al. A randomised double blind placebo controlled

phase 2 trial of adjunctive aspirin for tuberculous meningitis in HIV-uninfected adults. *Elife*.

2018;7.

18.   Kroesen VM, Rodriguez-Martinez P, Garcia E, et al. A Beneficial Effect of Low-

Dose Aspirin in a Murine Model of Active Tuberculosis. *Front Immunol*. 2018;9:798.

19.   Misra UK, Kalita J, Nair PP. Role of aspirin in tuberculous meningitis: a randomized

open label placebo controlled trial. *J Neurol Sci*. 2010;293(1-2):12-17.

20.   Schoeman JF, Janse van Rensburg A, Laubscher JA, Springer P. The role of aspirin in

childhood tuberculous meningitis. *J Child Neurol*. 2011;26(8):956-962.

21.   Tobin DM, Roca FJ, Oh SF, et al. Host genotype-specific therapies can optimize the

inflammatory response to mycobacterial infections. *Cell*. 2012;148(3):434-446.

22.   Peerlinck K, De Lepeleire I, Goldberg M, et al. MK-383 (L-700,462), a selective

nonpeptide platelet glycoprotein IIb/IIIa antagonist, is active in man. *Circulation*. 1993;88(4

Pt 1):1512-1517.

23.   Altieri DC, Mannucci PM, Capitanio AM. Binding of fibrinogen to human

monocytes. *J Clin Invest*. 1986;78(4):968-976.

24.   Armstrong PC, Kirkby NS, Chan MV, et al. Novel whole blood assay for phenotyping

platelet reactivity in mice identifies ICAM-1 as a mediator of platelet-monocyte interaction.

Blood. 2015;126(10):e11-18.

25.   Kullaya V, van der Ven A, Mpagama S, et al. Platelet-monocyte interaction in

Mycobacterium tuberculosis infection. *Tuberculosis (Edinb)*. 2018;111:86-93.

26.   Russell DG, Cardona PJ, Kim MJ, Allain S, Altare F. Foamy macrophages and the

progression of the human tuberculosis granuloma. *Nat Immunol*. 2009;10(9):943-948.

27.   Byrne ST, Denkin SM, Zhang Y. Aspirin and ibuprofen enhance pyrazinamide

treatment of murine tuberculosis. *J Antimicrob Chemother*. 2007;59(2):313-316.

28.   Byrne ST, Denkin SM, Zhang Y. Aspirin antagonism in isoniazid treatment of

tuberculosis in mice. *Antimicrob Agents Chemother*. 2007;51(2):794-795.

29.   Venkatasubramanian S, Tripathi D, Tucker T, et al. Tissue factor expression by

myeloid cells contributes to protective immune response against Mycobacterium tuberculosis

infection. *Eur J Immunol*. 2016;46(2):464-479.

30.   Weber C, Springer TA. Neutrophil accumulation on activated, surface-adherent

platelets in flow is mediated by interaction of Mac-1 with fibrinogen bound to alphaIIbbeta3

and stimulated by platelet-activating factor. *J Clin Invest*. 1997;100(8):2085-2093.

31.   Sanderson HM, Fox SC, Robbins RA, Losche W, Spangenberg P, Heptinstall S. Role

of GPIIb-IIIa in platelet-monocyte and platelet-neutrophil conjugate formation in whole

blood. *Platelets*. 1998;9(3-4):245-250.

32.   Kral JB, Schrottmaier WC, Salzmann M, Assinger A. Platelet Interaction with Innate

Immune Cells. *Transfusion Medicine and Hemotherapy*. 2016;43(2):78-88.

33.   Gaertner F, Ahmad Z, Rosenberger G, et al. Migrating Platelets Are Mechano-

scavengers that Collect and Bundle Bacteria. *Cell*. 2017;171(6):1368-1382 e1323.

34.   Li JL, Zarbock A, Hidalgo A. Platelets as autonomous drones for hemostatic and

immune surveillance. *J Exp Med*. 2017.

35.   Morrell CN, Aggrey AA, Chapman LM, Modjeski KL. Emerging roles for platelets as

immune and inflammatory cells. *Blood*. 2014;123(18):2759-2767.

Figure legends

Figure 1: *Mycobacterium marinum* induces coagulation and thrombosis around sites of

infection in zebrafish

A) Representative images of 5 DPI *Tg(cd41:GFP)* embryos infected with *M. marinum*-

tdTomato, showing thrombocyte accumulation (green) at sites of bacterial infection (red). B)

Quantification of total thrombocyte *Tg*(*cd41:GFP*) area within the tail of uninfected or WT

M. marinum-tdTomato infected embryos. Graph shows mean ± SEM. Statistical analysis

performed by T tests. C) Correlation between *M. marinum* bacterial burden and total

thrombocyte *Tg(cd41:GFP)* area within the tail of infected embryos. P and R^2 calculated

using linear regression. D) Representative 2 DPI, 3 DPI and 4 DPI images of *Tg(cd41:GFP)*

fish infected with *M. marinum*-tdTomato at 3 DPF. White arrowheads show thrombocyte

(green) association with areas of bacterial growth (red). Thrombocytes not indicated with an

arrowhead were circulating and not considered to be associated with bacteria.

Figure 2: Anti-platelet drugs reduce bacterial burden in *M. marinum* infection

A) Quantification of bacterial burden in embryos treated with aspirin normalized to DMSO

control. Data are combined results of two independent experiments. B) Quantification of

bacterial burden in embryos treated with tirofiban normalized to DMSO control. Data are

combined results of two independent experiments. C) Quantification of bacterial burden in

embryos treated with eptifibatide or DMSO control. Data are combined results of two

independent experiments. D) Quantification of bacterial growth by relative fluorescence in

7H9 broth culture supplemented with drugs as indicated.

All graphs show Mean ± SEM. Statistical analyses performed by T tests.

Figure 3: The anti-bacterial effect of anti-platelet drugs is thrombocyte dependent

A) Total fluorescence area of *M. marinum* bacteria in larvae injected with either control or

cmpl morpholino (MO) to deplete thrombocytes, and then treated with aspirin. Values are

normalized to DMSO-treated control MO larvae. Graphs show the combined results of 2

independent experiments. B) Total fluorescence area of *M. marinum* bacteria in larvae

injected with either control or cmpl morpholino (MO) to deplete thrombocytes, and then

treated with tirofiban (Tiro). Values are normalized to DMSO-treated control MO larvae.

Graphs show the combined results of 2 independent experiments. C) Quantification of

bacterial burden in in *itga2b* mutant embryos normalized to WT control. Data are combined

results of 3 independent experiments. D) Quantification of bacterial burden in *fga*$^{-/-}$ embryos

treated with tirofiban, normalized to *fga*$^{+/+}$ and *fga*$^{+/-}$controls. Data are combined results of 2

independent experiments.

All graphs show Mean ± SEM. Statistical analyses performed by ANOVA.

Figure 4: *Mycobacterium marinum* induces coagulation in zebrafish but inhibition of

coagulation does not affect the outcome of infection.

A) Representative images of a *Tg(fabp10a:fgb-EGFP)* embryo infected with *M. marinum*-

tdTomato by caudal vein injection, showing clot formation (green) at sites of infection (red)

at 3, 4, and 5 DPI. B) Quantification of clot formation in burden-matched ΔESX1 mutant-

infected *Tg(fabp10a:fgb-EGFP)* embryos normalized to WT *M. marinum* control. Data show

combined results from two independent experiments. C) Quantification of clotting in

warfarin-treated *Tg(fabp10a:fgb-EGFP)* embryos. D) Quantification of bacterial burden in

warfarin treated embryos, normalized to DMSO control. Data show combined results of two

independent experiments. All graphs show mean ± SEM, statistical analyses by T tests.

Figure 5: Thrombocytes increase mycobacterial burden independently of coagulation.

A) Representative images of 5 DPI *Tg(fabp10a:fgb-EGFP)* embryos infected with

M.marinum-tdTomato and treated with either DMSO or tirofiban. B) Quantification of

clotting relative to bacterial burden in embryos treated with tirofiban normalized to DMSO

control. Data are combined results of two independent experiments. C) Representative

images of *Tg(fabp10a:fgb-EGFP)* embryos, where clot formation can be visualized by GFP

fluorescence, infected with *M. marinum*-tdTomato (red) and treated with either DMSO or

warfarin. D) Quantification of bacterial burden in embryos treated with tirofiban, warfarin, or

tirofiban and warfarin, normalized to DMSO control. Data are combined results of 2

independent experiments. All graphs show mean ± SEM, statistical analysis by T test or

ANOVA where appropriate.

Figure 6: Thrombocyte activation compromises immunity through granuloma-associated

macrophages.

A) Quantification of total thrombocyte *Tg*(*cd41:GFP)* area within the tail WT *M. marinum*-

tdTomato infected embryos treated with either DMSO or tirofiban. B) Quantification of total

thrombocyte *Tg*(*cd41:GFP)* area within the tail of uninfected or ΔESX-1 *M. marinum*-

tdTomato infected embryos. C) Quantification of bacterial burden in embryos infected with

ΔESX1 *M. marinum*-tdTomato and treated with tirofiban. Combined data of 3 independent

experiments. D) Quantification of length of thrombocyte association with an infected

macrophage 5 DPI. Eptifibatide treatment was started at 4 DPI and was continued during

timelapse. Data points represent individual thrombocytes from four (two control and two

drug treated) infected fish. E) Quantification of distance thrombocyte traveled while

associated with a site of macrophage and bacterial accumulation. Data points represent

individual thrombocytes from four (two control and two drug treated) infected fish. F)

Representative tracks of thrombocytes (I,II,IV,V) within control and eptifibatide treated

embryos. Representative tracks of individual thrombocyte (blue), macrophage (red), and

internalized *M. marinum* (green) (III, VI). G) Representative images of caudal hematopoietic

tissue in 6 DPF *Tg(mfap4:tdTomato)* embryos, where macrophages are fluorescently labeled,

injected with clodronate or PBS liposomes at 4 DPF. H) Quantification of bacterial burden in

embryos infected with *M. marinum*-tdTomato, injected with clodronate liposomes and treated

with tirofiban from at 3 DPI. All graphs show mean ± SEM, statistical tests by T-tests.

Figure 7: Granuloma maturation and pathological thrombocyte activation have a

bidirectional relationship.

A) Quantification of bacterial burden at 3 DPI after continuous tirofiban treatment from 0

DPI. B) Quantification of bacterial burden at 5 DPI after overnight drug treatment initiated at

4 DPI. C) Representative images of bacterial granulomas chosen for analysis (bacteria are

white in greyscale images), and corresponding Oil Red O (ORO) staining (red-brown in color

images). D) Quantification of ORO pixel density relative to granuloma bacterial area, in

embryos treated with tirofiban, normalized to DMSO control. E) Representative images of

bacterial granulomas showing bacteria in red and TUNEL staining in green. F) Quantification

of TUNEL positive area within the largest granuloma of individual embryos. All graphs show

Mean ± SEM, statistical tests by T-tests. Data are combined results of two independent

experiments, except F) which represents a single experiment.

Supplementary Figure Legends

Figure S1

A) Representative images of thrombocyte accumulation at tail wound in 4 DPF

Tg(cd41:GFP) embryos , 4 hours post injury (hpi). B) Quantification of thrombocytes within

100 μm of the cut site 4 hpi in aspirin-treated embryos. C) Quantification of thrombocytes

within 100 μm of the cut site 4 hpi in tirofiban-treated embryos. D) Quantification of

thrombocytes within 100 µm of the cut site 4 hours after injury in *itga2b* WT, heterozygous

and knock-out embryos. Graphs show Mean ± SEM, statistical testing by T-test.

Supplementary Video 1

Video of 5 DPI *Tg(cd41:GFP)* embryo infected with *M. marinum*-tdtomato. Video captured

with a long pass GFP filter to capture thrombocytes and bacteria simultaneously. Arrow

indicates an example of a 5 minute thrombocyte-bacteria interaction.

Supplementary Video 2

Representative *Tg(fabp10a:fgb-EGFP)* embryo expressing EGFP-tagged fibrinogen beta

(FGB) infected with *M. marinum*-tdtomato. Imaged every 15 minutes from 3 days post

infection (DPI), until 6 DPI.

Supplementary Video 3

Representative timelapse of control *Tg(cd41:GFP; mfap4:tdtomato)* embryo infected with

M. marinum-cerulean, showing the interaction of thrombocytes (green), macrophages (red), and

bacteria (blue). Images were captured every 5 minutes for 4 hrs. Arrow shows successive short

'binding' events between thrombocytes and an infected macrophage.

Supplementary Video 4

Representative timelapse of eptifibatide treated *Tg(cd41:GFP; mfap4:tdtomato)* embryo

infected with *M. marinum*-cerulean, showing the interaction of thrombocytes (green),

macrophages (red), and bacteria (blue). Images were captured every 5 minutes for 5 hrs. Arrows

show simultaneous long 'binding' events between thrombocytes and infected macrophages.

Supplementary Video 5

Representative tracks of thrombocytes associated with infected macrophages, from timelapse

of control *Tg(cd41:GFP; mfap4:tdtomato)* embryo infected with *M. marinum*-cerulean. Images

were captured every 5 minutes for 4 hrs, and tracks were generated using the Manual Tracking

function in ImageJ.

Supplementary Video 6

Representative tracks of thrombocytes (blue and teal), showing the tracks of the macrophage

(green and purple) and internalized M. marinum (red and yellow) they are associated with.

From timelapse of control *Tg(cd41:GFP; mfap4:tdtomato)* embryo infected with *M. marinum*-

cerulean. Images were captured every 5 minutes for 4 hrs, and tracks were generated using the

Manual Tracking function in ImageJ.

Supplementary Video 7

Representative tracks of thrombocytes associated with infected macrophages, from timelapse

of eptifibatide treated *Tg(cd41:GFP; mfap4:tdtomato)* embryo infected with *M. marinum*-

cerulean. Images were captured every 5 minutes for 5 hrs, and tracks were generated using the

Manual Tracking function in ImageJ.

Supplementary Video 8

Representative tracks of thrombocytes (blue and teal), showing the tracks of the macrophage

(green and purple) and internalized M. marinum (red and yellow) they are associated with.

From timelapse of eptifibatide treated *Tg(cd41:GFP; mfap4:tdtomato)* embryo infected with

M. marinum-cerulean. Images were captured every 5 minutes for 4 hrs, and tracks were

generated using the Manual Tracking function in ImageJ.

A

3DPI   4DPI   5DPI

M. marinum FGB

B

Normalised FGB-GFP Area

0.0200

M.mar WT   *M. mar* ΔESX1

C

FGB-GFP Total Area

0.0073

DMSO   Warfarin

D

Normalised Bacteria Area

0.3129

DMSO   Warfarin