The effects of habits on motor skill learning

Nicola J. Popp[1], Atsushi Yokoi[2,3], Paul L. Gribble[1,4,5,6] & Jörn Diedrichsen[1,7,8*]

[1]The Brain and Mind Institute, University of Western Ontario, Canada,

[2]Graduate School of Frontier Biosciences, Osaka University, Japan,

[3]Center for Information and Neural Networks (CiNet), NICT, Japan,

[4]Department of Psychology, University of Western Ontario, Canada,

[5]Department of Physiology & Pharmacology, Schulich School of Medicine & Dentistry, University of Western Ontario, Canada,

[6]Haskins Laboratories, USA,

[7]Department of Statistical and Actuarial Sciences, University of Western Ontario, Canada,

[8]Department of Computer Science, University of Western Ontario, Canada

Acknowledgments

This work is supported by a James S. McDonnell Foundation Scholar award, a Natural Sciences and Engineering Council of Canada (NSERC) Discovery Grant (RGPIN-2016-04890) and the Canada First Research Excellence Fund (BrainsCAN) to J.D., a NSERC Discovery Grant (RGPIN 238338) and a Canadian Institutes of Health Research Grant (PJT-153447) to P.L.G. We thank Aaron L. Wong for helpful comments on earlier versions of this manuscript.

Correspondence address

Jörn Diedrichsen, The Brain and Mind Institute, University of Western Ontario, London, Canada. jdiedric@uwo.ca

Author contributions

N.J.P. and J.D. designed the experiment. N.J.P, A.Y. and J.D. programmed the experiment. N.J.P collected the data and analyzed the data together with J.D., N.J.P., J.D., and P.L.G. wrote the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Abstract

Skill learning involves the formation of stable motor patterns. In musical and athletic

training, however, these stable patterns can also impede the attainment of higher levels

of performance, and hence constitute a motor habit. We developed an experimental

paradigm to induce a specific motor pattern in a sequence production task and

investigated how it affected subsequent optimization over a 3-week training period.

Participants initially practiced small segments of 2 to 3 finger movements, which were

then combined to form longer sequences. This initial training induced a persistent

chunking behavior, with shorter inter-press-intervals within a chunk and longer ones at

chunk boundaries. We were able to induce chunking that was either beneficial or

detrimental to performance, and could show that the degree to which these detrimental

chunk structures were maintained, predicted lower levels of final performance. We also

identified two optimization processes by which participants overcame the detrimental

motor habits.

Introduction

Humans are capable of astonishing feats of motor skill in athletics, musical performance

and dance. But what does it take to become an expert? The first obvious factor is

practice: it is estimated that 10,000 hours of training are necessary to develop high-level

motor skills (Ericsson et al., 1993; Hayes, 2013). Perhaps the motor system simply

needs to gather a large amount of experience to optimize the motor commands that are

necessary to achieve skilled movement. Thus, given enough practice, motor expertise

may emerge automatically.

However, simply practicing for many hours will not automatically lead to expert

performance. There are numerous examples in which motor skill acquisition is slow or

fails (Haith & Krakauer, 2018). This is sometimes attributed to the formation of habits:

automatic (Hélie, Waldschmidt, & Ashby, 2010; Moors & De Houwer, 2006) and highly

entrenched behavioral patterns that resist change through retraining (Ashby, Ell, &

Waldron, 2003; Graybiel & Grafton, 2015; Hardwick, Forrence, Krakauer, & Haith, 2017;

Jager, 2003; Seger & Spiering, 2011).

Animal models have been integral to the study of habit formation and its neural

underpinnings (Jog, Kubota, Connolly, Hillegaart, & Graybiel, 1999; Robbins & Costa,

2017; Smith & Graybiel, 2014, 2016; Wickens, Horvitz, Costa, & Killcross, 2007).

However, the majority of animal experiments investigating this concept have focused on

habits in the context of action selection – i.e. choosing *what* action to perform. In

contrast, our paper addresses the question of habits in motor performance – i.e. habits

in *how* to perform a chosen action. For example, a tennis player could have a habit in

action selection, whereby she always chooses a forehand over a backhand to return a

serve. Independently, she could have a motor habit, whereby she executes the

forehand without rotating her hips.

Critical to the definition of a habit is that the behavior is maintained, even though

it is no longer adaptive (Adams, 1982; Dezfouli & Balleine, 2012; Dickinson, 1985). Most

experiments, therefore, demonstrate the existence of a habit by teaching subjects a

behavior under one reward contingency and show that it persists even when the reward

contingency switches (Ashby et al., 2003; Smith & Graybiel, 2013b). In a similar way,

we define motor habits here as a stable way of performing an action that is maintained,

even if it prevents optimal performance. This does not imply that motor habits always

have to be dysfunctional. Their automatic nature can be beneficial by increasing

processing speed (Hardwick et al., 2017) or by reducing cognitive load (Haith &

Krakauer, 2018; Hélie & Cousineau, 2011). Thus, habits can be either functional or

dysfunctional, but their defining criterion is that they are resistant to change even under

circumstance where a change would be beneficial.

To investigate the influence of habit formation on motor skill learning, we

introduce a novel experimental paradigm that enables us to induce beneficial and

detrimental motor habits and to test whether participants can overcome these habits

with practice. As an experimental model of skill acquisition, we used the discrete

sequence production task (DSP), in which participants perform an explicitly known

series of finger presses as fast as possible (Abrahamse, Ruitenberg, de Kleine, &

Verwey, 2013; Verwey, 2001). Learning in this task depends on both cognitive and

motor processes (Diedrichsen & Kornysheva, 2015; Wong, Lindquist, Haith, &

Krakauer, 2015). Initial performance relies strongly on forming a declarative memory of

the sequence and can, therefore, be sculpted through explicit instructions (de Kleine &

Verwey, 2009; Verwey, Abrahamse, & de Kleine, 2010; Verwey, Abrahamse, &

Jiménez, 2009). An important determinant of the structure of declarative memory is

"chunking" – the process by which participants separate a long sequence into smaller

subsets (Verwey, 1996; Verwey & Dronkert, 1996) to aid memorization and improve

performance by reducing memory capacity demands (Halford, Wilson, & Phillips, 1998;

Miller, 1956; Wymbs, Bassett, Mucha, Porter, & Grafton, 2012).

The structure of the declarative memory representation of a sequence may then

constrain subsequent motor optimization processes (Bo & Seidler, 2009; Seidler, Bo, &

Anguera, 2012). For example, it has been suggested that sequential movements may

be optimized within a chunk, but not across chunk boundaries (Ramkumar et al., 2016).

We hypothesized therefore that the initial cognitive chunking of the sequence influences

the learning of execution-related skills in subsequent motor training. That is, we tested

the hypothesis that cognitive chunking can evolve into a motor habit.

We trained participants to perform the same set of 7 sequences, each consisting

of 11 isometric keypresses. Training occurred on 14 separate days, spread over 3

weeks. In the induction phase (Fig. 1a) we imposed a specific chunk structure by

instructing participants to practice a set of 2-3 digit chunks. They then learned the 11-

digit sequences as being composed of four of the pre-trained 2-3 digit chunks. We

induced chunk structures that were designed to be either aligned or misaligned with

biomechanically easy or difficult finger transitions within the sequence and therefore

were predicted to be beneficial or detrimental to performance. Each participant learned

3 sequences using the misaligned chunk structure and 3 distinct sequences with the

aligned chunk structure (Fig. 1d). To test for patterns of spontaneous chunking, a

separate group of participants (control) was trained on the same sequences but did not

receive chunk training and instead practiced the complete sequences during the initial

training.

In the subsequent optimization phase (Fig. 1a), participants were instructed to

improve their performance through practice. During this phase, we did not make any

further mention of chunks (see Methods). Because participants performed the same

sequences (but differed in the chunk structures with which they had learned the

sequences), we could draw strong inferences about whether their performance was

dictated by biomechanical demands (which were identical across participants) or

whether it was affected by the chunk structure that was imposed during the induction

phase. Using a Bayesian model, we estimated changes in chunk structure. This allowed

us to investigate three questions: First, can explicit instructions at the beginning of

training lead to stable motor performance patterns? Second, to what degree are these

patterns maintained if they impede the participants' ability to reach skilled performance

– i.e. can these patterns be considered a motor habit? Finally, what are the optimization

processes that allow participants to overcome bad habits (misaligned chunk structures)

through practice?

102　　　　　　　　　　　　　　　　**Results**

Over 15 days we trained 32 participants to produce sequences of 11 isometric

keypresses from memory on a keyboard-like device. Participants were rewarded with

points for executing sequences as fast as possible while keeping the proportion of

incorrect keypresses in each block of trials below 15%. We maintained the participants'

motivation by gradually decreasing the movement time (MT) threshold at which they

received points.

We manipulated how participants memorized the sequences by splitting the

sequences into several chunks, each composed of 2-3 keypresses. We wanted to test

whether the different ways of chunking (hereafter "chunk structures") imposed in the

induction phase (Fig. 1a) would affect performance optimization in the subsequent two

weeks of training. On the first day, we trained participants to produce eleven 2-3 press

chunks in response to a visually presented letter. For example, "A" corresponded to the

chunk "3,2,1" (middle finger, index finger, thumb). At the end of the first day, participants

could reliably produce the chunks from memory with an average accuracy of 92.7%. On

days 2-4, we combined these chunks to form 7 different 11-press sequences. Each

sequence was associated with a specific character symbol (e.g. $ see Supp. Table 1).

At the end of day 4, participants were able to recall all sequences from memory using
the sequence cue with an accuracy of 92.5%.

*Figure 1. Experimental procedure. (**a**) Experimental timeline depicting the training at each stage. In the induction phase participants memorized chunks and sequences. In the optimization phase participants trained to perform these sequences as fast as possible from memory. In the last week of training, half of the participants were directly cued with the sequence, while the other half performed the sequences from memory. (**b**) Data from an independent dataset, in which participants performed all possible combinations of 2 and 3-digit transitions. Matrix indicates the median inter-press interval (IPI) to produce the transition between pairs of keypresses. (**c**) Top: Example sequence containing a 3-digit run and two digit repetitions. Bottom: The sequence was instructed using two possible chunk structures. In the aligned structure, the chunk boundaries fell between repetitions, in the misaligned structure the chunk boundary broke up the run. (**d**) The assignment of chunk structures to sequences was counterbalanced between participants.*

Chunk induction induces a stable motor pattern

To assess whether the imposed chunk structure influenced participants' motor behavior, we examined inter-press time intervals (IPIs). An increased IPI is commonly taken as a sign of a chunk boundary, as the cognitive processes (memory recall, action selection) involved in switching from one chunk to another require additional time (Verwey, 1999; Verwey et al., 2010). Hence, we would expect our participants to exhibit shorter IPIs between keypresses that belonged to a chunk imposed during day 1 (within-chunk IPIs) and larger IPIs for the boundaries between chunks (between-chunk IPIs). We indeed found significantly longer between-chunk IPIs compared to within-chunk IPIs in the first few days of training (Fig. 2a: days 2-4: $t_{(31)} = 7.728$, $p = 5.098\text{e-}09$).

In the optimization phase (day 5-14), we ceased to cue sequences using the alphabetic letters associated with the chunks. Instead, participants were asked to recall the sequences from memory in response to the symbolic sequence cues (e.g. "$"). From this point forward, no further reference to the imposed chunk structure was made. Across days 5-10, the difference between the within- and between-chunk IPIs remained stable; $t_{(31)} = 7.165$, $p = 2.351\text{e-}08$ (Fig. 2a). Importantly, this difference cannot be attributed to biomechanical difficulty of the finger transitions. The within-chunk IPIs for one group were the between-chunk IPIs for the other group and vice versa; IPIs that were within-chunk for all participants (e.g. the first and last IPI of a sequence) were excluded from this analysis. In summary, even though after day 4 we cued the sequences only with a single symbol, participants persisted in performing the sequences consistent with the chunk structures imposed early in training.

***Figure 2. Within- vs. between-chunk inter-press intervals (IPIs). (a)** Time course of IPIs that were within an instructed chunk (dashed line), or on the boundary between chunks (solid line). Asterisks indicate significant differences between average within- and between-chunk IPIs in the corresponding week (separated by dashed lines). Shaded area denotes between-subject standard error. **(b)** Difference of between- and within-chunk IPIs in the last week of training, split by whether participants had to recall the sequences from memory or were cued with the sequence numbers. Violin plots indicate distribution of individual participants, white circles indicate means.*

In the last four days of training (days 11-14) we tested whether the persistence of
the imposed chunk structure reflected a motor habit or whether it reflected memory
recall. We split each experimental group into two subgroups: half of the participants
continued to perform the sequences from memory, while the other half were cued using
the numbers (Fig. 1a) that indicated the necessary keypresses, therefore removing any
memory recall demands. Both the memory ($t_{(15)}$ = 4.865, *p* = 2.059e-04, Fig. 2b) and the
cued subgroup ($t_{(15)}$ = 3.403, *p* = 0.004) showed a significant difference between the
within- and between-chunk IPIs and there was no reliable difference between the two
subgroups in this effect ($t_{(30)}$ = -0.749, *p* = 0.460). Thus, removing the requirement for
memory recall did not abolish chunking. Because none of the subsequent analyses
showed any significant difference between the two subgroups, we will report their
combined results for the remainder of the article. Overall, these results suggest the

explicit chunk training early in learning established a stable performance pattern that

outlasted 10 days of subsequent practice.

Misaligned chunk structure impairs performance

To show that this initial instruction led to the emergence of a motor habit (as

defined above), we needed to not only show that this initial instruction induced a stable

temporal pattern of IPIs, but also that this pattern was maintained when it leads to

slower execution speeds than other patterns. We therefore designed chunk structures

that were predicted to be either beneficial or detrimental to performance. These

predictions were based on a separate experiment (see Methods), in which we trained 7

participants on all possible 2 and 3 keypress combinations over the course of 3 days

and measured their execution speed. Transitions between two adjacent fingers could be

performed faster than two repeated presses of the same finger ($t_{(6)}$ = 13.965, p =

8.404e-06; see Fig. 1b). Given that the 2-3 press sequences hardly taxed the cognitive

system, these results can be taken as a characterization of the biomechanical

constraints of our specific task (see Methods).

We used these results to design two different ways of separating the sequences into

chunks. In one case, we placed chunk boundaries so that they were **aligned** with digit

transitions that were performed more slowly (as measured in the independent dataset) –

i.e. they were preferentially placed between digit repetitions (Fig. 1c). The time required

to perform these difficult finger transition can therefore simultaneously be used to recall

a next chunk, which should benefit overall performance. In the **misaligned** chunk

structure, we placed chunk boundaries at digit transitions that can be performed quickly,

thereby breaking up transitions between adjacent fingers or runs (Fig. 1c). Participants

would, therefore, have to slow down their performance at these fast transitions to recall

the next chunk, which should slow overall performance. Each participant learned 3 of

the 7 sequences with a misaligned chunk structure and 3 sequences with an aligned

chunk structure, with the assignment counterbalanced across groups. For the last

remaining sequence, both ways of chunking were predicted to be equally fast, as both

possible chunk structures were aligned with the biomechanical requirements (neutral

chunk structure, Fig. 1d & Supp. Table 1).

We predicted that training on the misaligned chunk structure would lead to poorer

performance. To quantify performance, we used movement time (MT), the time between

the first finger press and the last finger release. Indeed, in the induction phase (days 2-

4), the sequences instructed with the misaligned chunk structure were performed slower

than the sequences instructed with the aligned chunk structure (one-sample t-test: $t_{(31)}$ =

2.693, p = 0.006; Fig. 3a). Hence, we were not only able to manipulate *how* participants

performed a sequence, but also *how well* they could perform it. This difference was

maintained in the second week of training (days 5-10: $t_{(31)}$ = 2.313, p = 0.014).

Importantly, this shows that the stable pattern of IPIs indeed constitutes a motor habit.

Figure 3. Change in chunk structure and performance for aligned and misaligned instructed sequences. (a) Differences in movement time (MT) between sequences instructed with an aligned or misaligned chunk structure. Asterisk indicates a significant difference from 0 (no difference). (b) Within- or between-chunk IPIs across training days, separated by whether they were in the aligned or misaligned instructed sequences. Error bars denote between-subject standard error.

Misaligned chunk structure is changed more rapidly

Interestingly, the difference in performance between the sequences instructed with the aligned compared to the misaligned chunk structure was no longer significant in the last four days of training (days 11-14: $t_{(31)}$ = 0.764, p = 0.225; Fig. 3a). This suggests that participants were able to overcome the "bad" habit of a misaligned chunk structure to some degree. To investigate this, we separated the IPI analysis (Fig. 2a) by whether the intervals came from sequences that were instructed using an aligned or misaligned structure. While the difference between within- and between-chunk IPIs for "aligned sequences" was stable over the entire training period, the difference for the "misaligned sequences" disappeared in the last four days of training (Fig. 3b). The three-way interaction between day x within/between x instruction (aligned or misaligned) was significant ($F_{(12,372)}$ = 19.790, p < 1e-16). Thus, in the last four days of training participants seemed to diverge from the misaligned chunk structure while maintaining the aligned chunk structure.

Tracking changes in chunking

A disadvantage of the above analysis, however, is that we cannot discern how participants restructured their chunking and whether they completely abandoned the misaligned chunk structure. For a clearer understanding of how participants changed their chunk structure, we used a Bayesian model to estimate the probability of each possible chunk structure, given the observed series of IPIs, on a trial-by-trial basis (Acuna et al., 2014). The state variable in this Hidden Markov Model indicates which of the 1023 possible chunk structures is present on each trial. Using the expectation-maximization (EM) algorithm (Dempster, Laird, & Rubin, 1977; Welch, 2003), we simultaneously estimated the 9 free parameters of the model (for details see Methods), and the posterior probability for each possible chunk structure on each trial. We accounted for the effects of biomechanical difficulty by regressing out the patterns of IPIs across finger transitions (Fig. 1b) from each block before modeling. Importantly, our version of the model could capture separate learning-related changes to the within- and between-chunk intervals (Fig. 4a). Our method, therefore, allowed us to estimate participants' chunk structure independently of the overall speed of performance. We confirmed this independence using simulated data (see Methods).

225   Figure 4b shows two examples of individual participants and sequences. In the

226   first panel, the participant chunked the sequence according to the initial instructions at

227   first, then inserted 1 or 2 additional chunk boundaries, and at the end of training

228   performed the sequence as a single chunk. In comparison, the other participant

229   maintained the instructed chunk structure for most of the training period.

Figure 4. Probabilistic chunking model fitted to example participant data. (a)
The change of within- and between-chunk IPIs were modeled using two separate
exponential functions across training. The density plot shows individual IPIs, with the
color indicating the probability of a between- (pink) or within-chunk interval (blue). ***(b)***
Posterior probability for two example participants (for one sequence per participant)
over the course of the experiment. Only the 4 most likely chunk structures out of the
1023 possible structures are shown. The color scale indicates the posterior probability
of a given chunk structure for each trial - with yellow indicating higher probabilities.
The dashed vertical lines indicate the boundaries between training phases (Days 2-4;
5-10 & 11-14). The black box (left) indicates the chunk boundaries as white lines
within the 11-press sequence (max. 10 boundaries) for the chosen chunk structures.
The first row indicates the instructed chunk structure (arrow). The other three rows
illustrate other chunk structures that were highly probable at some point during the
experiment. The distance measure expresses how many chunks need to be added or
removed to transform one structure (in this case the instructed chunk structure) into
the other.

Movement towards a single chunk structure

To characterize changes in chunk structure, we first defined a metric that quantified the difference between two chunking structures: we counted the number of chunk boundaries that differ – i.e. the number of chunks that needed to be split or merged to transform one chunk structure into the other (Fig. 4b - distance). We then used this measure to calculate the distance between the chunk structure used by the participant and three reference structures of interest: (1) the aligned-, (2) misaligned, and (3) a structure that consisted of a single chunk. These distances defined a coordinate system that enabled us to visualize changes in chunk structure. We then projected participants' estimated chunk structure into this space (Fig. 5a). On the horizontal axis, we plotted the expected distance of participants' chunk structure to the single-chunk structure. Given the definition of our distance, this measure simply counts the number of chunk boundaries. On the vertical axis, we plotted how close the estimated chunk structure was to the aligned and misaligned chunk structure.

Previous literature has suggested that participants group smaller chunks together with training (Kuriyama, Stickgold, & Walker, 2004; Ramkumar et al., 2016; Sakai, Kitaguchi, & Hikosaka, 2003; Song & Cohen, 2014; Verstynen et al., 2012; Verwey, 1996; Wymbs et al., 2012), a process that may help to improve performance (Abrahamse et al., 2013; Ramkumar et al., 2016; Verwey, 1999, 2001; Verwey et al., 2010; Verwey & Wright, 2014). In nearly all previous studies, however, the estimated number of chunks is biased by the overall movement speed. Using a modified probabilistic model (see Methods), we were able to disambiguate the two and critically test this assumption. We estimated the number of chunk boundaries for each participant averaged across sequences (the neutral sequence was excluded). Interestingly, on the 2nd day, participants separated sequences into more chunks than the 4 chunks we instructed (Fig. 5a, $t_{(31)}$ = 4.224, p = 0.0002). This tendency continued on day 3, where participants tended to subdivide the sequences into even smaller chunks (Fig. 5b; day 2 vs. 3: $t_{(31)}$ = 2.023, p = 0.052). After day three the number of chunk boundaries decreased as shown by a significant effect of day in a repeated measures ANOVA ($F_{(11,341)}$ = 11.710, p < 1e-16). However, even in the last phase of training, participants performed the sequences with an average of 2.9 chunk boundaries (we instructed 3

chunk boundaries). Thus, while there was a clear tendency towards merging chunks
after an initial increase, participants did not perform the sequence as a single chunk,
even after 3 weeks of practice.

Participants abandoned the misaligned chunk structure to a greater degree
Next, we probed how much participants diverged from the initial instructions.
Participants slowly changed their chunk structure for both aligned and misaligned
instructed sequences with training. The average distance to the instructed chunk
structure increased systematically over time (repeated measures ANOVA, effect of day,
$F_{(12,372)} = 7.055$, $p < 1e-16$, Fig. 5c).

Consistent with our IPI analysis (Fig. 3b), we observed that participants
abandoned the misaligned chunk structure to a greater degree than the aligned chunk
structure (Day x Instruction interaction: $F_{(12,372)} = 5.610$, $p < 1e-16$). In the last four days
of training, the sequences that were instructed with the misaligned chunk structure were
more dissimilar to the misaligned chunk structure than the sequences that were
instructed with the aligned chunk structure were to the aligned chunk structure: $t_{(31)} =$
2.294, $p = 0.029$ (Fig. 5c). Additionally, we found a significant Day x Instruction
interaction ($F_{(12,372)} = 2.215$, $p = 0.011$) for the distance to a single chunk (Fig. 5b),
suggesting a stronger tendency towards performing a sequence as a single chunk when
trained on the misaligned chunk structures. Together these results indicate that
participants changed their chunking behavior more readily for sequences that were
trained using the misaligned chunk structure than when trained using the aligned chunk
structure.

Despite the divergence from the misaligned chunk structure with training, our
analysis also revealed that participants did not overcome the influence of the instruction
completely. In the third week, sequences produced after training with a misaligned
chunk structure were still performed closer to the misaligned structure than to the
aligned structure ($t_{(31)} = 6.962$, $p < 1e-16$). This shows that even training on misaligned
chunk structures had a lasting influence on participants' motor behavior.

Figure 5. Changes in chunk structure with learning. ***(a)*** *The average chunk structure over 13 days of practice for aligned (red) and misaligned (blue) instructed sequences for the experimental participants. The results of the control group are shown in green. The horizontal axis represents the distance to the single-chunk structure, i.e. the number of chunk boundaries. The vertical axis shows the distance to the aligned or misaligned chunk structure. The crosses indicate the positions of the three reference structures (aligned, misaligned and single). Ellipses denote the between-subject standard error.* ***(b)*** *Average distance of participants' chunk structure to the single chunk structure across days.* ***(c)*** *Distance to the instructed chunk structure.* ***(d)*** *Day-by-day changes in chunk structure.* ***(e)*** *Trial-by-trial changes in chunk structures within each day. Error bars indicate between-subject standard error.*

Chunk structure "crystallizes" with training

Would longer training allow participants to completely overcome the influence of the instruction and to perform all sequences as a single chunk? Although experiments with longer training are necessary to provide a definitive answer, our data indicate that this process, if occurring, may take a very long time. The amount of change in the chunk structure for each sequence reduced dramatically in the last week of training, suggesting that a stable motor habit formed. This phenomenon is akin to the development of an invariant temporal and spectral structure in bird-song, a process that has been termed "crystallization" (Brainard & Doupe, 2002). To estimate crystallization, we calculated the distance between the chunk structures from one day to the next (Fig. 5d) and within each day from one trial to the next (Fig. 5e). The analysis was performed separately for each sequence and participant. Overall, both the day-to-day distance ($F_{(11,330)} = 18.794$, $p < 1e-16$) and the trial-by-trial distance decreased significantly across training days ($F_{(12,456)} = 13.245$, $p < 1e-16$). Therefore, participants appeared to settle onto a stable pattern in the last week. Consequently, additional training would likely only lead to slow changes in their chunk structure.

In summary, our analyses provide a clearer picture of how chunking changes with learning. Firstly, in line with previous research (Kuriyama et al., 2004; Ramkumar et al., 2016; Sakai et al., 2003; Song & Cohen, 2014; Verstynen et al., 2012; Verwey, 1996; Wymbs et al., 2012) participants gradually moved towards performing the sequence as a single chunk by dividing the sequence into fewer chunks. Secondly, participants diverged from the instructions over time with a quicker deviation from the misaligned chunk structure. Nevertheless, they did not completely overcome the initial instruction, nor did they perform the sequences as a single chunk at the end of training. Considering that the chunk structure crystallized in the last four days of training, these results demonstrate the formation of a stable motor habit that is still influenced by the initial instruction.

Spontaneously emerging chunk structures

To investigate how participants would spontaneously chunk the sequences, we tested an additional control group (N=8), which did not receive any explicit chunk training.

Rather, participants were presented with the entire sequences on the first day and had
to memorize them without any reference to chunks (see Methods for details). Even
though memorization was more difficult, the control group did not differ significantly from
the experimental groups in terms of their explicit knowledge on day 4 ($t_{(36)}$ = 1.288, p =
0.206), or in their overall MT across training (main effect of group: $F_{(1,38)}$ = 0.101, p =
0.753; interaction between group and day ($F_{(1,38)}$ = 1.387, p = 0.168).

Similar to the experimental groups, the control group initially subdivided the
sequences into small chunks and then slowly combined them into larger chunks. The
distance to a single chunk structure decreased significantly over days ($F_{(12,84)}$ = 17.977,
p < 1e-16), and reached a level that was not statistically different from the experimental
participants on the last day of training ($t_{(38)}$ = -0.940, p = 0.353). Interestingly, on the first
day, the control group performed the sequences closer to the misaligned chunk
structure than to the aligned chunk structure ($t_{(7)}$ = -2.799, p = 0.027). With training,
participants then moved closer to the aligned chunk structure, as indicated by a
significant change in the difference between the distance to the aligned and misaligned
chunk structure across days ($F_{(12,84)}$ = 5.303, p < 1e-16). The control group also showed
clear crystallization over time (see Figure 5d&e). Compared to the experimental groups,
control participants showed a higher day-to-day and trial-by-trial change in the
beginning of training, which then reduced more quickly (Group x Day interaction; day-to-
day: $F_{(11,330)}$ = 3.780, p = 4.003e-05; trial-by-trial: $F_{(12,456)}$ = 4.254, p = 2.167e-06). In
summary, the control group showed similar behavioral patterns to the experimental
participants, indicating that similar processes of habit formation are also at play in the
absence of explicit instructions.

Two optimization processes correlate with faster final performance
How did these changes in chunk structure determine how fast participants could
execute the sequences at the end of training? We first asked whether performing the
sequences using larger chunks would facilitate performance. For each participant, we
therefore regressed the MT for 6 sequences (last 4 days, excluding the neutral
sequence) against the corresponding distance to the single chunk structure (Fig. 6a).
The majority of the participants showed a positive relationship between the number of

chunks and MT: a one-sample t-test indicated that the individual slopes were

significantly greater than 0 (Fig. 6a, $t_{(31)}$ = 6.104, p = 4.560e-07). This significant

relationship was also found for the control participants (Fig. 6b, $t_{(7)}$ = 3.429, p = 0.006).

Thus, performing the sequences with fewer chunks led to better performance.

Secondly, we investigated whether performing the sequences in alignment with

the biomechanical constraints was also beneficial. We regressed the MT for 6

sequences in the last four days of training against the corresponding distance to the

aligned chunk structure. On average the individual slopes again were significantly

greater than 0, both for the experimental (Fig. 6c; $t_{(31)}$ = 2.220, p = 0.017), and control

group (Fig. 6d, $t_{(7)}$ = 2.720, p = 0.015). Finding a better way of chunking (for the same

number of chunk boundaries) therefore also improved performance.

Figure 6. Relationship between the distance to the aligned/single chunk structure and MT. (a) *Scatterplot between the normalized (per subj.) distance to a single chunk and normalized MT in the last four days of practice. A separate regression line is fitted to the 6 sequences for each participant. Red dots indicate sequences with aligned instructions, blue dots sequences with misaligned chunking instructions.* **(b)** *Same as* **a** *but for the control group.* **(c&d)** *same as a & b but for the normalized distance to the aligned chunk structure.*

To visualize the relationship between the chosen chunk structure and the MT in
the last four days of training, we plotted the MT and chunk structure for each sequence
and participant in the 2-dimensional space defined in Fig. 5a (Fig. 7). This visualization

clearly shows the performance benefit from being closer to a single chunk as well as

being closer to the aligned chunk structure.

Overall, these results suggest that the two optimization processes - joining

chunks and aligning the remaining chunk boundaries with biomechanical constraints -

positively influence participants' ultimate performance. Furthermore, sequences for

which participants could not develop a better way of chunking were performed

substantially slower.

Figure 7. Relationship between chunking and speed (days 11-14). The x-axis indicates the distance to a single chunk and the y-axis the relative distance to the two instructed chunk structures. Each data point indicates the average chunk structure and MT of a single sequence and participant in the last four days of training. The diameter of each circle represents the MT with larger circles indicating slower performance.

The amount of musical training did not systematically affect chunk behavior

370 One possible confound is the amount of musical training that participants' were exposed

371 to prior to the experiment. We found that participants with piano experience (55%)

372 performed the sequences faster compared to the non-experienced participants ($t_{(38)}$ =

373 6.297, p = 2.227e-07). This is perhaps not surprising, given the similarity of our task with

374 playing the piano. Furthermore, the number of practice years significantly correlated

375 with MT ($t_{(38)}$ = -3.850, p = 4.401e-04). Importantly, however, the amount of participants'

376 prior musical experience neither influenced the distance to the instructed chunk

377 structure in the last week of training ($t_{(30)}$ = -0.291, p = 0.773; practice years: $t_{(30)}$ =

378 0.059, p= 0.954) nor the distance to a single chunk ($t_{(38)}$ = -0.602, p = 0.551; practice

379 years: $t_{(38)}$ = -0.380, p = 0.706). Therefore, musical training did not seem to have a

380 qualitative influence on participants' chunking behavior.

Discussion

382  In this study, we utilized chunking as a tool to investigate the role of motor habits

383 in skill learning. We influenced the structure of the initial declarative sequence

384 representation by manipulating how participants memorized them (Park, Wilde, & Shea,

385 2004). By experimentally imposing two different chunk structures on the same physical

386 sequence, we could make causal inferences on the effects of cognitive chunking on

387 motor skill development. This is an important advance over previous observational

388 studies (Ramkumar et al., 2016; Wright, Rhee, & Vaculin, 2010; Wymbs et al., 2012),

389 which did not experimentally control how participants chose to chunk the sequence.

390  This paradigm yielded three main results. First, consistent with previous studies

391 (de Kleine & Verwey, 2009; Verwey et al., 2010, 2009; Verwey & Dronkert, 1996), our

392 data demonstrate that a stable chunking pattern can be induced through cognitive

393 manipulations during sequence learning. Importantly, participants did not completely

394 overcome this imposed chunk structure, even after 2 weeks of additional training.

395 Participants' chunk structure crystallized towards the end of training, making it unlikely

396 that the influence of the initial instruction would disappear completely with longer

397 practice. Finally, the chunking structure remained stable, even when the task changed

398 from a memory-guided to a stimulus-guided task. Thus, the initial instruction led to the

formation of specific motor patterns that were still clearly measurable after three weeks
of training.

Second, we tested whether this stable pattern of chunking could be considered a
motor habit. To do so, we designed two different ways of instructing the sequence, one
aligned and the other misaligned with its biomechanical requirements. This manipulation
either facilitated or impeded performance in the first two weeks of practice. We could
show that participants did not overcome the misaligned structure completely, even
though it was detrimental to their performance. Thus, the stable chunking pattern meets
the requirements (as laid out in our definition) for being called a motor habit. Therefore,
we believe that studying sequential chunking can provide valuable insights into the
neural systems underlying motor habits. Indeed, it has recently been suggested that
chunking plays an integral role in the formation and expression of habits (Dezfouli,
Lingawi, & Balleine, 2014; Graybiel, 2008) and is neurally represented in the dorsal
lateral striatum as action "start and stop signals" (Barnes, Kubota, Hu, Jin, & Graybiel,
2005; Graybiel, 1998; Jin, Tecuapetla, & Costa, 2014; Smith & Graybiel, 2013a, 2014).

Finally, our results also indicate that the "bad" habit was not completely
immutable. Participants were able to modify the misaligned chunk structure, and did so
more rapidly than the aligned chunk structure. As a consequence, the performance
detriment imposed by the misaligned instruction was no longer significant on the group
level in the last week of training.

We identified two ways by which participants overcame the limitation induced by
the bad habit. After initially breaking up the instructed sequences into 5 chunks on
average, participants then joined chunks together, decreasing the amount of additional
time spent on chunk boundaries. While previous research has suggested that the size
of chunks increases with training, these findings were usually conflated with the overall
speed of the action (Solopchuk, Alamia, Olivier, Ze, & Zénon, 2016; Song & Cohen,
2014; Wymbs et al., 2012). Using a Bayesian model to assess chunk structure
independent of performance, we could demonstrate a positive relationship between
chunk concatenation and execution speed, both in the experimental as well as in the
control group that developed a chunking strategy without explicit instructions. However,

our results also indicate that participants did not merge all sequences into a single

chunk after 3 weeks of training, but on average subdivided each sequence into 3-4

chunks. This suggests that the number of motor actions that can be joined in a single

chunk may be limited (Langan & Seidler, 2011; Ramkumar et al., 2016; Verwey &

Eikelboom, 2003; Verwey, Lammens, & Van Honk, 2002).

A second (and novel) finding was that participants also optimized performance by

rearranging chunk boundaries in a biomechanically efficient manner. Consistent with

our prediction based on the difficulty of individual digit transitions, placing chunk

boundaries at digit transitions that take more time to execute resulted in faster

performance for the full sequence. This optimization process was also observable in the

control group that memorized and practiced sequences on their own terms.

Conversely, we observed that sequences that were not chunked in line with

these strategies were performed slower. Therefore, if a more beneficial way of chunking

was not found, participants still showed a detriment, suggesting that other learning

mechanisms cannot fully make up for a persistent bad habit. Considering that

participants' behavior became highly invariant in the last week of practice, we predict

that some bad habit will remain and continue to influence participants' performance

even after prolonged training.

In many motor tasks, there are numerous strategies and processes that can lead

to excellent performance (Verstynen et al., 2012; Verwey et al., 2010). Examining

Figure 7, one can observe that the shortest MTs were achieved anywhere in the space

between the aligned and single chunk structure. Occasionally, good performance was

also reached in other positions in chunk space. Participants adopted quite idiosyncratic

chunk structures for each sequence at the end of training. This suggests that there may

be considerable inter-individual variability in which technique works best for reaching a

high level of performance. While we based our biomechanical constraint estimates on a

representative sample, it might not perfectly reflect the constraints experienced by each

participant. Alternatively, a number of ways of chunking may work approximately equally

well, such that the cost of changing an established habit may outweigh the small benefit

that could be gained from changing the structure. A similar observation can be made in

sports, where even top-ranked athletes use slightly different techniques to reach similar

levels of performance. This variation may reflect individual biomechanical differences

but also differences in instruction and training combined with subsequent habit

formation.

The establishment of a novel experimental paradigm to study motor habit

formation will allow us to explore ways to encourage learners to abandon or change a

current habit. While our attempt at accelerating this process by changing the task from a

memory-based to a stimulus-based task was ultimately not successful, there are many

other techniques that would be possible. In many disciplines, teachers have developed

ways to help students overcome bad habits. For instance, the Hanon piano exercise

helps students play difficult passages of a musical piece by breaking up learned

phrases into new chunks to explore different rhythms. Playing a passage slower than

intended has also been suggested to break bad habits (Chang, 2004). Overall, the

general advice from the diverse literature on learning piano is to diversify training and to

practice with careful awareness to prevent bad habits from forming (Sadnicka,

Kornysheva, Rothwell, & Edwards, 2018). This suggests that changes in context and

the exploration of novel ways of moving can aid performance and the abandonment of

bad habits.

While our experimental design enabled us to manipulate participants' habits in a

laboratory setting, sequence learning only captures a specific aspect of motor skill

acquisition. Nevertheless, similar persistence of habits has been observed in other

motor learning paradigms (Diedrichsen, White, Newman, & Lally, 2010). In bimanual

coordination, for instance, Park et al. (2013) showed that an acquired pattern stayed

remarkably stable even over 8 years of not performing the task.

The current study shows that motor habits can be cognitively induced and can

remain stable for extended time periods, even though they may prevent further

performance gains. Furthermore, the study provides the first insights into the learning

processes that are involved in overcoming a detrimental habit. Our experimental

paradigm allows the further study of how we can aid the abandonment of bad habits.

488   **Methods**

Participants

Forty neurologically healthy participants were recruited for the study (30 females; ages:

19 to 33). Thirty-two were randomly split into two experimental groups and the

remaining eight participants were assigned to the control group. All participants were

right-handed based on the Edinburgh Handedness Inventory and completed informed

consent. On average, participants had received 4.68 ($\pm$ 5.55) years of musical training,

with 55% percent having played the piano for more than half a year. The study protocol

was approved by the ethics board of the University of Western Ontario.

Apparatus

A custom-built five-finger keyboard was used. The keys were not depressible, but were

equipped with a force transducer (FSG-15N1A, Sensing and Control, Honeywell)

underneath each key which reliably measured participants' isometric force production

with a repeatability of <0.02N and a dynamic range of 16N (Wiestler & Diedrichsen,

2013; Wiestler, Waters-Metenier, & Diedrichsen, 2014; Yokoi et al., 2017). The signal

was amplified and sampled at 200 Hz.

Discrete sequence production task

We used a discrete sequence production task (DSP), in which participants had to

execute sequences of 2, 3, or 11 keypresses as fast as possible while keeping their

error rate under 15% within each block. A trial was termed erroneous if participants

pressed a wrong key anywhere within the sequence. No pause between presses was

required and thus some co-articulation between fingers emerged with faster execution.

A finger press was detected when the given finger produced a force above 3N.

Subsequently, a finger was detected as released when the force of the same finger fell

below 1.5N. In order for a subsequent finger to be registered as pressed the previous

finger had to be released. This rule prevented participants to press with more than 2

fingers at once. The force magnitude applied to each key was represented by 5 lines on

an LCD monitor, with the height of the line representing the force in the corresponding

finger. A white asterisk (memory-guided conditions) or digit (cued condition) for each

finger press was presented above the lines. Immediately after the press threshold was

reached, participants received visually and auditory feedback. If the press was correct,
the white cue changed its color to green and a sound was presented. If the press was
incorrect, the cue turned red and a lower-pitch sound was presented. After each trial,
participants received points based on their accuracy and movement time (MT; the time
between the first press and last release). Correct sequences performed faster than the
current MT threshold were rewarded with 1 point. MTs that were 20% faster than the
threshold were rewarded with 3 points. Incorrect presses or MTs exceeding the
threshold resulted in 0 points. At the end of each block, participants received feedback
on their error rate, median MT, points obtained during the block, and total points
obtained during the session. In order to maintain motivation, we adjusted the MT
threshold by lowering the threshold by 500ms after each block in which the participants
performed with an error rate of 15% or lower and had a median MT faster than the
current threshold. This manipulation resulted in an approximately stable overall success
rate of 61% SD: 13% (0.27% 1pt, 0.34 % 3pt) across the entire experiment.

Study design
To impose a particular way of chunking, we first had participants memorize and perform
smaller 2-3 press chunks. These chunks were then combined to form the training
sequences. All participants were trained on the same 7 sequences, each consisting of
11 digit presses (see suppl. Table 1). Each participant completed 14 training sessions in
total: one session per day across a 3-week period (excluding weekends). Each session
lasted approximately 1 hour, excluding the two initial sessions and the last session
which took 2 hours. Participants completed at least 10 blocks of 28 trials per training
day. Each block comprised 4 repetitions of each of the 7 sequences. Each trial started
with the visual presentation of the sequence to be executed and was completed once
the participants pressed the amount of presented numbers (irrespective of whether the
pressed keys were correct or incorrect).

To verify that the chunking behavior was influenced by the instruction, we used
two different ways of chunking. We split each sequence either into one 2-digit and three
3-digit chunks (2-3-3-3, misaligned) or into three 3-digit chunks and one 2-digit chunk
(3-3-3-2, aligned). Each participant practiced half of the sequences with one chunk

structure and the other half of the sequences with the other chunk structure. This
resulted in in two different sets of chunks (suppl. Table 2). The control group did not
receive any explicit chunking instructions.

Days 1-4: Chunk induction & initial sequence learning
<u>*Experimental group*</u>: At the beginning of training the experimental groups were pre-
trained on a specific set of eleven 2- or 3-digit chunks (2 two-press chunks and 9 three-
press chunks). Participants received one of two different sets of chunks (suppl. Table
2). Each chunk was consistently associated with a letter of the alphabet (A-K).
Participants were explicitly told to learn this association. Each chunk was presented
twice in succession. In half of the blocks, on the first trial of each chunk presentation,
the numbers corresponding to the finger presses accompanied the letter on the screen
while on the second trial participants had to recall the presses solely based on the letter
(numbers were interchanged with stars). This trial order was reversed on every second
block. To ensure that participants had memorized the chunks we added speeded recall
blocks at the end of days 1 and 2. After practicing the 2-3 press chunks on day 1 and at
the beginning of day 2, participants trained on the seven 11-press sequences. Each
sequence was associated with a symbol (e.g. $; suppl. Table 1). Each sequence was
presented twice in succession and participant had to perform the sequences from
memory using the sequence cue on one trial or with the help of the chunk letters on the
next trial. We tested participants' sequence knowledge with a self-paced recall block at
the end of days 2-4 (The first two participants did not perform the recall blocks).

<u>*Control group:*</u> The control group did not receive any chunk training but instead trained
directly on the seven 11-press sequences. On the first day, the control participants
practiced the sequences using the digits presented on the screen. We matched the
amount of training across groups by ensuring that all participants were required to
produce the same number of finger presses. On the first day, the control participants
were not aware that they would have to memorize the sequences later on. On days 2-4
they were then instructed to memorize the sequences using the same sequence cues
as the experimental groups and were subsequently tested on their sequence
knowledge. The rest of the experimental design was identical for all groups.

Days 5-10: Optimization - Memory Recall

For the days 5-10 of training participants practiced exclusively on the entire eleven-

press sequences and chunks were no longer mentioned or trained on. Each sequence

was presented twice in succession and participants had to recall the sequence from

memory on both trials using the sequence cue.

Days 11-14: Optimization - Memory recall or cued presentation

During the last four days of training, half of the experimental participants performed the

sequences from memory while for the other half and for the control participants we

removed the sequence cue and presented participants with the actual numbers that

corresponded to the sequences (Fig. 1a). Participants completed an additional

generalization test on day 15. The results of this test are not reported in this article.

Biomechanical baseline study

We conducted a separate study to determine the influence of biomechanical difficulty on

the finger transition speed. 7 participants (5 females, ages: 21-27) participated in this 3-

day study. Participants executed all possible two-finger transitions (25) and three-finger

transitions (125), each 8 times per day (each sequence was presented twice in a row).

Each day participants completed 8 blocks with 150 trials each. The setup and

motivational structure were identical to the main experiment. We found that on our

device, finger repetitions (e.g. 2-2) were executed more slowly than presses of

neighboring fingers (e.g. 2-1) To press the same finger twice, the force applied to the

key had to first exceed the press threshold (3N), then go below the release threshold

(1.5N) and then cross the press threshold again. This rapid alternation of forces takes

time to produce. In contrast, for two adjacent fingers, the second finger can be already

pressed before the finger is released (have already reached the press threshold),

making it easier to rapidly produce this force pattern. Even though participants improved

the overall speed from 622ms on the first to 522ms on the third day, the 5x5 pattern of

relative IPI was stable across both participants (average correlation r = 0.689) and days

(r = 0.894).

Aligned vs. misaligned chunk structures

To determine how to design our sequences and chunk structures to aid or impede

performance we used the finding from the biomechanical baseline study that finger

repetitions (e.g. 11) are performed slower than presses of adjacent fingers (e.g. 12).

We designed the sequences such that they would include both fast transitions

(runs e.g. 123) and slow finger repetitions (113; suppl. Table 1). Depending on which

chunk structure was instructed, these transitions would either fall on a chunk boundary

or lie within a chunk. We counterbalanced this within and between participants, meaning

that each participant trained on 3 sequences with the aligned chunk structure and 3 with

the misaligned chunk structure (suppl. Table 1). One control sequence was added

which included a within-chunk run for both groups.

Statistical Analysis

We recorded and analyzed the force traces for each finger. For each trial, we calculated

the reaction time (RT, time between presentation and first crossing of the threshold),

movement time (MT, time between first press and last release) and inter-press-intervals

(IPIs; time between force peaks of two consecutive presses). All analyses were

performed using custom-written code for MATLAB (the MathWorks). We excluded trials

that contained one or more incorrect presses from our analyses, as well as trials with an

MT or a press with an IPI three standard deviations above the mean. The data were

analyzed using mixed-effects analysis of variance (mixed ANOVA), Pearson's

correlation and paired and one sample t-tests. All t-tests were two-sided. A probability

threshold of $p<0.05$ for the rejection of the null hypothesis was used for all statistical

tests. For the regression analyses as well as for calculating the MT difference between

the sequences with misaligned and aligned instruction we normalized the data for each

participant by subtracting the mean performance for each day due to a wide range of

performance speeds.

Probabilistic model for estimating chunk structure

We used an extended version of a Bayesian model of chunking behavior,

developed by Acuna et al. (2014). The algorithm uses a Hidden Markov Model to

estimate the posterior probability that a specific chunk structure is active on a given trial.

As we had 10 digit transitions, each of which could either coincide with a chunk

boundary or not, we had to consider $2^{10}-1= 1023$ possible chunk structures. Between

trials, the hidden Markov process could either stay in the same chunk structure with

probability p or jump to any other chunk structure with probability $(1-p)/1022$. The IPIs

were modeled as a Gaussian random variable, with a different mean and variance,

depending on whether the transition was within- or between-chunks. In contrast to

Acuna et al.(Acuna et al., 2014), where learning effects were removed in a

preprocessing step using a single exponential, we modeled the learning within our

model with two separate exponentials for the IPI mean. This captured the faster

reduction in the between- compared to the within-chunk intervals (Fig. 2a). The

inclusion of separate learning curves for within- and between-chunk IPIs into the model

allowed us to estimate participants' chunk structure independently of the overall

performance speed. This is an important advance over previous methods that used a

constant cutoff value to distinguish between within- and between chunk intervals. For

these methods, faster performance would automatically decrease the number of chunk

boundaries detected. To confirm that our algorithm did not show this bias, we simulated

artificial data using parameter estimates for individual participants. We simulated

sequences that switched between 4 different chunk structures, each of which contained

4 chunks. Even though IPIs decreased by about 300ms with learning, the estimated

average number of chunks remained stable across the entire simulated experiment

(average distance to single chunk: 3.35).

The model did not use errors and IPIs covariance structure, as these did not

relate systematically to the imposed chunk structure even early in training. We used an

Expectation-Maximization algorithm to simultaneously estimate the posterior probability

of each chunk structure for each trial, as well as the 9 parameters of the model: 3

parameters each for the exponential curve for the within- and between-chunk IPIs, 1

variance parameter for each, and the transition probability p.

As a preprocessing step, we regressed the IPIs for each subject against the

average biomechanical profile, which was estimated as the average IPI profile for all

possible 2 digit-presses from our biomechanical baseline experiment (Fig. 1b). The

fitted values were removed from the IPIs. Removing temporal regularities that could be

modeled with biomechanics alone should result in chunking estimates that more closely

reflect cognitive and learning influences. Qualitatively comparable results were also

obtained using the raw IPIs, without biomechanical factors removed.

Expected distance

We quantified how much participants changed their chunking behavior over time by

calculating the expected distance between two estimated chunk structures. The

distance between two chunk structures, *d(i,j)*, was defined as how many of the 10

transitions would have to change from a chunk boundary to a non-boundary (and vice

versa) to transform one structure into the other (for an example, see Fig. 4b). A distance

of 0 would indicate no change and the average distance between two randomly chosen

chunk structures is 5. Because we did not know for certain which chunk structure

participants adopted in each trial, we calculated the expected distance. For this, we first

calculated a 1023 X 1023 matrix containing the distances between any chunk structure

i, and chunk structure *j*. From the posterior probability distribution, we could then derive

how likely each of these chunk structure changes was, *p(i,j)*. The expected value of the

distance was then simply calculated as

$E(d) = \sum_{i=1}^{1023} \sum_{j=1}^{1023} p(i,j)d(i,j).$

Code availability

Bayesian algorithm code available on GitHub:

https://github.com/jdiedrichsen/chunk_inference.

Custom MATLAB code is available from the corresponding author on request.

Data availability

The datasets generated during and analyzed during the current study are available from

the corresponding author on request.

Supplementary Table 1. Sequences and chunk structures for the experimental group. *Displayed are the 7 sequences used together with the associated sequence cue. The black vertical lines indicate the chunk boundaries that were imposed. Chunk structures were either aligned with the biomechanical requirements (red) or misaligned (blue). The last sequence (green) was included as a control sequence that was chunked either with a 3-3-3-2 or 2-3-3-3 structure but performance wise should lead to similar speeds as for both chunk structures the boundaries were placed at biomechanically slow transitions. This sequence was not included in the analyses. Half of the participants were instructed based on "Group1" assignment of aligned and misaligned chunk structures and the other half based on "Group2" assignment*

Sequence	Group 1	Group 2
?	1 2 3|5 1 3|3 2 1|3 4	1 2|3 5 1|3 3 2|1 3 4
+	4 3 5|5 1 2|3 5 3|3 2	4 3|5 5 1|2 3 5|3 3 2
%	4 3 5|5 1 2|3 5 3|3 2	4 3|5 5 1|2 3 5|3 3 2
$	3 2|1 1 3|5 1 2|3 4 3	3 2 1|1 3 5|1 2 3|4 3
&	3 2|1 1 3|5 1 3|4 4 3	3 2 1|1 3 5|1 3 4|4 3
@	3 4|1 1 3|5 1 3|4 4 3	3 4 1|1 3 5|1 3 4|4 3
# (control sequence)	3 4|1 2 3|5 1 2|3 4 3	3 4 1|2 3 5|1 2 3|4 3

|Chunk Boundary ━Aligned ━Misaligned

***Supplementary Table 2. Instructed chunks and sequences for the experimental group.** The tables depict the finger presses (1-5) that were associated with the chunk cues (A-K) and the chunks that were associated with the sequences cues (symbols). Half of the participants trained with the Group 1 chunks and the other half practiced the Group 2 chunks*

Chunks Group 1				Chunks Group 2			
Chunk	Presses			Chunk	Presses		
A	3	2		A	1	2	
B	3	4		B	4	3	
C	1	1	3	C	1	2	3
D	1	2	3	D	1	3	4
E	3	2	1	E	1	3	5
F	3	4	3	F	2	3	5
G	3	5	3	G	3	2	1
H	4	3	5	H	3	3	2
I	4	4	3	I	3	4	1
J	5	1	2	J	3	5	1
K	5	1	3	K	5	5	1

Sequences Group 1					Sequences Group 2				
Seq.	Chunks				Seq.	Chunks			
?	D	K	E	B	?	A	J	H	D
+	H	K	E	B	+	B	K	H	D
%	H	J	G	A	%	B	K	F	H
$	A	C	J	F	$	G	E	C	B
&	A	C	K	I	&	G	E	D	B
@	B	C	K	I	@	I	E	D	B
#	B	D	J	F	#	I	F	C	B

691   References

Abrahamse, E. L., Ruitenberg, M. F. L., de Kleine, E., & Verwey, W. B. (2013). Control of

automated behavior: insights from the discrete sequence production task. *Frontiers in*

Human Neuroscience, *7*(March), 1–16. https://doi.org/10.3389/fnhum.2013.00082

Acuna, D. E., Wymbs, N. F., Reynolds, C. A., Picard, N., Turner, R. S., Strick, P. L., … Kording, K. P.

(2014). Multifaceted aspects of chunking enable robust algorithms. *Journal of*
Neurophysiology, *112*(8), 1849–1856. https://doi.org/10.1152/jn.00028.2014

Adams, C. D. (1982). Variations in the sensitivity of instrumental responding to reinforcer
devaluation. *The Quarterly Journal of Experimental Psychology Section B*, *34*(2), 77–98.
https://doi.org/10.1080/14640748208400878

Ashby, F. G., Ell, S. W., & Waldron, E. M. (2003). Procedural learning in perceptual
categorization. *Memory and Cognition*, *31*(7), 1114–1125.
https://doi.org/10.3758/BF03196132

Barnes, T. D., Kubota, Y., Hu, D., Jin, D. Z., & Graybiel, A. M. (2005). Activity of striatal neurons
reflects dynamic encoding and recoding of procedural memories, *437*(October).
https://doi.org/10.1038/nature04053

Bo, J., & Seidler, R. D. (2009). Visuospatial working memory capacity predicts the organization
of acquired explicit motor sequences. *Journal of Neurophysiology*, *101*(6), 3116–3125.
https://doi.org/10.1152/jn.00006.2009

Brainard, M. S., & Doupe, A. J. (2002). What songbirds teach us about learning. *Nature*,
417(6886), 351–358. https://doi.org/10.1038/417351a

Chang, C. C. (2004). *Fundamentals of Piano Practice. Brain* (3rd ed.). CreateSpace Independent
Publishing Platform. Retrieved from http://www.pianopractice.org/

Dempster, A. P., Laird, N. M., & Rubin, D. B. (1977). Maximum likelihood from incomplete data
via the EM algorithm. *Journal of the Royal Statistical Society Series B Methodological*,
39(1), 1–38. https://doi.org/http://dx.doi.org/10.2307/2984875

Dezfouli, A., & Balleine, B. W. (2012). Habits, action sequences and reinforcement learning.
European Journal of Neuroscience, *35*(7), 1036–1051. https://doi.org/10.1111/j.1460-
9568.2012.08050.x

Dezfouli, A., Lingawi, N. W., & Balleine, B. W. (2014). Habits as action sequences: Hierarchical

action control and changes in outcome value. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *369*(1655). https://doi.org/10.1098/rstb.2013.0482

Dickinson, A. (1985). Actions and Habits: The Development of Behavioural Autonomy.

Philosophical Transactions of the Royal Society B: Biological Sciences, *308*(1135), 67–78.

https://doi.org/10.1098/rstb.1985.0010

Diedrichsen, J., & Kornysheva, K. (2015). Motor skill learning between selection and execution.

Trends in Cognitive Sciences, *19*(4), 227–233. https://doi.org/10.1016/j.tics.2015.02.003

Diedrichsen, J., White, O., Newman, D., & Lally, N. (2010). Use-Dependent and Error-Based

Learning of Motor Behaviors. *Journal of Neuroscience*, *30*(15), 5159–5166.

https://doi.org/10.1523/JNEUROSCI.5406-09.2010

Ericsson, K. A., Krampe, R. T., Tesch-romer, C., Ashworth, C., Carey, G., Grassia, J., … Schneider,

V. (1993). The role of deliberate practice in the acquisition of expert performance.

Psychological Review, *100*(3), 363–406. https://doi.org/10.1037//0033-295X.100.3.363

Graybiel, A. M. (1998). The basal ganglia and chunking of action repertoires. *Neurobiology of*

Learning and Memory, *70*(1–2), 119–136. https://doi.org/10.1006/nlme.1998.3843

Graybiel, A. M. (2008). Habits, Rituals, and the Evaluative Brain. *Annual Review of Neuroscience*,

31(1), 359–387. https://doi.org/10.1146/annurev.neuro.29.051605.112851

Graybiel, A. M., & Grafton, S. T. (2015). The Striatum: Where Skills and Habits Meet. *Cold Spring*

Harbor Perspectives in Biology, *7*(8), a021691.

https://doi.org/10.1101/cshperspect.a021691

Haith, A. M., & Krakauer, J. W. (2018). The multiple effects of practice: skill, habit and reduced

cognitive load. *Current Opinion in Behavioral Sciences*, *20*, 196–201.

https://doi.org/10.1016/j.cobeha.2018.01.015

Halford, G. S., Wilson, W. H., & Phillips, S. (1998). Processing capacity defined by relational

complexity: implications for comparative, developmental, and cognitive psychology. *The*

Behavioral and Brain Sciences, *21*(6), 803–864.

https://doi.org/10.1017/S0140525X98001769

Hardwick, R. M., Forrence, A. D., Krakauer, J. W., & Haith, A. M. (2017). Skill Acquisition and

Habit Formation as Distinct Effects of Practice. *BioRxiv*. https://doi.org/10.1101/201095

Hayes, J. R. (2013). *The Complete Problem Solver*. Taylor & Francis. Retrieved from

https://books.google.ca/books?id=uqttX7YEhIMC

Hélie, S., & Cousineau, D. (2011). The cognitive neuroscience of automaticity: Behavioral and

brain signatures. *Cognitive Sciences*, *6*(1), 25–43.

Hélie, S., Waldschmidt, J. G., & Ashby, F. G. (2010). Automaticity in rule-based and information-

integration categorization. *Attention, Perception, & Psychophysics*, *72*(4), 1013–1031.

https://doi.org/10.3758/APP.72.4.1013

Jager, W. (2003). Breaking ' bad habits ': a dynamical perspective on habit. *In Human Decision*

Making and Environmental Perception: Understanding and Assisting Human Decision

Making in Real-Life Settings (L. Hendrickx, W. Jager, and L. Steg, Eds.), Liber Amicorum for

Charles Vlek, University of Groningen, Groningen, Netherlands, (January 2003), 149–160.

Jin, X., Tecuapetla, F., & Costa, R. M. (2014). Basal ganglia subcircuits distinctively encode the

parsing and concatenation of action sequences. *Nature Neuroscience*, *17*(3), 423–430.

https://doi.org/10.1038/nn.3632

Jog, M. S., Kubota, Y., Connolly, C. I., Hillegaart, V., & Graybiel, A. M. (1999). Building neural

representations of habits. *Science (New York, N.Y.)*, *286*(5445), 1745–1750.

https://doi.org/10.1126/science.286.5445.1745

Kleine, E. De, & Verwey, W. B. (2009). Representations underlying skill in the discrete sequence

production task: effect of hand used and hand position. *Psychological Research*

Psychologische Forschung, *73*(5), 685–694. https://doi.org/10.1007/s00426-008-0174-2

Kuriyama, K., Stickgold, R., & Walker, M. P. (2004). Sleep-dependent learning and motor-skill

complexity. *Learning & Memory*, *11*(6), 705–713. https://doi.org/10.1101/lm.76304

Langan, J., & Seidler, R. D. (2011). Age differences in spatial working memory contributions to

visuomotor adaptation and transfer. *Behavioural Brain Research*, *225*(1), 160–8.

https://doi.org/10.1016/j.bbr.2011.07.014

Miller, G. A. (1956). The magical number seven, plus or minus two: some limits on our capacity

for processing information. *Psychological Review*, *63*(2), 81–97.

https://doi.org/10.1037/h0043158

Moors, A., & De Houwer, J. (2006). Automaticity: A theoretical and conceptual analysis.

Psychological Bulletin, *132*(2), 297–326. https://doi.org/10.1037/0033-2909.132.2.297

Park, J. H., Wilde, H., & Shea, C. H. (2004). Part-Whole Practice of Movement Sequences.

Journal of Motor Behavior, *36*(1), 51–61. https://doi.org/10.3200/JMBR.36.1.51-61

Park, S.-W., Dijkstra, T. M. H., & Sternad, D. (2013). Learning to never forget—time scales and

specificity of long-term memory of a motor skill. *Frontiers in Computational Neuroscience*,

7(September), 1–13. https://doi.org/10.3389/fncom.2013.00111

Ramkumar, P., Acuna, D. E., Berniker, M., Grafton, S. T., Turner, R. S., & Kording, K. P. (2016).

Chunking as the result of an efficiency computation trade-off. *Nature Communications*, *7*,

12176. https://doi.org/10.1038/ncomms12176

Robbins, T. W., & Costa, R. M. (2017). Habits. *Current Biology*, *27*(22), R1200–R1206.

https://doi.org/10.1016/j.cub.2017.09.060

Sadnicka, A., Kornysheva, K., Rothwell, J. C., & Edwards, M. J. (2018). A unifying motor control

framework for task-specific dystonia. *Nature Reviews Neurology*, *14*(2), 116–124.

https://doi.org/10.1038/nrneurol.2017.146

Sakai, K., Kitaguchi, K., & Hikosaka, O. (2003). Chunking during human visuomotor sequence

learning. *Experimental Brain Research*, *152*(2), 229–242. https://doi.org/10.1007/s00221-

003-1548-8

Seger, C. A., & Spiering, B. J. (2011). A critical review of habit learning and the Basal Ganglia.

Frontiers in Systems Neuroscience, *5*(August), 1–9.

https://doi.org/10.3389/fnsys.2011.00066

Seidler, R. D., Bo, J., & Anguera, J. A. (2012). Neurocognitive contributions to motor skill

learning: The role of working memory. *Journal of Motor Behavior*, *44*(6), 445–453.

https://doi.org/10.1080/00222895.2012.672348

Smith, K. S., & Graybiel, A. M. (2013a). A dual operator view of habitual behavior reflecting

cortical and striatal dynamics. *Neuron*, *79*(2), 361–374.

https://doi.org/10.1016/j.neuron.2013.05.038

Smith, K. S., & Graybiel, A. M. (2013b). Using optogenetics to study habits. *Brain Research*,

1511, 102–114. https://doi.org/10.1016/j.brainres.2013.01.008

Smith, K. S., & Graybiel, A. M. (2014). Investigating habits: strategies, technologies and models.

Frontiers in Behavioral Neuroscience, *8*(February), 1–17.

https://doi.org/10.3389/fnbeh.2014.00039

Smith, K. S., & Graybiel, A. M. (2016). Habit formation coincides with shifts in reinforcement

representations in the sensorimotor striatum. *Journal of Neurophysiology*, *115*(3), 1487–

1498. https://doi.org/10.1152/jn.00925.2015

Solopchuk, O., Alamia, A., Olivier, E., Ze, A., & Zénon, A. (2016). Chunking improves symbolic

sequence processing and relies on working memory gating mechanisms. *Learning &*

Memory, *23*(3), 108–112. https://doi.org/10.1101/lm.041277.115

Song, S., & Cohen, L. (2014). Impact of conscious intent on chunking during motor learning.

Learning & Memory (Cold Spring Harbor, N.Y.), *21*(9), 449–51.

https://doi.org/10.1101/lm.035824.114

Verstynen, T., Phillips, J., Braun, E., Workman, B., Schunn, C., & Schneider, W. (2012). Dynamic

Sensorimotor Planning during Long-Term Sequence Learning: The Role of Variability,

Response Chunking and Planning Errors. *PLoS ONE*, *7*(10), e47336.

https://doi.org/10.1371/journal.pone.0047336

Verwey, W. B. (1996). Buffer loading and chunking in sequential keypressing. *Journal of*
Experimental Psychology: Human Perception and Performance, *22*(3), 544–562.
https://doi.org/10.1037/0096-1523.22.3.544

Verwey, W. B. (1999). Evidence for a multistage model of practice in a sequential movement
task. *Journal of Experimental Psychology: Human Perception and Performance*, *25*(6),
1693–1708. https://doi.org/10.1037/0096-1523.25.6.1693

Verwey, W. B. (2001). Concatenating familiar movement sequences: The versatile cognitive
processor. *Acta Psychologica*, *106*(1–2), 69–95. https://doi.org/10.1016/S0001-
6918(00)00027-5

Verwey, W. B., Abrahamse, E. L., & de Kleine, E. (2010). Cognitive processing in new and
practiced discrete keying sequences. *Frontiers in Psychology*, *1*(JUL), 32.
https://doi.org/10.3389/fpsyg.2010.00032

Verwey, W. B., Abrahamse, E. L., & Jiménez, L. (2009). Segmentation of short keying sequences
does not spontaneously transfer to other sequences. *Human Movement Science*, *28*(3),
348–361. https://doi.org/10.1016/j.humov.2008.10.004

Verwey, W. B., & Dronkert, Y. (1996). Practicing a Structured Continuous Key-Pressing Task:
Motor Chunking or Rhythm Consolidation? *Journal of Motor Behavior*, *28*(1), 71–79.
https://doi.org/10.1080/00222895.1996.9941735

Verwey, W. B., & Eikelboom, T. (2003). Evidence for Lasting Sequence Segmentation in the
Discrete Sequence-Production Task. *Journal of Motor Behavior*, *35*(2), 171–181.
https://doi.org/10.1080/00222890309602131

Verwey, W. B., Lammens, R., & Van Honk, J. (2002). On the role of the SMA in the discrete
sequence production task: A TMS study. *Neuropsychologia*, *40*(8), 1268–1276.
https://doi.org/10.1016/S0028-3932(01)00221-4

Verwey, W. B., & Wright, D. L. (2014). Learning a keying sequence you never executed:

Evidence for independent associative and motor chunk learning. *Acta Psychologica*, *151*,

24–31. https://doi.org/10.1016/j.actpsy.2014.05.017

Welch, L. R. (2003). Hidden Markov Models and the Baum-Welch Algorithm. *IEEE Information*

Theory Society Newsletter, *53*(4), 1,10-13. https://doi.org/10.1023/A:1007425814087

Wickens, J. R., Horvitz, J. C., Costa, R. M., & Killcross, S. (2007). Dopaminergic Mechanisms in

Actions and Habits, *27*(31), 8181–8183. https://doi.org/10.1523/JNEUROSCI.1671-07.2007

Wiestler, T., & Diedrichsen, J. (2013). Skill learning strengthens cortical representations of

motor sequences. *ELife*, *2*(2), 1–20. https://doi.org/10.7554/eLife.00801

Wiestler, T., Waters-Metenier, S., & Diedrichsen, J. (2014). Effector-Independent Motor

Sequence Representations Exist in Extrinsic and Intrinsic Reference Frames. *Journal of*

Neuroscience, *34*(14), 5054–5064. https://doi.org/10.1523/JNEUROSCI.5363-13.2014

Wong, A. L., Lindquist, M. A., Haith, A. M., & Krakauer, J. W. (2015). Explicit knowledge

enhances motor vigor and performance: motivation versus practice in sequence tasks.

Journal of Neurophysiology, *114*(1), 219–232. https://doi.org/10.1152/jn.00218.2015

Wright, D. L., Rhee, J.-H., & Vaculin, A. (2010). Offline improvement during motor sequence

learning is not restricted to developing motor chunks. *Journal of Motor Behavior*, *42*(5),

317–24. https://doi.org/10.1080/00222895.2010.510543

Wymbs, N. F., Bassett, D. S., Mucha, P. J., Porter, M. A., & Grafton, S. T. (2012). Differential

Recruitment of the Sensorimotor Putamen and Frontoparietal Cortex during Motor

Chunking in Humans. *Neuron*, *74*(5), 936–946.

https://doi.org/10.1016/j.neuron.2012.03.038

Yokoi, A., Bai, W., Diedrichsen, J., Yokoi, X. A., Bai, W., & Diedrichsen, X. J. (2017). Restricted

transfer of learning between unimanual and bimanual finger sequences. *Journal of*

Neurophysiology, *117*(3), 1043–1051. https://doi.org/10.1152/jn.00387.2016