SEASONAL CHANGES IN BIRD ASSEMBLAGES OF A MAITENAL FOREST IN NORTH PATAGONIA

Víctor R. Cueto[1,2] and Cristian A. Gorosito[1]

[1]Centro de Investigación Esquel de Montaña y Estepa Patagónica (CIEMEP), CONICET – Universidad

Nacional de la Patagonia San Juan Bosco, Roca 780, U9200 Esquel, Chubut, Argentina.

[2]Corresponding author: vcueto@conicet.gov.ar

Running head: SEASONALITY IN PATAGONIAN BIRD ASSEMBLAGES

Abstract. We evaluate seasonal variations at the community level, analyzing changes in species richness,

species composition and total abundance, and at the species level, evaluating differences in breeding

and molting seasonality among bird species in a Maitenal forest of north Patagonia. The bird assemblage

showed a low seasonal variation in richness and total abundance, but a great change in species

composition between spring-summer and fall-winter. The change in species composition promoted few

seasonal variations in richness and total abundance, because they were compensated by the presence of

abundant species that visit the area in different seasons. At the species level, resident birds and short

distance migrants tended to begin breeding earlier than long distance migrants, and birds began to molt

body and flight feathers after breeding. Therefore, we found a low overlap of these two demanding

activities. Our results highlight the importance of bird movements in response to seasonal variations in

the availability of resources, which promote migration or local displacements of birds.

Resumen. Cambios estacionales en los ensambles de aves en un Maitenal del norte de Patagonia

Evaluamos las variaciones estacionales a nivel comunitario, analizando la riqueza de especies, la

composición de especies y la abundancia total, y a nivel de las especies, evaluando las diferencias en la

estacionalidad reproductiva y de muda entre las especies de aves en un Maitenal del norte de

Patagonia. El ensamble de aves mostró pocas variaciones estacionales en la riqueza y abundancia total,

pero un notable cambio en la composición de especies entre la primavera-verano y el otoño-invierno. El

cambio en la composición de especies promovió pocas variaciones estaciones en la riqueza y la

abundancia total, porque fueron compensadas por la abundancia de las especies que visitan el área en

las diferentes estaciones. A nivel de las especies, las aves residentes y migrantes de corta distancia

tendieron a comenzar la reproducción antes que las migrantes de larga distancia, y las aves comenzaron

la muda de plumas del cuerpo y del ala al finalizar la reproducción. Por lo cual encontramos una baja

superposición de estas dos demandantes actividades. Nuestros resultados remarcan la importancia del

movimiento de las aves en respuesta a las variaciones en la disponibilidad de recursos, que promueven

la migración o el desplazamiento local de las aves.

Keywords: Argentina, breeding season, Migration, Molting season, Seasonality, Species richness, Species

composition

INTRODUCTION

Seasonal changes in the richness and abundance of bird species assemblages are characteristics of

temperate forests (Wiens 1989), in part because some species migrate to avoid harsh climate conditions

and reduced food supply during winter months (Newton 2008). However, bird assemblages have been

studied mainly during the breeding season (Marra et al 2015), ignoring the role of seasonal changes in

richness and species composition as fundamental components of the structure of bird communities.

Several studies have been conducted on the seasonality of bird assemblages in South American

temperate regions (e.g., Capurro & Bucher 1986, Marone 1992, Rozzi et al. 1996, Cueto & Lopez de

Casenave 2000, Becerra & Grigera 2005, Ippi et al. 2009, Kelt et al. 2012), showing different patterns of

seasonal changes, independent of geographic location. In the Chaco Forest (Capurro & Bucher 1986),

Monte Desert (Marone 1992), Chilean Mediterranean Scrublands (Kelt et al. 2012) and northern

Patagonian Forest (Rozzi et al. 1996, Becerra & Grigera 2005), richness values were lower during

autumn-winter than in spring-summer, because of the migration of several species. Conversely, in the

Chilean southernmost Patagonian Forest (Ippi et al. 2009) and in the Argentinean Coastal Woodlands

(Cueto & Lopez de Casenave 2000) there was little variation in richness, because bird assemblages were

composed by few migrant species. Although bird migration in South America is not considered to be as

common as it is in the North Hemisphere (Elphick 2007), studies on avian seasonality have shown the

importance of migratory species in the structure of avian assemblages across different South American

ecosystems.

General aspects about the origin and biogeography of birds inhabiting the Patagonian Forest are well

known (Vuilleumier 1985). Also, there was an increase in the studies on bird communities in different

forest types (Rozzi et al. 1996, Becerra & Grigera 2005, Grigera & Pavic 2007, Lantschner et al. 2008, Ippi

et al. 2009). However, life history aspects of bird species have been less studied. For example, despite

previous studies on patterns of feather molt in the tropics of South America (e.g., Marini & Durães 2001,

Ryder & Wolfe 2009, Silveira & Marini 2012, Jahn et al. 2016), almost nothing is known about this aspect

of bird species in the Patagonian forest. Here, we evaluate seasonal variations in richness, species

composition and total abundance at community level, and differences in breeding and molting

seasonality among bird species at species level in a Maitenal forest of North Patagonia. Considering the

climate seasonality in Patagonia, with harsher weather conditions during fall-winter than spring-

summer, we expect lower richness and total abundance and no breeding and molting activity during

autumn and winter than in spring and summer seasons.

METHODS

Study site — We conducted our study at Cañadón Florido Ranch (42° 55' S, 71° 21' W), Chubut Province,

Argentina. The vegetation of the area corresponds to the Valdivian Forest Province of the Andean

Region (Morrone 2001). We studied bird assemblages in a forest dominated by *Maytenus boaria* trees,

commonly called Maitenal (Dimitri 1972). This forest is common in valleys and slope of hills in the

eastern portion of Patagonian forest in Chubut Province, and forms part of the ecotone between the

forest and the steppe (Dimitri 1972). The forest canopy is low, averaging a height of 5 m. The understory

is mainly dominated by *Berberis microphylla* shrubs. The climate is characterized by cold and wet winters

and mild but dry summers. Most precipitation falls as rain and snow during fall and winter (April -

September). Annual mean precipitation in the region is 484 mm. Summer and winter mean

temperatures are 17° and 2.5 °C, respectively.

Bird survey — We caught birds with mist nets within a 20-ha plot (Fig. 1) from October 2015 to March

2017. Nets were 12 m long with a 38 mm mesh size and were opened for 4–5 h after sunrise. We set two

groups of 10 nets within the plot. Each group was set in different forest patches, 200 m apart (Fig. 1).

Nets within each group were set 50 to 70 m apart and opened at least once per month. The order of net

group sampling was set randomly, with 10 or 15 days between sampling events, depending on weather

conditions (i.e., avoiding rainy or windy days). Due to adverse weather conditions, we sampled birds only

once during June 2016 and February 2017, and because of logistic problems, we did not sample birds

during April and July 2016. Our sampling effort consisted of 49 net-days and 2469 net-hours. In Table 1

we indicate the number of sampling days and the net hours of sampling on each month.

We banded all birds with aluminum leg bands and took five morphological measurements: body mass,

wing length, tail length, tarsus length, and bill length (from the anterior end of the nostril to the bill tip).

We used a digital scale (± 0.1 g) to record body mass, a wing ruler (± 1 mm) for wing and tail

measurements, and a digital caliper (± 0.01 mm) for tarsus and bill measurements. We scored body molt

intensity on a 5-point scale, from none to heavy (none = no feathers molting; trace = few feathers

molting; light = involving more than one feather tract; medium = half of body feathers molting; heavy =

most/all body feathers molting; Ralph et al. 1993). We scored remige molt by noting which primary

feather and secondary feather was molting on each wing (Ralph et al. 1993). When classifying flight

feather molt, we excluded birds with adventitious molt, following Wolfe et al. (2010), who considered

symmetrical molt of the first primary feather to indicate the start of a molt cycle. Thus, we classified

birds as being in remige molt when they were molting at least one primary feather on each wing. We

scored the level of brood patch development on a 5-point scale (0 = no patch; 1 = smooth; 2 =

vascularized; 3 = wrinkled; 4 = molting; Ralph et al. 1993). We scored development of the cloacal

protuberance on a 4-point scale (0 = none; 1 = small, 2 = medium; 3 = large; Ralph et al. 1993). We

determined sex of each individual based on characteristics of the plumage (Canevari et al. 1992),

morphological differences (e.g., Cueto et al. 2015, Pyle et al. 2015), or the presence of a brood patch for

females and a cloacal protuberance for males of species with sexually monomorphic plumage.

Climate patterns — To assess the potential effect of climate on bird assemblage seasonality, we

analyzed monthly variations on temperature and precipitation at the nearest weather station to our

study site. Data were available from the weather station "Río Percy" (42° 51´ S, 71° 25´ W, belonging to

Hidroeléctrica Futaleufú), located at 9 km from Cañadón Florido Ranch. We considered only mean

temperature, because maximum mean temperature and minimum mean temperature were strongly

correlated with this variable ($r = 0.97$ and $r = 0.96$, respectively).

Data analysis — Abundance of each species was estimated using capture rate as the number of captures

per 100 net-hours (Karr 1981). Although mist net sampling could underrepresent species richness

because of a lower probability to capture canopy or understory bird species (Mallory et al. 2004), we

considered that our estimation of richness is unbiased. First, the canopy height of the woodlands in our

study area is around 5 m and our nets have a height of 3 m. Second, we have been working in the study

area since November 2013 (see Bravo et al. 2017), and every time we visited the area, we recorded all

bird species observed or heard. With the exception of raptors (e.g., *Milvago chimango*, *Polyborus*

plancus, *Bubo magellanicus*) and bird species common in open habitats (e.g., *Hymenops perspicillatus*,

Lessonia rufa, *Muscisaxicola macloviana*), we captured all species that we had recorded in the forest of

our study area. Third, the species lists we recorded are similar to others reported in studies of the

eastern zone of north Patagonian forests (e.g., Ralph 1985, Grigera & Pavic 2007).

Assemblage structure was evaluated through seasonal changes in total abundance, species richness and

species composition. Total abundance was the sum of all new individuals captured each month (i.e., we

did not include recaptured individuals during each month, but were considered when recaptured

between months). Richness was estimated as the number of species captured each month. Species

composition variations among months was evaluated through Cluster Analysis (with the statistical

program InfoStat 2009). Months were classified based on the abundance of the species, using the Bray-

Curtis dissimilarity index, and the dendrogram was constructed with the UPGMA algorithm (Sneath &

Sokal 1973). We used the Cophenetic Correlation Coefficient to measure the agreement between the

dendrogram and the original dissimilarity matrix (Sneath & Sokal, 1973). To determine the level of

dissimilarity defining groups in the dendrogram, we considered the criteria of mean dissimilarity among

all month pairs (i.e., the average of all values in the original dissimilarity matrix; e.g., Holmes, 1981).

Seasonality in richness and total abundance were evaluated through correlation analysis with monthly

variations in temperature and precipitation, using Pearson Correlation tests.

Actively breeding birds were identified by presence of a smooth or vascularized brood patch in females

or a small, medium or large cloacal protuberance in males. At the assemblage level, we estimated the

average percentage of females and males of all species actively breeding in each month, and the average

percentage of individuals of all species molting body and wing feathers in each month. We estimated the

average abundance of females and males of each species actively breeding in each month, and the

average abundance of individuals of each species molting body and wing feathers in each month.

RESULTS

We captured 25 species and a total of 969 individuals were banded (Table 1). Little seasonal variation

was evident in richness and total abundance, although climate showed a seasonal pattern, with lower

temperatures during fall and winter and higher precipitations in winter (Fig. 2). Richness was higher in

spring (October, November and December), but during other seasons show few changes (Fig. 2b). Total

abundance did not show seasonality (Fig. 2c), the lowest value was recorded in September, and during

January and February of the summer 2017 there was an abrupt increase, with more than twice the

abundance recorded during the same months of the summer 2016 (Fig. 2c). Richness was not correlated

with monthly variations in temperature and precipitation (r = 0.21, P = 0.43, n =16 and r = -0.29, P = 0.27,

n= 16, respectively). Total abundance also was not correlated with temperature and precipitation

variations (r = 0.40, P = 0.13, n = 16 and r = 0.01, P = 0.97, n = 16, respectively). In contrast, species

composition show seasonality. Cluster Analysis with a high Cophenetic Correlation Coefficients (r = 0.79)

indicates that the dendrogram is an accurate representation of the structure of the original dissimilarity

matrices. Mean dissimilarity based on species abundance dissimilarity among months determined three

groups in the dendrogram (Fig. 3). One group was constituted by spring-summer months, the second

group by winter months, and a third group formed by March and September, i.e., transitional months

from summer to autumn (March) and from winter to spring (September).

More than 35 % of all captured individuals were *Elaenia albiceps* (Table 1). This species began to be

captured during the last days of October, with captures through March (Table 1). During the second

summer we had a high capture rate of *Elaenia albiceps* (January and February 2017, Table 1). The second

most abundant species was *Zonotrichia capensis* with more than 15% of all captured individuals (Table

1). We captured two subspecies: *Zonotrichia capensis choraules* and *Zonotrichia capensis australis*. The

former was present all year-round, although more abundant during fall (mainly in May, Table 1), while

the latter was abundant during fall and winter months (Table 1). The other two more abundant species

were *Turdus falcklandii* and *Phrygilus patagonicus* (Table 1), and both were present all year-round. In

summary, these four species represent more than 70% of the annual average total abundance in the

bird assemblage.

There were another five species present only during spring-summer and three during fall-winter. Spring-

summer visitors were *Phytotoma rara*, *Diuca diuca*, *Asthenes pyrrholeuca*, *Troglodytes aedon* and

Phrygilus gayi, and fall-winter visitors were *Colorhamphus parvirostris*, *Sephanoides sephaniodes* and

Mimus patagonicus (Table 1). The rest of the species were present all year-round or had no clear

seasonal pattern. For example, *Spinus barbatus* was very abundant only during spring, and was scarce in

summer and fall and absent during winter (Table 1).

The breeding season lasted from September to March (Table 2). Males were the first individuals

captured in breeding condition. The extreme case was a *Turdus falckandii* male captured in September

(Table 2). Males of the rest of the species in breeding condition began to be captured in October (Table

2), with the exception of *Elaenia albiceps*, *Diuca diuca* and *Troglodytes aedon* males, whose first

captures began in November (Table 2). Females in breeding condition were captured beginning in

November, but *Elaenia albiceps* and *Diuca diuca* females in breeding condition were first captured in

December and January, respectively (Table 2).

Even though almost all species began molting body and wing feathers during summer months (mainly in

February and March, Table 3), we recorded *Zenaida auriculata* and *Xolmis pyrope* molting feathers in

November and December, respectively (Table 3). Two migrant species, *Troglodytes aedon* and *Diuca*

diuca molted body and wing feathers (Table 3), but we did not capture any *Elaenia albiceps* molting.

There was little overlap between breeding and molting events (Fig. 4). The percentage of breeding

females was higher in December and January, while there was a similar percentage of breeding males

from October to February (Fig. 4). In contrast, the highest percentage of individuals molting body and

wing feathers was in March (Fig. 4).

DISCUSSION

The bird assemblage at Cañadón Florido did not show a seasonal pattern in richness and total

abundance, contrary to our expectation given the strong climatic seasonality in this temperate zone of

South America. Our results contrast with seasonal patterns of community parameters in temperate

forests of Patagonia at the same latitude of our study site (i.e., lower richness and bird abundance

during autumn-winter than in spring-summer, Rozzi et al. 1996), but are similar to that reported for the

most southern forest of South America (Ippi et al. 2009). These contrasting results could be related to

differences in location, forest composition, climatic conditions and bird assemblage composition. For

example, in the southern forests of Chile one third of the bird assemblage was composed by migratory

species (Ippi et al. 2009), but about a half of the species in forests further north in Chile were found to

be migrants (Jaksic & Feinsinger 1991). In our study site, a Maitenal forest of North Patagonia in

Argentina, 42 % of the species are migrants, but there are seven species that visit the zone during spring-

summer and four that are present in fall-winter. This seasonal turnover of species that visit Cañadón

Florido results in low seasonal variation of richness. Visitors during fall-winter could be a result of the

displacement of several species from mountain forests or from the steppe. For example, Rozzi et al.

(1996) suggested that *Sephanoides sephaniodes* moves to lowland forest in winter, and in our study area

we captured *Mimus patagonicus*, which is common in the steppe (Llanos et al. 2017). Moreover, the

presence of *Colorhamphus parvirostris* during fall-winter could be related to movements from highland

forest, similar to the displacement patterns of *Sephanoides sephaniodes*. *Colorhamphus parvirostris* is a

partial migrant that has a year-round presence in the northern portion of its breeding distribution

(Chesser & Marin 1994). Another partial migrant that visits our study site during fall-winter is *Zonotrichia*

capensis australis. A similar pattern was recorded by Keve & Kovács (1971) for this subspecies in

localities near El Bolsón and Bariloche (Argentina). *Zonotrichica capensis australis* breeds mainly in

southern Argentina and Chile (King 1974, Piloni 2002, Ortiz & Capllonch 2011), and moves to northern

latitudes in winter (Ortiz & Capllonch 2011, Sagario et al. 2014). During May we had a high capture rate

of this subspecies, but during the following fall and winter months its capture rate was low. Thus, these

birds could be on passage migration to lower latitudes in May. During spring and summer, we captured

very few individuals of this subspecies in breeding condition, so at least a small population of this

subspecies potentially breeds at our study site.

Total bird abundance did not show strong seasonal changes because it was compensated by spring-

summer and fall-winter visitors. During spring and summer, *Elaenia albiceps* was the most abundant

species in the Maitenal forest at Cañadón Florido ranch. This long distance migratory species (Bravo et

al. 2017) arrived during the last weeks of October and the last individuals were captured in late March.

Its high abundance during those months is also common in the eastern and western forests of Patagonia

(e.g., Ralph 1985, Grigera et al. 1994, Rozzi et al. 1996, Jiménez 2000, Ippi et al. 2009). During fall and

winter two subspecies of *Zonotrichia capensis* were the most abundant, and their high capture rates

compensated the reduction in abundance due to departure of spring-summer visitors. *Zonotrichia*

capensis choraules was resident all year-round, but its high capture rate was between March and

August, and during the same months the partial migrant *Zonotrichia capensis australis* was also very

abundant. *Zonotrichia capensis choraules* is resident in other localities of northern Patagonia (Keve &

Kovács 1971), and both subspecies form flocks of hundreds of individuals during fall-winter in northern

Patagonia (Keve & Kovács 1971) and in the region of the Monte desert (Sagario et al. 2014, Zarco &

Cueto 2017). During the second summer of our study (mainly January and February 2017), total

abundance increased notably. This increase was due to a high capture rate of *Elaenia albiceps* and

Turdus falcklandii during that summer. Fruit abundance of *Berberis microphylla* and *Maytenus boaria*

was higher in the summer 2017 than 2016 (Gorosito C.A. & Cueto V.R., unpublished data). As these two

species consume large quantities of fruit from those plant species (Amico & Aisen 2005), the high

abundance of fruits could be the reason for their high capture rate during the summer 2017. Abundance

of *Molothrus bonariensis* was low at Cañadón Florido ranch. This species is a generalist brood parasite

that exerts significant impacts on the breeding success of its hosts (Reboreda et al. 2003). We recorded

only five nests parasitized by *Molothrus bonariensis*, out of the 145 nests we found of 12 species

(Gorosito C.A., unpublished data). Notably, all parasitized nests were of *Diuca diuca*. Therefore, at our

study site, the effect of brood parasitism on breeding success of birds could be lower than that reported

for northern localities (Reboreda et al. 2003).

Resident birds and short distance migrants tend to begin breeding earlier than long distance migrants

(Murphy 1989). In Cañadón Florido we found the same pattern, since *Turdus falcklandii* males were in

breeding condition as early as September, and males of the other resident and several short distance

migrant species were in breeding condition in October. In contrast, captures of males in breeding

condition of the long distance migrant *Elaenia albiceps* began in November. Females showed the same

pattern than males, but differed over time. Females of resident and almost all short distance migrant

species were in breeding condition in November, but captures of *Elaenia albiceps* females in breeding

condition began in December. Similar differences in the beginning of reproduction between resident and

long distance migrants were found in species breeding in Monte desert (Mezquida 2002).

At Cañadón Florido, birds began to molt body and flight feathers in the last months of summer, with the

exception of *Elaenia albiceps*, which did not molt. This long distance migrant appears to molt in its

wintering areas (Pyle et al. 2015). For example, birds molting the outer primaries were captured in

Chapada dos Guimarães, Mato Grosso, Brazil (Guaraldo, A.C., personal communication). Feather molt is

an energetically expensive activity within a bird's annual cycle (Lindström et al. 1993, Murphy 1996).

Therefore, it is more common that breeding and molting overlap in the tropics (Foster 1974, Johnson et

al. 2012, de Araujo et al. 2017), whereas in temperate zones there is a low overlap between these two

demanding activities (Jenni & Winkler 1994, Murphy 1996, Jahn et al. 2017). At our study site, we found

this last pattern, as birds began to molt after breeding, and during a period of favorable climatic

conditions. Hence, at the end of the summer, food availability was presumably sufficient for birds to pay

the energetic cost of molting feathers.

In summary, the bird assemblage at Cañadón Florido showed lower seasonality in overall richness and

total abundance, because of a turnover in the composition of species between spring-summer and fall-

winter. These changes highlight the importance of bird movements in response to seasonal variation in

resource availability that promote migration or local displacements of birds.

ACKNOWLEDGMENTS

We thank the Roberts family for allowing us to work at Cañadón Florido. For constructive reviews of the

manuscript, we thank Alex Jahn and two anonymous referees. Research carried out at Cañadón Florido

was founded by the National Geographic Society (USA) and the Bergstrom Award of the Association of

Field Ornithologists (USA). Birds were captured with permission of Dirección de Fauna y Flora Silvestre,

Provincia del Chubut (Argentina).

REFERENCES

Amico, GC & MA Aizen (2005) Seed dispersal by birds in a temperate forest of southern South America:

who disperses to whom? *Ecología Austral* 15: 89–100.

Becerra, RM & D Grigera (2005) Dinámica estacional del ensamble de aves de un bosque norpatagónico

de lenga (*Nothofagus pumilio*) y su relación con la disponibilidad de sustratos de alimentación. *Hornero*

20: 131–139.

Bravo, SP, VR Cueto & CA Gorosito (2017) Migratory timing, rate, routes and wintering areas of White-

crested Elaenia (*Elaenia albiceps chilensis*), a key seed disperser for Patagonian forest regeneration. *PLoS*

ONE 12(2): e0170188.

Canevari, M, P Canevari, GR Carrizo, G Harris, J Rodríguez Mata & RJ Straneck (1991) *Nueva Guía de las*

Aves Argentinas. Fundación Acindar, Buenos Aires, Argentina.

Capurro, H.A. & EH Bucher (1986) Variación estacional en la comunidad de aves del bosque chaqueño de

Chamical. *Physis* 44: 1–6.

Chesser, RT & M Marín (1994) Seasonal distribution and natural history of the Patagonian Tyrant

(*Colorhamphus parvirostris*). *Wilson Bulletin* 106: 649–667.

Cueto, VR, SP Bravo, N Trujillo-Arias & GS Cabanne (2015) Sex determination by morphometry of adult

White-crested Elaenia (*Elaenia albiceps chilensis*). *Revista Brasileira de Ornitologia* 23: 18–24.

Cueto, VR & J Lopez de Casenave (2000) Seasonal changes in bird assemblages of coastal woodlands in

east-central Argentina. *Studies on Neotropical Fauna and Environment* 35: 173–177.

Dimitri, M.J. (1972) *La región de los bosques Andino-Patagónicos. Sinopsis general*. Colección Científica

del INTA, Buenos Aires, Argentina

de Araujo, HFP, AH Vieira-Filho, MR de V Barbosa, JAF Diniz-Filho & JMC da Silva JMC (2017) Passerine

phenology in the largest tropical dry forest of South America: effects of climate and resource availability.

Emu 117: 78–91.

Elphick, J (2007) *The atlas of bird migration: tracing the great journeys of the world's birds*. Natural

History Museum, London, UK.

Foster, MS (1974) A model to explain molt-breeding overlap and clutch size in some tropical birds.

Evolution 28: 182–190.

Grigera, D & C Pavic (2007) Ensambles de aves en un sitio quemado y en un sitio no alterado en un área

forestal del noroeste de la Patagonia argentina. *Hornero* 22: 29–37.

Grigera, D, CA Ubeda & S Cali (1994) Caracterización ecológica del ensamble de tetrápodos del Parque y

Reserva Nacional Nahuel Huapi, Argentina. *Revista Chilena de Historia Natural* 67: 273–298.

Holmes, RT (1981) Theoretical aspects of habitat use by birds. Pp 33–37 in Capen, DE (ed). *The use of*

multivariate statistics in studies of wildlife habitat. USDA Forest Service general technical report RM-87,

Vermont, USA.

INFOSTAT (2009) *InfoStat versión 2009*. Universidad Nacional de Córdoba, Córdoba, Argentina.

Ippi, S, CB Anderson, R Rozzi & CS Elphick (2009) Annual variation of abundance and composition in

forest bird assemblages on Navarino Island, Cape Horn Biosphere Reserve, Chile. *Ornitología Neotropical*

20: 231–245.

Jaksic, F M & P Feinsinger (1991) Bird assemblages in temperate forests of North and South America: a

comparison of diversity, dynamics, guild structure, and resource use. *Revista Chilena deHistoria Natural*

64: 491–510.

Jahn, AE, V Bejarano, M Benavides Guzmán, LM Brown, IC Carvalho Provinciato, J Cereghetti, VR Cueto, JI

Giraldo, V Gomez, MS Husak, H LePage, M MacPherson, MÂ Marini, MA Pizo, A Quickle, DV Roeder, JH

Sarasola & DT Tuero (2017) Molt while breeding? Lessons from New World *Tyrannus* flycatchers. *Journal*

of Ornithology 158: 1061–1072.

Jahn, AE, JI Giraldo, M MacPherson, DT Tuero, J Sarasola, J Cereghetti, DA Masson, MV Morales (2016)

Demographic variation in timing and intensity of feather molt in migratory Fork-Tailed Flycatchers

(*Tyrannus s. savana*). *Journal of Field Ornithology* 87: 143–154.

Jenni L & R Winkler (1994) *Moult and ageing in European passerines*. Academic Press, London, UK.

Jiménez, J E (2000) Effect of sample size, plot size, and counting time on estimates of avian diversity and

abundance in a Chilean rainforest. *Journal of Field Ornithology* 71: 66–87.

Johnson, EI, PC Stouffer, RO Bierregaard (2012) The phenology of molting, breeding and their overlap in

central Amazonian birds. *Journal of Avian Biology* 43: 141–154.

Karr, JA (1981) Surveying birds with mist nets. *Studies in Avian Biology* 6: 62–67.

Kelt, DA, A Engilis Jr, J Monárdez, R Walsh, PL Meserve & JR Gutiérrez (2012) Seasonal and Multiannual

Patterns in Avian Assemblage Structure and Composition in Northern Chilean Thorn-Scrub. *Condor* 114:

30–43.

Keve, A & A Kovács (1971) Notas taxonómicas sobre el Chingolo (*Zonotrichia capensis*) del Sud-oeste

Argentino. *Hornero* 11: 85–92.

King, JR (1974) Notes on geographical variation and the annual cycle in Patagonian populations of the

Rufous-collared Sparrow *Zonotrichia capensis*. *Ibis* 116: 74–83.

Llanos, FA, M Failla, GJ García, PM Giovine, M Carbajal, PM González, D Paz Barreto, P. Quillfeldt & JF

Masello (2011) Birds from the endangered Monte, the Steppes and Coastal biomes of the province of Río

Negro, northern Patagonia, Argentina. *Check List* 7: 782–797.

Lantschner, MV, V Rusch & C Peyrou (2008) Bird assemblages in pine plantations replacing native

ecosystems in NW Patagonia. *Biodiversity and Conservation* 17: 969–989.

Lindström, À, GH Visser & S Daan (1993) The energetic cost of feather synthesis is proportional to basal

metabolic rate. *Physiological Zoology* 66:490–510.

Mallory, EP, N Brokaw & SC Hess (2004) Coping with mist-net capture-rate bias: canopy height and

several extrinsic factors. *Studies in Avian Biology* 29:151–160.

Marini, MÂ & Durães R (2001) Annual patterns of molt and reproductive activity of passerines in south-

central Brazil. *Condor* 103: 767–775.

Marone, L (1992) Seasonal and year-to-year fluctuations of bird populations and guilds in the Monte

Desert, Argentina. *Journal of Field Ornithology* 63: 294–308.

Marra, PP, EB Cohen, SR Loss, JE Rutter & CM Tonra (2015) A call for full annual cycle research in animal

ecology. *Biology Letters* 11: 20150552.

Mezquida, ET (2002) Nidificación de ocho especies de Tyrannidae en la Reserva de Ñacuñán, Mendoza,

Argentina. *Hornero* 17: 31–40.

Morrone, JJ (2001) *Biogeografía de América Latina y el Caribe.* Manuales & Tesis de la Sociedad

Entomológica Aragonesa, Zaragoza, Spain.

Murphy, ME (1996) Energetics and nutrition of moult. Pp 158–198 in Carey, C (ed). *Avian energetics and*

nutritional ecology. Chapman and Hall, New York, USA.

Murphy, MT (1989) Life history variability in North American breeding tyrant flycatchers: phylogeny, size,

or ecology? *Oikos* 54: 3–14.

Newton, I (2008) *The migration ecology of birds*. Elsevier Press, London, UK.

Ortiz, D & P Capllonch (2011) La migración del Chingolo (*Zonotrichia capensis*) en Argentina. *Historia*

Natural 1: 105–109.

Piloni, G (2002) Revisión sistemática de las subespecies de *Zonotrichia capensis* (P.L.M. Muller) (Aves:

Emberizidae) y su distribución geográfica en la República Argentina. *Physis* 60: 1–23.

Pyle, P., A. Engilis Jr., and D.A. Kelt. 2015. *Manual for ageing and sexing landbirds of Bosque Fray Jorge*

National Park and North-central Chile, with notes on occurrence and breeding seasonality. Special

Publication of the Occasional Papers of the Museum of Natural Science, Louisiana State University,

Baton Rouge, USA.

Ralph, CJ (1985) Habitat association patterns of forest and steppe birds of northern Patagonia,

Argentina. *Condor* 87:471–483.

Ralph, CJ, GR Guepel, P Pyle, TE Martin & DF DeSante (1993) *Handbook of field methods for monitoring*

landbirds. US Forest Service general technical report PSW-GTR-144, Albany, USA.

Reboreda, JC, ME Mermoz, V Massoni, AA Astié, FL Rabuffetti (2003) Impacto del parasitismo de cría del

Tordo Renegrido (*Molothrus bonariensis*) sobre el éxito reproductivo de sus hospedadores. *Hornero* 18:

77–88.

Rozzi, R, JJ Armesto, A Correa, JC Torres-Mura & M Sallaberry (1996). Avifauna de bosques primarios

templados en islas deshabitadas del archipiélago de Chiloé, Chile. *Revista Chilena de Historia Natural* 69:

125–139.

Ryder, TB & JD Wolfe (2009) The current state of knowledge on molt and plumage sequences in selected

neotropical bird Families: a review. *Ornitología Neotropical* 20: 1–18.

Sagario, MC, VR Cueto & J Lopez de Casenave (2014) Movement patterns of three species of sparrows in

the central Monte desert: differences between and within species. *Emu* 114: 268–276.

Silveira, MB& Marini MÂ (2012) Timing, duration, and intensity of molt in birds of a Neotropical savanna

in Brazil. *Condor* 114: 435–448.

Sneath, PHA & RR Sokal (1973). *Numerical taxonomy. The principle and practice of numerical*

classification. WH Freeman & Co, San Francisco, USA.

Vuilleumier, F (1985) Forest Birds of Patagonia: Ecological Geography, Speciation, Endemism, and Faunal

History. *Ornithological Monographs* 36: 255–304.

Wiens, JA (1989) *The Ecology of Bird Communities*. Cambridge University Press, Cambridge, UK.

Wolfe, JD, TB Ryder & P Pyle (2010) Using molt cycles to categorize the age of tropical birds: an

integrative new system. Journal of Field Ornithology 81: 186–194.

Zarco, A & VR Cueto (2017) Winter flock structure in the central Monte desert, Argentina. *Ardea* 105:

89–97.

Table 1. Monthly abundance (individuals captured per 100 net hours) for each bird species captured during the 2015-2017 period in a Maitenal
forest at Cañadón Florido Ranch, Chubut Province, Argentina. For 'Resident status', 'Spr-Sum' indicates species that visit Cañadón Florido in
spring-summer and Fall-Win during fall-winter months. Species are listed in order of their percent contribution to average total abundance
during the study period. The symbol "−" denotes zero individuals captured. * percentage corresponds to both subspecies of *Zonotrichia capensis*
in each month.
415

Species	Resident status	Abudance (%)	2015						2016								2017	
			Oct	Nov	Dec	Jan	Feb	Mar	May	Jun	Ago	Sep	Oct	Nov	Dec	Jan	Feb	Mar
Elaenia albiceps	Spr-Sum	36.7	—	11.9	20.1	36.2	51.7	6.5	—	—	—	—	1.3	9.1	37.9	83.1	107.0	8.8
Zonotrichia capensis choraules	Resident	15.8*	1.8	0.2	1.4	1.0	4.5	14.2	24.0	1.8	3.3	—	2.2	2.0	1.5	—	9.3	3.1
Zonotrichia capensis australis	Fall-Win		—	—	0.3	—	—	7.8	48.0	9.1	8.9	0.7	1.3	—	0.5	—	—	12.5
Turdus falckandii	Resident	13.8	6.7	3.7	6.4	2.9	3.9	15.1	3.0	1.8	1.5	21.1	2.5	2.7	11.7	32.9	14.0	30.3
Phrygilus patagonicus	Resident	6.3	4.7	1.7	2.0	—	—	—	12.0	9.1	6.5	0.8	9.5	1.4	1.9	6.7	4.7	—
Spinus barbatus	Resident	4.7	5.8	5.0	2.3	0.7	—	1.9	1.0	—	—	—	8.7	13.9	5.2	—	—	1.4
Anairetes parulus	Resident	3.5	1.4	1.2	2.9	0.5	1.0	3.3	2.0	5.5	4.7	—	0.8	1.3	2.3	4.1	—	4.6
Phytotoma rara	Spr-Sum	3.3	2.0	—	—	1.3	1.1	0.8	—	—	—	—	3.0	4.1	1.4	0.7	18.6	—
Diuca diuca	Spr-Sum	2.0	1.3	2.4	1.5	1.3	0.6	3.8	—	—	0.7	—	—	7.7	0.5	—	—	—
Asthenes pyrrholeuca	Spr-Sum	2.0	3.0	0.9	1.1	1.9	—	—	—	—	—	—	1.7	5.0	6.2	—	—	—
Troglodytes aedon	Spr-Sum	1.7	—	1.9	1.8	4.6	0.6	1.0	—	—	—	—	—	1.3	1.9	0.7	—	3.1
Xolmis pyrope	Resident	1.5	—	0.6	1.3	2.7	—	—	2.0	—	2.7	0.6	1.5	3.2	—	0.7	—	—
Phrygilus gayi	Spr-Sum	1.1	5.7	2.2	1.5	0.7	—	—	—	—	—	—	—	—	1.0	—	—	—
Leptasthenura aegithaloides	Resident	1.1	0.7	0.5	0.2	1.4	1.1	0.8	—	3.6	0.6	—	0.7	—	0.5	0.7	—	—
Colorhamphus parvirostris	Fall-Win	1.0	—	—	—	—	—	—	4.0	1.8	4.0	0.4	—	—	—	—	—	—
Callipepla californica†	Resident	1.0	—	—	—	—	—	5.2	—	—	—	0.8	—	0.7	—	—	—	2.9
Zenaida auriculata	Resident	0.8	—	1.7	1.3	0.8	0.6	—	—	—	—	—	0.8	1.4	1.8	—	—	—
Colaptes pitius	Resident	0.8	—	0.8	0.5	—	—	1.9	1.0	1.8	0.7	—	—	0.7	0.5	—	—	—
Upucerthia saturatior	Resident	0.8	0.7	0.8	0.2	—	—	—	—	—	—	0.4	—	0.7	—	—	4.7	—
Curaeus curaeus	Resident	0.7	—	0.4	0.8	0.5	—	—	—	—	1.5	—	—	0.7	—	3.3	—	—
Sephanoides sephaniodes	Fall-Win	0.7	—	—	—	—	—	—	1.0	3.6	1.2	0.8	—	—	—	—	—	—
Mimus patagonicus	Fall-Win	0.5	—	—	—	—	—	—	—	5.5	—	—	—	—	—	—	—	—
Aphastura spinicauda	Resident	0.3	0.7	—	—	—	0.6	—	—	—	—	1.2	0.4	—	—	—	—	—
Sturnella loyca	Resident	0.1	—	1.2	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Agriornis lividus	Resident	0.1	—	—	—	—	—	—	1.0	—	—	—	—	—	—	—	—	—
Molothrus bonariensis	Spr-Sum	0.1	—	—	0.7	—	—	—	—	—	—	—	—	—	0.5	—	—	—
Number of sampling days			3	7	6	4	3	3	2	1	3	4	3	2	3	2	1	2
Net hours of sampling			160	346	315	149	125	107	100	55	143	278	240	110	180	85	23	51

418 Table 2. Average monthly abundance (individuals captured per 100 net-hours) of females and males in
419 breeding condition across bird species in a Maitenal forest at Cañadón Florido Ranch, Chubut Province,
420 Argentina. Species are listed following the order in Table 1. The symbol "—" denotes zero individuals
421 captured in breeding conditions.

Species	Sex	Sep	Oct	Nov	Dec	Jan	Feb	Mar
Elaenia albiceps	Female	—	—	—	1.2	5.0	4.7	0.5
	Male	—	—	2.6	6.9	16.3	20.9	—
Zonotrichia capensis choraules	Female	—	—	—	0.4	—	—	—
	Male	—	1.5	0.6	0.4	—	—	—
Zonotrichia capensis australis	Female	—	—	—	0.4	—	—	—
	Male	—	0.8	—	0.4	—	—	—
Turdus falcklandii	Female	—	—	0.1	1.9	1.8	—	0.5
	Male	0.2	2.7	1.2	2.9	0.9	—	1.4
Phrygilus patagonicus	Female	—	—	—	1.2	0.3	—	—
	Male	—	3.3	1.3	0.8	0.3	—	—
Spinus barbatus	Female	—	—	1.3	1.0	0.6	—	—
	Male	—	3.6	5.1	1.4	—	—	—
Anairetes parulus	Female	—	0.3	—	0.3	0.3	—	—
	Male	—	0.7	0.7	0.3	—	—	—
Phytotoma rara	Female	—	—	0.5	—	—	—	—
	Male	—	0.8	0.5	0.6	0.6	2.3	—
Diuca diuca	Female	—	—	—	—	0.3	—	—
	Male	—	—	4.5	0.5	—	—	—
Asthenes pyrrholeuca	Female	—	—	0.1	1.8	—	—	—
	Male	—	0.9	1.3	0.9	—	—	—
Troglodytes aedon	Female	—	—	1.4	0.2	—	—	—
	Male	—	—	0.9	0.9	0.6	2.3	—
Xolmis pyrope	Female	—	—	0.7	0.2	1.3	—	—
	Male	—	0.4	0.9	0.3	—	2.3	—
Phrygilus gayi	Female	—	—	—	0.3	0.3	—	—
	Male	—	0.6	0.9	0.9	—	—	—
Leptasthenura aegithaloides	Female	—	—	0.1	0.4	0.7	—	—
	Male	—	0.3	—	—	—	—	—
Upucerthia saturatior	Female	—	0.3	0.7	0.2	—	—	—
	Male	—	—	—	—	—	2.3	—
Curaeus curaeus	Female	—	—	0.1	0.3	0.3	—	—
	Male	—	—	0.5	0.2	0.6	—	—
Aphrastura spinicauda	Female	—	—	—	—	—	—	—
	Male	—	0.3	—	—	—	—	—

425 Table 3. Average monthly abundance of molting individuals (individuals captured per 100 net-hours) of
426 each bird species in a Maitenal forest at Cañadón Florido Ranch, Chubut Province, Argentina. Species are
427 listed following the order in Table 1. The symbol "−" denotes zero individuals captured in molting
428 conditions.
429

Species	Molt	Nov	Dec	Jan	Feb	Mar
Zonotrichia capensis choraules	Body	—	—	—	1.2	0.9
	Flight feathers	—	—	—	0.8	0.5
Zonotrichia capensis australis	Body	—	—	—	—	0.5
	Flight feathers	—	—	—	—	0.9
Turdus falcklandii	Body	—	—	—	—	9.6
	Flight feathers	—	—	2.7	1.6	5.7
Phrygilus patagonicus	Body	—	—	—	2.3	—
	Flight feathers	—	—	0.6	—	—
Spinus barbatus	Body	—	—	—	—	—
	Flight feathers	—	—	0.6	—	—
Anairetes parulus	Body	—	—	—	0.4	3.8
	Flight feathers	—	—	—	0.4	3.8
Phytotoma rara	Body	—	—	0.6	2.3	0.5
	Flight feathers	—	—	—	2.3	—
Diuca diuca	Body	—	—	—	0.4	0.9
	Flight feathers	—	—	—	0.4	1.4
Troglodytes aedon	Body	—	—	0.7	—	0.5
	Flight feathers	—	—	0.7	—	0.5
Xolmis pyrope	Body	—	—	—	—	—
	Flight feathers	—	0.2	—	—	—
Leptasthenura aegithaloides	Body	—	—	0.3	—	0.5
	Flight feathers	—	—	—	—	0.5
Callipepla califórnica	Body	—	—	—	—	2.9
	Flight feathers	—	—	—	—	2.4
Zenaida auriculata	Body	—	—	—	—	—
	Flight feathers	0.2	—	—	—	—
Upucerthia saturatior	Body	—	—	—	2.3	—
	Flight feathers	—	—	—	2.3	—
Curaeus curaeus	Body	—	—	0.6	—	—
	Flight feathers	—	—	0.6	—	—

430
431

Figure 1. Study site map showing the Maitenal forest areas (in grey) at Cañadón Florido Ranch, Chubut

Province, Argentina. The two ellipses indicate the zone where mist nets were installed in the forest area.

Figure 2. Monthly changes in mean temperature (solid line) and precipitation (dashed line) (a), richness

(b) and total abundance (individuals captured per 100 net hours) (c) in a Maitenal forest at Cañadón

Florido Ranch, Chubut Province, Argentina.

Figure 3. Cluster Analysis based on bird species abundance from a Maitenal forest (Cañadón Florido

Ranch, Chubut Province, Argentina), showing month groups. Dashed line indicates mean dissimilarity

among all month pairs.

Figure 4. Monthly changes in the percentage of individuals in breeding and molting condition in a

Maitenal forest at Cañadón Florido Ranch, Chubut Province, Argentina.

Figure 1.

Figure 2.

Figure 3.

Figure 4.