Microtubule plus-tips act as signaling hubs for positioning the cleavage furrow during cytokinesis.

Vikash Verma[1] and Thomas J. Maresca[1,2]

[1] Biology Department, University of Massachusetts, Amherst

[2] Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst

Correspondence to:
Thomas J. Maresca
Biology Department
University of Massachusetts, Amherst
Amherst, MA 01003
Phone: 413-545-0957
Fax: 413-545-3243
E-mail: tmaresca@bio.umass.edu

Character Count:

Running title: Plus-tips position the cleavage furrow

ABSTRACT

Cell division in animal cells culminates with the formation of a contractile ring that divides the

cytosol through formation of a cleavage furrow. Microtubules (MTs) are essential for furrow

positioning, but the molecular nature of MT-derived spatial signals is unresolved. In this study

essential cytokinesis regulators (the centralspindlin complex, aurora B kinase (ABK), and polo

kinase) were visualized in *Drosophila melanogaster* (*Dm*) cells and found localize to and track

MT plus-ends during cytokinesis. The RhoA GEF Pebble (*Dm* ECT2) did not robustly tip-track

but became enriched at MT plus-tips rapidly following cortical contact resulting in RhoA

activation and enrichment of myosin-regulatory light chain. Abrogation of cytokinesis regulator

tip-tracking by EB1 depletion or deletion of a novel EB1-interaction motif (hxxPTxh) in the

centralspindlin component RacGAP50C resulted in higher incidences of cytokinesis failure. We

propose that EB1-dependent, MT plus-tip-based signaling hubs recruit cortical *Dm* ECT2 upon

contact to locally activate RhoA.

INTRODUCTION

Cell division ends with the formation of an actomyosin contractile ring, positioned

midway between the segregated chromosomes. Constriction of the ring generates a cleavage

furrow that physically divides the cytosol, a process known as cytokinesis. Establishing and

maintaining the position of the cleavage furrow between daughter nuclei is vital for embryonic

development, tissue and stem cell maintenance, and preserving ploidy (Li, 2007). Furthermore,

cytokinesis failure yields tetraploid daughter cells and has been shown to potentiate

chromosomal instability, metastatic cellular behaviors, and tumorigenesis in mice (Fujiwara et

al., 2005). Therefore, a cell must ensure positioning, maintenance, and completion of the

cleavage furrow with high fidelity. In animal cells, the spindle apparatus specifies the position of

the cleavage furrow during anaphase (Rappaport, 1961), and it is widely accepted that the

cleavage plane is positioned via microtubule (MT)-dependent mechanisms (Rappaport, 1996);

however, there is lack of consensus regarding the nature and molecular composition of the MT-

derived positioning signals and how they are delivered. Three prominent models, namely astral

stimulation, central spindle stimulation, and polar relaxation have been proposed to contribute to

defining the cleavage plane (Bringmann and Hyman, 2005; Canman et al., 2003; Cao and

Wang, 1996; Crest et al., 2012; D'Avino et al., 2005; Glotzer, 2004; Hutterer et al., 2009;

Mogilner and Manhart, 2016; Nguyen et al., 2014; von Dassow, 2009). The astral stimulation

model proposes that positive furrow specifying signals are delivered to the cortex along

equatorial astral MTs positioned in between the sister chromatids (Devore et al., 1989). Based

on the assessments that equatorial astral MTs are stabilized relative to polar MTs and that

MKLP1 is a plus-end directed motor, Odell and Foe (OF) developed a computational model for

furrow positioning positing that equatorial astral MTs become stabilized at anaphase onset (Foe

and von Dassow, 2008) and, as a result, persist long enough to serve as tracks for the motor

protein MKLP1 to deliver a positive furrowing signal to the equatorial cortex (Odell and Foe,

2008). However, this assumption is not universal, and some reports indicate that contact

between the cortex and the astral MTs is important for furrow initiation, and not the stabilization

or dynamic state of MTs (Strickland et al., 2005a; von Dassow, 2009). In addition, there is

evidence that MKLP1 motor activity is dispensable for cleavage furrow initiation (Matuliene and

Kuriyama, 2002; Minestrini et al., 2003). Thus, neither the stabilization of equatorial MTs nor

MKLP1 motility are universally accepted features of the astral stimulation model.

According to the central spindle stimulation model, a diffusive positive signal originating

from the central spindle or midzone (an overlapping MT zone in the middle of the cell) specifies

the cleavage plane. Many signaling proteins, including the chromosome passenger complex

(CPC), which is comprised of four proteins: survivin, borealin, INCENP, and ABK (Carmena et

al., 2012), localize to the central spindle during anaphase (Cooke et al., 1987). After anaphase

onset, CPC has been shown to produce an ABK activity gradient emanating from the midzone

MTs (Afonso et al., 2014; Fuller et al., 2008; Kelly et al., 2007; Sampath et al., 2004), that is

generally accepted to provide a critical spatio-temporal clue for furrow positioning. An

accompanying proposition for this model is that ABK facilitates the formation of centralspindlin

oligomers at the equatorial plasma membrane to initiate cleavage furrow formation (Basant et

al., 2015).

In favor of the polar relaxation model, it has been proposed that MT-derived inhibitory

signals impede contractility near the polar cortex by blocking RhoA activity (Murthy and

Wadsworth, 2008). In *C. elegans*, the nature of this inhibitory signal has recently been shown to

be dependent upon TPXL-1 mediated activation of aurora A kinase (Mangal et al., 2018).

Another study in *Drosophila* and human cells found that inhibitory signals originate from a

kinetochore-derived PP1-Sds22 phosphatase activity gradient during anaphase (Kunda et al.,

2012; Rodrigues et al., 2015).

In addition to MTs, spatio-temporal regulation of cytokinesis is controlled by a complex

interplay of cyclin-dependent kinase 1 (CDK1), ABK, and polo kinase. High CDK1 activity during

metaphase results in phosphorylation of the centralspindlin component MKLP1, which prevents

its association with the MTs (Glotzer, 2009; Mishima et al., 2004). Once the checkpoint is

satisfied and cyclin B levels drop, reduced CDK1 activity results in dephosphorylation of

MKLP1, thereby allowing it to bind to MTs where it becomes highly enriched in the midzone.

MKLP1 has also been shown to localize at the MT plus-ends and the cell cortex (Nishimura and

Yonemura, 2006; Vale et al., 2009). However, the functional significance of MKLP1 localization

to MT plus-ends is not clear. Binding and enrichment of MKLP1 on the central spindle is further

regulated by ABK, which stabilizes the midzone localization of the centralspindlin complex via

phosphorylation of MKLP1 (Douglas et al., 2010). In many cell types, midzone localization of the

CPC is dependent on the plus-end directed motor MKLP2 (Cesario et al., 2006; Gruneberg et

al., 2004; Kitagawa et al., 2013; Nguyen et al., 2014), while its cortical localization depends on

actin binding by INCENP (Landino et al., 2017; Landino and Ohi, 2016). Although, the role of

ABK at the central spindle is well documented; its function at the equatorial cortex is less clear

and, like MKLP1, it has also been reported to localize to MT plus-tips following anaphase onset

(Vale et al., 2009).

Polo kinase plays a central role in cytokinesis, where it localizes to the midzone and has

also been shown to regulate many functions during mitosis (Brennan et al., 2007; Burkard et al.,

2009; Carmena et al., 2014; Llamazares et al., 1991; Petronczki et al., 2008). Polo kinase

phosphorylates the centralspindlin component MgcRacGAP (Ebrahimi et al., 2010), to recruit

the RhoGEF, ECT2, to the midzone (Burkard et al., 2009; Petronczki et al., 2007; Somers and

Saint, 2003). ECT2 produces RhoA-GTP at the membrane, which promotes cortical contractility

via activation of downstream actin and myosin regulatory pathways (Bement et al., 2005; Jordan

and Canman, 2012; Yuce et al., 2005). Interestingly, it was recently reported that cytokinesis

can occur normally in the absence of midzone-localized ECT2 through mechanisms that signal

to plasma membrane-associated ECT2 (Kotynkova et al., 2016).

PRC1 (Feo in *Drosophila*) and Kinesin-4 (Klp3A in *Drosophila*) are necessary for Polo

recruitment to the central spindle in *Drosophila* and mammalian cells (D'Avino et al., 2007; Neef

et al., 2007), suggesting that this mode of Polo recruitment to the central spindle might be

evolutionary conserved. However, depletion of Feo does not result in cleavage furrow initiation

or ingression defects (D'Avino et al., 2007), indicating that Polo recruitment to the central

spindle is not necessary for furrow positioning. Nevertheless, global Polo kinase activity is

essential for cytokinesis as it is required for cleavage furrow initiation (Brennan et al., 2007;

Lenart et al., 2007; Petronczki et al., 2007). Taking together, these data suggest that Polo

kinase and ECT2 localization to the midzone may be dispensable for furrow initiation.

In this study, we examined MT-based signaling to the process of cleavage furrow

positioning in *Drosophila melanogaster* (*Dm*) S2 cells. We applied live-cell TIRF microscopy to

quantitatively characterize the spatio-temporal dynamics of lynchpin cytokinesis regulators: *Dm*

MKLP1 (Pavarotti in *Dm*), MgcRacGAP (RacGAP50C/Tumbleweed in *Dm*), ABK, Polo kinase,

Dm ECT2 (Pebble in *Dm*), and active RhoA. Live-cell imaging revealed that *Dm* MKLP1,

RacGAP50C, ABK, and Polo each localize to and track astral MT plus-tips within minutes of

anaphase onset before becoming patterned onto a subset of equatorial astral MTs. We

observed rapid recruitment and subsequent amplification of cortical Rho-GEF Pebble (*Dm*

ECT2) at sites contacted by MT plus-tips, which then resulted in localized activation of RhoA at

the cortex, and accumulation of myosin-regulatory light chain (MRLC). Therefore, these

specialized MT plus-tips were deemed ''cytokinesis signaling TIPs'' (referred to hereafter as CS-

TIPs). CS-TIP assembly and signaling required global, but not midzone-localized ABK activity

while Polo kinase activity was not required for CS-TIP assembly, but was necessary for CS-

TIPs to recruit ECT2 and activate RhoA. Cortical contact by CS-TIPs was required to maintain

signaling to the contractile machinery as localized RhoA activation, and MRLC enrichment were

lost concomitant with disassembly of CS-TIP components. Interestingly, localization of *Dm*

MKLP1 and Polo kinase as well as efficient furrow initiation and high-fidelity cytokinesis required

the plus-tip tracking protein EB1. Plus-tip tracking by the centralspindlin component

RacGAP50C required a newly identified EB1-interaction motif (hxxPTxh), the deletion of which

resulted in increased rates of cytokinesis failure comparable to EB1 depletion. We propose that

CS-TIPs act as signaling hubs that deliver furrow positioning signals to the cortex via EB1-

dependent MT plus-tip localization and tracking of centralspindlin, ABK, and Polo kinase.

Cortical contact by CS-TIPs recruits *Dm* ECT2, thereby activating RhoA to aid in positioning the

cleavage furrow during cytokinesis.

RESULTS

The centralspindlin complex, ABK, and Polo kinase localize to astral MT plus-tips

following anaphase onset and become patterned onto equatorial astral MTs over time.

It has long been known that the centralspindlin complex and CPC are highly enriched in

the midzone during cytokinesis (Glotzer, 2009); however, previous studies in *Drosophila* and

mammalian cells as well as *Xenopus* embryos have reported MT plus-tip localization of MKLP1

and MgcRacGAP (centralspindlin complex) as well as the CPC component ABK (Breznau et al.,

2017; Nishimura and Yonemura, 2006; Vale et al., 2009). While there has been significant

investigation of midzone populations of cytokinesis regulators, very little attention has been

given to MT plus-end localized components. Thus, we sought to further investigate the spatio-

temporal dynamics of the tip-localization properties of centralspindlin and ABK by live-cell TIRF

microscopy of dividing *Drosophila* S2 cells expressing fluorescently tagged Pavarotti (*Dm*

MKLP1), RacGAP50C (*Dm* MgcRacGAP), and ABK.

We began by imaging MKLP1, a motor protein that belongs to Kinesin-6 family (also

known as Pavarotti in *Drosophila*). In metaphase, *Dm* MKLP1 diffusely localized throughout the

cytosol without evident spindle MT localization **(Supplementary Movie 1)**; however, within 2-3

minutes of anaphase onset, *Dm* MKLP1 localized to and tip-tracked on the plus-ends of astral

MTs (CS-TIPs) **(Figure 1A; Supplementary Figure 1E; Supplementary Movie 1)**. In

agreement with prior observations (Vale et al., 2009), *Dm* MKLP1 localized uniformly to astral

MT plus-tips, both polar and equatorial, before being lost from the polar tips and becoming

preferentially patterned onto equatorial tips ~10 minutes following anaphase onset **(Figure 1A;**

Supplementary Movie 1). We next visualized FP-tagged RacGAP50C since homodimers of

MKLP1 and MgcRacGAP interact to form a functional centralspindlin complex (Mishima et al.,

2002; Pavicic-Kaltenbrunner et al., 2007). In agreement with a recent report of MgcRacGAP

localization to the MT plus-ends in *Xenopus laevis* (Breznau et al., 2017*)*, we also observed MT

plus-end localization and tip-tracking activity of RacGAP50C during cytokinesis **(Figure 1B;**

Supplementary Movie 2). Like *Dm* MKLP1, RacGAP50C uniformly decorated polar and

equatorial astral MT plus-ends within ~2 minutes of anaphase onset before becoming patterned

onto equatorial MTs **(Figure 1B; Supplementary Movie 2).** We also observed both

RacGAP50C and *Dm* MKLP1 accumulation on midzone MTs and the equatorial cortex **(Figure**

1, A and B; Supplementary Movies 1 and 2). Thus, the two components of the centralspindlin

complex, *Dm* MKLP1 and RacGAP50C, exhibit the same dynamic localization pattern on

midzone MTs, the cortex, and astral MT plus-tips as cells progressed through anaphase and

cytokinesis. Consistent with previous reports (Landino et al., 2017; Vale et al., 2009), we also

observed ABK localization on the polar and equatorial CS-TIPs, midzone, and the equatorial

cortex **(Figure 1C).** Interestingly, ABK localization on the CS-TIPs was transient (~1 minute) in

comparison to centralspindlin components, which typically lasted for 2-10 minutes on polar MTs

and >10 minutes on equatorial MTs.

Polo kinase becomes enriched on midzones during cytokinesis and its activity is

required for furrow initiation and successful cytokinesis (Brennan et al., 2007; Burkard et al.,

2009; Carmena et al., 1998; Lenart et al., 2007; Petronczki et al., 2007). We next applied TIRF

microscopy to Polo-EGFP expressing cells to examine its spatio-temporal dynamics during

cytokinesis. As expected, Polo localized to the overlapping zone of the midzone MT array

(Figure 1D; Supplementary Movie 3). However, Polo kinase also localized to the CS-TIPs

(Figure 1D; Supplementary Movie 3) within ~2-3 minutes of anaphase onset before becoming

preferentially patterned onto equatorial astral tips ~10 minutes post anaphase onset. Polo

kinase uniformly localized along the length of MTs as cells advanced into late telophase, which

was not surprising since Polo kinase associates with interphase MTs in *Drosophila* cells

(D'Avino et al., 2007). Thus, prior to telophase, Polo kinase exhibited an identical localization

pattern as the centralspindlin complex. In cells expressing Polo-EGFP we often observed

inward moving zones of Polo-EGFP exclusion **(Figure 1D; Supplementary Movie 3)**. Since

MTs could not penetrate these exclusion zones and were pushed inward as the zones

expanded, we interpreted the Polo-EGFP exclusion zones as indicative of local regions of

actomyosin contractility. Cortical contact by Polo-positive MT plus-tips consistently triggered

local contractility at both the equatorial region and in the vicinity of polar astral CS-TIPs prior to

patterning **(Figure 1D; Supplementary Movie 3)** suggesting that CS-TIPs may spatially trigger

contractility upon cortical contact.

CS-TIPs recruit cortical *Dm* ECT2 (Pebble) upon contact and activate RhoA

The observation of Polo-EGFP exclusion zones led us to postulate that CS-TIPs induced

cortical contractility upon contact. Since cortical contractility is triggered by the GTP-bound state

of the small GTPase RhoA (Bement, 2000; Glotzer, 2004; Jordan and Canman, 2012), we next

examined the behavior of the RhoA-GEF ECT2 (Pebble in *Drosophila*) by visualizing dividing

cells co-expressing *Dm* ECT2-EGFP and Tag-RFP-α-tubulin by TIRF microscopy. During

interphase *Dm* ECT2 localized to the nucleus, but a cortical pool was also evident that remained

throughout mitosis. Following anaphase onset, *Dm* ECT2 localized to midzone MTs and

became enriched on the nearby equatorial cortex **(Figure 2A; Supplementary Movie 4)**. Unlike

the CS-TIP components, *Dm* ECT2 did not robustly tip-track on astral MTs, rather, cortical *Dm*

ECT2 co-localized with CS-TIPs within 10.5 +/- 1.0 seconds of contact **(Figure 2B)**, which was

followed by amplification of *Dm* ECT2 recruitment that peaked 1.5 ± 0.1 minutes after the initial

cortical contact by CS-TIPs **(Figure 2C; Supplementary Movie 4)**.

We next hypothesized that CS-TIP-mediated recruitment of *Dm* ECT2 should result in

local activation of RhoA. TIRF-based visualization of cells co-expressing TagRFP-T-α-tubulin

and Rhotekin, a reporter for active RhoA-GTP (Bement et al., 2005; Benink and Bement, 2005),

N-terminally tagged with EGFP clearly revealed that a subset of astral MT plus-tips locally

activated RhoA within seconds of contacting the cortex (**Figure 2, B and D; Supplementary**

Movie 5). The Rhotekin (active RhoA) signal initially appeared within 14.3 +/- 1.7 seconds of

cortical contact by the astral MT plus-tips and local enrichment peaked within 2.5 ± 0.2 minutes

of first contact **(Figure 2C).** Due to the high spatial and temporal resolution of the live-cell TIRF

microscopy, both MT plus-tip-based and midzone-derived RhoA activation signals could be

visualized in Rhotekin-expressing cells. The two activation signals often arrived simultaneously,

although MT plus-tip signaling sometimes preceded evident midzone-localized RhoA activation

(Figure 2D, Supplementary Movie 5). We further confirmed co-localization of Rhotekin (active

RhoA) with Polo-positive CS-TIPs **(Figure 2E; Supplementary Movie 6)** and *Dm* MKLP1,

demonstrating that CS-TIPs were responsible for localized RhoA activation.

During cytokinesis, activation of RhoA results in assembly and contraction of an

actomyosin ring. We reasoned that if indeed CS-TIPs were involved in RhoA activation near the

cortex, then it would result in myosin accumulation in their vicinity. To test this hypothesis, we

made a stable cell line co-expressing Polo-EGFP (an indicator of CS-TIPs) and myosin

regulatory light chain (MRLC) tagged with TagRFP-T-α-tubulin. Dual-color TIRF imaging of this

cell line revealed a strong enrichment of MRLC near the Polo-positive CS-TIPs **(Figure 2F;**

Supplementary Movie 7) that peaked within 5.5 ± 0.54 minutes of CS-TIPs contacting the

cortex **(Figure 2C).** MRLC enrichment also mediated localized cortical contractility as the CS-

TIPs were clearly pulled inward when the cortical contacts persisted over minutes leading to

coalescence of the CS-TIPs with a band of enriched MRLC localized several microns interior to

the initial CS-TIP contact points at the cortex (**Figure 2F; Supplementary Movie 7**). CS-TIP

signaling was required to maintain RhoA activation and cortical contractility as polar sites of

Rhotekin and MRLC enrichment were lost upon equatorial patterning and disassembly of polar

CS-TIPs **(Figure 2, E and F; Supplementary Movies 6 and 7)**. Identical results for the

establishment and maintenance of RhoA activation and MRLC enrichment were attained in cells

expressing *Dm* MKLP1. Altogether, these results overwhelmingly support the conclusion that

CS-TIPs recruit the Rho-GEF, *Dm* ECT2, to activate cortical RhoA, which results in localized

myosin accumulation and induction of cortical contractility.

Investigating the contribution of midzone components (ABK, polo kinase, kinesin-4,

MKLP2) to CS-TIP assembly.

Given the central role both ABK and Polo kinase play in cytokinesis, we proceeded to

investigate the contribution of ABK and Polo kinase activities to CS-TIP assembly. Wash-in

experiments with the *Drosophila* specific ABK inhibitor, Binucleine 2, and the Polo kinase

inhibitor BI 2536 (Eggert et al., 2004; Lenart et al., 2007; Smurnyy et al., 2010) were conducted

to isolate cytokinesis-specific contributions of each kinase to CS-TIP assembly. Polo-EGFP or

Dm MKLP1-EGFP cells were continuously imaged from metaphase/anaphase onset and

Binucleine 2 was directly introduced into the imaging chamber when CS-TIPs became evident.

Interestingly, wash-in of Binucleine 2 resulted in a complete loss of Polo kinase and *Dm* MKLP1

from MT plus-tips within 2.5 ± 0.43 minutes of introducing the inhibitor **(Figure 3, A and B;**

Supplementary Movies 8 and 9). However, wash-in experiments with the Polo kinase inhibitor

BI-2536, which we and others have shown effectively inhibits *Dm* Polo kinase (Carmena et al.,

2014; Kachaner et al., 2017; Lenart et al., 2007), did not affect the CS-TIP localization of either

Polo kinase or *Dm* MKLP1 **(Figure 3, C and D, Supplementary Movies 10 and 11)**. Taken

together, the data demonstrate that ABK activity is required for centralspindlin and Polo kinase

to localize to CS-TIPs while Polo kinase activity is not.

PRC1 (Feo in *Drosophila*) is a well-characterized midzone MT component that

preferentially bundles anti-parallel MTs (Bieling et al., 2010). PRC1 forms a complex with

kinesin-4 that regulates the midzone MT length and its stabilization (Bieling et al., 2010; Hu et

al., 2011; Subramanian et al., 2013; Zhu and Jiang, 2005). Interestingly, PRC1, which on its

own coats individual MTs and bundles anti-parallel MTs, becomes enriched at MT plus-ends

when associated with Kinesin-4 (Zhu and Jiang, 2005). *Dm* Kinesin-4 was next depleted from

Polo-EGFP or MKLP1-EGFP expressing cells to investigate if the PRC1-Kinesin-4 complex

contributes to CS-TIP assembly. In agreement with prior work (D'Avino et al., 2007), Polo kinase

failed to localize to midzone MTs in cells with ≥ 97% of *Dm* Kinesin-4 (Klp3A in *Drosophila*)

depleted **(Figure 3E; Supplementary Figure 1B; Supplementary Movie 12)**; however, Polo

kinase assembled normally onto CS-TIPs **(Figure 3E)**. *Dm* Kinesin-4 was next depleted from

cells expressing *Dm* MKLP1-EGFP and neither midzone MT localization nor CS-TIP localization

was affected **(Supplementary Figure 1A)**. These data support the conclusion that *Dm* MKLP1

and Polo kinase do not require the PRC1-Kinesin-4 complex to assemble onto CS-TIPs.

During cytokinesis, ABK localizes to midzone MTs, astral MT plus-tips, and the cortex;

however, the contribution of different ABK pools to furrow formation is unclear. In many model

systems, including *Drosophila*, CPC localization to the midzone is mediated by MKLP2 (Subito

in *Drosophila*) (Cesario et al., 2006; Gruneberg et al., 2004; Kitagawa et al., 2013; Nguyen et

al., 2014). To investigate whether midzone-localized ABK (mABK) contributed to CS-TIP

assembly, *Dm* MKLP2 was depleted from cells expressing MKLP1-EGFP. Depletion of *Dm*

MKLP2 resulted in a significant reduction in midzone levels of active phosphorylated ABK

(Figure 3F; Supplementary Figure 1C), but *Dm* MKLP1 still localized robustly to CS-TIPs and

cells completed cytokinesis normally **(Figure 3G),** (Ye et al., 2015). The results from the

Binucleine 2 wash-ins and *Dm* MKLP2 depletions, taken together, support the conclusion that

while global ABK activity is essential for CS-TIP assembly, the mABK pool is largely

dispensable **(Figure 3, A, B, F, and G)**. Since cortical ABK becomes evident following CS-TIP

assembly **(Figure 1C)**, we favor the interpretation that a cytoplasmic pool of active ABK

promotes assembly of CS-TIPs after anaphase onset**.**

ABK and polo kinase activities are required for RhoA activation by CS-TIPs

After establishing the core components of CS-TIPs and their function in *Dm* ECT2

recruitment, and RhoA activation, we next examined the specific roles of ABK and Polo kinase

in CS-TIP-mediated activation of RhoA by conducting wash-in experiments with Binucleine 2

and BI 2536 for specific inhibition of ABK and Polo kinase respectively. Dual color TIRF imaging

of anaphase cells expressing TagRFP-T-Rhotekin and EGFP-α-tubulin revealed a dramatic loss

of active RhoA from the vicinity of astral MT plus-tips within 3.16 +/- 0.77 minutes of introducing

Binucleine 2 **(Figure 4, A and E, Supplementary Movie 13)**. Identical results were attained in

cells co-expressing TagRFP-T-Rhotekin and the CS-TIP component *Dm* MKLP1-EGFP **(Figure**

4C, Supplementary Movie 14). Loss of localized Rhotekin and *Dm* MKLP1 from MT plus-tips

occurred concomitantly and within ~ 2-3 minutes of Binucleine 2 addition, demonstrating that

ABK activity is required for both CS-TIP assembly and downstream activation of RhoA.

Polo kinase activity is required for targeting ECT2 to the midzone by promoting a

physical association between ECT2 and centralspindlin (Burkard et al., 2009; Petronczki et al.,

2007). Thus, we next examined the effects of Polo kinase inhibition on downstream signaling to

RhoA activation by CS-TIPs. Introduction of BI 2536 to anaphase cells expressing TagRFP-T-

Rhotekin and EGFP-α-Tubulin resulted in a dramatic reduction in active RhoA enrichment near

the astral MT plus-tips within ~4 minutes of adding the inhibitor **(Figure 4, B and E;**

Supplementary Movie 15). The same experiment was also conducted in cells co-expressing

Dm MKLP1-EGFP and TagRFP-T-Rhotekin to simultaneously visualize CS-TIPs and RhoA

activation. In agreement with our prior observations (**Figure 2C**), Polo kinase inhibition did not

affect the CS-TIP localization of *Dm* MKLP1-GFP; however, Rhotekin signal was lost in the

vicinity of the *Dm* MKLP1 positive MT plus-tips **(Figure 4D; Supplementary Movie 16)**. Thus,

Polo kinase activity is required for CS-TIP signaling, but not for CS-TIP assembly. We reasoned

that since Polo kinase activity is required for CS-TIP-mediated activation of RhoA, then Polo

inhibition should alter the localization of *Dm* ECT2 in response to physical contact by CS-TIPs.

Indeed, Polo kinase inhibition resulted in the loss of localized cortical *Dm* ECT2 from the CS-TIP

contact points **(Figure 4F, Supplementary Movie 17).**

Satellite MT arrays analogous to midzone and astral MTs assemble in proximity to the

cortex

Over the course of visualizing hundreds of dividing cells by TIRF microscopy we

frequently noticed compelling MT structures that assembled in close proximity to the cortex as

cells progressed through cytokinesis. We referred to these structures as satellite MT arrays

because they formed de-novo and were not evidently linked to the spindle apparatus.

Interestingly, two classes of satellite arrays were observed, which were deemed midzone and

astral brush satellites because their behavior and organization were analogous to conventional

midzone and astral MTs respectively.

Similar to prior observations, we noted that *Dm* MKLP1 on astral MT plus-tips bundled

both parallel and anti-parallel MTs (Vale et al., 2009). Midzone satellites were comprised of anti-

parallel bundles of equatorial astral MTs that crossed paths by growing towards the cortex, often

buckling upon contact, and eventually becoming bundled by *Dm* MKLP1 and likely other cross-

linking factors such as PRC1 **(Supplementary Movie 1).** Midzone satellites assembled in close

proximity to the cortex and the ensuing cortical contractility further bundled the MTs and

enriched *Dm* MKLP1 (and Polo kinase) in the overlap regions as MTs were moved inward. The

process resulted in midzone satellites being in direct contact with the contractile machinery

throughout cytokinesis **(Figure 5A; Supplementary Movie 1).** In many cases satellite midzones

where eventually incorporated into the conventional midzone, which was typically microns away

from the cortex when contractility began.

Astral brush satellites formed de novo from cytosolic nucleation centers producing

dynamic MTs that became organized over time into branched, spindle pole-like structures with

MTs oriented toward the cortex **(Figure 5B).** While exhibiting astral-like properties, astral

brushes typically assembled ~5-10 microns away from the spindle apparatus and the MTs

comprising astral brushes did not emanate from spindle poles or centrosome. Furthermore, the

MT plus-tips of astral brushes were nearly always CS-TIPs that robustly activated RhoA,

especially prior to patterning of CS-TIPs onto equatorial MTs **(Figure 5B).**

EB1 is required for robust MT plus-tip localization of CS-TIP components and contributes

to furrow initiation and successful completion of cytokinesis

Having established that CS-TIPs act as signaling hubs to activate RhoA via recruitment

of cortical *Dm* ECT2, we next turned our attention to the mechanism by which CS-TIP

components tip-track on astral MTs. The MT plus-end-tracking protein, EB1 is known to

autonomously tip-track, and interact with other tip-tracking proteins to recruit them to

polymerizing MT plus-ends (Akhmanova and Steinmetz, 2008; Duellberg et al., 2014; Honnappa

et al., 2009; Strickland et al., 2005b). To test if EB1 was required for the MT plus-tip localization

of CS-TIPs, EB1 was depleted by RNAi in cells co-expressing TagRFP-T-α-tubulin and either

Dm MKPL1-EGFP; or Polo-EGFP. While >80% of control cells had robust CS-TIPs, EB1

depletions, which reproducibly yielded >90% knock down efficiencies **(Supplementary Figure**

1D), significantly compromised tip-tracking of both Polo kinase and *Dm* MKLP1 as <10% of

EB1-depleted cells exhibited MT plus-tip localization of CS-TIP components **(Figure 6, A and**

B; Supplementary Movie 18). If CS-TIPs make a functional contribution to the efficiency of

cytokinesis then perturbing CS-TIPs via EB1 depletion should result in cytokinesis defects. EB1

was depleted from cells expressing Tag-RFP-T-α-tubulin, EGFP-Rhotekin and subjected to

overnight imaging. Interestingly, there was a > 4-fold increase, relative to control cells, in failure

to initiate furrow ingression following anaphase onset **(Figure 6C).** Furrow initiation defects

have been previously reported in dividing sea urchin eggs upon inhibitory antibody injection

against EB1 or dominant negative EB1 truncation (Strickland et al., 2005b). Cytokinesis failure

results in binucleated cells and, accordingly, analysis of fixed cells from EB1 depletion

experiments revealed that EB1-depleted cells exhibited ~3-fold increase in the number of

binucleated cells with respect to control cells **(Figure 6D).** Thus, EB1 depletion significantly

compromised CS-TIP localization and tracking, and resulted in functional consequences as cells

did not initiate furrow ingression and failed cytokinesis at 3-4-fold higher rates than control cells.

These results are in agreement with a prior observation in HeLa cells where depletion of EB1

and EB3 caused a 5-fold increase in cytokinesis failure (Ferreira et al., 2013)

MT plus-tip localization of RacGAP50C requires a new type of EB1-interaction motif that

contributes to the efficiency of cytokinesis

Our characterization of the EB1-dependent nature of CS-TIP tracking is consistent with a

recent work in *Xenopus laevis* embryonic cells describing the plus-tip tracking behavior of the

centralspindlin component MgcRacGAP via a SxIP EB1-interaction motif during cytokinesis

(Breznau et al., 2017). The SxIP motif in *X. laevis* MgcRacGAP is not conserved beyond

Xenopus species despite the fact that our work and others demonstrate that centralspindlin

plus-tip tracking is conserved in *Drosophila* and human cells. Our group recently identified

unconventional EB1-interaction motifs in the *Drosophila* chromokinesin NOD (deemed "PT"

motifs) (Ye et al., 2018, In press) that are similar to a newly characterized MT plus-tip

localization sequence (Kumar et al., 2017; Manatschal et al., 2016; Stangier et al., 2018). From

our work and others, we propose a PT motif consensus of hxxPTxh ("h", hydrophobic a.a.; x,

any amino acid). Sequence alignments revealed a highly conserved, possibly with the exception

of *C. elegans,* putative hxxPTxh motif in the C-terminal region of MgcRacGAPs (**Figure 6E)**.

After confirming RacGAP50C localization on CS-TIPs **(Figure 1B; Supplementary Figure 1F),**

we generated a C-terminal truncation of RacGAP50C lacking the hxxPTxh motif (a.a. 530-536;

IFGPTIV), but retaining the ability to associate with *Dm* MKLP1 via its N-terminus (Mishima et

al., 2002; Pavicic-Kaltenbrunner et al., 2007). ΔC-RacGAP50C resulted in a severe reduction in

tip-tracking efficiency; however, its localization to the midzone and equatorial cortex was

unaffected **(Figure 6F; Supplementary Movie 19).**

While the ΔC-RacGAP50C truncation was informative in dissecting RacGAP50's tip-

tracking activity, attributing any observed cytokinesis defects in ΔC-RacGAP50C-expressing

cells to loss of CS-TIPs was complicated by the fact that ΔC-RacGAP50C prematurely truncates

the GAP domain and may; therefore, compromise its GAP activity, which has been shown to be

required for efficient cytokinesis (Canman et al., 2008; D'Avino et al., 2004; Minoshima et al.,

2003). Therefore, we next generated a more targeted deletion mutant of RacGAP50C in which

only 4 amino acids (*PTIV*[533-536]) were deleted from the hxxPTxh motif. The PTIV deletion does

not remove any amino acids that are required for GAP activity and structural simulations (Biasini

et al., 2014) did not reveal a significant change in the structure of the deletion mutant

(Supplementary Figure 2A). Since attempts to make stable cell lines expressing the ΔPTIV

mutant were unsuccessful, which is indicative of the mutant having a dominant negative effect,

cells were examined following transient transfections. Like the ΔC-RacGAP50C mutant, ΔPTIV-

RacGAP50C localized normally to midzone MTs and the equatorial cortex **(Figure 6G;**

Supplementary Movie 20). However, the tip-tracking activity of RacGAP50C-ΔPTIV was

compromised as the mutant exhibited tip-tracking in only ~23% of cells that were transfected

compared to ~80% of cells expressing WT RacGAP50C **(Figure 6H**). The fact that 23% of cells

expressing ΔPTIV-RacGAP50C-EGFP exhibited tip-tracking compared to <10% of EB1

depleted cells may be a consequence of homo-dimerization of the mutant with the endogenous

RacGAP50C that still possesses an intact hxxPTxh motif. Nonetheless, the fact that the mutant

acted in a dominant negative manner led us to closely examine the effects of its expression on

cytokinesis following transient transfection.

To observe the functional consequences of PTIV deletion from RacGAP50C on

cytokinesis, TagRFP-T-α-tubulin cells that were transected with ΔPTIV-RacGAP50C-EGFP

were fixed ~4 days post-transfection to quantify the number of binucleated cells. Since there

was variability in the baseline levels of binucleated cells between experiments, the fold change

in binucleated transfected cells (+EGFP) relative to binucleated non-transfected cells (-EGFP)

cells on the same coverslip was measured for each experiment. Importantly, there was no

increase in binucleated cells transfected with WT RacGAP50C-EGFP (+EGFP) compared to

non-transfected cells (-EGFP) in the same population. However, transfection of ΔPTIV-

RacGAP50C (+EGFP) resulted in a > 3.5-fold increase in binucleated cells compared to the

baseline level of binucleated cells in non-transfected cells (-EGFP) on the same coverslip

(Figure 6I). Taken together our data support the conclusions that furrow positioning signals

travel to the cortex in an EB1-dependent manner on growing astral MT plus-ends and that this

mode of delivery requires a novel and well-conserved EB1-interaction motif (hxxPTxh) in

RacGAP50C that, when deleted, results in an increased incidence of cytokinesis failure.

DISCUSSION

In this study high resolution live-cell TIRF microscopy was employed to study the

molecular mechanism of furrow establishment in *Drosophila* cells. Our data demonstrate that

key cytokinesis regulators (centralspindlin*,* ABK, and Polo kinase) localize to and track MT plus-

ends within 2-3 minutes of anaphase onset. Localization of *Dm* MKLP1 and ABK has been

observed previously in *Drosophila* cells but signaling capacity and regulation of these plus-end

populations have not been characterized. Furthermore, the behavior of centralspindlin during

cytokinesis is well-conserved as MKLP1 and MgcRacGAP localize to MT plus-tips after

anaphase onset in human cells and *Xenopus* embryos (Breznau et al., 2017; Nishimura and

Yonemura, 2006) thereby warranting further investigation. The application of high spatio-

temporal resolution TIRF-based imaging to dividing *Drosophila* S2 cells in this study represents

an ideal approach and model system for studying a conserved, but poorly characterized

phenomenon.

Here we show that CS-TIP assembly is regulated in a cell cycle-specific manner. CS-

TIPs assemble during a "sub-phase" of mitosis that has been referred to as cytokinesis phase

(C-phase) (Canman et al., 2000). Once the checkpoint is satisfied, and CDK1 activity is

reduced, centralspindlin binds to MTs, which is prevented during metaphase because of high

CDK1 activity and phosphorylation of MKLP1 by CDK1 (Figure 7, A and B). Within minutes of

anaphase onset, *Dm* MKLP1, RacGAP50C, ABK, and Polo kinase uniformly assembled on the

polar and equatorial astral MT plus-tips (CS-TIPs), (**Figure 1 and 7B**). However, within ~10

minutes CS-TIPs were lost from most of the polar astral MTs and patterned specifically on

equatorial MTs (**Figure 1, 3, and 7C**). Upon physical contact, CS-TIPs recruit cortical *Dm*

ECT2, which leads to local activation of RhoA and actomyosin ring assembly (**Figure 2 and**

7C). We further provide evidence that global ABK activity is absolutely required for CS-TIPs

assembly, while Polo kinase activity is required for signaling. Cleavage furrow positioning

signals via CS-TIPs require EB1 interaction, which is mediated by a conserved hxxPTxh motif in

RacGAP50C (**Figure 6 and 7D**).

CS-TIP- vs midzone-based signaling in positioning the cleavage furrow

MTs specify the site of furrow ingression during cytokinesis via local activation of the

small GTPase RhoA. The positioning of the contractile region has been variously proposed to

involve positive equatorial cues derived from the central spindle, astral MTs, or both; and

inhibitory factors acting at poles (Canman et al., 2003; Mogilner and Manhart, 2016). While

inferred, direct evidence of RhoA activation by astral MTs is lacking. Here we provide evidence

that cortical *Dm* ECT2 and active RhoA (Rhotekin) localize to CS-TIPs within seconds of contact

(**Figure 2**). *Dm* ECT2 amplification at CS-TIP cortical contact points may occur via a poorly

understood positive feedback mechanism (Bement et al., 2015; Graessl et al., 2017; Tyson et

al., 2003). Sustained CS-TIP signaling over a minutes time-scale further recruits myosin

(MRLC) - eventually resulting in localized cortical contractility (**Figure 2D**). Due to the high

spatial and temporal resolution of the TIRF-based imaging assays we employed, both MT plus-

tip-proximal and midzone-derived RhoA activation signals could be readily visualized in

Rhotekin-expressing cells. The CS-TIPs and midzone-derived RhoA activation signals typically

occurred simultaneously, although CS-TIPs signaling sometimes preceded evident midzone

RhoA activation (**Figure 2B**). The arrival of a midzone gradient-derived activation signal to the

cortex would be inexorably tied to variable features of the midzone MT array such as distance

from the cortex and localization/levels of any gradient-producing midzone component. To the

contrary, we propose that CS-TIP-based signaling would be more consistent than midzone-

mediated signaling since CS-TIPs depend primarily on MT polymerization rates, lengths, and

the orientation of astral MT plus-ends, which may be derived from centrosomes or spindle

poles, towards the cortex – features that are highly reproducible between anaphase spindles. In

fact, the CS-TIP mode of signaling could play a more critical role in larger cells such as those in

early embryonic divisions where the midzone MT array is long distances from the cortex.

Although it is worth noting that in large cells amplification of the astral MT array (Mitchison et al.,

2012) similar to the satellite astral brushes observed here (**Figure 5, A and B**) would also be

required for astral MTs to act over very long distances.

A midzone derived ABK gradient has been proposed for cleavage furrow positioning

(Fuller et al., 2008; Tan and Kapoor, 2011) ; and in addition to ABK, Polo kinase has also been

shown to induce furrow initiation by recruiting RhoGEF, ECT2, to the central spindle (Petronczki

et al., 2007), indicating that both ABK and Polo activity is required for furrow positioning.

However, a complete understanding of how ABK and/or Polo kinase activate RhoA to initiate

furrowing is still lacking. We show that CS-TIPs assembly and signaling was unaffected by

treatments that mis-localize midzone pools of active ABK (***Dm* MKLP2 depletion, Figure 3, F**

and G) and Polo kinase (***Dm* Kinesin-4 depletion, Figure 3E)**. While global inhibition of either

ABK or Polo kinase activities leads to severe cytokinesis defects, furrow positioning and

cytokinesis was not evidently impacted in this study by mis-localization of midzone pools of ABK

or Polo kinase **(Figure 3)**. Thus, midzone-based ABK and polo kinase activity gradients are

either dispensable for furrow positioning in *Drosophila* or redundant with CS-TIP-based furrow

positioning cues and/or other pathways. Our observation further supports a recent report where

artificial targeting of ECT2 was sufficient to restore cytokinesis, and mutations that prevented

midzone localization of ECT2 didn't result in any apparent cytokinesis defects in most of the

cells (Kotynkova et al., 2016).

Establishment and maintenance of cleavage furrow positioning signals

The formation of actomyosin ring and contractility typically lasts for minutes, while the

furrow must be positioned correctly, it must also be maintained for successful cytokinesis.

During our high-resolution imaging, we often observed ''satellite MT arrays'' that were in close

proximity to the cortex and assembled after contractility had been established. Near the

equatorial cortex, midzone satellites were built from equatorial astral MTs that became bundled

and forced into anti-parallel MT arrays by the contractile machinery. Importantly, once

contractility was established, the midzone satellites maintained direct physical contact with the

cortex such that any midzone-based cues would no longer have to act over long distances to

signal to the cortex. The other type of satellite MT array - astral brushes – also assembled after

contractility had begun, but via apparent branching MT nucleation from astral MTs. The astral

brushes, in which MT plus-ends were oriented toward the cortex, locally activated RhoA by

creating a high density of new CS-TIPs. We posit that midzone and astral brush MT satellites

could fulfill two critical roles during cytokinesis as autonomous signaling centers that can 1) act

at distances far from their analogous structures (midzones and poles/centrosomes) in the

spindle proper, and 2) maintain contractility after it has been established.

Role of EB1 and a conserved hxxPTxh motif in RacGAP50C in cleavage furrow

positioning

We found that localization of the centralspindlin complex on the CS-TIPs is dependent

on EB1. This is consistent with recent work in *Xenopus laevis* embryonic cells describing the

plus-tip tracking behavior of the centralspindlin component MgcRacGAP via a SxIP EB1-

interaction motif during cytokinesis (Breznau et al., 2017). However, the SxIP motif in *X. laevis*

MgcRacGAP is not conserved beyond *Xenopus* species despite the fact that centralspindlin

plus-tip tracking is well conserved (Nishimura and Yonemura, 2006; Vale et al., 2009). We

recently identified unconventional EB1-interaction motifs ("PT" motifs) in the *Drosophila*

chromokinesin NOD (Ye et al., 2018, In press) that are similar to a newly characterized MT tip

localization sequence (Kumar et al., 2017). Interestingly, sequence alignments revealed a highly

conserved putative hxxPTxh motif in the C-terminal region of the MgcRacGAPs. Depletion of

EB1, truncation of RacGAP50C C-terminus, and deletion of PTIV from the putative hxxPTxh

motif from the RacGAP50C yielded the same phenotypes - loss of plus-end localization and

tracking of centralspindlin in a significant majority of cells and an increase incidence of

cytokinesis failure. Our results are consistent with earlier reports where injection of an inhibitory

antibody against EB1 or dominant negative EB1 truncation into dividing sea urchin eggs

resulted in delayed furrow ingression (Strickland et al., 2005b), and reported finding of EB1 and

EB3 depletion in HeLa cells that caused a 5-fold increase in cytokinesis failure (Ferreira et al.,

2013). EB1 depletion in *Drosophila* S2 cells has also been reported to cause spindle defects

(Rogers et al., 2002), however we did not observe any obvious spindle morphology defects.

Although, we reproducibly achieved ≥ 90% EB1 depletion, this discrepancy could result from

different levels of EB1 depletion. While similar differences in the extent of mitotic phenotypes

were reported in the recent knockout studies of EB proteins in HeLa cells (McKinley and

Cheeseman, 2017; Yang et al., 2017), these studies didn't focus on cytokinesis, so it will be

worthwhile to investigate whether the hxxPTxh motif is required for centralspindlin plus-tip

localization and efficient cytokinesis in other cell types.

CS-TIP assembly, signaling, and patterning onto equatorial MTs

The observation that *Dm* MKLP1, RacGAP50C, ABK, and Polo localize to the plus-ends

of polar and equatorial astral MT tips before becoming patterned onto equatorial astral MTs

raises an important question: how do CS-TIPs selectively disassemble from the polar astral

MTs? Since we have shown that CS-TIPs activate cortical contractility, the patterning

mechanisms that disassemble polar CS-TIPs would be akin to a polar relaxation mechanism. A

recent study in *C. elegans* found that TPXL-1 (*C. elegans* TPX-2 homologue) mediated

activation of Aurora A kinase was responsible for clearing anillin and F-actin (Mangal et al.,

2018) from the polar cortex. A polar Aurora A kinase activity gradient that regulates kinetochore-

MT attachments has been visualized in *Drosophila* and mammalian cells (Chmatal et al., 2015;

Ye et al., 2015) although its role in C-phase warrants further study. Also in *Drosophila*, a

kinetochore based PP1 phosphatase activity gradient derived from a PP1-SDS22 complex was

proposed for the loss of actin and inactivation of ezrin/radixin/moesin proteins at the polar cortex

(Kunda et al., 2012; Rodrigues et al., 2015). It is entirely possible that phosphatase and/or

kinase activity gradients in *Drosophila* contribute to the loss of CS-TIPs from polar astral MTs.

Alternatively, a chromosome based RanGTP gradient has been shown to locally reduce Anillin

at the cortex (Kiyomitsu and Cheeseman, 2013), and could also impede CS-TIPs-based

signaling from the polar astral MTs as the chromosomes approach the polar cortex in late

anaphase prior to nuclear envelope reformation. RanGTP regulates many functions during cell

division, including nuclear-cytoplasmic transport and spindle assembly (Kalab et al., 2006;

Kalab et al., 2002; Wu et al., 2013), and because of this, it is difficult to isolate specific temporal

Ran-dependent functions during cytokinesis. A small molecule inhibitor, importazole (2,4-

diaminoquinazoline) has been shown to specifically disrupt RanGTP and importin-β interaction

in *Xenopus* egg extracts and cultured cells (Soderholm et al., 2011), and could potentially be

used to isolate specific temporal functions during cytokinesis. However, cytokinesis can occur in

the absence of chromatin (Alsop and Zhang, 2003; Rappaport, 1996). While we presently favor

the hypothesis that polar- or chromosome/kinetochore-derived activity gradients regulate CS-

TIP patterning, a more detailed investigation is necessary to resolve the question of how polar

CS-TIPs are selectively disassembled as cells progress through cytokinesis.

MATERIALS AND METHODS

Cloning and site directed mutagenesis

The *Drosophila* MKLP1 gene (known as Pavarotti in *Drosophila*, CG1258) and ECT2 (kwon as

Pebble in *Drosophila*, CG8114) were PCR amplified from cDNA clones RE22456 and SD01796

respectively with a 5′ SpeI site and a 3′ XbaI site. The resulting PCR products of the individual

genes were then inserted into the 5′ SpeI and 3′ XbaI sites of the pMT/V5 His-B vector

(Invitrogen) containing in-frame EGFP gene at the 3' end, cloned between 5' XbaI and 3' SacII

sites, and the CENPC promoter at the 5' end, cloned between single Kpn1 site. The regulatory

light chain of the non-muscle type 2 myosin (known as Spaghetti squash in *Drosophila*,

CG3595) was amplified from the cDNA clone LD14743. The resulting PCR product was cloned

between 5' SpeI and 3' XbaI of the pMT vector containing in-frame RFP gene at the 3' end and

Mis12 promoter at the 5'end. EGFP-Rhotekin was PCR amplified from the pCS2-eGFP-rGBD

plasmid (gift from the Bement lab (UW-Madison) via Wadsworth lab (UMass Amherst)). The

resulting PCR product was cloned between 5' XhoI and 3'ApaI sites of the pMT vector

containing CENPC promoter at the 5'end cloned between single KpnI site. A second version of

Rhotekin was also created in the pMT vector where we swapped the EGFP fluorophore with

RFP. Polo kinase under its own promoter was cloned between 5' KpnI and 3' SpeI sites of the

pMT vector containing in-frame EGFP gene, cloned between 5' SpeI and 3' ApaI sites. The

RacGAP50C (also known as Tumbleweed in *Drosophila*, CG13345) gene was amplified from

genomic DNA of *Drosophila*. The resulting PCR product was cloned in a similar fashion as

described for MKLP1. A four-amino acid deletion ($PTIV^{533-536}$) in the RacGAP50C gene was

generated by Q5 site directed mutagenesis kit (NEB, MA, USA), following manufacturer's

protocol. C-terminal truncation of the RACGAP50C gene was created by amplifying amino acids

(1-526) from the full length RacGAP50C (a.a. 1-625) gene. The resulting PCR product was

inserted into the 5′ SpeI and 3′ XbaI sites of the pMT vector containing in-frame EGFP gene at

the 3' end, and the CENPC promoter at the 5' end. See Table 1 for the primers used in cloning.

***Drosophila Schneider (S2)* cell culture and generation of transient/stable cell line**

All cell lines were grown at 25 °C in Schneider's medium (Life Technologies, Carlsbad, CA),

containing 10% heat-inactivated fetal bovine serum (FBS) and 0.5X antibiotic-antimycotic

cocktail (Gibco, Life Technologies, NY, USA). Cell lines were generated by transfecting the DNA

constructs containing the gene of interest with Effectene transfection reagent (Qiagen, Hilden,

Germany), following the manufacturer's protocol. After 4 days of transfection, expression of

EGFP/RFP tagged proteins was checked by fluorescence microscopy. To make a stable cell

line, cells were selected in the presence of Blasticidin S HCl (Thermo Fisher Scientific,

Waltham, MA) and/or Hygromycin (Sigma-Aldrich) until there was no observable cell death.

Thereafter, cell lines were either frozen down or maintained in the S2 media @ 25 °C without

Blasticidin and/or Hygromycin B. Transient cell lines were imaged after 4-5 days of transfection,

and they were not subjected to antibiotics selection. ABK-GFP and mCherry-α-tubulin cell line,

was a generous gift from Dr. Eric Griffis, University of Dundee.

RNA interference (RNAi) experiments

Around 500 base pairs of DNA templates containing T7 promoter sequence at the 5' end for *Dm*

MKLP2 (also known as Subito in *Drosophila*, CG12298), EB1 (CG3265), and *Dm* Kinesin-4

(also known as Klp3A in Drosophila, CG8590) were generated by PCR. Double-stranded RNAs

(dsRNAs) were synthesized from the respective DNA templates at 37 °C using the T7 RiboMax

Express Large Scale RNA Production System (Promega Corp., Madison, WI), following the

manufacturer's protocol. For RNAi experiments, cells at about 25% confluency were incubated

in a 35 x 10 mm tissue culture dish for an hour. Thereafter, media was carefully aspirated off the

dish and 1 ml of serum-free S2 media containing 20 *µ*g of dsRNA was added to the dish. After 1

hour, 1 ml of fresh S2 media containing FBS was added to the dish and incubated for 2-4 days

at 25 °C. See Table 2 for primer sequences.

Immunofluorescence

Drosophila S2 cells were plated on an acid-washed Concanavalin A (Sigma-Aldrich) coated

coverslips for 30 minutes, then cells were quickly rinsed with 1X BRB80 buffer. Subsequently,

cells were fixed with 10% paraformaldehyde for 10 minutes. Cells were then permeabilized with

phosphate-buffered saline (PBS) containing 1% Triton X-100 for 8 min, rinsed 3 times with PBS

plus 0.1% Triton X-100, and blocked with 5% boiled donkey serum ((Jackson Immuno Research

Laboratories, Inc., West Grove, PA) for 60 minutes. All primary antibodies were diluted in 5%

boiled donkey serum. Anti-Phospho-aurora A/B/C (Cell Signaling Technology, Danvers, MA)

was used at a concentration of 1:1000, and anti-tubulin antibody (DM1α Sigma-Aldrich) at

1:2000. All secondary antibodies (Jackson Immuno Research Laboratories, Inc., West Grove,

PA) were diluted in boiled donkey serum at 1:200. After secondary antibody incubation,

coverslips were washed 3 times with PBS plus 0.1% Triton X-100, followed by incubation with

4′,6′-diamidino-2-phenylindole (DAPI) at a concentration of 1:1000 for 10 minutes, and 3

additional washes with PBS plus 0.1% Triton X-100. Coverslips were sealed in a mounting

media containing 20 mM Tris, pH 8.0, 0.5% N-propyl gallate, and 90% glycerol. Four-color Z-

series consisting of ~30 planes at 0.2-μm intervals were acquired for GFP, RFP, Cy5, and DAPI

channels on a TIRF-Spinning Disk Confocal system assembled on a Nikon Ti-E microscope

equipped with a 100X 1.4 NA DIC Apo Oil immersion objective, two Hamamatsu ORCA-Flash

4.0 LT digital CMOS camera (C11440), 4 laser lines (447, 488, 561, 641), and a MetaMorph

software. Fluorescence intensities were obtained by drawing a 50 x 100 grid manually around

the maximum-intensity projection of the Z-series images. The local background was estimated

by placing the 50 x 100 region to a nearby place. To obtain the ratio intensities, identical regions

were drawn manually on the *Dm* MKLP1-EGFP channel and transferred to a phosphorylated

ABK (pABK) channel.

Quantitative Western blotting

Equal amount of proteins was loaded into a 10 or 8% SDS-PAGE gel. After running the gel,

proteins were transferred to a nitrocellulose membrane using the Trans-Blot Turbo transfer

system (Bio-Rad Laboratories, Inc., Hercules, CA) for 10 minutes. Subsequently, the membrane

was incubated in 5% milk (V/V made in Tris-buffered saline with 0.1% Tween) for one hour.

Following blocking, the membranes was incubated with their respective primary antibodies (α-

EB1 - 1:5000, α-Klp3A - 1:500, α-Tubulin – 1: 10,000) for 1 hour, followed by 3 times wash and

secondary antibody incubation at 1:5000 dilution for 1 hour. Following secondary antibody

incubation, membrane was washed 3 times in Tris-buffered saline with 0.1% Tween (TBST) and

imaged with a G:BOX system controlled by GeneSnap software (Syngene, Cambridge, U.K.).

Images were further quantified to estimate the knock down efficiency with the help of Fiji/Image

J software (Schindelin et al., 2012). To obtain the intensity values, identical regions were drawn

over all the bands and their integrated intensity was recorded. Intensity values were normalized

to their respective loading controls (Tubulin) to estimate the knockdown efficiency.

Fluorescence and Live-Cell Total internal reflection fluorescence (TIRF) microscopy

Around 500 to 700 µl of cells, expressing the gene of interest were seeded on a 35mm glass

bottom dish with 20mm bottom well (Cellvis, CA, USA) coated with Concanavalin A for 30-60

minutes. Before imaging 2 ml of fresh S2 media containing FBS was added to the dish. Live-cell

TIRF movies were acquired on a Nikon Ti-E microscope equipped with a 100X 1.4 NA DIC Apo

Oil immersion objective, a Hamamatsu ORCA-Flash 4.0 LT digital CMOS camera (C11440), 4

laser lines (447, 488, 561, 641), and a MetaMorph software. Live-cell TIRF movies of all the

constructs (MKLP1-EGFP and Tag-RFP-α-tubulin, RacGAP50C-EGFP and Tag-RFP-α-tubulin,

Polo Kinase-EGFP, ABK-GFP and Tubulin mCherry, EGFP-Rhotekin and Tag-RFP-α-tubulin,

RFP-Rhotekin and *Dm* MKLP1-EGFP, RFP-Rhotekin and MRLC-EGFP, *Dm* ECT2-EGFP and

Tag-RFP-α-tubulin) were acquired at the following settings: 561 exposure time 200ms, 488

exposure time 200ms, temperature 25 °C, and frame acquisition rate 5 Sec. To observe

cytokinesis defects, overnight imaging was performed on control and EB1 depleted cells. Cells

were seeded on a Concanavalin A coated 35mm glass bottom dish as described above. A total

of 76 time-points were acquired for 50 stage positions at the interval of 4 minutes @ 25 °C on a

Nikon Eclipse Ti-E microscope (Nikon, Tokyo, Japan) equipped with 40X/1.30 Oil Nikon Plan

Fluor DIC N2 objective (Nikon), Andor iXon$_3$ EMCCD camera (Andor Technology, Belfast, U.K.),

and Metamorph software (Molecular Devices, Sunnyvale, CA). Images were further analyzed

with the help of MetaMorph software.

Chemical perturbation experiments

Cells expressing *Dm* MKLP1-EGFP; Tag-RFP-α-Tubulin, Polo-EGFP, Rhotekin-EGFP; Tag-

RFP-α-Tubulin, Rhotekin-RFP; *Dm* MKLP1-EGFP, *Dm* ECT2-EGFP; Tag-RFP-α-Tubulin were

subjected to ABK or Polo inhibition. Cells were seeded on a Concanavalin A coated 35mm

glass bottom dish with 20mm bottom well (Cellvis, CA, USA) for 30-60 minutes. Before imaging

2 ml of fresh S2 media containing FBS was added to the dish. Live-cell TIRF movies were

acquired on a Nikon Ti-E microscope as described above. The whole field of view under the

microscope was scanned to find mitotic cells. Mitotic cells expressing the gene of interest were

imaged at the interval of 20-30 seconds until the anaphase onset. After anaphase onset, when

the CS-TIPs were visible, S2 media in the glass bottom dish was exchanged with S2 media

containing Binucleine 2 (40µM) for ABK inhibition or BI 2536 (1µM) for Polo inhibition.

Thereafter, imaging was continued at 5 second intervals. Images were further analyzed with the

help of Metamorph software (CA, USA).

GFP/RFP intensity quantitation

GFP/RFP intensities before and after the addition of inhibitor (Binucleine 2 or BI 2536) were

quantified to estimate the effects of inhibitor on CS-TIPs patterning and signaling. Identical

regions were drawn over the GFP/RFP puncta to measure the fluorescence intensity. The local

background was estimated by placing the same region to a nearby place. Background intensity

values were subtracted from GFP/RFP values to estimate the actual GFP/RFP fluorescence

intensity. Histograms and dot plots were generated using Prism software (Graphpad, CA, USA),

and figures were assembled using Adobe Illustrator software (Adobe, CA, USA).

Table 1. Primers for cloning

MKLP1 FW	TTCAATTGCAAATGGTACCTACTAGTATGAAGGCAGTACCCAGGAC
MKLP1 RV	TCCTCGCCCTTGCTCACCATTCTAGAGATTTTCGACTTCTTGCTGC
RacGAP50C FW	GTTTCAATTGCAAATGGTACCTACTAGTATGGCGCTCTCCGCATTGGC
RacGAP50C RV	CTCCTCGCCCTTGCTCACCATTCTAGATTTCTTGTGCGCAGATGCCG
ΔC-RacGAP50 RV	CTCCTCGCCCTTGCTCACCATTCTAGAGTTATCGATCGGCATTAGCAC
ΔPTIV-RacGAP50C FW	GGCTACTCCACACCAGATCC
ΔPTIV-RacGAP50C RV	GCCAAAGATGAGCGAGATGTTATCG
Polo Kinase FW	GGGGTACCGAAAAGCTGTATCGAGTCGC

Polo Kinase RV	GGACTAGTTGTGAACATCTTCTCCAGCA
Tag RFP Rhotekin FW	GGCATGGACGAGCTGTACAAGACTAGTTCCGGACTCAGATCTCGAG
Tag RFP Rhotekin RV	GGGATAGGCTTACCTTCGAACCGCGGTTAGCCTGTCTTCTCCAGCACCTG
EGFP Rhotekin FW	TTCAAGTTTCAATTGCAAATGGTACCACCATGGTGAGCAAGGGCGA
EGFP Rhotekin RV	TAGGCTTACCTTCGAACCGCGGGCCCGCCTGTCTTCTCCAGCACCT
MRLC FW	TTCAATTGCAAATGGTACCTACTAGTATGTCATCCCGTAAGACCGC
MRLC RV	AGCTCTTCGCCCTTAGACACTCTAGACTGCTCATCCTTGTCCTTGG

2 **Table 2. Primers for RNAi**

Subito FW	TAATACGACTCACTATAGGGCTGAAGCTAATCAATGGCAGC
Subito RV	TAATACGACTCACTATAGGGTTTTCTGAACTGTACTGGCCG
EB1 FW	GAATTAATACGACTCACTATAGGGAGAATGGCTGTAAACGTCTACTCCACAAATGTG
EB1 RV	GAATTAATACGACTCACTATAGGGAGATGCCCGTGCTGTTGGCACAGGCGTTTA

ACKNOWLEDGMENTS

Authors would like to acknowledge Patricia Wadsworth, UMass and Alex Mogilner, NYU. We

are also grateful to Eric Griffis, University of Dundee, for the kind gift of ABK cell line; and Bill

Bement, UW-Madison for the kind gift of Rhotekin plasmid via Patricia Wadsworth. This work

was supported by an NIH grant (5 R01 GM107026) to TJM.

AUTHOR CONTRIBUTIONS

T.J.M conceived the project. T.J.M and V.V designed and performed the experiments, analyzed

the data, and wrote the paper.

REFERENCES

Afonso, O., I. Matos, A.J. Pereira, P. Aguiar, M.A. Lampson, and H. Maiato. 2014. Feedback
control of chromosome separation by a midzone Aurora B gradient. *Science*. 345:332-
336.
Akhmanova, A., and M.O. Steinmetz. 2008. Tracking the ends: a dynamic protein network
controls the fate of microtubule tips. *Nat Rev Mol Cell Biol*. 9:309-322.
Alsop, G.B., and D. Zhang. 2003. Microtubules are the only structural constituent of the spindle
apparatus required for induction of cell cleavage. *J Cell Biol*. 162:383-390.
Basant, A., S. Lekomtsev, Y.C. Tse, D. Zhang, K.M. Longhini, M. Petronczki, and M. Glotzer.
2015. Aurora B kinase promotes cytokinesis by inducing centralspindlin oligomers that
associate with the plasma membrane. *Dev Cell*. 33:204-215.
Bement, W.M. 2000. Rho GTPase activity zones and transient contractile arrays. *BioEssays*.
28:983-993.
Bement, W.M., H.A. Benink, and G. von Dassow. 2005. A microtubule-dependent zone of active
RhoA during cleavage plane specification. *J. Cell Biol.* 170:91-101.
Bement, W.M., M. Leda, A.M. Moe, A.M. Kita, M.E. Larson, A.E. Golding, C. Pfeuti, K.C. Su,
A.L. Miller, A.B. Goryachev, and G. von Dassow. 2015. Activator-inhibitor coupling
between Rho signalling and actin assembly makes the cell cortex an excitable medium.
Nat Cell Biol. 17:1471-1483.
Benink, H.A., and W.M. Bement. 2005. Concentric zones of active RhoA and Cdc42 around
single cell wounds. *J Cell Biol*. 168:429-439.
Biasini, M., S. Bienert, A. Waterhouse, K. Arnold, G. Studer, T. Schmidt, F. Kiefer, T. Gallo
Cassarino, M. Bertoni, L. Bordoli, and T. Schwede. 2014. SWISS-MODEL: modelling
protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids
Res*. 42:W252-258.
Bieling, P., I.A. Telley, and T. Surrey. 2010. A minimal midzone protein module controls
formation and length of antiparallel microtubule overlaps. *Cell*. 142:420-432.
Brennan, I.M., U. Peters, T.M. Kapoor, and A.F. Straight. 2007. Polo-like kinase controls
vertebrate spindle elongation and cytokinesis. *PLOS ONE*. 2:e409.
Breznau, E.B., M. Murt, T.L. Blasius, K.J. Verhey, and A.L. Miller. 2017. The MgcRacGAP SxIP
motif tethers Centralspindlin to microtubule plus ends in Xenopus laevis. *J Cell Sci*.
130:1809-1821.
Bringmann, H., and A.A. Hyman. 2005. A cytokinesis furrow is positioned by two consecutive
signals. *Nature*. 436:731-734.
Burkard, M.E., J. Maciejowski, V. Rodriguez-Bravo, M. Repka, D.M. Lowery, K.R. Clauser, C.
Zhang, K.M. Shokat, S.A. Carr, M.B. Yaffe, and P.V. Jallepalli. 2009. Plk1 self-
organization and priming phosphorylation of HsCYK-4 at the spindle midzone regulate
the onset of division in human cells. *PLoS Biol*. 7:e1000111.
Canman, J.C., L.A. Cameron, P.S. Maddox, A. Straight, J.S. Tirnauer, T.J. Mitchison, G. Fang,
T.M. Kapoor, and E.D. Salmon. 2003. Determining the position of the cell division plane.
Nature. 424:1074-1078.
Canman, J.C., D.B. Hoffman, and E.D. Salmon. 2000. The role of pre- and post-anaphase
microtubules in the cytokinesis phase of the cell cycle. *Curr Biol*. 10:611-614.
Canman, J.C., L. Lewellyn, K. Laband, S.J. Smerdon, A. Desai, B. Bowerman, and K. Oegema.
2008. Inhibition of Rac by the GAP activity of centralspindlin is essential for cytokinesis.
Science. 322:1543-1546.
Cao, L.G., and Y.L. Wang. 1996. Signals from the spindle midzone are required for the
stimulation of cytokinesis in cultured epithelial cells. *Mol Biol Cell*. 7:225-232.

Carmena, M., M.O. Lombardia, H. Ogawa, and W.C. Earnshaw. 2014. Polo kinase regulates the localization and activity of the chromosomal passenger complex in meiosis and mitosis in Drosophila melanogaster. *Open Biol*. 4:140162.

Carmena, M., M.G. Riparbelli, G. Minestrini, A.M. Tavares, R. Adams, G. Callaini, and D.M. Glover. 1998. Drosophila polo kinase is required for cytokinesis. *J Cell Biol*. 143:659-671.

Carmena, M., M. Wheelock, H. Funabiki, and W.C. Earnshaw. 2012. The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nat Rev Mol Cell Biol*. 13:789-803.

Cesario, J.M., J.K. Jang, B. Redding, N. Shah, T. Rahman, and K.S. McKim. 2006. Kinesin 6 family member Subito participates in mitotic spindle assembly and interacts with mitotic regulators. *J Cell Sci*. 119:4770-4780.

Chmatal, L., K. Yang, R.M. Schultz, and M.A. Lampson. 2015. Spatial Regulation of Kinetochore Microtubule Attachments by Destabilization at Spindle Poles in Meiosis I. *Curr Biol*. 25:1835-1841.

Cooke, C.A., M.M. Heck, and W.C. Earnshaw. 1987. The inner centromere protein (INCENP) antigens: movement from inner centromere to midbody during mitosis. *J Cell Biol*. 105:2053-2067.

Crest, J., K. Concha-Moore, and W. Sullivan. 2012. RhoGEF and positioning of rappaport-like furrows in the early Drosophila embryo. *Curr Biol*. 22:2037-2041.

D'Avino, P.P., V. Archambault, M.R. Przewloka, W. Zhang, K.S. Lilley, E. Laue, and D.M. Glover. 2007. Recruitment of Polo kinase to the spindle midzone during cytokinesis requires the Feo/Klp3A complex. *PLoS One*. 2:e572.

D'Avino, P.P., M.S. Savoian, and D.M. Glover. 2004. Mutations in sticky lead to defective organization of the contractile ring during cytokinesis and are enhanced by Rho and suppressed by Rac. *J Cell Biol*. 166:61-71.

D'Avino, P.P., M.S. Savoian, and D.M. Glover. 2005. Cleavage furrow formation and ingression during animal cytokinesis: a microtubule legacy. *J Cell Sci*. 118:1549-1558.

Devore, J.J., G.W. Conrad, and R. Rappaport. 1989. A model for astral stimulation of cytokinesis in animal cells. *J Cell Biol*. 109:2225-2232.

Douglas, M.E., T. Davies, N. Joseph, and M. Mishima. 2010. Aurora B and 14-3-3 coordinately regulate clustering of centralspindlin during cytokinesis. *Curr Biol*. 20:927-933.

Duellberg, C., M. Trokter, R. Jha, I. Sen, M.O. Steinmetz, and T. Surrey. 2014. Reconstitution of a hierarchical +TIP interaction network controlling microtubule end tracking of dynein. *Nat Cell Biol*. 16:804-811.

Ebrahimi, S., H. Fraval, M. Murray, R. Saint, and S.L. Gregory. 2010. Polo kinase interacts with RacGAP50C and is required to localize the cytokinesis initiation complex. *J Biol Chem*. 285:28667-28673.

Eggert, U.S., A.A. Kiger, C. Richter, Z.E. Perlman, N. Perrimon, T.J. Mitchison, and C.M. Field. 2004. Parallel chemical genetic and genome-wide RNAi screens identify cytokinesis inhibitors and targets. *PLoS Biol*. 2:e379.

Ferreira, J.G., A.J. Pereira, A. Akhmanova, and H. Maiato. 2013. Aurora B spatially regulates EB3 phosphorylation to coordinate daughter cell adhesion with cytokinesis. *J Cell Biol*. 201:709-724.

Foe, V.E., and G. von Dassow. 2008. Stable and dynamic microtubules coordinately shape the myosin activation zone during cytokinetic furrow formation. *J Cell Biol*. 183:457-470.

Fujiwara, T., M. Bandi, M. Nitta, E.V. Ivanova, R.T. Bronson, and D. Pellman. 2005. Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null cells. *Nature*. 437:1043-1047.

Fuller, B.G., M.A. Lampson, E.A. Foley, S. Rosasco-Nitcher, K.V. Le, P. Tobelmann, D.L. Brautigan, P.T. Stukenberg, and T.M. Kapoor. 2008. Midzone activation of aurora B in anaphase produces an intracellular phosphorylation gradient. *Nature*. 453:1132-1136.

Glotzer, M. 2004. Cleavage furrow positioning. *J. Cell Biol.* 164:347-351.

Glotzer, M. 2009. The 3Ms of central spindle assembly: microtubules, motors and MAPs. *Nat Rev Mol Cell Biol*. 10:9-20.

Graessl, M., J. Koch, A. Calderon, D. Kamps, S. Banerjee, T. Mazel, N. Schulze, J.K. Jungkurth, R. Patwardhan, D. Solouk, N. Hampe, B. Hoffmann, L. Dehmelt, and P. Nalbant. 2017. An excitable Rho GTPase signaling network generates dynamic subcellular contraction patterns. *J Cell Biol*. 216:4271-4285.

Gruneberg, U., R. Neef, R. Honda, E.A. Nigg, and F.A. Barr. 2004. Relocation of Aurora B from centromeres to the central spindle at the metaphase to anaphase transition requires MKlp2. *J Cell Biol*. 166:167-172.

Helmke, K.J., R. Heald, and J.D. Wilbur. 2013. Interplay between spindle architecture and function. *Int Rev Cell Mol Biol*. 306:83-125.

Honnappa, S., S.M. Gouveia, A. Weisbrich, F.F. Damberger, N.S. Bhavesh, H. Jawhari, I. Grigoriev, F.J. van Rijssel, R.M. Buey, A. Lawera, I. Jelesarov, F.K. Winkler, K. Wuthrich, A. Akhmanova, and M.O. Steinmetz. 2009. An EB1-binding motif acts as a microtubule tip localization signal. *Cell*. 138:366-376.

Hu, C.K., M. Coughlin, C.M. Field, and T.J. Mitchison. 2011. KIF4 regulates midzone length during cytokinesis. *Curr Biol*. 21:815-824.

Hutterer, A., M. Glotzer, and M. Mishima. 2009. Clustering of centralspindlin is essential for its accumulation to the central spindle and the midbody. *Curr Biol*. 19:2043-2049.

Jordan, S.N., and J.C. Canman. 2012. Rho GTPases in animal cell cytokinesis: an occupation by the one percent. *Cytoskeleton (Hoboken)*. 69:919-930.

Kachaner, D., D. Garrido, H. Mehsen, K. Normandin, H. Lavoie, and V. Archambault. 2017. Coupling of Polo kinase activation to nuclear localization by a bifunctional NLS is required during mitotic entry. *Nat Commun*. 8:1701.

Kalab, P., A. Pralle, E.Y. Isacoff, R. Heald, and K. Weis. 2006. Analysis of a RanGTP-regulated gradient in mitotic somatic cells. *Nature*. 440:697-701.

Kalab, P., K. Weis, and R. Heald. 2002. Visualization of a Ran-GTP gradient in interphase and mitotic Xenopus egg extracts. *Science*. 295:2452-2456.

Kelly, A.E., S.C. Sampath, T.A. Maniar, E.M. Woo, B.T. Chait, and H. Funabiki. 2007. Chromosomal enrichment and activation of the aurora B pathway are coupled to spatially regulate spindle assembly. *Dev Cell*. 12:31-43.

Kitagawa, M., S.Y. Fung, N. Onishi, H. Saya, and S.H. Lee. 2013. Targeting Aurora B to the equatorial cortex by MKlp2 is required for cytokinesis. *PLoS One*. 8:e64826.

Kiyomitsu, T., and I.M. Cheeseman. 2013. Cortical dynein and asymmetric membrane elongation coordinately position the spindle in anaphase. *Cell*. 154:391-402.

Kotynkova, K., K.-C. Su, S.C. West, and M. Petronczki. 2016. Plasma membrane association but not midzone recruitment of RhoGEF Ect2 is essential for cytokinesis. *Cell reports*. 17:2672-2686.

Kumar, A., C. Manatschal, A. Rai, I. Grigoriev, M.S. Degen, R. Jaussi, I. Kretzschmar, A.E. Prota, R. Volkmer, R.A. Kammerer, A. Akhmanova, and M.O. Steinmetz. 2017. Short Linear Sequence Motif LxxPTPh Targets Diverse Proteins to Growing Microtubule Ends. *Structure*. 25:924-932 e924.

Kunda, P., N.T. Rodrigues, E. Moeendarbary, T. Liu, A. Ivetic, G. Charras, and B. Baum. 2012. PP1-mediated moesin dephosphorylation couples polar relaxation to mitotic exit. *Curr Biol*. 22:231-236.

Landino, J., S.R. Norris, M. Li, E.R. Ballister, M.A. Lampson, and R. Ohi. 2017. Two mechanisms coordinate the recruitment of the chromosomal passenger complex to the plane of cell division. *Mol Biol Cell*. 28:3634-3646.

Landino, J., and R. Ohi. 2016. The timing of midzone stabilization during cytokinesis depends on myosin II activity and an interaction between INCENP and actin. *Curr Biol*. 26:698-706.

Lenart, P., M. Petronczki, M. Steegmaier, B. Di Fiore, J.J. Lipp, M. Hoffmann, W.J. Rettig, N. Kraut, and J.M. Peters. 2007. The small-molecule inhibitor BI 2536 reveals novel insights into mitotic roles of polo-like kinase 1. *Curr Biol*. 17:304-315.

Li, R. 2007. Cytokinesis in development and disease: variations on a common theme. *Cell. Mol. Life Sci*. 64:3044-3058.

Llamazares, S., A. Moreira, A. Tavares, C. Girdham, B.A. Spruce, C. Gonzalez, R.E. Karess, D.M. Glover, and C.E. Sunkel. 1991. polo encodes a protein kinase homolog required for mitosis in Drosophila. *Genes Dev*. 5:2153-2165.

Manatschal, C., A.M. Farcas, M.S. Degen, M. Bayer, A. Kumar, C. Landgraf, R. Volkmer, Y. Barral, and M.O. Steinmetz. 2016. Molecular basis of Kar9-Bim1 complex function during mating and spindle positioning. *Mol Biol Cell*.

Mangal, S., J. Sacher, T. Kim, D.S. Osorio, F. Motegi, A.X. Carvalho, K. Oegema, and E. Zanin. 2018. TPXL-1 activates Aurora A to clear contractile ring components from the polar cortex during cytokinesis. *J Cell Biol*. 217:837-848.

Matuliene, J., and R. Kuriyama. 2002. Kinesin-like protein CHO1 is required for the formation of midbody matrix and the completion of cytokinesis in mammalian cells. *Mol Biol Cell*. 13:1832-1845.

McKinley, K.L., and I.M. Cheeseman. 2017. Large-Scale Analysis of CRISPR/Cas9 Cell-Cycle Knockouts Reveals the Diversity of p53-Dependent Responses to Cell-Cycle Defects. *Dev Cell*. 40:405-420 e402.

Minestrini, G., A.S. Harley, and D.M. Glover. 2003. Localization of Pavarotti-KLP in living Drosophila embryos suggests roles in reorganizing the cortical cytoskeleton during the mitotic cycle. *Mol Biol Cell*. 14:4028-4038.

Minoshima, Y., T. Kawashima, K. Hirose, Y. Tonozuka, A. Kawajiri, Y.C. Bao, X. Deng, M. Tatsuka, S. Narumiya, W.S. May, Jr., T. Nosaka, K. Semba, T. Inoue, T. Satoh, M. Inagaki, and T. Kitamura. 2003. Phosphorylation by aurora B converts MgcRacGAP to a RhoGAP during cytokinesis. *Dev Cell*. 4:549-560.

Mishima, M., S. Kaitna, and M. Glotzer. 2002. Central spindle assembly and cytokinesis require a kinesin-like protein/RhoGAP complex with microtubule bundling activity. *Devel. Cell*. 2:41-54.

Mishima, M., V. Pavicic, U. Gruneberg, E.A. Nigg, and M. Glotzer. 2004. Cell cycle regulation of central spindle assembly. *Nature*. 430:908-913.

Mitchison, T., M. Wuhr, P. Nguyen, K. Ishihara, A. Groen, and C.M. Field. 2012. Growth, interaction, and positioning of microtubule asters in extremely large vertebrate embryo cells. *Cytoskeleton (Hoboken)*. 69:738-750.

Mogilner, A., and A. Manhart. 2016. Agent-based modeling: case study in cleavage furrow models. *Mol Biol Cell*. 27:3379-3384.

Murthy, K., and P. Wadsworth. 2008. Dual role for microtubules in regulating cortical contractility during cytokinesis. *J Cell Sci*. 121:2350-2359.

Neef, R., U. Gruneberg, R. Kopajtich, X. Li, E.A. Nigg, H. Sillje, and F.A. Barr. 2007. Choice of Plk1 docking partners during mitosis and cytokinesis is controlled by the activation state of Cdk1. *Nat Cell Biol*. 9:436-444.

Nguyen, P.A., A.C. Groen, M. Loose, K. Ishihara, M. Wuhr, C.M. Field, and T.J. Mitchison. 2014. Spatial organization of cytokinesis signaling reconstituted in a cell-free system. *Science*. 346:244-247.

Nishimura, Y., and S. Yonemura. 2006. Centralspindlin regulates ECT2 and RhoA accumulation
at the equatorial cortex during cytokinesis. *J Cell Sci.* 119:104-114.
Odell, G.M., and V.E. Foe. 2008. An agent based model contrasts opposite effects of dynamic
and stable microtubules on cleavage furrow positioning. *J. Cell Biol.* 183:471-483.
Pavicic-Kaltenbrunner, V., M. Mishima, and M. Glotzer. 2007. Cooperative assembly of CYK-
4/MgcRacGAP and ZEN-4/MKLP1 to form the centralspindlin complex. *Mol Biol Cell.*
18:4992-5003.
Petronczki, M., M. Glotzer, N. Kraut, and J.-M. Peters. 2007. Polo-like kinase1 triggers the
initiation of cytokinesis in human cells by promoting recruitment of the RhoGEF Ect2 to
the central spindle. *Devel. Cell.* 12:713-725.
Petronczki, M., P. Lenart, and J.M. Peters. 2008. Polo on the Rise-from Mitotic Entry to
Cytokinesis with Plk1. *Dev Cell.* 14:646-659.
Rappaport, R. 1961. Experiments concerning the cleavage stimulus in sand dollar eggs. *J Exp*
Zool. 148:81-89.
Rappaport, R. 1996. Cytokinesis in animal cells. Cambridge University Press, Cambridge, U.K.
386 pp.
Rodrigues, N.T., S. Lekomtsev, S. Jananji, J. Kriston-Vizi, G.R. Hickson, and B. Baum. 2015.
Kinetochore-localized PP1-Sds22 couples chromosome segregation to polar relaxation.
Nature. 524:489-492.
Rogers, S.L., G.C. Rogers, D.J. Sharp, and R.D. Vale. 2002. Drosophila EB1 is important for
proper assembly, dynamics, and positioning of the mitotic spindle. *J Cell Biol.* 158:873-
884.
Sampath, S.C., R. Ohi, O. Leismann, A. Salic, A. Pozniakovski, and H. Funabiki. 2004. The
chromosomal passenger complex is required for chromatin-induced microtubule
stabilization and spindle assembly. *Cell.* 118:187-202.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C.
Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image
analysis. *Nat Methods.* 9:676-682.
Smurnyy, Y., A.V. Toms, G.R. Hickson, M.J. Eck, and U.S. Eggert. 2010. Binucleine 2, an
isoform-specific inhibitor of Drosophila Aurora B kinase, provides insights into the
mechanism of cytokinesis. *ACS Chem Biol.* 5:1015-1020.
Soderholm, J.F., S.L. Bird, P. Kalab, Y. Sampathkumar, K. Hasegawa, M. Uehara-Bingen, K.
Weis, and R. Heald. 2011. Importazole, a small molecule inhibitor of the transport
receptor importin-beta. *ACS Chem Biol.* 6:700-708.
Somers, G.W., and R. Saint. 2003. A RhoGEF and Rho family GTPase-activating protein
complex links the contractile ring to cortical microtubules at the onset of cytokinesis.
Devel. Cell. 4:29-39.
Stangier, M.M., A. Kumar, X. Chen, A.M. Farcas, Y. Barral, and M.O. Steinmetz. 2018.
Structure-Function Relationship of the Bik1-Bim1 Complex. *Structure.* 26:607-618 e604.
Strickland, L.I., E.J. Donnelly, and D.R. Burgess. 2005a. Induction of cytokinesis is independent
of precisely regulated microtubule dynamics. *Mol Biol Cell.* 16:4485-4494.
Strickland, L.I., Y. Wen, G.G. Gundersen, and D.R. Burgess. 2005b. Interaction between EB1
and p150glued is required for anaphase astral microtubule elongation and stimulation of
cytokinesis. *Curr Biol.* 15:2249-2255.
Subramanian, R., S.C. Ti, L. Tan, S.A. Darst, and T.M. Kapoor. 2013. Marking and measuring
single microtubules by PRC1 and kinesin-4. *Cell.* 154:377-390.
Tan, L., and T.M. Kapoor. 2011. Examining the dynamics of chromosomal passenger complex
(CPC)-dependent phosphorylation during cell division. *Proc Natl Acad Sci U S A.*
108:16675-16680.

Tyson, J.J., K.C. Chen, and B. Novak. 2003. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory and signaling pathways in the cell. *Curr Opin Cell Biol*. 15:221-231.

Vale, R.D., J.A. Spudich, and E.R. Griffis. 2009. Dynamics of myosin, microtubules, and Kinesin-6 at the cortex during cytokinesis in Drosophila S2 cells. *J Cell Biol*. 186:727-738.

von Dassow, G. 2009. Concurrent cues for cytokinetic furrow induction in animal cells. *Trends in Cell Bio*. 19:165-173.

Wu, Z., Q. Jiang, P.R. Clarke, and C. Zhang. 2013. Phosphorylation of Crm1 by CDK1-cyclin-B promotes Ran-dependent mitotic spindle assembly. *J Cell Sci*. 126:3417-3428.

Yang, C., J. Wu, C. de Heus, I. Grigoriev, N. Liv, Y. Yao, I. Smal, E. Meijering, J. Klumperman, R.Z. Qi, and A. Akhmanova. 2017. EB1 and EB3 regulate microtubule minus end organization and Golgi morphology. *J Cell Biol*. 216:3179-3198.

Ye, A.A., J. Deretic, C.M. Hoel, A.W. Hinman, D. Cimini, J.P. Welburn, and T.J. Maresca. 2015. Aurora A Kinase Contributes to a Pole-Based Error Correction Pathway. *Curr Biol*. 25:1842-1851.

Yuce, O., A. Piekny, and M. Glotzer. 2005. An ECT2-centralspindlin complex regulates the localization and function of RhoA. *J. Cell Biol.* 170:571-582.

Zhu, C., and W. Jiang. 2005. Cell cycle-dependent translocation of PRC1 on the spindle by Kif4 is essential for midzone formation and cytokinesis. *Proc Natl Acad Sci U S A*. 102:343-348.

FIGURE LEGENDS

Figure 1. *Dm* MKLP1, RacGAP50C, ABK, and polo kinase localize to CS-TIPs and the

midzone. (A) Selected still frames from the live-cell TIRF movies showing *Dm* MKLP1-EGFP

(green) localization to the plus-tips of polar and equatorial MTs (red) within 2.30 minutes of

anaphase onset. *Dm*-MKLP1-EGFP gets patterned on the equatorial MTs after ~10 minutes (the

last panel). (B) Selected still frames from the live-cell TIRF movies showing RacGAP50C-EGFP

(green) localization to the plus-tips of polar and equatorial MTs (red) within 1.33 minutes of

anaphase onset. The last panel shows patterning of RacGAP50C-EGFP mainly on the

equatorial MTs. (C) ABK (red) localizes to the plus-tips of polar and equatorial MTs (green)

within 30 seconds of anaphase onset. The last panel shows ABK localization to the equatorial

and midzone MTs. (D) Polo localization on the MTs (red regions) within minutes of anaphase

onset and then patterning on the equatorial MT tips. Outline of the cell cortex at anaphase onset

with the position of CS-TIPs that appeared over the time-lapse marked with red circles (bottom

panel). The membrane invaginates where CS-TIPs contact the cortex as evident in the overlays

of the cell peripheries at later time-points (blue – 11:00; and orange – 20:00) relative to the cell

boundary at anaphase onset (green). Zero time-points indicate anaphase onset, the next time-

points indicate decoration of proteins on the polar and equatorial MTs (pre-patterning), and the

last time-points indicate localization of cytokinesis proteins mainly on the equatorial MTs

(patterning) in all the figures. Time: mins:secs. Scale bars, 10 µm.

Figure 2. CS-TIPs, upon cortical contact recruit *Dm* ECT2 (Pebble), which then leads to

RhoA activation and myosin accumulation. (A) Representative TIRF micrographs showing

Dm ECT2 (green) localization to the CS-TIPs (red). *Dm* ECT2 localization to the CS-TIPs (time-

point 00.00) is followed by its auto-amplification (later time-points) near the equatorial cortex.

Red arrow heads indicate *Dm* ECT2 localization to the CS-TIPs and its amplification. (B) Dot

plots showing first appearance of *Dm* ECT2 (10.48 ± 1.03), n=21 and Rhotekin (14.25 ± 1.68),

n=16 after MTs contacting the cortex. Error bars: SEM. (C) Dot plots showing average peak

accumulation timings of *Dm* ECT2 (1.5 ± 0.14), n=18; RhoA (2.5 ± 0.2), n=18; and Myosin (5.5 ±

0.54), n=19, Error Bars: SEM. (D) The active RhoA reporter Rhotekin is enriched near CS-TIPs

within minutes (3:58) of anaphase onset (AO) and tip-signaling can precede detectable midzone

RhoA activation (9:44). Red ovals denote polar signaling TIPs prior to patterning onto equatorial

MTs. (E) Polo-positive CS-TIPs activate cortical RhoA (Rhotekin) upon contact during C-Phase.

(F) Polo-positive CS-TIPs trigger localized cortical contractility as evidenced by MRLC

accumulation. Time: mins:secs. Color wedge displays pixel values from 150-1200. Scale bars,

10 µm (B, C, and D), 5µm (A).

Figure 3. CS-TIPs require ABK activity, but not Polo kinase activity or midzone-localized

CPC. (A, B) Polo and *Dm* MKLP1 are rapidly lost from CS-TIPs following wash-in of an ABK

inhibitor, Binucleine 2. Zero time-points indicate anaphase onset, 04:35 and 05:20 time-points

indicate the time of Binucleine 2 addition. (C, D) Polo and *Dm* MKLP1 remain localized to the

CS-TIPs following wash-in of a Polo kinase inhibitor, BI 2536. Zero time-points indicate

anaphase onset, 03:53 and 03:01 time-points indicate the time of BI 2536 addition. (E) Control

cells show normal localization of polo-EGFP on the CS-TIPs and midzone (left). Depletion of

Dm Kinesin-4 (Klp3A) results in loss of polo kinase from the midzone, but not from the CS-TIPs

(Right). The first image is a selected frame from a wide-field movie, while last two images are

selected frames from a live-cell TIRF movie for both control and Klp3A depletion. (F) Depletion

of *Dm* MKLP2 leads to loss of active phosphorylated ABK (pABK) at the midzone (left panel),

yet *Dm* MKLP1 assembles on CS-TIPs normally (right panel). Time: mins:secs. Scale bars, 5

µm (C,D, and E), 10 µm (A, B, and F).

Figure 4. ABK and polo kinase activities are required for RhoA activation. (A, C) Active

RhoA (Rhotekin) is lost from CS-TIPs following wash-in of an ABK inhibitor, Binucleine 2. (B, D)

Active RhoA (Rhotekin) is lost from CS-TIPs following wash-in of Polo kinase inhibitor (BI 2536),

but *Dm* MKLP1 remain on the CS-TIPs. (E) Bar graphs show normalized Rhotekin intensity

before and after the addition of ABK specific inhibitor, Binucleine 2 or Polo inhibitor, BI 2536.

Control cells were not treated with either Binucleine 2 or BI 2536. For quantitation of Rhotekin

intensity after the addition of Binucleine 2 or BI 2536, a representative frame after ~3-4 minutes

was chosen. Both cortical and equatorial regions were quantified. n=4, Error bars: SD. (F) *Dm*

ECT2 is lost from CS-TIPs following wash-in of Polo kinase inhibitor (BI 2536). Zero time-points

indicate the time of Binucleine 2 or BI 2536 addition to the imaging chamber in all the figures.

Red ovals indicate active RhoA (Rhotekin) accumulation near the CS-TIPs. Red circles indicate

active *Dm* ECT2 accumulation near the CS-TIPs. Time: mins:secs. Scale bars, 5 μm.

Figure 5. Satellite MT arrays provide a signaling platform near the cortex and midzone.

(A) Representative frames from a TIRF time-lapse movie of *Dm* MKLP1-EGFP (Supplementary

Video 1), (top panel). Inset shows enlarged midzone region (bottom panel). Two CS-TIP

bundling events (labeled 1 and 2) occurring at the equatorial cortex. The CS-TIPs start at

oblique angles, become organized into antiparallel bundles with localized *Dm* MKLP1, and are

pushed inward. (B) Representative frames from a TIRF time-lapse movie of Tag-RFP-α-tubulin

and EGFP-Rhotekin (left). Inset 1 and 2 show enlarged satellite MT arrays where Rhotekin gets

enriched (right). Time: mins:secs. Scale bar, 5 μm.

Figure 6. A ''PTIV'' motif in RacGAP50C is required for localization to MT plus-tips and

efficient cytokinesis. (A) TIRF micrographs showing loss of CS-TIPs assembly in cells

depleted of EB1 (bottom panel); control cells show normal CS-TIPs localization (top panel). (B)

Quantitation of CS-TIPs assembly in control and EB1 depleted cells (observed in both Polo and

Dm MKLP1-EGFP-expressing cells; 2 experiments each, n = 38 and 41 for control and EB1

depleted cells respectively, Error bars: SD). (C) Quantitation of cells that fail to initiate furrowing

in control and EB1 RNAi cells. n = 120 and 110 for control and EB1 RNAi cells respectively,

Error bars = SD. (D) Quantitation of binucleated cells in control and EB1 RNAi, n = 410 and 452

for control and EB1 RNAi cells respectively, Error bars = SD. (E) The putative hxxPTxh motif

(FGPTIV) in RacGAP50C is highly conserved although key residues in the motif are different in

C. elegans. In the red "PT" consensus is highlighted, "h" is a hydrophobic amino acid. (F, G)

Deletion of C-terminus or ''PTIV motif'' in RacGAP50C result in loss of CS-TIPs assembly, but

not midzone or cortical localization. (H) Quantitation of CS-TIPs assembly in cells expressing

WT (n=21), ΔPTIV (n=26), or ΔC-RacGAP50-EGFP (n=20), 3 experiments each, Error bars: SD.

(I) Bar charts showing fold increase in binucleated cells in cells expressing ΔPTIV-RagGAP50C-

EGFP or WT- RacGAP50C-EGFP, n ≥ 250 for each condition, Error Bars: SD.

Figure 7. Model for the establishment of the cleavage furrow by CS-TIPs during

cytokinesis. (A) Phosphorylation of MKLP1 by CDK1 during metaphase impedes its

association with the spindle MTs. (B) As cell progresses towards anaphase, CDK1 activity

drops, resulting in loss of MKLP1 phosphorylation and binding of MKLP1 to the spindle MTs.

Within 2-3 minutes of anaphase onset, centralspindlin complex, ABK, and Polo kinase decorate

the CS-TIPs. CS-TIPs after making a cortical contact, recruits Rho-GEF, ECT2, thereby

activating RhoA in the vicinity. (C) CS-TIPs disassemble from the polar MT tips, and

preferentially pattern on the equatorial MT tips, resulting in local activation RhoA near the

equatorial cortex and assembly of actomyosin ring. (D) CS-TIPs assembly is achieved by the

interaction between centralspindlin component, RacGAP50C and +TIP end-binding protein 1,

(EB1). ECT2 recruitment to the CS-TIPs and its auto-amplification (shown by curved arrow and

question mark) most likely involves a mechanism that requires positive feedback.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. (A) Still-images from live-cell TIRF microscopy showing that *Dm*

Kinesin-4 (Klp3A) depletion doesn't affect CS-TIPs or midzone localization of *Dm* MKLP1 (red).

(B, D) Western blots showing depletion of *Dm* Kinesin-4 (Klp3A) and EB1 by RNAi in cells

expressing Polo-EGFP and MKLP1-EGFP respectively. (C) Quantitation of pABK intensity in

control and *Dm* MKLP2 depleted cells, n=25, Error bar: SEM. (E) Kymograph of MKLP1 tip

tracking (left) and microtubule polymerization (middle) generated from time-lapse TIRF

microscopy (Supplementary Movie 1). The last image (right) shows the merge picture of MKLP1

and microtubule, Scale bar, 2μm, time: 1.5 min. (F) Still-images from live-cell TIRF microscopy

showing RacGAP50C-EGFP (green) localization to the CS-TIPs (red) in control RNAi cells. Red

ovals/circles highlight RacGAP50C localization to the CS-TIPs. Time: mins:secs. Scale bar, 5

μm.

Supplementary Figure 2. (A) Homology structure modelling of WT-RacGAP50C and ΔPTIV-

RacGAP50C. Structures were generated using Rac GTPase-activating protein 1 (2ovj. 1. A) as

a template in SWISS-MODEL. Arrow heads indicate the position of hxxPTxh motif (FGPTIV) in

Drosophila RacGAP50C.

SUPPLEMENTARY VIDEOS

Video 1. *Dm* MKLP1 localizes to the CS-TIPs of polar and equatorial MTs. Video shows co-

expression of *Dm* MKLP1-EGFP (green) and Tag-RFP-T-α-tubulin (red) in S2 cells. *Dm* MKLP1-

EGFP decorates the polar and equatorial CS-TIPs within ~ 2-3 minutes of anaphase onset

before becoming patterned onto equatorial MTs (~7-10 minutes). *Dm* MKLP1-EGFP is also

visible on the growing MT plus-tips and the central spindle. Frames were acquired at 10-sec.

intervals on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per

second. Time: mins:secs. Scale bar, 5 μm.

Video 2. **RacGAP50C localizes to the CS-TIPs of polar and equatorial MTs**. Video shows

co-expression of RacGAP50C-EGFP (green) and Tag-RFP-T-α-tubulin (red) in S2 cells.

RacGAP50C-EGFP decorates the CS-TIPs within ~2 minutes of anaphase onset before

becoming patterned onto equatorial MTs. RacGAP50C-EGFP is also visible on the growing MT

plus-tips and the central spindle. Frames were acquired at 5-sec. intervals after anaphase onset

on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per second Time:

mins:secs. Scale bar, 5 μm.

Video 3. Polo kinase localizes to the CS-TIPs and triggers cortical contractility. Video

shows the expression of Polo-EGFP in S2 cells. Polo-EGFP localizes to the CS-TIPs, central

spindle, and interphase MTs. Cortical contractility and membrane invagination is clearly visible

in the regions where CS-TIPs contact the cortex. Frames were acquired at 15-sec. intervals on

a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per second. Time:

mins:secs. Scale bar, 5 μm.

Video 4. *Dm* ECT2 (Pebble) accumulation on the CS-TIPs causes its auto-amplification

near the cortex. Video shows co-expression of *Dm* ECT2-EGFP (green) and Tag-RFP-T-α-

tubulin (red) in *Drosophila* S2 cells. *Dm* ECT2 localization to the CS-TIPs at t = 3.30, and its

amplification between t = 3.30 and 8.30 is clearly visible around the cortex. Accumulation of *Dm*

ECT2-EGFP on midzone MTs and cortex is also visible. Frames were acquired at 5-sec

intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is

20 frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 5. CS-TIPs activate RhoA. Video shows co-expression of EGFP-Rhotekin (green) and

Tag-RFP-T-α-tubulin (red) in *Drosophila* S2 cells. RhoA activation near the CS-TIPs is visible

between t = 02:01 and 5:31. Cortical localization of Rhotekin-EGFP precedes its midzone

localization. The midzone localization of Rhotekin-EGFP is visible between t = 5:31 and 07:31.

Frames were acquired at 5-sec. intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF

microscope. The playback rate is 10 frames per second.Time: mins:secs. Scale bar, 5 μm.

Video 6. Polo-positive CS-TIPs activate RhoA. Video shows co-expression of RFP-Rhotekin

(red) and Polo-EGFP (green) in *Drosophila* S2 cells. In this movie, Polo-EGFP serves as a CS-

TIP marker. RhoA activation near the polo positive CS-TIPs is visible between t= 01:36 and

6:36. Frames were acquired at 5-sec. intervals after anaphase onset on a Nikon Eclipse Ti-E -

TIRF microscope. The playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5

μm.

Video 7. CS-TIPs trigger cortical contractility by myosin accumulation. Video shows co-

expression of myosin light regulatory chain (MRLC)-RFP (red) and Polo-EGFP (green) in

Drosophila S2 cells. Myosin accumulation and cortical contractility near the polo positive CS-

TIPs are visible between t = 02:00 and 13:16. Frames were acquired at 5-sec. intervals on a

Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per second. Time:

mins:secs. Scale bar, 5 μm.

Video 8. ABK is required for CS-TIPs assembly. Video shows the expression of Polo-EGFP

in S2 cells. Addition of ABK inhibitor, Binucleine 2, between t = 03:15 and 04:35 results in

disassembly of CS-TIPs. Frames were acquired at 15-sec. intervals on a Nikon Eclipse Ti-E -

TIRF microscope. The playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5

μm.

Video 9. ABK is required for CS-TIPs assembly. Video shows the expression of *Dm* MKLP1-

EGFP in S2 cells. Addition of ABK inhibitor, Binucleine 2, between t = 04:10 and 6:10 results in

disassembly of CS-TIPs. Frames were acquired at 10-sec. intervals on a Nikon Eclipse Ti-E -

TIRF microscope. The playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5

μm.

Video 10. Polo kinase is not required for CS-TIPs assembly. Video shows the expression of

Polo-EGFP in S2 cells. Addition of Polo inhibitor, BI 2536, between t = 02:45 and 03:53, doesn't

result in CS-TIPs disassembly. Frames were acquired at 15-sec. intervals after anaphase onset

on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per second. Time:

mins:secs. Scale bar, 5 μm.

Video 11. Polo kinase is not required for CS-TIPs assembly. Video shows co-expression of

Dm MKLP1-EGFP (green) and Tag-RFP-T-α-tubulin (red) in S2 cells. Addition of Polo inhibitor,

BI 2536, between t = 01:20 and 02:50, doesn't result in CS-TIPs disassembly. Frames were

acquired at 5-sec. intervals on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10

frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 12. Polo Kinase doesn't localize to the midzone, but assembles normally on the

CS-TIPs in Klp3A depleted cells. Video shows the expression of Polo-EGFP in Klp3A

depleted cells in *Drosophila* S2 cells. Polo-EGFP doesn't localize to the midzone, but

assembles normally on the CS-TIPs. Frames were acquired at 5-sec. intervals on a Nikon

Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per second. Time: mins:secs.

Scale bar, 5 μm.

Video 13. ABK is required for CS-TIP signaling. Video shows co-expression of EGFP-

Rhotekin (green) and Tag-RFP-T-α-tubulin (red) in *Drosophila* S2 cells. RhoA activation near

the CS-TIPs is visible between t = 02:00 to 3:55. Addition of ABK inhibitor, Binucleine 2,

between t = 03:55 and 5:08 results in loss of Rhotekin from the CS-TIPs within ~3 minutes.

However, midzone Rhotekin signals take ~6.30 minutes to disappear. Frames were acquired at

5-sec. intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF microscope. The playback

rate is 10 frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 14. Polo kinase activity is required for CS-TIPs signaling. Video shows co-expression

of EGFP-Rhotekin and Tag-RFP-T-α-tubulin (red) in *Drosophila* S2 cells. RhoA activation near

the CS-TIPs is visible between t = 00:00 to 01:40. Addition of Polo inhibitor, BI 2536, between t

= 01:40 and 2:51 results in loss of Rhotekin from the CS-TIPs within ~3 minutes. Frames were

acquired at 5-sec. intervals on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10

frames per second. Time: mins:secs. Scale bar, 5 µm.

Video 15. ABK is required for CS-TIPs signaling. Video shows the expression of MKLP1-

EGFP (green) and Rhotekin-RFP (red) in *Drosophila* S2 cells. RhoA is visible around the cortex

at t=1.00, but it gets reorganized near the CS-TIPs by t=3:58. Addition of ABK inhibitor,

Binucleine 2, between t = 04:48 and 6:30 results in loss of Rhotekin from the CS-TIPs within ~3

minutes. Frames were acquired at 5-sec. intervals after anaphase onset on a Nikon Eclipse Ti-E

- TIRF microscope. The playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5

µm.

Video 16. Polo kinase activity is required for CS-TIPs signaling but not for assembly.

Video shows the expression of MKLP1-EGFP (green) and RFP-Rhotekin (red) in *Drosophila* S2

cells. RhoA accumulation near the CS-TIPs is visible after t = 01:02. Addition of Polo inhibitor,

BI 2536, between t = 01:27 and 02:55 results in loss of Rhotekin from the CS-TIPs within ~4.30

minutes, however MKLP1 localization to the cortex is still visible. Frames were acquired at 5-

sec. intervals on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is 10 frames per

second. Time: mins:secs. Scale bar, 5 µm.

Video 17. Polo kinase activity is required for ECT2 accumulation near the cortex. Video

shows co-expression of *Dm* ECT2-EGFP (green) and Tag-RFP-T α-tubulin (red) in *Drosophila*

S2 cells. ECT2 localization on the CS-TIPs is visible after t = 04:29. Addition of Polo inhibitor, BI

2536, between t = 04:34 and 05:49 results in loss of ECT2 from the CS-TIPs within ~3 minutes.

Frames were acquired at 5-sec. intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF

microscope. The playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 18. EB1 is required for CS-TIPs assembly. Video shows co-expression of *Dm* MKLP1-

EGFP (green) and Tag-RFP-T-α-tubulin (red) in *Drosophila* S2 cells. In contrast to control RNAi

cells, EB1 RNAi in cells expressing *Dm* MKLP1-EGFP results in loss of CS-TIPs assembly.

Frames were acquired at 5-sec intervals on a Nikon Eclipse Ti-E - TIRF microscope. The

playback rate is 10 frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 19. C-terminal truncation of RacGAP50C results in loss of CS-TIPs assembly. Video

shows co-expression of C-terminal truncation mutant of RacGAP50C-EGFP (green) and Tag-

RFP-T-α-tubulin (red) in *Drosophila* S2 cells. In contrast to full length RacGAP50C, C-terminal

truncation of RacGAP50C results in loss CS-TIPs assembly. Frames were acquired at 5-sec.

intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF microscope. The playback rate is

10 frames per second. Time: mins:secs. Scale bar, 5 μm.

Video 20. Deletion of ''PTIV'' motif in RacGAP50C results in loss of CS-TIPs assembly.

Video shows co-expression of ''PTIV'' deletion mutant of RacGAP50C-EGFP (green) and Tag-

RFP-T-α-tubulin (red) in *Drosophila* S2 cells. In contrast to full length RacGAP50C, ''PTIV''

deletion mutant of RacGAP50C results in loss CS-TIPs assembly. Frames were acquired at 5-

sec. intervals after anaphase onset on a Nikon Eclipse Ti-E - TIRF microscope. The playback

rate is 10 frames per second. Time: mins:secs. Scale bar, 5 μm.

Figure 1

Figure 2

A

Tubulin

Dm ECT2

Overlay

00:00   02:15   03:20   05:30

B

C

D

Tubulin

Rhotekin (Active RhoA)

Overlay

3:58 (post-AO)   9:44 (post-AO)

Pre-Patterning   Post-Patterning

E

Rhotekin   Polo Kinase   Overlay

F

MRLC   Polo Kinase   Merge

Figure 3

Figure 4

Figure 5

Figure 6

A

Ctrl. RNAi

EB1 RNAi

00:10

02:41

Tubulin   *Dm* MKLP1   Merge

B

n=38

n=41

% cells with CS-TIPs

Ctrl. RNAi   EB1 RNAi

C

% Cells that fail to initiate furrow

Ctrl. RNAi   EB1 RNAi

D

% Binucleates

Ctrl. RNAi   EB1 RNAi

E

Dm RacGAP50C   528   SI I FGPTIV GYS
Hs MgcRacGAP   497   AK VFGPTIV AHA
Mm MgcRacGAP   498   AK VFGPTIV AHT
Xt MgcRacGAP   496   SK VFGPTLV GHA
Sp MgcRacGAP   486   AK VFGPTII GYG

Ce CYK-4   572   AR MVAPAVM GHP

hxxPTxh

F

Tubulin

ΔC-RacGAP

Merge

00:00   01:00   02:50

G

Tubulin

ΔPT-RacGAP

Merge

00:00   00:38   02:53

H

% Cells with CS-TIPs

WT   ΔPT   ΔC

RacGAP50C

I

Fold increase in Binucleates

-GFP   +GFP   -GFP   +GFP

Δ PT   WT

RacGAP50C

Figure 7

Supplementary Figure 1

A Klp3A RNAi
Tubulin *Dm* MKLP1 Merge
Pre-patterning
Post-patterning

B RNAi:
Ctrl. Klp3a
Klp3a
Tubulin
Anti-Klp3A and anti-tubulin
Western blot
(97% Depletion)

C
Normalized pABK Intensity
Ctrl. RNAi *Dm* MKLP2 RNAi

D RNAi:
Ctrl. EB1
EB1
Tubulin
Anti-EB1and anti-tubulin
Western blot
(92% Depletion)

E Time
Distance
1.5min
2µm
MKLP1 MT Merge

F Anaphase onset Pre-Patterning Post-Patterning
Tubulin
RacGAP50C
Merge
00:00 01:40 10:10

Supplementary Figure 2

FGPTIV

FGΔPTIV

WT-RacGAP50C

ΔPTIV-RacGAP50C