**Nano-particles constructed from *Chaetomium cochliodes* CTh05 against**

***Magnaporthe oryzae* causing rice blast**

**JiaoJiao Song [a*], Somdej Kanokmedhakul [b], Kwanjai Kanokmedhkul[b], Supattar**

**Poeaim[c] and Kasem Soytong[a]**

[a]Department of Plant Production Technology, Faculty of Agricultural Technology, King

Mongkut's Institute of Technology Ladkrabang, Ladkrabang, Bangkok, Thailand;

[b]Natural Products Research Unit, Department of Chemistry and Centre of Excellence for

Innovation in Chemistry , Faculty of Science, Khon Kaen University, Khon Kaen 40000,

Thailand;

[c]Department of Biology, Faculty of Science, King Mongkut's Institute of Technology

Ladkrabang, Ladkrabang, Bangkok, Thailand.

**Running Head:** Nano-particles from *Chaetomium* against Rice Blast

**\*Corresponding Author:** Jiaojiao Song; **E-mail:** misssongjiaojiao@gmail.com

**Co-Authors:** Somdej Kanokmedhakul:    somdej@kku.ac.th

19                 Kwanjai Kanokmedhkul:    kwanjai@kku.ac.th

20                 Supattar Poeaim:    poeaim@hotmail.com

21                 Kasem Soytong:    ajkasem@gmail.com

**ABSTRACT**   The effective isolate of *Chaetomium cochliodes* CTh05 actively against *Magnaporthe oryzae* isolate PO1 causing rice blast were tested on rice var RD57.   The symptoms of blast on rice var RD57 was isolated and confirmed by morphological and molecular phylogeny which identified as *Magnaporthe oryzae*. Morphology and phylogenic identification was also confirmed *C. cochliodes* CTh05. *Magnporthe oryzae* isolate PO1 proved to be aggressive isolate to cause blast of rice var. Rd57. Bi-culture test proved that *C. cochliodes* CTh05 can be suppressed the growth of *M. oryzae* PO1. The fungal metabolites extracted from *C. cochliodes* CTh05 (CCoE, CCoM and CCoH) expressed antifungal against *M. oryzae* isolate P01 (rice blast) with the $ED_{50}$ of 85, 144 and 374 ppm, respectively. Nano-particles derived from *C. cochliodes* CTh05 extracts (nanoCCoE, nanoCCoM and nanoCCoH) exhibited significantly antifungal activity against *M. oryzae* PO1 wih the $ED_{50}$ values of 9, 16 and 33 ppm, respectively. Further investigation is to formulate the nanoparticles from active compounds of *C. cochliodes* CTh05 for plant immunity and to be applied in the rice fields.

**IMPORTANCE** Rice blast is an economic important disease which caused by *Magnaporthe oryzae*. It causes yield loss and economic damage wherever rice is grown in Asia. It becomes the main disease due to wide distribution and most infection under favorable condition. The farmers have been applied several kinds of chemical fungicides that leading to the pathogen become resistant to those fungicides, and causes environmental pollution which the toxic chemicals residue in soil, water and surrounding environment. The significance of this research is to discover a new agricultural input as nanoparticles from *Chaetomium cochliodes* CTh05 to control the blast pathogen which will develop the nanoproduct to elicit immunity in rice, environmental protection and food safety.

**KEYWORDS:** Nano-particles, *Chaetomium cochliodes* CTh05, rice blast, *Magnaporthe oryzae*

**INTRODUCTION** Rice blast caused by *Magnaporthe oryzae* (Anamorph; *Pyricularia oryzae*). It is one of the first recorded as rice fever disease in China as early as 1637. Rice blast expanded through Asia, Latin America, and Africa and is now found in over 85 countries worldwide (1, 2).   *M. oryzae* infects all parts of plants causing yield losses leading to shortage in many developing countries in recent years (1). Valent (3) stated that disease is epidemic in all continents wherever rice is grown and yield loss due to blast can be as high as 50% (4). The search for new effective disease control is needed to reduce or eliminate the chemical fungicides which causes the environment problem. Nanotechnology in agriculture becomes a new method to build up and restructure the materials at molecular level. Molecular nanotechnology involves constructing organic materials into defined structures, atom by atom or molecule by molecule (5). The application of nanotechnology in agriculture have increased attention in recent years (6). Researchers have actively investigated the synthesis of organic nanomaterials including different kinds of nanoparticle for biological properties (7, 8). The application of nanotechnology in agriculture are being explored (5) and could be helped the solution of overuse toxic chemicals faced by agriculture (9). Nanoparticles contain bioactive substances from natural products that can enable fast effective penetration through plant cuticles and tissues (10). It can provide efficient pest management in agriculture (11) and formulate to contain pesticides in colloidal suspension or powder to apply in agriculture (9), and increase the stability of active compounds (10). Application of natural products from *Chaetomium* spp. has been proven to be antifungal activity against several plant pathogens (12). Tann and Soytong (13) reported that nano particles constructed from *Chaetomium globosum* KMITL-N0805 expressed antifungal activity against *Curvularia lunata*, leaf spot of rice var. Sen Pidoa, and the tested nanoparticles can be controlled leaf spot of rice at 60 days. Furthermore, Tann and Soytong (14) reported nano-products derived from *Chaetomium cupreum* can

be applied to reduce the incidence of rice leaf spot. This research was conducted to evaluate nano-particles
constructed from *Chaetomium cochliodes* CTh05 for biological activity against *Magnaporthe oryzae* causing
rice blast

**RESULTS**

**Isolate of rice blast pathogen and *Chaetomium* sp.** The symptoms of blast was isolated and
morphological identified as *Magnaporthe oryzae* isolate PO1 as seen in Fig. 1. Culture on PDA grown to full
plate (9 cm) in10 days. Mycelia are septated and hyaline, producing conidiophores and 3 cells conidia.   *C.*
*cochliodes* CTh05 from previous experiment (15), it was cultured on PDA for 3 weeks and shows olive-green to
brown, producing perithecia and subglobose asci, one ascus has eight ascospores (Fig. 2).

**FIG1** *Magnaporthe oryzae* PO1, A= pure culture in PDA; B= conidia; C = mycelium and conidia.

**FIG 2** *Chaetomium cochliodes* CTh05, A=Colony, B= Perithecia, C=Asci, D=Ascospores.

Molecular phylogenic identification was confirm species. Phylogenic tree was clearly demonstrated that the
rice blast pathogen as *Magnporthe oryzae* compared to database from Genbank which closedly related to
MF668691, MF669475, and followed by FN555115, FN555121 and KY825257. *Sordaria fimicola* KY930619
was out group (Fig. 3).  Phylogenic confirmation of *Chaetomium cochliodes* CTh05 was clearly shown in
phylogenic tree compared to *Colletotrichum gloeospoiodes* AY714052 as outgroup (Fig. 4). Data from Genbank
can be reliable confirmed CTh05 as *Chaetomium cochliodes* when compared to HQ316565, and followed by
KT895345, JN209868 and JN209864.  *Magnporthe oryzae* isolate PO1 was conformed as the pathogenic
isolate to rice var. Rd57 by pathogenicity test. The inoculated wounds showed the lesions as roundish to
elongated, necrotic, gray spots about 1-2 mm in diameter within 5 days as seen in Fig. 5.

**FIG 3** Phylogenic tree of *Magnporthe oryzae* (PO1)    **FIG 4** Phylogenic tree of *Chaetomium cochliodes* CTh05

**FIG 5** Pathogenicity test on rice leaf

Left = Control

Right   =Treatment

**FIG 6** Bi-culture test between *Magnaporthe oryzae* isolate PO1 and *Chaetomium cochliodes* CTh05, Left = Chaetomium culture, Middle = Bi-culture plates, and Right = Magnaporthe culture

**Bi-culture Test.** *M. oryzae* isolate PO1 was inhibited by *C. cochliodes* CTh05. The colony diameter of the
antagonist averaged of 4.35 cm while the control was 9.00 cm, the colony growth inhibition was 51.66 % in 10
days. Thereafter incubation period was extended to 30 days and the colony inhibition averaged over 90 % (Fig.
6).

**Bioactivity test of fungal metabolites from *Chaetomium cochliodes* CTh05.** Result showed that CCoE
gave significantly highest spore inhibition of 88 %, followed by CCoM and CCoH with spore inhibition of 80.74
and 67.57 %, respectively at 1,000 ppm in 12 days. The fungal metabolites CCoE, CCoM and CCoH showed
antifungal activity against *M. oryzae* isolate PO1 for spore inhibition with the $ED_{50}$ values of 85, 144, 374 ppm
(Table 1). Interestingly, at 15 ppm nanoCCoH, nanoCCoE and nanoCCoM gave significantly inhibited spore
production.   NanoCCoE gave significantly inhibited spore production of 68 % while nanoCCoM and
nanoCCoH at 47 and 34 %, respectively in 12 days. The nanoparticles, nanoCCoE, nanoCCoM and nanoCCoH
showed effective antifungal activity against *M. oryzae* PO1 (rice blast) with $ED_{50}$ values of 9.41, 16.51 and
33.41 ppm, respectively (Table 2). It observed that the spores were abnormally shape and broken cells after
treated with nanoparticles of *C. cochliodes* CTh05 and spores were normal shape in the non-treated control
(Figure 7).

**TABLE 1** Fungal metabolites of *Chaetomium cochliodes* CTh05 against *Magnaporthe oryzae* PO1 at 12 days

| metabolites | Concentration (ppm) | Colony diameter (cm)[2,3] | Growth inhibition (%)[2,3] | Number of spores [2,3]($10^5$) | Spore Inhibition (%)[2,3] | $ED_{50}$(ppm) |
|---|---|---|---|---|---|---|
| CCoH | 0 | 5.00[a] | - | 20.75[a] | - | |
| | 10 | 4.88[bc] | 2.25[ij] | 20.25[a] | 2.56[j] | |
| | 50 | 4.73[d] | 5.25[gh] | 17.25[b] | 16.86[h] | 374.43 |
| | 100 | 4.43[f] | 11.25[e] | 14.25[c] | 31.23[g] | |
| | 500 | 3.91[i] | 21.75[b] | 10.00[e] | 52.18[de] | |
| | 1000 | 3.68[j] | 26.25 [a] | 6.75[fg] | 67.57[c] | |
| CCoE | 0 | 5.00[a] | - | 20.75[a] | - | |
| | 10 | 4.94[ab] | 1.00[j] | 17.25[b] | 16.70[h] | |
| | 50 | 4.82[c] | 3.50[i] | 12.75[cd] | 38.64[f] | 85.87 |
| | 100 | 4.67[d] | 6.50[g] | 9.00[ef] | 56.69[d] | |
| | 500 | 4.02[h] | 19.50[c] | 5.00[gh] | 76.01[b] | |
| | 1000 | 3.64[j] | 27.00[a] | 2.50[h] | 88.03[a] | |
| CCoM | 0 | 5.00[a] | - | 20.75[a] | - | |
| | 10 | 4.86[c] | 2.75[i] | 18.75[ab] | 9.68[i] | |
| | 50 | 4.73[d] | 4.00[hi] | 14.00[c] | 32.58[g] | 144.23 |
| | 100 | 4.53[e] | 9.25[f] | 11.00[de] | 46.95[e] | |
| | 500 | 4.11[g] | 17.75[d] | 7.00[fg] | 66.40[c] | |
| | 1000 | 3.91[i] | 21.75[b] | 4.00[h] | 80.74[b] | |
| C.V.(%) | | 0.74 | 7.22 | 10.79 | 6.27 | - |

[1]/Average of four replications. Means followed by a common letter are not significantly different at P=0.05 level;

[2]/Average of four replications. Means followed by a common letter are not significantly different at P=0.01 level;

[3]/Inhibition(%)=R1-R2/R1x100 where R1 was colony diameter of pathogen in control and R2 was colony diameter of pathogen in treated plates.

**TABLE 2** Nano particles derived from *Chaetomium cochliodes* CTh05 against *Magnaporthe oryzae* PO1 at 12 days

| Nano-particles | Concentration (ppm) | Colony diameter (cm)[1] | Growth inhibition (%)[1,3] | Number of spores [2,3]($10^5$) | Spore Inhibition (%)[1,3] | $ED_{50}$ (ppm) |
|---|---|---|---|---|---|---|
| Nano-CCoH | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.96[ab] | 0.75[e] | 67.00[ab] | 2.78[i] | |
| | 5 | 4.88[bc] | 2.25[de] | 56.00[bcd] | 18.98[f] | 33.41 |
| | 10 | 4.82[cd] | 3.50[cd] | 50.75[cd] | 26.63[e] | |
| | 15 | 4.76[d] | 4.75[c] | 45.00[de] | 34.87[cd] | |
| Nano-CCoE | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.93[ab] | 1.25[e] | 60.00[abc] | 13.23[fg] | |
| | 5 | 4.87[bc] | 2.50[de] | 49.00[cd] | 28.99[de] | 9.47 |
| | 10 | 4.78[d] | 4.25[cd] | 36.00[e] | 47.99[b] | |
| | 15 | 4.58[e] | 8.25[b] | 22.00[f] | 68.26[a] | |
| Nano-CCoM | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.87[bc] | 2.50[de] | 64.00[ab] | 7.36[gh] | |
| | 5 | 4.77[d] | 3.75[cd] | 51.25[cd] | 26.00[e] | 16.51 |
| | 10 | 4.56[e] | 8.75[b] | 44.00[de] | 36.40[c] | |
| | 15 | 3.88[f] | 22.25[a] | 36.00[e] | 47.97[b] | |
| C.V.(%) | | 1.18 | 25.04 | 11.03 | 13.91 | - |

[1]/Average of four replications. Means followed by a common letter are not significantly different at P=0.05 level.

[2]/Average of four replications. Means followed by a common letter are not significantly different at P=0.01 level.

[3]/Inhibition(%)=R1-R2/R1x100 where R1 was colony diameter of pathogen in control and R2 was colony diameter of pathogen in treated plates.

**FIG 7** Normal spores of rice blast pathogen (A) and abnormal spores (B) of *Magnaporthe oryzae* isolate PO1
after treated with nano-particles derived *Chaetomium cochliodes* CTh05 (B).

**DISCUSSION**

The symptoms of blast on rice var RD57 was isolated and confirmed morphological and phylogenic identified as
*Magnaporthe oryzae. Magnporthe oryzae* isolate PO1 proved to be aggressive isolate to cause blast of rice var.
RD57. *Chaetomium cochliodes* CTh05 was confirmed by morphology and phylogenic identification and it was
actively effective against *Magnaporthe oryzae* isolate PO1 causing rice blast. Bi-culture test proved that *C.*
*cochliodes* CTh05 can suppress the growth of *M. oryzae* PO1. Previous report of Soytong (16) stated that *C.*
*cochliodes* proved to be antagonized *Drechslera oryzae* (brown leaf spot of rice var Pittsanulok 2). It
significantly inhibited colony growth and spores of tested pathogen in bi-culture test. Similar report stated that
*Chaetomium cupreum* CC3003 inhibited *C. -lunata* (leaf spot of rice) in bi-culture test (14). The fungal
metabolites of *C. cochliodes* CTh05 (CCoE, CCoM and CCoH) expressed antifungal activity against *M. oryzae*
isolate PO1 (rice blast) to inhibit spore production which the $ED_{50}$ of 85, 144, 374 ppm. Soytong (16) reported
that metabolites from *C. cochliodes* gave significant inhibition on the spore production of *D. oryzae* (brown leaf
spot of rice) that the ED50 value was 66.45 ppm. Tann and Soytong (14) reported that crude hexane, EtOAc and
MeOH extracts of *C. cupreum* C3003 inhibited spore production of *C. lunata* (leaf spot of rice) with the $ED_{50}$
values of 6.41, 0.83 and 7.81 ppm, respectively.

The secondary metabolites from *C. cochliodes* CTh05 had been reported as the azaphilone types,
cochliodones C and D, chaetoviridines E and F, *epi*-chaetoviridine A, and chaetoviridine A. The
chaetoviridines E showed antimalarial activity against *Plasmodium falciparum*. Whereas, cochliodones C,
chaetoviridines E and F expressed antimycobacterial activity against *Mycobacterium tuberculosis.* Moreover,
chaetoviridines E and F exhibited cytotoxicity against the KB, BC1, and NCI-H187 cell lines (15). This research
finding the fungal metabolites from *C. cochliodes* CTh05 actively against *M. oryzae* P01 causing rice blast
pathogen.

Nano-particles derived from *C. cochliodes* CTh05 (nanoCCoE, nanoCCoM and nanoCCoH) showed
significant antifungal activity against *M. oryzae* PO1 with $ED_{50}$ values of 9, 16 and 33 ppm, respectively. Similar
report of Tann and Soytong (13) stated that nano-CGH, nano-CGE, and nano-CGM derived from *Chaetomium*
*globosum* KMITL-N0805 showed antifungal activity against *Curvularia lunata* (leaf spot of rice) with $ED_{50}$
values of 1.21, 1.19, and 1.93 ppm, respectively.

Dar and Soytong (17) stated that the fungal metabolites from *Chaetoimum globosum* and *Chaetoimum*
*cupreum* which were reacted with polylactic acid and electropun at 25-30 kv, nanomaterial from *C. globosum*
was 241 nanometers,and *C. cupreum* was 171 nanometers. Furthermore, the research finding showed abnormal
spores of *M. oryzae* after treated with nanoparticles from *C. cochliodes* CTh05 which was also reported by Tann

and Soytong (13) that nanoparticles from *C. globosum* can be disrupted and distorted the pathogen cells, and loss
of pathogenicity.  With this, Singh *et al*. (18) stated that nanotechnology in agriculture is revolutionized by
innovating new techniques to control diseases. In this research findings revealed that nano-particles constructed
from fungal metabolites gave the ability for plant disease control. As said by Sharon *et al*. (19) that
nanotechnology is helped in agriculture and reduced environmental pollution by applying nano particles with the
ability to increase the efficiency of pesticide application at lower dose.
151
**MATERIALS AND METHODS**
**Isolation of pathogen and pathogenicity test.** The blast specimens were collected from blast symptom on
leaves of rice var. RD 57. Isolation was done using tissue transplanting technique. Pure culture was maintained
in rice flour agar (RFA) medium. Morphological identification was done under binocular compound microscope
and followed the work of Ou (20). The pathogen is deposited at culture collection, Biocontrol Research Unit,
Faculty of Agricultural Technology, KMITL, Bangkok, Thailand.
158   **Antagonistic fungus.** *Chaetomium cochliodes* CTh05 used in this study which is deposited at culture
159 collection, Biocontrol Research Unit, Faculty of Agricultural Technology, KMITL, Bangkok, Thailand. that
160 reported to be produced antimicrobial activity against malaria (*Plasmopara falciparum*) and tuberculosis
161 (*Mycobacterium tuberculosis*) as well as cancer cell lines (15). This isolate was cultured and incubated in potato
162 dextrose agar (PDA) at room temperature (27-30 C), and morphological identified according to von Arx (21) and
163 Soytong (22).
164   **Molecular identification.** The genomic DNA were separately extracted from *Magnaporthe oryzae* PO1
and *Chaetomium cochliodes* CTh05. Each fungus was cultured in potato dextrose broth (PDB) media for 3 days.
The mycelia were freeze-dried, genomic DNA was extracted by modified CTAB (Cetyl trimethyl ammonium
bromide) method, cleaned with 25mM EDTA by centrifugation. 100 mg of mycelia were crushed in liquid
nitrogen. The fungal cells were lysed in CTAB buffer,  β-mercaptoethanol, then incubated at 65 ℃ for 1h in
mixing tubes every 15 min. The lysate were extracted with an equal volume of chloroform/isoamyl alcohol
(24:1), then centrifuged at 14,000 rpm for 5 min at 4 ℃. The aqueous phase was added 2 μl Rnase (20 μg/ml), and
incubated 30 min at 37 ℃, then mixed with 50 μl 10% CTAB. The genomic DNA was precipitated in isopropanol
and centrifuged at 4 ℃ for 20 min at 14,000 rpm. The resulting pellets were washed twice with 70% and 95%
ethanol, air dried and dissolved in 100 μl TE buffer at 37 ℃, overnight. The quality and quantity of extracted
DNA were monitored by electrophoresis in a 1% agarose gel. Quantification was performed with comparison to
the known dilution of lambda phage DNA. Polymerase chain reaction (PCR) was performed by ITS ribosomal
DNA regions which amplified by PCR using the universal primers, ITS1 (5'-TCCGTAGGTGAACCTGCGG-3')
and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (23). The 25 μl reaction mixture contained 2.5 μl 10 × PCR
buffer, 0.625 μl each dNTP (1.25mM), 0.5 μl MgCl$_2$, 1 μl of each primer (20pmol/ μl), 2 ng of DNA and 0.2 μl of
Taq DNA polymerase (1 U) were performed. PCR condition for the ITS regions were programed with initial
denaturation at 95 ℃ for 5 min, followed by 35 cycles of 95 ℃ for 1 min., 56 ℃ for 1 min., 72 ℃ for 2 min., and
a final extension at 72 ℃ for 5 min. The amplified products (5 μl) were visualized on 1% (w/v) agarose gel for
the presence of a single amplified band. The amplified products were sequenced and aligned with comparison to
the sequences in the GenBank by basic local alignment search tool (BLAST) analysis (24) at National Center for
Biotechnology Information (NCBI) databases. The sequences of closely related organisms were downloaded to
construct the phylogenetic trees, aligned through CLUSTALW using MEGA version 6.0 software (25) and a
phylogenetic tree was designed by neighbor-joining method using the same software.

**Bi-culture test.** The test was done by followed the method of Soytong and Quimio (26). *C. cochliodes*
CTh05 and rice blast pathogen were transferred each 0.3 m culture agar plug to PDA at opposite sites. *C.*
*cochliodes* CTh05 and rice blast pathogen were separately cultured where each isolate on PDA served as the
controls. All plates were incubated at room temperature for 30 days. Data were collected as colony diameter (cm)
and sporulation and subjected to analysis of variance and compared using Duncan Multiple's Range Test
(DMRT) at P=0.05 and 0.01.

**Fungal metabolites from *Chaetomium cochliodes* CTh05.** Biomass culture of *Chaetomium cochliodes*
CTh05 was extracted with hexane, ethyl acetate and methanol using method described Phonkerd *et al.* (15) to
give crude hexane (CCoH), ethyl acetate (CCoE) and methanol (CCoM) extracts. . The experiment was set up as
two factor factorial experiment in CRD with four replications. Factor A represented crude extracts and factor B
represented various concentrations of 0, 10, 50, 100, 500 and 1,000 ppm. The culture agar plug of 3 mm was
transferred to the middle PDA plate in each treatment, then incubated at room temperature (27-30 C) for 15 days.
Data were collected as colony diameter (cm) and counted number of spores using haemacytometer.  Data were
statistically computed analysis of variance (ANOVA). Treatment means were compared using DMRT at P=0.05
and 0.01. The effective dose ($ED_{50}$) was computed using probit analysis program.

**Nano-particles.** The crude extracts from *C. cochliodes* CTh05, CCoH, CCoE and CCoM were separately
incorporated into polylactic acid (PLA)-based nanoparticles through electrospinning by followed the method of
Dar and Soytong (17) to yield nanoCCoH, nanoCCoM and nanoCCoE, respectively. Nano CCoH, CCoM and
CCoE were tested to inhibit the rice blast pathogen using two factorial in CRD with four replications. Factor A
was nanoparticles, and factor B was 0, 3, 5, 7, 10 and 15 ppm. CRD was designed for the experiment which
repeated four times. Data were collected as colony diameter (cm) and counted number of spores using
haemacytometer, and statistically calculated analysis of variance (ANOVA). Means were compared using
DMRT at P=0.05 and 0.01. The effective dose ($ED_{50}$) was calculate using probit analysis program.

**ACKNOWLEDGMENT**

I would like to acknowledge the King Mongkut's Institute of Technology Ladkrabang (KMITL) to offer a
PhD Scholarship and research fund is supported by Faculty of Agricultural Technology, KMITL and further
research fund [KREF126004] supported by KMITL, Bangkok, Thailand. The financial support from Thailand
Research Fund (Grant No RTA5980002) is also gratefully acknowledged.

**REFERENCES**

1. Wang GL, Valent B. (eds.). 2009. Advances in
Genetics, Genomics and Control of Rice Blast
Disease. DOI 10.1007/978-1-4020-9500-9 4.
2. TeBeest DO, Guerber C, Ditmore M. 2007.
Rice blast. The Plant Health Instructor. DOI:
10.1094/PHI-I-2007-0313-07.Reviewed 2012.
3. Valent B. 2004. Plant disease: Underground
life for rice foe. Nature 431:516-517.
4. Gnanamanickam SS, 2009. Biological Control
of Rice Diseases, Progress in Biological
Control. DOI 10.1007/978-90-481-2465-7 2.

5. Soutter W. 2012. Nanotechnology in
agriculture. AZoNano.com Publishers.
Available at
http://www.azonano.com/article.aspx?Article
ID=3141#1%204. Accessed on May 25, 2013.
6. Li X, Xu H, Chen ZS, Chen G. 2011.
Biosynthesis of nanoparticles by
microorganisms and their applications. Journal
of Nanomaterials 16:270974.
7. Elibol OH, Morisette DD, Denton JP, Bashir R.
2003. Integrated nanoscale silicon sensors

using top-down fabrication. Applied Physics
Letters 83:4613-4615.

8. Salata OV. 2004. Applications of nanoparticles
in biology and medicine. Journal of
Nanobiotechnology 2:3.

9. Ditta A. 2012. How helpful is nanotechnology
in agriculture? Advances in Natural
Sciences:Nanoscience and Nanotechnology
3:033002.

10. Perlatti B, de Souza Bergo PL, das Graas
Fernandes da Silva MF, Fernandes JB, Forim
MR. 2013. Polymeric nanoparticle-based
insecticides: a controlled release purpose for
agrochemicals. In: Insecticides – Development
of Safer and More Effective Technologies. S.
Trdan (ed.). In Tech Open Access, Croatia. Ch
pp. 20.

11. Rai M, Ingle A. 2012. Role of nanotechnology
in agriculture with special reference to
management of insect pests. Appl Microbiol
Biotechnol 94:287-293.

12. Soytong K, Kanokmedhakul S,
Kukongviriyapan V, Isobe M. 2001.
Application of Chaetomium species
(Ketomium) as a new broad spectrum
biological fungicide for plant disease control: a
review article. Fungal Diversity 7:1-15.

13. Tann H, Soytong K. 2016a. Effects of
nanoparticles loaded with *Chaetomium*
*globosum* LMITL0805 extracts against leaf
spot of rice var Sen Pidoa. Malays. Appl. Biol.
45:1-7.

14. Tann H, Soytong K. 2016b. Biological Control
of Brown Leaf Spot Disease Caused by
*Curvularia lunata* and Field Application
Method on Rice Variety IR66 in
Cambodia.AGRIVITA Journal of Agricultural
Science 39:111-117.

15. Phonkerd N, Kanokmedhakul S,
Kanokmedhakul K, Soytong K, Prabpai S,
Kongsearee P. 2008. Bis-spiro-Azaphilones
and Azaphilones from the Fungi *Chaetomium*

*cochliodes* VTh01 and *C. cochliodes* CTh05.
Tetrahedron 64:9636-9645.

16. Soytong K. 2014. Bio-formulation of
*Chaetomium cochliodes* for controlling brown
leaf spot of rice. International Journal of
Agricultural Technology 10:321-337.

17. Dar J, Soytong K. 2014. Construction and
characterization of copolymer nanomaterials
loaded with bioactive compounds from
*Chaetomium* species. Journal of Agricultural
Technology 10: 823-831.

18. Singh S, Singh BK, Yadav SM, Gupta AK.
2015. Applications of Nanotechnology in
Agricultural and their Role in Disease
Management. Research Journal of Nanoscience
and Nanotechnology 5:1-5.

19. Sharon M, Choudhary AK, Kumar R. 2010.
Nanotechnology in agricultural diseases and
food safety. J. Phytol 2:83-92.

20. Ou SH. 1985. Rice Diseases. Commonwealth
Mycological Institute. pp. 380.

21. Arx JA, von Guarro J, Figuera MJ. 1986. The
Ascomycete genus *Chaetomium*. Beihefte
zurNovaHedwigia 84:1-162.

22. Soytong K. 1989. Antagonism of *Chaetomium*
*cupreum* to *Pyricularia oryzae*: a case study to
biocontrol of a rice blast disease. Thai
Phytopathology 9: 28-33.

23. White TJ, Bruns T, Lee S, Taylor J. 1990.
Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. In
PCR Protocols: a guide to methods and
applications (M. A. Innis, D. H. Gelfand, J. J.
Sninsky and T. J. White, eds). Academic Press
315-322.

24. Altschul SF, Madden TL, Schaffer AA, Zhang
JH, Zhang Z, Miller W, Lipman DJ. 1997.
Gapped BLAST and PSI-BLAST: a new
generation of protein database search programs.
Nucleic Acids Res 25:3389-3402.

25. Tamura K, Dudley J, Nei M, Kumar S. 2007.
MEGA4: Molecular evolutionary genetics

analysis (MEGA) software version 4.0. Mol.
Bio. Evol. 24:1596-1599.
26. Soytong K, Quimio TH. 1989. Antagonism of
*Chaetomium globosum* to the rice blast
pathogen, *Pyricularia oryzae*. Kasetsart
Journal (Natural Science) 23:198-203.

**FIG1** *Magnaporthe oryzae* PO1, A= pure culture in PDA; B= conidia; C = mycelium and conidia.

**FIG 2** *Chaetomium cochliodes* CTh05, A=Colony, B= Perithecia, C=Asci, D=Ascospores.

**FIG 3** Phylogenic tree of *Magnporthe oryzae* (PO1)

FIG 4 Phylogenic tree of *Chaetomium cochliodes* CTh05

FIG 5 Pathogenicity test on rice leaf

Left = Control

Right = Treatment

FIG 6 Bi-culture test between *Magnaporthe oryzae* isolate PO1 and *Chaetomium cochliodes* CTh05, Left = Chaetomium culture, Middle = Bi-culture plates, and Right = Magnaporthe culture

FIG 7 Normal spores of rice blast pathogen (A) and abnormal spores (B) of *Magnaporthe oryzae* isolate PO1 after treated with nano-particles derived *Chaetomium cochliodes* CTh05 (B).

**TABLE 1** Fungal metabolites of *Chaetomium cochliodes* CTh05 against *Magnaporthe oryzae* PO1 at 12 days

| metabolites | Concentration (ppm) | Colony diameter (cm)[2,3] | Growth inhibition (%)[2,3] | Number of spores [2,3](10$^5$) | Spore Inhibition (%)[2,3] | ED$_{50}$(ppm) |
|---|---|---|---|---|---|---|
| CCoH | 0 | 5.00$^a$ | - | 20.75$^a$ | - | |
| | 10 | 4.88$^{bc}$ | 2.25$^{ij}$ | 20.25$^a$ | 2.56$^j$ | |
| | 50 | 4.73$^d$ | 5.25$^{gh}$ | 17.25$^b$ | 16.86$^h$ | 374.43 |
| | 100 | 4.43$^f$ | 11.25$^e$ | 14.25$^c$ | 31.23$^g$ | |
| | 500 | 3.91$^i$ | 21.75$^b$ | 10.00$^e$ | 52.18$^{de}$ | |
| | 1000 | 3.68$^j$ | 26.25 $^a$ | 6.75$^{fg}$ | 67.57$^c$ | |
| CCoE | 0 | 5.00$^a$ | - | 20.75$^a$ | - | |
| | 10 | 4.94$^{ab}$ | 1.00$^j$ | 17.25$^b$ | 16.70$^h$ | |
| | 50 | 4.82$^c$ | 3.50$^i$ | 12.75$^{cd}$ | 38.64$^f$ | 85.87 |
| | 100 | 4.67$^d$ | 6.50$^g$ | 9.00$^{ef}$ | 56.69$^d$ | |
| | 500 | 4.02$^h$ | 19.50$^c$ | 5.00$^{gh}$ | 76.01$^b$ | |
| | 1000 | 3.64$^j$ | 27.00$^a$ | 2.50$^h$ | 88.03$^a$ | |
| CCoM | 0 | 5.00$^a$ | - | 20.75$^a$ | - | |
| | 10 | 4.86$^c$ | 2.75$^i$ | 18.75$^{ab}$ | 9.68$^i$ | |
| | 50 | 4.73$^d$ | 4.00$^{hi}$ | 14.00$^c$ | 32.58$^g$ | 144.23 |
| | 100 | 4.53$^e$ | 9.25$^f$ | 11.00$^{de}$ | 46.95$^e$ | |
| | 500 | 4.11$^g$ | 17.75$^d$ | 7.00$^{fg}$ | 66.40$^c$ | |
| | 1000 | 3.91$^i$ | 21.75$^b$ | 4.00$^h$ | 80.74$^b$ | |
| C.V.(%) | | 0.74 | 7.22 | 10.79 | 6.27 | - |

[1]/Average of four replications. Means followed by a common letter are not significantly different at P=0.05 level;
[2]/Average of four replications. Means followed by a common letter are not significantly different at P=0.01 level;
[3]/Inhibition(%)=R1-R2/R1x100 where R1 was colony diameter of pathogen in control and R2 was colony diameter of pathogen in treated plates.

**TABLE 2** Nano particles derived from *Chaetomium cochliodes* CTh05 against *Magnaporthe oryzae* PO1 at 12 days

| Nano-particles | Concentration (ppm) | Colony diameter (cm)[1] | Growth inhibition (%)[1,3] | Number of spores [2,3]($10^5$) | Spore Inhibition (%)[1,3] | $ED_{50}$ (ppm) |
|---|---|---|---|---|---|---|
| Nano-CCoH | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.96[ab] | 0.75[e] | 67.00[ab] | 2.78[i] | |
| | 5 | 4.88[bc] | 2.25[de] | 56.00[bcd] | 18.98[f] | 33.41 |
| | 10 | 4.82[cd] | 3.50[cd] | 50.75[cd] | 26.63[e] | |
| | 15 | 4.76[d] | 4.75[c] | 45.00[de] | 34.87[cd] | |
| Nano-CCoE | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.93[ab] | 1.25[e] | 60.00[abc] | 13.23[fg] | |
| | 5 | 4.87[bc] | 2.50[de] | 49.00[cd] | 28.99[de] | 9.47 |
| | 10 | 4.78[d] | 4.25[cd] | 36.00[e] | 47.99[b] | |
| | 15 | 4.58[e] | 8.25[b] | 22.00[f] | 68.26[a] | |
| Nano-CCoM | 0 | 5.00[a] | - | 69.00[a] | - | |
| | 3 | 4.87[bc] | 2.50[de] | 64.00[ab] | 7.36[gh] | |
| | 5 | 4.77[d] | 3.75[cd] | 51.25[cd] | 26.00[e] | 16.51 |
| | 10 | 4.56[e] | 8.75[b] | 44.00[de] | 36.40[c] | |
| | 15 | 3.88[f] | 22.25[a] | 36.00[e] | 47.97[b] | |
| C.V.(%) | | 1.18 | 25.04 | 11.03 | 13.91 | - |

[1]/Average of four replications. Means followed by a common letter are not significantly different at P=0.05 level. [2]/Average of four replications. Means followed by a common letter are not significantly different at P=0.01 level. [3]/Inhibition(%)=R1-R2/R1x100 where R1 was colony diameter of pathogen in control and R2 was colony diameter of pathogen in treated plates.