Oxygen Restriction Generates Difficult-To-Culture Pathogens

Lasse Kvich [2], Blaine Fritz [2], Stephanie Crone [2], Kasper N. Kragh [2], Mette Kolpen [1], Majken Sønderholm [2], Mikael Andersson [3], Anders Koch [3,4], Peter Ø. Jensen [1,2*], Thomas Bjarnsholt [1,2*]

[1] Department of Clinical Microbiology, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark.

[2] Costerton Biofilm Center, Department of Immunology and Microbiology, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.

[3] Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark.

[4] Department of Infectious Diseases, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark.

* Correspondence:

Professor Thomas Bjarnsholt
Costerton Biofilm Center
Department of Immunology and Microbiology
University of Copenhagen
Blegdamsvej 3b, Room 22.2.22
DK-2200 Copenhagen N
Denmark.
Tlf: +45 2065 9888
tbjarnsholt@sund.ku.dk

Associated Professor Peter Ø. Jensen
Costerton Biofilm Center
Department of Immunology and Microbiology
University of Copenhagen
Blegdamsvej 3b, Room 24.1.14
DK-2200 Copenhagen N
Denmark.
Tlf: +45 3031 9073
peter.oestrup.jensen@regionh.dk

ORCID: LK 0000-0003-3226-3633, BF 0000-0002-2085-894X, KK 0000-0002-7662-9118, MK 0000-0001-5575-0384, MS 0000-0001-9576-5775, AK 0000-0001-9205-1048, PJ 0000-0003-1648-7186, TB 0000-0002-8003-7414

Keywords:

P. aeruginosa biofilm / anoxia / biofilm / reactive oxygen species / difficult-to-culture / viable but non-culturable

Abstract: 233 words

Document: 4604 words

Abstract

Induction of a non-culturable state has been demonstrated for many bacteria. In a clinical

perspective, the lack of growth due to these non-culturable bacteria can have major consequences

for the diagnosis and treatment of patients. Here we show how anoxic conditioning (restriction of

molecular oxygen, O_2) generates difficult-to-culture (DTC) bacteria during biofilm growth.

A significant subpopulation of *Pseudomonas aeruginosa* entered a DTC state after anoxic

conditioning, ranging from five to 90 % of the total culturable population, in both planktonic and

biofilm models. Anoxic conditioning also generated DTC subpopulations of *Staphylococcus aureus*

and *Staphylococcus epidermidis* (89 and 42 % of the total culturable population, respectively).

Growth of the DTC populations were achieved by substituting O_2 with 10 mM NO_3^- as an

alternative electron acceptor for anaerobic respiration or, in the case of *P. aeruginosa*, by adding

sodium pyruvate or catalase as scavengers against reactive oxygen species (ROS) during aerobic

respiration. An increase in normoxic plating due to addition of catalase suggests the molecule

hydrogen peroxide as a possible mechanism for induction of DTC *P. aeruginosa.* Anoxic

conditioning also generated a true viable but non-culturable (VBNC) population of *P. aeruginosa*

that was not resurrected by substituting O_2 with NO_3^- during anaerobic respiration.

These results demonstrate that habituation to an anoxic micro-environment could complicate

diagnostic culturing of bacteria, especially in the case of chronic infections where oxygen is

restricted due to the host immune response.

Importance

Diagnostics of bacteria from chronic infections by standard culture-based methods is challenging.

Bacteria in a non-culturable state may contribute to the lack of culturing from these infections.

Many stressors are known to induce a non-culturable state, among others the absence of molecular

oxygen, which is evident in chronic infections due to high rates of oxygen consumption by the host

response. In this study, we have shown that *Pseudomonas aeruginosa*, *Staphylococcus aureus* and

Staphylococcus epidermidis can enter a non-culturable state after oxygen restriction. Regrowth was

not possible using conventional normoxic plating where oxygen served as electron acceptor.

Instead, regrowth was enabled during anoxic conditions with added nitrate as alternative electron

acceptor. In the case of *P. aeruginosa* we were able to counteract this non-culturable state by adding

scavengers for reactive oxygen species during normoxic incubation. This suggest a link between

reactive oxygen species and a non-culturable state and it also renders a solution to circumvent this

problem in a clinical diagnostic setting. Our findings show that bacteria can habituate to their

environment and that it has to be taken into consideration especially when culturing clinical samples

e.g. from chronic infections.

Introduction

Clinical laboratories often culture bacteria with enriched media, developed to support maximal

growth of particular pathogens. However, even these bacterial human pathogens can enter a growth-

restricted, transient state resulting in loss of culturability, especially following antibiotic treatment

(Pasquaroli et al., 2013). Non-culturable bacteria are apparent by microscopic identification in

clinical cases presenting sign of infection, but no positive cultures. These cases are especially

prevalent during chronic infection, where lack of culturability prevents proper diagnosis and

treatment (Høiby et al., 2010;Costerton et al., 2011;Høiby et al., 2015).

Two well defined, yet difficult to distinguish, non-culturable states have been described for

bacteria: viable but non-culturable (VBNC) and persisters (Oliver, 2010;Wood et al., 2013;Fisher et

al., 2017;Kim et al., 2018). Several environmental stresses have been associated with non-

culturable states, e.g. oxidative stress, change in temperature or pH, nutrient starvation, change in

osmotic concentrations and presence of heavy metals or antibiotics (Oliver, 2010;Ayrapetyan et al.,

2015). Oxygen starvation, referred to as anoxic conditioning in this paper, has also shown to induce

a non-culturable state in batch cultures of *Pseudomonas aeruginosa*, but supplementation with

nitrate (NO_3^-) as an alternative electron acceptor during anoxic plating restored culturability

(Binnerup, 1993). The lack of growth during normoxic plating suggests that anoxic conditioning

sensitizes a subpopulation of *P. aeruginosa* to O_2 or its toxic derivatives, such as reactive oxygen

species (ROS). ROS are continuously created during aerobic respiration by incomplete reduction of

O_2, which is toxic to cells if not scavenged (Fenchel and Finlay, 2008). Presence of ROS has, in

some studies, been shown to generate VBNC bacteria (Oliver, 2010;Noor, 2015), but most research

has focused on *Escherichia coli* strains or *Vibrio* spp., while the effects of ROS on *P. aeruginosa*

and other facultative pathogens are not well characterized.

Evidence of anoxic zones and anaerobic bacterial activity in chronic infections suggests that

colonizing bacteria, such as *P. aeruginosa,* experience anoxic conditioning (Hassett et al.,

2002;Worlitzsch et al., 2002;Kolpen et al., 2014). High rates of O_2 consumption by

polymorphonuclear leukocytes generate local, anoxic zones and likely play a major role in O_2

depletion during chronic infection (Høiby et al., 2015). In addition, other cells of the host may also

consume O_2, increasing the likelihood of bacteria experiencing anoxia (Worlitzsch et al., 2002).

Many chronic infections are thought to contain bacteria in the biofilm mode of growth (Costerton et

al., 2003) and endogenous O_2 depletion inside the biofilm (Sønderholm et al., 2018), along with

intense O_2 consumption by the host immune response (Worlitzsch et al., 2002;Kolpen et al.,

2010;Trunk et al., 2010;Kragh et al., 2014;James et al., 2016;Jensen et al., 2017), may therefore

increase the number of VBNC bacteria.

This study aimed to determine whether anoxic conditioning generates VBNC cells of *P. aeruginosa*

during biofilm growth. Furthermore, we investigated whether ROS are involved in the loss of

culturability.

Materials and Method

Bacterial strains

Pseudomonas aeruginosa (PAO1, ATCC 15692), a catalase A deficient *Pseudomonas aeruginosa*

strain (Δ*katA* PAO1) (Hassett et al., 1999), *Staphylococcus epidermidis* ATCC 14990,

Staphylococcus aureus NCTC 8325-4 (methicillin susceptible) (Frees et al., 2004), *Staphylococcus*

aureus USA300 JE2 (MRSA) (Fey et al., 2013), E*scherichia coli* CFT073 (Welch et al., 2002) and

a clinical strain of *Enterococcus faecalis* from the Department of Clinical Microbiology,

Copenhagen University Hospital – Rigshospitalet, Denmark were used in this study.

Agar plates and media

Primarily lysogeny broth (LB) agar plates were applied in this study. Lysogeny broth (pH 7.5)

consisted of 5 g/L yeast extract (Oxoid, Roskilde, Denmark), 10 g/L tryptone (Oxoid), 10 g/L NaCl

(Merck, USA). All plates contained 2 % agarose. Plates used for anoxic growth were supplemented

with 10 mM KNO_3 (Sigma-Aldrich, USA) to serve as alternative electron acceptor during anoxic

plating, referred to as NO_3^- throughout the paper. All agar plates and media in this study were

supplied by the Panum Institute Substrate Department (Copenhagen, DK).

Anoxic growth

Experiments investigating growth under anoxic conditions were performed in an anaerobic chamber

(Concept 400 Anaerobic Workstation, Ruskinn Technology Ltd, UK). The gas atmosphere

consisted of $N_2/H_2/CO_2$ (ratio - 80:10:10). Anoxic chamber environment was confirmed with an

oxygen sensor (HQ40d multi, HACH Company, USA). All media and chemical solutions used in

anaerobic experiments were equilibrated in the anaerobic chamber 3 days prior to experiment. In

the case of solutions requiring refrigeration, a minimal volume was applied (< 1 mL), sealed with

Parafilm M, and thoroughly shaken upon entry into anaerobic chamber for quick gas equilibration.

Direct viable count with LIVE/DEAD staining

LIVE/DEAD staining was applied to estimate the proportion of viable and non-viable cells in 24-hour-old batch cultures and in 16-day-old anoxically conditioned batch cultures of *P. aeruginosa*. The dyes consisted of two fluorescent nucleic acid stains, the green fluorescent stain (live cells) SYTO9 (Invitrogen, USA) and the red fluorescent stain (dead cells) propidium iodide (PI, Sigma-Aldrich, USA). SYTO9 penetrates both intact and damaged membranes while PI only stains damaged cells, thereby creating an opportunity to discriminate between live and dead cells (Li et al., 2014). Bacterial suspensions were vortexed thoroughly (1 min) and sonicated (5 min degas + 5 minutes sonication) before staining. To stain the cells, one µL of PI and SYTO9 was added to 1 mL of bacterial suspension and incubated for 15 minutes at room temperature. Suspensions of each biological replicate were then filtered through a 0.2 µm black Whatman, Nuclepore Trach-Etch Membrane (Sigma-Aldrich, Denmark) and visualized with confocal laser scanning microscopy using a Zeiss LSM 710 with a 63×/1.4 (numerical aperture) objective (Zeiss, Germany). Fifteen random fields (135µm x 135 µm) were examined for each filter (n=3). Enumeration of live (green) and dead (red) bacteria were done with the IMARIS software package (Bitplane AG, Schwitzerland). Briefly, individual particles in each field were identified and counted as bacteria. Particles that were stained with both PI and SYTO9 were considered non-viable and thus counted as dead cells. Conversion of enumerated viable and dead cells to bacterial counts per milliliter was performed as described (Boulos et al., 1999) to compare them with CFU/mL. Briefly, the numbers of particles per mL were estimated using the following equation $\frac{particles}{mL} = A * (\frac{B}{C})/D$. Where A is the amount of particles per field, B is the surface of filtration (mm), C is the area of the microscopic field and D the volume of the sample filtered. Normoxic and anoxic CFU/mL was carried out simultaneously to estimate the proportion of VBNC cells. CFU/mL was determined in the same way as described for liquid batch cultures.

Bead-embedded inoculum of *P. aeruginosa*

Preparation of alginate beads with *P. aeruginosa* was carried out according to a method described

by Sønderholm *et al*. (Sønderholm et al., 2017). Subsequently, beads were divided (10 beads per

vial) into vials (Oximate Vial, PerkinElmer Inc., USA) containing 15 mL LB medium. Vials were

sealed with Parafilm M (oxygen permeable) and incubated at 37°C on an orbital shaker at 180 rpm.

Anoxic and normoxic CFU counts were determined from beads and from the surrounding

suspension from the same vials on each sampling day (day 3, 6, 8, 12, 19 and 21). Two beads were

sampled per biological replicate (n=4). Before determination of CFU, the beads were washed twice

with 0.9 % NaCl to remove non-attached cells and transferred to 1.5 mL microcentrifuge tubes

(Sigma-Aldrich, USA). One-hundred microliters of 0.1 M sodium carbonate (Na_2CO_3) followed by

100 µL of 0.04 M citric acid was added to the tubes to dissolve the beads. The suspensions were

then sonicated (5 min degas + 5 minutes sonication; Bransonic ultrasonic cleaner 2510, Emerson

Electric, USA) before ten-fold dilution series were performed in 0.9 % NaCl. CFU was determined

by plating three, 10 µL-drops per dilution per replicate. Anoxic CFU determinations were

performed inside an anaerobic chamber on LB plates supplemented with 10 mM NO_3^-. The same

dilution series were applied to normoxic CFU determination outside the anaerobic chamber on LB

plates with 10 mM NO_3^-. Plates were incubated 2 days before counting CFU.

Filter biofilms with *P. aeruginosa*

This protocol was adapted to grow reproducible biofilms under anoxic and normoxic conditions.

The method has previously been described by Bjarnsholt *et al.* (2015). The filter biofilms were kept

on the same LB plates throughout the experiment. *P. aeruginosa* was propagated from frozen stock

and grown overnight in 20 mL LB medium at 37°C on an orbital shaker at180 rpm. Cultures were

adjusted to an optical density of 0.05 (OD_{600}; UV spectrophotometer UV-1800 UV-VIS, Shimadzu

corporation, JP) and 10 µL was transferred to the cellulose nitrate membrane filters (25 mm in

diameter, GE Healthcare Life Sciences, UK). Plates were incubated under normoxic and anoxic

conditions and kept in plastic bags with wet paper to avoid dehydration. Two filters were sampled

per biological replicate (n=4) on each sampling day (day 1, 3, 7, 15 and 17). Filters were removed,

placed in 10 mL tubes containing 5 mL 0.9 % NaCl, vortexed thoroughly (one min.) and sonicated

(5 min degas + 5 minutes sonication). CFU/mL was determined as previously described.

Liquid batch cultures of *P. aeruginosa*

P. aeruginosa was propagated from frozen stock and grown overnight in 20 mL LB media at 37°C

and orbitally shaken at 180 rpm. Cultures were adjusted to an optical density (OD$_{600}$) of 0.1 in glass

vials (Oximate Vial) with a final volume of 20 mL LB. Vials were left to incubate at 37°C and

orbitally shaken at 180 rpm. To create a normoxic environment, half of the vials were incubated

with Parafilm M on top (normoxic conditioning), while the rest were incubated with a lid on top

creating an anoxic environment (anoxic conditioning). Anoxic and normoxic plating was carried out

from normoxic and anoxic conditioned liquid batch cultures (referred to as batch cultures

throughout the paper) on each sampling day (day 1, 3, 5, 9, 11, 14, 16, 18, 21 and 28). Normoxic

plating was carried out on LB plates with and without 10 mM NO$_3^-$ to test whether presence of NO$_3^-$

affected the number of CFU. Two mL (2 x 1 mL = 2 technical replicates) were sampled per

biological replicate (n=4) on each sampling day. CFU/mL was determined as previously described.

Colonies of *P. aeruginosa*

P. aeruginosa was propagated from frozen stock and grown overnight in 20 mL LB media at 37°C

on an orbital shaker at 180 rpm. The cultures were then streaked onto LB plates supplemented with

10 mM NO$_3^-$. Plates were incubated under normoxic and anoxic conditions and kept in plastic bags

with wet paper to avoid dehydration. A one-μl loop was used to sample colony material from each

biological replicate (n=3) on each sampling day (day 6, 9, 13, 15 and 20). Colonies were transferred

to 1.5mL microcentrifuge tubes (Sigma-Aldrich, Denmark) with 0.5 mL 0.9 % NaCl, vortexed

thoroughly (1 min) and sonicated (5 min degas + 5 minutes sonication). CFU/mL was determined

as previously described.

Reactive oxygen species (ROS)

To elucidate whether growth of anoxically conditioned bacteria was restricted by creation of ROS

during aerobic respiration, sodium pyruvate and catalase (Sigma-Aldrich, USA) was tested as ROS

scavenger in LB plates with 10 mM NO_3^-. Sodium pyruvate (0.3 %) was tested during anoxic and

normoxic plating for 24-hour-old batch cultures and anoxically conditioned 16-day-old batch

cultures of *P. aeruginosa*. Catalase (~50,000 units of catalase per L medium) was tested during

anoxic and normoxic plating for anoxically conditioned 16-day-old batch cultures of *P. aeruginosa*.

Furthermore, CFU/mL was also determined from 16-day-old anoxically conditioned batch cultures

of *P. aeruginosa* and a catalase A deficient *P. aeruginosa* (Δ*katA* PAO1) to test the influence of

ROS. Lastly, creation of ROS was measured along with optical density (OD) at day 1, 8 and 16 for

normoxically and anoxically conditioned batch cultures of *P. aeruginosa*. Creation of ROS was

measured by adding 5 µM 2′,7′dichlorodihydrofluorescein diacetate (DCHF-DA; Sigma) to the

bacterial suspension. This concentration has previously been used to detect ROS for *P. aeruginosa*

(Jensen et al., 2014). Two hundred microliter of anoxically conditioned cultures (OD_{600} adjusted to

0.1 in fresh LB medium) was added to each well of black 96-well microtiter plates with transparent

flat bottom (16503, Thermo Fisher Scientific, Rochester, NY). OD and ROS was measured over 15

hours at 27°C with a VICTOR Multilabel Plate Reader (Perkin Elmer, MA). All experiments

consisted of 3-4 biological replicates.

Filter biofilms with other pathogens

To determine if other bacteria behaved in the same way as *P. aeruginosa*, we investigated the effect

of anoxic conditioning on a selected group of pathogens (*Staphylococcus epidermidis* ATCC 14990,

Staphylococcus aureus NCTC 8325-4 (methicillin susceptible), *Staphylococcus aureus* USA300

JE2 (MRSA), E*scherichia coli* CFT073 and a clinical strain of *Enterococcus faecalis*). The filter

biofilm method was used as described earlier. For the methicillin susceptible strain, CFU was

determined in the same way as described for the filters with *P. aeruginosa*. For the remaining

strains, CFU was determined at day 1 and 9 on LB plates supplemented with 10 mM $NO_3^- \pm O_2$.

Furthermore, CFU was determined on LB plates supplemented with 10 mM NO_3^- and 0.3 % sodium

pyruvate to evaluate the effect of ROS. A minimum of three biological replicates was performed

and CFU was carried out in the same way as described earlier.

Statistics

To evaluate the difference in bacterial growth between incubation conditions (normoxic and anoxic

conditioning), a linear regression was used with the difference between the logarithmically

transformed values for normoxic and anoxic plating as outcome and with day (categorical) and

interaction between day and plating condition (binary) as explanatory variables in SAS Genmod

Procedure. The p-value of the interaction term was used as the p-value for the difference in bacterial

growth. Log difference was calculated as ($\log_{10}[CFU/mL]_{anoxic}$ - $\log_{10}[CFU/mL]_{normoxic}$). The mean

and standard error of the mean (SEM) were calculated for recovering bacteria and plotted using

GraphPad Prism 6.1 (GraphPad Software, La Jolla, USA). From the ratio between anoxic (CFU -

O_2) and normoxic (CFU +O_2) colony counts it was possible to calculate the fraction of DTC

bacteria [% DTC bacteria $= \frac{(CFU -O_2)-(CFU +O_2)}{(CFU -O_2)}$ x 100]. Data that were not part of the long-term

experiments were instead analyzed with a 1-way ANOVA followed by Tukey's multiple

comparison tests or an unpaired t-test. A p-value ≤ 0.05 was considered statistically significant. The

tests were performed with either Prism 6.1 (GraphPad Software, La Jolla, USA) or SAS v.9.4 (SAS

Institute Inc., Cary, NC, USA).

Results

Direct viable counts with LIVE/DEAD staining revealed a VBNC *P. aeruginosa* population.

Direct viable counts were carried out for 16-day-old, anoxically conditioned batch cultures of *P.*

aeruginosa to investigate if anoxic conditioning could generate a VBNC population. Findings were

compared to normoxic and anoxic plating (plating methods) performed on LB plates supplemented

with 10 mM NO_3^- (Fig. 1B). Anoxic plating yielded a significantly (p = 0.023) higher count of

colony forming units per milliliter (CFU/mL) than normoxic plating (0.85 log values ± 0.04 SD).

When comparing bacterial plate counts with direct viable counting, significantly more viable cells

were measured compared to normoxic (2.31 log values ± 0.47 SD, p < 0.0001) and anoxic plating

(1.46 log values ± 0.5 SD, p = 0.0009). The cells represented by this difference were considered

VBNC. Direct viable counts were also performed on 24-hour-old batch cultures to investigate

whether findings of VBNC *P. aeruginosa* were restricted to anoxically conditioned cells. No

difference in bacterial counts was observed (Fig. 1A).

Anoxic conditioning generated oxygen intolerant subpopulations of *P. aeruginosa* in biofilm

and planktonic models.

To examine if biofilm growth affected the subsequent normoxic and anoxic plating, *P. aeruginosa*

was grown in an alginate-bead biofilm model (Sønderholm et al., 2017). Plate counts were

performed on LB plates supplemented with 10 mM NO_3^-. Plates were incubated under normoxic

and anoxic conditions and plate counts were compared from the beads (biofilm) and the

surrounding media (planktonic) of the same vials over a period of 21 days (Fig. 2A and 2B,

respectively). The log difference was significantly (p = 0.003) higher for biofilm than planktonic

cells (Fig. 2C). The cells represented by this difference were considered to be in a difficult-to-

culture (DTC) state and ranged from five to 54 % of the entire population from day six to 19 (Fig.

S1).

P. aeruginosa was also grown anoxically or normoxically on a filter biofilm model using LB plates

supplemented with 10 mM NO_3^- over a period of 17 days. CFU/mL was then determined using

anoxic and normoxic plating, as above (Fig. 3A and 3B, respectively). The log difference between

plating methods was significantly (p = 0.01) higher for anoxically conditioned biofilms than

normoxically conditioned biofilms (Fig. 3C). The fraction of DTC *P. aeruginosa* ranged from six to

23 % from day three to 17 (Fig. S1).

Similar results were obtained for anoxically and normoxically conditioned planktonic batch cultures

of *P. aeruginosa* over a period of 28 days (Fig. 4A and 4B, respectively). The log difference

between plating methods was significantly (p < 0.0001) higher for anoxically conditioned batch

cultures (Fig. 3C). The fraction of DTC bacteria ranged from 60 to 90 % from day nine to 21 (Fig.

S1).

The effect of anoxic conditioning was further investigated on colonies of *P. aeruginosa* grown on

LB plates supplemented with 10 mM NO_3^- under anoxic and normoxic conditions over a period of

20 days (Fig. 5A and 5B, respectively). The log difference between plating methods was

significantly higher (p < 0.0001) for anoxically conditioned colonies (Fig. 5C). The fraction of DTC

P. aeruginosa ranged from 12 to 46 % from day six to 20 (Fig. S1).

10 mM NO_3^- in LB plates did not affect CFU/mL in batch cultures of *P. aeruginosa*.

To ensure that presence of NO_3^- did not affect the number of CFU generated on LB plates, CFU/mL

was determined anoxically (LB plates + 10 mM NO_3^-) and normoxically (LB plates ± 10 mM NO_3^-)

from 24-hour-old batch cultures of *P. aeruginosa*. No difference was observed between type of

plating (p = 0.93): anoxic log CFU/mL + NO_3^- = 9.96 (± 0.22 SD), normoxic log CFU/mL + NO_3^- =

9.91 ($\pm$ 0.10 SD) and normoxic log CFU/mL - NO_3^- = 9.91 ($\pm$ 0.14 SD). Furthermore, there was no

effect of NO_3^- supplementation on normoxic plating in the prolonged experiment with 28-day-old

batch cultures (Fig. 4A and 4B). It was not possible to detect growth of *P. aeruginosa* on LB plates

under anoxic conditions without NO_3^-, but growth was observed when these plates subsequently

were placed under normoxic conditions.

Oxidative stress restricted the growth of anoxically conditioned *P. aeruginosa* when re-grown

in a normoxic environment.

Bacteria from 16-day-old anoxically and normoxically conditioned batch cultures of *P. aeruginosa*

were plated on 10 mM NO_3^- LB plates $\pm$ sodium pyruvate and $\pm$ catalase to investigate whether

formation of ROS contributed to the DTC state induced by anoxic conditioning. Presence of sodium

pyruvate as a ROS scavenger (Mizunoe et al., 1999) significantly (p = 0.04) increased normoxic

CFU/mL compared to plating without sodium pyruvate (Fig. 6B). This was not the case in 24-hour-

old batch cultures of *P. aeruginosa* (p = 0.62), indicating that the effect was restricted to anoxically

conditioned cells (Fig. 6A). Presence of catalase also significantly (p = 0.03) improved normoxic

plate counts compared to normoxic plating without catalase (Fig. 6B). These results show that

hydrogen peroxide is involved in the induction of a DTC state created by anoxic conditioning. To

add proof to this hypothesis, a similar experiment was carried out with a catalase deficient *P.*

aeruginosa mutant ($\Delta katA$ PAO1), which is more susceptible to oxidative stress (Hassett et al.,

1999;Jensen et al., 2014). The log difference between plating methods for the $\Delta katA$ mutant and

reference strain was significantly (p = 0.00002, unpaired t-test) different. These log difference

values were 3.02 $\pm$ 0.05 SD and 1.61 $\pm$ 0.07 SD for $\Delta katA$ PAO1 and the reference strain,

respectively (Fig. 6C). Moreover, measurement of ROS and optical density (OD) was carried out

for *P. aeruginosa* at day 1, 8 and 16 from anoxically and normoxically conditioned batch cultures.

These results showed creation of ROS during normoxic growth. The longer PAO1 was anoxically

conditioned; the lower was the following normoxic growth, as measured by OD. This effect was

less pronounced for normoxically-conditioned cells (Fig.7 A-C).

Anoxic conditioning affects *Staphylococcus aureus* and *Staphylococcus epidermidis*, but not

Escherichia coli* and *Enterococcus faecalis

The effect of anoxic conditioning was then tested on a selected group of pathogens to determine if

this phenomenon was restricted to *P. aeruginosa*. *S. aureus* (methicillin susceptible) was tested as

described in the *P. aeruginosa* filter biofilm setup. Plate counts for anoxically and normoxically

conditioned filters was determined (Fig. 8A and 8B, respectively). The log difference between

plating method was significantly higher ($p < 0.001$) when plate counts were performed from

anoxically conditioned filter biofilms than from normoxically conditioned filter biofilms (Fig. 8C).

The fraction of DTC *S. aureus* ranged from three to 89 % from day one to day 17 (Fig. S1). Since

both *P. aeruginosa* and *S. aureus* demonstrated effects of anoxic conditioning, a smaller experiment

was initiated to test other pathogens. The effect was determined on LB plates supplemented with 10

mM NO_3^- ± sodium pyruvate to investigate if ROS were involved in the lack of growth. Both *S.*

aureus (MRSA) and *S. epidermidis* showed effects of anoxic conditioning, resulting in an

significant ($p = 0.01$ and $p = 0.03$, respectively) increase in CFU/mL during anoxic plating

compared to normoxic plating (Fig. S2). The fraction of DTC *S. aureus* (MRSA) and *S. epidermidis*

was 36 and 42 % at day 9, respectively. There was no observed effect of sodium pyruvate during

normoxic plating for these two strains. In the case of *E. coli* and *E. faecalis,* there was no observed

effect of anoxic conditioning (Fig. S3).

Discussion

DTC and VBNC *P. aeruginosa*

In the current study, we demonstrate that plate counting with *P. aeruginosa* is significantly affected

by anoxic conditioning when cultured in the presence of atmospheric oxygen levels. A DTC state

was created after only a few days of anoxic conditioning in both biofilm and planktonic models.

DTC *P. aeruginosa* was observed in the beads from the bead biofilm model, but not in the

surrounding suspension, despite that the cultures had full access to atmospheric oxygen. This

indicates that oxygen restriction within a biofilm generates a DTC subpopulation. The DTC state,

detected as increased growth during anoxic plating with NO_3^-, has previously been generated in

planktonic *P. aeruginosa* by energy starvation after cultivation without O_2 as electron acceptor for

aerobic respiration (Binnerup, 1993). *In vitro* biofilms of *P. aeruginosa* may contain internal anoxic

zones (Walters et al., 2003;Kolpen et al., 2016), so we hypothesized that anoxia inside the biofilm

contributes to the induction of DTC *P. aeruginosa*. Accordingly, DTC *P. aeruginosa* was only

observed in the filter biofilm model and the colony model when the model was anoxically

conditioned. Interestingly, we found that 99.68 % of the population was VBNC with the

LIVE/DEAD assay after anoxic conditioning. LIVE/DEAD staining is based upon membrane

permeability and is only an approximation of true viability. Cells that were stained both red and

green (resulting in yellow) were counted as dead cells but may still be viable.

Approximately 50 % of the population in the alginate beads was DTC even though they were kept

in normoxically-conditioned vials. In comparison, the fraction of DTC cells was approximately 90

% in the anoxically conditioned batch cultures. The difference (90 % vs. 50 %) may be explained

by the fact that a majority of the bacteria in the beads are peripherally located where they have

increased access to oxygen compared to the center of the beads (Sønderholm et al., 2017). In

comparison, bacteria from the anoxically conditioned batch cultures were fully deprived of O_2. In

the present study, the effect of anoxic conditioning was restricted to an intermediate period. This is

probably due to the static methodological setup. Dynamic processes, such as entry of nutrients and

removal of waste, occur during infection and are not modeled here (Brown et al., 2008).

Nevertheless, this is not the first time a "resuscitation window" has been described for non-

culturable bacteria. Similar studies have shown that VBNC bacteria can only be detected in an

intermediate period of culturing (Pinto et al., 2015).

ROS restricts growth of anoxically conditioned *P. aeruginosa*

The DTC fraction of bacteria in this study was O_2 intolerant given that growth only was achievable

when NO_3^- served as an alternative electron acceptor during anaerobic respiration. This led us to

investigate whether this phenomena was due to oxidizing properties of ROS created by incomplete

reduction of O_2 during aerobic respiration (Fenchel and Finlay, 2008). Sodium pyruvate increased

counts for anoxically conditioned *P. aeruginosa* during normoxic plating (roughly 43 % of the DTC

population) and we believe that it was due to its properties as a ROS scavenger (Mizunoe et al.,

1999). Accordingly, 0.3 % sodium pyruvate has been used to resuscitate non-culturable populations

of *S. aureus* (Pasquaroli et al., 2013). Catalase also increased counts during normoxic plating

(roughly 50 % of the DTC population), indicating that hydrogen peroxide is involved in the

induction of a DTC state. The effect of ROS was further confirmed by the reduced growth of a

catalase A deficient *P. aeruginosa* mutant ($\Delta katA$ PAO1) after anoxic conditioning. Finally,

measurements of ROS along with OD showed that anoxically conditioned batch cultures of *P.*

aeruginosa created more ROS when regrown under atmospheric conditions the longer they were

deprived from oxygen, resulting in an accumulation of ROS and a reduction in growth represented

by lower OD measurements. These results indicate that accumulation of ROS during aerobic

respiration has an impact on normoxic plate counting when *P. aeruginosa* has been anoxically

conditioned. Furthermore, it seems that hydrogen peroxide is involved in this loss of culturing. It

has been suggested that VBNC cells cannot be resuscitated by addition of ROS scavengers and that

"revived" cells in the presence of ROS scavengers are only injured cells and not VBNC cells (Pinto

et al., 2015). It was not possible to determine whether DTC bacteria in this study were in an injured

state.

S. epidermidis **and** *S. aureus* **also become DTC after anoxic conditioning**

Additional experiments were conducted to elucidate whether anoxic conditioning generated DTC

sub-populations in other facultative pathogens. Filter biofilms of methicillin susceptible *S. aureus*

showed the same effects of anoxic conditioning as *P. aeruginosa*. These experiments were also

repeated for *S. epidermidis*, *S. aureus* (MRSA), *E. coli* and *E. faecalis*. Both *S. epidermidis* and *S.*

aureus were significantly (p = 0.01 and p = 0.03, respectively) affected by anoxic conditioning,

whereas *E. coli* and *E. faecalis* were not. The results in this study points toward an intermediate

period where anoxic conditioned bacteria are benefitted by anoxic incubation with nitrate as

alternative electron acceptor. We cannot rule out that we missed the period where *E. coli* and *E.*

faecalis were DTC, given that they were tested at selected days. No effect of sodium pyruvate was

found during normoxic plating for any of these tested organisms, suggesting that ROS is not

involved in the loss of culturability for these pathogens, but additional ROS scavengers should be

tested.

Conclusion

A non-culturable state can be induced by several physiological stresses and our knowledge in this

area is expanding, though far from fully resolved. In this study, we demonstrate that anoxic

conditioning generates a VBNC subpopulation in *P. aeruginosa* batch-cultures. Furthermore, we

demonstrate that anoxic conditioning generates DTC sub-populations of *P. aeruginosa*, *S. aureus*

and *S. epidermidis* during biofilm growth. The DTC population was only able to grow under anoxic

conditions in the presence of NO_3^- as an alternative electron acceptor. In the case of *P. aeruginosa,*

this phenomenon was explained by creation of lethal amounts of ROS during aerobic respiration

and the bacteria's inability to neutralize it.

Data availability

The data generated in the current study are available from the corresponding author on reasonable

request.

Funding information

This study was supported by grants from The Lundbeck foundation (R105-A9791) to TB and by a

Novo-Nordisk Tandem (ref.nr. NNF16OC0023482) grant to MK. The funders had no role in study

design, data collection and interpretation, or the decision to submit the work for publication

Author contributions

LK performed the majority of the experiments. MK, MS, BFG, SC and KNK also performed

experiments. TB, PØJ, KNK AK, and LK conceived and designed experiments. LK wrote the

manuscript. MA and PØJ performed statistics. All authors analyzed data. All authors contributed to

and corrected the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

Ayrapetyan, M., Williams, T.C., Baxter, R., and Oliver, J.D. (2015). Viable but Nonculturable and Persister Cells Coexist Stochastically and Are Induced by Human Serum. *Infect Immun* 83**,** 4194-4203.

Binnerup, S.J., And J. Sorensen. (1993). Long-term oxidant deficiency in Pseudomonas aeruginosa PAO303 results in cells which are non-culturable under aerobic conditions. *FEMS Microbiol. Ecol.* 13**,** 79-84.

Boulos, L., Prevost, M., Barbeau, B., Coallier, J., and Desjardins, R. (1999). LIVE/DEAD BacLight : application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water. *J Microbiol Methods* 37**,** 77-86.

Brown, S.A., Palmer, K.L., and Whiteley, M. (2008). Revisiting the host as a growth medium. *Nat Rev Microbiol* 6**,** 657-666.

Costerton, J.W., Post, J.C., Ehrlich, G.D., Hu, F.Z., Kreft, R., Nistico, L., Kathju, S., Stoodley, P., Hall-Stoodley, L., Maale, G., James, G., Sotereanos, N., and Demeo, P. (2011). New methods for the detection of orthopedic and other biofilm infections. *FEMS Immunol Med Microbiol* 61**,** 133-140.

Costerton, W., Veeh, R., Shirtliff, M., Pasmore, M., Post, C., and Ehrlich, G. (2003). The application of biofilm science to the study and control of chronic bacterial infections. *J Clin Invest* 112**,** 1466-1477.

Fenchel, T., and Finlay, B. (2008). Oxygen and the spatial structure of microbial communities. *Biol Rev Camb Philos Soc* 83**,** 553-569.

Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles, K.W. (2013). A genetic resource for rapid and comprehensive phenotype screening of nonessential Staphylococcus aureus genes. *MBio* 4**,** e00537-00512.

Fisher, R.A., Gollan, B., and Helaine, S. (2017). Persistent bacterial infections and persister cells.

Nat Rev Microbiol 15**,** 453-464.

Frees, D., Chastanet, A., Qazi, S., Sørensen, K., Hill, P., Msadek, T., and Ingmer, H. (2004). Clp

ATPases are required for stress tolerance, intracellular replication and biofilm formation in

Staphylococcus aureus. *Mol Microbiol* 54**,** 1445-1462.

Hassett, D.J., Cuppoletti, J., Trapnell, B., Lymar, S.V., Rowe, J.J., Yoon, S.S., Hilliard, G.M.,

Parvatiyar, K., Kamani, M.C., Wozniak, D.J., Hwang, S.H., Mcdermott, T.R., and Ochsner,

U.A. (2002). Anaerobic metabolism and quorum sensing by Pseudomonas aeruginosa

biofilms in chronically infected cystic fibrosis airways: rethinking antibiotic treatment

strategies and drug targets. *Adv Drug Deliv Rev* 54**,** 1425-1443.

Hassett, D.J., Ma, J.F., Elkins, J.G., Mcdermott, T.R., Ochsner, U.A., West, S.E., Huang, C.T.,

Fredericks, J., Burnett, S., Stewart, P.S., Mcfeters, G., Passador, L., and Iglewski, B.H.

(1999). Quorum sensing in Pseudomonas aeruginosa controls expression of catalase and

superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. *Mol*

Microbiol 34**,** 1082-1093.

Høiby, N., Bjarnsholt, T., Givskov, M., Molin, S., and Ciofu, O. (2010). Antibiotic resistance of

bacterial biofilms. *Int J Antimicrob Agents* 35**,** 322-332.

Høiby, N., Bjarnsholt, T., Moser, C., Bassi, G.L., Coenye, T., Donelli, G., Hall-Stoodley, L., Hola,

V., Imbert, C., Kirketerp-Møller, K., Lebeaux, D., Oliver, A., Ullmann, A.J., and Williams,

C. (2015). ESCMID guideline for the diagnosis and treatment of biofilm infections 2014.

Clin Microbiol Infect 21 Suppl 1**,** S1-25.

James, G.A., Ge Zhao, A., Usui, M., Underwood, R.A., Nguyen, H., Beyenal, H., Delancey Pulcini,

E., Agostinho Hunt, A., Bernstein, H.C., Fleckman, P., Olerud, J., Williamson, K.S.,

Franklin, M.J., and Stewart, P.S. (2016). Microsensor and transcriptomic signatures of

oxygen depletion in biofilms associated with chronic wounds. *Wound Repair Regen* 24**,**

373-383.

Jensen, P.Ø., Briales, A., Brochmann, R.P., Wang, H., Kragh, K.N., Kolpen, M., Hempel, C.,

Bjarnsholt, T., Høiby, N., and Ciofu, O. (2014). Formation of hydroxyl radicals contributes

to the bactericidal activity of ciprofloxacin against Pseudomonas aeruginosa biofilms.

Pathog Dis 70**,** 440-443.

Jensen, P.Ø., Kolpen, M., Kragh, K.N., and Kühl, M. (2017). Microenvironmental characteristics

and physiology of biofilms in chronic infections of CF patients are strongly affected by the

host immune response. *Apmis* 125**,** 276-288.

Kim, J.S., Chowdhury, N., Yamasaki, R., and Wood, T.K. (2018). Viable but non-culturable and

persistence describe the same bacterial stress state. *Environ Microbiol*.

Kolpen, M., Hansen, C.R., Bjarnsholt, T., Moser, C., Christensen, L.D., Van Gennip, M., Ciofu, O.,

Mandsberg, L., Kharazmi, A., Doring, G., Givskov, M., Høiby, N., and Jensen, P.Ø. (2010).

Polymorphonuclear leucocytes consume oxygen in sputum from chronic *Pseudomonas*

aeruginosa pneumonia in cystic fibrosis. *Thorax* 65**,** 57-62.

Kolpen, M., Kühl, M., Bjarnsholt, T., Moser, C., Hansen, C.R., Liengaard, L., Kharazmi, A.,

Pressler, T., Høiby, N., and Jensen, P.Ø. (2014). Nitrous oxide production in sputum from

cystic fibrosis patients with chronic Pseudomonas aeruginosa lung infection. *PLoS One* 9**,**

e84353.

Kolpen, M., Mousavi, N., Sams, T., Bjarnsholt, T., Ciofu, O., Moser, C., Kuhl, M., Hoiby, N., and

Jensen, P.O. (2016). Reinforcement of the bactericidal effect of ciprofloxacin on

Pseudomonas aeruginosa biofilm by hyperbaric oxygen treatment. *Int J Antimicrob Agents*

47**,** 163-167.

Kragh, K.N., Alhede, M., Jensen, P.Ø., Moser, C., Scheike, T., Jacobsen, C.S., Seier Poulsen, S.,

Eickhardt-Sørensen, S.R., Trostrup, H., Christoffersen, L., Hougen, H.P., Rickelt, L.F.,

Kühl, M., Høiby, N., and Bjarnsholt, T. (2014). Polymorphonuclear leukocytes restrict

growth of Pseudomonas aeruginosa in the lungs of cystic fibrosis patients. *Infect Immun* 82**,**

4477-4486.

Li, L., Mendis, N., Trigui, H., Oliver, J.D., and Faucher, S.P. (2014). The importance of the viable

but non-culturable state in human bacterial pathogens. *Front Microbiol* 5**,** 258.

Mizunoe, Y., Wai, S.N., Takade, A., and Yoshida, S. (1999). Restoration of culturability of

starvation-stressed and low-temperature-stressed Escherichia coli O157 cells by using

H2O2-degrading compounds. *Arch Microbiol* 172**,** 63-67.

Noor, R. (2015). Mechanism to control the cell lysis and the cell survival strategy in stationary

phase under heat stress. *Springerplus* 4**,** 599.

Oliver, J.D. (2010). Recent findings on the viable but nonculturable state in pathogenic bacteria.

FEMS Microbiol Rev 34**,** 415-425.

Pasquaroli, S., Zandri, G., Vignaroli, C., Vuotto, C., Donelli, G., and Biavasco, F. (2013).

Antibiotic pressure can induce the viable but non-culturable state in Staphylococcus aureus

growing in biofilms. *J Antimicrob Chemother* 68**,** 1812-1817.

Pinto, D., Santos, M.A., and Chambel, L. (2015). Thirty years of viable but nonculturable state

research: unsolved molecular mechanisms. *Crit Rev Microbiol* 41**,** 61-76.

Sønderholm, M., Koren, K., Wangpraseurt, D., Jensen, P.Ø., Kolpen, M., Kragh, K.N., Bjarnsholt,

T., and Kühl, M. (2018). Tools for studying growth patterns and chemical dynamics of

aggregated Pseudomonas aeruginosa exposed to different electron acceptors in an alginate

bead model. *NPJ Biofilms Microbiomes* 4**,** 3.

Sønderholm, M., Kragh, K.N., Koren, K., Jakobsen, T.H., Darch, S.E., Alhede, M., Jensen, P.Ø.,

Whiteley, M., Kühl, M., and Bjarnsholt, T. (2017). Pseudomonas aeruginosa Aggregate

Formation in an Alginate Bead Model System Exhibits In Vivo-Like Characteristics. *Appl*

Environ Microbiol 83.

Trunk, K., Benkert, B., Quack, N., Munch, R., Scheer, M., Garbe, J., Jansch, L., Trost, M.,

Wehland, J., Buer, J., Jahn, M., Schobert, M., and Jahn, D. (2010). Anaerobic adaptation in

Pseudomonas aeruginosa: definition of the Anr and Dnr regulons. *Environ Microbiol* 12,

1719-1733.

Walters, M.C., 3rd, Roe, F., Bugnicourt, A., Franklin, M.J., and Stewart, P.S. (2003). Contributions

of antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of

Pseudomonas aeruginosa biofilms to ciprofloxacin and tobramycin. *Antimicrob Agents*

Chemother 47, 317-323.

Welch, R.A., Burland, V., Plunkett, G., 3rd, Redford, P., Roesch, P., Rasko, D., Buckles, E.L.,

Liou, S.R., Boutin, A., Hackett, J., Stroud, D., Mayhew, G.F., Rose, D.J., Zhou, S.,

Schwartz, D.C., Perna, N.T., Mobley, H.L., Donnenberg, M.S., and Blattner, F.R. (2002).

Extensive mosaic structure revealed by the complete genome sequence of uropathogenic

Escherichia coli. *Proc Natl Acad Sci U S A* 99, 17020-17024.

Wood, T.K., Knabel, S.J., and Kwan, B.W. (2013). Bacterial persister cell formation and dormancy.

Appl Environ Microbiol 79, 7116-7121.

Worlitzsch, D., Tarran, R., Ulrich, M., Schwab, U., Cekici, A., Meyer, K.C., Birrer, P., Bellon, G.,

Berger, J., Weiss, T., Botzenhart, K., Yankaskas, J.R., Randell, S., Boucher, R.C., and

Doring, G. (2002). Effects of reduced mucus oxygen concentration in airway Pseudomonas

infections of cystic fibrosis patients. *J Clin Invest* 109, 317-325.

Figure legends

Fig. 1 – Direct viable counting reveals a viable but non-culturable population of *Pseudomonas*
***aeruginosa* (PAO1) after anoxic conditioning**

Bacterial counts per milliliter were determined with plate counting and direct viable counting from

24-hour-old batch cultures (A) and from 16-day-old anoxic conditioned batch cultures (B) of PAO1.

Bacteria were stained with LIVE/DEAD staining to estimate the proportion of viable ("live") and

non-viable ("dead") cells. There was significantly more viable bacterial counts when applying

direct viable counting in comparison to normoxic plating ($p < 0.0001$, one-way ANOVA test) and

anoxic plating ($p = 0.0009$, one-way ANOVA). Anoxic plating yielded significantly higher bacterial

counts ($p = 0.023$, one-way ANOVA test) than normoxic plating. Symbols with error bars indicate

the mean $\pm$ SEM (n = 3). + NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates.

Fig. 2 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was

generated in the bead biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 21 days from the

beads (A) and the surrounding suspension (B). Symbols with error bars indicate the mean $\pm$ SEM (n

= 4). + NO$_3^-$ refers to the addition of 10 mM KNO$_3$ to LB agar plates. The log difference (C)

represents the difference in mean log CFU/mL between plating methods from the suspension and

the beads, respectively. The log difference was significantly higher (p = 0.003, linear regression) in

the beads than the surrounding suspension. Symbols with error bars indicate the mean + confidence

intervals.

Fig. 3 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was

generated in the filter biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 17 days from

anoxically (A) and normoxically (B) conditioned filters. Symbols with error bars indicate the mean

$\pm$ SEM (n = 4). + NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates. The log difference

(C) represents the difference in mean log CFU/mL between plating methods from the normoxically

and anoxically conditioned filters, respectively. The log difference was significantly higher (p =

0.01, linear regression) in the anoxically conditioned filters than the normoxically conditioned.

Symbols with error bars indicate the mean + confidence intervals.

Fig. 4 – An oxygen intolerant subpopulation of Pseudomonas aeruginosa (PAO1) was

generated in the planktonic batch cultures

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 28 days from

anoxically (A) and normoxically (B) conditioned batch cultures. Symbols with error bars indicate

the mean $\pm$ SEM (n = 4). $\pm$ NO$_3^-$ refers to the addition of 10 mM KNO$_3$ to LB agar plates. The log

difference (C) represents the difference in mean log CFU/mL between plating methods from the

normoxically and anoxically conditioned batch cultures, respectively. The log difference was

significantly higher (p < 0.0001, linear regression) in the anoxically conditioned batch cultures than

the normoxically conditioned. Symbols with error bars indicate the mean + confidence intervals.

Fig. 5 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was generated in colonies

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 20 days from anoxically (A) and normoxically (B) conditioned colonies. Symbols with error bars indicate the mean $\pm$ SEM (n = 3). +NO$_3^-$ refers to the addition of 10 mM KNO$_3$ to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL between plating methods from the normoxically and anoxically conditioned colonies, respectively. The log difference was significantly higher (p < 0.0001, linear regression) in the anoxically conditioned colonies than the normoxically conditioned. Symbols with error bars indicate the mean + confidence intervals.

Fig. 6 – Reactive oxygen species generates a DTC subpopulation of *Pseudomonas aeruginosa*

(PAO1)

Normoxic and anoxic colony forming units per milliliter (CFU/mL) were determined for 24-hour-

old batch cultures (A) and 16-day-old anoxically conditioned batch cultures (B) of PAO1.

Normoxic CFU/mL was determined ± 0.3% sodium pyruvate (A and B), while addition of catalase

was only tested for 16-day-old batch cultures. Symbols with error bars indicate the mean ± SEM (n

= 4). +NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates. There was a significant

difference between normoxic plating ± 0.3 % sodium pyruvate (p = 0.04, one-way ANOVA test)

and normoxic plating ± catalase (p = 0.03, one-way ANOVA test). Normoxic and anoxic

determination of CFU/mL were determined for 16-day-old anoxically conditioned batch cultures of

PAO1 and *ΔkatA* PAO1 (C). Symbols with error bars indicate the mean ± SEM (n = 3). Significant

difference between anoxic and normoxic plating with both PAO1 and *ΔkatA* PAO1 (p = 0.0014 and

p < 0.0001, respectively, one-way ANOVA test).

Fig. 7 - Increase in reactive oxygen species affects growth of anoxically conditioned

Pseudomonas aeruginosa **(PAO1) over time**

Determination of reactive oxygen species (ROS) and optical density (OD) at day 1 (A), 8 (B) and

16 (C) for anoxically ($-O_2$) and normoxically ($+O_2$) conditioned PAO1. ROS was measured as

counts per second (CPS) when 2′,7′dichlorodihydrofluorescein diacetate was converted to 2′7′-

dichlorofluorescein. Growth was measured with optical density (OD_{600}) simultaneously to ROS

measurement. Data is generated from 99 continuously measurements presented as a solid line with

dots representing ± SEM (n = 4).

Fig. 8 - An oxygen intolerant subpopulation of *Staphylococcus aureus* (methicillin susceptible)

was generated in the filter biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of *Staphylococcus aureus* over

17 days from anoxically (A) and normoxically (B) conditioned filters. Symbols with error bars

indicate the mean $\pm$ SEM ($n = 4$). $+NO_3^-$ refers to the addition of 10 mM KNO_3 to LB agar plates.

The log difference (C) represents the difference in mean log CFU/mL between plating methods

from the normoxically and anoxically conditioned filters, respectively. The log difference was

significantly higher ($p < 0.001$, linear regression) in the anoxically conditioned filters than the

normoxically conditioned. Symbols with error bars indicate the mean + confidence intervals.