Oxygen restriction generates difficult-to-culture pathogens

Running title: Oxygen restriction generates DTC pathogens

Authors: Lasse Kvich [b], Blaine G. Fritz [b] Stephanie Crone [b], Kasper N. Kragh [b], Mette Kolpen [a], Majken Sønderholm [b],

Mikael Andersson [c], Anders Koch [c,d], Peter Ø. Jensen [a,b*], Thomas Bjarnsholt [a,b*#]

* Corresponding authors

Author affiliation:

[a] Department of Clinical Microbiology, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark.

[b] Costerton Biofilm Center, Department of Immunology and Microbiology, Faculty of Health and Medical Sciences,

University of Copenhagen, Denmark.

[c] Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark.

[d] Department of Infectious Diseases, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark.

ORCID: LK 0000-0003-3226-3633, BF 0000-0002-2085-894X, KK 0000-0002-7662-9118, MK 0000-0001-5575-0384, MS 0000-0001-9576-5775, AK 0000-0001-9205-1048, PJ 0000-0003-1648-7186, TB 0000-0002-8003-7414

Corresponding authors:

Professor Thomas Bjarnsholt

Costerton Biofilm Center

Department of Immunology and Microbiology

University of Copenhagen

Blegdamsvej 3b, Room 22.2.22

DK-2200 Copenhagen N

Denmark.

Tlf: +45 2065 9888

tbjarnsholt@sund.ku.dk

Associated Professor Peter Ø. Jensen

Costerton Biofilm Center

Department of Immunology and Microbiology

University of Copenhagen

Blegdamsvej 3b, Room 24.1.14

DK-2200 Copenhagen N

Denmark.

Tlf: +45 3031 9073

peter.oestrup.jensen@regionh.dk

Abstract: 245 words

Document: 5958 words

Abstract

Induction of a non-culturable state has been demonstrated for many bacteria. In a clinical perspective, the lack of growth due to these non-culturable bacteria can have major consequences for the diagnosis and treatment of patients. Here we show how anoxic conditioning (restriction of molecular oxygen, O_2) generates difficult-to-culture (DTC) bacteria during biofilm growth. A significant subpopulation of *Pseudomonas aeruginosa* entered a DTC state after anoxic conditioning, ranging from five to 90 % of the total culturable population, in both planktonic and biofilm models. Anoxic conditioning also generated DTC subpopulations of *Staphylococcus aureus* and *Staphylococcus epidermidis*. Growth of the DTC populations were achieved by substituting O_2 with 10 mM NO_3^- as an alternative electron acceptor for anaerobic respiration or, in the case of *P. aeruginosa*, by adding sodium pyruvate or catalase as scavengers against reactive oxygen species (ROS) during aerobic respiration. An increase in normoxic plating due to addition of catalase suggests the molecule hydrogen peroxide as a possible mechanism for induction of DTC *P. aeruginosa.* Anoxic conditioning also generated a true viable but non-culturable (VBNC) population of *P. aeruginosa* that was not resurrected by substituting O_2 with NO_3^- during anaerobic respiration. Moreover, bacterial detection in clinical samples was improved significantly by supplementing 10 mM NO_3^- to LB plates and incubating under anoxic conditions. These results demonstrate that habituation to an infectious anoxic micro-environment complicates diagnostic culturing of bacteria, especially in the case of chronic infections where oxygen is restricted due to the host immune response.

Keywords:

anoxia / **biofilm** / *P. aeruginosa* / **reactive oxygen species** / **difficult-to-culture**

Importance

Diagnostics of bacteria from chronic infections by standard culture-based methods is challenging. Bacteria in a non-culturable state may contribute to the lack of culturing from these infections. Many stressors are known to induce a non-culturable state, among others the absence of molecular oxygen, which is evident in chronic infections due to high rates of oxygen consumption by the host response. In this study, we have shown that *Pseudomonas aeruginosa, Staphylococcus aureus and Staphylococcus epidermidis* can enter a difficult-to-culture state after oxygen restriction. Regrowth was not possible using conventional normoxic plating where oxygen served as electron acceptor. Instead, regrowth was enabled during anoxic conditions with added nitrate as alternative electron acceptor. Our findings show that bacteria can habituate to their environment and that it has to be taken into consideration especially when culturing clinical samples e.g. from chronic infections.

Introduction

Clinical laboratories often culture bacteria with enriched media, developed to support growth of particular pathogens. However, these human bacterial pathogens can enter a growth-restricted, transient state resulting in loss of culturability, especially following antibiotic treatment (1). Non-culturable bacteria are apparent by microscopic identification in clinical cases presenting sign of infection, but no positive cultures. These cases are especially prevalent during chronic infection, where lack of culturability prevents proper diagnosis and treatment (2-4).

Two well defined, yet difficult to distinguish, non-culturable states have been described for bacteria, viable but non-culturable (VBNC) and persisters (5-8). Several shared environmental stresses has been described to induce these non-culturable states, e.g. oxidative stress, change in temperature or pH, nutrient starvation, change in osmotic concentrations and presence of heavy metals or antibiotics (6, 9). Oxygen starvation, referred to as anoxic conditioning in this paper, have also shown to induce a non-culturable state in batch cultures of *Pseudomonas aeruginosa*, but supplementation with nitrate (NO_3^-) as an alternative electron acceptor during anoxic plating restored culturability (10). The lack of growth during normoxic plating suggests that anoxic conditioning sensitizes a subpopulation of *P. aeruginosa* to O_2 or its toxic derivatives, such as reactive oxygen species (ROS). ROS are continuously created during aerobic respiration by incomplete reduction of O_2, which is toxic to cells if not scavenged (11). Presence of ROS has, in some studies, been shown to generate VBNC bacteria (6, 12), but most research has focused on *Escherichia coli* strains or *Vibrio* spp., while the effects of ROS on *P. aeruginosa* are not well characterized.

Evidence of anoxic zones and anaerobic bacterial activity in chronic infections suggests that colonizing bacteria, such as *P. aeruginosa,* experience anoxic conditioning (13-15). High rates of O_2 consumption by polymorphonuclear leukocytes generate local, anoxic zones and likely play a

major role in O_2 depletion during chronic infection (4). In addition, other cells of the host may also

consume O_2, increasing the likelihood of bacteria experiencing anoxia (13). Many chronic

infections are thought to contain bacteria in the biofilm mode of growth (16) and endogenous O_2

depletion inside the biofilm (17), along with intense O_2 consumption by the host immune response

(13, 18-21), may therefore increase the number of VBNC bacteria.

This study aimed to determine whether anoxic conditioning generates VBNC cells of *P. aeruginosa*

during biofilm growth. Furthermore, we investigated whether ROS are involved in the loss of

culturability. Finally, we evaluated the efficiency of two novel, anoxic growth media to grow

bacteria from clinical samples with suspected biofilm infection.

Results

Direct viable counts with LIVE/DEAD staining revealed a VBNC *P. aeruginosa* population.

Direct viable counts were carried out for 16-day-old, anoxically conditioned batch cultures of *P. aeruginosa* to investigate if anoxic conditioning could generate a VBNC population. Findings were compared to normoxic and anoxic plating (plating methods) performed on LB plates supplemented with 10 mM NO_3^- (Fig. 1B). Anoxic plating yielded a significantly (p = 0.002) higher count of colony forming units per milliliter (CFU/mL) than normoxic plating (0.85 log values ± 0.04 SD). When determining bacterial plate counts with direct viable counting, a significant higher number of viable cells could be calculated compared to normoxic (2.31 log values ± 0.47 SD, p = 0.002) and anoxic plating (1.46 log values ± 0.5 SD, p = 0.008). The cells represented by this difference were considered VBNC. Direct viable counts were also performed on 24-hour-old batch cultures to investigate whether findings of VBNC *P. aeruginosa* was restricted to anoxically conditioned cells. No difference in bacterial counts was observed (Fig. 1A).

Anoxic conditioning generated oxygen intolerant subpopulations of *P. aeruginosa* in biofilm and planktonic models.

To examine if biofilm growth affected the subsequent normoxic and anoxic plating, *P. aeruginosa* was grown in an alginate-bead biofilm model (22). Plate counts were performed on LB plates supplemented with 10 mM NO_3^-. Plates were incubated under normoxic and anoxic conditions and plate counts were compared from the beads (biofilm) and the surrounding media (planktonic) of the same vials over a period of 21 days (Fig. 2A and 2B, respectively). The log difference was significantly (p = 0.003) higher for biofilm than planktonic cells (Fig. 2C). The cells represented by this difference were considered to be in a difficult-to-culture (DTC) state and ranged from five to 54 % of the entire population from day six to 19 (Fig. S1).

P. aeruginosa was also grown anoxically or normoxically on a filter biofilm model using LB plates

supplemented with 10 mM NO_3^- over a period of 17 days. CFU/mL was then determined using

anoxic and normoxic plating, as above (Fig. 3A and 3B, respectively). The log difference between

plating methods was significantly (p = 0.01) higher for anoxically conditioned biofilms than

normoxically conditioned biofilms (Fig. 3C). The fraction of DTC *P. aeruginosa* ranged from six to

23 % from day three to 17 (Fig. S1).

Similar results were obtained for anoxically and normoxically conditioned planktonic batch cultures

of *P. aeruginosa* over a period of 28 days (Fig. 4A and 4B, respectively). The log difference

between plating methods was significantly (p < 0.0001) higher for anoxically conditioned batch

cultures (Fig. 3C). The fraction of DTC bacteria ranged from 60 to 90 % from day nine to 21 (Fig.

S1).

The effect of anoxic conditioning was further investigated on colonies of *P. aeruginosa* grown on

LB plates supplemented with 10 mM NO_3^- under anoxic and normoxic conditions over a period of

20 days (Fig. 5A and 5B, respectively). The log difference between plating methods was

significantly higher (p < 0.0001) for anoxically conditioned colonies (Fig. 5C). The fraction of DTC

P. aeruginosa ranged from 12 to 46 % from day six to 20 (Fig. S1).

10 mM NO_3^- in LB plates did not affect CFU/mL in batch cultures of *P. aeruginosa*.

To ensure that presence of NO_3^- did not affect the number of CFU generated on LB plates, CFU/mL

was determined anoxically (LB plates + 10 mM NO_3^-) and normoxically (LB plates ± 10 mM NO_3^-)

from 24-hour-old batch cultures of *P. aeruginosa*. No difference was observed between type of

plating (p = 0.93): anoxic log CFU/mL + NO_3^- = 9.96 (± 0.22 SD), normoxic log CFU/mL + NO_3^- =

9.91 (± 0.10 SD) and normoxic log CFU/mL - NO_3^- = 9.91 (± 0.14 SD). Furthermore, there was no

effect of NO_3^- supplementation on normoxic plating in the prolonged experiment with 28-day-old

batch cultures (Fig. 4A and 4B). It was not possible to detect growth of *P. aeruginosa* on LB plates under anoxic conditions without NO_3^-, but growth was observed when these plates were placed under normoxic conditions.

Oxidative stress restricted the growth of anoxically conditioned *P. aeruginosa* when re-grown in a normoxic environment.

Bacteria from 16-day-old anoxically and normoxically conditioned batch cultures of *P. aeruginosa* were plated on 10 mM NO_3^- LB plates ± sodium pyruvate and ± catalase to investigate whether formation of ROS contributed to the DTC state induced by anoxic conditioning. Presence of sodium pyruvate as a ROS scavenger (23) significantly ($p = 0.04$) increased normoxic CFU/mL compared to plating without sodium pyruvate (Fig. 6B). This was not the case in 24-hour-old batch cultures of *P. aeruginosa* ($p = 0.62$), indicating that the effect was restricted to anoxically conditioned cells (Fig. 6A). Presence of catalase also significantly ($p = 0.03$) improved normoxic plate counts compared to normoxic plating without catalase (Fig. 6B). These results show that hydrogen peroxide is involved in the induction of a DTC state created by anoxic conditioning. To add prove to this hypothesis, a similar experiment was carried out with a catalase deficient *P. aeruginosa* mutant (Δ*katA* PAO1), which is more susceptible to oxidative stress (24, 25). The log difference between plating methods for the Δ*katA* mutant and reference strain was significantly ($p = 0.00002$, unpaired t-test) different. These log difference values were $3.02 ± 0.05$ SD and $1.61 ± 0.07$ SD for Δ*katA* PAO1 and the reference strain, respectively (Fig. 6C). Moreover, measurement of ROS and optical density (OD) was carried out for *P. aeruginosa* at day 1, 8 and 16 from anoxically and normoxically conditioned batch cultures. These results showed creation of ROS during normoxic growth. The longer PAO1 was anoxically conditioned; the lower was the following normoxic

growth, as measured by OD. This effect was less pronounced for normoxically-conditioned cells

(Fig.7 A-C).

Anoxic conditioning affects *Staphylococcus aureus* and *Staphylococcus epidermidis* but not

Escherichia coli* and *Enterococcus faecalis

The effect of anoxic conditioning was then tested on a selected group of pathogens to determine if

this phenomenon was restricted to *P. aeruginosa*. *S. aureus* (methicillin susceptible) was tested as

described in the *P. aeruginosa* filter biofilm setup. Plate counts for anoxically and normoxically

conditioned filters was determined (Fig. 8A and 8B, respectively). The log difference between

plating method was significantly higher ($p < 0.001$) when plate counts were performed from

anoxically conditioned filter biofilms than from normoxically conditioned filter biofilms (Fig. 8C).

The fraction of DTC *S. aureus* ranged from three to 89 % from day one to day 17 (Fig. S1). Since

both *P. aeruginosa* and *S. aureus* demonstrated effects of anoxic conditioning, a smaller experiment

was initiated to test other pathogens. The effect was determined on LB plates supplemented with 10

mM NO_3^- ± sodium pyruvate to investigate if ROS were involved in the lack of growth. Both *S.*

aureus (MRSA) and *S. epidermidis* showed effects of anoxic conditioning, resulting in an

significant ($p = 0.01$ and $p = 0.03$, respectively) increase in CFU/mL during anoxic plating

compared to normoxic plating (Fig. S2). There was no observed effect of sodium pyruvate during

normoxic plating for these two strains. In the case of *E. coli* and *E. faecalis,* there was no observed

effect of anoxic conditioning (Fig. S3).

Implementation of a supplementary diagnostic media increased findings from samples where

biofilm infections could be expected.

To investigate the effect of a DTC state on the outcome of culturing from patient samples, LB and

ABTG plates supplemented with 10 mM NO_3^- were introduced as a supplementary anoxic growth

media in addition to standard culturing practices at the Department of Clinical Microbiology,

Rigshospitalet, Denmark. A total of 135 samples were consecutively collected from 73 patients.

Only samples from sterile parts of the body, e.g. soft tissues and bones, were included. Figure S4 A

and B show the distribution of bacterial findings from the conventional method compared with LB

and ABTG plates supplemented with 10 mM NO_3^-. The conventional method used several growth

media, but it was not possible to separate findings of species according to growth media. When

comparing LB plates supplemented with 10 mM NO_3^- and the conventional method, an equal

number of culture positive samples (45 vs. 50) were identified, resulting in 27 and 28 patients with

culture positive findings, respectively (Fig. S4 B). Interestingly, 19 microorganisms from 16

patients were found exclusively on LB plates supplemented with 10 mM NO_3^- and in 5 cases (6.9

%), only NO_3^- supplemented LB plates were culture positive (Table 1). Addition of LB plates

supplemented with NO_3^- to the conventional method resulted in changed microbiological findings

for 21.9 % of the patients (p = 0.00003). The clinical significance of these observations was not

possible to investigate due to Danish legislation regarding access to patient files. The use of ABTG

plates supplemented with 10 mM NO_3^- did not provide significant benefit for growing

microorganisms from clinical samples (Fig. S4 A).

Discussion

DTC and VBNC *P. aeruginosa*

In the current study, we demonstrate that plate counting with *P. aeruginosa* is significantly affected

by anoxic conditioning when cultured in the presence of atmospheric oxygen levels. A DTC state

was created after only a few days of anoxic conditioning in both biofilm and planktonic models.

DTC *P. aeruginosa* was observed in the beads from the bead biofilm model, but not in the

surrounding suspension, despite that the cultures had full access to atmospheric oxygen. This

indicates that oxygen restriction within a biofilm generates a DTC subpopulation. The DTC state,

detected as increased growth during anoxic plating with NO_3^-, has previously been generated in

planktonic *P. aeruginosa* by energy starvation after cultivation without O_2 as electron acceptor for

aerobic respiration (10). *In vitro* biofilms of *P. aeruginosa* may contain internal anoxic zones (26,

27), so we hypothesized that anoxia inside the biofilm contributes to the induction of DTC *P.*

aeruginosa. Accordingly, DTC *P. aeruginosa* was only observed in the filter biofilm model and the

colony model when the model was anoxically conditioned. Interestingly, we found that 99.68 % of

the population was VBNC with the LIVE/DEAD assay after anoxic conditioning. LIVE/DEAD

staining is based upon membrane permeability and is only an approximation of true viability. Cells

that were stained both red and green (resulting in yellow) were counted as dead cells but may still

be viable.

Approximately 50 % of the population in the alginate beads was DTC even though they were kept

in normoxically-conditioned vials. In comparison, the fraction of DTC cells was approximately 90

% in the anoxically conditioned batch cultures. The difference (90 % vs. 50 %) may be explained

by the fact that a majority of the bacteria in the beads are peripherally located where they have

increased access to oxygen compared to the center of the beads (22). In comparison, bacteria from

the anoxically conditioned batch cultures were fully deprived of O_2. In the present study, the effect

of anoxic conditioning was restricted to an intermediate period. This is probably due to the static

methodological setup. Dynamic processes, such as entry of nutrients and removal of waste, occur

during infection and are not modeled here (28). Nevertheless, this is not the first time a

"resuscitation window" has been described for non-culturable bacteria. Similar studies have shown

that VBNC bacteria can only be detected in an intermediate period of culturing (29).

ROS restricts growth of anoxically conditioned *P. aeruginosa*

The DTC fraction of bacteria in this study was O_2 intolerant given that growth only was achievable

when NO_3^- served as an alternative electron acceptor during anaerobic respiration. This led us to

investigate whether this phenomena was due to oxidizing properties of ROS created by incomplete

reduction of O_2 during aerobic respiration (11). Sodium pyruvate increased counts for anoxically

conditioned *P. aeruginosa* during normoxic plating (roughly 43 % of the DTC population) and we

believe that it was due to its properties as a ROS scavenger (23). Accordingly, 0.3 % sodium

pyruvate has been used to resuscitate non-culturable populations of *S. aureus* (1). Catalase also

increased counts during normoxic plating (roughly 50 % of the DTC population), indicating that

hydrogen peroxide is involved in the induction of a DTC state. The effect of ROS was further

confirmed by the reduced growth of a catalase A deficient *P. aeruginosa* mutant ($\Delta katA$ PAO1)

after anoxic conditioning. Finally, measurements of ROS along with OD showed that anoxically

conditioned batch cultures of *P. aeruginosa* created more ROS when regrown under atmospheric

conditions the longer they were deprived from oxygen, resulting in an accumulation of ROS and a

reduction in growth represented by lower OD measurements. These results indicate that

accumulation of ROS during aerobic respiration has an impact on normoxic plate counting when *P.*

aeruginosa has been anoxically conditioned. Furthermore, it seems that hydrogen peroxide is

involved in this loss of culturing. It has been suggested that VBNC cells cannot be resuscitated by

addition of ROS scavengers and that "revived" cells in the presence of ROS scavengers are only

injured cells and not VBNC cells (29). It was not possible to determine whether DTC bacteria in

this study were in an injured state.

***S. epidermidis* and *S. aureus* also become VBNC after anoxic conditioning**

Additional experiments were conducted to elucidate whether anoxic conditioning generated DTC

sub-populations in other facultative pathogens. Filter biofilms of methicillin susceptible *S. aureus*

showed the same effects of anoxic conditioning as *P. aeruginosa*. These experiments were also

repeated for *S. epidermidis*, *S. aureus* (MRSA), *E. coli* and *E. faecalis*. Both *S. epidermidis* and *S.*

aureus were significantly (p = 0.01 and p = 0.03, respectively) affected by anoxic conditioning,

whereas *E. coli* and *E. faecalis* were not. No effect of sodium pyruvate was found during normoxic

plating for any of these tested organisms, suggesting that ROS is not involved in the loss of

culturability for these pathogens, but additional ROS scavengers should be tested.

Efficacy of LB plates supplemented with NO_3^- in a clinical setting

LB plates supplemented with NO_3^- were tested in a clinical setting and enabled detection of bacteria

in almost as many patients as the conventional method (27 vs. 28, respectively). Findings of

microorganisms (n=19 in 16 patients) exclusively detected on LB plates supplemented with NO_3^-

were primarily "low-virulent" bacteria. These samples often also contained "highly virulent"

bacteria. "Highly virulent" bacteria grew equally well on both LB plates supplemented with NO_3^-

and with the conventional method. With the exception of *Helcococcus kunzii*, all identified bacteria

detected exclusively by LB plates supplemented with NO_3^- were known to harbor nitrate reductase,

enabling reduction of NO_3^-. Furthermore, according to the literature, all identified bacteria are able

to produce biofilm and have been associated with chronic infections (Table 1). LB plates

supplemented with NO_3^- detected growth in 6.9 % of patients (mostly Staphylococcal species),

whereas the conventional method did not. These organisms are often associated with chronic

infections, although they are considered "low virulent" (30). Using LB plates with 10 Mm NO_3^- as a

supplementary growth media to the conventional method changed the microbiological findings for

16 patients (p = 0.00003). These findings indicate that anoxic LB agar plates supplemented with

NO_3^- may be useful as a supplementary medium for the cultivation of "low-virulent" bacteria from

chronic infections.

Conclusion

A non-culturable state can be induced by several physiological stresses and our knowledge in this area is expanding, though far from fully resolved. In this study, we demonstrate that anoxic conditioning generates a VBNC subpopulation in *P. aeruginosa* batch-cultures. Furthermore, we demonstrate that anoxic conditioning generates DTC sub-populations of *P. aeruginosa*, *S. aureus* and *S. epidermidis* during biofilm growth. The DTC population was only able to grow under anoxic conditions in the presence of NO_3^- as an alternative electron acceptor. In the case of *P. aeruginosa,* this phenomenon was explained by creation of lethal amounts of ROS during aerobic respiration and the bacteria's inability to neutralize it. Finally, LB plates supplemented with 10 mM NO_3^- were an effective, anoxic growth medium for resuscitation of anoxically conditioned bacteria and may be useful as a supplemental growth media for "low-virulent" anoxic conditioned bacteria in a clinical setting.

Materials and Method

Bacterial strains

Pseudomonas aeruginosa (PAO1, ATCC 15692), a catalase A deficient *Pseudomonas aeruginosa* strain (Δ*katA* PAO1) (25), *Staphylococcus epidermidis* ATCC 14990, *Staphylococcus aureus* NCTC 8325-4 (methicillin susceptible) (31), *Staphylococcus aureus* USA300 JE2 (MRSA) (32), E*scherichia coli* CFT073 (33) and a clinical strain of *Enterococcus faecalis* from the Department of Clinical Microbiology, Copenhagen University Hospital – Rigshospitalet, Denmark were used in this study.

Agar plates and media

Primarily lysogeny broth (LB) agar plates were applied in this study. Lysogeny broth (pH 7.5) consisted of 5 g/L yeast extract (Oxoid, Roskilde, Denmark), 10 g/L tryptone (Oxoid), 10 g/L NaCl

(Merck, USA). Both LB and minimum media plates were tested as supplementary anoxic growth

media for clinical samples. Minimum media consisted of Btrace media buffered with 10% A-10

phosphate buffer and supplemented with 0.5 % (w/v) glucose (Panum Institute Substrate

Department, DK), referred to as ABTG media throughout the paper. All plates contained 2 %

agarose. Plates used for anoxic growth were supplemented with 10 mM KNO_3 (Sigma-Aldrich,

USA) to serve as alternative electron acceptor during anoxic plating, referred to as NO_3^- throughout

the paper. All agar plates and media in this study were supplied by the Panum Institute Substrate

Department (Copenhagen, DK).

Anoxic growth

Experiments investigating growth under anoxic conditions were performed in an anaerobic chamber

(Concept 400 Anaerobic Workstation, Ruskinn Technology Ltd, UK). The gas atmosphere

consisted of $N_2/H_2/CO_2$ (ratio - 80:10:10). Anoxic chamber environment was confirmed with an

oxygen sensor (HQ40d multi, HACH Company, USA). All media and chemical solutions used in

anaerobic experiments were equilibrated in the anaerobic chamber 3 days prior to experiment. In

the case of solutions requiring refrigeration, a minimal volume was applied (< 1 mL), sealed with

Parafilm M, and thoroughly shaken upon entry into anaerobic chamber for quick gas equilibration.

Direct viable count with LIVE/DEAD staining

LIVE/DEAD staining was applied to estimate the proportion of viable and non-viable cells in 24-

hour-old batch cultures and in 16-day-old anoxically conditioned batch cultures of *P. aeruginosa*.

The dyes consisted of two fluorescent nucleic acid stains, the green fluorescent stain (live cells)

SYTO9 (Invitrogen, USA) and the red fluorescent stain (dead cells) propidium iodide (PI, Sigma-

Aldrich, USA). SYTO9 penetrates both intact and damaged membranes while PI only stains

damaged cells, thereby creating an opportunity to discriminate between live and dead cells (34).

Bacterial suspensions were vortexed thoroughly (1 min) and sonicated (5 min degas + 5 minutes

sonication) before staining. To stain the cells, one µL of PI and SYTO9 was added to 1 mL of

bacterial suspension and incubated for 15 minutes at room temperature. Suspensions were then

filtered through a 0.2 µm black Whatman, Nuclepore Trach-Etch Membrane (Sigma-Aldrich,

Denmark) and visualized with confocal laser scanning microscopy using a Zeiss LSM 710 with a

63×/1.4 (numerical aperture) objective (Zeiss, Germany). Fifteen random fields (135µm x 135 µm)

were examined for each filter. Enumeration of live (green) and dead (red) bacteria were done with

the IMARIS software package (Bitplane AG, Schwitzerland). Cells that were stained with both PI

and SYTO9 were considered non-viable and thus counted as dead cells. Normoxic and anoxic

CFU/mL was carried out simultaneously to estimate the proportion of VBNC cells. CFU/mL was

determined in the same way as described for liquid batch cultures. Conversion of enumerated viable

and dead cells to bacterial counts per milliliter was performed as described (35) to compare them

with CFU/mL.

Bead-embedded inoculum of *P. aeruginosa*

Preparation of alginate beads with *P. aeruginosa* was carried out according to a method described

by Sønderholm *et al*. (22). Subsequently, beads were divided (10 beads per vial) into vials (Oximate

Vial, PerkinElmer Inc., USA) containing 15 mL LB medium. Vials were sealed with Parafilm M

and incubated at 37°C on an orbital shaker at 180 rpm. Anoxic and normoxic CFU counts were

determined from beads and from the surrounding suspension from the same vials on each sampling

day (day 3, 6, 8, 12, 19 and 21). Two beads were sampled per biological replicate (4 biological

replicates in total). Before determination of CFU, the beads were washed twice with 0.9 % NaCl to

remove non-attached cells and transferred to 1.5 mL microcentrifuge tubes (Sigma-Aldrich, USA).

One-hundred microliters of 0.1 M sodium carbonate (Na_2CO_3) followed by 100 µL of 0.04 M citric

acid was added to the tubes to dissolve the beads. The suspensions were then sonicated (5 min

degas + 5 minutes sonication; Bransonic ultrasonic cleaner 2510, Emerson Electric, USA) before

367   ten-fold dilution series were performed in 0.9 % NaCl. CFU was determined by plating three, 10

368   µL-drops per dilution per replicate. Anoxic CFU determinations were performed inside an

369   anaerobic chamber on LB plates supplemented with 10 mM NO_3^-. The same dilution series were

370   applied to normoxic CFU determination outside the anaerobic chamber on LB plates with 10 mM

371   NO_3^-. Plates were incubated 2 days before counting CFU.

Filter biofilms with *P. aeruginosa*

This protocol was adapted to grow reproducible biofilms under anoxic and normoxic conditions.

The method has previously been described by Bjarnsholt *et al.* (2015). The filter biofilms were kept

on the same LB plates throughout the experiment. *P. aeruginosa* was propagated from frozen stock

and grown overnight in 20 mL LB medium at 37°C on an orbital shaker at180 rpm. Cultures were

adjusted to an optical density of 0.05 (OD_{600}; UV spectrophotometer UV-1800 UV-VIS, Shimadzu

corporation, JP) and 10 µL was transferred to the cellulose nitrate membrane filters (25 mm in

diameter, GE Healthcare Life Sciences, UK). Plates were incubated under normoxic and anoxic

conditions and kept in plastic bags with wet paper to avoid dehydration. Two filters were sampled

per biological replicate (4 biological replicates in total) on each sampling day (day 1, 3, 7, 15 and

17). Filters were removed, placed in 10 mL tubes containing 5 mL 0.9 % NaCl, vortexed thoroughly

(one min.) and sonicated (5 min degas + 5 minutes sonication). CFU/mL was determined as

previously described.

Liquid batch cultures of *P. aeruginosa*

P. aeruginosa was propagated from frozen stock and grown overnight in 20 mL LB media at 37^{o}C

and orbitally shaken at 180 rpm. Cultures were adjusted to an optical density (OD_{600}) of 0.1 in glass

vials (Oximate Vial) with a final volume of 20 mL LB. Vials were left to incubate at 37°C and

orbitally shaken at 180 rpm. To create a normoxic environment, half of the vials were incubated

with Parafilm M on top (normoxic conditioning), while the rest were incubated with a lid on top

creating an anoxic environment (anoxic conditioning). Anoxic and normoxic plating was carried out

from normoxic and anoxic conditioned liquid batch cultures (referred to as batch cultures

throughout the paper) on each sampling day (day 1, 3, 5, 9, 11, 14, 16, 18, 21 and 28). Normoxic

plating was carried out on LB plates with and without 10 mM NO_3^- to test whether presence of NO_3^-

affected the number of CFU. Two mL (2 x 1 mL = 2 technical replicates) were sampled per

biological replicate (4 biological replicates in total) on each sampling day. CFU/mL was determined

as previously described.

Colonies of *P. aeruginosa*

P. aeruginosa was propagated from frozen stock and grown overnight in 20 mL LB media at 37°C

on an orbital shaker at 180 rpm. The cultures were then streaked onto LB plates supplemented with

10 mM NO_3^-. Plates were incubated under normoxic and anoxic conditions and kept in plastic bags

with wet paper to avoid dehydration. A one-μl loop was used to sample colony material from each

biological replicate (3 biological replicates in total) on each sampling day (day 6, 9, 13, 15 and 20).

Colonies were transferred to 1.5mL microcentrifuge tubes (Sigma-Aldrich, Denmark) with 0.5 mL

0.9 % NaCl, vortexed thoroughly (1 min) and sonicated (5 min degas + 5 minutes sonication).

CFU/mL was determined as previously described.

Reactive oxygen species (ROS)

To elucidate whether growth of anoxically conditioned bacteria was restricted by creation of ROS

during aerobic respiration, sodium pyruvate and catalase (Sigma-Aldrich, USA) was tested as ROS

scavenger in LB plates with 10 mM NO_3^-. Sodium pyruvate (0.3 %) was tested during anoxic and

normoxic plating for 24-hour-old batch cultures and anoxically conditioned 16-day-old batch

cultures of *P. aeruginosa*. Catalase (~50,000 units of catalase per L medium) was tested during

anoxic and normoxic plating for anoxically conditioned 16-day-old batch cultures of *P. aeruginosa*.

Furthermore, CFU/mL was also determined from 16-day-old anoxically conditioned batch cultures

of *P. aeruginosa* and a catalase A deficient *P. aeruginosa* (Δ*katA* PAO1) to test the influence of

ROS. Lastly, creation of ROS was measured along with optical density (OD) at day 1, 8 and 16 for

normoxically and anoxically conditioned batch cultures of *P. aeruginosa*. Creation of ROS was

measured by adding 5 µM 2′,7′dichlorodihydrofluorescein diacetate (DCHF-DA; Sigma) to the

bacterial suspension. This concentration has previously been used to detect ROS for *P. aeruginosa*

(24). Two hundred microliter of anoxically conditioned cultures (OD_{600} adjusted to 0.1 in fresh LB

medium) was added to each well of black 96-well microtiter plates with transparent flat bottom

(16503, Thermo Fisher Scientific, Rochester, NY). OD and ROS was measured over 15 hours at

27°C with a VICTOR Multilabel Plate Reader (Perkin Elmer, MA). All experiments consisted of 3-

4 biological replicates.

Filter biofilms with other pathogens

To determine if other bacteria behaved in the same way as *P. aeruginosa*, we investigated the effect

of anoxic conditioning on a selected group of pathogens (*Staphylococcus epidermidis*,

Staphylococcus aureus (methicillin susceptible), *Staphylococcus aureus* USA300 JE2 (MRSA),

E*scherichia coli* CFT073 and a clinical strain of *Enterococcus faecalis*). The filter biofilm method

was used as described earlier. For the methicillin susceptible strain, CFU was determined in the

same way as described for the filters with *P. aeruginosa*. For the remaining strains, CFU was

determined at day 1 and 9 on LB plates supplemented with 10 mM NO_3^- ± O_2. Furthermore, CFU

was determined on LB plates supplemented with 10 mM NO_3^- and 0.3 % sodium pyruvate to

evaluate the effect of ROS. A minimum of three biological replicates was performed and CFU was

carried out in the same way as described earlier.

Implementation of supplementary plates

LB and ABTG plates, supplemented with 10 mM NO_3^-, were tested as a supplementary anoxic

growth media at the Department of Clinical Microbiology, Rigshospitalet, Denmark. A total of 135

plates of each were applied to test if NO_3^- would improve microbiological findings from clinical

samples where potential biofilm infections could be suspected. The plates were implemented in the

daily routine on tissue and bone samples from sterile sites of the body. Plates were incubated up to

7 days in an anaerobic chamber. Microbiological findings were performed by inspection of plates at

day 3, 5 and 7, and confirmation of species was performed with Matrix-assisted laser desorption

ionization time-of-flight mass spectrometry (MALDI-TOF MS) on a Microflex LT instrument

(Bruker Daltonik GmbH, Germany). Protein profile was obtained by the software FlexControl 3.3

(Bruker Daltonik GmbH), analyzed with FlexAnalysis 3.3 (Bruker Daltonik GmbH). The database

used to match spectra was Bruker Taxonomy (7311 MSPs). Results were estimated per patient and

compared with findings from conventional methods applied at the Department of Clinical

Microbiology, Rigshospitalet. The conventional method consisted of determination of species from

solid agar plates under normoxic and anoxic incubation. Normoxic incubation was performed on

solid agar plates ['Blue plates', a modified Conradi-Drigalski medium containing 10 g/L detergent,

1 g/L $Na_2S_2O_3 \cdot H_2O$, 0.1 g/L bromothymol blue, 9 g/L lactose, 0.4 g/L glucose, pH 8.0], [Blood agar

plates, 5% horse blood, pH 7.4] and in serum bouillons [Basal culture medium supplemented with

defibrinated horse blood and horse serum]. Anoxic incubation was performed on solid agar plates

['Chocolate plates', a modified Reyn and Bentzon medium containing defibrinated horse blood,

ascitic fluid in a broth-agar base consisting of 2.4 % of Danish AKI agar in beefheart broth with 1

% of peptone ("Orthana" special), 0.3% of NaCl and 0.2% of $Na_2HPO_4 \cdot 12H_2O$] (36), ['anaerobic

plates', prepared as chocolate medium and supplemented with vitamin K and cysteine] and in

thioglycollate broths [anaerobic culture medium for sterility]. All media applied at the clinic were

produced and delivered by Statens Serum Institut (Copenhagen, Denmark). Microbiological

findings by the conventional method were also confirmed with MALDI-TOF MS.

Ethics

A comparison of culture-methods for quality assurance is according to Danish legislation not a health research project, as defined by the 'Danish Act on Research Ethics Review of Health Research Projects'. Hence, comparison of culture methods on clinical samples could be initiated without approval from The Committees on Health Research Ethics in the Capital Region of Denmark.

Statistics

To evaluate the difference in bacterial growth between incubation conditions (normoxic and anoxic conditioning), a linear regression was used with the difference between the logarithmically transformed values for normoxic and anoxic plating as outcome and with day (categorical) and interaction between day and plating condition (binary) as explanatory variables in SAS Genmod Procedure. The p-value of the interaction term was used as the p-value for the difference in bacterial growth. Log difference was calculated as ($\log_{10}$[CFU/mL]$_{anoxic}$ - $\log_{10}$[CFU/mL]$_{normoxic}$). The mean and standard error of the mean (SEM) were calculated for recovering bacteria and plotted using GraphPad Prism 6.1 (GraphPad Software, La Jolla, USA). From the ratio between anoxic (CFU -O_2) and normoxic (CFU +O_2) colony counts it was possible to calculate the fraction of DTC bacteria [% DTC bacteria $= \frac{(CFU\ -O_2) - (CFU\ +O_2)}{(CFU\ -O_2)}$ x 100]. Data that were not part of the long-term experiments were instead analyzed with a 1-way ANOVA followed by Tukey's multiple comparison tests, Fisher's exact test or an unpaired t-test. McNemar's exact test was used to test how many patients would get a change in culture response if LB plates with NO_3^- were used as a supplementary growth media to the conventional method applied at Rigshospitalet. A p-value $\leq$ 0.05 was considered statistically significant. The tests were performed with either Prism 6.1 (GraphPad Software, La Jolla, USA) or SAS v.9.4 (SAS Institute Inc., Cary, NC, USA).

Data availability

The data generated in the current study are available from the corresponding author on reasonable

request.

Funding information

This study was supported by grants from The Lundbeck foundation to TB and by a UC-CARE

(University of Copenhagen–Center for Antimicrobial Research) grant to MK. The funders had no

role in study design, data collection and interpretation, or the decision to submit the work for

publication

Acknowledgments

We would like to thank the laboratory technicians at the Department of Clinical Microbiology,

Rigshospitalet, for helping us with the collection of samples.

Author contributions

 LK performed the majority of the experiments. MK, MS, BFG, SC and KNK also performed

experiments. TB, PØJ, KNK AK, and LK conceived and designed experiments. LK wrote the

manuscript. MA and PØJ performed statistics. All authors analyzed data. All authors contributed to

and corrected the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. Pasquaroli S, Zandri G, Vignaroli C, Vuotto C, Donelli G, Biavasco F. 2013. Antibiotic pressure can induce the viable but non-culturable state in *Staphylococcus aureus* growing in biofilms. J Antimicrob Chemother 68:1812-7.

2. Høiby N, Bjarnsholt T, Givskov M, Molin S, Ciofu O. 2010. Antibiotic resistance of bacterial biofilms. Int J Antimicrob Agents 35:322-32.

3. Costerton JW, Post JC, Ehrlich GD, Hu FZ, Kreft R, Nistico L, Kathju S, Stoodley P, Hall-Stoodley L, Maale G, James G, Sotereanos N, DeMeo P. 2011. New methods for the detection of orthopedic and other biofilm infections. FEMS Immunol Med Microbiol 61:133-40.

4. Høiby N, Bjarnsholt T, Moser C, Bassi GL, Coenye T, Donelli G, Hall-Stoodley L, Hola V, Imbert C, Kirketerp-Møller K, Lebeaux D, Oliver A, Ullmann AJ, Williams C. 2015. ESCMID guideline for the diagnosis and treatment of biofilm infections 2014. Clin Microbiol Infect 21 Suppl 1:S1-25.

5. Fisher RA, Gollan B, Helaine S. 2017. Persistent bacterial infections and persister cells. Nat Rev Microbiol 15:453-464.

6. Oliver JD. 2010. Recent findings on the viable but nonculturable state in pathogenic bacteria. FEMS Microbiol Rev 34:415-25.

7. Kim JS, Chowdhury N, Yamasaki R, Wood TK. 2018. Viable but non-culturable and persistence describe the same bacterial stress state. Environ Microbiol doi:10.1111/1462-2920.14075.

8. Wood TK, Knabel SJ, Kwan BW. 2013. Bacterial persister cell formation and dormancy. Appl Environ Microbiol 79:7116-21.

9.   Ayrapetyan M, Williams TC, Baxter R, Oliver JD. 2015. Viable but Nonculturable and

Persister Cells Coexist Stochastically and Are Induced by Human Serum. Infect Immun

83:4194-203.

10.   Binnerup SJ, and J. Sørensen. 1993. Long-term oxidant deficiency in *Pseudomonas*

aeruginosa PAO303 results in cells which are non-culturable under aerobic conditions.

FEMS Microbiol Ecol 13:79-84.

11.   Fenchel T, Finlay B. 2008. Oxygen and the spatial structure of microbial communities. Biol

Rev Camb Philos Soc 83:553-69.

12.   Noor R. 2015. Mechanism to control the cell lysis and the cell survival strategy in stationary

phase under heat stress. Springerplus 4:599.

13.   Worlitzsch D, Tarran R, Ulrich M, Schwab U, Cekici A, Meyer KC, Birrer P, Bellon G,

Berger J, Weiss T, Botzenhart K, Yankaskas JR, Randell S, Boucher RC, Doring G. 2002.

Effects of reduced mucus oxygen concentration in airway Pseudomonas infections of cystic

fibrosis patients. J Clin Invest 109:317-25.

14.   Hassett DJ, Cuppoletti J, Trapnell B, Lymar SV, Rowe JJ, Yoon SS, Hilliard GM,

Parvatiyar K, Kamani MC, Wozniak DJ, Hwang SH, McDermott TR, Ochsner UA. 2002.

Anaerobic metabolism and quorum sensing by *Pseudomonas aeruginosa* biofilms in

chronically infected cystic fibrosis airways: rethinking antibiotic treatment strategies and

drug targets. Adv Drug Deliv Rev 54:1425-43.

15.   Kolpen M, Kühl M, Bjarnsholt T, Moser C, Hansen CR, Liengaard L, Kharazmi A, Pressler

T, Høiby N, Jensen PØ. 2014. Nitrous oxide production in sputum from cystic fibrosis

patients with chronic *Pseudomonas aeruginosa* lung infection. PLoS One 9:e84353.

16.   Costerton W, Veeh R, Shirtliff M, Pasmore M, Post C, Ehrlich G. 2003. The application of

556   biofilm science to the study and control of chronic bacterial infections. J Clin Invest

557   112:1466-77.

558   17.   Sønderholm M, Koren K, Wangpraseurt D, Jensen PØ, Kolpen M, Kragh KN, Bjarnsholt T,

559   Kühl M. 2018. Tools for studying growth patterns and chemical dynamics of aggregated

560   *Pseudomonas aeruginosa* exposed to different electron acceptors in an alginate bead model.

561   NPJ Biofilms Microbiomes 4:3.

562   18.   Trunk K, Benkert B, Quack N, Munch R, Scheer M, Garbe J, Jansch L, Trost M, Wehland J,

563   Buer J, Jahn M, Schobert M, Jahn D. 2010. Anaerobic adaptation in *Pseudomonas*

564   *aeruginosa*: definition of the Anr and Dnr regulons. Environ Microbiol 12:1719-33.

565   19.   James GA, Ge Zhao A, Usui M, Underwood RA, Nguyen H, Beyenal H, deLancey Pulcini

566   E, Agostinho Hunt A, Bernstein HC, Fleckman P, Olerud J, Williamson KS, Franklin MJ,

567   Stewart PS. 2016. Microsensor and transcriptomic signatures of oxygen depletion in

568   biofilms associated with chronic wounds. Wound Repair Regen 24:373-83.

569   20.   Jensen PØ, Kolpen M, Kragh KN, Kühl M. 2017. Microenvironmental characteristics and

570   physiology of biofilms in chronic infections of CF patients are strongly affected by the host

571   immune response. Apmis 125:276-288.

572   21.   Kragh KN, Alhede M, Jensen PØ, Moser C, Scheike T, Jacobsen CS, Seier Poulsen S,

573   Eickhardt-Sørensen SR, Trostrup H, Christoffersen L, Hougen HP, Rickelt LF, Kühl M,

574   Høiby N, Bjarnsholt T. 2014. Polymorphonuclear leukocytes restrict growth of

575   *Pseudomonas aeruginosa* in the lungs of cystic fibrosis patients. Infect Immun 82:4477-86.

576   22.   Sønderholm M, Kragh KN, Koren K, Jakobsen TH, Darch SE, Alhede M, Jensen PØ,

577   Whiteley M, Kühl M, Bjarnsholt T. 2017. *Pseudomonas aeruginosa* Aggregate Formation in

an Alginate Bead Model System Exhibits In Vivo-Like Characteristics. Appl Environ

Microbiol 83.

23.   Mizunoe Y, Wai SN, Takade A, Yoshida S. 1999. Restoration of culturability of starvation-

stressed and low-temperature-stressed *Escherichia coli* O157 cells by using H2O2-

degrading compounds. Arch Microbiol 172:63-7.

24.   Jensen PØ, Briales A, Brochmann RP, Wang H, Kragh KN, Kolpen M, Hempel C,

Bjarnsholt T, Høiby N, Ciofu O. 2014. Formation of hydroxyl radicals contributes to the

bactericidal activity of ciprofloxacin against *Pseudomonas aeruginosa* biofilms. Pathog Dis

70:440-3.

25.   Hassett DJ, Ma JF, Elkins JG, McDermott TR, Ochsner UA, West SE, Huang CT,

Fredericks J, Burnett S, Stewart PS, McFeters G, Passador L, Iglewski BH. 1999. Quorum

sensing in *Pseudomonas aeruginosa* controls expression of catalase and superoxide

dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. Mol Microbiol

34:1082-93.

26.   Kolpen M, Mousavi N, Sams T, Bjarnsholt T, Ciofu O, Moser C, Kühl M, Høiby N, Jensen

PØ. 2016. Reinforcement of the bactericidal effect of ciprofloxacin on *Pseudomonas*

aeruginosa biofilm by hyperbaric oxygen treatment. Int J Antimicrob Agents 47:163-7.

27.   Walters MC, 3rd, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. 2003. Contributions of

antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of

Pseudomonas aeruginosa biofilms to ciprofloxacin and tobramycin. Antimicrob Agents

Chemother 47:317-23.

28.   Brown SA, Palmer KL, Whiteley M. 2008. Revisiting the host as a growth medium. Nat

Rev Microbiol 6:657-66.

29.   Pinto D, Santos MA, Chambel L. 2015. Thirty years of viable but nonculturable state

research: unsolved molecular mechanisms. Crit Rev Microbiol 41:61-76.

30.   Rogers KL, Fey PD, Rupp ME. 2009. Coagulase-negative staphylococcal infections. Infect

Dis Clin North Am 23:73-98.

31.   Frees D, Chastanet A, Qazi S, Sørensen K, Hill P, Msadek T, Ingmer H. 2004. Clp ATPases

are required for stress tolerance, intracellular replication and biofilm formation in

Staphylococcus aureus. Mol Microbiol 54:1445-62.

32.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A

genetic resource for rapid and comprehensive phenotype screening of nonessential

Staphylococcus aureus genes. MBio 4:e00537-12.

33.   Welch RA, Burland V, Plunkett G, 3rd, Redford P, Roesch P, Rasko D, Buckles EL, Liou

SR, Boutin A, Hackett J, Stroud D, Mayhew GF, Rose DJ, Zhou S, Schwartz DC, Perna NT,

Mobley HL, Donnenberg MS, Blattner FR. 2002. Extensive mosaic structure revealed by

the complete genome sequence of uropathogenic *Escherichia coli*. Proc Natl Acad Sci U S

A 99:17020-4.

34.   Li L, Mendis N, Trigui H, Oliver JD, Faucher SP. 2014. The importance of the viable but

non-culturable state in human bacterial pathogens. Front Microbiol 5:258.

35.   Boulos L, Prevost M, Barbeau B, Coallier J, Desjardins R. 1999. LIVE/DEAD BacLight :

application of a new rapid staining method for direct enumeration of viable and total

bacteria in drinking water. J Microbiol Methods 37:77-86.

36.   Møller V, Reyn A. 1965. A new solid medium for the isolation of *Neisseria Gonorrhoeae*.

Bull World Health Organ 32:471-6.

37.   Lentino JR. 2004. Infections Associated with Prosthetic Knee and Prosthetic Hip. Curr

Infect Dis Rep 6:388-392.

38.   Müller D, Gabriel J. 1999. Bacterial degradation of the herbicide bromoxynil by

Agrobacterium radiobacter in biofilm. Folia Microbiol (Praha) 44:377-9.

39.   Rodby RA, Glick EJ. 1991. *Agrobacterium radiobacter* peritonitis in two patients

maintained on chronic peritoneal dialysis. Am J Kidney Dis 18:402-5.

40.   Stanger KM, Albert F, Kneser U, Bogdan C, Horch R. 2013. Management of chronic

osteomyelitis of the tibia with life-threatening complications under negative pressure wound

therapy and isolation of *Helcococcus kunzii*. Int Wound J doi:10.1111/iwj.12133.

41.   Perez-Jorge C, Cordero J, Marin M, Esteban J. 2012. Prosthetic joint infection caused by

Helcococcus kunzii. J Clin Microbiol 50:528-30.

42.   Cui B, Smooker PM, Rouch DA, Deighton MA. 2015. Effects of erythromycin on the

phenotypic and genotypic biofilm expression in two clinical *Staphylococcus capitis*

subspecies and a functional analysis of Ica proteins in *S. capitis*. J Med Microbiol

doi:10.1099/jmm.0.000059.

43.   Frid P, Tornes K, Nielsen O, Skaug N. 2009. Primary chronic osteomyelitis of the jaw--a

microbial investigation using cultivation and DNA analysis: a pilot study. Oral Surg Oral

Med Oral Pathol Oral Radiol Endod 107:641-7.

44.   Achermann Y, Goldstein EJ, Coenye T, Shirtliff ME. 2014. *Propionibacterium acnes*: from

commensal to opportunistic biofilm-associated implant pathogen. Clin Microbiol Rev

27:419-40.

45.   Donelli G, Vuotto C, Cardines R, Mastrantonio P. 2012. Biofilm-growing intestinal

anaerobic bacteria. FEMS Immunol Med Microbiol 65:318-25.

46.   Seng P, Bayle S, Alliez A, Romain F, Casanova D, Stein A. 2015. The microbial

epidemiology of breast implant infections in a regional referral centre for plastic and

reconstructive surgery in the south of France. Int J Infect Dis 35:62-66.

47.   Al-Shuneigat J, Cox SD, Markham JL. 2005. Effects of a topical essential oil-containing

650   formulation on biofilm-forming coagulase-negative staphylococci. Lett Appl Microbiol

651   41:52-5.

652   48.   Announ N, Mattei JP, Jaoua S, Fenollar F, Sati H, Chagnaud C, Roudier J, Guis S. 2004.

653   Multifocal discitis caused by *Staphylococcus warneri*. Joint Bone Spine 71:240-2.

654   49.   Weckwerth PH, de Mattias Franco AT, de Magalhaes Lopes CA, Santos FD, Weckwerth

655   AC, Vivan RR, Duarte MA. 2014. Bacterial pathogens related to chronic suppurative otitis

656   media in individuals with cleft palate: bacteriological culture and polymerase chain reaction.

657   Cleft Palate Craniofac J 51:145-53.

658   50.   Fukuda TT, Shoji; Kota, Kohichi; Iwaku, Masaaki; Hoshino, Etsuro. 2004. Aerotolerance of

659   *Peptostreptococcus anaerobius* biofilms. Microbial Ecology in Health & Disease Vol. 16

660   p205.

661   51.   Ambalam P, Kondepudi KK, Nilsson I, Wadstrom T, Ljungh A. 2012. Bile stimulates cell

662   surface hydrophobicity, Congo red binding and biofilm formation of Lactobacillus strains.

663   FEMS Microbiol Lett 333:10-9.

Table

Table 1 – Bacterial finding in a clinical setting on LB plates supplemented with 10 mM KNO$_3^-$

Bacteria exclusively detected on lysogeny broth (LB) agar plates supplemented with 10 mM KNO$_3^-$ (n = 19 in 16 patients). In five cases (five patients), growth was only detected with LB + NO$_3^-$. Bacteria marked with * were not possible to identify with MALDI-TOF MS to the genus or species level. NA means not applicable, this is due to the failure of proper identification. Ability to reduce nitrate was available from the Kyoto Encyclopedia of Genes and Genomes (KEGG) database under the nitrogen metabolism - reference pathway (http://www.genome.jp/kegg-bin/show_pathway?map=map00910&show_description=show). +/- refers to the ability of the organism to perform or not perform either nitrate reductase, biofilm or be associated with chronic infections.

Bacteria found exclusively on LB supplemented with NO$_3^-$.	Numbers detected	Growth only detected with LB + NO$_3^-$.	Nitrate reductase	Biofilm formation	Possible infection (chronic)	Literature
Staphylococcus epidermidis	4	2	+	+	+	(30, 37)
Rhizobacterium (Agrobacterium) radiobacter	1	-	+	+	+	(38, 39)
Helcococcus kunzii	1	-	-	+	+	(40, 41)
Staphylococcus capitis	1	-	+	+	+	(42, 43)
Propionebacterium acnes	2	-	+	+	+	(43, 44)
Finegoldia magna	4	-	+	+	+	(45, 46)
Staphylococcus warneri	2	2	+	+	+	(47, 48)
Peptostreptococcus anaerobius	1	-	+	+	+	(49, 50)
Lactobacillus rhamnosus	1	-	+	+	+	(51)
Propionebacterium sp. *	1	-	NA	NA	NA	-
Cocci *	1	1	NA	NA	NA	-

Figures

Fig. 1 – Direct viable counting reveals a viable but non-culturable population of *Pseudomonas aeruginosa* (PAO1) after anoxic conditioning

Bacterial counts per milliliter were determined with plate counting and direct viable counting from 24-hour-old batch cultures (A) and from 16-day-old anoxic conditioned batch cultures (B) of PAO1. Bacteria were stained with LIVE/DEAD staining to estimate the proportion of viable ("live") and non-viable ("dead") cells. There was significantly more viable bacterial counts (p = 0.049, one-way ANOVA test) when applying direct viable counting in comparison to normoxic plating. Anoxic plating yielded significantly higher bacterial counts (p = 0.001, one-way ANOVA test) than normoxic plating. Symbols with error bars indicate the mean ± SEM (n = 3). + NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates.

Fig. 2 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was generated in the bead

biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 21 days from the beads (A) and the

surrounding suspension (B). Symbols with error bars indicate the mean ± SEM (n = 4). + NO_3^- refers to the addition of

10 mM KNO_3 to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL between plating

methods from the suspension and the beads, respectively. The log difference was significantly higher (p = 0.003, linear

regression) in the beads than the surrounding suspension. Symbols with error bars indicate the mean + confidence

intervals.

Fig. 3 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was generated in the filter

biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 17 days from anoxically (A) and

normoxically (B) conditioned filters. Symbols with error bars indicate the mean $\pm$ SEM (n = 4). + NO_3^- refers to the

addition of 10 mM KNO_3 to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL

between plating methods from the normoxically and anoxically conditioned filters, respectively. The log difference was

significantly higher (p = 0.01, linear regression) in the anoxically conditioned filters than the normoxically conditioned.

Symbols with error bars indicate the mean + confidence intervals.

Fig. 4 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was generated in the planktonic

batch cultures

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 28 days from anoxically (A) and

normoxically (B) conditioned batch cultures. Symbols with error bars indicate the mean ± SEM (n = 4). ± NO_3^- refers to

the addition of 10 mM KNO_3 to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL

between plating methods from the normoxically and anoxically conditioned batch cultures, respectively. The log

difference was significantly higher (p < 0.0001, linear regression) in the anoxically conditioned batch cultures than the

normoxically conditioned. Symbols with error bars indicate the mean + confidence intervals.

Fig. 5 – An oxygen intolerant subpopulation of *Pseudomonas aeruginosa* (PAO1) was generated in colonies

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of PAO1 over 20 days from anoxically (A) and

normoxically (B) conditioned colonies. Symbols with error bars indicate the mean $\pm$ SEM (n = 3). $+NO_3^-$ refers to the

addition of 10 mM KNO_3 to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL

between plating methods from the normoxically and anoxically conditioned colonies, respectively. The log difference

was significantly higher (p < 0.0001, linear regression) in the anoxically conditioned colonies than the normoxically

conditioned. Symbols with error bars indicate the mean + confidence intervals.

Fig. 6 – Reactive oxygen species generates a DTC subpopulation of *Pseudomonas aeruginosa* (PAO1)

Normoxic and anoxic colony forming units per milliliter (CFU/mL) were determined for 24-hour-old batch cultures (A) and 16-day-old anoxically conditioned batch cultures (B) of PAO1. Normoxic CFU/mL was determined ± 0.3% sodium pyruvate (A and B), while addition of catalase was only tested for 16-day-old batch cultures. Symbols with error bars indicate the mean ± SEM (n = 4). $+NO_3^-$ refers to the addition of 10 mM KNO_3 to LB agar plates. There was a significant difference between normoxic plating ± 0.3 % sodium pyruvate (p = 0.04, one-way ANOVA test) and normoxic plating ± catalase (p = 0.03, one-way ANOVA test). Normoxic and anoxic determination of CFU/mL were determined for 16-day-old anoxically conditioned batch cultures of PAO1 and *ΔkatA* PAO1 (C). Symbols with error bars indicate the mean ± SEM (n = 3). Significant difference between anoxic and normoxic plating with both PAO1 and *ΔkatA* PAO1 (p = 0.0014 and p < 0.0001, respectively, one-way ANOVA test).

Fig. 7 - Increase in reactive oxygen species affects growth of anoxically conditioned *Pseudomonas aeruginosa*

(PAO1) over time

Determination of reactive oxygen species (ROS) and optical density (OD) at day 1 (A), 8 (B) and 16 (C) for anoxically

($-O_2$) and normoxically ($+O_2$) conditioned PAO1. ROS was measured as counts per second (CPS) when

2′,7′dichlorodihydrofluorescein diacetate was converted to 2′7′-dichlorofluorescein. Growth was measured with optical

density (OD_{600}) simultaneously to ROS measurement. Data is generated from 99 continuously measurements presented

as a solid line with dots representing $\pm$ SEM (n = 4).

Fig. 8 - An oxygen intolerant subpopulation of *Staphylococcus aureus* (methicillin susceptible) was generated in the filter biofilm model

Normoxic and anoxic colony forming units per milliliter (CFU/mL) of *Staphylococcus aureus* over 17 days from anoxically (A) and normoxically (B) conditioned filters. Symbols with error bars indicate the mean ± SEM (n = 4). $+NO_3^-$ refers to the addition of 10 mM KNO_3 to LB agar plates. The log difference (C) represents the difference in mean log CFU/mL between plating methods from the normoxically and anoxically conditioned filters, respectively. The log difference was significantly higher ($p < 0.001$, linear regression) in the anoxically conditioned filters than the normoxically conditioned. Symbols with error bars indicate the mean + confidence intervals.

Supplemental figure legends

Fig. S1 – Percentage distribution of DTC bacteria over time in different growth models

The percentage of DTC *Pseudomonas aeruginosa* after anoxic conditioning in the bead biofilm model (A), filter biofilm model (B), batch cultures (C), colonies (D). The percentage of DTC *Staphylococcus aureus* after anoxic conditioning in the filter biofilm model (E).

S2 - An oxygen intolerant subpopulation of S*taphylococcus aureus* (MRSA) and *Staphylococcus epidermidis* was generated in the filter biofilm model.

Normoxic and anoxic (± sodium pyruvate) determination of CFU/mL were determined for 1 and 9-day-old anoxically (A) and normoxically (B) conditioned filters with *Staphylococcus aureus* (MRSA) and anoxically (C) and normoxically (D) conditioned filters with *Staphylococcus epidermidis*. Significant difference between anoxic and normoxic plating ± sodium pyruvate at day 9 (p = 0.01 and p < 0.01, respectively, one-way ANOVA test) for *S. aureus*. Significant difference between anoxic and normoxic plating at day 9 (p = 0.003, one-way ANOVA test) for *S. epidermidis*. Symbols with error bars indicate the mean ± SEM (n = 3). +NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates.

S3 – Anoxic conditioning did not affect E*scherichia coli* or *Enterococcus faecalis* during growth in the filter biofilm model.

Normoxic and anoxic (± sodium pyruvate) determination of CFU/mL were determined for 1- and 9-day-old anoxically (A) and normoxically (B) conditioned filters with *E. coli* and anoxically (C) and normoxically (D) conditioned filters with *E. faecalis*. No difference between types of plating (one-way ANOVA test). Symbols with error bars indicate the mean ± SEM (n = 3). +NO_3^- refers to the addition of 10 mM KNO_3 to LB agar plates.

Fig. S4 – Implementation of a supplementary diagnostic media at the Department of Clinical Microbiology, Rigshospitalet.

Percentage distribution of culture positive findings in patient samples between the conventional method, LB agar plates and ABTG plates supplemented with 10 mM NO_3^- (A). LB plates supplemented with 10 mM NO_3^- and the conventional method found significantly more positive samples in comparison to ABTG plates supplemented with 10 mM NO_3^- (p = 0.0003 and p = 0.0001, respectively, Fisher's exact test). Number of patients with culture positive findings by conventional method and on anoxic LB plates supplemented with 10 mM NO_3^- (B).