Re-programming of *Pseudomonas syringae* pv. *actinidiae* gene expression during early stages of infection of kiwifruit

Peter McAtee[1], Lara Brian[1], Ben Curran[1,2], Otto van der Linden[1], Niels Nieuwenhuizen[1], Rebecca Henry-Kirk[1], Simona Nardozza[1], Erik H. A. Rikkerink[1], Cris G. Print[3], Andrew C. Allan[1,2], Matthew D. Templeton[1,2,4,*]

[*]*Correspondence: Matthew Templeton,* matt.templeton@plantandfood.co.nz*, The New Zealand Institute for Plant & Food Research Limited, Auckland, New Zealand*

Running Title: Re-programming of *Psa* gene expression during infection

Keywords: RNA-seq, type III effectors, pathogenicity, secreted proteins

Word count

Abstract

Background: *Pseudomonas syringae* is a widespread bacterial species complex that includes a number of significant plant pathogens.   Amongst these, *P. syringae* pv. *actinidiae* (*Psa*) initiated a worldwide pandemic in 2008 on cultivars of *Actinidia chinensis* var. *chinensis*.   To gain information about the expression of genes involved in pathogenicity we have carried out transcriptome analysis of *Psa* during the early stages of kiwifruit infection.

Results: Gene expression in *Psa* was investigated during the first five days after infection of kiwifruit plantlets, using RNA-seq.   Principal component and heatmap analyses showed distinct phases of gene expression during the time course of infection.   The first phase was an immediate transient peak of induction around three hours post inoculation (HPI) that included genes that code for a Type VI Secretion System and nutrient acquisition (particularly phosphate).   These genes are probably involved in immediate adaption to the surface of the plant.   This was followed by a significant commitment, between 3 and 24 HPI, to the induction of genes encoding the Type III Secretion System (T3SS) and Type III Secreted Effectors (T3SE).   Expression of these genes collectively accounted for 6.3% of the bacterial transcriptome at this stage.   There was considerable variation in the expression levels of individual T3SEs but all followed the same temporal expression pattern, with the exception of HopAS1, which peaked in expression at 48 HPI.   As infection progressed over the time course of five days, there was an increase in the expression of genes with roles in sugar, amino acid and sulfur transport and the production of alginate and

colanic acid.   These are both polymers that are major constituents of extracellular

polysaccharide substances (EPS) and are involved in biofilm production.

Conclusions: The results from this study indicate that there is a complex remodeling of the

transcriptome during the early stages of infection, with at least three distinct phases of

coordinated gene expression.   These include genes induced during the immediate contact

with the host, those involved in the initiation of infection, and finally those responsible for

nutrient acquisition.

Background

Pseudomonas syringae is a widespread bacterial species complex that comprises plant epiphytes and pathogens, as well as being found in non-plant environments such as waterways [1, 2]. Each pathovar of *P. syringae* has a relatively narrow host range related to the specific effector and secondary metabolite profile encoded by its accessory genome. Effectors are proteins that are secreted into plant cells via the Type III Secretion system (T3SS) that function to repress the host defense response [3]. The kiwifruit vine (*Actinidia* Lindl spp.) disease pathogen *P. syringae* pv. *actinidiae* (*Psa*) was first identified in Japan in 1984 [4, 5] and was subsequently found in Korea in the 1990s [6]. Both these strains caused canker symptoms, but did not spread from their country of origin. In 2008, a particularly virulent canker-causing strain of *Psa* was reported in Italy and it quickly decimated plantings of *A. chinensis* var. *chinensis* cultivars, particularly 'Hort16A', 'Hongyang' and 'Jin Tao' [7]. This strain was found in other kiwifruit-growing regions including New Zealand, Chile and China by 2010 [8].

Whole genome sequence analysis was carried out on over 25 strains of *Psa* representing isolates from all locations where *Psa* had been reported. Phylogenetic analysis of the core genome indicated that the canker-causing isolates formed three clades. The first clade comprised the initial isolates from Japan, the second those collected in the 1990s from Korea and the third the pandemic outbreak strains from Italy, New Zealand, Chile and China [9-11]. Isolates within these clades are designated as biovars [12]. The core genome of isolates from the pandemic clade (biovar 3) differed by very few Single

Nucleotide Polymorphisms (SNPs) suggesting that this is a clonal population; however, the

isolates from New Zealand, Italy, Chile and China each possessed a different member of a

family of integrative conjugative elements [9-11]. More recently a comprehensive

phylogenetic analysis of eighty *Psa* isolates has shown the origin of the pandemic strains to

be China [13]. Two new biovars of *Psa* have been recently discovered in Japan [14, 15],

and thus the location of the source population of *Psa* biovars has yet to be conclusively

determined.

The three canker-causing biovars each had a surprisingly varied accessory genome with

different complements of genes encoding effectors and toxins [11, 16]. Many of these

genes are encoded on putative mobile genetic elements. While bioinformatic analysis has

identified genes that might be unique to the recent outbreak clade, little is known about the

expression of these and other genes that might have a crucial role in pathogenicity.

Surprisingly there are few RNA-seq data on the early stages of infection of plants by

pathogenic bacteria, including *P. syringae*.

Several transcriptome studies have been carried out on different *P. syringae* pathovars

[17, 18]. The most comprehensive *in planta* analysis has been of *P. syringae* pv. *syringae*

(*Pss*) B728a. This pathovar is a particularly successful epiphyte as well as a pathogen of

bean (*Phaseolus vulgaris* L.). Global analysis of the transcriptome as an epiphyte,

pathogen, and under various stress conditions was carried out using a microarray

covering >5000 coding sequences [19, 20]. The transcript profiles indicated that success

as an epiphyte is enabled by flagellar motility, swarming motility based on surfactant

production, chemosensing, and chemotaxis. This could indicate active relocation primarily

on the leaf surface.   Occupation of an epiphytic niche was accompanied by high transcript

levels for phenylalanine degradation, which may help to counteract phenylpropanoid-based

plant defenses [19]. In contrast, intercellular or apoplastic colonization led to the high-level

expression of genes for γ-aminobutyric acid (GABA) metabolism (degradation of GABA

would attenuate GABA repression of virulence) and the synthesis of phytotoxins, syringolin

A and two additional secondary metabolites.   Perhaps surprisingly the T3SS and T3SEs

were not found to be strongly induced in the apoplast [19].   Subsequent analysis of several

regulatory mutants illustrated a central role for GacS, SalA, RpoN, and AlgU in global

regulation in *Pss* B728a *in planta* and a high degree of plasticity in these transcriptional

regulators' responses to distinct environmental signals [20].

More recently a comprehensive analysis of gene expression by *P. syringae* pv. *tomato*

DC3000 (*Pto*) has been carried out on wild-type Arabidopsis and several defense gene

mutants [21].   T3SS and T3SEs genes were upregulated *in planta,* as were transporter

genes.   A key finding was that Arabidopsis perturbs iron homeostasis in *Pto* [21].

To gain additional information about the expression of genes involved in pathogenicity,

we have carried out transcriptome analysis of *Psa* grown *in vitro* on minimal media and *in*

planta during the early stages of kiwifruit infection, using an RNA-seq approach.   Our

analysis showed that there are at least three distinct coordinated phases of gene expression

and has resulted in the discovery of several uncharacterized genes that may have a role in

pathogenicity.

Results

Infection assay and RNA-seq time course

Infection assays of kiwifruit tissue-cultured plantlets with *Psa* were performed in flood-inoculated tissue culture vessels under sterile conditions. This method gave consistent and reproducible infection rates, with visible leaf-spot symptoms progressively developing from day 5. A time course was carried out to assess the rate of infection and to measure and distinguish the relative populations of apoplastic and leaf surface-colonizing (epiphytic) bacteria. Bacterial counts in the apoplast (surface sterilized samples) rose rapidly during the first six days post inoculation and reached a plateau at approximately 10^8 colony-forming units (CFU) thereafter (Figure 1). Total bacterial counts (non-surface sterilized samples), which included both apoplastic and epiphytic bacteria, rose from 10^5 to 10^8 CFU during the time course. The results suggest that for the first two days of infection, the majority of live cells were located epiphytically on the surface of the plant, but that the proportion of apoplastic colonizing bacteria progressively rose from days 2 to 6 so that the apoplast became the predominant niche from that time on.

A time course of five days (120 hours) was selected for the RNA-seq analysis, with a focus on very early time points to identify genes induced in the first stages of contact with the plant surface and subsequent infection. It was postulated that key genes responsible for the initiation of infection would be induced at the early stages of contact with the plant surface. Obtaining significant numbers of bacterial reads from infected plants at the early

stages of infection is extremely challenging. For this reason, leaves were not surface

sterilized before RNA extraction.

RNA-seq expression profile

Trimmed reads were mapped onto the complete *Psa* ICMP 18884 genome (CP011972.2

and CP011973) [22]. All 27 control uninfected treatments showed no reads mapping to the

genome. A principal component analysis (PCA) was carried out on the inoculated samples

to assess overall similarity, and the three biological replicates showed little variance within

each time point (Figure 2A). PCA also demonstrated that each of the *Psa*-infected tissue

samples belonged to one of three major phases that closely aligned to the post inoculation

period and that were distinct from the *in vitro* control. The component groupings included

the *in vitro* control, an early phase of infection (1.5 and 3 hours post infection, HPI), a mid-

phase of infection (6, 12, and 24 HPI), and a late phase of infection (48, 72, 96, and 120

HPI) (Figure 2B).

Heat map analysis with k-means clustering

Comparison of expression profiles is a powerful tool that can be used to identify and discover

genes under the same regulatory regime. Furthermore, it was postulated that novel genes

that showed similar expression profiles to known genes involved in pathogenicity might also

have a role in causing disease. To identify such genes, similarities in the expression values

for each gene were determined by first normalizing expression against its maximum value

and then clustering by k-means analysis [23]. This analysis was restricted to those genes

displaying <u>R</u>eads <u>P</u>er <u>K</u>ilobase per <u>M</u>illion (RPKM) values over 50 for at least one time-point,

to eliminate lowly expressed genes from the analysis [24]. Of the 5985 predicted gene

models in the core and accessory genomes of *Psa*, 269 genes did not display evidence of

being expressed at any sample point, 1473 had no sample point with an RPKM above 50,

and 4243 genes had at least one sample point with an RPKM value above 50 (Table S1).

Hierarchical Clustering on Principal Components (HCPC) using the remaining 4243 genes

was used to partition a k-means analysis of genes into 13 clades (clusters) based on their

expression profiles (Figure 3). These were further consolidated into six groups based on

their broader expression patterns (Table 1). Of these 4243 genes, 1137 were constitutively

expressed, 1323 genes were down-regulated *in planta,* and a further 815 did not show

significant differential expression (Table 1). The remaining 968 genes were up-regulated

in planta compared with *in vitro* and thus likely to have the most direct relevance to

pathogenicity. Of the upregulated genes, there were three distinct groupings that differed

in their temporal patterns and level of gene expression over the time course. These groups

corresponded to 107 genes induced in the early (1.5 and 3 HPI) time points (early phase),

followed by a group of 311 genes highly induced between 3 and 24 HPI (mid phase). The

latter group included the majority of the T3SS and T3SE genes controlled by the HrpL

regulon. Finally, 550 genes increased in their expression towards the late (48-120 HPI) time

points (late phase). These three phases of gene expression were similar to the groupings

identified by PCA analysis (Figure 1). Expression profiles were subsequently evaluated in

more detail for genes with known or as yet undetermined roles in pathogenicity.

Early phase of infection was characterized by the induction of a type VI secretion system and nutrient adaptation

Approximately 100 genes were up-regulated immediately upon contact with the host in the early phase of infection (1.5-3.0 HPI). These were found in clade 12 from the clustering analysis (Figure 3, Table 1). The majority of these genes were annotated as being involved in nutrient acquisition (Table S2). This probably reflects the adaptation to the surface of a leaf, where nutrients are more scarce [25]. Genes that were particularly highly expressed included those predicted to be involved in phosphate and iron transport. In addition, some genes involved in the degradation of cell wall polymers, including a polygalacturonase (IYO_008325), were in this group. However, few genes predicted to have a direct role in pathogenicity were found. Two of the 43 annotated chemotactic response genes (chemoreceptors) found in the *Psa* genome were highly expressed during the early phase (Table S2), but these two are not amongst those previously functionally characterized [26]. These chemoreceptors could have a role in locating stomata or other potential sites of entry into the plant. Another set of genes that was highly induced in this phase encodes a putative Type VI Secretion System (T6SS). Effectors secreted through the T6SS have a variety of roles usually associated with killing both prokaryotic and eukaryotic cells [27]. Roles for T6SS effectors as virulence factors for animal pathogens have been well documented; however, there is as yet no evidence for an equivalent function in plant pathogens [28, 29]. Alternatively, the T6SS induced by *Psa* may have a role under field conditions in the antagonism of competing epiphytic microbes on the leaf surface.

Mid-phase of infection was characterized by expression of T3SS and T3SEs

Genes from three clades (6, 10 and 11) from the clustering analysis show increased expression 3-24 HPI (Table 1). Of these the most striking is a large transcriptional commitment to the induction of the T3SS apparatus and the expression of T3SEs, which were among the most highly upregulated genes within these time points (Figure 3; Table S3). Transcripts encoding for T3SS and T3SEs rose from 1.5 HPI, peaking between 3 and 12 HPI before falling to about half maximal levels for the remainder of the time course. Between 3 and 12 HPI these genes collectively accounted for 6.3% of the total reads (Table S3). Expression of HrpA1 was by far the highest of all T3SS genes, accounting for over 50% of these reads. The HrpA protein comprises the needle of the T3SS apparatus. For plant pathogens the needle is much longer than that of animal pathogens because of the need to penetrate the host cell wall, thus presumably requiring higher expression of the corresponding gene [30].

The *Psa* biovar 3 genome has 40 genes encoding T3SEs, and 35 of these are predicted to encode full-length proteins [11], including one additional T3SE (HopBN1) recently identified (http://pseudomonas-syringae.org/). The expression profile of T3SEs during the mid-phase of infection followed that of the T3SS transcripts, rising rapidly after 1.5 HPI, with a maximum between 3 and 12 HPI, and then falling for the rest of the time course to around 20-40% of the highest level. The expression levels of each effector varied considerably: most were relatively abundant, in particular HopAU1, HopS2, HopAO2, HopAZ1, HopZ5 and HopF2, AvrRmp1, AvrB4 and AvrPto5 peaked at over 1000 RPKM (Table S4). However, several other effectors were weakly expressed during all the early phases of infection < 150

RPKM, such as HopAH1 and HopBB1-2 (Table S4). This may be due to lack of a role for

these particular genes in the infection of kiwifruit, expression at the later stages of disease

development (after 120 HPI), or a role in the infection of tissues other than leaves.   The

effector that displayed the most distinct temporal expression profile was HopAS1.   This

effector had low expression in the second phase of infection and peaked at 48 HPI.   *Pto*

strains that are pathogenic on *Arabidopsis thaliana* carry a truncated version of this effector,

which is widely distributed in *P. syringae* [31]. Full-length versions caused effector-triggered

immunity in almost all ecotypes of Arabidopsis, explaining why it effectively operates as a

barrier to infection in this non-host. In contrast, deletion of the full-length version of HopAS1

reduced virulence of *Pto* on tomato, suggesting it has a virulence function on this natural

host. Both the *Pto* and *Psa* orthologs of this effector have a putative HrpL box situated

upstream of their putative start sites. In between lies a short uncharacterised potential open

reading frame which could be an "unrecognised" effector chaperone. The HopAS1 effector

is also found in *P. syringae* pv. *phaseolicola,* where it was not found to be differentially

expressed in response to induction of the HrpL TTSS regulatory system [32]. Unfortunately

there are no strong clues about the possible biochemical function of HopAS1. It is one of

the largest effectors (over 1300 residues, third largest *Psa* effector). Automated searching

of the conserved domain database at NCBI identified just one tentative match (Bit score 51;

E-value $5.7e^{-6}$) to a 330 residue portion of a heterodimerization domain in the N-terminus of

the chromosome maintenance protein superfamily [33, 34]. This could indicate that the

function of this effector is not focused on post-translational modification of plant defence

proteins (as the majority of *P. syringae* effectors appear to be), but that it targets host

chromatin remodelling processes associated with later stages of the defence response, and

could be an explanation for its unusual expression profile in *Psa*. Recently HopAS1 was

shown to be one of only six T3SEs from *Pto* able to bind to yeast plasma membrane, binding

to several different phospho–inositol derivatives [35]. Unfortunately this research appears to

have been performed with the truncated version of this gene from *Pto* DC3000. In contrast,

the full-length HopAS1 from *Psa* could not be localised when expressed in *Nicotiana*

benthamiana, but this may be because it triggers cell death in that host [36].

T3SS and T3SEs are under the control of the HrpL regulon and hence their co-regulation

would be expected.   Several other genes that do not code for T3SEs also possess HrpL

boxes 5´ to their start site. These include genes that encode a putative lytic transglycosylase

(IYO_006775), M20 peptidase (IYO_027210), ApbE involved in thiamine biosynthesis

(IYO_010630), a phosphatidylserine decarboxylase (IYO_025425), and an indole acetic

acid-lysine ligase (IAAL, IYO_002060, Table S5). IAAL is found adjacent to a gene encoding

a multidrug and toxic compound extrusion protein (MATE, IYO_002055) on the chromosome

of many *P. syringae* and *P. savastanoi* pathovars.   Some *P. savastanoi* pathovars have an

additional plasmid-associated IAAL copy linked with indole acetic acid (IAA) production and

gall formation.   The proteins encoded by these genes are 92% identical, and the plasmid-

located copy has been expressed heterologously and functionally characterized [37].   IAAL

is postulated to convert free IAA into less active conjugate forms [38].   Heterologous

expression of IAAL in tobacco and potato led to abnormal developmental changes [39].

Transcript levels of *Psa iaal* were induced early in infection and, in contrast to T3SS and

T3SEs, remained high throughout the infection period; however, the adjacent *mate* gene did

not appear to be highly expressed during this time period (Figure 4).   In *Pto* DC3000 it has

been shown that *iaaI* can be both transcribed independently and co-transcribed with *mate*

as an operon [40].

Other sets of genes that were strongly expressed during the mid-phase of infection (3-

12 HPI) included four co-located genes on two operons that code for a diguanylate cyclase

and two transcription factors, and thus may have a regulatory role (IYO_012110-25) (Table

S6).   Another set of four genes in two operons contains proteins involved in metal transport

(IYO_003310-25); included in these is the highly expressed copper resistance/binding

protein CopZ (IYO_003325).   Very high expression of the chemotaxis protein IYO_006420

was also observed; while not a membrane-bound chemoreceptor, it is predicted to contain

a 4-helix bundle, which is a common chemoreceptor sensor domain [26].   This protein is

predicted to be structurally similar to di-iron binding proteins (Pfam 09537), suggesting an

alternate role in iron acquisition as opposed to chemoreception.

Late phase of infection was driven by nutrient acquisition and EPS production

A total of 550 genes were upregulated in the later phase of infection (groups 7-9).   Ninety

genes increased over 5-fold in expression between 1.5 and 120 HPI (Table S7).   Of these

genes, 14 were annotated to be involved in alginate and colanic acid biosynthesis and

polymer export.   Alginate is a hygroscopic polymer composed of D-mannuronic acid

residues interspersed with L-guluronic acid residues with various degrees of acetylation [41].

This polymer has an important role in biofilm production and is well characterized in *P.*

aeruginosa [42].   *P. syringae* is also known to produce alginate, but its role in pathogenicity

is less well understood [43].   Recently it has been shown that alginate accumulates in high

amounts in the sub-stomatal spaces in *Psa*-infected leaves of kiwifruit (Sutherland et al.,

unpublished).   A further 26 genes in this grouping were annotated as having a role in

metabolite transport.   This strongly suggests that as early-stage infection progresses there

is a widespread induction of genes involved in metabolite transport and nutrient acquisition.

These transporters are distinct from those observed in the early phase of leaf colonization.

Expression of secondary metabolite pathways during infection

Several predicted secondary metabolite biosynthesis pathways have been identified in *Psa*

using either antiSMASH 3.0 [44], or by similarity to known biosynthetic pathways (Table S8).

Three of these pathways, for achromobactin, pyoverdine and yersiniabactin, are involved in

iron accumulation.   *Psa* biovar 3 produces fluorescent compounds, i.e. pyoverdine, when

grown on King's B medium, but to a lesser extent than other *Psa* biovars [8]. The genes that

code for this pathway appear to be poorly expressed *in planta* and on minimal media (Table

S8).   Genes coding for the alternative iron siderophores yersiniabactin and achromobactin

are present in *Psa* but were also expressed at low levels *in planta* (Table S8).   It has

recently been postulated that plants are able to interfere with iron homeostasis in pathogenic

bacteria, which may explain the apparent lack of expression of these pathways *in planta*

[21].   Alternatively, *Psa* may be using a different mechanism for acquiring iron.

Psa-infected kiwifruit leaves show a distinct chlorotic halo which is presumably the result

of the diffusion of a phytotoxin [8]. In addition to the three pathways with roles in iron

absorption, there were four other secondary metabolite pathways identified in the *Psa* biovar

3 genome that have potential roles in pathogenicity and might account for leaf chlorosis.

Psa biovar 3 possesses gene clusters involved in the biosynthesis of mangotoxin, a novel

non-ribosomal peptide (NRP; IYO_003775-003830), an unknown metabolite (IYO_026725-

026760), and an unknown compound synthesized from chorismate; the last-named pathway

is plasmid-borne (Table S8).   The genes involved in mangotoxin biosynthesis, NRP, and

the unknown metabolite did not appear to be significantly induced during the early stages of

infection, although genes in the NRP pathway were constitutively expressed between 50

and 100 RPKM throughout the infection time course (Table S8).   While BLAST/antiSMASH

searches did not identify likely products of either of the unknown biosynthetic pathways,

many *Pseudomonas* spp. produce surfactive molecules to wet the leaf surface to aid motility

[45].   In addition, the apoplast is a relatively dry environment that pathogens often modify

to increase the relative humidity.   For example, syfA - an NRP from *Pss* - produces an

extremely hygroscopic molecule that facilitates wetting of surfaces including the leaf surface

and apoplast [46-48].

The uncharacterized biosynthetic pathway on the plasmid of *Psa* biovar 3 has two

operons, and is adjacent to a LuxR receptor [49]. The first operon codes for a chorismate-

utilizing enzyme and a glutamine amidotransferase (annotated as anthranilate synthase I

and II) [11].   The second operon codes for the biosynthesis and secretion of a putative

aromatic, but uncharacterized, compound that was strongly induced *in planta* after 12 HPI

and remained steady for the remainder of the time course (Table S8, Figure 5).   The

plasmid-localized secondary metabolite pathway is not widespread in *P. syringae* but

interestingly is also present in the vascular pathogen *Xylella fastidiosa*, the causal agent of

Pierce's disease, and some root-associated *Pseudomonas* species [11].

Proteins secreted through the type II secretion system

In addition to the translocation of proteins through specialized structures such as the T3SS

and the T6SS, bacteria also use the Sec or Tat systems to secrete proteins into the

periplasm for the Type II secretion system (T2SS) to export [50]. This system will target

proteins *in planta* to the apoplast, as opposed to the cytoplasmic location of the T3SEs.

This is an important function since the plant apoplast has a number of largely constitutive

antimicrobial defenses such as phytoanticipins, hydrolytic enzymes and enzyme inhibitors

that may need to be inactivated to facilitate colonization. Recently analysis of the *Pto*

secretome identified a protease inhibitor *Cip1* as playing a role in virulence of *Pto* on

tomato [51]. We were therefore interested to see if there were T2SS proteins upregulated

in the early and mid-phases of infection.

Type II secreted proteins (T2SP) can be identified by their canonical secretory leader

sequence using SignalP [52]. Five hundred and thirty-nine proteins were predicted to be

secreted. Of these proteins, 21 were induced in the early phase of infection (clade12).

Of the significantly induced genes (ratio RPKM 3HPI/RPKM *in vitro* >5), the majority are

predicted subunits of membrane-bound complexes with a role in nutrient transport (Table

S9). All of these had annotations assigned.

Twenty-six proteins with predicted leader sequences were present in the mid phase of

infection (Table S9). Of those strongly expressed compared to *in vitro* growth, four were

annotated as hypothetical proteins.   However, two of these predicted gene products have

similarity to enzyme inhibitors.   IYO_001870 has homology to a superfamily of vertebrate

lysozyme inhibitors and IYO_009660 contains a region with homology to Pfam domain

13670, present in some putative protease inhibitors.   Both these proteins may have a role

in neutralizing the apoplast and are candidates for further functional analysis.   Forty-two

non-annotated secreted proteins identified in the *Pto* genome were screened for the ability

to inhibit the tomato C14 defense-related protease and one, Cip1, was shown to be an

inhibitor [51]. Of these, 37 had orthologs (95% sequence identify) in *Psa* biovar 3 but only

seven were clearly differentially expressed *in planta*.   Interestingly, the *Psa* ortholog of

Cip1 (IYO_021465) was not differentially expressed during the time course in this study.

Discussion

RNA-seq was used to investigate the early stages of infection of kiwifruit plantlets by *Psa* biovar 3. This biovar is highly virulent on kiwifruit, with apoplastic CFU reaching a plateau from 6 days post inoculation. PCA and clustering analysis revealed three phases of gene expression *in planta* during early stages of colonization by *Psa*. The first was a rapid transient phase that occurred immediately upon contact with the plant. Included in these genes was a T6SS, which might have a role in pathogenesis, similarly to that in animal systems [27]. Alternatively, the T6SS may have a role in competition against epiphytic bacteria. Interestingly none of the other genes in this group had a predicted function that could have a direct role in pathogenicity. This suggests that these early expressed *Psa* genes play a role in rapid adaptation to the plant surface, since most of the bacterial counts were on the surface of the plant rather than the apoplast at this stage. Two chemotactic receptors were also highly expressed in this early phase. These are strong candidates for a role in sensing stomata, hydathodes and other points of entry for the pathogen.

In contrast, the mid phase of infection, which occurred between 3 and 12 HPI, included the T3SS and majority of the T3SEs. These genes were the most upregulated at these time points, accounting for over 6% of the transcripts detected. Similar results were observed for *Pto* colonization of Arabidopsis [21]. This is in contrast to the data observed for *Pss* during the early stages of infection of bean, where a large induction of either T3SS or T3SEs was not observed. This might be due to different infection strategies of the two

pathogens: *Pss* is regarded as a stronger epiphyte than other *P. syringae* pathovars, because the phylogroup it belongs to (II) has a greater focus on toxin production, and its members typically have fewer effectors than other *P. syringae* phylogroups [1]. The difference could alternatively be attributed to the different experimental approaches employed [19].   Levels of individual *Psa* T3SE gene expression varied considerably during the time course.   However, the temporal expression pattern was largely consistent between effectors, with most (25/30) fitting into the mid-phase gene expression clusters 6, 10 and 11.   The most notable exception was HopAS1, which peaked later in expression around 48 HPI, as opposed to 3-12 HPI for most other effectors.

The roles of the T3SS and T3SE in repressing the induced host defense response are increasingly well understood.   Less well understood is the repression of constitutive plant defenses in the apoplast, the inactivation of which is an essential prerequisite to the establishment of the T3SS.   These defenses include phytoanticipins, cell wall degrading enzymes, proteases and enzyme inhibitors.   Furthermore, the apoplast is a relatively dry space that needs to be humidified to optimize colonization [47, 48]. Two resident proteins in the conserved effector locus, HopM and AvrE, appear to be important in establishing the right humidity conditions in other *P. syringae* hosts [53], and in *Psa* these effectors both follow the mid-peak expression profile but show only average expression levels. This study has also identified two predicted proteins that may have a role in neutralizing the apoplast. One was a predicated lysozyme inhibitor (IYO_001870) and the other a predicted protease inhibitor (IYO_009660).   It is likely, however, that there are further genes to be discovered

that play a role in neutralizing the apoplast, including the production of potential

surfactants.

The final phase comprised of genes whose expression progressively increased over

the five-day (120 h) time course.   Included were a raft of genes coding for proteins

involved in nutrient acquisition such as transporters.   Notably, these were different

transporters from those induced at the very early phase of infection.   There was also

strong induction of genes involved in alginate and colanic acid production.   These

compounds are a large component of the extracellular polysaccharide substances (EPS)

and known virulence factors of *Pseudomonas* [41, 43].   Their precise role in pathogenicity

is not known, but they have been postulated to protect the bacteria from adversity, in this

case plant defenses, and also to enhance adhesion to solid surfaces. Indeed alginate

synthesis, along with ice nucleation, auxin synthesis and auxin inactivation by IAAL, is

common among the canonical *P. syringae* lineages that have been traced back to a last

common ancestor (LCA) 150-180 million years ago [54]. Another component predicted

to be derived from the LCA is the tripartite pathogenicity island structure consisting of

the *hrp*/*hrc* gene cluster flanked by both the Conserved Effector Locus and an

exchangeable effector locus (EEL). *Psa* shows this tripartite structure, albeit that the

EEL is further away from the other two pathogenicity islands. The heat map analysis did

not highlight any obvious differences in expression patterns between the effectors

located on these three pathogenicity islands.

While effectors are well known to play a role in plant defense suppression, the role of

many other genes expressed during infection is far less certain.   This study has identified

a number of non-effector genes that were strongly induced *in planta* and are likely to be

having a role in establishing infection.   The relative importance of these will need to be

ascertained using either gene knockouts or TraDIS (Transposon Directed Insertion-site

Sequencing) [55, 56].

Conclusions

The results from this study indicate that there is a complex remodeling of the bacterial

transcriptome during the early stages of infection, with at least three distinct phases of

coordinated gene expression.   The first includes genes induced during the immediate

contact with the host.   These were dominated by the expression of a T6SS and genes

annotated as involved in nutrient transport.   The second phase was dominated by genes

predicted to have roles in initiating infection and includes the T3SS and T3SEs.   Included

in this group are novel proteins that may have roles in neutralizing constitutive defenses in

the apoplast.   The final phase includes genes involved in nutrient transport and biofilm

formation.

Experimental Procedures

Infection assays

Actinidia chinensis Planch. var. *chinensis* 'Hort16A' plantlets, grown from axillary buds on Murashige and Skoog rooting medium without antibiotics in a 400-mL clear plastic tub with a sealed lid, were purchased from Multiflora (http://www.multiflora.co.nz/home.htm). Plantlets were grown at 20°C under fluorescent lights with a 16 h on/8 h off regime and used within a month of purchase. For inoculation an overnight shake culture of *Psa* ICMP 18884 was grown in liquid Lysogeny Broth (LB) [57] at 20°C and 180 rpm shaking. The cell density was determined by measuring the absorbance at 535 nm. Cells were washed in 10 mM $MgSO_4$ and resuspended at a cell density of 10^7 CFU/mL. The surfactant Silwet L-77 (Cat VIS-30, Lehle Seeds, Round Rock, TX, USA) was added to the inoculum to a concentration of 0.0025% (v/v) to facilitate leaf wetting. The inoculation method was modified from the method developed for Arabidopsis [58]. Containers with 'Hort16A' plantlets were filled with the inoculum fully submerging the plantlets and left for three minutes. Containers were drained, the lid replaced, then incubated in a controlled climate room at 20°C with a light/dark cycle of 16 h on/8 h off.

Growth assay

Leaf samples were taken at different times post inoculation as appropriate. Each sample consisted of four leaf discs, taken with a 1-cm diameter cork borer, from four different leaves. All four discs were taken from the same tub. To estimate CFU in the apoplast discs were

surface sterilized in 70% (v/v) ethanol for 30 s and subsequently washed in sterile Milli-Q

water. Samples for estimation of total bacteria were not surface sterilized. Leaf discs

were placed in Eppendorf tubes containing three stainless steel ball bearings and 300 µL 10

mM $MgSO_4$, and macerated in a bead crusher for 2 min at maximum speed (Storm 24 Bullet

Blender, Next Advance, Averill Park, NY, USA). A dilution series of the leaf homogenates

was made in sterile 10 mM $MgSO_4$ until a dilution of 10^{-8}. The dilution series was plated in

5-µL droplets on LB medium supplemented with both 12.5 µg/mL nitrofurantoin and 40

µg/mL cephalexin. After 72 hours of incubation at 20°C CFU were counted for the lowest

possible dilution(s), which was calculated back to the CFU per cm^2 of leaf area.

RNA extractions

RNA was extracted from *Psa* ICMP 18884 grown to late log phase at 18°C on Hoitnik and

Sinden minimal media [59]. Cells were harvested and total RNA extracted using an Ambion

RNA extraction kit (Thermo Fisher, Waltham, MA, USA). RNA was extracted from 1-month-

old *A. chinensis* var. *chinensis* 'Hort16A' plantlets propagated from tissue culture infected

with *Psa* as described above after 1.5, 3, 6, 12, 24, 48, 72, 96 and 120 HPI. Each time point

consisted of three biological replicates. Three pots were used for each time point and each

biological replicates consisted of three combined plantlets sampled across each of the three

pots (Figure S1). Mock-inoculated plants were used as controls for each time point. RNA

was extracted using the Spectrum™ Plant Total RNA Kit (Sigma-Aldrich, Milwaukee, WI,

USA). Sequencing libraries were constructed from total RNA using the Ribo-Zero Plant

procedure (Illumina, San Diego, CA, USA).

Bioinformatics and differential expression analysis

Sequencing was performed using HiSeq2000 (Illumina) by Macrogen (www.macrogen.com).

Raw RNA reads (100 bp paired end reads) were trimmed, quality filtered (≥Q20) and their

adaptors removed using Trimmomatic v0.36 [60]. The cleaned reads were mapped to the

Psa ICMP 18884 gene models [22] using the Bowtie2 v2.25 aligner in conjunction with

samtools [61]. To ensure the best alignments were made against each gene model, a

mapping quality > 10 was used.

Analysis of mapped reads was done using the statistical software R (version 3.4.3). K-

means cluster analysis of expression data was done using the R packages ggplot2 (v2.2.1)

[62], FactoMineR (v1.39) [63] and FactoExtra (v1.05) [64]. Principal component analysis was

done using the R package DESeq2 (v1.18.1) [65].

Abbreviations

CFU: Colony-forming units; EEL: Exchangeable effector locus; GABA: γ-aminobutyric acid; HPCP: Hierarchical clustering on principal components; HPI: Hours post inoculation; EPS: Extracellular polysaccharide substances; LB: Lysogeny broth; LCA: Last common ancestor; PCA: Principal component analysis; *Psa*: *Pseudomonas syringae* pv. *actinidiae*; *Pss*: *Pseudomonas syringae* pv. *syringae;* Pto: NRP: Non-ribosomal peptide; *Pseudomonas syringae* pv. *tomato*; RPKM: Reads per kilobase per million; SNP: Single nucleotide polymorphism; T2SS: Type II secretion system; T2SP: Type II secreted proteins; T3SS: Type III Secretion System; T3SE: Type III Secreted Effectors; T6SS: Type VI Secretion System; TraDIS: Transposon directed insertion-site sequencing.

Acknowledgements

We would like to thank Dr Joanna K. Bowen and Amali Thrimawithana (Plant and Food Research) for critically reviewing this manuscript.

Funding

This work was funded by a grant from the New Zealand Ministry for Business, Innovation and Employment (C11X1205).

Availability of data and materials

The RNA-seq experiment is described in BioProject PRJNA472664 with separate BioSamples for each time-point (SAMN09240241-97), reads can be downloaded from the

540   Sequence Read Archive SRP148711 [66].

Author contributions

543   BC, ACA and CGP conceived of and designed the experiments.   BC, RH-K and OvdL

544   carried out the experiments and generated the data. PM, LB, EHR and MDT analyzed the

545   data.   MDT, PM, NN, SN, EHR, and ACA wrote the paper.

Competing interests

548   The authors declare that they have no competing interests.

Ethics approval and consent to participate

551   All experiments using *Psa* were carried out with the permission of the Ministry for Primary

552   Industries, New Zealand (CTO approval 12-05-17) and The Environmental Protection

553   Authority, New Zealand (APP202231).

Author details

556   [1]The New Zealand Institute for Plant and Food Research Limited, Auckland, New Zealand

557   [2]School of Biological Sciences, University of Auckland, Auckland, New Zealand

558   [3]Department of Molecular Medicine and Pathology, University of Auckland, New Zealand

559   [4]Bio-Protection Research Centre, New Zealand

References

1. Baltrus DA, Nishimura MT, Romanchuk A, Chang JH, Mukhtar MS, Cherkis K et al. Dynamic evolution of pathogenicity revealed by sequencing and comparative genomics of 19 *Pseudomonas syringae* isolates. PLoS Pathog. 2011;7(7):e1002132.

2. Monteil CL, Cai R, Liu H, Mechan Llontop ME, Leman S, Studholme DJ et al. Nonagricultural reservoirs contribute to emergence and evolution of *Pseudomonas syringae* crop pathogens. The New phytologist. 2013.

3. Büttner D. Behind the lines–actions of bacterial type III effector proteins in plant cells. FEMS Microbiology Reviews. 2016;40(6):894-937.

4. Serizawa S, Ichikawa T, Takikawa Y, Tsuyumu S, Goto M. Occurrence of bacterial canker of kiwifruit in Japan description of symptoms, isolation of the pathogen and screening of bactericides. Japanese Journal of Phytopathology. 1989;55(4):427-36.

5. Takikawa Y, Serizawa S, Ichikawa T, Tsuyumu S, Goto M. *Pseudomonas syringae* pv. *actinidiae* pv. nov the causal bacterium of canker of Kiwifruit in Japan. Japanese Journal of Phytopathology. 1989;55(4):437-44.

6. Koh Y CB, Chung H, Lee D. Outbreak and spread of bacterial canker in kiwifruit. Korean Journal of Plant Pathology. 1994;1068-72.

7. Balestra GM, Mazzaglia A, Quattrucci A, Renzi M, Rossetti A. Current status of bacterial canker spread on kiwifruit in Italy. Australasian Plant Disease Notes. 2009;434.

8. Everett KR, Taylor RK, Romberg MK, Rees-George J, Fullerton RA, Vanneste JL et al. First report of *Pseudomonas syringae* pv. *actinidiae* causing kiwifruit bacterial canker in New Zealand. Australasian Plant Disease Notes. 2011;667-71.

9. Butler MI, Stockwell PA, Black MA, Day RC, Lamont IL, Poulter RTM. *Pseudomonas syringae* pv. *actinidiae* from recent outbreaks of kiwifruit bacterial canker belong to different clones that originated in China. Plos One. 2013;8(2):18.

10. Mazzaglia A, Studholme DJ, Taratufolo MC, Cai RM, Almeida NF, Goodman T et al. *Pseudomonas syringae* pv. *actinidiae* (Psa) Isolates from recent bacterial canker of Kiwifruit outbreaks belong to the same genetic lineage. Plos One. 2012;7(5):11.

11. McCann HC, Rikkerink EHA, Bertels F, Fiers M, Lu A, Rees-George J et al. Genomic analysis of the Kiwifruit pathogen *Pseudomonas syringae* pv. *actinidiae* provides insight into the origins of an emergent plant disease. PLoS Pathog. 2013;9(7):e1003503.

12. Cunty A, Poliakoff F, Rivoal C, Cesbron S, Saux ML, Lemaire C et al. Characterization of *Pseudomonas syringae* pv. *actinidiae* (*Psa*) isolated from France and assignment of Psa biovar 4 to a *de novo* pathovar: *Pseudomonas syringae* pv. *actinidifoliorum* pv. nov. Plant Pathol. 2015;64(3):582-96.

13. McCann HC, Li L, Liu Y, Li D, Hui P, Zhong C et al. The origin and evolution of a pandemic lineage of the Kiwifruit pathogen *Pseudomonas syringae* pv. *actinidiae* Genome Biology and Evolution. 2017;9(4):932-44.

14. Fujikawa T, Sawada H. Genome analysis of the kiwifruit canker pathogen Pseudomonas syringae pv. actinidiae biovar 5. Scientific Reports. 2016;621399.

15. Sawada H, Kondo K, Nakaune R. Novel biovar (biovar 6) of *Pseudomonas syringae* pv. *actinidiae* causing bacterial canker of kiwifruit (Actinidia deliciosa) in Japan. Jpn J Phytopathol. 2016;82101-15.

16. Marcelletti S, Ferrante P, Petriccione M, Firrao G, Scortichini M. *Pseudomonas syringae* pv. *actinidiae* draft genomes comparison reveal strain-specific features involved in adaptation and virulence to Actinidia species. Plos One. 2011;6(11):17.

17. Filiatrault MJ, Stodghill PV, Bronstein PA, Moll S, Lindeberg M, Grills G et al. Transcriptome analysis of

604		*Pseudomonas syringae* identifies new genes, noncoding RNAs, and antisense activity. Journal of Bacteriology.
605		2010;192(9):2359-72.
606	18.	Filiatrault MJ, Stodghill PV, Wilson J, Butcher BG, Chen H, Myers CR et al. *CrcZ* and *CrcX* regulate carbon source
607		utilization in *Pseudomonas syringae* pathovar *tomato* strain DC3000. RNA Biology. 2013;10(2):245-55.
608	19.	Yu X, Lund SP, Scott RA, Greenwald JW, Records AH, Nettleton D et al. Transcriptional responses of
609		*Pseudomonas syringae* to growth in epiphytic versus apoplastic leaf sites. Proc Natl Acad Sci U S A.
610		2013;110(5):E425-34.
611	20.	Yu X, Lund SP, Greenwald JW, Records AH, Scott RA, Nettleton D et al. Transcriptional analysis of the global
612		regulatory networks active in *Pseudomonas syringae* during leaf colonization. mBio. 2014;5(5):e01683-14.
613	21.	Nobori T, Velásquez AC, Wu J, Kvitko BH, Kremer JM, Wang Y et al. Transcriptome landscape of a bacterial
614		pathogen under plant immunity. Proceedings of the National Academy of Sciences. 2018;115(13):E3055-E64.
615	22.	Templeton MD, Warren BA, Andersen MT, Rikkerink EHA, Fineran PC. Complete DNA sequence of
616		*Pseudomonas syringae* pv. *actinidiae*, the causal agent of Kiwifruit canker disease. Genome Announcements.
617		2015;3(5).
618	23.	Liu P, Si Y: Cluster analysis of RNA-sequencing data. In: Statistical Analysis of Next Generation Sequencing
619		Data. Edited by Datta S, Nettleton D. Cham: Springer International Publishing; 2014: 191-217.
620	24.	Lee S, Seo CH, Lim B, Yang JO, Oh J, Lim M et al. Accurate quantification of transcriptome from RNA-Seq data
621		by effective length normalization Nucleic Acids Research. 2011;39(2):e9.
622	25.	Mercier J, Lindow SE. Role of leaf surface sugars in colonization of plants by bacterial epiphytes. Applied and
623		Environmental Microbiology. 2000;66(1):369-74.
624	26.	McKellar JLO, Minnell JJ, Gerth ML. A high-throughput screen for ligand binding reveals the specificities of
625		three amino acid chemoreceptors from *Pseudomonas syringae* pv. *actinidiae*. Molecular Microbiology.
626		2015;96(4):694-707.
627	27.	Kostiuk B, Unterweger D, Provenzano D, Pukatzki S. T6SS intraspecific competition orchestrates *Vibrio*
628		*cholerae* genotypic diversity. Int Microbiol. 2017;20(3):130-7.
629	28.	Ho Brian T, Dong Tao G, Mekalanos John J. A view to a kill: The bacterial type VI secretion system. Cell Host &
630		Microbe. 2014;15(1):9-21.
631	29.	Russell AB, Peterson SB, Mougous JD. Type VI secretion system effectors: poisons with a purpose. Nat Rev
632		Micro. 2014;12(2):137-48.
633	30.	Roine E, Wei W, Yuan J, Nurmiaho-Lassila EL, Kalkkinen N, M. R et al. Hrp pilus: an hrp-dependent bacterial
634		surface appendage produced by *Pseudomonas syringae* pv. *tomato* DC3000. Proc Natl Acad Sci USA.
635		1997;94(7):3459-64.
636	31.	Sohn KH, Saucet SB, Clarke CR, Vinatzer BA, O'Brien HE, Guttman DS et al. HopAS1 recognition significantly
637		contributes to Arabidopsis nonhost resistance to *Pseudomonas syringae* pathogens. The New phytologist.
638		2012;193(1):58-66.
639	32.	Mucyn TS, Yourstone S, Lind AL, Biswas S, Nishimura MT, Baltrus DA et al. Variable suites of non-effector
640		genes are co-regulated in the type III secretion virulence regulon across the *Pseudomonas syringae* phylogeny.
641		PLoS pathogens. 2014;10(1):e1003807.
642	33.	Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S et al. CDD/SPARCLE: functional classification of proteins
643		via subfamily domain architectures. Nucleic Acids Research. 2017;45(Database issue):D200-D3.
644	34.	Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY et al. CDD: NCBI's conserved domain
645		database. Nucleic Acids Res. 2015;43(Database issue):D222-6.
646	35.	Weigele BA, Orchard RC, Jimenez A, Cox GW, Alto NM. A systematic exploration of the interactions between
647		bacterial effector proteins and host cell membranes. Nature Communications. 2017;8532.

36.   Choi S, Jayaraman J, Segonzac C, Park H-J, Park H, Han S-W et al. *Pseudomonas syringae* pv. *actinidiae* type III
effectors localized at multiple cellular compartments activate or suppress innate Immune responses in
Nicotiana benthamiana. Frontiers in Plant Science. 2017;8(2157).

37.   Glass NL, Kosuge T. Cloning of the gene for indoleacetic acid-lysine synthetase from *Pseudomonas syringae*
subsp. *savastanoi*. Journal of Bacteriology. 1986;166(2):598-603.

38.   Ostrowski M, Mierek-Adamska A, Porowińska D, Goc A, Jakubowska A. Cloning and biochemical
characterization of indole-3-acetic acid-amino acid synthetase PsGH3 from pea. Plant Physiology and
Biochemistry. 2016;1079-20.

39.   Spena A, Prinsen E, Fladung M, Schulze SC, Van Onckelen H. The indoleacetic acid-lysine synthetase gene of
Pseudomonas syringae subsp. *savastanoi* induces developmental alterations in transgenic tobacco and potato
plants. Mol Gen Genet. 1991;227(2):205-12.

40.   Castillo-Lizardo MG, Aragón IM, Carvajal V, Matas IM, Pérez-Bueno ML, Gallegos M-T et al. Contribution of the
non-effector members of the HrpL regulon, iaaL and matE, to the virulence of *Pseudomonas syringae* pv.
tomato DC3000 in tomato plants. BMC Microbiology. 2015;15165.

41.   Franklin MJ, Nivens DE, Weadge JT, Howell PL. Biosynthesis of the *Pseudomonas aeruginosa* extracellular
polysaccharides, alginate, pel, and psl. Frontiers in Microbiology. 2011;2167.

42.   Ghafoor A, Hay ID, Rehm BHA. Role of exopolysaccharides in *Pseudomonas aeruginosa* biofilm formation and
architecture. Applied and Environmental Microbiology. 2011;77(15):5238-46.

43.   Markel E, Stodghill P, Bao Z, Myers CR, Swingle B. AlgU controls expression of virulence genes in *Pseudomonas*
syringae pv. *tomato* DC3000. Journal of Bacteriology. 2016;198(17):2330-44.

44.   Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R et al. antiSMASH 3.0—a comprehensive resource for
the genome mining of biosynthetic gene clusters. Nucleic Acids Research. 2015;43(W1):W237-W43.

45.   Alsohim AS, Taylor TB, Barrett GA, Gallie J, Zhang X-X, Altamirano-Junqueira AE et al. The biosurfactant
viscosin produced by *Pseudomonas fluorescens* SBW25 aids spreading motility and plant growth promotion.
Environmental Microbiology. 2014;16(7):2267-81.

46.   Hockett KL, Burch AY, Lindow SE. Thermo-regulation of genes mediating motility and plant interactions in
Pseudomonas syringae. PLOS ONE. 2013;8(3):e59850.

47.   Burch AY, Shimada BK, Browne PJ, Lindow SE. Novel high-throughput detection method to assess bacterial
surfactant production. Applied and Environmental Microbiology. 2010;76(16):5363-72.

48.   Burch AY, Shimada BK, Mullin SWA, Dunlap CA, Bowman MJ, Lindow SE. *Pseudomonas syringae* coordinates
production of a motility-enabling surfactant with flagellar assembly. Journal of Bacteriology.
2012;194(6):1287-98.

49.   Patel HK, Ferrante P, Covaceuszach S, Lamba D, Scortichini M, Venturi V. The kiwifruit emerging pathogen
Pseudomonas syringae pv. actinidiae does not produce AHLs but possesses three luxR solos. PLoS One.
2014;9(1):e87862.

50.   Green ER, Mecsas J. Bacterial secretion systems – an overview. Microbiology spectrum.
2016;4(1):10.1128/microbiolspec.VMBF-0012-2015.

51.   Shindo T, Kaschani F, Yang F, Kovács J, Tian F, Kourelis J et al. Screen of non-annotated small secreted proteins
of *Pseudomonas syringae* reveals a virulence factor that inhibits Tomato immune proteases. PLoS pathogens.
2016;12(9):e1005874.

52.   Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from
transmembrane regions. Nature Methods. 2011;8785.

53.   Xin XF, Nomura K, Aung K, Velasquez AC, Yao J, Boutrot F et al. Bacteria establish an aqueous living space in
plants crucial for virulence. Nature. 2016;539(7630):524-9.

54.　Xin X-F, Kvitko B, He SY. *Pseudomonas syringae*: what it takes to be a pathogen. Nature Reviews Microbiology.
2018.

55.　Mesarich CH, Rees-George J, Gardner PP, Ghomi FA, Gerth ML, Andersen MT et al. Transposon insertion
libraries for the characterization of mutants from the kiwifruit pathogen Pseudomonas syringae pv. actinidiae.
PLOS ONE. 2017;12(3):e0172790.

56.　Barquist L, Mayho M, C. C, Cain AK, Boinett CJ, Page AJ et al. The TraDIS toolkit: sequencing and analysis for
dense transposon mutant libraries. Bioinformatics. 2016;32(7):1109-11.

57.　Bertani G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. J Bacteriol.
1951;62(3):293-3000.

58.　Ishiga Y, Ishiga T, Uppalapati SR, Mysore KS. Arabidopsis seedling flood-inoculation technique: a rapid and
reliable assay for studying plant-bacterial interactions. . 2011;7:32. Plant Methods. 2011;732.

59.　Hoitink HAJ, Sinden SL. Partial purification and properties of chlorosis inducing toxins of *Pseudomonas
phaseolicola* and *Pseudomonas glycinea*. Phytopathology. 1970;601236-7.

60.　Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics.
2014;30(15):2114-20.

61.　Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA
sequences to the human genome. Genome Biol 2009;10(3):R25.

62.　Warnes GR, Bolker B, Bonebakker L, Gentleman R, Wolfgang Huber W, Liaw A et al. gplots: various R
programming tools for plotting data. R package version 301 2016.

63.　Le S, Josse J, F. H. FactoMineR: An R package for multivariate analysis. Journal of Statistical Software.
2008;25(1):1-18.

64.　Kassambara A, Mundt F. factoextra: extract and visualize the results of multivariate data analyses. R package
version 105. 2017.

65.　Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for　RNA-seq data with
DESeq2. Genome Biology. 2014;15(12):550.

66.　Sayers EW, Barrett T, Benson DA, Bolton E, Bryant SH, Canese K et al. Database resources of the National
Center for Biotechnology Information. Nucleic Acids Res. 2012;40(Database issue):D13-25.

Table 1. k-means clustering of 4243 genes Psa into 13 clades. HPI, hours post infection.

Group	k-means clade	Number of genes	Description	Expression Phase
1	1 & 13	1137	Constitutively expressed genes	N/A
2	2,3, & 4	1323	Genes down-regulated *in planta*	N/A
3	5	815	Little differential expression compared to *in vitro*	N/A
4	6, 10 & 11	311	Genes upregulated (3-24 HPI) *in planta*	Mid
5	7,8, & 9	550	Genes upregulated late (48-120 HPI) *in planta*	Late
6	12	107	Early upregulated (1.5-3 HPI) genes *in planta*	Early

Figure Legends

Figure 1. Time course of *Pseudomonas syringae* pv. *actinidiae* (*Psa*) infection of kiwifruit plantlets over 14 days. CFU, colony forming units; DPI, days post inoculation.   (●) surface-sterilized (■) non-surface sterilized.   The experiment was duplicated and each had four technical replicates.   Error bars represent SE, n=2.   Zero-time controls had no *Psa* present.

Figure 2. Principal component analysis plot (PCA) showing the clustering of vst (variance stabilizing transformation) transcriptomic data. In the left plot data points are colored by treatment time point (1.5 HPI, 3 HPI, 6 HPI, 12 HPI, 24 HPI, 48 HPI, 72 HPI, 96 HPI and 120 HPI). In the right plot data points are colored by infection phase (*in vitro*, Early (1.5-3 HPI), Mid (6-24 HPI), Late (48-120 HPI)). HPI, hours post infection.

Figure 3. Heatmap and k-means clustering showing the expression of *Pseudomonas syringae* pv. *actinidiae* (*Psa*) genes in 'Hort16A' kiwifruit plantlets post infection. Similar expression profiles were clustered into 13 distinct groups by k-means. Line graphs displaying the prototype mean expression of each cluster are included on the right.

Figure 4. Expression of genes encoding *IAAl* and *Mate*.   *DNAA* was included as a constitutively expressed control.

Figure 5. Expression of genes from the plasmid-borne operon coding for a putative aromatic compound.

Figure S1. Experimental design of RNA-seq sampling.

753   Figure 1.

766   Figure 2.

768   A   B

769
770

771   Figure 3.

774   Figure 4.

775
776

777   Figure 5.

781   Supplementary Figures

782   Figure S1. RNA-seq experimental design

802 Table S1. RPKM for all genes

804 Table S2. Early induced genes ranked by the ratio of expression at 3 HPI compared with *in*
805 *vitro* (cutoff 5-fold). HPI, hours post infection.
806

Gene ID	Gene Annotation	3 HPI/ *in vitro*
IYO_011995	phosphate ABC transporter substrate-binding protein	148.8
IYO_012000	phosphate ABC transporter permease	43.7
IYO_019585	thioredoxin	31.4
IYO_012010	phosphate ABC transporter ATP-binding protein	30.8
IYO_018790	magnesium transporter CorA	30.8
IYO_027385	ABC transporter substrate-binding protein	29.5
IYO_027390	GntR family transcriptional regulator	25.0
IYO_015395	hypothetical protein	23.9
IYO_006115	amino acid ABC transporter substrate-binding protein	23.3
IYO_027380	ABC transporter permease	21.5
IYO_028665	phosphate-binding protein	20.4
IYO_018545	acid phosphatase	19.5
IYO_000970	ammonia channel protein	18.7
IYO_006555	chemotaxis protein	17.4
IYO_021410	short-chain dehydrogenase	17.2
IYO_013555	type VI secretion effector protein (Hcp)	16.3
IYO_013560	EvpB family type VI secretion protein	15.9
IYO_028645	transcriptional regulator PhoU	14.3
IYO_002185	peptidase M19	13.6
IYO_010675	phosphatase	12.6
IYO_021050	ABC transporter substrate-binding protein	12.5
IYO_013550	Type VI secretion protein	12.2
IYO_013565	type VI secretion protein	11.7
IYO_001685	MFS transporter	11.6
IYO_021420	polysaccharide deacetylase	10.9
IYO_020035	ABC transporter substrate-binding protein	10.8
IYO_027365	metallophosphatase	10.4
IYO_014740	sugar ABC transporter substrate-binding protein	10.2
IYO_002190	hydrocarbon binding protein	9.8
IYO_020310	hypothetical protein	9.7
IYO_013545	type VI secretion system protein ImpG	9.5
IYO_006130	glutamine ABC transporter ATP-binding protein	9.4
IYO_028615	transcriptional regulator PhoB	9.2
IYO_004270	chemotaxis protein	9.1
IYO_001975	nitrogen regulation protein NR(I)	9.1
IYO_002205	electron transfer flavoprotein subunit alpha	8.8
IYO_025185	urease accessory protein UreG	8.7

IYO_021405	3-oxoacyl-ACP reductase	8.5
IYO_026920	ABC transporter permease	8.5
IYO_006120	amino acid ABC transporter permease	7.5
IYO_002210	electron transfer flavoprotein subunit beta	7.4
IYO_027370	iron ABC transporter substrate-binding protein	7.3
IYO_027375	ABC transporter permease	6.8
IYO_018365	MFS transporter	6.6
IYO_014745	xylose isomerase	6.3
IYO_018070	ATPase	6.2
IYO_006260	acetyltransferase	6.0
IYO_027495	sarcosine oxidase subunit alpha	5.9
IYO_001120	hypothetical protein	5.8
IYO_026150	acyl carrier protein	5.6
IYO_004585	branched-chain amino acid ABC transporter substrate-binding protein	5.6
IYO_009210	quercetin 2,3-dioxygenase	5.5
IYO_014735	xylose transporter	5.5
IYO_025180	urease accessory protein UreF	5.5
IYO_021415	MFS transporter	5.2

807
808

809 Table S3.   Genes most highly upregulated in the mid phase of the infection time course
810 (3-24 hours post infection).   Genes are ranked on the ratio of maximum Reads Per
811 Kilobase per Million reads over that time period to *in vitro* expression (cutoff 5-fold).
812

Gene ID	Gene Annotation	Ratio
IYO_006750	type III effector HrpW	257.8
IYO_022020	hemolysin	177.0
IYO_006820	type III secretion protein	118.7
IYO_006755	Shc Hop M1 (disrupted)	104.3
IYO_006865	type III secretion system protein	77.2
IYO_006790	HrpA1	77.0
IYO_004052	HopS2	76.4
IYO_006825	type III secretion protein	67.6
IYO_006875	type III secretion protein	63.0
IYO_006880	type III secretion protein	54.1
IYO_006795	type III secretion protein HrpZ	52.4
IYO_012110	Ais protein	52.2
IYO_006905	RNA polymerase sigma factor HrpL	50.3
IYO_022025	glycerol acyltransferase	47.3
IYO_004050	type III chaperone ShcO1	44.3
IYO_006830	secretin	44.3
IYO_028770	LysR family transcriptional regulator	44.2
IYO_003325	copper resistance protein CopZ	41.4
IYO_006910	type III effector HrpK	39.6
IYO_012005	phosphate ABC transporter permease	37.7
IYO_002060	IAA lysine ligase	36.0
IYO_006420	chemotaxis protein	35.7
IYO_014395	lytic transglycosylase	31.2
IYO_006890	type III secretion protein	30.5
IYO_014235	hypothetical protein	30.4
IYO_006860	type III secretion system protein SsaR	30.3
IYO_006870	type III secretion system protein	29.4
IYO_006815	type III secretion protein	29.0
IYO_005735	glycoside hydrolase	25.9
IYO_006900	type III secretion protein HrpJ	25.2
IYO_003725	type III chaperone protein ShcF	23.8
IYO_006800	type III secretion protein	23.3
IYO_020425	AvrPto5	23.3
IYO_006810	type III secretion protein	22.4
IYO_006805	type III secretion protein	22.1
IYO_006885	ATP synthase	21.2
IYO_006855	type III secretory protein EscS	20.9

IYO_006765	type III chaperone ShcE	18.7
IYO_006760	HopM1	18.2
IYO_014245	membrane protein	18.0
IYO_003570	avrD1	18.0
IYO_029290	type III chaperone protein ShcF	17.8
IYO_006850	type III secretion system protein	17.5
IYO_029040	type III secretion chaperone CesT	16.9
IYO_008282	HopZ5	16.8
IYO_009200	hypothetical protein	16.7
IYO_005160	HopI1	16.6
IYO_012125	diguanylate cyclase	16.1
IYO_006895	type III secretion protein HrpI	15.5
IYO_003600	AvrB4	15.5
IYO_022140	SAM-dependent methyltransferase	14.6
IYO_012120	AraC family transcriptional regulator	13.9
IYO_002045	hypothetical protein	13.8
IYO_017375	phosphonate/organophosphate ester transporter subunit	13.2
IYO_018555	HopAZ1	12.9
IYO_012115	XRE family transcriptional regulator	12.2
IYO_006745	HopAA1-1	11.9
IYO_028535	NADP transhydrogenase subunit alpha	11.7
IYO_008285	HopH1	11.5
IYO_008065	AvrRpm1	11.4
IYO_010805	LuxR family transcriptional regulator	11.4
IYO_029795	HopAU1	10.9
IYO_006770	AvrE1	10.8
IYO_003720	HopAO2	10.7
IYO_000845	HopY1	10.7
IYO_012140	protein tolQ	10.3
IYO_006250	tail protein	10.1
IYO_022695	alkaline phosphatase	9.8
IYO_009265	serine/threonine protein phosphatase	9.4
IYO_016255	Ais protein	9.2
IYO_024150	HopR1	9.0
IYO_009660	hypothetical protein	8.9
IYO_027435	DNA polymerase III subunit epsilon	8.4
IYO_012610	MarR family transcriptional regulator	8.4
IYO_013150	HopBN1	8.0
IYO_002040	hypothetical protein	8.0
IYO_027360	transcriptional initiation protein Tat	8.0
IYO_012030	nitrite reductase	7.7

IYO_013145	type III chaperone protein ShcF	7.7
IYO_028955	nitrate ABC transporter ATP-binding protein	7.7
IYO_003727	HopBB1-1	7.7
IYO_012145	biopolymer transporter TolR	7.5
IYO_029288	AvrRpm2 (frameshifts)	7.5
IYO_003315	metal ABC transporter ATPase	7.5
IYO_028380	type III chaperone protein ShcA	7.5
IYO_000385	dodecin flavoprotein	7.5
IYO_006845	type III secretion system protein	7.4
IYO_028960	sulfonate ABC transporter permease	7.3
IYO_014240	hypothetical	7.2
IYO_023505	chemotaxis protein	7.1
IYO_028540	NAD(P) transhydrogenase	7.0
IYO_005855	UDP-N-acetylglucosamine 2-epimerase	6.8
IYO_001870	hypothetical protein	6.8
IYO_013690	membrane protein	6.7
IYO_010630	thiamine biosynthesis protein ApbE	6.6
IYO_003680	HopAF1	6.6
IYO_023400	energy transducer TonB	6.4
IYO_011020	chemotaxis protein	6.4
IYO_029045	HopZ3	6.3
IYO_023390	biopolymer transporter ExbB	6.2
IYO_024520	voltage-gated chloride channel protein	5.9
IYO_009335	Fe-S oxidoreductase	5.8
IYO_024535	hypothetical protein	5.8
IYO_004060	hypothetical protein	5.7
IYO_021665	MFS transporter	5.7
IYO_016185	UDP-4-amino-4-deoxy-L-arabinose-oxoglutarate aminotransferase	5.5
IYO_020420	iron ABC transporter permease	5.5
IYO_022135	InaA protein	5.5
IYO_007455	membrane protein	5.3
IYO_016195	UDP-4-amino-4-deoxy-L-arabinose formyltransferase	5.3
IYO_022030	ACP phosphodiesterase	5.2
IYO_006775	lytic transglycosylase	5.2
IYO_004045	lipoprotein	5.1
IYO_018725	membrane protein	5.1
IYO_012605	fusaric acid resistance protein	5.1
IYO_004240	hypothetical protein	5.0
IYO_014250	chemotaxis protein CheY	5.0

Table S4.   Expression levels of individual *Psa* effectors over the infection time course.
Effectors are ranked by the highest level of expression between 3 and 12 HPI in Reads
Per Kilobase per Million reads. Effectors likely to be disrupted or pseudogenes were not
included. HPI, hours post infection. ND = not determined.

Gene ID	Effector	Heat map group	RPKM									
			in vitro	1.5 HPI	3 HPI	6 HPI	12 HPI	24 HPI	48 HPI	72 HPI	96 HPI	120 HPI
IYO_029795	HopAU1	11	346.4	970.6	3772.5	2876.6	3196.1	2141.7	1138.8	952.1	619.7	946.2
IYO_008065	AvrRpm1	11	123.9	365.5	1293.2	867.8	1415.6	940.2	538.8	542.9	526.2	693.6
IYO_020425	AvrPto5	11	58.8	331	1198.8	920.8	1368.6	770.7	472.4	424.3	281.7	487.2
IYO_018555	HopAZ1	11	102.1	226.8	995	616.3	1320.1	1178.3	479.8	682.5	464.8	788
IYO_004052	HopS2	11	15	181.7	1145.3	751.5	762.2	494.6	324.2	239.9	129	226.5
IYO_003720	HopAO2	11	102.7	346.6	1102.7	664.4	821.5	560.7	357.2	378.4	228.5	356.7
IYO_003600	AvrB4	11	65.6	326.8	925.5	696.7	1015.7	621.4	332.6	431.6	251.2	465.3
IYO_013150	HopBN1	11	105.3	179.4	633	468.2	843.7	622.5	327.7	440.3	232.6	342.3
IYO_008282	HopZ5	11	43.4	314.1	662.1	546.9	727.8	426.2	222	186.1	141.9	440.9
IYO_003570	avrD1	6	38.2	99.4	432.1	412.6	687.1	492.6	240.1	419.9	264.3	703.8
IYO_024217	HopF2	11	187.9	225.4	658.8	479.4	649.8	563.5	288	400.6	213.4	403
IYO_003657	HopAW1	11	195	166.1	401.6	375.5	539	402.8	208.9	254.2	187	361.9
IYO_006735	HopN1	7	200.4	301.6	380.7	510.9	478.6	467.8	383.2	394.1	347.3	314.1

IYO_008285	HopH1	11	43.1	147.6	493.5	241.2	462.3	284.1	135.7	158.6	80.9	177.6
IYO_000845	HopY1	11	42.8	121.7	426.6	279.2	459.5	317	172.9	238.8	115.9	204.8
IYO_005160	HopI1	11	22.3	96.9	370	246.3	367	324.7	169.6	217.6	143.6	181.2
IYO_006760	HopM1	11	20.3	106.4	368.9	286.3	347.5	186.1	98.1	102.9	69.2	114.3
IYO_003727	HopBB1 -1	11	47.3	103.6	362.9	266.3	232.3	173.2	76.2	107	143.6	100.6
IYO_003680	HopAF1	6	46	90.6	176.3	145.4	302.4	321.1	145.9	186.5	154.3	213.1
IYO_029045	HopZ3	11	47.3	89.7	248.7	179.1	300.1	206.6	111.2	116.5	84.1	135.8
IYO_006770	AvrE1	11	27	73.9	291.1	176.3	164.6	129.1	57.6	53.6	44.9	47.8
IYO_003525	HopQ1	11	53.6	96.9	165	163.3	232.5	146.8	86.5	131.6	79.6	168.7
IYO_003530	HopD1	7	53.5	99.3	212.3	184	229.5	216.6	156.4	170.4	124.3	156.3
IYO_024150	HopR1	11	24.1	58.5	217.6	103	94	68.1	77.2	112.8	45.6	94.9
IYO_012225	HopAE1	11	31.2	53.1	133.8	95.1	142.7	174.9	115.4	111.7	71.9	121.5
IYO_003675	HopBB1 -2	6	32.1	40.7	129.6	67.4	148.5	104.5	55.6	50.1	21.9	136.4
IYO_027420	HopAS1	9	24.9	11.4	12.3	17.1	21.7	60.3	129.9	109.4	38.4	53.4
IYO_023985	HopAH1	1	53.5	111.4	111.3	110.1	81.9	93.5	70.5	82.1	57.6	42.3
IYO_006745	HopAA1 -1	11	9.2	27.7	107.2	60.6	109.3	80.5	32.5	27.7	12.8	31
IYO_003635	HopX3*	5	53.1	93.4	67.9	70.6	65.6	33.7	63.6	107.1	59.7	88.2
	HopAM1 -1	ND										
	HopAM1 -2	ND										

821   Table S5.   Expression of non-effector genes with HrpL boxes. Genes are ranked by the
822   ration of expression at 12 HPI compared with *in vitro* expression. HPI, hours post infection.
823
824

Gene ID	Gene Annotation	Cluster	12 HPI/*in vitro*
IYO_002060	IAA lysine ligase	11	36.0
IYO_006775	lytic transglycosylase	11	5.2
IYO_010630	thiamine biosynthesis protein ApbE	11	5
IYO_027210	peptidase M20	11	3.5
IYO_025425	phosphatidylserine decarboxylase	11	3.4
IYO_002055	multidrug transporter Mate	7	2.5
IYO_000225	AraC transcription factor	8	1.5
IYO_008215	Transporter	1	1.2

825
826

827 Table S6.   Genes expressed in the mid phase of infection that do not encode the T3SS or
828 T3SEs.   Effectors were ranked by the highest level of expression between 3 and 24 HPI
829 compared with *in vitro* (cutoff 5-fold). HPI, hours post infection.
830

Gene ID	Gene Annotation	Max 3-12 / *in vitro*
IYO_022020	hemolysin	177.0
IYO_012110	Ais protein	52.2
IYO_022025	glycerol acyltransferase	47.3
IYO_028770	LysR family transcriptional regulator	44.2
IYO_003325	copper resistance protein CopZ	41.4
IYO_012005	phosphate ABC transporter permease	37.7
IYO_028385	transposase	36.1
IYO_006420	chemotaxis protein	35.7
IYO_014235	hypothetical protein	30.4
IYO_005735	glycoside hydrolase	25.9
IYO_014245	membrane protein	18.0
IYO_009200	hypothetical protein	16.7
IYO_012125	diguanylate cyclase	16.1
IYO_022140	SAM-dependent methyltransferase	14.6
IYO_012120	AraC family transcriptional regulator	13.9
IYO_002045	hypothetical protein	13.8
IYO_017375	phosphonate/organophosphate ester transporter subunit	13.2
IYO_012115	XRE family transcriptional regulator	12.2
IYO_028535	NADP transhydrogenase subunit alpha	11.7
IYO_010805	LuxR family transcriptional regulator	11.4
IYO_012140	protein tolQ	10.3
IYO_006250	tail protein	10.1
IYO_022695	alkaline phosphatase	9.8
IYO_009265	serine/threonine protein phosphatase	9.4
IYO_016255	Ais protein	9.2
IYO_009660	hypothetical protein IYO_009660	8.9
IYO_027435	DNA polymerase III subunit epsilon	8.4
IYO_012610	MarR family transcriptional regulator	8.4
IYO_002040	hypothetical protein	8.0
IYO_027360	transcriptional initiation protein Tat	8.0
IYO_012030	nitrite reductase	7.7
IYO_028955	nitrate ABC transporter ATP-binding protein	7.7
IYO_012145	biopolymer transporter TolR	7.5
IYO_003315	metal ABC transporter ATPase	7.5
IYO_000385	dodecin flavoprotein	7.5

IYO_028960	sulfonate ABC transporter permease	7.5
IYO_014240	Hypothetical protein	7.3
IYO_023505	chemotaxis protein	7.2
IYO_028540	NAD(P) transhydrogenase	7.1
IYO_005855	UDP-N-acetylglucosamine 2-epimerase	7.0
IYO_001870	hypothetical protein	6.8
IYO_013690	membrane protein	6.8
IYO_010630	thiamine biosynthesis protein ApbE	6.7
IYO_023400	energy transducer TonB	6.6
IYO_011020	chemotaxis protein	6.4
IYO_023390	biopolymer transporter ExbB	6.4
IYO_024520	voltage-gated chloride channel protein	6.2
IYO_009335	Fe-S oxidoreductase	5.9
IYO_024535	hypothetical protein	5.8
IYO_004060	hypothetical protein IYO_004060	5.8
IYO_021665	MFS transporter	5.7
IYO_016185	UDP-4-amino-4-deoxy-L-arabinose-oxoglutarate aminotransferase	5.7
IYO_020420	iron ABC transporter permease	5.5
IYO_022135	InaA protein	5.5
IYO_007455	membrane protein	5.5
IYO_016195	UDP-4-amino-4-deoxy-L-arabinose formyltransferase	5.3
IYO_022030	ACP phosphodiesterase	5.3
IYO_004045	lipoprotein	5.2
IYO_018725	membrane protein	5.1
IYO_012605	fusaric acid resistance protein	5.1
IYO_004240	hypothetical protein	5.1
IYO_014250	chemotaxis protein CheY	5.0

833 Table S7. Late upregulated genes. Genes were ranked based on the ratio of expression at
834 120 compared with 1.5 HPI. HPI, hours post infection.
835

Gene ID	Gene Annotation	120HPI RPKM/1.5 HPI RPKM
IYO_016130	ABC transporter	119.9
IYO_016135	acyl-CoA synthetase	93.1
IYO_006065	glycosyl transferase	75.6
IYO_013825	alkanesulfonate monooxygenase	51.2
IYO_006070	GDP-mannose dehydrogenase	51.1
IYO_009605	YqcI/YcgG family protein	39.1
IYO_023405	biopolymer transporter ExbD	34.3
IYO_026605	monooxygenase	33.2
IYO_017250	energy transducer TonB	28.3
IYO_017245	biopolymer transporter ExbB	28.0
IYO_027905	transporter	26.9
IYO_016105	acyl-CoA dehydrogenase	24.8
IYO_005580	hypothetical protein	24.2
IYO_027910	aliphatic sulfonates transport ATP-binding subunit	23.7
IYO_006055	alginate biosynthesis protein	22.7
IYO_006050	alginate regulatory protein	20.0
IYO_009255	sulfonate ABC transporter ATP-binding protein	19.9
IYO_006015	mannose-1-phosphate guanylyltransferase	19.3
IYO_018210	calcium-binding protein	19.2
IYO_008315	lipoprotein	18.3
IYO_016100	acyl-CoA dehydrogenase	18.2
IYO_016120	ABC transporter permease	16.7
IYO_026615	N5,N10-methylene tetrahydromethanopterin reductase	16.5
IYO_015950	polar amino acid ABC transporter permease	15.6
IYO_006060	hemolysin D	15.3
IYO_006040	alginate O-acetyltransferase	15.1
IYO_026620	methionine ABC transporter substrate-binding protein	15.0
IYO_026625	ABC transporter	15.0
IYO_027920	ABC transporter substrate-binding protein	14.1
IYO_015300	ABC transporter permease	13.1
IYO_026675	sulfonate ABC transporter ATP-binding protein	12.7
IYO_006030	poly(beta-D-mannuronate) O-acetylase	12.3
IYO_011010	catalase	12.0
IYO_016095	5,10-methylene tetrahydromethanopterin reductase	11.7
IYO_014785	sugar ABC transporter	11.4
IYO_026610	acyl-CoA dehydrogenase	11.0
IYO_006025	alginate O-acetyltransferase	11.0

IYO_012440	hypothetical protein	10.7
IYO_020560	peptidase M4	10.4
IYO_011310	NAD(P)H-dependent FMN reductase	10.2
IYO_024090	porin	10.1
IYO_003290	hypothetical protein	10.0
IYO_010385	lipoprotein	10.0
IYO_027985	hypothetical protein	9.8
IYO_006045	poly(beta-D-mannuronate) C5 epimerase	9.6
IYO_006035	poly(beta-D-mannuronate) lyase	9.6
IYO_014780	sugar ABC transporter substrate-binding protein	9.5
IYO_016140	monooxygenase	9.1
IYO_017240	biopolymer transporter ExbD	9.1
IYO_011305	lysine transporter LysE	8.7
IYO_001790	taurine transporter ATP-binding subunit	8.7
IYO_026775	alpha/beta hydrolase	8.6
IYO_024095	ABC transporter substrate-binding protein	8.5
IYO_027980	ABC transporter permease	8.1
IYO_026630	ABC transporter permease	7.8
IYO_001460	prophage PssSM-01	7.7
IYO_011840	hemolysin D	7.7
IYO_013055	aldolase	7.4
IYO_026680	taurine dioxygenase	7.2
IYO_014465	hypothetical protein	6.9
IYO_014710	lipoprotein	6.8
IYO_017510	lipoprotein	6.8
IYO_001820	hypothetical protein	6.8
IYO_027915	alkanesulfonate transporter permease subunit	6.7
IYO_016110	branched-chain amino acid ABC transporter ATP-binding protein	6.6
IYO_009290	LTXXQ domain-containing protein	6.6
IYO_006935	hypothetical protein	6.5
IYO_016115	ABC transporter permease	6.3
IYO_009230	sulfurtransferase	6.1
IYO_013050	nitrate ABC transporter substrate-binding protein	6.0
IYO_020620	hypothetical protein	5.8
IYO_011220	Fis family transcriptional regulator	5.8
IYO_029660	coenzyme F390 synthetase (plasmid)	5.6
IYO_027965	sulfate ABC transporter ATP-binding protein	5.4
IYO_005875	hypothetical protein	5.4
IYO_016125	ABC transporter permease	5.4
IYO_001465	prophage PssSM-01	5.3
IYO_011375	class V aminotransferase	5.3

IYO_009250	ABC transporter permease	5.3
IYO_001810	ribonucleotide reductase	5.2
IYO_004495	hypothetical protein	5.1
IYO_001805	transposase	5.1
IYO_028000	diguanylate cyclase	5.0
IYO_009615	serine dehydratase	5.0

838   Table S8.   Expression levels of secondary metabolite gene clusters.   Reads Per

839   Kilobase per Million reads were means plus/minus standard deviation for each gene

840   across all time points.

Secondary metabolite pathway	Function	Gene members	Induction levels *in planta*
Novel Non-ribosomal peptide synthetase	unknown	IYO_003775-003830	Constitutive expression *in planta* (average RPKM 67 +/- 43)
Pyoverdine	Iron chelation	IYO_010820-010910	Constitutive expression *in planta* (average RPKM 41 +/- 43)
Achromobactin	Iron chelation	IYO_013460-013515	Constitutive expression *in planta* (average RPKM 17 +/- 9)
Yersiniabactin	Iron chelation	IYO_013840-013910	Constitutive expression *in planta* (average RPKM 15 +/-19)
Unknown	unknown	IYO_026725-026760	Weak constitutive *in planta* (expression 48 +/-16 RPKM)
Mangotoxin	Inhibitor of ornithine deacetylase	IYO_028470-028715	Weak expressed *in planta* (average RPKM 13 +/- 9)
Plasmid-borne pathway	unknown	IYO_029645-029685	Induced late *in planta* (see figure 5)

Table S9. RPKM values of genes encoding proteins predicted to be secreted via T2SS. HPI, hours post infection.

# ID	Function	Phase	*in vitro*	1.5 HPI	3 HPI	12 HPI	24 HPI	8 HPI	72 HPI	96 HPI	120 HPI
IYO_011995	phosphate ABC transporter substrate-binding protein	Early	39.5	3366.6	5876.1	3269.3	2774.8	1220.1	1143.4	1114.5	1774.3
IYO_019585	thioredoxin	Early	59.4	1525.9	1863.2	983.2	433.9	374.4	329.6	350.7	469.3
IYO_027385	ABC transporter substrate-binding protein	Early	10.4	222.5	306.4	105.9	117.6	54.4	68.2	7.9	58.9
IYO_006115	amino acid ABC transporter substrate-binding protein	Early	117.7	3408.6	2742.5	843.8	829.8	596.2	750.2	622.4	722.3
IYO_028665	phosphate-binding protein	Early	33.6	759.5	683.8	278.3	194.3	145.9	138.5	166.4	210.9
IYO_000970	ammonia channel protein	Early	43.5	1336.6	811.4	274.2	268.1	231.2	409.6	259.8	426.5
IYO_021410	short-chain dehydrogenase	Early	6.6	142.6	113.3	26.3	33.4	19.3	4.4	19.6	12.1
IYO_010675	phosphatase	Early	32.6	228.2	410.1	158.8	97.9	151.1	56.9	80.3	89.7
IYO_021050	ABC transporter substrate-binding protein	Early	55.0	945.8	685.8	281.4	281.5	212.0	245.3	245.3	224.0
IYO_020035	ABC transporter substrate-binding protein	Early	131.3	1169.0	1415.7	583.5	416.4	253.4	327.7	249.7	292.0
IYO_014740	sugar ABC transporter substrate-binding protein	Early	59.2	490.2	604.3	243.3	163.4	143.6	89.5	131.6	96.6
IYO_020310	hypothetical protein	Early	24.9	380.2	242.5	139.5	139.0	187.8	112.7	43.1	75.4
IYO_004585	branched-chain amino acid ABC transporter substrate-binding protein	Early	87.8	956.2	489.1	174.2	232.2	130.6	255.0	172.8	338.3
IYO_025190	protein hupE	Early	25.0	198.5	123.6	36.9	61.1	38.8	90.6	89.1	123.2
IYO_006385	porin	Early	1383.7	7387.9	6833.1	1618.0	1721.0	1130.5	1055.6	984.5	945.0
IYO_020485	glycine/betaine ABC transporter substrate-binding protein	Early	33.5	134.6	129.0	35.6	46.6	43.4	44.4	37.1	32.5
IYO_008325	polygalacturonase	Early	164.0	369.3	564.7	150.2	140.3	107.1	97.9	83.8	107.4
IYO_021455	Methylamine utilization protein MauL	Early	94.4	582.6	317.5	319.1	399.1	422.6	247.9	279.5	147.7
IYO_004580	urea ABC transporter permease	Early	14.9	74.3	43.4	20.5	16.3	35.5	30.8	19.4	22.0
IYO_026915	amino acid ABC transporter substrate-binding protein	Early	50.4	426.4	139.6	38.5	65.7	44.0	87.4	35.9	98.4
IYO_006365	sugar ABC transporter substrate-binding protein	Early	1316.7	2310.1	3559.7	671.4	762.4	398.4	474.9	375.7	335.4
IYO_024670	hypothetical protein part of ICE	Early	34.2	23.9	18.9	59.0	31.6	29.2	22.7	8.5	38.9
IYO_006805	type III secretion protein	Mid	57.1	309.6	1263.6	1201.0	751.4	414.5	501.0	276.3	450.1
IYO_002045	hypothetical protein	Mid	73.0	63.8	30.4	1006.5	102.8	120.8	106.7	104.0	50.3

IYO_009660	hypothetical protein	Mid	31.6	11.5	0.0	280.3	84.6	121.0	73.3	66.4	81.0
IYO_002040	hypothetical protein	Mid	60.1	51.1	27.5	479.1	77.9	74.5	104.8	106.2	117.1
IYO_001870	hypothetical protein	Mid	510.6	318.1	478.1	2793.5	2242.8	1524.5	1518.2	1974.0	1162.6
IYO_006020	alginate O-acetyltransferase	Mid	88.2	49.0	27.9	423.9	197.5	278.0	383.9	259.6	417.2
IYO_023395	TonB-dependent receptor	Mid	25.2	19.7	20.4	98.7	101.4	46.8	37.8	47.0	83.2
IYO_008760	sorbosone dehydrogenase	Mid	105.4	49.9	40.3	391.5	269.6	218.2	157.1	176.8	125.4
IYO_018720	sorbosone dehydrogenase	Mid	60.3	37.9	25.8	224.0	111.0	97.4	102.4	122.0	93.4
IYO_027210	peptidase M20	Mid	53.5	35.4	114.4	189.8	132.1	98.8	88.0	68.1	119.3
IYO_022715	phospholipid-binding protein	Mid	8230.8	2550.5	2568.7	29098.2	18992.7	15528.5	13209.7	16287.6	10178.4
IYO_004060	hypothetical protein	Mid	1959.4	7898.0	5063.3	6887.1	5478.9	4687.6	4501.3	3573.6	3898.1
IYO_019200	BNR/Asp-box repeat-containing protein	Mid	56.0	117.1	157.2	196.2	133.8	88.1	59.6	80.2	66.9
IYO_004055	membrane protein	Mid	34.6	39.5	114.7	86.9	62.9	37.4	39.6	39.0	33.7
IYO_006835	type III secretion protein	Mid	312.2	275.5	532.0	847.3	672.8	399.1	381.4	330.7	383.1
IYO_020600	ABC transporter substrate-binding protein	Mid	76.1	47.3	60.0	198.5	143.9	138.9	98.6	106.3	119.6
IYO_022515	toluene tolerance protein	Mid	116.7	73.1	109.9	251.8	129.2	148.1	174.9	121.3	128.6
IYO_006575	superoxide dismutase	Mid	178.1	46.6	35.6	375.8	202.9	114.6	173.8	176.6	223.3
IYO_022005	hypothetical protein	Mid	45.1	33.6	34.6	88.2	69.7	55.2	0.0	17.9	66.5
IYO_027595	phosphorylcholine phosphatase	Mid	50.4	33.8	36.3	82.4	97.7	53.8	62.8	63.3	64.2
IYO_011990	hypothetical protein	Mid	1900.4	539.1	489.4	3491.9	1831.7	1693.3	1220.1	1023.3	836.5
IYO_027840	ABC transporter substrate-binding protein	Mid	70.8	93.0	65.3	56.8	99.4	36.2	24.1	37.9	42.6
IYO_017485	hypothetical protein	Mid	300.2	114.9	115.4	411.4	188.4	259.0	298.3	216.1	202.3
IYO_010560	cytochrome C	Mid	86.6	54.4	56.6	92.6	76.9	66.5	80.4	56.4	49.9
IYO_014770	hypothetical protein	Mid	208.1	169.6	157.6	200.6	190.5	206.4	197.7	172.8	410.9
IYO_011885	type III effector	Mid	25.2	6.4	3.6	7.2	7.8	8.5	6.0	3.4	2.9