Associations between Afrotropical bats, parasites, and microbial symbionts

Lutz HL[1,2], Jackson EW[3], Dick, CW[2,4], Webala PW[5], Babyesiza WS[6], Kerbis

Peterhans JC[2,7], Demos TC[2], Patterson BD[2], Gilbert JA[1,8]

[1] Department of Surgery, University of Chicago, Chicago, Illinois, USA.

[2] Integrative Research Center, Field Museum of Natural History, Chicago,

Illinois, USA

[3] Department of Microbiology, Cornell University, Ithaca, New York, USA

[4]Department of Biology, Western Kentucky University, Bowling Green, Kentucky,

USA

[5] Department of Forestry and Wildlife Management, Maasai Mara University,

Narok, Kenya

[6] Department of Wildlife Management, Sokoine University of Agriculture,

Morogoro, Tanzania

[7] Department of Biological Sciences, Roosevelt University, Chicago, Illinois, USA

[8] Biosciences Division, Argonne National Laboratory, Argonne, Illinois, USA

Corresponding authors:

Holly L. Lutz & Jack A. Gilbert

Department of Surgery

University of Chicago

5851 S. Maryland Avenue

Chicago, IL 60637 - 1508

hlutz@surgery.bsd.uchicago.edu | gilbertjack@uchicago.edu

ABSTRACT

Bats are among the most diverse animals on the planet and harbor numerous

bacterial, viral, and eukaryotic symbionts. The interplay between bacterial

community composition and parasitism in bats is not well understood and may

have important implications for studies of similar systems. Here we present a

comprehensive survey of dipteran and haemosporidian parasites, and

characterize the gut, oral, and skin microbiota of Afrotropical bats. We identify

significant correlations between bacterial community composition of the skin and

dipteran ectoparasite prevalence across four major bat lineages, as well as links

between the oral microbiome and malarial parasitism, suggesting a potential

mechanism for host selection and vector-borne disease transmission in bats.

Mirroring recent studies of host-microbiome co-speciation in mammals, we find

a weak correlation between chiropteran phylogenetic distances and bacterial

community dissimilarity across the three anatomical sites, suggesting that host

environment is more important than shared ancestry in shaping the composition

of associated bacterial communities.

Keywords: microbiome, malaria, vector-borne disease, Afrotropics, Chiroptera

SIGNIFICANCE

Animals rely on bacterial symbionts for numerous biological functions, such as

digestion and immune system development. Increasing evidence suggests that

host-associated microbes may play a role in mediating parasite burden. This

study is the first to provide a comprehensive survey of bacterial symbionts from

multiple anatomical sites across a broad taxonomic range of Afrotropical bats,

demonstrating significant associations between the bat microbiome and parasite

prevalence. This study provides a framework for future approaches to systems

biology of host-symbiont interactions across broad taxonomic scales, which will

allow for the recognition of the interdependence between microbial symbionts

and vertebrate health in the study of wild organisms and their natural history.

\body

INTRODUCTION

Humans and other animals rely on bacterial symbionts for numerous

biological functions, such as digestion and immune system development (1, 2).

Many studies have found significant associations between host phylogeny (shared

common ancestry) and bacterial community composition (3, 4), while others

have identified spatiotemporal variables as significant drivers of host-microbe

associations over the course of individual lifespans (5-7). The influence of

microbes on their hosts may be context dependent, such that the presence of a

particular microbe may be beneficial under one set of ecological conditions and

harmful under another. Thus, patterns of association between vertebrates and

bacterial symbionts provide a unique lens through which to explore evolutionary

and ecological phenomena.

Recognition of the interdependence between microbial symbionts and

vertebrate health has led to a growing paradigm shift in the study of wild

organisms and their natural history. Vertebrate species not only exhibit inherent

life history characteristics, but serve as hosts to myriad bacteria, archaea, viruses,

fungi, and eukaryotic organisms that abound in their environments. Many

relationships between eukaryotic parasites and hosts have ancient origins, and

the same may be true for host-microbial associations. Indeed, it is possible that

bacterial symbionts of vertebrate hosts interact with eukaryotic parasites, viruses,

or fungal symbionts in ways that could ultimately shape host evolution (8). For

example, evidence from human and anthropophilic mosquito interactions

suggests that the skin microbiome can influence vector feeding preference,

thereby affecting transmission patterns of mosquito-borne pathogens (such as

WNV, yellow fever, dengue, malaria, etc.), and ultimately imposing selective

pressures on human populations - indeed, positive selection of malaria-protective

genes can be seen in the human genome (9). Despite the potential significance of

such interactions between hosts, microbes, and pathogen-transmitting vectors,

they have not been well studied in most wild vertebrate systems.

Bats (Mammalia: Chiroptera) are an important system for comparison of

the relative contributions of evolutionary and ecological factors driving host-

symbiont associations. In addition to being one of the most speciose orders of

mammals (second only to the order Rodentia), bats frequently live in large

colonies, are long-lived, and volant, granting them access to a wide geographic

range relative to their non-volant mammalian counterparts. The associations of

diverse eukaryotic parasites (e.g. dipteran insects, haemosporidia, helminths)

within numerous bat lineages have been well-characterized (10-13). Furthermore,

bats have received increasing attention due to their role as putative vectors of

human pathogens (*e.g.* Ebola, Marburg, SARS (14, 15)). Indeed, numerous

serological surveys have supported the role of Afrotropical bats as reservoirs for a

number of viruses (16-18). Taken together, these features make bats an appealing

and tractable model for studying the interaction of bacterial symbionts and non-

bacterial parasites and pathogens.

In this study, we conduct the first broad-scale study of Afrotropical bat-

associated microbes. We test associations between bacterial community

composition in the gastrointestinal tract, skin, and oral cavities from nine

families and nineteen genera of bats. We pair this information with host-parasite

associations between bats and ectoparasites in the superfamily Hippoboscoidea

(obligate hematophagous dipteran insects), and haemosporidian (malarial)

parasites putatively vectored by these hippoboscoid insects. Using a combination

of machine learning, network theory, and negative binomial distribution models,

we test the hypothesis that host-associated bacterial communities predict

prevalence of parasitism by obligate dipteran and malarial parasites.

RESULTS

1) Ectoparasite and malarial parasite prevalence among Afrotropical bats

Sampling was conducted across 20 sites in Kenya and Uganda from July-

August of 2016. Sites ranged from sea level to ~2500m in elevation (Fig. 1; Table

S1). We collected gut, oral, and skin samples for bacterial community

characterization from a total of 495 individual bats, comprising 9 families, 19

genera, and 28 recognized species. Bat families with the greatest representation

included Hipposideridae (n = 80), Miniopteridae (n = 116), Rhinolophidae (n =

88), and Pteropodidae (n = 106). All host and parasite vouchers are accessioned

at the Field Museum of Natural History (Chicago, IL, USA) (Table S2).

Miniopterid bats experienced the highest prevalence of both ectoparasitism (*M.*

minor, 89%) and malarial parasitism (*M. minor*, 67%) (Table 1). Bats with

similarly high ectoparasite prevalence at the host species level included

Rhinolophus eloquens (79% prevalence), *Stenonycteris lanosus* (62%), and

Triaenops afer (60%). Unlike miniopterid bats, these bat species did not harbor

any detectable malarial parasites (Table 1).

2) Bacterial richness of bat skin drastically exceeds that of gut or oral

communities

Across all samples, 51,136 Exact Sequence Variants (ESVs) were identified

using Deblur (19). Gut microbial communities exhibited the lowest overall

diversity (9,804 ESVs), followed by oral (13,629 ESVs), and skin (46,904 ESVs),

the latter being significantly greater than gut or oral ($p < 2.2e\text{-}16$, Kruskal-Wallis;

Bonferroni corrected p-value $p < 1e\text{-}113$, Dunn's test) (Fig. 2A). Aggregate mean

observed ESVs by host genus were 70, 93, and 531 for gut, oral, and skin samples,

respectively (Table 2). As with observed ESV richness counts, the Shannon index

of bat skin microbial communities was significantly greater than that of either gut

or oral microbiota ($p < 2.2e\text{-}16$, Kruskal-Wallis; Bonferroni corrected p-value $p <$

$1e\text{-}119$, Dunn's Test) (Fig. 2B). Based on weighted UniFrac distances, measures of

intraspecific beta dispersion revealed a continuum of dissimilarities across all

host species (Fig. 3). Mean beta dispersion among anatomical sites differed

significantly ($p < 1.2e\text{-}7$, Kruskal-Wallis; Bonferroni corrected p-value $p < 0.01$,

Dunn's Test). Meausres of intraspecific beta dispersion among unweighted

UniFrac and Bray-Curtis distances also showed a continuum of dissimilarities

across host species, and exhibited significant differences in mean beta dispersion

across anatomical sites (Fig. S1).

3) Microbial communities significantly correlate with geographic locality,

anatomical site, and host taxonomy, but not host phylogeny

Permutational analysis of variance (PERMANOVA) identified geographic

locality, host taxonomy, and anatomical sampling site (gut, oral, skin) as

significant factors explaining variation in three independent measures of

microbial beta diversity (Bray-Curtis, unweighted UniFrac, and weighted

UniFrac) ($p < 0.001$, ADONIS) (Table 4). Secondary analysis of sites by elevation

revealed that bats at higher elevations tended to host increased alpha diversity

across gut, oral, and skin microbiomes ($p < 2e\text{-}16$, linear regression) (Fig. S2). In

general, gut microbiota were dominated by Proteobacteria (Enterobacteraceae)

and Firmicutes (Bacillaceae). Oral microbiota were dominated by Proteobacteria

(Neisseriaceae, Pasteurellaceae). The oral microbiota of several insect bat

families (Miniopteridae, Nycteridae, Rhinolophidae) were enriched for

Firmicutes in the Mycoplasmataceae family, while the oral microbiota of fruit

bats (Pteropodidae) were enriched for Firmicutes in the Streptococaccea family.

Similar to gut and oral microbiota, skin also showed a high relative abundance of

Proteobacteria (Moraxellaceae, Enterobacteraceae) and Firmicutes (Bacillaceae),

with a pronounced increase in relative abundance of Actinobacteria and

Bacteroidetes (Fig. 4).

Linear regression analyses of host phylogenetic distances and microbial

community dissimilarity (unweighted UniFrac (uf) and weighted UniFrac (wuf)

distances) revealed weak correlations for gut (uf: $R^2 = 0.013$, $p < 0.05$; wuf: $R^2 =$

0.002, $p = 0.752$; $R^2 = 0.0007$, $p = 0.2643$; $R^2 = 0.0015$, $p = 0.522$), oral ($R^2 =$

0.009; $p < 0.05$), and skin ($R^2 = 0.024$; $p < 0.005$) microbiota and host

evolutionary relatedness (Fig. S3).

4) The microbiome is associated with parasitism in African bats

To test for significant associations between bacterial communities and

eukaryotic parasites (obligate ectoparasitic dipteran insects, and obligate

endoparasitic malarial parasites), we employed a combination of machine

learning techniques, network analyses, and negative binomial distribution

models (see methods). PERMANOVA analysis identified ectoparasite status and

malarial infection status as significant predictors of bacterial beta diversity

dissimilarity among skin and oral microbiota, respectively ($p < 0.001$, ADONIS).

Tests of three independent measures of beta diversity (weighted UniFrac,

unweighted UniFrac, and Bray-Curtis) produced congruent results, with the

exception of oral microbiome, which was not significantly predictive of malarial

infection based on unweighted UniFrac analysis (Table 3).

Supervised machine learning analyses (random forests; see methods)

produced models that could classify the anatomical source of microbial

communities and the host genus of gut, oral, and skin microbial samples with

reasonable accuracy (ratio of baseline to observed classification error ≥2; *i.e.*

random forest models performed at least twice as well as random). Random

forest models also performed well when classifying ectoparasite status based on

skin bacterial community composition, but less well for classification of malarial

status based on oral bacterial community composition (Table 5).

Following the application of statistical and machine learning approaches,

we employed network analyses to characterize the co-occurrence topology of

microbial communities (in terms of the relative abundance of co-occurring ESVs)

across the skin microbiota of our four most well-sampled bat families

(Hipposideridae (n = 80), Miniopteridae (n = 116), Rhinolophida (n = 88), and

Pteropodidae (n = 106)). Network analyses produced strikingly consistent results,

revealing a significant decrease in cluster size (p < 0.05, Mann-Whitney-

Wilcoxon rank sum test) and median node degree (p < 0.05, t test), as well as

reduced network connectivity for parasitized bats from three of the four bat

families examined (Fig. 5; Fig. S4).

5) Bacterial taxa on skin correlated with presence or absence of obligate dipteran

ectoparasites

Negative binomial distribution (*e.g*. DESeq) models applied to skin

microbiota in four well-sampled bat families (Hipposideridae, Miniopteridae,

Rhinolophidae, Pteropodidae) identified a number of ESVs that were

significantly associated with either ectoparasitized or non-ectoparasitized bats

(Fig. 6). Overall, we identified 89 and 24 ESVs significantly associated with

parasitized and non-parasitized bats, respectively (Table S3). Bacterial classes

with the greatest representation among significant results were Actinobacteria

(16 families), Gammaproteobacteria (11 families), Bacilli (5 families), and

Alphaproteobacteria (3 families). ESVs significantly enriched in parasitized bats

from at least three out of four bat families included Mycobacteraceae

(Actinobacteria), and Xanthomonadaceae (Gammaproteobacteria). ESVs

significantly enriched in parasitized bats from at least two out of four bat families

included Hyphomicrobiaceae (Alphaproteobacteria), Alcaligenaceae

(Betaproteobacteria), Moraxellaceae (Gammaproteobacteria), Planococcaceae

(Bacilli), Flavobacteraceae (Flavobacteria), Halobacteraceae (Halobacteria), and

Chitinophagaceae (Saprospirae) (Fig. 6).

DISCUSSION

The bacterial diversity we observed among gut, oral, and skin microbiota

of bats fall within ranges similarly observed in other vertebrate groups (3, 20-23).

Although few studies have simultaneously compared gut, oral, and skin

microbiota from the same individuals, our data refelct an apparent trend in the

literature of skin bacterial diversity among vertebrates significantly

outnumbering gut or oral bacterial diversity (24-27). Our data corroborate the

findings of Nishida and Ochman (3), revealing no relationship between

chiropteran phylogeny and gut bacterial community dissimilarity. We found the

same absence of phylogenetic signal among oral and skin microbial communities.

As suggested in other studies of volant vertebrates (bats and birds), convergent

adaptations driven by the evolution of flight may be influencing the nature and

composition of microbial communities in both bats and birds (28-30).

Microbial community specificity can be assessed as a function of

intraspecific variation in dissimilarity (beta dispersion), where low dispersion

suggests a tight and perhaps co-evolutionary link between hosts and symbionts,

whereas high dispersion suggests more random associations between hosts and

symbionts (31). Measures of beta dispersion among bats revealed a continuum

for all three anatomical sites, with oral bacterial commuities showing lower levels

of beta dispersion (for weighted UniFrac distances) than gut or skin communities

(Fig. 3). This continuum suggests a possible gradient of host-symbiont specificity

across different bat species that may be influenced by evolutionary history or host

ecology. Given that we found no association between bacterial community

dissimilarity and host phylogenetic distance, variation in beta dispersion is more

likely a reflection of host ecology than evolutionary history.

Similar to recent studies in North American bats (32), we found sampling

locality to be a significant factor influencing skin, gut and oral microbial

composition (Table 4). Furthermore, we observed an apparent trend in

increasing Shannon diversity and observed ESV richness along an elevational

gradient that was most pronounced for skin microbiota (Fig. S2). A positive

correlation between bacterial richness and elevation has been observed in studies

of amphibian skin (33) and montane soil, and this pattern may be the result of

climatological and other abiotic factors (*e.g.* pH) found along elevational

gradients (34, 35).

We found the general composition of gut microbiota in East African bats

to be similar to that of Neotropical bats, with Proteobacteria being the dominant

bacterial phylum present (36). Regardless of diet (insectivorous or frugivorous),

the distal bat gut is dominated by bacteria in the family Enterobacteriaceae

(Phylum: Proteobacteria), though fruit bats do have an increased relative

abundance of bacteria in the family Clostridiaceae (Phylum: Firmicutes) relative

to insectivorous bats. In their study of neotropical bats, Phillips et al. (37) noted

an increased relative abundance of Lactobacillales in frugivorous bats, and we

note a similar pattern among pteropodid fruit bats in this study, which exhibited

a slightly higher proportion of Streptococcaceae (Order: Lactobacillales) relative

to insectivorous bats. Overall, the domination of the chiropteran gut by

Proteobacteria differs markedly from other mammalian gut microbiomes, which

are generally dominated by Firmicutes (21, 38, 39).

Among most bat families, the oral microbiome was dominated by

Pasteurellaceae (Phylum: Proteobacteria), and in some cases a high relative

abundance of bacteria in the families Mycoplasmataceae (in nycterids),

Neisseriaceae (in vespertilionids and rhinonycterids), and Streptococcaceae (in

pteropodids) was also observed. Although the oral microbiome has received less

attention than that of the gut, several studies have found diverse Pasteurellaceae

and Neiserria lineages present in the oral microbiota of animals, including

domestic cats (20) and marine mammals (40). Pasteurellaceae lineages have also

recently been documented in the oral microbiota of Tasmanian devils (23, 41). In

humans, Pasteurallaceae (genera *Haemophilus* and *Aggregatibacter*) and

Neisseriaceae (genera *Neisseria*, *Kingella*, and *Eikenella*) play an important role

in the formation supragingival plaque (22). Though these bacterial groups are

present in lower proportions in other animals relative to bats, their presence in a

broad range of host taxa suggest a conserved evolutionary niche.

Our analysis identified links between ectoparasitism, malarial parasitism,

and bacterial communities on the skin and in oral cavities, respectively. Network

analyses identified consistent, stable, and species-rich clusters of bacteria on the

skin of non-ectoparasitized bats, compared to relatively disconnected and

apparently transient bacteria on the skin of bats harboring ectoparasites. This

result mirrors that found in human-mosquito interactions, in which individuals

with lower bacterial diversity on the skin are significantly more attractive to

blood-seeking mosquitoes than individuals with higher diversity (42). In humans,

skin bacteria play a known role in attracting mosquitoes via their production of

volatile organic compounds (VOCs), and studies have shown that variation in

skin microbial community composition can increase or decrease human

attractiveness to blood-seeking mosquitoes (42-44). Similar mechanisms may be

at play in the bat-ectoparasite system, particularly given the phylogenetic

proximity of hippobscoid bat parasites to mosquitoes.

Several bacterial families exhibited significant associations with presence

of ectoparasitism in bats based on DESeq analyses. Bacteria found across

multiple host families included (but were not limited to) Alcaligenaceae,

Chitinophagaceae, Flavobacteriaceae, Moraxellaceae, Mycobacteriaceae

(*Mycobacterium* spp.), and Xanthomonadaceae. In many cases, these bacterial

families were associated with parasitism in some bat families, and absence of

parasitism in others, suggesting a potential mechanism by which ectoparasites

might be distinguishing between "correct" and "incorrect" hosts. As suggested by

human-mosquito interaction studies (42, 43, 45), bacteria positively associated

with increased rates of blood-feeding dipteran host selection may be producing

VOCs on which the insects rely to identify their hosts. Bacteria that are negatively

associated with such insects may be consuming the products of the former, or

may be producing VOCs of their own that mask those of the former (suggested by

Verhulst et al. (42)). To better understand the mechanisms underlying these

correlations in wild populations, future experiments should consider including

sampling of VOCs *in vivo*.

PERMANOVA analyses identified associations between the oral

microbiome and malarial parasite prevalence among bats in the family

Miniopteridae, although these associations were less robust than those of the skin

bacteria and ectoparasitism. Upon further exploration of this potential

association, we identified a single bacterial ESV in the genus *Actinobacillus* (99%

similar to *A. porcinus* based on NCBI blastn search) as significantly reduced in

malaria-free bats (baseMean 7.61, -24.2 log2FoldChange, p = 1.7E-20). Network

analyses indicated no significant differences in connectivity or node degree

distribution (results not shown). Because no other bat groups experienced rates

of malarial parasitism adequate for statistical analyses, we were unable to explore

this relationship further. Future studies that incorporate greater sampling of

malaria-positive species may reveal more robust microbial associations, as have

been documented in numerous experiments with controlled rodent and human

malaria infections (45-47).

Although we cannot ascertain causality of differences in the microbial

composition of skin in this study, our results support the hypothesis that these

differences may provide a mechanism by which ectoparasites can locate or

distinguish hosts. Alternatively, observed differences in microbial composition

could result from microbial transfer from parasites to hosts. Given the known

effect of locality and apparent absence of host phylogenetic signal in microbial

community composition of skin, one possible explanation is that local

environmental variables play a greater role in determining host-bacteria

associations in bats. Indeed, in North America, multiple bat species have been

found to share many bacterial genera with soil and plant material (32). Thus,

local conditions and bacterial composition of bat roosts are likely playing an

important role in driving the composition of skin bacteria, and via mechanisms

similar to the camouflage hypothesis, could subsequently influence which

individuals become parasitized.

METHODS

1) Sampling

Sampling for this study was conducted from the eastern coast of Kenya to the

northern border of Uganda during August-October 2016 (Fig. 1; Table S1, S2).

Eight families and nineteen genera of bats (order: Chiroptera) were collected as

part of bird and small mammal biodiversity inventories. All sampling was

conducted in accordiance with the Field Museum of Natural History IACUC and

voucher specimens are accessioned at the Field Museum of Natural History

(Table S2). Blood samples were collected and screened for haemosporidia and

haemosporidian taxonomy was assigned using previously described molecular

methods (13). Following blood sampling, ectoparasites were removed with

forceps and placed directly into 95% EtOH; ectoparasites taxonomy was assigned

based on morphological features. For the purposes of analysis with microbiome

data, ectoparasite and malarial status were each scored separately as 1 (present)

or 0 (absent). Gut, skin, and oral samples were taken for each bat for microbial

analyses. Gut samples consisted of fecal material collected directly from the distal

end of the colon using sterilized tools, and preserved on Whatman® FTA® cards

for microbiome analyses. For oral microbiome analyses, we preserved both

buccal swabs in lN$_2$ and tongue biopsies in 95% ethanol (EtOH). Comparison of

ESV diversity obtained from paired subsets of each sample type revealed greater

diversity recovered from tongue biopsies (data not shown); tongues were

therefore used for characterization of oral microbiomes in this study. Lastly, skin

samples from five regions of the body (ear, wing membrane, tail membrane,

chest, back) were collected and pooled in 95% EtOH using sterile Integra®

Miltex® 5mm biopsy punches. The goal of sampling from five body regions was

to maximize bacterial diversity recovered from the external skin surface of each

individual. We based our storage media selections on the recent study by Song et

al. (48). Host sequencing and phylogenetic methods are described in Fig. S2.

2) Microbiome sequencing, characterization, and parasite association

DNA extractions were performed on gut, tongue, and skin samples using the

MoBio PowerSoil 96 Well Soil DNA Isolation Kit (Catalog No. 12955-4, MoBio,

Carlsbad, CA, USA). We used the standard 515f and 806r primers (49-51) to

amplify the V4 region of the 16S rRNA gene, using mitochondrial blockers to

reduce amplification of host mitochondrial DNA. Sequencing was performed

using paired-end 150 base reads on an Illumina HiSeq sequencing platform.

Following standard demultiplexing and quality filtering using the Quantative

Insights Into Microbial Ecology pipeline (QIIME2) (52) and vsearch8.1 (53),

ESVs were identified using the Deblur method (19) and taxonomy was assigned

using the Greengenes Database (May 2013 release; http://**greengenes**.lbl.gov).

Libraries containing fewer than 1000 reads were removed from further analyses.

Negative controls all contained fewer than 1000 reads and were filtered at this

step. We did not rarefy the data, based on the recommendations of McMurdie

and Holmes (54). Data were then subset for analyses according to sample type,

host genus, and locality (or some combination thereof). Site-specific analyses

were only performed for sites from which five or more individual bats were

sampled. We calculated alpha diversity for each sample type (gut, oral, skin)

using the Shannon index, and measured species richness based on actual

observed diversity. Significance of differing mean values for each diversity

calculation was determined using the Kruskal-Wallis rank sum test, followed by a

post-hoc Dunn test with bonferroni corrected p-values. Three measures of beta

diversity (unweighted UniFrac, weighted UniFrac, and Bray-Curtis) were

calculated using relative abundances of each ESV (calculated as ESV read depth

over total read depth per library). Significant drivers of communitity similarity

were identified using the ADONIS test with Bonferroni correction for multiple

comparisons using the R package Phyloseq (55). Complete code for microbiome

analyses can be found at http://github.com/hollylutz/BatMP.

3) Machine learning and network analyses

A supervised machine learning approach was used to produce random forests

(RF) for the classification of different variables. RFs were constructed using 500

decision trees and subsets of ESV data via the supervised_learning.py script

implemented in QIIME (52). We tested the ability of RFs to accurately classify 1)

anatomical site (using all data), 2) host genus (using gut, oral, or skin microbial

data separately), 3) ecotparasite status (using skin microbial data), and 4)

malarial status (using oral microbial data). RF performance was assessed by

comparing the out-of-bag estimated error (OOB) with baseline (random) error. If

the ratio of OOB to baseline error was less than or equal to two, the model was

considered to perform reasonably well, as it performed at least twice as well as

random (56). To reconstruct microbial networks for skin and oral bacterial

communities within bat family groupings (which were further sub-divided into

parasitized or non-parasitized), we utilized the R package Sparse Inverse

Covariance Estimation for Ecological Association Inference (SPIEC-EASI) (57).

All network datasets were filtered to contain only ESVs that appeared in at least

three individuals within each respective dataset. Network results produced with

SPIEC-EASI were summarized using the R packages CAVnet (58) and igraph

(59). Network stability was assessed by sequentially removing network nodes

(ordered by betweeness centrality and degree) and observing natural connectivity

(*i.e.* eigenvalue of the graph adjacency matrix) as nodes are removed. To

determine which, if any, bacterial ESVs were significantly associated with

ectoparasite or malarial prevalence, we performed analyses based on the negative

binomial distribution of ESVs relative abundance, utilizing the R package

DESeq2 (60). False discovery rate (FDR) was calculated using the Benjamini-

Hochberg method (default method in DESeq), and *p*-values were adjusted

accordingly.

ACKNOWLEDGMENTS

We thank the Kenya Wildlife Service and the Uganda Wildlife Authority for

permission to conduct research in national parks. For logistical support and

assistance in the field, we thank Mike Bartonjo of the National Museums of

Kenya, Phausia Kweyu of Karatina University, Dr. Robert Kityo, Solomon

Sebuliba, and Cissy Akoth of the Makerere University Zoological Museum, Drs.

Brian Amman, Jonathan Towner, and Rebecca Tiller of the Centers for Disease

Control and Prevention, and Lauren Lutz. We thank Neil Gottel for his knowledge

and assistance with laboratory processing of microbial samples, and other

members of the Gilbert Lab, including Alyson Yee, Cesar Cardona, Thomas

Kuntz, Drs. Bea Penalver, Melissa Dsouza, and Naseer Sangwan for the assistance

with bacterial 16s analyses.

AUTHOR CONTRIBUTIONS

H.L.L. designed the research and wrote the first draft; H.L.L., E.W.J., C.W.D., T.C.D.

analyzed data; H.L.L., P.W.W., W.B.S., J.C.K. conducted field research; J.A.G., B.D.P.

provided funding and research support; all authors interpreted results and contributed to

writing.

REFERENCES

481
482

1.   Human Microbiome Project C (2012) Structure, function and diversity of

the healthy human microbiome. *Nature* 486(7402):207-214.

2.   Thaiss CA, Zmora N, Levy M, & Elinav E (2016) The microbiome and

innate immunity. *Nature* 535(7610):65-74.

3.   Nishida AH & Ochman H (2018) Rates of gut microbiome divergence in

mammals. *Mol Ecol* 27(8):1884-1897.

4.   Moeller AH, *et al.* (2014) Rapid changes in the gut microbiome during

human evolution. *Proc Natl Acad Sci U S A* 111(46):16431-16435.

5.   Kundu P, Blacher E, Elinav E, & Pettersson S (2017) Our Gut Microbiome:

The Evolving Inner Self. *Cell* 171(7):1481-1493.

6.   Li X, *et al.* (2017) Composition of Gut Microbiota in the Gibel Carp

(Carassius auratus gibelio) Varies with Host Development. *Microb Ecol*

74(1):239-249.

7.   Kolodny 0, *et al.* (2017).

8.   McFall-Ngai M, *et al.* (2013) Animals in a bacterial world, a new

imperative for the life sciences. *Proc Natl Acad Sci U S A* 110(9):3229-

3236.

9.   Kwiatkowski DP (2005) How malaria has affected the human genome and

what human genetics can teach us about malaria. *Am. J. Hum. Genet.*

77:171-190.

10.   Dittmar K, Morse, Solo F., Dick, Carl W., Patterson, Bruce D. (2015) Bat

fly evolution from the Eocene to the present (Hippoboscoidea, Streblidae

and Nycteribiidae). *Parasite Diversity and Diversification: Evolutionary*

Ecology Meets Phylogenetics, ed S. Morand BRK, D. T. J. Littlewood

(Cambridge University Press, Cambridge, U. K.).

11.   Dick CW, Patterson, B. D. (2006) *Micromammals and Macroparasites:*

From Evolutionary Ecology to Management (Springer, Kato Bunmeisha,

Japan).

12.   Schaer J, *et al.* (2013) High diversity of West African bat malaria parasites

and a tight link with rodent Plasmodium taxa. *Proceedings of the National*

Academy of Sciences 100:17415-17419.

13.   Lutz HL*, et al.* (2016) Diverse sampling of East African haemosporidians

reveals chiropteran origin of malaria parasites in primates and rodents.

Mol Phylogenet Evol 99:7-15.

14.   Bausch DG, Schwarz, L. (2014) Outbreak of Ebola virus disease in Guinea:

where ecology meets economy. *PLoS Negl Trop Dis* 8(7):e3056.

15.   Hayman DT, Bowen, R.A., Cryan, P.M., McCracken, G.F., O'Shea, T.J.,

Peel, A.J., Gilbert, A., Webb, C.T., Wood, J.L. (2013) Ecoogy of zoonotic

infectious diseases in bats: current knowledge and future directions.

Zoonoses and Public Health 60:2-21.

16.   Towner JS*, et al.* (2009) Isolation of genetically diverse Marburg viruses

from Egyptian fruit bats. *PLoS Pathog* 5(7):e1000536.

17.   Olival KJ & Hayman DT (2014) Filoviruses in bats: current knowledge and

future directions. *Viruses* 6(4):1759-1788.

18.   Amman BR, *et al.* (2015) A Recently Discovered Pathogenic

528   Paramyxovirus, Sosuga Virus, is Present in Rousettus aegyptiacus Fruit

529   Bats at Multiple Locations in Uganda. *J Wildl Dis* 51(3):774-779.

530   19.   Amir A, *et al.* (2017) Deblur Rapidly Resolves Single-Nucleotide

531   Community Sequence Patterns. *mSystems* 2(2).

532   20.   Sturgeon A, Pinder SL, Costa MC, & Weese JS (2014) Characterization of

533   the oral microbiota of healthy cats using next-generation sequencing. *Vet J*

534   201(2):223-229.

535   21.   Ley RE, *et al.* (2008) Evolution of mammals and their gut microbes.

536   *Science* 320(5883):1647-1651.

537   22.   Mark Welch JL, Rossetti BJ, Rieken CW, Dewhirst FE, & Borisy GG (2016)

538   Biogeography of a human oral microbiome at the micron scale. *Proc Natl*

539   *Acad Sci U S A* 113(6):E791-800.

540   23.   Brix L, Hansen MJ, Kelly A, Bertelsen MF, & Bojesen AM (2015)

541   Occurrence of Pasteurellaceae Bacteria in the Oral Cavity of the

542   Tasmanian Devil (Sarcophilus Harrisii). *J Zoo Wildl Med* 46(2):241-245.

543   24.   Grice EA & Segre JA (2011) The skin microbiome. *Nat Rev Microbiol*

544   9(4):244-253.

545   25.   Ursell LK, *et al.* (2012) The interpersonal and intrapersonal diversity of

546   human-associated microbiota in key body sites. *J Allergy Clin Immunol*

547   129(5):1204-1208.

548   26.   Costello EK, *et al.* (2009) Bacterial community variation in human body

549   habitats across space and time. *Science* 326(5960):1694-1697.

550   27.   Chiarello M, Villéger, S., Bouvier, C., Bettarel, Y., Bouvier, T. (2015) High

551   diversity of skin-associated bacterial communities in marine fishes is

552   promoted by their high variability among body parts, individuals and

553   species. *FEMS Microbiol Ecol* 91(7):1-12.

554   28.   Caviedes-Vidal E, McWhorter, T. J., Lavin, S. R., Chediak, J. G., Tracy, C.

555   R., Karasov, W. H. (2007) The digestive adaptation of flying vertebrates:

556   High intestinal paracellular absorption compensates for smaller guts.

557   *PNAS* 104(48):19132-19137.

558   29.   Price ER, Brun A, Caviedes-Vidal E, & Karasov WH (2015) Digestive

559   adaptations of aerial lifestyles. *Physiology (Bethesda)* 30(1):69-78.

560   30.   Caviedes-Vidal E*, et al.* (2008) Paracellular absorption: a bat breaks the

561   mammal paradigm. *PLoS One* 3(1):e1425.

562   31.   Thomas T*, et al.* (2016) Diversity, structure and convergent evolution of

563   the global sponge microbiome. *Nat Commun* 7:11870.

564   32.   Avena CV*, et al.* (2016) Deconstructing the Bat Skin Microbiome:

565   Influences of the Host and the Environment. *Front Microbiol* 7:1753.

566   33.   Muletz Wolz CR, Yarwood SA, Campbell Grant EH, Fleischer RC, & Lips

567   KR (2018) Effects of host species and environment on the skin

568   microbiome of Plethodontid salamanders. *J Anim Ecol* 87(2):341-353.

569   34.   Bryant JA, Lamanna, C., Morlon, H., Kerkhoff, A. J., Enquist, B. J., Green,

570   J. L. (2008) Microbes and mountainsides: contrasting elevational patterns

571   of bacterial and plan diversity. *PNAS* 105:11505-11511.

572   35.   Singh D, *et al.* (2014) Strong elevational trends in soil bacterial community
573   composition on Mt. Halla, South Korea. *Soil Biology and Biochemistry*
574   68:140-149.

575   36.   Carrillo-Araujo M, *et al.* (2015) Phyllostomid bat microbiome composition
576   is associated to host phylogeny and feeding strategies. *Front Microbiol*
577   6:447.

578   37.   Phillips CD, *et al.* (2012) Microbiome analysis among bats describes
579   influences of host phylogeny, life history, physiology and geography. *Mol*
580   *Ecol* 21(11):2617-2627.

581   38.   Eckburg PB, Bik, E. M., Bernstein, C. N., Purdom, E., Dethlefsen, L.,
582   Sargent, M., Gill, S. R., Nelson, K. E., Relman, D. A. (2005) Diversity of
583   human intestinal microbial flora. *Science* 308(5728):1635-1638.

584   39.   Yildirim S, *et al.* (2010) Characterization of the fecal microbiome from
585   non-human wild primates reveals species specific microbial communities.
586   *PLoS One* 5(11):e13963.

587   40.   Bik EM, *et al.* (2016) Marine mammals harbor unique microbiotas shaped
588   by and yet distinct from the sea. *Nat Commun* 7:10516.

589   41.   Gutman N, Hansen MJ, Bertelsen MF, & Bojesen AM (2016)
590   Pasteurellaceae bacteria from the oral cavity of Tasmanian devils
591   (Sarcophilus Harrisii) show high minimum inhibitory concentration
592   values towards aminoglycosides and clindamycin. *Lett Appl Microbiol*
593   62(3):237-242.

594   42.   Verhulst NO, *et al.* (2011) Composition of human skin microbiota affects
595   attractiveness to malaria mosquitoes. *PLoS One* 6(12):e28991.

43.   Busula AO, Takken W, JG DEB, Mukabana WR, & Verhulst NO (2017)

Variation in host preferences of malaria mosquitoes is mediated by skin

bacterial volatiles. *Med Vet Entomol* 31(3):320-326.

44.   Verhulst NO, *et al.* (2009) Cultured skin microbiota attracts malaria

mosquitoes. *Malar J* 8:302.

45.   Robinson A, *et al.* (2018) Plasmodium-associated changes in human odor

attract mosquitoes. *Proc Natl Acad Sci U S A* 115(18):E4209-E4218.

46.   De Moraes CM, *et al.* (2014) Malaria-induced changes in host odors

enhance mosquito attraction. *Proc Natl Acad Sci U S A* 111(30):11079-

11084.

47.   de Boer JG, *et al.* (2017) Odours of Plasmodium falciparum-infected

participants influence mosquito-host interactions. *Sci Rep* 7(1):9283.

48.   Song SJ, Amir, A., Metcalf, J. L., Amato, K. R., Xu, Z. Z., Humphrey, G.,

Knight, R. (2016) Preservation methods differ in fecal microbiome

stability, affecting suitability for field studies. *mSystems* 1(3):1 - 12.

49.   Caporaso JG, Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C.

A., Turnbaugh, P. J., Fierer, N., Knight, R. (2011) Global patterns of 16S

rRNA diversity at a depth of millions of sequences per sample. *PNAS*

108:4516-4522.

50.   Caporaso JG, *et al.* (2012) Ultra-high-throughput microbial community

analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 6(8):1621-

1624.

51.   Kozich JJ, Westcott, S. L., Baxter, N. T., Highlander, S. K., Schloss, P. D.

(2013) Development of a dual-index sequencing strategy and curation

pipeline for analyzing amplicon sequence data on teh MiSeq Illumina

sequencing platform. *Applied and Environmental Microbiology*

79(17):5122-5120.

52.   Caporaso JG, Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D.,

624   Costello, E. K., Fierer, N., Gonzalez Peña, A., Goodrich, E. K., Gordon, J. I.,

625   Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E.,

626   Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J.,

627   Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J.,

628   Yatsunenko, T., Zaneveld, J., Knight, R. (2010) QIIME allows analysis of

629   high-throughput community sequencing data. *Nature Methods* 7(5):335 -

630   336.

631   53.   Rognes T, Flouri T, Nichols B, Quince C, & Mahe F (2016) VSEARCH: a

632   versatile open source tool for metagenomics. *PeerJ* 4:e2584.

633   54.   McMurdie PJ, Holmes, S. (2014) Waste not, want not: Why rarefying

634   microbiome data is inadmissable. *PLoS Comput Biol* 10(4):1-12.

635   55.   McMurdie PJ & Holmes S (2013) phyloseq: an R package for reproducible

636   interactive analysis and graphics of microbiome census data. *PLoS One*

637   8(4):e61217.

638   56.   Breiman L (2001) Random Forests. *Machine Learning* 45:5-32.

639   57.   Kurtz ZD, *et al.* (2015) Sparse and compositionally robust inference of

640   microbial ecological networks. *PLoS Comput Biol* 11(5):e1004226.

641   58.   Cardona C (2017) CAVNet: Creation Analysis and Visualization of

642   Networks).

643   59.   Csardi G, Nepusz, T. (2006) The Igraph Software Package for Complex

644   Network Research. *InterJournal, Complex Systems* 1695.

645   60.   Love MI, Huber W, & Anders S (2014) Moderated estimation of fold

646   change and dispersion for RNA-seq data with DESeq2. *Genome Biol*

647   15(12):550.

648   61.   Teeling EC, Springer, M. S., Madsen, O., Bates, P., O'Brien, S. J., Murphy,

649   W. J. (2005) A molecular phylogeny for bats illuminates biogeography and

650   fossil record. *Science* 307:580-584.

FIGURE LEGENDS

Figure 1. Sampling localities and elevation, grouped by district. Colors

correspond to elevation, and white numbers and size of points correspond to

number of bats collected.

Figure 2. Alpha diversity of Exact Sequence Variants (ESVs) by anatomical sites,

including (A) Observed richness, (B) Shannon index of diversity, (C) ESVs shared

between anatomical sites. Asterisks indicate significant differences between

groups (Dunn's Test, Bonferroni corrected p-value $p < 0.0001$).

Figure 3. Intraspecific variation across anatomical sites measured as beta

dispersion of weighted UniFrac distances. Dotted lines indicate mean dispersion

for anatomical groupings; numbers in parentheses indicate sample size per bat

species. White and gray boxes correspond to the chiropteran suborders

Yangochiroptera (microbats) and Yinpterochiroptera (fruits bats and kin),

respectively.

Figure 4. (A) Relative abundance of top 6 bacterial phyla grouped by anatomical

site, with each bar corresponding to individual libraries. (B) Relative abundance

of the most prevelant eight bacterial families across all anatomical sites, grouped

by bat family. Phylogeny based on Teeling et al. (61).

Figure 5. (A) Distribution of skin microbial network clusters for parasitized and

non-parasitized bats, grouped by bat family (* indicates signifiance at $p < 0.005$,

Kruskal-Wallis) (B) Visualization of skin bacterial networks (based on

Fruchterman-Reingold algorithm); colored nodes correspond to unique clusters

of co-occurring ESVs within each network.

Figure 6. Log2fold change in relative abundance of skin-associated ESVs from the

four most-sampled bat families. ESVs shown were found to be significantly

associated with ectoparasite status in bats based on analysis of negative binomial

distributions of relative abundance (Banjamini-Hochberg FDR corrected p-value

$p < 0.05$). Positive values correspond to ESVs found to be enriched on parasitized

bats, and negative values correspond to ESVs found to be enriched on non-

parasitized bats. Gray bars highlight ESVs in bacterial families that were enriched

in parasitized bats for three out of four bat families.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

Distance of samples from group centroid

FIGURE 5

FIGURE 6

Table 1. Bat sampling, ectoparasite prevalence (n_{ecto}), and malarial parasite prevalence (n_{haem}) and identification.

Bat family	Bat species	n_{bats}	n_{ecto} (%)	n_{haem} (%)
Emballonuridae	*Coleura afra*	11	2 (18)	0
Hipposideridae	*Hipposideros caffer*	47	18 (38)	0
	Hipposideros camerunensis	2	0	0
	Hipposideros ruber	21	16 (76)	0
	Macronycteris vittatus	10	0	0
Miniopteridae	*Miniopterus africanus*	22	13 (59)	11 (50)
	Miniopterus natalensis	54	16 (30)	13 (24)
	Miniopterus rufus	22	20 (61)	20 (91)
	Miniopterus minor	18	16 (89)	12 (67)
Molossidae	*Chaerephon bivittatus*	14	0	0
	Otomops harrisoni	33	1 (3)	0
Nycteridae	*Nycteris arge*	3	0	0
	Nycteris thebaica	7	1 (14)	0
	Nycteris sp.	6	0	0
Pteropodidae	*Epomophorus labiatus*	2	0	0
	Epomophorus wahlbergi	11	0	3 (27)
	Micropteropus pusillus	4	0	0
	Myonycteris angolensis	4	0	0
	Rousettus aegyptiacus	48	24 (50)	0
	Stenonycteris lanosus	37	23 (62)	0
Rhinolophidae	*Rhinolophus clivosus*	43	8 (19)	0
	Rhinolophus eloquens	24	19 (79)	0
	Rhinolophus hildebrandti	4	1 (25)	0
	Rhinolophus landeri	14	0	3 (21)
	Rhinolophus sp.	3	0	
Rhinonycteridae	*Triaenops afer*	10	6 (60)	0
Vespertilionidae	*Myotis tricolor*	9	8 (89)	3 (33)
	Neoromicia nana	1	0	0
	Neoromicia sp.	3	0	
	Pipistrellus sp.	1	0	0
	Scotoecus hindei	4	1 (25)	0
	Scotophilus dingani	3	0	0
Total		495	193	65

Table 2. Alpha diversity of microbial communities across anatomical sites within each host genus, measured by Shannon Index of diversity (S-I) and observed sOTU richness (obs); n corresponds to number of libraries included in each calculation (following quality filtering).

| Host Family | Host Genus | Fecal | | | Oral | | | Skin | | |
		S-I	obs	n_{fecal}	S-I	obs	n_{oral}	S-I	obs	n_{skin}
Emballonuridae	*Chaerephon*	1.16	52	12	1.39	57	14	3.57	547	14
Hipposideridae	*Hipposideros*	1.70	79	65	2.01	155	52	4.95	439	74
	Macronycteris	1.82	74	9	2.12	110	9	4.94	883	7
Miniopteridae	*Miniopterus*	1.41	70	92	1.55	87	74	4.12	403	114
Molossidae	*Coleura*	1.59	52	11	0.38	41	11	4.01	566	11
	Otomops	0.88	53	26	0.35	22	26	3.88	288	33
Nycteridae	*Nycteris*	1.60	80	10	1.62	78	14	4.48	807	14
Pteropodidae	*Epomophorus*	1.44	49	11	1.42	46	11	3.78	566	13
	Micropteropus	1.90	39	3	2.21	39	4	2.30	84	3
	Myonycteris	1.14	117	4	1.29	195	5	5.21	1246	4
	Rousettus	1.62	93	32	1.95	84	34	4.90	1207	34
	Stenonycteris	1.55	61	41	1.72	97	38	4.59	855	33
Rhinolophidae	*Rhinolophus*	1.34	62	58	1.95	81	59	4.71	543	79
Rhinonycteridae	*Triaenops*	1.69	82	9	1.28	414	9	4.03	508	10
Vespertilionidae	*Myotis*	1.62	54	1	1.33	72	6	5.41	771	3
	Neoromicia	2.13	65	4	1.47	37	4	3.76	267	4
	Pipistrellus	1.05	NA	1	NA	NA	0	4.80	360	2
	Scotoecus	1.86	92	4	1.97	17	3	4.20	360	4
	Scotophilus	1.23	64	3	0.38	96	1	4.08	459	2
Mean		1.51	69	n_{fecal} 396	1.47	96	n_{orall} 375	4.30	587	n_{skin} 458

Table 3. Nonparametric permutational multivariate analysis of variance using distance matrices (via ADONIS), with distance matrices among sources of variation partitioned by host taxonomy (species nested within genus), ectoparasite status, malarial infection status, and locality included as strata to constrain permutation across this variable; * indicates p-value < 0.05.

Site	Partition Variable	Weighted UniFrac			Unweighted UniFrac			Bray-Curtis		
		F	R2	Pr(>F)	F	R2	Pr(>F)	F	R2	Pr(>F)
Fecal	(Host genus (species))	4.27	0.162	0.001*	3.15	0.120	0.001*	2.89	0.110	0.001*
	Ectoparasite status	0.47	0.001	0.912	1.42	0.004	0.048*	1.40	0.004	0.097
	Malarial status	1.34	0.004	0.21	1.33	0.004	0.077	1.98	0.005	0.011*
Oral	(Host genus (species))	6.82	0.279	0.001*	3.50	0.143	0.001*	6.69	0.274	0.001*
	Ectoparasite status	0.51	0.001	0.836	1.41	0.004	0.057	1.00	0.003	0.447
	Malarial status	2.78	0.008	0.015*	1.17	0.003	0.2	1.98	0.006	0.019*
Skin	(Host genus (species))	7.68	0.329	0.001*	3.98	0.170	0.001*	5.60	0.240	0.001*
	Ectoparasite status	2.42	0.006	0.01*	1.54	0.004	0.02*	2.07	0.005	0.001*
	Malarial status	0.92	0.002	0.513	1.02	0.002	0.363	1.06	0.003	0.32

Table 4. Permutational multivariate analysis of variance using distance matrices, with distance matrices among sources of variation partitioned by host taxonomy (species nested within genus), locality, and anatomical site.

Partition Variable	Weighted UniFrac			Unweighted UniFrac			Bray-Curtis		
	SumSq	F	Pr(>F)	SumSq	F	Pr(>F)	SumSq	F	Pr(>F)
Anatomical site	10.67	198.01	0.001*	56.52	82.90	0.001*	38.2	36.97	0.001*
Host Genus	3.77	13.09	0.001*	25.54	7.02	0.001*	85.30	15.06	0.001*
Locality	1.56	11.00	0.001*	20.62	11.34	0.001*	23.85	8.42	0.001*
Host Genus:species	1.39	4.08	0.001*	11.20	2.59	0.001*	25.25	1.33	0.001*

Table 5. Supervised machine learning results, showing random forest model performance with respect to different classification variables and input data sets (fecal, oral, skin microbiome). Model performance is assessed by measuring the ratio of Out-of-bag estimated error (OOB) to baseline error.

Classification variable	Input Data	Baseline error	OOB error	Baseline:OOB
Anatomical site	All data	0.68	0.14	4.8
Host Genus	Skin	0.75	0.17	4.3
Host Genus	Oral	0.78	0.24	3.2
Host Genus	Gut	0.77	0.35	2.2
Ectoparasite Status	Skin	0.53	0.27	2.0
Malarial Status (Miniopteridae only)	Oral	0.46	0.38	1.2