1   Title

2   The fungal-specific Hda2 and Hda3 proteins regulate morphological switches

3   in the human fungal pathogen *Candida albicans*.

Running title:

Hda2 and Hda3 control *C. albicans* morphogenesis

Misty R. Peterson[a], Robert Jordan Price[a], Sarah Gourlay[a], Alisha May[a*], Jennifer

Tullet[a] and Alessia Buscaino[a#]

[a] : Kent Fungal Group, School of Biosciences, University of Kent, Canterbury, United

Kingdom

[#]Address correspondence to Alessia Buscaino, A.Buscaino@kent.ac.uk

[*:] Present address: Department of Life Sciences, Imperial College London, South

Kensington Campus, London, SW7 2AZ

Word Count Main Text: 3902

Word Count Material and Methods: 1880

ABSTRACT

The human fungal pathogen *Candida albicans* is responsible for millions of infections

annually. Due to the few available anti-fungal drugs and the increasing incidence of

drug resistance, the number of *C. albicans* infections is dramatically increasing.

Morphological switches, such as the white-opaque switch and the yeast-hyphae

switch, are key for the development of *C. albicans* pathogenic traits. Lysine

deacetylases are emerging as important regulators of morphological switches. Yet,

targeting lysine deacetylases for drug development is problematic due to the high

homology between the fungal and human proteins that could result in toxicity. Here

we provide evidence that the fungal specific proteins Hda2 and Hda3 interact with

the lysine deacetylase Hda1. By combining phenotypic analyses with genome-wide

transcriptome analyses, we demonstrate that Hda2 and Hda3 control *C. albicans*

morphological switches. Under nutrient-rich conditions, deletion of *HDA2* or *HDA3*

leads to moderate overexpression of the master regulator of white-opaque switching

WOR1 and increase switching frequency. Under hyphae inducing conditions,

deletion of *HDA2* and *HDA3* block hyphae development. However, deletion of *HDA2*

and *HDA3* does not affect hyphae-formation and virulence *in vivo*. We propose that

Hda2 and Hda3 are good targets for the development of anti-fungal drugs to be used

in combination therapy.

KEYWORDS

INTRODUCTION

Fungal pathogens are a leading cause of human mortality causing over 1.5 million

deaths per year (1). *C. albicans* is a commensal organism that colonises the mouth,

gastrointestinal and reproductive tract of healthy individuals without causing any

harm. Still, *Candida albicans* is also the most common human fungal pathogen and

the principal causal agent of mycotic death (2). This is because, in immune-

compromised patients, *C. albicans* can invade vital organs and cause serious, life-

threating systemic infections associated with a mortality rate up to 70 % (2). The

ability to transition between different morphological forms in response to changing

environments is a key virulence trait in *C. albicans*.

For example, *C. albicans* cells can reversibly switch between white and opaque

forms (3). White and opaque cells are genetically identical, yet they differ in cellular

morphology, colony shape, gene expression profile and mating behaviour (4). In

addition, white cells are more virulent in a murine model of systemic infection (5, 6)

whereas opaque cells preferentially colonise the skin (7). White-opaque switching is

under the control of the master regulator Wor1, a transcription factor whose

expression is necessary and sufficient for opaque cell formation (8–11). Stochastic

increases in Wor1 levels drives the transition from the white to the opaque phase.

Furthermore, Wor1 expression produces a direct positive feedback loop by binding

its own promoter and turning on its own expression (8, 9, 11). Switching is also

regulated by the mating type locus as opaque formation occurs predominantly in a or

α cells (12).

C. albicans virulence also depends on its ability to convert between yeast and hyphal

morphology: yeast cells are critical for colonisation, early infection and

dissemination, while hyphal growth is responsible for tissue invasion and chronic

infections (13). Hyphal morphogenesis is coupled with virulence, as several of the

genes, that are specifically expressed in hyphae, encode virulence factors (14–17).

Hyphal morphogenesis is a complex and highly orchestrated process, and *C.*

albicans uses multiple redundant pathways to integrate host signals and promote

hyphae development. Indeed, *C. albicans* filamentation can be induced by many

environmental cues such mammalian serum, body temperature, hypoxia and CO_2

concentration which reflects the variety of signals sensed by the fungus in the

different microenvironments encountered in the host. In yeast cells, hyphae

morphogenesis is inhibited by the DNA-binding repressor Nrg1 that, together with

Tup1, blocks expression of a subset of filament-specific genes (18). Upon hyphal

induction, filamentous growth is promoted by Efg1 and Flo8, two transcription

regulators essential for hyphal development and virulence [14–16]. During the

initiation phase of hyphae morphogenesis, Nrg1-mediated repression is cleared via

NRG1 transcriptional repression and Nrg1 protein degradation. During hyphae

maintenance, chromatin remodelling of hyphae promoters prevents Nrg1 binding

despite its increased protein levels (19).

Due to the close evolutionary relationship between fungi and the human host,

effective treatment for *C. albicans* infections is hindered by the limited number of

sufficiently divergent potential drug targets. There are only three classes of

antifungal drugs effective for the treatment of systemic fungal infections and their

clinical utility is limited by the rapid emergence of drug resistance (1, 20). Hence,

there is an immediate and urgent need to develop alternative treatments. A potential

strategy is to target *C. albicans* morphological plasticity.

Lysine deacetylases (KDACs, also known as HDACs) act as global regulators of

gene expression by catalysing the removal of acetyl functional groups from the lysine

residues of histones and non-histone proteins (21). KDACs can favour transcriptional

repression by deacetylating lysine residues on histone tails allowing chromatin

compaction and/or preventing binding of bromodomain-containing transcriptional

activators (21). KDACs can also activate transcription by deacetylation of non-

histone proteins (22). As a consequence, deletion or inhibition of KDACs often

results in the upregulation and downregulation of an approximately equivalent

number of genes (23). KDACs are highly conserved across eukaryotes and can be

phylogenetically divided into three main classes: the Rpd3 and Hos2-like (class I)

enzymes, the Hda1-like (class II) enzymes and the Sir2-like (class III) enzymes. The

class I and class II enzymes are related, sharing a conserved central enzymatic

domain. The class III enzymes are nicotinamide adenine dinucleotide (NAD)

dependent. KDACs lack intrinsic DNA-binding activity and are recruited to target

genes via incorporation into large multiprotein complexes or direct association with

transcriptional activators and repressors (24).

In *C. albicans*, the lysine deacetylase Hda1 (class II) is an important regulator of

morphological switches. Hda1 controls white-opaque switching as deletion of the

HDA1 gene increases switching rates from white to opaque (25, 26). In response to

serum, N-acetylglucosamine or nutrient limitation, Hda1 also controls the yeast to

hyphae switch by deacetylating Yng2, a component of the histone acetyltransferase

NuA4 complex, blocking maintenance of hyphal growth (19). However, the Hda1

pathway is not required for hyphae elongation in hypoxia or in the presence of

elevated CO_2 because of the presence of redundant pathways (27, 28). As a result,

the Hda1-mediated hyphae maintenance pathway contributes, but it is not absolutely

required, for virulence *in vivo* (27). These results suggest that Hda1 is a good target

for antifungal drugs development to be used in combination therapies.

KDACs are promising druggable targets: one KDAC inhibitor is currently used for

cancer treatment and several other KDAC inhibitors are in clinical trials (29, 30).

However effective targeting of Hda1 for anti-fungal drug development is impaired by

the high sequence similarities between Hda1 and its human orthologs and

consequently a likeliness for high toxicity.

In *Saccharomyces cerevisiae*, Hda1 assembles with two non-catalytic subunits,

Hda2 and Hda3, essential for Hda1 deacetylation activity both *in vivo* and *in vitro*

(31). Interestingly, no metazoan homologous of Hda2 and Hda3 have been

identified. Hda2 and Hda3 are similar in sequence and share a similar protein

organisation with an N-terminal DNA binding domain (DBD) and a C-terminal coil-coil

domain (CCD). The DBD domain, similar in structure to the helicase domain of the

SWI-SNF chromatin remodellers, is sufficient to bind DNA *in vitro*. The CCD domains

act as a scaffold for the assembly of the Hda1 complex (32).

Here for the first time, we analyse the role of *C. albicans* Hda2 and Hda3. We

demonstrate that the Hda1 complex is conserved in *C. albicans* as Hda2 and Hda3

interact with Hda1 *in vivo*. Our analyses demonstrated that, in yeast-inducing

conditions, deletion of *HDA2* and *HDA3* leads to transcriptional upregulation of

WOR1 and that Hda2 and Hda3 inhibit white and opaque switching. In contrast,

under hyphae-inducing conditions, Hda2 and Hda3 regulate the yeast to hyphae

switch. This functional rewiring is linked to a reduced expression of components of

the Hda1 complex. Our study demonstrates that Hda2 and Hda3 control key

morphological switches in *C. albicans*. Therefore, we propose that Hda2 and Hda3

are attractive targets for the development of novel antifungal drugs.

RESULTS

The Hda1 complex is conserved in *C. albicans*

S. cerevisiae Hda2 and Hda3 are critical components of the Hda1 complex as they

are required for the Hda1-mediated histone deacetylation activity (31). BLAST

analyses reveal that the *C. albicans* genome contains two genes encoding proteins

with homology to *S. cerevisiae* Hda2 (C3_03670W) and Hda3 (CR_09490W_A).

Structural alignments predict that Hda2 and Hda3 have a similar protein organisation

with an N-terminal DNA binding domain (DBD) and a C-terminal coil-coil domain

(CCD) (Fig 1A and 1B). To explore the potential functional relationship between *C.

albicans* Hda2, Hda3 and Hda1, we assessed whether we could detect a physical

interaction between these proteins. To this end, we generated strains expressing at

the endogenous locus an epitope-tagged Hda1 protein (Hda1-HA) together with

either Hda2-GFP or Hda3-GFP. Western analyses show that the Hda1-HA and

Hda2-GFP tagged proteins are expressed at high levels while we could not detect

Hda3-GFP in whole cell extract indicating that the protein is expressed at low levels

(Fig 1C). Immuno-precipitation (Ip) of Hda1-HA with a highly specific anti-HA

antibody demonstrates that Hda2 strongly interacts with Hda1. We could also detect

an interaction between Hda1 and Hda3, despite the low level of expression of Hda3

(Fig 1C). Thus, Hda2 and Hda3 physically interact with Hda1. To delineate the

function of *C. albicans* Hda2 and Hda3, we generated homozygous deletions

mutants for *HDA2* (*hda2 Δ/Δ*) and *HDA3* (*hda3 Δ/Δ*) genes using a recyclable Clox

system (Fig 1D) (33). The Clox system allows for the generation of a homozygous

deletion mutant lacking any marker genes and therefore genetically identical to the

parental strain except for the deleted gene. This allows for the direct comparison of

phenotypes between mutant and parental strains. Growth analyses reveal that Hda2

and Hda3 are not required for survival and fitness as the *hda2 Δ/Δ* and *hda3 Δ/Δ*

strains grow similarly to the wild-type (WT) control both on solid and in liquid media

(Fig 1E and 1F). Therefore, *C. albicans* Hda2 and Hda3 are *bonafide* components of

the Hda1 complex that are not required for cell fitness and survival under optimal

growth conditions.

Global gene expression changes in the absence of Hda2 and Hda3

To gain insights about the function of Hda2 and Hda3 in conditions stimulating yeast

growth, we analysed changes in gene expression by strand-specific RNA-

sequencing (RNA-seq) upon deletion of the *HDA2* and *HDA3* genes. As a

comparison, we also performed RNA-seq on strains with the *HDA1* gene deleted.

Deletion of *HDA2* results in 577 genes significantly differentially regulated (348

upregulated and 229 downregulated, q < 0.05; Dataset S1). These gene expression

changes are highly similar to the ones observed in *hda1 Δ/Δ* (Pearson correlation

coefficient r = 0.7, Dataset S1) (Fig 2A). Deletion of the *HDA3* gene has a less

profound effect on gene expression as only 78 genes are significantly differentially

expressed (45 upregulated and 33 downregulated, q < 0.05; Dataset S1). Despite

this difference, changes in *hda3 Δ/Δ* cells correlate with *hda1 Δ/Δ* and *hda2 Δ/Δ* gene

expression changes (Pearson correlation coefficient r = 0.66 and 0.46, respectively)

(Fig 2A).

To reveal the cellular pathways regulated by Hda2 and Hda3 in *C. albicans*, we

performed Gene Set Enrichment Analysis (GSEA) of the RNA-seq datasets (34). To

this end, transcript profiles of *hda2 Δ/Δ* or *hda3 Δ/Δ* isolates were ranked according

to their differential expression and this list was compared to gene sets identified in

other experimental analyses (35) and (Sellam et al, personal communications). This

allows the identification of statistically significant gene sets enriched in the top (up-

regulated) or the bottom (down-regulated genes) of the ranked list (34). The network

of similar gene sets was visualized using Cytoscape where nodes represent gene

sets, and lines connect nodes sharing a significant number of genes (36).

GSEA detected enrichment for rRNA and ribosome biogenesis genes and genes

involved in transport (Fig 2B). Enrichment was also found in gene sets important for

virulence-promoting function in *C. albicans*. This includes genes differentially

expressed during *C. albicans*-host interactions with mouse macrophages and in

response to drugs (Fig 2C).

Modulation of several gene sets enriched in *hda2 Δ/Δ* and/or *hda3 Δ/Δ* mutants, for

example, down-regulation of genes required for protein synthesis, are part of stress

response in *C. albicans* (37). Accordingly, we subjected the *hda2 Δ/Δ* and *hda3 Δ/Δ*

mutants to a phenotypic analysis by applying a set of different stress conditions (38).

For most conditions, we did not observe any difference between the WT and the

mutant strains. However, lack of Hda2 or Hda3 leads to sensitivity to copper, sodium

chloride and sodium nitroprusside (SNP) with salicylhydroxamic acid (SHAM).

Strains mutant in these proteins also show resistance to rapamycin (Fig 3) indicating

a role for Hda2 and Hda3 in specific stress responses.

211   Hda2 and Hda3 inhibit white-opaque switching

The GSEA analysis identifies white-opaque switching as a process potentially

regulated by Hda2 and Hda3 as genes upregulated in the opaque state are also

upregulated upon deletion of *HDA2* and *HDA3* (Fig 4A and 4B). Wor1 is the master

regulator of white-opaque phenotypic switching and a stochastic increase in Wor1

levels drives the transition from the white to the opaque phase (8, 9, 11). Therefore,

we asked whether *WOR1* expression levels were increased in *hda2 Δ/Δ* and *hda3*

Δ/Δ isolates compared to WT cells.

The RNA sequencing analysis demonstrates that deletion of *HDA2* or *HDA3* leads to

transcriptional upregulation of *WOR1* (Fig 4C). This result suggests that, in WT cells,

Hda2 and Hda3 repress *WOR1* transcription and inhibit white-opaque switching. To

test this hypothesis, we generated *MTL a/a* homozygous *hda2 Δ/Δ* and *hda3 Δ/Δ*

mutants and measured the frequency of white to opaque conversion using

quantitative switching assays. Briefly, cells were plated on Phloxine B plates and the

frequency of opaque colonies or colonies containing at least one opaque sector was

scored. This analysis demonstrates that deletion of both *HDA2* and *HDA3* increases

the frequency of white-opaque switching (Fig 4D). Thus, Hda2 and Hda3 control

white-opaque switching, a process that is linked to *WOR1* overexpression.

Hda2 and Hda3 contribute to filamentous growth but not virulence

To investigate the role of Hda2 and Hda3 under hyphae-inducing conditions, we

performed RNA-seq analyses of WT, *hda2 Δ/Δ* and *hda3 Δ/Δ* strains grown in RPMI,

a medium which mimics human physiological conditions and therefore strongly

induces hyphal growth. As a control, RNA-seq analyses were also performed in *hda1*

Δ/Δ strain. GSEA analysis of the gene expression changes of WT cells grown in

yeast-inducing conditions (YPD) and hyphae-inducing conditions (RPMI) confirmed

the validity of our experimental approach as genes reported to be upregulated in

hyphae compared to yeast are less expressed in YPD compared to RPMI while

genes with a yeast-specific expression are expressed at higher levels in YPD than

RPMI (Fig 5A).

In hyphae-inducing conditions, deletions of *HDA2* and *HDA3* genes result in 350 and

484 significantly differentially expressed genes, respectively (q< 0.05; Dataset S1).

Gene expression changes in *hda2 Δ/Δ* and *hda3 Δ/Δ* correlate with gene expression

changes observed in *hda1 Δ/Δ* although less profoundly than in yeast-inducing

conditions (Pearson correlation coefficient, r = 0.55 and 0.38, respectively; Fig 5B).

The RNA-seq analysis reveals that filamentation is the major pathway misexpressed

in the deletion mutants as hyphae-responsive genes are differentially expressed in

hda2 Δ/Δ and *hda3 Δ/Δ* isolates compared to WT cells (Fig 5C). qRT-PCR analyses

of two hyphal growth markers, *HWP1* and *ALS3*, confirmed the RNA-seq results as,

in hyphae inducing conditions, expression of *HWP1* and *ALS3* is downregulated in

hda2 Δ/Δ and *hda3 Δ/Δ* but not in WT cells (Fig 5D).

Collectively, our results suggest that Hda2 and Hda3 are important for hyphal

growth. To test this hypothesis, we assessed the morphology of WT, *hda2 Δ/Δ* and

hda3 Δ/Δ strains upon growth in two different hyphae-inducing media (RPMI and

Spider). While WT cells form hyphae efficiently, hyphal growth is defective in *hda2*

Δ/Δ and *hda3 Δ/Δ* cells on both solid and in liquid media (Fig 6A and B). This

phenotype was rescued by reintroduction of the *HDA3* gene (Fig 6C). Therefore,

Hda2 and Hda3 are important regulators of the yeast to hyphae switch.

To test whether the hyphae-defective phenotype observed in *hda2 Δ/Δ* and *hda3 Δ/Δ*

isolates is sufficient to impair hyphal growth *in vivo*, we performed killing assays

using the nematode *Caenorhabditis elegans* as an infection system (39, 40). These

analyses revealed no differences in hyphae formation or percentage of killing

between WT, *hda2 Δ/Δ* and *hda3 Δ/Δ* strains (Fig 6D). Therefore, although Hda2 and

Hda3 are critical for hyphae-formation in specific media, additional redundant

pathways can compensate for the lack of Hda2 and Hda3 in a more complex *in vivo*

situation.

Stability of Hda1 protein is regulated by environmental changes

Comparison of the gene expression profiles in yeast-promoting conditions (YPD 30

^{o}C) and hyphae-promoting conditions (RPMI 37 ^{o}C) reveals that deletion of *HDA1*

leads to different transcriptional changes in different environments (Fig 7A). In yeast

inducing conditions, 2046 genes are significantly differentially expressed upon

deletion of Hda1 compared to WT cells, suggesting that Hda1 acts as a global

regulator of gene expression (Fig 7A and Dataset S1). In contrast, in hyphae

inducing conditions, 507 genes are significantly differentially expressed with hyphal

growth being one of the major pathways that is altered (Fig 7B and Dataset S1).

These results suggest that Hda1 function is rewired during the yeast to hyphae

switch. We hypothesise that this change in function could be due to a differential

gene expression profile of *HDA1* in yeast compared to hyphae. While analysis of the

RNA-seq dataset reveals that *HDA1*, *HDA2* and *HDA3* transcription levels are similar

across these two conditions, Western analyses clearly demonstrate that levels of

Hda1 and Hda3 proteins, but not Hda2, are lower in hyphal cells compared to yeast

cells (Fig 7C). Therefore, the functional rewiring of the Hda1 complex upon an

environmental change is linked to a reduced expression of this subunit.

DISCUSSION

The role of the Hda1 complex in regulating morphological switches in *C.

albicans*

Results presented in this study show that the fungal-specific proteins Hda2 and

Hda3 are important regulators of morphological switches in *C. albicans*. We propose

that this regulation is mediated via the Hda1 complex. This hypothesis is supported

by our observation that Hda1, Hda2 and Hda3 physically interact and by the

published results demonstrating that Hda1 controls both the white-opaque and the

yeast-hyphae morphological switches (19, 25, 26). Based on the data presented

here, it is possible that Hda2 and Hda3 control the activity of the Hda1 complex by

different non-mutually exclusive mechanisms. First, Hda2 and Hda3 could mediate

Hda1 recruitment to target sites regulating their chromatin acetylation state and their

transcriptional activity. Indeed, in *S. cerevisiae*, Hda2 and Hda3 DNA-binding

domains are sufficient to bind DNA *in vitro* (32). The structural alignment presented

in this study demonstrates that, similar to *S. cerevisiae*, *C. albicans* Hda2 and Hda3

contain a DNA binding domains with structure resembling the helicase fold found in

the SWI2/SNF2-type chromatin-remodeling ATPase (32). We hypothesise that Hda2

and Hda3 could target Hda1 to key genomic locations leading to transcriptional

downregulation of associated genes. In support of this hypothesis, we have found

that *WOR1*, the master regulator of white-opaque switching, is moderately

upregulated in cells deleted for *HDA2* and *HDA3* genes. A similar upregulation is

observed in *hda1 Δ/Δ* isolates. We propose that this upregulation is sufficient to

activate the Wor1 positive feedback loop promoting white-opaque switching. Indeed,

hda2 Δ/Δ and hda3 *Δ/Δ* cells undergo white-opaque switching more frequently than

WT cells.

Alternatively, interactions between Hda1, Hda2 and Hda3 may induce a

conformational change promoting Hda1 deacetylase activity. In support of this

hypothesis, it has been established that the *in vitro* catalytic activity of *S. cerevisiae*

Hda1 depends on Hda2 and Hda3 (31). This regulation could be critical for

controlling deacetylation of non-histone substrates. For example, *C. albicans* Hda1

controls the yeast to hyphae switch by deacetylating a non-histone substrate, the

NuA4 component Yng2 (19). We hypothesise that interaction between Hda1, Hda2

and Hda3 causes a conformational change in Hda1 allowing deacetylation of Yng2.

Chromatin modifiers are ideal sensors of changing environments as they can

respond to external stimuli by rapidly and reversibly changing the transcriptional

state of many genes simultaneously. Our results indicate that the activity of the Hda1

complex is rewired in different environmental conditions. While in yeast cells, Hda1

acts as a globular regulator of gene expression, in hyphae-inducing conditions Hda1

function is dedicated to the yeast-hyphae switch. This functional rewiring is

accompanied by changes in the protein levels of Hda1 and Hda3. These findings

suggest a model through which hyphae-specific function is achieved by diminishing

the concentration of key proteins of the Hda1 complex. We did not observe any

changes in the RNA levels of *HDA1* and *HDA3* and therefore we hypothesise that

translation efficiency and/or stability of these two proteins is differentially regulated in

different environments.

Chromatin modifiers are often embedded in multiprotein complexes associated with

several non-catalytic subunits that regulate their targeting to substrates or their

catalytic activity (24). Our results highlight how Hda2 and Hda3, the non-catalytic

subunits of the Hda1 complex, play important roles in regulating morphological

switches in response to environmental changes.

Hda2 and Hda3 as potential targets for anti-fungal therapy

The yeast to hyphae switch is central to *C. albicans* virulence and pathogenesis. Our

results signify that while Hda2 and Hda3 promote hyphae formation under specific

growth conditions (RPMI and Spider media at 37 ^{o}C), they are dispensable for

hyphae formation and virulence in the *C. elegans* infection system. We hypothesise

that this is due to the presence of redundant pathways that stimulate hyphae

formation. Indeed, it has been shown that hyphae induction is a highly orchestrated

process integrating several different environmental signals (13). The finding that, *in*

vivo, lack of Hda2 and Hda3 does not impair hyphae formation is in agreement with

the observation that strains mutant for the *YNG2* gene, the target of Hda1, also do

not impair hyphae growth *in vivo* while they are defective for hyphae formation *in*

vitro (27).

There is an urgent need to develop new anti-fungal drugs due to emergence of

fungal strains resistant to currently available drugs. We propose that Hda2 and Hda3

could be targets for novel antifungal drugs to be used in combination therapy.

Combination therapy offers several advantages compared to single-drug therapy.

This is because it allows for widening of the spectrum and potency of drug activity

and it can also lead to reduction in the dosage of individual agents preventing

emergence of antifungal resistance (41). Several features make Hda2 and Hda3

attractive targets for antifungal drug development. First and most importantly, the

proteins are present in fungi but absent in humans minimising potential toxicity.

Secondly, drugs targeting Hda2 and Hda3 could have a broad spectrum of activity

against a variety of human fungal pathogens as Hda2 and Hda3 orthologs are

present in other human fungal pathogens, such as *C. glabrata* (CAGL0H01331g and

CAGL0G09867g) and *A. fumigatus* (Hda2: Afu5g03390).

KDACs are emerging as promising candidates for drug development and several

small molecules inhibiting KDAC activity are currently in clinical trials as potential

anti-cancer therapeutics. We propose that inhibition of Hda2 and Hda3 will impair *C.*

albicans Hda1 activity but not human KDACs reducing the risk of toxicity.

MATERIAL AND METHODS

Growth conditions

Strains are listed in Table 1. Yeast cells were cultured in rich medium (YPD)

containing extra adenine (0.1 mg/ml) and extra uridine (0.08 mg/ml), complete SC

medium (Formedium) or RPMI medium (Sigma-Aldrich). When indicated, media

were supplemented with 30 µg/ml doxycycline. For analysis in stress conditions,

YPD agar media was supplemented with 9 nM rapamycin, 2 M sodium chloride, 1

mM SNP with 1 mM SHAM, 7 mM copper sulphate, 300 mM lithium chloride, .01 %

sodium dodecyl sulfate (SDS), 15mM caffeine, 1.5 M sorbitol, 18 mM cycloheximide,

1.5 mM cobalt, 25 mM hydroxyurea, 3 % ethanol, 20 µM Calcofluor White, 4.5 mM

hydrogen peroxide, 0.75 mM EDTA and 2 µM Cerulenin (42). Cells were grown at 30

°C or 37 °C as indicated.

Plasmid construction

Oligos and plasmids used in this study are listed in Table 2 and 3. Plasmid ABp133

contains the *C. albicans HDA1* gene cloned in frame to a C-terminal HA tag. To

generate this plasmid, the full length *HDA1* gene was amplified from plasmid ABp88

(synthesised by GeneArt) with oligos Abo408 and Abo409, containing recognition

sites for *Xma*I. The digested PCR product was cloned into plasmid pHA-NAT

(ABp17)(43) digested with *Xma*I. Cloning was confirmed by PCR and Sanger

sequencing. *HDA3* was cloned in the pNIM plasmid (ABp111) (44) to generate

plasmid ABp177. For this purpose, the full length *HDA3* gene was amplified from

plasmid ABp152 (synthesised by GeneArt) with oligos ABo624 and ABo625,

containing *Xho*I and *Bg*lII restriction sites. ABo625 also supplied a stop codon

upstream the restriction site. This PCR fragment was cloned in pNIM digested with

*Sal*I and *Bg*lII. Cloning was confirmed by PCR and Sanger sequencing.

Construction of *C. albicans* mutants

Deletions of *HDA2* and *HDA3* were generated in the SN152 or BWP17 background

using the *Clox* system for gene disruption (33) using long-oligos PCR, the LAL (loxP-

ARG4-loxP) and NAT1-Clox (loxP-NAT1-MET3p-cre-loxP) plasmids as templates.

During all selections for *Clox* transformants media was supplemented with 2.5 mM

methionine and 2.5 mM cysteine to repress the *MET3* promoter and minimize Cre-

*lox*P mediated recombination. Nourseothricin resistant (Nou^R) transformants were

selected using 200 µg/ml nourseothricin (Melford). *HDA2* and *HDA3* gene deletions

were confirmed by PCR and markers were resolved by allowing Cre expression in

medium lacking methionine and cysteine as previously described (33).

C-terminal GFP tagging of *HDA2* and *HDA3* genes at the endogenous loci was

performed by long oligos PCR using plasmid pGFP-His1 (ABp11) as a template (45).

C-terminal HA tagging of Hda1 was performed by long oligos PCR using plasmid

ABp133 as a template and transformation into BWP17 (ABy215), HDA2-GFP

(ABy532) and HDA3-GFP (ABy376).

For rescue experiments, *HDA3* re-integrated at the *ADH* locus by digesting ABp177

with *Kpn*I and *Sac*II and transforming the product into the *hda3 ∆/∆* (Aby33) deletion

strain. Correct integration was confirmed by colony PCR. Transformations of *C.*

albicans strains were performed by electroporation using the protocol described in

(46) or by lithium acetate transformation (47) of competent cells (48) with a *S.*

cerevisiae method adapted to *C. albicans* (49).

Hyphal growth induction and quantification

Cultures were grown overnight (~16 hours) in 5 ml liquid YPD at 30 ^{o}C. Overnight

cultures were diluted 1:100 in YPD media and grown ~5 hours at 30 ^{o}C. 50 – 100

cells were plated to RPMI or Spider agar and incubated for ~6 days at 37 °C. For

doxycycline induction, the drug was added to the media at 30 µg/ml in 20 ml plates.

Phenotypes were documented by imaging with a Leica MZFLIII microscope at 10x to

16x magnification A minimum of 150 colonies were counted for each strain. For

liquid filamentation assays, overnight cultures were grown throughout the day and

from this stationary culture, 1 mL was harvested and washed with distilled water

before resuspending in 10 mL pre-warmed final media (YPD (control) or RPMI).

These cultures were grown at 37 °C (control at 30 °C) for 15-17 hours before

imaging on an Olympus IX81 inverted microscope at 60x magnification. Samples

were evaluated on four separate days and a minimum of 400 cells were counted per

sample.

White-opaque switching essay

Quantitative switching assays were performed as previously described (12) with

modifications. Briefly, strains were streaked from frozen stocks on YPD plates and

grown at 30 °C for 2 days. Single colonies were picked, resuspended in sterile water

and spread onto synthetic complete agar containing 5 μg/ml Phloxine B (Sigma-

Aldrich). Formation of opaque colonies or sectors was scored after 9 days.

Experiments were done in biological duplicate or triplicate on at least three separate

days. A minimum of 400 total colonies were counted for each strain.

The ggplot package in R studio was used to construct violin plots. Unpaired t-tests

were performed to test for significant differences between wild-type and mutant

strains.

Structural modelling

The model of Hda3 DBD and CCS domain was produced using Phyre2 in intensive

mode (50) and visualised using PyMol (The PyMOL Molecular Graphics System,

Version 1.8 Schrödinger, LLC.)

Whole cell extracts

Preparation of whole cell extracts was performed as described (51). Briefly,

overnight YPD cultures were diluted in YPD or RPMI and grown to OD_{600} = ~0.8 at

30 °C. Cells were harvested, resuspended in 200 µl lysis buffer (0.1 M NaOH, 0.05

M EDTA, 2 % SDS, 2% β-mercaptoethanol) and heated for 10 minutes at 95 °C

before adding 5 µl of 4 M Acetic acid and incubating for further 10 minutes at 90 °C.

Extracts were mixed with 50 µl loading buffer (0.25 M Tris-HCl pH 6.8, 50 %

Glycerol, 0.05 % Bromophenol Blue), incubated at 96 °C for 5 minutes and

centrifuged at 13000 rpm for 5 minutes. Supernatants were collected and analysed

by SDS-PAGE and Western blot analyses.

Immunoprecipitation

Immunoprecipitation was performed as described (52) with modifications. 1 L YPD

cultures (OD_{600} = 1) were harvested at 4000 rpm. Cell were washed 3 times in cold

water and resuspended in 1/5th volume (water/cells). Cell pellets were ground in

liquid nitrogen using a mortar and pestle for 30 minutes and resuspended in 10 ml of

cold lysis buffer (50 mM HEPES-NaOH pH 7.5, 150 mM NaCl, 5 mM EDTA, 0.1 %

NP-40, 5 mM DTT, 1x Roche EDTA-free protease inhibitor cocktail and 0.2 mM

PMSF added fresh before use). The cells were solubilized for 30 minutes with

rotation at 4 °C. Following centrifugation, supernatant samples were incubated with 4

µl of magnetic beads pre-coupled with anti-HA antibody (Sigma-Aldrich) for 2 hours

with rotation at 4 °C. Beads were washed four times in lysis buffer and analysed by

SDS-PAGE and Western blot.

Antibodies information

The following antibodies were used for Western analyses. Anti-HA antibody

(#11666606001, Sigma-Aldrich) diluted at 1:1000, anti-GFP antibody (Roche

#1184460001) diluted at 1:5000, anti-actin (Cooper Lab, Washington University, St.

Louis, Mo., USA) diluted at 1:5000.

RNA extraction

Overnight YPD cultures were diluted in YPD and grown to OD_{600} = ~0.8. Cells were

pelleted, washed once with sterile water and resuspended in pre-warmed YPD (30

ºC) or RPMI (37 ºC) for 90 minutes. RNA extraction was performed using a yeast

RNA extraction kit (E.Z.N.A. Isolation Kit RNA Yeast, Omega Bio-Tek) following

the manufacturer's instructions with the following modifications: (1) 30 ^{o}C

incubation in SE Buffer/2-mercaptoethanol/lyticase solution time was increased to 90

minutes; (2) lysis was performed with bead mill at top speed for 30 minutes at 4^{o}C.

RNA was treated with DNAse I and RNA quality was checked by electrophoresis

under denaturing conditions in 1% agarose, 1× HEPES, 6% Formaldehyde

(Sigma-Aldrich). RNA concentration was measured using a NanoDrop ND-1000

Spectrophotometer. cDNA synthesis was performed using qPCRBIO cDNA

Synthesis Kit (PCR Biosystems) following manufacturer's instructions.

High-throughput RNA sequencing

Strand-specific cDNA Illumina Barcoded Libraries were generated from 1 µg of

total RNA extracted from WT, *hda1 Δ/Δ*, *hda2 Δ/Δ* and *hda3 Δ/Δ* and sequenced

with an Illumina iSeq2000 platform. Illumina library preparation and deep-

sequencing was performed by the Genomics Core Facility at EMBL (Heidelberg,

Germany). RNA sequencing was performed in duplicates. Raw reads were

analysed following the RNA deep sequencing analysis pipeline using the Galaxy

platform (https://usegalaxy.org/). Downstream analysis of differential expressed

genes was performed with R Studio (https://www.rstudio.com/). Scatter Plot

matrices, using Pearson correlation coefficients, were generated with the ggplot

package. Heatmaps were generated with the pheatmap package and Pearson

correlation for clustering. Heatmaps show the log2 fold changes of differentially

expressed genes in *hda1 Δ/Δ*, *hda2 Δ/Δ* or *hda3 Δ/Δ* compared to wild-type

expression. Venn diagrams were generated using the FunRich programme(53).

RNA sequencing data are deposited into ArrayExpress (accession number: E-

MTAB-6920).

qRT-PCR

qRT-PCR was performed in the presence of SYBR Green (Bio-Rad) on a Bio-Rad

CFXConnect Real-Time System. Data was analysed with Bio-Rad CFX Manager 3.1

software and Microsoft Excel. Enrichment was calculated over actin. Histograms

represent data from three biological replicates. Error bars: standard deviation of

three biological replicates generated from 3 independent cultures of the same strain.

Functional analysis and modelling of transcriptional profiles

Gene set enrichment analysis (GSEA) (34, 54) was performed using the GSEA

PreRanked tool to determine whether a ranked gene list exhibited statistically

significant bias in their distribution within defined gene sets (55) and (*Sellam et al*,

personal communication). The weighted enrichment statistics were calculated on

10512 gene sets each containing 5-1000 genes, and the false discovery rate (FDR)

was calculated from 1000 permutations. Selected results graphs are shown. Since

enrichment profiles can exhibit correlations with hundreds of overlapping gene sets,

Cytoscape 3.6 (http://www.cytoscape.org) (56) and the Enrichment Map Pipeline

Collection plug-ins (http://apps.cytoscape.org/apps/enrichmentmappipelinecollection)

were used to further organise and visualise the GSEA. Enrichment maps were

calculated using default parameters.

C. albicans-C. elegans **pathogenesis assay**

The *C. elegans glp-4/sek-1* strain was used as described previously (39, 57). Briefly,

C. elegans were propagated on nematode growth medium (NGM) on lawns of *E. coli*

OP50. The *C. albicans-C. elegans* pathogenesis assay was performed as previously

described (40). Briefly, 100 µl of *C. albicans* cells from an overnight culture were

spread into a square lawn on a 10 cm plate containing brain-heart infusion (BHI)

agar and kanamycin (45 µg/ml). These were incubated for approximately 20 hours at

30 °C. Synchronized adult *C. elegans glp-4/sek-1* nematodes grown at 25°C were

carefully washed from NGM plates using sterile M9 buffer. Approximately 100 to 200

washed animals were then added to the centre of the *C. albicans* lawns and the

plates were incubated at 25°C for 4 hours. Worms were then carefully washed into a

15 ml tube using 10 ml of sterile M9, taking care to minimize the transfer of yeast.

Worms were washed four or five times with sterile M9. 30-40 worms were then

transferred into three wells of a six-well tissue culture plate (Corning, Inc.) containing

2 ml of liquid medium (80% M9, 20% BHI) and kanamycin (45 µg/ml). Worms were

scored daily into one of three categories: alive, dead with hyphae piercing the cuticle,

and dead without hyphae piercing the cuticle. Worms were considered to be dead if

they did not move in response to mechanical stimulation with a pick. Dead worms

were removed from the assay. *C. elegans* survival was examined by using the

Kaplan-Meier method and differences were determined by using the log-rank test

using OASIS 2 tool (58). Differences in the number of worms with *C. albicans* hyphal

formation were determined by using a one-way ANOVA with Dunnett's test for

multiple comparisons. The *C. elegans* pathogenesis assay presented here is an

average of three independent biologic replicates. A p-value of <0.05 was considered

statistically significant.

FUNDING INFORMATION

A.B and R.J.P: Medical Research Council [MR/M019713/1]; S.G: [BB/L008041/1].

Funding for open access charge: University of Kent.

ACKNOWLEDGMENTS

We thank GeneCore at EMBL for performing Illumina library preparation and

sequencing. We thank Dr Sellam (Biotechnology Research Institute, National

Research Council of Canada) and Dr Traven (Monash University) for help with the

GSEA analysis. We are grateful to Dr Anderson (The Ohio State University) for the

plasmid allowing deletion of mating type cassette and Prod Alistair Brown (University

of Aberdeen) for the Clox system. We thank members of the Kent Fungal Group for

useful discussions and critical reading of the manuscript.

REFERENCES

1.   Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG, White TC. 2012.

550   Hidden Killers: Human Fungal Infections. Sci Transl Med 4:165rv13-165rv13.

551   2.   Pfaller MA, Diekema DJ. 2010. Epidemiology of invasive mycoses in North

552   America. Crit Rev Microbiol 36:1–53.

553   3.   Slutsky B, Staebell M, Anderson J, Risen L, Pfaller M, Soll DR. 1987.

554   "White-opaque transition": a second high-frequency switching

555   system in Candida albicans. J Bacteriol 169:189–97.

556   4.   Bennett RJ, Johnson a D. 2005. Mating in Candida albicans and the search

557   for a sexual cycle. Annu Rev Microbiol 59:233–55.

558   5.   Kvaal C, Lachke SA, Srikantha T, Daniels K, McCoy J, Soll DR. 1999.

559   Misexpression of the opaque-phase-specific gene PEP1 (SAP1) in the white

phase of Candida albicans confers increased virulence in a mouse model of

cutaneous infection. Infect Immun 67:6652–62.

6.   Kvaal CA, Srikantha T, Soll DR. 1997. Misexpression of the white-phase-

specific gene WH11 in the opaque phase of Candida albicans affects switching

and virulence. Infect Immun 65:4468–75.

7.   Lachke SA, Lockhart SR, Daniels KJ, Soll DR. 2003. Skin facilitates Candida

albicans mating. Infect Immun 71:4970–6.

8.   Huang G, Wang H, Chou S, Nie X, Chen J, Liu H. 2006. Bistable expression of

WOR1, a master regulator of white-opaque switching in Candida albicans.

Proc Natl Acad Sci 103:12813–12818.

9.   Srikantha T, Borneman AR, Daniels KJ, Pujol C, Wu W, Seringhaus MR,

Gerstein M, Yi S, Snyder M, Soll DR. 2006. TOS9 Regulates White-Opaque

Switching in Candida albicans. Eukaryot Cell 5:1674–1687.

10.   Zordan RE, Miller MG, Galgoczy DJ, Tuch BB, Johnson AD. 2007. Interlocking

Transcriptional Feedback Loops Control White-Opaque Switching in Candida

albicans. PLoS Biol 5:e256.

11.   Zordan RE, Galgoczy DJ, Johnson AD. 2006. Epigenetic properties of white-

opaque switching in Candida albicans are based on a self-sustaining

transcriptional feedback loop. Proc Natl Acad Sci U S A 103:12807–12.

12.   Miller MG, Johnson AD. 2002. White-opaque switching in Candida albicans is

controlled by mating-type locus homeodomain proteins and allows efficient

mating. Cell 110:293–302.

13.   Noble SM, Gianetti BA, Witchley JN. 2017. Candida albicans cell-type

switching and functional plasticity in the mammalian host. Nat Rev Microbiol

15:96–108.

14. Staab JF, Bradway SD, Fidel PL, Sundstrom P. 1999. Adhesive and

mammalian transglutaminase substrate properties of Candida albicans Hwp1.

Science 283:1535–8.

15. Almeida RS, Brunke S, Albrecht A, Thewes S, Laue M, Edwards JE, Filler SG,

Hube B. 2008. The Hyphal-Associated Adhesin and Invasin Als3 of Candida

albicans Mediates Iron Acquisition from Host Ferritin. PLoS Pathog

4:e1000217.

16. Phan QT, Myers CL, Fu Y, Sheppard DC, Yeaman MR, Welch WH, Ibrahim

AS, Edwards JE, Filler SG. 2007. Als3 Is a Candida albicans Invasin That

Binds to Cadherins and Induces Endocytosis by Host Cells. PLoS Biol 5:e64.

17. Weissman Z, Kornitzer D. 2004. A family of Candida cell surface haem-binding

proteins involved in haemin and haemoglobin-iron utilization. Mol Microbiol

53:1209–1220.

18. Braun BR, Kadosh D, Johnson AD. 2001. NRG1, a repressor of filamentous

growth in C.albicans, is down-regulated during filament induction. EMBO J

20:4753–4761.

19. Lu Y, Su C, Wang A, Liu H. 2011. Hyphal Development in Candida albicans

Requires Two Temporally Linked Changes in Promoter Chromatin for Initiation

and Maintenance. PLoS Biol 9:e1001105.

20. Denning DW, Hope WW. 2010. Therapy for fungal diseases: opportunities and

priorities. Trends Microbiol 18:195–204.

21.   Kurdistani SK, Grunstein M. 2003. Histone acetylation and deacetylation in

607   yeast. Nat Rev Mol Cell Biol 4:276–284.

608   22.   Glozak MA, Sengupta N, Zhang X, Seto E. 2005. Acetylation and

609   deacetylation of non-histone proteins. Gene 363:15–23.

610   23.   Haberland M, Montgomery RL, Olson EN. 2009. The many roles of histone

611   deacetylases in development and physiology: Implications for disease and

612   therapy. Nat Rev Genet 10:32–42.

613   24.   Falkenberg KJ, Johnstone RW. 2014. Histone deacetylases and their inhibitors

614   in cancer, neurological diseases and immune disorders. Nat Rev Drug Discov

615   13:673–691.

616   25.   Srikantha T, Tsai L, Daniels K, Klar AJS, Soll DR. 2001. The Histone

617   Deacetylase Genes HDA1 and RPD3 Play Distinct Roles in Regulation of

618   High-Frequency Phenotypic Switching in Candida albicans. Society 183:4614–

619   4625.

620   26.   Hnisz D, Schwarzmüller T, Kuchler K. 2009. Transcriptional loops meet

621   chromatin: a dual-layer network controls white-opaque switching in Candida

622   albicans. Mol Microbiol 74:1–15.

623   27.   Lu Y, Su C, Solis N V., Filler SG, Liu H. 2013. Synergistic regulation of hyphal

624   elongation by hypoxia, CO_2, and nutrient conditions controls the virulence of

625   Candida albicans. Cell Host Microbe 14:499–509.

626   28.   Kadosh D, Lopez-Ribot JL. 2013. Candida albicans: Adapting to Succeed. Cell

627   Host Microbe 14:483–485.

29.   Tzogani K, van Hennik P, Walsh I, De Graeff P, Folin A, Sjöberg J, Salmonson

T, Bergh J, Laane E, Ludwig H, Gisselbrecht C, Pignatti F. 2018. EMA Review

of Panobinostat (Farydak) for the Treatment of Adult Patients with Relapsed

and/or Refractory Multiple Myeloma. Oncologist 23:631–636.

30.   Simó-Riudalbas L, Esteller M. 2015. Targeting the histone orthography of

cancer: drugs for writers, erasers and readers. Br J Pharmacol 172:2716–

2732.

31.   Wu J, Carmen a a, Kobayashi R, Suka N, Grunstein M. 2001. HDA2 and

HDA3 are related proteins that interact with and are essential for the activity of

the yeast histone deacetylase HDA1. Proc Natl Acad Sci U S A 98:4391–6.

32.   Lee JH, Maskos K, Huber R. 2009. Structural and Functional Studies of the

Yeast Class II Hda1 Histone Deacetylase Complex. J Mol Biol 391:744–757.

33.   Shahana S, Childers DS, Ballou ER, Bohovych I, Odds FC, Gow N a R, Brown

AJP. 2014. New Clox Systems for rapid and efficient gene disruption in

Candida albicans. PLoS One 9:e100390.

34.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,

Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-

wide expression profiles. Proc Natl Acad Sci U S A 102:15545–50.

35.   Sellam A, van het Hoog M, Tebbji F, Beaurepaire C, Whiteway M, Nantel A.

2014. Modeling the transcriptional regulatory network that controls the early

hypoxic response in Candida albicans. Eukaryot Cell 13:675–90.

36.   Merico D, Isserlin R, Stueker O, Emili A, Bader GD. 2010. Enrichment Map: A

Network-Based Method for Gene-Set Enrichment Visualization and

Interpretation. PLoS One 5:e13984.

37. Enjalbert B, Nantel A, Whiteway M. 2003. Stress-induced gene expression in

Candida albicans: absence of a general stress response. Mol Biol Cell

14:1460–7.

38. Tuch BB, Mitrovich QM, Homann OR, Hernday AD, Monighetti CK, De La

Vega FM, Johnson AD. 2010. The transcriptomes of two heritable cell types

illuminate the circuit governing their differentiation. PLoS Genet 6:e1001070.

39. Breger J, Fuchs BB, Aperis G, Moy TI, Ausubel FM, Mylonakis E. 2007.

Antifungal Chemical Compounds Identified Using a C. elegans Pathogenicity

Assay. PLoS Pathog 3:e18.

40. Pukkila-Worley R, Peleg AY, Tampakakis E, Mylonakis E. 2009. Candida

albicans hyphal formation and virulence assessed using a Caenorhabditis

elegans infection model. Eukaryot Cell 8:1750–8.

41. Lewis RE, Kontoyiannis DP. 2001. Rationale for combination antifungal

therapy. Pharmacotherapy 21:149S–164S.

42. Homann OR, Dea J, Noble SM, Johnson AD. 2009. A phenotypic profile of the

Candida albicans regulatory network. PLoS Genet 5.

43. Gerami-nejad M, Forche A, Mcclellan M, Berman J. 2012. Analysis of protein

function in clinical C . albicans isolates 5314.

44. Park Y, Park Y, Morschhauser J, Morschhauser J. 2005. Tetracycline-

Inducible Gene Expression and Gene Deletion in Candida albicans.

Microbiology 4:1328–1342.

45. Gerami-Nejad M, Berman J, Gale CA. 2001. Cassettes for PCR-mediated

construction of green, yellow, and cyan fluorescent protein fusions inCandida

albicans. Yeast 18:859–864.

46. De Backer MD, Maes D, Vandoninck S, Logghe M, Contreras R, Luyten

WHML. 1999. Transformation of Candida albicans by Electroporation. Yeast

15:1609–1618.

47. Schiestl RH, Gietz RD. 1989. High efficiency transformation of intact yeast

cells using single stranded nucleic acids as a carrier. Curr Genet 16:339–346.

48. Knop M, Siegers K, Pereira G, Zachariae W, Winsor B, Nasmyth K, Schiebel

E. 1999. Epitope tagging of yeast genes using a PCR-based strategy: more

tags and improved practical routines. Yeast 15:963–972.

49. Walther A, Wendland J. 2003. An improved transformation protocol for the

human fungal pathogen Candida albicans. Curr Genet 42:339–343.

50. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2

web portal for protein modeling, prediction and analysis. Nat Protoc 10:845–

58.

51. von der Haar T. 2007. Optimized protein extraction for quantitative proteomics

of yeasts. PLoS One 2:e1078.

52. Buscaino A, White SA, Houston DR, Lejeune E, Simmer F, de Lima Alves F,

Diyora PT, Urano T, Bayne EH, Rappsilber J, Allshire RC. 2012. Raf1 Is a

DCAF for the Rik1 DDB1-Like Protein and Has Separable Roles in siRNA

Generation and Chromatin Modification. PLoS Genet 8:e1002499.

53. Pathan M, Keerthikumar S, Chisanga D, Alessandro R, Ang C-S, Askenase P,

Batagov AO, Benito-Martin A, Camussi G, Clayton A, Collino F, Di Vizio D,

Falcon-Perez JM, Fonseca P, Fonseka P, Fontana S, Gho YS, Hendrix A,

Hoen EN-'t, Iraci N, Kastaniegaard K, Kislinger T, Kowal J, Kurochkin I V.,

Leonardi T, Liang Y, Llorente A, Lunavat TR, Maji S, Monteleone F, Øverbye

A, Panaretakis T, Patel T, Peinado H, Pluchino S, Principe S, Ronquist G,

Royo F, Sahoo S, Spinelli C, Stensballe A, Théry C, van Herwijnen MJC,

Wauben M, Welton JL, Zhao K, Mathivanan S. 2017. A novel community

driven software for functional enrichment analysis of extracellular vesicles

data. J Extracell Vesicles 6:1321455.

54.   Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J,

Puigserver P, Carlsson E, Ridderstråle M, Laurila E, Houstis N, Daly MJ,

Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES,

Hirschhorn JN, Altshuler D, Groop LC. 2003. PGC-1alpha-responsive genes

involved in oxidative phosphorylation are coordinately downregulated in human

diabetes. Nat Genet 34:267–73.

55.   Sellam A, Hogues H, Askew C, Tebbji F, van Het Hoog M, Lavoie H,

Kumamoto C a, Whiteway M, Nantel A. 2010. Experimental annotation of the

human pathogen Candida albicans coding and noncoding transcribed regions

using high-resolution tiling arrays. Genome Biol 11:R71.

56.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

Schwikowski B, Ideker T. 2003. Cytoscape: A Software Environment for

Integrated Models of Biomolecular Interaction Networks. Genome Res

13:2498–2504.

57.   Moy TI, Ball AR, Anklesaria Z, Casadei G, Lewis K, Ausubel FM. 2006.

Identification of novel antimicrobials using a live-animal infection model. Proc

Natl Acad Sci U S A 103:10414–10419.

58.   Han SK, Lee D, Lee H, Kim D, Son HG, Yang J-S, Lee S-J V., Kim S. 2016.

OASIS 2: online application for survival analysis 2 with features for the

analysis of maximal lifespan and healthspan in aging research. Oncotarget

7:56147–56152.

59.   Synnott JM, Guida A, Mulhern-Haughey S, Higgins DG, Butler G. 2010.

Regulation of the Hypoxic Response in Candida albicans. Eukaryot Cell

9:1734–1746.

60.   Marcil A, Gadoury C, Ash J, Zhang J, Nantel A, Whiteway M. 2008. Analysis of

PRA1 and its relationship to Candida albicans-macrophage interactions. Infect

Immun 76:4345–4358.

61.   Noble SM, Johnson AD. 2005. Strains and Strategies for Large-Scale Gene

Deletion Studies of the Diploid Human Fungal Pathogen Candida albicans

Eukaryot Cell 4: 298-309

62.   Wilson RB, Davis D, Mitchell AP. 1999. Rapid Hypothesis Testing with

Candida albicans through Gene Disruption with Short Homology Regions

Journal of bacteriology 181: 1868-74

740
FIGURE LEGENDS

Figure 1. The Hda1 complex is conserved in *C. albicans*

(A) Domain organisation of *C. albicans* Hda2 and Hda3 proteins **(B)** *Left:* Structural

alignment of *C. albicans* DBD3 domain (red) with *S. cerevisiae* DBD3 domain

(yellow); *Right:* Structural modelling of *C. albicans* CCD3 **(C)** Co-

Immunoprecipitation of Hda1 with Hda2 and Hda3. Hda1-HA Immunoprecipitation

(Ip) analysed with anti-HA or with anti-GFP to detect Hda2 and Hda3. **(D)** Schematic

of *Clox* gene disruption strategy used to construct *hda2Δ/Δ* and *hda3Δ/Δ* mutants.

(E) Serial dilution assay of WT, *hda2Δ/Δ* and *hda3Δ/Δ* mutants on solid YPD media

at 30 °C. **(F)** Growth curves of WT, *hda2Δ/Δ* and *hda3Δ/Δ* isolates in YPD liquid

media at 30 °C.

Figure 2. Global gene expression changes in the absence of Hda2 and Hda3

(A) Pearson correlation matrix of gene expression changes observed in *hda1 Δ/Δ,*

hda2 Δ/Δ and *hda3 Δ/Δ* grown in YPD at 30 ^{o}C. r = Pearson correlation coefficient. p

= p-value. **(B)** The network of functional groups of genes regulated by Hda2 and

Hda3 constructed by GSEA and Enrichment Map. Blue circles represent down

regulated while orange circles depict upregulated gene sets which are linked in the

network by grey lines. The diameter of the circles varies based upon the number of

transcripts within each set. **(C)** Example enrichment plots for selected genes sets

differentially expressed in *hda2 Δ/Δ* (left) and *hda3 Δ/Δ* (right). Ketoconazole_up: set

of genes upregulated in *C. albicans* cells grown in the presence of ketoconazole

(59); Phagocytosis_up: gene set upregulated following engulfment by primary Bone

Marrow Derived Macrophages (60). The x-axis shows genes ranked according to

their expression in the mutants from up-regulated (left) to down-regulated (right)

genes. Black vertical lines mark individual genes. The cumulative value of the

enrichment score (y-axis) is represented by the green line. A positive normalised

enrichment score (NES) indicates enrichment in the up-regulated group of genes in

hda2 Δ/Δ and *hda3 Δ/Δ*.

Figure 3. Phenotyping of *hda2* Δ/Δ and *hda3* Δ/Δ strains

Serial dilution assay showing growth of *hda2* Δ/Δ and *hda3* Δ/Δ mutants on solid

YPD media with additives as indicated and incubation at 30 °C for 2 - 4 days. **(A)**

Additives affecting *hda2* Δ/Δ and *hda3* Δ/Δ growth relative to wildtype strain. Mutants

are resistant to rapamycin and sensitive to SNP and SHAM, sodium chloride and

copper sulphate. **(B)** Conditions eliciting normal growth of mutant strains relative to

matching wildtype.

Figure 4. Hda2 and Hda3 inhibit white-opaque switching

(A) Enrichment plots for genes differentially expressed in *hda1* Δ/Δ, *hda2* Δ/Δ and

hda3 Δ/Δ in comparison to genes upregulated in opaque cells (38). The x-axis shows

genes ranked according to their expression in the mutants from up-regulated (left) to

down-regulated (right) genes. Black vertical lines mark individual genes. The

cumulative value of the enrichment score (y-axis) is represented by the green line. A

positive normalised enrichment score (NES) indicates enrichment in the up-regulated

group of genes in *hda1* Δ/Δ, *hda2* Δ/Δ and *hda3* Δ/Δ. **(B)** Heat map depicting the log2

fold change in *hda1* Δ/Δ, *hda2* Δ/Δ and *hda3* Δ/Δ isolates compared to WT for the

Opaque_up gene set. **(C)** Log2 fold change values and significance for *WOR1* gene

expression in *hda1* Δ/Δ, *hda2* Δ/Δ and *hda3* Δ/Δ compared to WT. **(D)** *Left*:

Percentage of sector colonies in WT, *hda2* Δ/Δ and *hda3* Δ/Δ isolates. ** = p value ≤

.01. *Right*: Representative image of cells grown on Phloxine B agar. O = opaque; W

= white.

Figure 5. Gene expression profiling in hyphae-inducing conditions in WT, *hda1*

***Δ/Δ, hda2* Δ/Δ and *hda3* Δ/Δ isolates**

(A) The network representing changes in gene expression in WT cells grown in

yeast-inducing conditions (YPD) versus hyphae-inducing conditions (RPMI)

constructed by GSEA and Enrichment Map. Blue circles represent down regulated

gene sets, while orange depicts upregulated gene sets which are linked in the

network by grey lines that indicate function. The diameter of the circles varies based

upon the number of transcripts within each set. **(B)** Pearson correlation matrix of

gene expression changes observed in *hda1 Δ/Δ*, *hda2 Δ/Δ* and *hda3 Δ/Δ* grown in in

hyphae growth media (RPMI) at 37 ^{o}C. r = Pearson correlation coefficient. p = p-

value. **(C)** *Left:* Heat map depicting the log2 fold change in WT cells grown in

hyphae-inducing conditions (RPMI at 37 ^{o}C) versus WT cells grown in yeast-inducing

conditions (YPD at 30 ^{o}C). Gene known to be involved in hyphae formation or biofilm

formation are indicated. *Right:* Log2 fold changes of hyphae-induced and repressed-

genes in *hda1 Δ/Δ, hda2 Δ/Δ* and *hda3 Δ/Δ* isolates. Cell were grown in RPMI at 37

^{o}C. **(D)** Quantitative reverse transcriptase PCR (qRT-PCR) analyses to measure

HWP1 and *ALS3* transcript levels in WT, *hda1 Δ/Δ, hda2 Δ/Δ* and *hda3 Δ/Δ* isolates

grown in yeast (YPD at 30 ^{o}C) or hyphae (RPMI at 37 ^{o}C) inducing conditions.

Transcripts levels are visualized relative to *ACT1* transcript levels. Error bars in each

panel: standard deviation of three biological replicates.

Figure 6. Hda2 and Hda3 contribute to filamentation growth but not virulence

(A) *Left:* Representative images of colony morphology of WT, *hda1 Δ/Δ, hda2 Δ/Δ*

and *hda3 Δ/Δ* grown on hyphal inducing Spider and RPMI agar at 37 °C. *Right:*

Quantification of colony morphologies. **(B)** Quantification of cellular morphologies of

WT, *hda1 Δ/Δ, hda2 Δ/Δ* and *hda3 Δ/Δ* grown in liquid RPMI media at 37 °C. **(C)**

Rescue experiment of colony morphology upon genomic integration of the *HDA3*

gene in the *hda3 Δ/Δ* mutant background (*hda3 Δ/Δ+ HDA3*). WT, heterozygous

(*HDA3/ hda3* Δ) and homozygous *hda3* Δ/Δ isolates were included as a control. Cell

were grown on hyphal inducing RPMI agar at 37 °C. **(D)** *Left:* Representative images

of dead *C. elegans* with and without hyphae-mediated killing. *Middle:* Survival curve

of worms incubated with WT, *hda2* Δ/Δ and *hda3* Δ/Δ strains over a 72 hour period.

Right: The percentage of total worms dead (black bars) and worms killed by hyphae

piercing the cuticle (grey bars) after the 72 hour incubation period. Error bars

represent the standard deviation of three independent biological replicates.

Figure 7. Stability of the Hda1 protein is regulated by environmental changes

(A) Venn diagram of genes differentially expressed in *hda1*Δ/Δ relative to WT in

yeast and hyphal growth conditions. **(B)** Enrichment plots for genes differentially

expressed in *hda1* Δ/Δ relative to genes downregulated (Hyphae_Lee_DN) or

upregulated (Hyphae_Lee_up) in hyphae-growth condition in Lee's media . The x-

axis shows genes ranking according to their expression in *hda1*Δ/Δ from up-

regulated (left) to down-regulated (right) genes. Black vertical lines mark individual

genes. The cumulative value of the enrichment score (y-axis) is represented by the

green line. A positive normalised enrichment score (NES) indicates enrichment in the

up-regulated group of genes while a negative NES indicates prevalence of the genes

in the down-regulated group. **(C)** HA and GFP Western blots of whole protein extract

from strains: Hda1-HA, Hda2-GFP and Hda3-GFP. Actin is shown as a loading

control. Cells were grown in yeast (YPD 30 °C) or hyphae-inducing (RPMI 37 °C)

conditions. **(D)** Model showing how decreased Hda1 and Hda3 protein levels leading

to the functional rewiring of the Hda1 complex in *C. albicans* in different

environments. Under yeast growth condition, the Hda1 complex functions as a global

regulator of gene expression due to the high level of Hda1 and Hda3. Under hyphae

growth conditions, decreased levels of Hda1 and Hda3 leads to specialisation of the

Hda1 complex controlling only filamentous growth.

Table 1. *C.albicans* strains used in this study.

Strain Number	Description	Genotype	Figure	Source
ABy_54	SN152	*MTL a/alpha ura3Δ-iro1Δ::imm434/URA3-IRO1 his1Δ/his1Δ arg4Δ/arg4Δ leu2Δ/leu2Δ*	1E-F, 2A-C, 3A-B, 4A-C, 5A-D, 6A- B, 6D, 7A-B	(61)
ABy_66	BWP17	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG*	1F, 6B-C	(62)
ABy_179	*hda1 Δ/Δ*	*MTL a/a arg4Δ/arg4Δ his1Δ/his1Δ leu2Δ/leu2Δ URA3/ura3Δ::λimm434IRO1 /iro1Δ::λimm434 hda1Δ::C.d.HIS1/hda1Δ::C. m.LEU2*	2A-C, 3A-B, 4A-C, 5A-D, 6A-B, 6D, 7A-B	(26)
ABy_191	HDA3/Δ	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA3/hda3Δ::LAL*	6C	This study
ABy_331	BWP17 *hda3Δ/Δ*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG hda3Δ/hda3Δ*	1F, 6B-C	This study
ABy_347	*hda2Δ/Δ*	*MTL a/alpha ura3Δ-iro1Δ::imm434/URA3-IRO1 his1Δ/his1Δ arg4Δ/arg4Δ leu2Δ/leu2Δ hda2Δ/hda2Δ*	1E-F, 2A-C, 3A-B, 4A-D, 5A-C, 6B, 6D, 6A	This study
ABy_376	*HDA3:HDA3-GFP*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA3::HIS/HDA3-GFP*	1C, 7C	This study
ABy_393	BWP17 *MTLαΔ*	*MTLa/MTLalphaΔ ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG*	4D	This study

ABy_460	SN152 *hda3Δ/Δ*	*MTL a/alpha ura3Δ-iro1Δ::imm434/URA3-IRO1 his1Δ/his1Δ arg4Δ/arg4Δ leu2Δ/leu2Δ hda3Δ/hda3Δ*	1E, 2A-C, 3A-B, 4A-C, 5A-D, 6A-B, 6D	This study
ABy_472	*ADH1/ pNIM-HDA3:adh1Δ*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG hda3Δ/hda3Δ* pNIM-HDA3::adh1Δ/ADH1	6C	This study
ABy_532	*HDA2:HDA2-GFP*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA2::HIS/HDA2-GFP*	1C, 7C	This study
ABy_536	*HDA2:HDA2-GFP HDA1:HDA1-HA*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA2::HIS/HDA2-GFP HDA1::NAT/HDA1-HA*	1C	This study
ABy_539	*HDA3:HDA3-GFP HDA1:HDA1-HA*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA3::HIS/HDA3-GFP HDA1::NAT/HDA1-HA*	1C	This study
ABy_547	*hda2Δ/Δ MTLαΔ*	*MTL a/ MTLalphaΔ ura3Δ-iro1Δ::imm434/URA3-IRO1 his1Δ/his1Δ arg4Δ/arg4Δ leu2Δ/leu2Δ hda2Δ/hda2Δ*	4D	This study
ABy_551	*hda3Δ/Δ MTLαΔ*	*MTL a/ MTLalphaΔ ura3Δ-iro1Δ::imm434/URA3-IRO1 his1Δ/his1Δ arg4Δ/arg4Δ leu2Δ/leu2Δ hda3Δ/hda3Δ*	4D	This study
ABy_563	*HDA1:HDA1-HA*	*MTL a/alpha ura3Δ::Δimm434/ura3Δ::Δimm434 his1::hisG/his1::hisG arg4::hisG/arg4::hisG HDA1::NAT/HDA1-HA*	1C, 7C	This study

847 **Table 2**. Oligos used in this study.

Number	Primer	Sequence	Figure	Description
ABo_408	HDA1_Xma_fw1	TAAGCACCCGGGatgtcgactggtcaagaagaa	1C, 7C-D	Isolate synthetic Hda1 from p88
ABo_409	HDA1_Xma_rev1	TAAGCACCCGGGatctt	1C, 7C-D	Isolate synthetic Hda1

		cggaagaggagtagtc		from p88
ABo_351	Pr_sir2_adh1	CGCACTCACGTAAAC ACTT	1C, 6C, 7C-D	Check Hda1 in p17; confirm reintegration of Hda3_pNIM cassette at ADH1 locus
ABo_412	HDA1end_chkseq	GGAAATAGTTCGAAC GGTGG	1C, 7C-D	Check Hda1 in p17; sequencing p133
ABo_452	F1gaHDA1_Haintg	TCCGAAATTCATTATT AAGGAATTATATAGA AGCTACCATTTTCAC ATCTATTATCATTTTC CTTTTTAAGAatgtcgact ggtcaagaagaaca	1C, 7C-D	Isolate Hda1-HA integration cassette from p133; long oligo
ABo_453	R1gaHDA1_Haintg	CAGATCTATATCTATT CTCTTTCTTTCTTTTT TTTTGGTTTTTTGTTG TTGTTGTTGTTGTTTC TACTCGAAgtaaaacga cggccagtgaattc	1C, 7C-D	Isolate Hda1-HA integration cassette from p133; long oligo
ABo_179	PR1_NAT	CTGTATCTATAAGCA GTATCATCC	1C, 7C-D	Check presence of Hda1-HA integration cassette at native locus
ABo_417	dwnstrmHDA1rev_ check	CTCGATGCCTGATTT GGATG	1C, 7C-D	Check presence of Hda1-HA integration cassette at native locus
ABo_458	F1HDA2_GFP	TGTTAGAAAACTCTG GCTCGGGTGCCAATA ATAGACAAAATAATC GTATTAGTCGAGGTG CAACACCTCTTGGTG GTGGTtctaaaggtgaaga attatt	1C, 7C-D	Isolate GFP tagging cassette from p11 with flanking integration sequences for Hda2; long oligo
ABo_459	R2HDA2_GFP	GCATGTATTTACAAA TTTTTGGATAAGAAA AAGTAGCATATGGAA ACACAAAACCAAGAA AGAAATCATGgaattcc ggaatatttatgagaaac	1C, 7C-D	Isolate GFP tagging cassette from p11 with flanking integration sequences for Hda2; long oligo
ABo_460	F1HDA3_GFP	CAATGCTTTTACATTT TTAAATGATACTAAAT ATTTGAAAAAGAGGA AAAATCGAGGAATAA CTCCTAAAGGTGGTG GTtctaaaggtgaagaattatt	1C, 7C-D	Isolate GFP tagging cassette from p11 with flanking integration sequences for Hda3; long oligo
ABo_461	R2HDA3_GFP	CTTATCATTTACATAA TTAAAAAAACAAAAA ACAAGCTAATCTTAT GTTTATGTGGGGGCC ACATTTTCTgaattccgg aatatttatgagaaac	1C, 7C-D	Isolate GFP tagging cassette from p11 with flanking integration sequences for Hda3; long oligo
ABo_145	PF1_his1	GTTCCAGCAGATGGC	1C, 7C-D	Check Hda3-GFP or

		GAGTAC		Hda2-GFP tagging
ABo_261	Pr4_hda2chk_rev	GTAATATCTGATCAG AACCTTT	1C, 7C-D	Check Hda2-GFP tagging
ABo_262	Pr4_hda3chk_rev	TCGTTAATCAAAATTA TACACTC	1C, 7C-D	Check Hda3-GFP tagging
ABo_229	HDA2D_Clox_Fw1	TCTATTTTCAAGAAGT TAGACCCATTCTTGG AATAATTATACTTGCA AGAGAAGGCATTGAA ATTGCATTACGGCCA GTGAATTGTAATA	1D	Isolate deletion cassette from p80 or p83 with flanking integration sequences for Hda2; long oligo
ABo_230	HDA2D_Clox_rev1	GCATGTATTTACAAA TTTTTGGATAAGAAA AAGTAGCATATGGAA ACACAAACCAAGAA AGAAATCATGTCGGA ATTAACCCTCACTAA	1D	Isolate deletion cassette from p80 or p83 with flanking integration sequences for Hda2; long oligo
ABo_165	Pr1_chck_arg4	AGTGTGGAAAGAAGA GATGC	1D	Check presence of p80 cassette in Hda2 or Hda3 heterozygote
ABo_423	HDA3_Natcloxchk	CCGGTGCTATGGTTA GATTG	1D	Check presence of p83 cassette in hda2ΔΔ or hda3ΔΔ
ABo_261	Pr4_hda2chk_rev	GTAATATCTGATCAG AACCTTT	1D	Check presence of p80 cassette in Hda2 heterozygote and hda2ΔΔ
ABo_413	HDA2chk_internalf w	CAGCAGGTAGACTTG ATG	1D	Check presence/absence of Hda2 in hda2ΔΔ unresolved/resolved
ABo_261	Pr4_hda2chk_rev	GTAATATCTGATCAG AACCTTT	1D	Check presence/absence of Hda2 in hda2ΔΔ unresolved/resolved
ABo_231	HDA3D_Clox_Fw1	GTTCTTAATATTTGTA ACTTTTCCAACTTAAA ATAATTATTGCATATT GCACTAAAACTAAAA CTACTATAAATacggcc agtgaattgtaata	1D	Isolate deletion cassette from p80 or p83 with flanking integration sequences for Hda3; long oligo
ABo_232	HDA3D_Clox_Rev 1	CAATCTTATCATTTAC ATAATTAAAAAAACAA AAAACAAGCTAATCT TATGTTTATGTGGGG GCCACATTTTCTtcgga attaaccctcactaa	1D	Isolate deletion cassette from p80 or p83 with flanking integration sequences for Hda3; long oligo
ABo_262	Pr4_hda3chk_rev	TCGTTAATCAAAATTA TACACTC	1D	Check presence of p80 or p83 cassette in Hda3 heterozygote or hda3ΔΔ
ABo_415	HDA3chk_internalf w	CAACAAGAAcTGTGG AACATG	1D	Check presence/absence of Hda3 in hda3ΔΔ

				unresolved/resolved
ABo_416	HDA3chk_internalrev	GGTGGTTCTATAAATCCCGG	1D	Check presence/absence of Hda3 in hda3ΔΔ unresolved/resolved
ABo_462	Pf_MTLa1_Chk	TTGAAGCGTGAGAGGCAGGAG	4D	Check presence of MATa
ABo_463	Pr_MTLa1_Chk	GTTTGGGTTCCTTCTTTCTCATTC	4D	Check presence of MATa
ABo_464	Pf_MTLalpha1_Chk	TTCGAGTACATTCTGGTCGCG	4D	Check presence of MATα
ABo_465	Pr_MTLalpha1_Chk	TGTAAACATCCTCAATTGTACCCGA	4D	Check presence of MATα
ABo_511	Pf_SAT1	GGTGGCGGAAACATTGGATG	4D	Check presence of Sat1
ABo_512	Pr_SAT1	TCAATGCCGCCGAGAGTAAA	4D	Check presence of Sat1
ABo_624	HDA3_XhoI_fw	TAAGCACTCGAGATGgatttaaggaaaattttg	6C	Isolate Hda3 from ABp152 with XhoI and BglII sites (for cloning into ABp111)
ABo_625	HDA3_stpBglII_rv	TTAACGAGATCTttatttaggagttattcctcgat	6C	Isolate Hda3 from ABp152 with XhoI and BglII sites (for cloning into ABp111: pNIM)
ABo_416	HDA3chk_internalrev	GGTGGTTCTATAAATCCCGG	6C	Confirm cloning of p177 (Hda3_pNIM)
ABo_514	pNIM_Rev2	CAGTTTGGTTCAGCACCTTG	6C	Confirm cloning of ABp_177 (Hda3_pNIM)
LD_515		CCATCATAAAATGTCGAGCGTC	6C	Sequencing of ABp_177 (Hda3_pNIM)
ABo_350	Pf_sir2_adh1	ctctatcactgatagggagtgg	6C	Confirm reintegration of ABp_177 (Hda3_pNIM) cassette at ADH1 locus
ABo_174	Act1_Fw2	CTACGTTTCCATTCAAGCTGTT	5D	qRT-PCR for ACT1
ABo_176	Act1_Rev3	AAACTGTAACCACGTTCAGACA	5D	qRT-PCR for ACT1
ABo_469	qALS3_Fw1	CCTATTCCAACAACTACAAT	5D	qRT-PCR for ALS3
ABo_470	qALS3_Rev1	TATTGAGTCAGTTGGATTAG	5D	qRT-PCR for ALS3
ABo_471	qHWP1_Fw1	CCAGTTACTTCTGGATCATC	5D	qRT-PCR for HWP1
ABo_472	qHWP1_Rev1	TCGGTACAAACACTGTTAGA	5D	qRT-PCR for HWP1

850   **Table 3**. Plasmids used in this study.

ABp_Number	Plasmid	Figure	Description/Source	Source
ABp_11	pGFP-His1	1C	GFP tagging vector	(45)
ABp_17	pHA_*NAT1*	1C	HA tagging vector	(43)
ABp_80	LAL (loxP-ARG4-loxP)	1D	*Arg4* substitution products	(33)
ABp_83	NAT1-Clox (loxP-NAT1-MET3p-cre-loxP)	1D	*Nat* substitution products	(33)
ABp_88	HDA1_synthetic	1C	Hda1 synthetic	GeneArt
ABp_111	pNIM	6C	Tetracycline inducible integration to ADH1 locus	(44)
ABp_133	HDA1synthetic_pHA_NAT	1C	Source of Hda1-HA integration cassette	This study
ABp_136	Sat1 flipper with MTLαKO flanking regions	4D	Deletion of MATα	Gift from Matthew Anderson Lab, Ohio State
ABp_152	HDA3 synthetic	6C	HA-HDA3 synthetic: 6x CUG->TCA	GeneArt
ABp_177	HDA3-pNIM	6C	Tetracycline inducible integration of Hda3 to ADH1 locus	This study

A

Hda2 (orf19.6952)
669 aa

Hda3 (orf19.7344)
835 aa

DBD2   CCD2

I: 20 %

DBD3   CCD3

B

■ *C. albicans DBD3* ■ *S. cerevisiae DBD3* ■ *C. albicans CCD3*

C

	Input			Ip		
Hda1-HA	+	-	+	+	-	+
Hda2-GFP	+	+	-	+	+	-

Hda1-HA

Hda2-GFP

	Input			Ip		
Hda1-HA	+	-	+	+	-	+
Hda3-GFP	+	+	-	+	+	-

Hda1-HA

Hda3-GFP

D

HDA2/HDA3

Disruption 1st allele

LoxP   ARG4   LoxP

Disruption 2nd allele

LoxP   ARG4   LoxP

NatR   Cre Rec

Cre induction

LoxP

E

N/S

hda2 Δ/Δ

hda3 Δ/Δ

WT

F

N/S

N/S

■ *hda2 Δ/Δ* ■ *WT*   ■ *hda3 Δ/Δ* ■ *WT*

Peterson et al_Fig2

A

Yeast growth conditions

hda1 Δ/Δ		
r= 0.7 p= 2.2e-16	*hda2* Δ/Δ	
r= 0.66 p= 2.2e-16	r= 0.46 p= 2.2e-16	*hda3* Δ/Δ

Pearson Correlation

1   0   −1

B

hda2 Δ/Δ

RNA and Ribosome Biogenesis

Transporters

hda3 Δ/Δ

RNA and Ribosome Biogenesis

Transporters

○ upregulated   ○ downregulated

C

Response to Drugs

hda2 Δ/Δ
Ketoconazole_UP

NES= + 2.55

hda3 Δ/Δ
Ketoconazole_UP

NES= + 3.10

■ upregulated ■ downregulated

Host Interaction

hda2 Δ/Δ
Phagocytosis_up

NES= + 2.63

hda3 Δ/Δ
Phagocytosis_up

NES= + 2.40

A

B

C

WOR1 (C1_10150W)

Strain	Log2 Fold change	p	q
hda1 Δ/Δ	3.95	3.86 e-32	5.92 e-29
hda2 Δ/Δ	2.46	4.61 e-13	2.67 e-10
hda3 Δ/Δ	2.50	1.46 e-13	2.62 e-10

D

A

WT YPD vs RPMI

Hyphae

Yeast

○ upregulated   ○ downregulated

Pearson Correlation

−1

B

HYPHAE MEDIUM

hda1 Δ/Δ		
r= 0.55 p= 2.2e-16	*hda2* Δ/Δ	
r= 0.38 p= 2.2e-16	r= 0.65 p= 2.2e-16	*hda3* Δ/Δ

C

WT   *hda1* Δ/Δ   *hda2* Δ/Δ   *hda3* Δ/Δ

Hyphal Growth

Biofilm

Lo2 Fold Changes

−3

D

qRT-PCR: HWP1

expression relative to *ACT1*

YEAST — WT, *hda1* Δ/Δ, *hda2* Δ/Δ, *hda3* Δ/Δ
HYPHAE — WT, *hda1* Δ/Δ, *hda2* Δ/Δ, *hda3* Δ/Δ

qRT-PCR: ALS3

expression relative to *ACT1*

YEAST — WT, *hda1* Δ/Δ, *hda2* Δ/Δ, *hda3* Δ/Δ
HYPHAE — WT, *hda1* Δ/Δ, *hda2* Δ/Δ, *hda3* Δ/Δ