Identification of Adomavirus Virion Proteins

Nicole L. Welch[1], Michael J. Tisza[1], Gabriel J. Starrett[1], Anna K. Belford[1], Diana V. Pastrana[1],

Yuk-Ying S. Pang[1], John T. Schiller[1], Ping An[2], Paul G. Cantalupo[2], James M. Pipas[2], Samantha

Koda[3], Kuttichantran Subramaniam[3], Thomas B. Waltzek[3], Chao Bian[4], Qiong Shi[4], Zhiqiang

Ruan[4], Terry Fei Fan Ng[5], and Christopher B. Buck[1]*

[1]Lab of Cellular Oncology, NCI, NIH, Bethesda, MD, 20892 USA

[2]Dept. of Biological Sciences, Univ. of Pittsburgh, Pittsburgh, PA 15260 USA

[3]Dept. of Infectious Disease and Pathology, Univ. of Florida, Gainesville, FL, 32611 USA

[4]Shenzhen Key Lab of Marine Genomics, Guangdong Provincial Key Lab of Molecular Breeding
in Marine Economic Animals, BGI Academy of Marine Sciences, BGI, Shenzhen, Guangdong
518083, China

[5]College of Veterinary Medicine, University of Georgia, Athens, GA 30602 USA

*Corresponding author: buckc@mail.nih.gov

Abstract

Adenoviruses, papillomaviruses, and polyomaviruses are collectively known as small DNA
tumor viruses. Although it has long been recognized that small DNA tumor virus oncoproteins
and capsid proteins show a variety of structural and functional similarities, it is unclear whether
these similarities reflect descent from a common ancestor, convergent evolution, horizontal gene
transfer among virus lineages, or acquisition of genes from host cells. Here, we report the
discovery of a dozen new members of an emerging virus family, the *Adomaviridae*, that unite a
papillomavirus/polyomavirus-like replicase gene with an adenovirus-like virion maturational
protease. Adomaviruses were initially discovered in a lethal disease outbreak among endangered
Japanese eels. New adomavirus genomes were found in additional commercially important fish
species, such as tilapia, as well as in reptiles. The search for adomavirus sequences also revealed
an additional candidate virus family, which we refer to as xenomaviruses, in mollusk datasets.
Analysis of native adomavirus virions and expression of recombinant proteins showed that the
virion structural proteins of adomaviruses are homologous to those of both adenoviruses and
another emerging animal virus family called adintoviruses. The results pave the way toward
development of vaccines against adomaviruses and suggest a framework that ties small DNA
tumor viruses into a shared evolutionary history.

Author Summary

In contrast to cellular organisms, viruses do not encode any universally conserved genes. Even
within a given family of viruses, the amino acid sequences encoded by homologous genes can
diverge to the point of unrecognizability. Although members of an emerging virus family, the
Adomaviridae, encode replicative DNA helicase proteins that are recognizably similar to those of
polyomaviruses and papillomaviruses, the functions of other adomavirus genes have been
difficult to identify. Using a combination of laboratory and bioinformatic approaches, we
identify the adomavirus virion structural proteins. The results link adomavirus virion protein
operons to those of other midsize non-enveloped DNA viruses, including adenoviruses and
adintoviruses.

Introduction

Polyomaviruses, papillomaviruses, and adenoviruses are historically defined as small DNA
tumor viruses (Pipas 2019). Although members of a fourth animal-tropic non-enveloped DNA
virus family, the *Parvoviridae*, are not known to cause tumors they share a number of biological
features with traditional small DNA tumor viruses. Each of the four virus families encodes non-
enveloped virion proteins with similar pentameric single-β-jellyroll core folds and members of
each of the four families express functionally similar oncogenes that inactivate cellular tumor
suppressor proteins (Figure 1)(de Souza, Iyer et al. 2010, Krupovic and Koonin 2017). An
emerging group of animal viruses called adintoviruses appears to represent a candidate fifth
family with similarities to small DNA tumor viruses
https://www.biorxiv.org/content/10.1101/697771v3.

Figure 1: Maps of representative virus genomes. Genes are color-coded based on
known or inferred functions. Polyomaviruses, papillomaviruses, and adomaviruses have
circular double-stranded DNA genomes that were linearized for display. Parvoviruses
have linear single-stranded DNA (ssDNA) genomes, while adintoviruses and adenoviruses
have linear dsDNA genomes. Symbols and abbreviations: white lollipop,
retinoblastoma/pocket protein (Rb) interaction motif; gray lollipop, domain with predicted
fold similar to DNAJ chaperone proteins; AEP, domain resembling archael-eukaryotic
primase small catalytic subunit; PLA2, domain with predicted fold similar to
phospholipase A2. See main text for other gene names.

Polyomaviruses, papillomaviruses, and parvoviruses are proposed to have descended from
circular Rep-encoding single-stranded DNA (CRESS) virus ancestors (Koonin, Dolja et al.
2015). This model explains the phylogenetic relationships of the replicative superfamily 3
helicase (S3H) and rolling circle "nickase" endonuclease domain of CRESS virus and small
DNA tumor virus replicase genes (Kazlauskas, Varsani et al. 2019), but it does not account for
possible similarities between the virion proteins and accessory genes of the "-oma" families and
adenoviruses. Achieving a better understanding of the relationships between small DNA tumor
virus families has the potential to guide comparative studies of these common human pathogens.

In 2011, a previously unknown circular dsDNA virus was discovered in a lethal disease outbreak
among Japanese eels (*Anguilla japonica*)(Mizutani, Sayama et al. 2011, Okazaki, Yasumoto et
al. 2016). Two related viruses have since been isolated from Taiwanese marbled eels (*Anguilla*

marmorata) and a giant guitarfish (*Rhynchobatus djiddensis*)(Wen, Chen et al. 2015, Dill,
Camus et al. 2018). In contrast to the eel viruses, which encode S3H replicase proteins that
closely resemble the large tumor antigens (LT) of fish polyomaviruses, the guitarfish virus
encodes a distinct S3H replicase, called EO1, that is distant from LT and is instead more closely
related to the E1 replicases of papillomaviruses. The name "adomaviruses" has been applied to
this emerging family, connoting the fact that the three known species each encode homologs of
the adenain virion-maturational proteases of *ad*enoviruses as well as poly*oma*virus and
papill*oma*virus S3H homologs.

The primary goal of this study is to identify the adomavirus virion proteins and to uncover
possible evolutionary relationships to the virion proteins of small DNA tumor viruses. To
discover additional adomavirus species, we conducted metagenomic sequencing studies and
developed a pipeline to detect small DNA tumor virus-related sequences in the NCBI Sequence
Read Archive (SRA). Bioinformatic methods were used to predict which adomavirus ORFs
might represent virion proteins and the predictions were confirmed through functional expression
in cell culture. The results pave the way toward development of preventive vaccines against
pathogenic adomaviruses.

Results

Detection of additional adomaviruses

A post-mortem metagenomics analysis of an aquarium-bred Amazon red discus cichlid
(*Symphysodon discus*) exhibiting lethargy and inflamed skin lesions revealed a complete circular
adomavirus genome (Figure 2). Histopathological analysis of skin lesions from the discus
specimen showed no evidence of intranuclear inclusions or other obvious histopathology.

Figure 2: Genome maps of previously unknown adomaviruses. Genes were named
based on conventions originally developed for Japanese eel adomavirus (NC_015123).
Alpha adomaviruses (top half of figure) are defined by the presence of an adomavirus-
specific S3H replicase, designated EO1. A row of fragmentary adomavirus sequences from
RNA datasets are shown in the middle of the figure. Beta adomaviruses (bottom half of
figure) encode a polyomavirus-like large tumor antigen (LT). In some cases, repetitive or
GC-rich patches (particularly near the 3' end of the LO2-3 ORF) could not be resolved
using available short-read datasets. Coverage gaps are represented as white bars on the
ruler. A poorly conserved set of accessory genes upstream of Alpha adomavirus EO1 genes
show varying degrees of similarity to the S-adenosyl methionine-binding pocket of cellular
SET proteins, which function as histone lysine-methyltransferases. Adomavirus SET
homologs are highly divergent from all previously described eukaryotic and viral SET
genes. The same is true for adomavirus EO2-4 and EO3 segments that encode homologs of
the catalytic small subunit of archaeal eukaryotic primases (AEPs, hatched boxes).

Figure supplement 1: a table of accession numbers and Linnaean designations of hosts

Figure supplement 2: examples of the annotation process

124 An adomavirus from an apparently healthy green arowana (*Scleropages formosus*) was first
125 identified in TBLASTN searches of the NCBI Whole-Genome Shotgun (WGS) database as a set
126 of short contigs with similarity to Japanese eel adomavirus proteins. A complete adomavirus
127 genome was characterized by Sanger sequencing of overlapping PCR products using DNA left
128 over from the original fin snip used for the WGS project (Bian, Hu et al. 2016).

129 The WGS TBLASTN survey also revealed a 4 kb contig with a sequence resembling adomavirus
130 LT in a dataset for western softhead grenadier fish (*Malacocephalus occidentalis*) and a nearly
131 complete adomavirus genome in a dataset for a skin biopsy of a leatherback sea turtle
132 (*Dermochelys coriacea*). Genome sequences for the two viruses were completed using parent
133 SRA datasets.

134 A pipeline using DIAMOND (a faster alternative to BLASTX (Buchfink, Xie et al. 2015)) was
135 developed to screen SRA datasets for fish, amphibians, and reptiles. SRA datasets rich in reads
136 encoding adomavirus-like protein sequences were subjected to *de novo* assembly. This approach
137 resulted in the identification of seven additional adomavirus genomes (Figure 2). Notably,
138 adomavirus sequences were found in genome sequencing datasets for farmed tilapia
139 (*Oreochromis niloticus*), which represent a $7.5 billion per year global aquaculture industry, and
140 in the most extensively aquacultured fish in developing countries, the mirror carp (*Cyprinus
141 carpio*)(Bacharach, Mishra et al. 2016, Belton, Little et al. 2018). Adomavirus-like fragments
142 were also detected in transcriptomic datasets for brook salamander (*Calotriton asper*) and two
143 closely related species of anole lizard (genus *Anolis*).

144 The WGS search for adomavirus-like S3H sequences also led to the discovery of a divergent
145 class of circular DNA elements that we tentatively designate "xenomaviruses," connoting the
146 exotic nature of their predicted S3H and virion proteins (Figure 2 Figure supplement 2, Figure 3,
147 and Figure 4). Intriguingly, a conserved xenomavirus ORF shows distant predicted structural
148 similarity to the L1 and VP1 penton proteins of papillomaviruses and polyomaviruses. The
149 inferred replicase gene of a partial xenomavirus sequence detected in a dataset for pink abalone
150 (*Haliotis corrugata*) encodes a domain with predicted structural similarity to the nickase domain
151 of porcine circovirus 2 (a CRESS virus). The identification of additional viruses in this class
152 could shed light on the evolutionary interrelationships between small DNA tumor viruses and
153 CRESS viruses.

Figure 3: Genome maps of candidate xenomaviruses. Brackets indicate segments where remote similarities were detected in DELTA-BLAST, HHpred, and Phyre[2] searches. The functions of these segments remain hypothetical.

Adomavirus phylogeny

Adomavirus sequences can be divided into two groups based on their replicative S3H genes. A group we designate Alpha is defined by the presence of an EO1 S3H replicase gene that yields moderate hits (BLASTP E-values ~1e-9) for papillomavirus E1 proteins. Adomavirus group Beta is defined by the presence of an S3H replicase similar (E-values ~1e-60) to the LT proteins of polyomaviruses. A network display of BLASTP relationships is shown in Figure 4. The Alpha and Beta groupings are recapitulated in phylogenetic analyses of adomavirus LO8 (Adenain) homologs (Figure 4 Figure supplement 2).

Figure 4: Sequence similarity network (BLASTP E-value threshold 1e-9) for S3H proteins from indicated virus groups.

170   Figure supplement 1: an interactive version of the figure that can be viewed using
171   Cytoscape software https://cytoscape.org

172   Figure supplement 2: phylogenetic tree of LO8 proteins

173 The adomavirus LT-like proteins share a full range of familiar polyomavirus-like features,
174 including an N-terminal DnaJ domain, a potential retinoblastoma-interaction motif (LXCXE or
175 LXXLFD)(An, Saenz Robles et al. 2012, Gouw, Michael et al. 2018). Several Alpha adomavirus
176 EO1 proteins encode a C-terminal domain with predicted similarity to DNA-binding RFX-type
177 winged helices (pfam02257). The RFX-like domain is conserved at the C-terminus of S3H
178 proteins found in larger dsDNA (for example, vaccinia virus D5 YP232992), some parvoviruses
179 (for example, bovine parvovirus NS1 NP_041402), and xenomaviruses. HHpred analyses
180 indicate that, like other small DNA tumor virus replicases, both classes of adomavirus S3H
181 proteins encode a central nicking endonuclease-like origin binding domain (Hickman, Ronning
182 et al. 2002, Iyer, Koonin et al. 2005, Koonin, Dolja et al. 2015, Kazlauskas, Varsani et al. 2019).

183 **Bioinformatic prediction of adomavirus virion proteins**

184 To determine whether predicted adomavirus LO proteins are expressed from spliced or unspliced
185 ORFs, RNAseq data published by Wen and colleagues (Wen, Chen et al. 2015) were analyzed to
186 determine the splicing patterns of marbled eel adomavirus transcripts. Splice acceptors
187 immediately upstream of the inferred ATG initiator codons of LO4, LO5, LO6, and LO7
188 proteins were extensively utilized (Figure 5 Figure supplement 5). Messenger RNAs encoding
189 the adenovirus late genes carry a tripartite leader (TPL) that has been shown to enhance
190 translation late in the adenovirus life cycle (Logan and Shenk 1984). A similar three-exon leader
191 sequence was detected upstream of the marbled eel adomavirus LO genes (Figure 1).

192 HHpred searches were performed to detect remote similarities between the predicted structures
193 of adomavirus proteins and known protein structures. Adomavirus LO4 proteins have a predicted
194 C-terminal trimeric coiled-coil domain. This relatively generic predicted fold gives a large and
195 diverse range of hits in HHpred searches, including the coiled-coil domains of various viral fiber
196 proteins (for example, avian reovirus σC, 97%). Negative-stain EM images indicate that
197 adomavirus virions do not have a vertex fiber (Mizutani, Sayama et al. 2011, Wen, Chen et al.
198 2015, Dill, Camus et al. 2018). Intriguingly, LO4 proteins yielded low probability (~50%)
199 HHpred hits for adenovirus pIX, a trimeric coiled coil protein that serves as a "cement" that
200 smooths the triangular facets of the adenovirus virion. The hypothesis that LO4 is a pIX homolog
201 could hypothetically account for the smooth appearance of adomavirus virion facets in negative-
202 stain EM.

203 A C-terminal segment of some LO5 sequences, as well as alignments of multiple LO5
204 sequences, yielded moderate (~60% probability) HHpred hits for CvsA1_340L protein (single-
205 jellyroll vertex penton) of *Paramecium bursaria* chlorella virus 1 (PDB:6NCL_a6). A screen
206 shot of a typical HHpred result is shown in Figure 2 Figure supplement 2.

207 Alignments of LO6 ORFs show high-probability (95%) HHPred hits for a 37 amino acid
208 hydrophobic segment of adenovirus pX, a minor virion core protein that is thought to participate
209 in condensation of the viral chromatin (Nemerow, Stewart et al. 2012). LO6 alignments also
210 showed moderate probability hits (Figure 2 Figure supplement 2) for adenovirus pVI, which is

believed to play a role in destabilizing cellular membranes during the infectious entry process
(Moyer, Besser et al. 2015). The results suggest that LO6 might be a fused homolog of
adenovirus pVI and pX virion core proteins.

HHpred searches using LO7 sequences did not produce interpretable results, with the single
exception of the LO7 sequence of grenadier adomavirus, which gives a moderate-probability hit
for the V20 double-jellyroll hexon major capsid protein of Sputnik virophage (Figure 2 Figure
supplement 2).

In addition to offering a convenient way to summarize aggregate BLASTP interrelationships
(e.g., Figure 4), all-against-all sequence similarity network analysis can be a useful method for
discovering distant similarities between highly divergent groups of proteins (Iranzo, Krupovic et
al. 2017). In one noteworthy example, network analyses were recently used to detect remote
sequence similarities between small DNA tumor virus S3H replicases and CRESS virus
replicases (Kazlauskas, Varsani et al. 2019). In contrast to traditional analyses using
phylogenetic trees, it is possible for network analyses to detect individual pairs of sequences in
separate clusters that both happen to have preserved the primary sequence of a common ancestor.
We performed low-stringency network analyses to further investigate possible remote sequence
similarities between adenovirus, adintovirus, and candidate adomavirus virion proteins.

Networks for adomavirus LO4 (inferred fiber or cement protein) and LO7 (inferred double-
jellyroll hexon major capsid protein) showed few or no connections to adenovirus or adintovirus
virion protein sequences, even at a BLASTP E-value threshold of 1e-1. In contrast, LO8
(adenain) proteins yielded informative networks at E-value thresholds of 1e-5 (Figure 5). At less
stringent E-value thresholds (1e-2) similarities between adomavirus LO5 (inferred single-
jellyroll vertex penton) and inferred adintovirus penton protein sequences emerged. PSI-BLAST
searches using LO5 alignments confirmed the apparent sequence similarities (E-values ~1e-6) to
predicted adintovirus penton proteins found in arthropod and coral datasets (e.g., GBM63801
EFA12278 LSMT01002030). Although adenovirus pVI did not cluster with LO6 (inferred virion
core protein) or proposed adintovirus virion core proteins (Cupiennin, GasderminX, PLA2X) at
an E-value threshold of 1e-2, Alpha adomavirus LO6 proteins clustered with adintovirus
cupiennin and Beta adomavirus LO6 proteins clustered with adenovirus pX proteins.

Figure 5: Sequence similarity network analysis of predicted minor virion proteins. The Adenain (LO8) network was constructed with a BLASTP E-value threshold of 1e-5. Penton (LO5) and virion core protein (LO6) networks used a threshold of 1e-2.

Figure supplement 1: an interactive version of the figure that can be viewed using Cytoscape software https://cytoscape.org

Experimental confirmation of predicted adomavirus virion proteins

The bioinformatic results suggest that the LO4-8 operon encodes syntenic homologs of adenovirus and adintovirus virion proteins. To experimentally test this prediction, marbled eel adomavirus was grown in eel kidney cell culture (Wen, Chen et al. 2015) and virions were purified using Optiprep gradient ultracentrifugation. Virion-enriched gradient fractions were separated on SDS-PAGE gels and bands were subjected to mass spectrometric analysis (Figure 6). The analysis identified prominent bands in the Coomassie-stained gel as LO4, LO5, LO7, and LO8. The relative intensities of bands identified as LO5 and LO7 are consistent with the prediction that the two proteins constitute penton and hexon subunits, respectively. Lower molecular weight bands showed hits for LO6, suggesting that this protein is present in virions in an LO8 (adenain) cleaved form. LO6 proteins were found to encode potential adenain cleavage motifs ((MIL)XGXG or L(LR)GG) (Ruzindana-Umunyana, Imbeault et al. 2002). A list of protein modifications observed in the mass spectrometric analysis is shown in Figure 6 Figure supplement 1.

Figure 6: SDS-PAGE analysis of marbled eel adomavirus virions purified from infected EK-1 cells, with bands annotated by mass spectrometric analysis. Clarified lysates of infected cells were ultracentrifuged through an Optiprep gradient. Peak virion-containing fractions were selected then subjected to size exclusion chromatography over agarose resin. The sample was then subjected to TCA precipitation and run on an SDS-PAGE gel. Thirteen gel bands were individually excised, trypsin-digested, and analyzed on a Q Exactive HF Mass Spectrometer.

Figure supplement 1: post-translational modifications observed in mass spectrometric results

Expression of recombinant virion proteins

Adenovirus penton proteins can spontaneously assemble into 12-pentamer subviral particles that may serve as decoy pseudocapsids in vivo (Vragniau, Hubner et al. 2017). Similarly, recombinant polyomavirus and papillomavirus penton proteins can spontaneously assemble into icosahedral virus-like particles (VLPs) that closely resemble native virions. We are not aware of any reports of production of full-size (i.e., hexon+penton) adenovirus VLPs. To investigate the behavior of recombinant adomavirus virion proteins, codon-modified marbled eel adomavirus LO1-LO8 expression plasmids were transfected individually into human 293TT cells (Buck, Pastrana et al. 2004). Optiprep ultracentrifugation was used to separate virus-like particles (VLPs) from smaller solutes. A human papillomavirus type 16 (HPV16) L1/L2 expression plasmid was used as a positive control for VLP formation (Buck, Thompson et al. 2005). Cells transfected with adomavirus LO4, LO5, or LO7 expression constructs each produced particles that migrated into the core fractions of Optiprep gradients, whereas cells transfected with LO1,

LO2-3, LO6, or LO8 alone did not show evidence of particle formation (Figure 7). Negative-
stain EM analysis showed that the LO4, LO5, and LO7 particles were irregular (Figure 7 Figure
supplement 1). In co-transfections of various combinations of LO genes, it was found that
inclusion of LO6 inhibited the formation of LO5 and LO7 particles but did not impair the
formation of LO4 particles (Figure 8). The fact that over-expression of LO6 can antagonize
particle formation supports the bioinformatic prediction that LO6 is a minor virion component
that directly interacts with LO5 and LO7.

Figure 7: SDS-PAGE analysis VLPs assembled in cells transfected with individual
LO expression constructs. 293TT cells were transfected with individual codon modified
marbled eel LO expression plasmids indicated at the top of the image. The cells were
lysed, subjected to nuclease digestion and a clarifying 5000 x g spin. Soluble material
was ultracentrifuged through Optiprep gradients. Core gradient fractions with peak VLP
content were subjected to SDS-PAGE analysis.

Figure supplement 1: Negative-stain electron microscopy of recombinant particles.

Figure 8: SDS-PAGE gel showing that co-expression of LO6 antagonizes LO5 and
LO7 (but not LO4) particle formation. 293TT cells were transfected with the indicated
combination of plasmids and subjected to ultracentrifugal separation through Optiprep
gradients.

Individually expressed LO5 (penton) and LO7 (hexon) particle preparations showed dsDNA
signal in Quant-iT PicoGreen assays (Invitrogen), indicating the presence of nuclease-resistant
encapsidated DNA within the purified particles. Optiprep-purified particle preparations from
cells co-transfected with LO4, LO5, and LO7 were subjected to an additional round of nuclease
digestion with salt-tolerant Benzonase endonuclease (Sigma) followed by agarose gel filtration
to remove the nuclease and any residual digested DNA fragments. Nuclease-treated/gel filtered
particles typically contained roughly seven nanograms of DNA per microgram of total protein,
confirming the presence of nuclease-resistant nonspecific cellular DNA within the particles. The
observation is reminiscent of findings for recombinant papillomavirus VLPs (Buck, Thompson et
al. 2005).

Discussion

We have identified a dozen new representatives of the emerging virus family *Adomaviridae*.
Four of the new sequences are associated with terrestrial vertebrates, extending the known host
range beyond fish. Phylogenetic analyses reveal two adomavirus lineages that appear to have
independently co-evolved with host animals. This observation suggests that the two adomavirus
lineages both infected the first jawed vertebrates roughly half a billion years ago. Within both
adomavirus lineages, there are members associated with commercially important fish species.

Vaccine immunogens comprised of recombinant VLPs have been highly successful in humans
(Schiller and Lowy 2015). In particular, vaccines against HPVs have proven remarkably
immunogenic, even after a single dose. Our identification of adomavirus virion proteins and
demonstration of their ability to assemble into roughly spherical DNA-containing particles
should facilitate the development of recombinant subunit vaccines against these viruses, some of
which are known to cause severe disease in fish.

The adomavirus virion protein genes, penton (LO5), core (LO6), hexon (LO7), and adenain
(LO8), appear to be syntenic homologs of adenovirus and adintovirus virion protein genes. The
results also suggest that LO4 may be a homolog of adenovirus pIX, a trimeric coiled coil protein
that cements the facets of the adenovirus virion. At a primary sequence level, adomavirus virion
proteins more closely resemble adintovirus virion proteins, rather than adenovirus virion proteins
(Figure 5). These results tie the *Adomaviridae* into a broad consortium of eukaryotic virus
families (Koonin, Krupovic et al. 2015).

In unicellular eukaryotes, non-enveloped midsize (10-50 kb) dsDNA viruses have been shown to
have a remarkable degree of genetic modularity (Koonin, Dolja et al. 2015, Yutin, Shevchenko et
al. 2015) https://www.biorxiv.org/content/10.1101/697771v3. The pairing of related virion
proteins in adenoviruses, adintoviruses, and adomaviruses with entirely different classes of DNA
replicase genes thus has ample precedence in non-animal eukaryotes. It will be important to
apply emerging higher-throughput search algorithms, such as Mash Screen (Ondov, Starrett et al.
2019) and Cenote-Taker (Tisza, Pastrana et al. 2019), to exhaustively search for each of the
overlapping hallmark genes of this virus supergroup in genomic, transcriptomic, and
metagenomic surveys, particularly datasets for terrestrial vertebrates.

Materials and Methods

Sample Collection and cell culture

A red discus cichlid (*Symphysodon discus*) was purchased at a pet shop in Gainesville, Florida.
The fish was moribund and showed erythematous skin lesions. Propagation of the discus
adomavirus in cell culture was attempted by overlaying skin tissue homogenates on Grunt Fin
(GF) and Epithelioma Papulosum Cyprini cell lines (ATCC). Neither cytopathic effects nor
qPCR-based detection of viral replication were observed during two blind passages of 14 days
each.

Dr. Chiu-Ming Wen generously provided EK-1 cells (a Japanese eel kidney line) infected with
the Taiwanese marbled eel adomavirus (Wen, Chen et al. 2015). The virus was propagated by
inoculation of supernatants from the infected culture into uninfected EK-1 cells cultured at room
temperature in DMEM with 10 % fetal calf serum. Human embryonic kidney-derived 293TT
cells were cultured as previously described (Buck, Pastrana et al. 2004).

Viral genome sequencing

For the discus adomavirus, total DNA was extracted from a skin lesion and subjected to deep sequencing. Marbled eel adomavirus virions were purified from lysates of infected EK-1 cells using Optiprep gradient ultracentrifugation (Peretti, FitzGerald et al. 2015). DNA extracted from Optiprep gradient fractions was subjected to rolling circle amplification (RCA, TempliPhi, GE Health Sciences). The marbled eel adomavirus RCA products and discus total DNA were prepared with a Nextera XT DNA Sample Prep kit and sequenced using the MiSeq (Illumina) sequencing system with 2 × 250 bp paired-end sequencing reagents. In addition, the marbled eel adomavirus RCA product was digested with *AclI* and *EcoRI* restriction enzymes and the resulting early and late halves of the viral genome were cloned separately into the *AclI* and *EcoRI* restriction sites of pAsylum+. The sequence of the cloned genome was verified by a combination of MiSeq and Sanger sequencing. The clones are available upon request.

For the arowana adomavirus, overlapping PCR primers were designed based on WGS accession numbers LGSE01029406, LGSE01031009, LGSE01028643, LGSE01028176, and LGSE01030049 (Bian, Hu et al. 2016). PCR products were subjected to primer-walking Sanger sequencing.

Discovery of viral sequences in NCBI databases

Papillomavirus E1 sequences were downloaded from PaVE https://pave.niaid.nih.gov (Van Doorslaer, Li et al. 2017). Polyomavirus LT sequences were downloaded from PyVE https://ccrod.cancer.gov/confluence/display/LCOTF/Polyomavirus (Buck, Van Doorslaer et al. 2016). Parvovirus NS1 proteins and S3H proteins of CRESS viruses and virophage-like viruses were compiled from multiple databases, including RefSeq, WGS, and TSA, using TBLASTN searches. Adenovirus, virophage, and bacteriophage PolB sequences were downloaded from GenBank nr using DELTA-BLAST searches (Boratyn, Schaffer et al. 2012) with Alpha or Beta adintovirus PolB proteins as bait.

SRA datasets for fish, amphibians, and reptiles were searched using DIAMOND (Buchfink, Xie et al. 2015) or NCBI SRA Toolkit (http://www.ncbi.nlm.nih.gov/books/NBK158900/) in TBLASTN mode using adomavirus protein sequences as the subject database or query, respectively. Reads with similarity to the baits were collected and subjected to BLASTX searches against a custom library of viral proteins representing adomaviruses and other small DNA tumor viruses. SRA datasets of interest were subjected to de novo assembly using the SPAdes suite (Bankevich, Nurk et al. 2012, Nurk, Meleshko et al. 2017) or Megahit (Li, Liu et al. 2015, Li, Luo et al. 2016). Contigs encoding virus-like proteins were identified by TBLASTN searches against adomavirus protein sequences using Bowtie (Langmead and Salzberg 2012). The candidate contigs were validated using the CLC Genomics Workbench 12 align to reference function.

Predicted protein sequences were automatically extracted from contigs of interest using getorf (http://bioinfo.nhri.org.tw/cgi-bin/emboss/getorf)(Rice, Longden et al. 2000). Sequences were clustered using EFI-EST (https://efi.igb.illinois.edu/efi-est/)(Gerlt, Bouvier et al. 2015, Zallot, Oberg et al. 2018) and displayed using Cytoscape v3.7.1 (Shannon, Markiel et al. 2003). Multiple sequence alignments were constructed using MAFFT (https://toolkit.tuebingen.mpg.de/#/tools/mafft)(Kuraku, Zmasek et al. 2013, Katoh, Rozewicki

et al. 2019). Individual or aligned protein sequences were subjected to HHpred searches
(https://toolkit.tuebingen.mpg.de/#/tools/hhpred)(Hildebrand, Remmert et al. 2009, Meier and
Soding 2015, Zimmermann, Stephens et al. 2017) against PDB, Pfam-A, NCBI Conserved
Domains, and PRK databases.

Contigs were annotated using Cenote-Taker (Tisza, Pastrana et al. 2019) with an iteratively
refined library of conserved adintovirus protein sequences. Maps were drawn using MacVector
17. Phylogenetic analyses were performed using Phylogeny.fr with default settings (Dereeper,
Guignon et al. 2008).

Marbled eel adomavirus transcript analysis, late ORF expression, and virion purification

RNAseq reads reported by Wen et al (Wen, Chen et al. 2015) were aligned to the marbled eel
adomavirus genome using HISAT2 version 2.0.5 (Kim, Langmead et al. 2015) with the
following options: "--rna-strandness FR --dta --no-mixed --no-discordant". Integrated Genome
Viewer (IGV) version 2.4.9 (Robinson, Thorvaldsdottir et al. 2017) was used to determine splice
junctions and their depth of coverage. Additional validation was performed by visual inspection
using CLC Genomics Workbench 12.

Codon-modified expression constructs encoding the marbled eel adomavirus LO1-LO8 proteins
were designed according to a modified version of a previously reported algorithm
(https://github.com/BUCK-LCO-NCI/Codmod_as_different_as_possible)(Pastrana, Buck et al.
2004). 293TT cells were transfected with LO expression constructs for roughly 48 hours. Cells
were lysed in a small volume of PBS with 0.5% Triton X-100 or Brij-58 and Benzonase
Dnase/Rnase (Sigma)(Buck and Thompson 2007). After one hour of maturation at neutral pH,
the lysate was clarified at 5000 x g for 10 min. The clarified lysate was loaded onto a 15-27-33-
39-46% Optiprep gradient in PBS with 0.8 M NaCl. Gradient fractions were collected by bottom
puncture of the tube and screened by PicoGreen dsDNA stain (Invitrogen), BCA, or SDS-PAGE
analysis. Electron microscopic analysis was performed by spotting several microliters of
Optiprep fraction material (or, in some instances, particles exchanged out of Optiprep using
agarose gel filtration) onto carbon film copper grids, followed by staining with 0.5% uranyl
acetate.
433
Mass Spectrometry

Optiprep-purified marbled eel adomavirus virions were precipitated with trichloroacetic acid. A
1 ml sample was treated with 100 µl of 0.15% deoxycholic acid and incubated at room
temperature for 10 minutes. 100 µl of 100% TCA was then added and the sample was vortexed
and incubated on ice for 30 minutes. Following the incubation, the sample was centrifuged at
10,000 x g for 10 minutes at 4°C. The supernatant was removed, and the remaining pellet was
washed with ice-cold acetone to remove residual TCA. The protein pellet was solublized with
NuPAGE Sample Buffer + 5% BME (Sigma) and run on a 10-12% Bis-Tris MOPS gel
(Thermo). The protein bands were visualized using InstantBlue (Expedeon). Thirteen gel bands
were individually excised and placed into 1.5 ml Eppendorf tubes. The gel bands were sent to the
National Cancer Institute in Fredrick, Maryland where they were de-stained, digested with
trypsin, and processed on a Thermo Fisher Q Exactive HF Mass Spectrometer. Thermo Proteome
Discoverer 2.2 software was used for initial protein identification. The uninterpreted mass
spectral data were also searched against *Anguilla* proteins (Swiss-Prot and TrEMBL database

containing 105,268 proteins), Bos taurus proteins (Swiss-Prot and TrEMBL database containing
48,288 proteins), a common contaminants database (cRAPome), and translated marbled eel
adomavirus ORFs. Further analysis was conducted using Protein Metrics Biopharma software to
identify modifications missed in initial analyses.

Ethics Statement

All animal tissue samples were received as diagnostic specimens collected for pathogen testing
and disease investigation purposes.

Data Availability

GenBank accession numbers for sequences deposited in association with this study are
BK010891 BK010892 BK011012 BK011013 BK011014 BK011015 BK011016 BK011017
BK011018 BK011019 BK011020 BK011021 BK012039 BK012040 BK012041 MF946549
MF946550 MH282863.

Acknowledgments

The authors are indebted to Eugene Koonin and Natalya Yutin for their generous guidance and
for the spirited discussions that inspired us to pursue this study. We are particularly grateful to
them for sharing their observation that adomaviruses encode a recognizable adenain homolog
and their discovery of adomavirus sequences in the arowana WGS datasets. The authors are also
grateful to Lisa Jenkins for her extensive technical guidance on analyzing mass spectrometric
data. We thank Karl Münger for useful discussions, including advice about oncogene sequence
motifs.
469
470

References

An, P., M. T. Saenz Robles and J. M. Pipas (2012). "Large T antigens of polyomaviruses:
amazing molecular machines." Annu Rev Microbiol **66**: 213-236.
Bacharach, E., N. Mishra, T. Briese, M. C. Zody, J. E. Kembou Tsofack, R. Zamostiano, A.
Berkowitz, J. Ng, A. Nitido, A. Corvelo, N. C. Toussaint, S. C. Abel Nielsen, M. Hornig, J. Del
Pozo, T. Bloom, H. Ferguson, A. Eldar and W. I. Lipkin (2016). "Characterization of a Novel
Orthomyxo-like Virus Causing Mass Die-Offs of Tilapia." mBio **7**(2): e00431-00416.
Bankevich, A., S. Nurk, D. Antipov, A. A. Gurevich, M. Dvorkin, A. S. Kulikov, V. M. Lesin, S.
I. Nikolenko, S. Pham, A. D. Prjibelski, A. V. Pyshkin, A. V. Sirotkin, N. Vyahhi, G. Tesler, M.
A. Alekseyev and P. A. Pevzner (2012). "SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing." J Comput Biol **19**(5): 455-477.
Belton, B., D. Little and S. Bush. (2018). "Let them eat carp: Fish farms are helping to fight
hunger." from https://theconversation.com/let-them-eat-carp-fish-farms-are-helping-to-fight-
hunger-90421.
Bian, C., Y. Hu, V. Ravi, I. S. Kuznetsova, X. Shen, X. Mu, Y. Sun, X. You, J. Li, X. Li, Y. Qiu,
B. H. Tay, N. M. Thevasagayam, A. S. Komissarov, V. Trifonov, M. Kabilov, A. Tupikin, J.

Luo, Y. Liu, H. Song, C. Liu, X. Wang, D. Gu, Y. Yang, W. Li, G. Polgar, G. Fan, P. Zeng, H.
Zhang, Z. Xiong, Z. Tang, C. Peng, Z. Ruan, H. Yu, J. Chen, M. Fan, Y. Huang, M. Wang, X.
Zhao, G. Hu, H. Yang, J. Wang, J. Wang, X. Xu, L. Song, G. Xu, P. Xu, J. Xu, S. J. O'Brien, L.
Orban, B. Venkatesh and Q. Shi (2016). "The Asian arowana (Scleropages formosus) genome
provides new insights into the evolution of an early lineage of teleosts." Sci Rep **6**: 24501.
Boratyn, G. M., A. A. Schaffer, R. Agarwala, S. F. Altschul, D. J. Lipman and T. L. Madden
(2012). "Domain enhanced lookup time accelerated BLAST." Biol Direct **7**: 12.
Buchfink, B., C. Xie and D. H. Huson (2015). "Fast and sensitive protein alignment using
DIAMOND." Nat Methods **12**(1): 59-60.
Buck, C. B., D. V. Pastrana, D. R. Lowy and J. T. Schiller (2004). "Efficient intracellular
assembly of papillomaviral vectors." J Virol **78**(2): 751-757.
Buck, C. B. and C. D. Thompson (2007). "Production of papillomavirus-based gene transfer
vectors." Curr Protoc Cell Biol **Chapter 26**: Unit 26.21.
Buck, C. B., C. D. Thompson, Y. Y. Pang, D. R. Lowy and J. T. Schiller (2005). "Maturation of
papillomavirus capsids." J Virol **79**(5): 2839-2846.
Buck, C. B., K. Van Doorslaer, A. Peretti, E. M. Geoghegan, M. J. Tisza, P. An, J. P. Katz, J. M.
Pipas, A. A. McBride, A. C. Camus, A. J. McDermott, J. A. Dill, E. Delwart, T. F. Ng, K.
Farkas, C. Austin, S. Kraberger, W. Davison, D. V. Pastrana and A. Varsani (2016). "The
Ancient Evolutionary History of Polyomaviruses." PLoS Pathog **12**(4): e1005574.
de Souza, R. F., L. M. Iyer and L. Aravind (2010). "Diversity and evolution of chromatin
proteins encoded by DNA viruses." Biochim Biophys Acta **1799**(3-4): 302-318.
Dereeper, A., V. Guignon, G. Blanc, S. Audic, S. Buffet, F. Chevenet, J. F. Dufayard, S.
Guindon, V. Lefort, M. Lescot, J. M. Claverie and O. Gascuel (2008). "Phylogeny.fr: robust
phylogenetic analysis for the non-specialist." Nucleic Acids Res **36**(Web Server issue): W465-
469.
Dill, J. A., A. C. Camus, J. H. Leary and T. F. F. Ng (2018). "Microscopic and Molecular
Evidence of the First Elasmobranch Adomavirus, the Cause of Skin Disease in a Giant
Guitarfish, Rhynchobatus djiddensis." MBio **9**(3).
Gerlt, J. A., J. T. Bouvier, D. B. Davidson, H. J. Imker, B. Sadkhin, D. R. Slater and K. L.
Whalen (2015). "Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST): A web tool for
generating protein sequence similarity networks." Biochim Biophys Acta **1854**(8): 1019-1037.
Gouw, M., S. Michael, H. Samano-Sanchez, M. Kumar, A. Zeke, B. Lang, B. Bely, L. B.
Chemes, N. E. Davey, Z. Deng, F. Diella, C. M. Gurth, A. K. Huber, S. Kleinsorg, L. S.
Schlegel, N. Palopoli, K. V. Roey, B. Altenberg, A. Remenyi, H. Dinkel and T. J. Gibson (2018).
"The eukaryotic linear motif resource - 2018 update." Nucleic Acids Res **46**(D1): D428-D434.
Hickman, A. B., D. R. Ronning, R. M. Kotin and F. Dyda (2002). "Structural unity among viral
origin binding proteins: crystal structure of the nuclease domain of adeno-associated virus Rep."
Mol Cell **10**(2): 327-337.
Hildebrand, A., M. Remmert, A. Biegert and J. Soding (2009). "Fast and accurate automatic
structure prediction with HHpred." Proteins **77 Suppl 9**: 128-132.
Iranzo, J., M. Krupovic and E. V. Koonin (2017). "A network perspective on the virus world."
Commun Integr Biol **10**(2): e1296614.
Iyer, L. M., E. V. Koonin, D. D. Leipe and L. Aravind (2005). "Origin and evolution of the
archaeo-eukaryotic primase superfamily and related palm-domain proteins: structural insights
and new members." Nucleic Acids Res **33**(12): 3875-3896.

Katoh, K., J. Rozewicki and K. D. Yamada (2019). "MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization." Brief Bioinform **20**(4): 1160-1166.
Kazlauskas, D., A. Varsani, E. V. Koonin and M. Krupovic (2019). "Multiple origins of
prokaryotic and eukaryotic single-stranded DNA viruses from bacterial and archaeal plasmids."
Nat Commun **10**(1): 3425.
Kim, D., B. Langmead and S. L. Salzberg (2015). "HISAT: a fast spliced aligner with low
memory requirements." Nat Methods **12**(4): 357-360.
Koonin, E. V., V. V. Dolja and M. Krupovic (2015). "Origins and evolution of viruses of
eukaryotes: The ultimate modularity." Virology **479-480**: 2-25.
Koonin, E. V., M. Krupovic and N. Yutin (2015). "Evolution of double-stranded DNA viruses of
eukaryotes: from bacteriophages to transposons to giant viruses." Ann N Y Acad Sci **1341**: 10-
24.
Krupovic, M. and E. V. Koonin (2017). "Multiple origins of viral capsid proteins from cellular
ancestors." Proc Natl Acad Sci U S A **114**(12): E2401-E2410.
Kuraku, S., C. M. Zmasek, O. Nishimura and K. Katoh (2013). "aLeaves facilitates on-demand
exploration of metazoan gene family trees on MAFFT sequence alignment server with enhanced
interactivity." Nucleic Acids Res **41**(Web Server issue): W22-28.
Langmead, B. and S. L. Salzberg (2012). "Fast gapped-read alignment with Bowtie 2." Nat
Methods **9**(4): 357-359.
Li, D., C. M. Liu, R. Luo, K. Sadakane and T. W. Lam (2015). "MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn graph."
Bioinformatics **31**(10): 1674-1676.
Li, D., R. Luo, C. M. Liu, C. M. Leung, H. F. Ting, K. Sadakane, H. Yamashita and T. W. Lam
(2016). "MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced
methodologies and community practices." Methods **102**: 3-11.
Logan, J. and T. Shenk (1984). "Adenovirus tripartite leader sequence enhances translation of
mRNAs late after infection." Proc Natl Acad Sci U S A **81**(12): 3655-3659.
Meier, A. and J. Soding (2015). "Automatic Prediction of Protein 3D Structures by Probabilistic
Multi-template Homology Modeling." PLoS Comput Biol **11**(10): e1004343.
Mizutani, T., Y. Sayama, A. Nakanishi, H. Ochiai, K. Sakai, K. Wakabayashi, N. Tanaka, E.
Miura, M. Oba, I. Kurane, M. Saijo, S. Morikawa and S. Ono (2011). "Novel DNA virus isolated
from samples showing endothelial cell necrosis in the Japanese eel, Anguilla japonica." Virology
412(1): 179-187.
Moyer, C. L., E. S. Besser and G. R. Nemerow (2015). "A Single Maturation Cleavage Site in
Adenovirus Impacts Cell Entry and Capsid Assembly." J Virol **90**(1): 521-532.
Nemerow, G. R., P. L. Stewart and V. S. Reddy (2012). "Structure of human adenovirus." Curr
Opin Virol **2**(2): 115-121.
Nurk, S., D. Meleshko, A. Korobeynikov and P. A. Pevzner (2017). "metaSPAdes: a new
versatile metagenomic assembler." Genome Res **27**(5): 824-834.
Okazaki, S., S. Yasumoto, S. Koyama, S. Tsuchiaka, Y. Naoi, T. Omatsu, S. Ono and T.
Mizutani (2016). "Detection of Japanese eel endothelial cells-infecting virus in Anguilla japonica
elvers." J Vet Med Sci **78**(4): 705-707.
Ondov, B. D., G. J. Starrett, A. Sappington, A. Kostic, S. Koren, C. B. Buck and A. M. Phillippy
(2019). "Mash Screen: high-throughput sequence containment estimation for genome discovery."
Genome Biol **20**(1): 232.

Pastrana, D. V., C. B. Buck, Y. Y. Pang, C. D. Thompson, P. E. Castle, P. C. FitzGerald, S.
Kruger Kjaer, D. R. Lowy and J. T. Schiller (2004). "Reactivity of human sera in a sensitive,
high-throughput pseudovirus-based papillomavirus neutralization assay for HPV16 and HPV18."
<u>Virology</u> **321**(2): 205-216.
Peretti, A., P. C. FitzGerald, V. Bliskovsky, C. B. Buck and D. V. Pastrana (2015). "Hamburger
polyomaviruses." <u>J Gen Virol</u> **96**(Pt 4): 833-839.
Pipas, J. M. (2019). "DNA Tumor Viruses and Their Contributions to Molecular Biology." <u>J</u>
<u>Virol</u> **93**(9).
Rice, P., I. Longden and A. Bleasby (2000). "EMBOSS: the European Molecular Biology Open
Software Suite." <u>Trends Genet</u> **16**(6): 276-277.
Robinson, J. T., H. Thorvaldsdottir, A. M. Wenger, A. Zehir and J. P. Mesirov (2017). "Variant
Review with the Integrative Genomics Viewer." <u>Cancer Res</u> **77**(21): e31-e34.
Ruzindana-Umunyana, A., L. Imbeault and J. M. Weber (2002). "Substrate specificity of
adenovirus protease." <u>Virus Res</u> **89**(1): 41-52.
Schiller, J. T. and D. R. Lowy (2015). "Raising Expectations For Subunit Vaccine." <u>J Infect Dis</u>
211(9): 1373-1375.
Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.
Schwikowski and T. Ideker (2003). "Cytoscape: a software environment for integrated models of
biomolecular interaction networks." <u>Genome Res</u> **13**(11): 2498-2504.
Tisza, M. J., D. V. Pastrana, N. L. Welch, B. Stewart, A. Peretti, G. J. Starrett, Y.-Y. S. Pang, A.
Varsani, S. R. Krishnamurthy, P. A. Pesavento, D. H. McDermott, P. M. Murphy, J. L. Whited,
B. Miller, J. M. Brenchley, S. P. Rosshart, B. Rehermann, J. Doorbar, B. A. Ta'ala, O.
Pletnikova, J. Troncoso, S. M. Resnick, A. M. Segall and C. B. Buck (2019). "Discovery of
several thousand highly diverse circular DNA viruses." 555375.
Van Doorslaer, K., Z. Li, S. Xirasagar, P. Maes, D. Kaminsky, D. Liou, Q. Sun, R. Kaur, Y.
Huyen and A. A. McBride (2017). "The Papillomavirus Episteme: a major update to the
papillomavirus sequence database." <u>Nucleic Acids Res</u> **45**(D1): D499-d506.
Vragniau, C., J. M. Hubner, P. Beidler, S. Gil, K. Saydaminova, Z. Z. Lu, R. Yumul, H. Wang,
M. Richter, P. Sova, C. Drescher, P. Fender and A. Lieber (2017). "Studies on the Interaction of
Tumor-Derived HD5 Alpha Defensins with Adenoviruses and Implications for Oncolytic
Adenovirus Therapy." <u>J Virol</u> **91**(6).
Wen, C. M., M. M. Chen, C. S. Wang, P. C. Liu and F. H. Nan (2015). "Isolation of a novel
polyomavirus, related to Japanese eel endothelial cell-infecting virus, from marbled eels,
Anguilla marmorata (Quoy & Gaimard)." <u>J Fish Dis</u>.
Yutin, N., S. Shevchenko, V. Kapitonov, M. Krupovic and E. V. Koonin (2015). "A novel group
of diverse Polinton-like viruses discovered by metagenome analysis." <u>BMC Biol</u> **13**: 95.
Zallot, R., N. O. Oberg and J. A. Gerlt (2018). "'Democratized' genomic enzymology web tools
for functional assignment." <u>Curr Opin Chem Biol</u> **47**: 77-85.
Zimmermann, L., A. Stephens, S. Z. Nam, D. Rau, J. Kübler, M. Lozajic, F. Gabler, J. Söding,
A. N. Lupas and V. Alva (2017). "A Completely Reimplemented MPI Bioinformatics Toolkit
with a New HHpred Server at its Core." <u>J Mol Biol</u> **S0022-2836**(17): 30587-30589.