tDCS over the left prefrontal cortex does not affect time-trial self-paced cycling

performance: Evidence from oscillatory brain activity and power output.

Darías Holgado*[1,2], Thomas Zandonai[2], Luis F. Ciria[2], Mikel Zabala[1], James Hopker[3], Daniel

Sanabria*[2]

[1]Department of Physical Education and Sport, Faculty of Sport Sciences, University of Granada,

[2]Mind, Brain and Behaviour Research Centre, Department of Experimental Psychology, University of Granada,

[3]School of Sport and Exercise Sciences, Endurance Research group, University of Kent, UK.

Corresponding Author

[a]Mind, Brain and Behaviour Research Centre

University of Granada

Campus Universitario de Cartuja s/n

18011 Granada (Spain)

Telf : +34 958247875

dariashn@ugr.es; daniel@ugr.es

Abstract

Transcranial direct current stimulation (tDCS) is a non-invasive brain stimulation technique, which is increasing in popularity amongst sport scientists. The rationale of using tDCS as a tool in sport is that stimulating brain areas related to exercise performance would make athletes to boost their physical performance. Here, in a pre-registered, (https://osf.io/rf95j/), we tested the hypothesis that tDCS over the left dorsolateral prefrontal cortex (DLPFC) influences performance in a 20' time-trial self-paced exercise and electroencephalographic (EEG) oscillatory brain activity. 36 trained male cyclists completed a 20' time-trial self-paced exercise in three separate sessions, corresponding to three stimulation conditions: anodal, cathodal and sham. tDCS was administered before each test during 20' at a current intensity of 2.0 mA. In each session, power output, heart rate, rate of perceived exertion (RPE) and EEG (at baseline and during exercise) was measured. We found that neither anodal, cathodal nor sham improved performance, affected heart rate, RPE or EEG activity. Our data suggest that the effects of tDCS on endurance performance should be taken with caution, with the brain region stimulated being an important factor.

Introduction

Self-paced exercise refers to a physical activity in which the effort needs to be evenly distributed and monitored in order to complete the task without reaching premature exhaustion [1]. Performance in self-paced exercise is undoubtedly related to the functioning of peripheral body systems, such as the muscles, heart, lungs etc., as well as the brain. In this respect, self-pacing during exercise is a challenging cognitive task [2], as it requires constant control and monitoring of internal (e.g., heart rate) and external inputs (e.g., a bump on the road while cycling), while maintaining the goals of the task (e.g. completing a set distance as fast as possible). In other words, self-paced exercise can be regarded as an executive task, with high demands of self-control, goal-monitoring and inhibition [3].

Research in cognitive neuroscience has long pointed to the prefrontal cortex as a key brain area involved in executive processing [4]. Interestingly, the few neuroimaging studies testing participants while exercising have shown activation of the prefrontal cortex, together with the expected sensorio-motor recruitment [5], which reinforces the hypothesis of the crucial role of executive processing on self-paced exercise. The rationale of the present study was therefore that stimulation of the prefrontal cortex via transcranial direct current will affect self-paced exercise performance.

Transcranial direct current stimulation (tDCS) is a non-invasive electrical brain stimulation technique that is able to induce cortical changes by depolarizing (anodal) or hyperpolarizing (cathodal) a neuron's resting membrane potential [6]. Recently, there have been an increasing interest in the use of tDCS to enhance endurance performance [7]. For example, Angius et al. [8] and Vitor-Costa et al. [9] found an increased time to exhaustion (TTE) in a cycling test after acute stimulation of the primary motor cortex (M1). Angius et al. [8] attributed that performance enhancement to a reduction of the perceived effort (RPE), although Vitor-Costa et al. [9] did not find such a reduction perceived exertion. These apparently contradictory results leave open the question of whether tDCS affects people's RPE when stimulating the motor cortex. Meanwhile, Okano et al. [10] found improved cycling performance (greater peak power output) in the anodal condition than in the sham condition after stimulating the temporal cortex of ten trained cyclists. The authors argued that their anodal condition might have influenced activity in the insular cortex, which has been linked to autonomic regulation and to self-perception and awareness of body sensations [11]. Most of research on the effect of tDCS on endurance performance has hitherto been focused on activation or inhibition of the motor and temporal cortices. To the best of our knowledge, only two studies have targeted the prefrontal cortex. Lattari et al. [12] found increased exercise tolerance in a TTE at 100% of the peak power after stimulating the left dorsolateral prefrontal cortex for 20' in eleven physically active women. This improvement was not accompanied by a reduction in the RPE. Meanwhile, Borducchi et al. [13] found an improvement in cognitive performance and mood in elite athletes of different sport modalities (n = 10) after ten days of anodal stimulation over the left dorsolateral prefrontal cortex, which, according the authors, may contribute to

performance gains, greater well-being and faster recovery. However, due to the lack of a control

condition (Borducchi et al.) and small sample sizes in their studies (like in almost every previous study

on tDCS and sport performance), the above results should be considered with caution.

The present (pre-registered, https://osf.io/rf95j/) research is novel as it is the first to directly test the

hypothesis that stimulation of the prefrontal cortex would affect performance in a 20' time-trial self-

paced exercise bout in trained male cyclists. More precisely, we expected that activation via anodal

stimulation would improve performance, whilst inhibition of the prefrontal cortex via cathodal

stimulation would impair performance (compared to a sham condition). The indexes of physical

performance were the power output during exercise and the RPE after the self-paced exercise.

Additionally, we asked participants to perform an executive task [14] after the exercise. The purpose was

to test the hypothesis that any change on physical performance produced by the tDCS over the

prefrontal cortex would modulate the subsequent (known [15]) effect of exercise on inhibitory control.

This is in line with the idea of a bi-directional relationship between exercise, brain and cognition [15],

i.e., brain and cognitive functioning influences exercise performance and vice versa. Brain electrical

activity was measured at rest, during exercise, and during the cognitive task by recording

electroencephalography (EEG) in order to examine the effects of tDCS at brain level. Even though the

literature over the effect of tDCS on EEG is scarce and inconclusive [16], we anticipated an increase in

the alpha and beta band after stimulation in the anodal condition compared to cathodal and sham

condition.

Results

The intervention was well tolerated and participants only reported common side effects such as a

tingling (anodal: 22%, cathodal: 8% and sham: 11%), or itchy sensation in the scalp (anodal: 30%,

cathodal: 8% and sham: 16%).

Exercise performance

The average power output during the time trial self-paced exercise was not significantly different

$(F(2,34) = 0.31, p > 0.05)$ between conditions (see Fig. 1): Anodal (235 W [95%CI 222 - 249 W];

Cathodal (235 W [95%CI 222 - 248 W] and Sham (234 W [95%CI 220 - 248 W].

The heart rate signal for three participants was lost during the 20' time-trial self-paced exercise,

consequently they were removed from the subsequent analysis (n= 33). The average heart rate during

the time trial was not significantly different $(F(2,34) = 1.02, p > 0.05)$ between conditions: Anodal

(161 beats min−1 [95%CI 157 - 166 beats min−1]; Cathodal (162 beats min−1 [95%CI 158 - 167

beats min−1] and Sham (162 beats min−1 [95%CI 157 - 167 beats min−1].

Post time-trial sRPE did not show any significant differences between conditions: Anodal (17.02

[95%CI 16.5 - 17.5]; Cathodal (17 [95%CI 16.8 - 17.4] and Sham (17.02 [95%CI 16.5 - 17.5], F(2,34)

= 1.69; p > 0.05.

Electrical brain activity (EEG)

Due to excessive noise in the EEG signal, five participants were not included in the EEG analysis (n=

31). The analysis of tonic spectral power (see Fig. 2) did not provide any significant difference (all ps

> 0.05) between conditions (anodal, cathodal and sham), and for each period of time (baseline-pre;

baseline-post, warm-up, self-paced exercise and recovery).

The event-related spectral perturbation (stimulus-locked) analysis in the flanker task (see Fig. 3) did

not reveal any main effect of condition for the congruent or incongruent trial (both ps > 0.05).

Executive task

A main effect of stimulus was reported in the flanker task, with participants being less accurate (M=

98 vs 91 % correct responses; F(2,34) = 13.17, p < 0.01) and slower (423 vs 515 ms; F(2,34) = 182.39,

p < 0.05) in the incongruent stimulus compared to the congruent stimulus. There were no significant

differences between conditions for the congruent and incongruent target, for RT and accuracy (Fs < 1,

all ps > 0.05).

Discussion

To the best of our knowledge, this is the first study testing the influence of prefrontal cortex tDCS'

stimulation on self-paced exercise and brain activity during exercise. The main finding of this study

was that 20' anodal or cathodal tDCS' stimulation (relative to sham) over the left dorsolateral

prefrontal cortex did not affect exercise performance or brain electrical activity. Moreover, neither

sRPE, EEG or cognitive performance were affected by the stimulation.

Our findings indicated that anodal or cathodal tDCS applied over the left dorsolateral prefrontal cortex

before exercise did not modulate exercise performance during a 20' time-trial self-paced exercise. This

finding contrast the results of the two previous studies testing the effect of tDCS over the same brain

area [12,13], as well as previous studies reporting positive effects of tDCS. In our view, several reasons

might explain the mixed results (including ours) present in the literature: 1) the experimental protocol.

While Lattari et al. [12], Angius et al. [8] and Vitor-Costa et al. [9] used a TTE, and Okano et al.[10] a

maximal incremental test, Barwood et al.[17] and our study used a time-trial self-paced exercise test.

Due to TTE tests and test based on time-trial self-paced present different physiological and

psychological demands, these divergent results might be partially explained by the protocol chosen[18].

2) The selection of the electrode montage. To date, M1 is the most recurrent stimulated area. It is
suggested that M1 stimulation is likely to increase power output during exercise or to reduce RPE for
a given force or power [8]. It could then be argued that tDCS would only affect physical performance
when applied to M1. However, there is a previous study targeting the left prefrontal cortex finding
significant results [12]; and another stimulating the temporal cortex, [10], which may jeopardize the
hypothesis that only stimulation of M1 leads to positive results. 3) Sample size. To date, the vast
majority of the tDCS' studies in the sport science area present low samples sizes. A statistically
significant effect from a study with a small sample size (and low statistical power) may easily reflect a
false positive[19]. 4) Effect size. It may well be possible that the actual size of the effect (if any) of tDCS
on physical (cycling in this particular case) performance is rather small, so that with our sample size
we might have not been able to find it. However, note that we a priori calculated the size of our sample
according to the previous literature and to practical/applied interest. 5) Stimulation intensity. During
the process of writing this manuscript, an interesting paper has been published[20] showing that the
intensity of 2mA (the common tDCS intensity used in previous sport science studies) does not seem
enough to affect neuronal circuits. The authors argued that at least a current of 4.5 mA is necessary to
affect neural circuits because a significant fraction of the current is lost due to skin and soft tissue and
the resistance of the skull. Thus, the use of 2mA in our study (like in every other tDCS-exercise study
published to date) may well explain why we did not obtain a significant effect of tDCS in brain
oscillatory activity or physical performance. 6) Last, but not least, we believe that the "file drawer
effect" (i.e., the tendency to only publish positive outcomes) might be biasing the literature to positive
findings[21].

The hypothesis that anodal would increase EEG amplitude was not confirmed in the present study.
After the 20' stimulation, the EEG spectral power was similar across all condition for each period of
time. This null effect could be again explained by the low intensity of the stimulation. Vöröslakos et
al.[20] found that currents between 4-6 mA should be delivered to modulate EEG amplitude.

The rationale of including the flanker task after the cycling self-paced exercise was that any change in
physical performance and brain activity via tDCS would modulate the subsequent influence of cycling
on cognitive (inhibition) performance. The lack of differences in physical exertion, RPE and EEG
between the three experimental conditions make reasonable to have found no difference in RT or
accuracy as a function of tDCS.

Conclusions

tDCS is an increasingly popular technique used within a wide range of settings, from treatment of
neurological disorders, to attempting to improve exercise performance. Our data, however, add further
to the mixed evidence in this area, challenging the idea that an acute session of tDCS can improve

physical performance. At this point, we believe that research on this topic will benefit from further
empirical and meta-analytical research, in order to accumulate evidence on whether an acute session
of tDCS affect sport performance or not.

Methods

Following institutional ethical approved by the University of Granada Ethics Committee
(287/CEIH/2017), a randomized, sham-controlled, within-subject experimental design was conducted
on male trained and triathletes cyclists. All experimental procedures were designed to comply with the
Declaration of Helsinki. Before being recruited, participants provided written informed consent having
previously read a participant information sheet. All data were entered in a case report form, and
subsequently in a computerized database and stored at the Mind, Brain and Behaviour Research
Centre (MBBRC) of the University of Granada. Exclusion criteria was the presence of symptomatic
cardiomyopathy, metabolic disorders such as obesity (BMI >30) or diabetes, chronic obstructive
pulmonary disease, epilepsy, therapy with b-blockers and medications that would alter cardiovascular
function, hormonal therapy, smoking, and neurological disorders, as well as the presence of implanted
metal devices (e.g., pacemakers, metal plates, wires).

The method and planned analyses of this study were pre-registered on the Open Science Framework
(OSF). This was done on June 29, 2017, and can be found at https://osf.io/rf95j/. The raw files can be
found in OSF.

Previous research has shown that the effect of tDCS on exercise performance is small-medium.
Additionally, we considered that a medium effect would be appropriate in terms of the potential future
practical application of the findings from this type of research to elite cyclists. Therefore, according to
the G*Power software[22], 36 participants were required for a power of .8 and a medium effect size,
(partial eta-squared $\eta^2 = 0.13$) for a 3 conditions (anodal, cathodal, sham) design. During the data
collection, two of the participants could not complete the three experimental sessions and were
replaced by two further participants. Accordingly, data collection stopped when complete datasets
(successful completion of all three condition) were obtained for 36 endurance trained cyclists and
triathletes. The physiological characteristic of the participants are (mean and SD): age = 27 (6.8)
years, weight = 70.1 (9.5) Kg; VO_{2max} = 54 (6.13) ml.min^{-1}.kg^{-1}.

Participants visited the MBBRC four times (one screening visit and three experimental sessions).
Participants initially attended the MBBRC for a screening visit. After verifying that the participants
met the inclusion criteria, they performed a maximal incremental exercise test in order to identify their
maximal oxygen consumption using a standard laboratory protocol[23]. After completing the maximal
incremental test, participants performed a 10' time-trial self-paced exercise test in order to familiarise
themselves with the protocol to be used in subsequent visits. The shorter duration of the

familiarization test (with respect to the proper experimental self-paced exercise) might be seen as a

limitation of our study. Nevertheless, that was motivated for two reasons: 1) our participants were

experienced cyclists used to performing self-paced exercise, and given their expertise, the purpose was

that of familiarize them with the laboratory setting testing procedure; 2) we were aware that the 10'

test was performed after the maximal incremental exercise test and participants were already fatigued.

After the screening visit, participants attended the lab on three separate occasions to perform the 20'

time-trial acute self-paced exercise (all procedures were the same, except for the stimulation

condition). Participants were asked to refrain from drinking alcohol (48 h abstinence) and caffeine (24

h abstinence) and instructed not to perform any exhaustive exercise in the 48 h before each

experimental session. Participants were also asked to keep their pre-exercise meal the same for every

session. The experimental sessions were completed at the same time of the day to avoid diurnal

variations. EEG was recorded throughout the session, except for the stimulation period. Before the

beginning of the stimulation, we recorded 5' EEG with open-eyes as a baseline measure. After the

baseline measure, we delivered 20 minutes' of tDCS stimulation (anodal, cathodal or sham, depending

on the counterbalanced order). Next, we repeated the 5' baseline EEG measure with open-eyes. After

that, participants performed the 20' self-paced exercise preceded by 5' warm-up (at 120 watts) on the

cycle ergometer (SRM, Julich, Germany). During the data collection, the SRM broke and we had to

replace it for a Phantom 5 ergometer (CyleOps, Madison, USA). The Phantom 5 measure the power

output using an on-board power meter PowerTap (PowerTap, Madison, USA) with power accuracy of

+/- 1.5%. Every participant completed the time-trial self-paced exercise on the same ergometer:

seventeen participants completed the trial on the SRM and nineteen on the Phantom 5. Participants

were instructed to achieve the highest average power possible during time-trial self-paced exercise and

were freely able to change gearing and cadence throughout. Participants were aware of the elapsed

time, but they did not have feedback on performance (wattage and heart rate) during, or after the self-

paced exercise. Heart rate was measured continuously throughout the protocol (V800, Polar Electro,

Finland). Immediately after exercise, we asked the participant to rate their session RPE (sRPE)[24].

Finally, participants completed a 5' cool-down and the executive task. The interval between the

different sessions was at least 48h to allow the full recovery and to minimize carryover effects.

Stimulation was delivered using battery powered DC stimulators (Newronika S.r.l, Milan, Italy) and

delivered through a saline soaked pair of surface sponge electrodes (5 x 5 cm). For the anodal

(increased excitability) or cathodal (decreased excitability) we targeted the prefrontal cortex. The

anode or cathode electrode was placed over F3 area according to the international EEG 10-20 system

[25]. The opposite electrode was placed over the contralateral shoulder area in order to avoid the delivery

of current on the participant's scalp. Current was set at 2 mA and was delivered for 20', which has

previously been shown to provoke cortical changes [26]. The sham stimulation (control) was similar to

the anodal and cathodal stimulation but the device only provided 2mA for 30s after which is was

turned off without the participant's awareness. This method replicates the sensory feelings

experienced in the tDCS trial (i.e. itching and tingling sensations) and cannot be distinguished from it,

whether the stimulation is continued or stopped [27]. The EEG cap was kept over the sponges during

stimulation period, but the EEG activity was not recorded. At the end of the session (after completing

the cognitive task), participants answered a questionnaire regarding their experience during and after

the tDCS sessions [28].

Participants completed a modified flanker task [14], via use of computer software (E-Prime, Psychology

Software Tools, Pittsburgh, PA, USA), to assess inhibitory control, a form of executive processing

after the self-paced exercise. Here, the flanker task involves the response to the direction of a central

arrow surrounded by other arrows pointing in the same or opposite direction. Congruent trials consist

of a central target arrow being flanked by other arrows that faced the same direction (e.g., <<<<< or

>>>>>). The incongruent trials consist of the target arrow being flanked by other arrows that faced the

opposite directions (e.g., <<><< or >><>>). Participants pressed a button with their left index finger

when the target arrow (regardless of condition) faced to the left (e.g., '<') and a button with their right

index finger when the target arrow faced to the right (e.g., '>'). Each trial started with the presentation

of a cross (fixation point) that remained on a steady until the appearance of the target arrows 2 seconds

later. The target was presented in the middle of the screen for 150 ms and a response window of 1350

ms was allowed. The next trial started 1500 ms after the response. Total task duration was

approximately 7'. Participants completed one block of 160 trials with equal probability for congruent

and incongruent trials, randomized across task conditions. A brief familiarization of the task was

included in the screening visit. For each stimulus, the RT (in ms) and response accuracy (percentage of

correct responses) were recorded.

EEG were recorded at 1000 Hz using a 30-channel actiCHamp System (Brain Products GmbH,

Munich, Germany) with active electrodes positioned according to the 10-20 EEG International System

and referenced to the Cz electrode. The cap was adapted to individual head size, and each electrode

was filled with Signa Electro-Gel (Parker Laboratories, Fairfield, NJ) to optimize signal transduction.

Participants were instructed to avoid body movements as much as possible, and to keep their gaze on

the centre of a computer screen during the measurement. Electrode impedances were kept below 10

kΩ. EEG pre-processing was conducted using custom Matlab scripts and the EEGLAB and Fieldtrip

Matlab toolboxes. EEG data were resampled at 500 Hz, bandpass filtered offline from 1 and 40 Hz to

remove signal drifts and line noise, and re-referenced to a common average reference. Horizontal

electrooculograms (EOG) were recorded by bipolar external electrodes for the offline detection of

ocular artefacts. Independent component analysis was used to detect and remove EEG components

reflecting eye blinks.

All analyses were completed using statistical nonparametric permutation tests with a Monte Carlo approach. These tests do not make any assumption of the underlying data distribution, are unbiased, and as efficient and powerful as parametric statistics. When statistical significance ($p < 0.05$) was found, values were corrected by the false discovery rate method. The effect of experimental condition (anodal, cathodal, sham) on self-paced exercise power output, heart rate and RPE were analysed using a within-subject design condition.

Spectral power was analysed using a within-participants' design with the factor of stimulation (anodal, cathodal, sham). Each period (Baseline, Warming Up, Exercise, Cooling Down) was tested separately for significance. In the absence of strong a priori hypotheses over the frequency range and channels which tDCS may induce a change, we use a stepwise, cluster-based, non-parametric permutation test [29].

For the cognitive task, we analysed the event-related spectral perturbation main effects of stimulation (anodal, cathodal, sham) for each stimulus (congruent, incongruent) by applying the cluster-based approach [30]. In order to reduce the possibility that the type II error rate was inflated by multiple comparisons correction, we set an a priori criteria of collapsing data into four frequency bands: Theta (4–8 Hz), Alpha (8–14 Hz), lower Beta (14–20 Hz) and upper Beta 1 (20–40 Hz). To avoid an overlap with behavioural responses, we also limited the time windows of interest to the first 500 ms after the stimuli onset (based on average behavioural response times) for target and standard trials, respectively. This analysis is similar to other researches published by our lab and further detailed information of the method can be found in the papers [29].

Practical applications

The use of tDCS is increasing in popularity in sport science

tDCS over the left prefrontal cortex does not improve performance in trained cyclists

tDCS does not seem to change EEG activity at rest or during exercise

References

1. Roelands, B., De Koning, J., Foster, C., Hettinga, F. & Meeusen, R. Neurophysiological determinants of theoretical concepts and mechanisms involved in pacing. *Sports Medicine* **43,** 301–311 (2013).

2. Walsh, V. Is sport the brain's biggest challenge? *Current Biology* **24,** R859–R860 (2014).

3.   Brick, N. E., MacIntyre, T. E. & Campbell, M. J. Thinking and action: a cognitive perspective on

self-regulation during endurance performance. *Frontiers in Physiology* **7,** (2016).

4.   Diamond, A. Executive Functions. *Annual review of clinical psychologyPsychol.* **64,** 135–168

(2014).

5.   SCHNEIDER, S., ASKEW, C. D., ABEL, T., MIERAU, A. & STRÜDER, H. K. Brain and

Exercise. *Medicine & Science in Sports & Exercise* **42,** 600–607 (2010).

6.   Brunoni, A. R. *et al.* Clinical Research with Transcranial Direct Current Stimulation (tDCS):

Challenges and Future Directions. *Brain stimulation* **5,** 175–195 (2013).

7.   Angius, L., Hopker, J. & Mauger, A. R. The ergogenic effects of transcranial direct current

stimulation on exercise performance. *Frontiers in Psychology* **8,** (2017).

8.   Angius, L. *et al.* Bilateral extracephalic transcranial direct current stimulation improves endurance

performance in healthy individuals. *Brain Stimulation: Basic, Translational, and Clinical*

Research in Neuromodulation **0,** (2017).

9.   Vitor-Costa, M. *et al.* Improving Cycling Performance: Transcranial Direct Current Stimulation

Increases Time to Exhaustion in Cycling. *PLoS One* **10,** e0144916 (2015).

10. Okano, A. H. *et al.* Brain stimulation modulates the autonomic nervous system, rating of

perceived exertion and performance during maximal exercise. *British journal of sports medicine*

49, 1213–8 (2015).

11. Kimmerly, D. S., O'Leary, D. D., Menon, R. S., Gati, J. S. & Shoemaker, J. K. Cortical regions

associated with autonomic cardiovascular regulation during lower body negative pressure in

humans. *The Journal of Physiology* **569,** 331–345 (2005).

12. Lattari, G. A. M., Eduardo de Oliveira, Bruno Soares Oliveira, Bruno Ribeiro Ramalho de Mello

Pedreiro, Rodrigo Cunha Machado, Sérgio Neto. Effects of transcranial direct current stimulation

on time limit and ratings of perceived exertion in physically active women. *Neuroscience Letters*

662, 12–16 (2017).

13. Borducchi, D. M. M. *et al.* Transcranial Direct Current Stimulation Effects on Athletes' Cognitive

Performance: An Exploratory Proof of Concept Trial. *Frontiers in Psychiatry* **7,** 183 (2016).

14. Eriksen, B. A. & Eriksen, C. W. Effects of noise letters upon the identification of a target letter in

a nonsearch task. *Perception & Psychophysics* **16,** 143–149 (1974).

15. Loprinzi, P. D., Herod, S. M., Cardinal, B. J. & Noakes, T. D. Physical activity and the brain: A

review of this dynamic, bi-directional relationship. *Brain Research* **1539,** 95–104 (2013).

16. Mangia, A. L., Pirini, M. & Cappello, A. Transcranial direct current stimulation and power

spectral parameters: a tDCS/EEG co-registration study. *Frontiers in human neuroscience* **8,** 601

(2014).

17. Barwood, M. J. *et al.* The Effects of Direct Current Stimulation on Exercise Performance, Pacing

and Perception in Temperate and Hot Environments. *Brain Stimulation* **9,** 842–849 (2016).

18. Marino, F. E. The limitations of the constant load and self-paced exercise models of exercise

physiology. *Comparative Exercise Physiology* **8,** 3–9 (2012).

19. Button, K. S. *et al.* Power failure: why small sample size undermines the reliability of

neuroscience. *Nature reviews. Neuroscience* **14,** 365–376 (2013).

20. Vöröslakos, M. *et al.* Direct effects of transcranial electric stimulation on brain circuits in rats and

humans. *Nature Communications* **9,** 1–17 (2018).

21. Simonsohn, U., Nelson, L. D. & Simmons, J. P. P-curve: a key to the file-drawer. *Journal of*

experimental psychology. General **143,** 534–547 (2014).

22. Faul, F., Erdfelder, E., Buchner, A. & Lang, A.-G. Statistical power analyses using G*Power 3.1:

tests for correlation and regression analyses. *Behavior research methods* **41,** 1149–60 (2009).

23. Lucía, A. *et al.* Frequency of the V·O 2max Plateau Phenomenon in World-Class Cyclists.

International Journal of Sports Medicine **27,** 984–992 (2006).

24. Borg, G. Psychophysical bases of perceived exertion. *Medicine & Science in Sports & Exercise*

14, 377–381 (1982).

25. Schönfeldt-lecuona, C., Herwig, U., Satrapi, P. & Schönfeldt-lecuona, C. Using the International

10-20 EEG System for Positioning of Transcranial Magnetic Stimulation Using the International

10-20 EEG System for Positioning of Transcranial Magnetic Stimulation. *Brain topography* **16,**

95–99 (2003).

26. Nitsche, M. A. & Paulus, W. Sustained excitability elevations induced by transcranial DC motor

cortex stimulation in humans. *Neurology* **57,** 1899–1901 (2001).

27. Kessler, S. K., Turkeltaub, P. E., Benson, J. G. & Hamilton, R. H. Differences in the experience of

active and sham transcranial direct current stimulation. *Brain Stimulation* **5,** 155–162 (2012).

28. Aparicio, L. V. M. *et al.* A Systematic Review on the Acceptability and Tolerability of

Transcranial Direct Current Stimulation Treatment in Neuropsychiatry Trials. *Brain Stimulation*

9, 671–681 (2016).

29. Holgado, D. *et al.* Tramadol effects on physical performance and sustained attention during a 20-

min indoor cycling time-trial: A randomised controlled trial. *Journal of Science and Medicine in*

Sport (2017). doi:10.1016/j.jsams.2017.10.032

30. Pesarin, F., Salmaso, L. & Wiley InterScience (Online service). *Permutation tests for complex*

data : theory, applications, and software. (Wiley, 2010).

Acknowledgment

The authors thank Newronika S.L. for lending their tDCS device. The company was not involved in the study.

This project was supported by grants from from the Spanish Ministerio de Economía, Industria y Competitividad-PSI2016-75956-P- to Daniel Sanabria and Mikel Zabala, a predoctoral grant from the Spanish Ministerio de Economía, Industria y Competitividad (BES-2014-069050) to Luis Ciria, and a Spanish "Ministerio de Educación, Cultura y Deporte" predoctoral grant (FPU14/06229) to Darías Holgado.

Author contributions

371 DH, TZ, JH, MZ and DS designed the study; DH and TZ collected the data; DH and LC analysed the

372 data; DH wrote the manuscript under the supervision of DS; DH and LC prepared the figures. All

373 authors reviewed the manuscript before submission.

374 Additional information

375 **Competing interests:** The author(s) declare no competing interests.

376 **Data availability**: EEG data can be found in http://doi.org/10.5281/zenodo.1254077

377 Figure Legends

Figure 1. Power output (watts) profile for the 20' self-paced exercise. Data are means and 95%CI.

Figure 2. A) Differences in brain power spectrum as a function of tDCS condition. Average EEG

power spectrum across participants among anodal (blue lines), cathodal (red line) and sham (black

lines) condition at baseline pre, baseline post and exercise period. The shaded lines denote the average

tonic spectral power for each participant and condition (given that there were not significant

differences between conditions, the lines tend to overlap). B) Parametric F-test colormap comparing

the relative power across frequency (x-axes) and channels (y-axes). Note that the analysis of the other

periods (warmup and recovery) did not yield significant between-intensity differences.

Figure 3. Event-related spectral perturbation of flanker task. Time-locked spectral power averaged

over all electrodes for each condition. Each panel illustrates time-frequency power across time (x-

axes) and frequency (y-axes) for the congruent and incongruent stimuli (blue: decreases; red:

increases). Dashed vertical line represents stimulus onset.