ESSENTIAL ROLES OF THE *SPPRA* FRUCTOSE-PHOSPHATE

PHOSPHOHYDROLASE OPERON IN CARBOHYDRATE METABOLISM AND

VIRULENCE EXPRESSION BY *STREPTOCOCCUS MUTANS*

Lin Zeng and Robert A. Burne[#]

Department of Oral Biology, University of Florida, Gainesville, Florida, USA

Running title: Adaptation to sugar-phosphate stress

Key words: fructose metabolism, sugar-phosphate stress, xylitol, biofilm, sugar-

phosphate phosphohydrolase

[#]Corresponding Author:

1395 Center Drive, Gainesville, Florida 32610

Phone: 352 273-8850

FAX: 352 273-8829

Email: rburne@dental.ufl.edu

Abstract

The dental caries pathogen *Streptococcus mutans* can ferment a variety of sugars to produce organic acids. Exposure of *S. mutans* to certain non-metabolizable carbohydrates (e.g., xylitol) impairs growth and can cause cell death. Recently, the presence of a sugar-phosphate stress in *S. mutans* was demonstrated using a mutant lacking 1-phosphofructokinase (FruK) when fructose was present in the environment. Here we studied an operon in *S. mutans, sppRA,* which was highly expressed in the *fruK* mutant. Biochemical characterization of a purified, recombinant SppA protein indicated that it possessed hexose-phosphate phosphohydrolase activity, with preferences for fructose-1-phosphate (F-1-P) and fructose-6-phosphate (F-6-P). SppA activity was stimulated by Mg^{2+} and Mn^{2+}, but inhibited by NaF. SppR, a DeoR-family regulator, repressed the expression of the *sppRA* operon in the absence of fructose-derived metabolites. Constitutive expression of *sppA,* via a plasmid, greatly alleviated fructose-induced stress in a *fruK* mutant, and enhanced resistance to xylitol. Furthermore, accumulation of F-1-P not only induced *sppA* expression, it significantly altered biofilm maturation, increased cell lysis and enhanced extracellular DNA release. Finally, by identifying three additional putative phosphatases that are capable of promoting sugar-phosphate tolerance, we show that *S. mutans* is capable of mounting a sugar-phosphate stress response by modulating the levels of certain glycolytic intermediates, functions that are interconnected with the ability of the organism to manifest key virulence behaviors.

Importance

Streptococcus mutans is a major etiologic agent for dental caries, primarily due to its ability to form biofilms on tooth surface and to convert carbohydrates into organic acids. We have discovered a two-gene operon in *S. mutans* that regulates fructose metabolism by controlling the levels of fructose-1-phosphate, a potential signaling compound that affects bacterial behaviors. With fructose becoming increasingly common and abundant in the human diet, we reveal the ways fructose may alter bacterial development, stress tolerance, and microbial ecology in the oral cavity to promote oral diseases.

Introduction

Most bacteria use the sugar:phosphotransferase system (PTS) to internalize mono- and di-saccharides, a process that uses phosphoenolpyruvate (PEP) to energize the transport and concomitant phosphorylation of the carbohydrates (1). While certain components of the PTS are vital to the growth and persistence of many bacteria, the accumulation in cells of phosphorylated carbohydrates can lead to sugar-phosphate stress, a condition that often manifests in growth inhibition or cell death (2-4). A number of mechanisms have been proposed to contribute to sugar-phosphate stress: (i) futile cycling, in which the cells expend energy to internalize and phosphorylate the sugar, but the sugar is then dephosphorylated and expelled (5); (ii) reduction of essential metabolic intermediates arising from an inability to internalize other carbohydrates (6); (iii)

creation of toxic metabolic byproducts, such as methylglyoxal (7, 8), or (iv)

deleterious effects of specific sugar phosphates that act as allosteric modifiers of

enzymes or regulatory proteins, causing dysregulation of metabolism (9). The

majority of research on sugar-phosphate stress has been conducted with Gram-

negative, model organisms, with far fewer studies carried out with known

pathogens or Gram-positive bacteria (10).

Streptococcus mutans, a primary dental caries pathogen, is a Gram-positive,

lactic acid-producing bacterium that depends on, and is particular adept at,

transporting and fermenting carbohydrates for energy production and growth.

The most studied sugar-phosphate stress in *S. mutans* is that induced by xylitol,

a non-metabolizable pentitol that is internalized as xylitol-5-phosphate (X-5-P) by

a fructose-specific PTS, but is subsequently dephosphorylated and released from

the cells (11-13). Although the exact nature of the process remains to be

clarified, including the enzyme(s) responsible for removing the phosphate group

from X-5-P, it is generally believed that both the futile cycling of xylitol and the

presence of X-5-P in cells are detrimental. While futile cycling wastes energy, it

has been proposed that X-5-P is an inhibitor of phosphofructokinase that causes

a block in glycolysis (9). The observed effects of xylitol intake by *S. mutans*

include reductions in high-energy metabolites, such as fructose-1,6-bisphosphate

(F-1,6-bP) and glucose-6-phosphate (G-6-P), as well as impaired glucose-PTS

activity, decreased rates of acid production (14), and inhibition of the formation of

extracellular insoluble polysaccharides from sucrose (15, 16). Given its ability to

negatively affect a caries pathogen, xylitol is widely used as a non- or anti-

cariogenic additive in products that include chewing gum and candies, despite

the fact that spontaneous emergence of xylitol-resistant mutants can occur in

habitual users of xylitol-containing products (17, 18). It has been shown that the

fructose-PTS permease encoded by the *fruI* gene is responsible for transporting

the majority of xylitol, and a *fruI*-deficient mutant lost sensitivity to presence of

xylitol (18). Another example of sugar-phosphate-related stress in *S. mutans* is

one induced by 2-deoxyglucose (2-DG), an analog of glucose that can be

internalized and phosphorylated by the PTS, but cannot be metabolized. Uptake

of 2-DG by *S. mutans* reduces the transport of carbohydrates via the PTS (19)

and the multiple sugar metabolism (*msm*) system, an ATP-binding cassette

(ABC) transporter for melibiose, raffinose, and certain other carbohydrates that

are found in the human diet (20).

Fructose is transported into *S. mutans* via multiple PTS permeases and further

metabolism requires phosphofructokinase enzymes, including a 1-

phosphofructokinase and 6-phosphofructokinase that convert fructose-1-

phosphate (F-1-P) and fructose-6-phosphate (F-6-P), respectively, into F-1,6-bP

that can be further catabolized via glycolysis (3). We recently engineered a point

mutant in *S. mutans* strain UA159 (*fruK*M28stop, strain FruK-13) that prevents

the translation of an intact 1-phosphofructokinase (FruK) enzyme (3). Due

apparently to the accumulation of F-1-P in FruK-13, a severe growth phenotype

is evident when fructose is present. FruK-13 also displayed significant growth

defects when cultured in media containing glucose, sucrose, mannose,

galactose, lactose or certain amino sugars, demonstrated a broad influence of F-

1-P on carbohydrate metabolism and growth of *S. mutans* (3). RNA deep

sequencing (RNA-Seq) was used to define changes in the transcriptome arising

from loss of the FruK enzyme. The expression of 394 genes, including genes

encoding ribosomal proteins, proteins required for biosynthesis of nucleotides

and amino acids, and components of the PTS, were altered in FruK-13 compared

to the parental strain (3). Among the most up-regulated genes in the *fruK* mutant

were two open reading frames (ORF) in an apparent operonic arrangement,

SMU.507 and SMU.508; designated *sppR* and *sppA*, respectively. SppR is a

DeoR-type transcription regulator and SppA is a predicted hydrolase of the HAD

(haloacid dehalogenase) family IIB that includes enzymes such as trehalose-6-

phosphatase, plant and cyanobacterial sucrose-phosphatases, and a large

subfamily of bacterial Cof-like hydrolases; Cof being an *E. coli* phosphatase.

While both *sppR* and *sppA* are well conserved in *S. mutans* strains, homologous

systems comprised of both genes in tandem have only been identified in

Streptococcus gordonii and *Streptococcus sanguinis.* In this communication, we

investigated the regulation and function of the *sppRA* operon and explored the

contribution of its gene products to fitness and cariogenic potential of *S. mutans*,

particularly in the context of regulation of F-1-P levels.

Results

Overexpression of *sppA* enhances growth of FruK-13 in the presence of

fructose. In a previous study with a mutant lacking a 1-phosphofructokinase

(*fruK*), FruK-13, an operon encoding a DeoR-like regulator (SMU.507, *sppR*) and

a predicted hydrolase of the HAD superfamily (SMU.508, *sppA*) was significantly

up-regulated (3). As FruK-13 displays the most severe growth defect on fructose,

likely due to accumulation of F-1-P, it was posited that certain enzymes encoded

by *S. mutans* could eliminate these potentially toxic intermediates. The putative

sugar-phosphate phosphatase encoded by *sppA* was cloned into an expression

vector pIB184 behind a constitutive promoter and the resultant plasmid, pIB508,

was introduced into FruK-13. After confirming enhanced expression of *sppA* by

plasmid pIB508 (Fig. 1A), these resultant strains were compared in their growth

phenotypes in a fructose-based medium. Compared to the vector-only control

strain FruK-13/pIB184, which grew poorly on a TV-fructose medium, FruK-

13/pIB508 grew at a rate comparable to the wild-type strain UA159 (Fig. 1B),

providing a strong support for the hypothesis that SppA could function as a

sugar-phosphate phosphatase that decreased the levels of F-1-P in the *fruK*

genetic background.

SppA has hexose-phosphate phosphohydrolase activity. Multiple computer

algorithms predicted that SppA is a hydrolase of the HAD superfamily. To study

the function of SppA *in vitro*, the coding sequence of *sppA* was cloned into

pQE30 and expression was induced in *E. coli* to produce an N-terminally His-

tagged recombinant protein, designated His-SppA. His-SppA has 274 amino

acids, excluding the His-tag, and migrated as a single 37-kDa band in SDS-

polyacrylamide gel electrophoresis (SDS-PAGE). The protein was relatively

stable in a variety of buffers and pH values, although addition of 0.5 M NaCl and

at least 10% of glycerol was required to retain activity during longer-term storage

(>1 month) at 4°C. To test the kinetics of SppA in removing the phosphate group

from presumptive substrates, a set of phosphate-free buffers were tested to carry

out these experiments (see Materials and Methods for detail), and the phosphate

released in each reaction was monitored using a Malachite green reagent kit.

Substrate specificity. Using 2.5 mM p-nitrophenylphosphate (pNPP) as a

general colorimetric substrate for phosphatases, phosphatase activity by His-

SppA was detected at 37°C in 50 mM HEPES buffer at pH 7.5. SppA activity on

pNPP could be stimulated by the addition of millimolar concentrations of $MgCl_2$

(data not shown). We then conducted a number of preliminary experiments to

gain insights into optimal conditions for activity of His-SppA on a number of sugar

phosphates. Then, as determined by an end-point assay performed under

optimized conditions (Fig. 2A), the preferred substrate for His-SppA was found to

be F-1-P, followed by F-6-P. Very little activity was observed when F-1,6-bP, G-

6-P, or GlcN-6-P were used as substrates, and SppA had no activity on GlcNAc-

6-P under the conditions tested.

Effects of metal ions. Many phosphatases are affected by the presence of

metal ions (21), thus the activity of His-SppA was studied by adding to the

reactions millimolar levels of $CaCl_2$, $CuSO_4$, $MgCl_2$, $MnCl_2$, and $ZnCl_2$. $CaCl_2$,

$CuSO_4$, and $ZnCl_2$ at these levels had no detectable impact to the activity of His-

SppA on F-1-P as a substrate, but the presence of $MgCl_2$ and $MnCl_2$ significantly

enhanced the release of phosphate by the purified, recombinant enzyme (Fig. 2).

By testing both Mg^{2+} and Mn^{2+} together, it was determined that optimal enzymatic

activities could be elicited when $MgCl_2$ was present at 5 mM, along with 1 mM

$MnCl_2$.

pH. The optimal pH for the activities of His-SppA was determined to be about

6.5, and the rate of phosphate release significantly decreased when the pH

values of the reaction deviated from this point, particularly at values of 6.0 and

below (Fig. 2C). Since *S. mutans* maintains ΔpH values of 0.5 to 1.0 units above

the external environment (22), SppA is probably most active in cells growing in a

moderately acidic environment, such as what would occur when *S. mutans* is

actively metabolizing carbohydrate(s).

Fluoride. Sodium fluoride (NaF) is an inhibitor of many phosphatases. The

presence of 10 mM NaF significantly reduced (by >50%) the rate of phosphate

release from F-1-P by His-SppA (Fig. S1 in the supplemental material).

Reaction kinetics on F-1-P and F-6-P. Finally, using the optimal pH and

metal ion conditions described above, V_{max} and K_m of His-SppA were measured

using F-1-P or F-6-P. The purified enzyme was incubated with increasing

concentrations of each substrate (see Materials and Methods for details) at 37°C,

and release of phosphate over time was monitored by periodic sampling of the

reactions. The release rate of phosphate for each reaction was then plotted

against the substrate levels to estimate rate and substrate affinity (Fig. 2D). For

F-1-P, the V_{max} was 9.50 ± 0.38 μmol/(min x mg protein), and the K_m was 1.90 ±

0.21 mM. For F-6-P, the V_{max} was 0.52 ± 0.03 μmol/(min x mg protein) and the

K_m was 4.83 ± 0.51 mM. Thus, under these conditions, F-1-P is a preferred

substrate for SppA, whereas F-6-P can be dephosphorylated by SppA, albeit less

efficiently. These data are consistent with idea that *S. mutans* may use SppA to

limit accumulation of F-1-P, which can be further phosphorylated (by FruK) into

F-1,6-bP. However, the data may also reflect a preference of the organism to

internalize then further metabolize F-6-P. These parameters, including V_{max} and

K_m, are also in line with what has been reported for other members of the HAD

family phosphatases in bacteria (5, 23, 24).

SppR is a negative regulator of *sppA* that responds to fructose-phosphate

signals. SppR is predicted to be a member of the DeoR family, a group of

transcription regulators that often respond allosterically to metabolic

intermediates including phosphorylated carbohydrates. To begin understanding

the mode of regulation by SppR, a deletion mutant of *sppR* was constructed and

the expression of *sppA* was measured using quantitative RT-PCR (qRT-PCR).

The results showed a significant enhancement (8.95-fold ± 1.03-fold) in mRNA

levels of *sppA* due to the loss of SppR, when both the wild type and the mutant

were growing in the rich medium BHI.

Since higher expression of both *sppR* and *sppA* was noted using RNA-Seq of

the *fruK* mutant (FruK-13), which should have elevated levels of F-1-P, it was

hypothesized that SppR negatively regulates the *spp* operon, with SppR being

allosterically regulated by fructose phosphates, including F-1-P. To test this

hypothesis, the promoter activity of the operon (P*sppR*) was examined by

utilizing a promoter-reporter gene fusion P*sppR::cat* that was integrated in single

copy at a distant site on the chromosome. Expression of the *cat* fusion was

measured in cells growing on various carbohydrates. A UA159 derivative

carrying the P*sppR::cat* fusion was cultivated in TV medium supplemented with

10 mM glucose to mid-exponential phase, then pulsed for 1 h with 25 mM or 50

mM fructose to increase the levels of fructose-derived metabolites. In addition, 25

mM or 50 mM of glucose, xylitol, or sucrose were also used to treat these cells.

The CAT activities measured (Fig. 3A) on cells pulsed with fructose showed

significant, concentration-dependent increases compared to cells pulsed with

glucose. A slight reduction in CAT activity was noted in cells pulsed with 50 mM

xylitol, and sucrose had little to no impact on the expression of the *spp* promoter

fusion. When compared to glucose, treatment of cells with fructose should result

in accumulation of F-1-P, adding support to the idea that F-1-P is an allosteric

effector capable of relieving the repression by SppR. On the other hand, xylitol

treatment could potentially reduce the capacity of the PTS and carbon flux,

reducing the levels of glycolytic metabolites, including F-1-P and F-6-P. The lack

of response from sucrose treatment could be due to the preference by the

bacterium to utilize the glucose moiety, as our recent study has indicated that

free fructose is expelled by *S. mutans* after sucrose is transported and

phosphorylated on the glucose moiety by ScrA, and the resultant sucrose-6-

phosphate (S-6-P) is cleaved by ScrB (25).

We have previously shown that fructose internalization via FruI likely results in

creation of F-1-P, whereas internalization of fructose by EII^{Lev} (*levDEFG*) yields

F-6-P (3). To test if transport of substrates via different PTS permeases could

affect the regulation of expression of the *spp* operon, the P*sppR::cat* fusion was

placed in the background of mutants deficient in the fructose-PTS permeases,

levD or *fruI*, or in a mutant lacking the primary glucose-PTS permease (*manL*)

(26). A mutant defective in production of catabolite control protein A (CcpA), a

major regulator of genes involved in carbon flow in *S. mutans*, was also included

as CcpA can regulate the expression of certain PTS permease genes (27). The

wild-type control and the mutant strains were grown in TV-glucose to mid-

exponential phase, then pulsed with 50 mM fructose to induce the P*sppR::cat*

fusion. In support of our hypothesis that F-1-P is required for inducing expression

of *spp*, loss of *fruI* abolished fructose-dependent induction of CAT activity,

whereas the *levD* mutant behaved like the wild type (Fig. 3B). Interestingly, a

deletion of the glucose-PTS gene *manL* resulted in significantly increased

expression of the P*sppR::cat* fusion, consistent with our recent study that showed

interconnected regulation of fructose and glucose PTS permeases in *S. mutans*

(3). The *ccpA* mutant showed no difference in response to fructose.

To gauge the overall impact of the regulation by SppR on the *sppRA* operon

and the potential that depletion of F-1-P associated with *sppA* induction could

function in feedback regulation of the operon, we assayed the expression of

P*sppR::cat* fusion in strains deficient in *sppR* and *sppA*, respectively. The results

(Fig. 3B) showed nearly a 2-log increase in CAT activities in the *sppR* mutant

regardless of the sugars tested, and no change in the *sppA* mutant. Together,

these results suggest that SppR exerts a strong, negative regulation over the *spp*

operon, resulting in low levels of expression of *sppA* even when fructose is

present, at least in an otherwise wild-type genetic background.

SppR binds to the *spp* promoter and to F-6-P. To demonstrate the direct

interaction of SppR with its own promoter region, a recombinant His-SppR was

engineered and overexpressed in *E. coli*. The purified His-SppR was used in an

electrophoretic mobility shift assay (EMSA) that included a biotin-labeled DNA

fragment containing the 250-bp intergenic region immediately upstream of the

sppR coding sequence. As shown in Fig. 4A, the presence of 0.5 µM His-SppR

resulted in significant shifting of the DNA probe. However, when the reaction was

repeated with the addition of 5 mM sugar-phosphates, including F-1-P, F-6-P, F-

1,6-bP and G-6-P, there was little effect on the interactions of His-SppR with the

probe.

As an alternative to EMSA, we also performed a thermal shift assay in which

increasing temperature induces unfolding of His-SppR, which is monitored using

a protein dye that fluoresces when bound to unfolded polypeptides (28). The

melting temperature (Tm) of proteins usually increases in the presence of

compounds that are bound by the protein. We added to the melting reaction

various amounts of glucose, G-6-P, fructose, F-1-P, F-6-P, F-1,6-bP, GlcN, GlcN-

6-P, GlcNAc and GlcNAc-6-P. While most sugar and sugar phosphates had no

impact to the unfolding of His-SppR, addition of F-6-P increased the Tm in a

dose-responsive manner, indicating an interaction between F-6-P and His-SppR.

By plotting the Tm values of the reactions against the concentrations of F-6-P

used, it was determined that F-6-P associates with His-SppR with a *Kd* of 3.6 ±

0.15 mM (Fig. 4B). Considering the results of our genetic analysis, we posit that

the interactions between SppR and fructose-derived metabolites modulate the

repression of *spp* expression. It is not clear why F-1-P did not elicit an increase in

Tm in this system. In the future, we plan to utilize a spectrum of binding assays

and non-tagged SppR proteins to further evaluate the possible allosteric effectors

that modulate the DNA-binding affinity of SppR *in vivo.*

Overexpression of SppA enhances resistance to xylitol. Having

demonstrated that enhanced expression of SppA from pIB508 abolished the

sugar-phosphate stress in strain FruK-13, we tested if overexpression of SppA

could also alleviate stress caused by the presence of xylitol. Strains

UA159/pIB184 and UA159/pIB508 were created and their growth was monitored

in TV containing 0.5% glucose or fructose, with or without 1% xylitol. Without

addition of xylitol, both strains grew similarly well in TV glucose or fructose (data

not shown). When 1% xylitol was added to the TV-glucose medium,

UA159/pIB184 displayed a reduced growth rate compared to when the strain was

grown in TV with glucose alone. Interestingly, UA159/pIB508 showed

significantly faster growth than UA159/pIB184 in the presence of both glucose

and xylitol, indicating enhanced resistance to xylitol as a result of increased

expression of SppA (Fig. 5A; see Fig. 1A for *sppA* mRNA levels). In contrast,

addition of 1% xylitol to TV-fructose had no discernable effect on the growth

phenotypes of either strain (data not shown); a result consistent with the notion

that xylitol is internalized via a fructose-specific PTS (FruI) (18). In this case, our

interpretation is that fructose competes better than xylitol for uptake, so less

xylitol is internalized and there is less of an impact on growth.

Since deletion of *sppR* enhanced the expression of *sppA* (Fig. 3B), we

compared resistance to xylitol by the wild type and its otherwise-isogenic *sppR*

and *sppA* mutants. The *sppR* mutant showed significantly improved growth in

TV-glucose supplemented with 1% of xylitol, similar to UA159/pIB508 that

overexpressed *sppA* (Fig. 5B). However, loss of *sppA* had no effect on xylitol

resistance, as the *sppA* mutant grew similarly to the wild type. These results

agree with our earlier findings that the expression of *sppA* is likely maintained at

minimal levels, perhaps to avoid futile cycling. We have attempted, but have

failed, to demonstrate the capacity of SppA to dephosphorylate xylitol-5-

phosphate (X-5-P)(see Text S1 in the supplemental material), in large part due to

the difficulty producing sufficient amounts of the compound; it is not commercially

available.

Accumulation of F-1-P enhances eDNA release and affects biofilm

formation. *S. mutans* depends on the development of biofilms for its virulence,

but little information is available on the effects of fructose on *S. mutans* biofilms.

Wild-type cultures in exponential phase were grown in BHI and used to inoculate

a biofilm medium (BM) constituted with 2 mM sucrose and 18 mM glucose

(BMGS) or 18 mM fructose (BMFS), as well as a number of combinations of both

glucose and fructose totaling 18 mM hexose. After 24 h or 48 h of incubation, the

supernates of the culture were removed and development of biofilms was

visualized and quantified by staining with 0.1% crystal violet. As shown in Fig.

6A, growth on fructose (BMFS) changed the accumulation of biomass by *S.*

mutans on a polystyrene surface, resulting in a thinner but more uniform biofilm,

as opposed to a patchy coverage by the biofilms formed on BMGS. Elution of

crystal violet also indicated that greater amounts of dye were retained by the

biofilms formed on fructose than glucose (Fig. 6A). This effect was not likely a

result of reduced growth caused by a lack of glucose, as similar phenotypes were

visible by the presence of as little as 3 mM fructose (data not shown). Also, to

differentiate between effects of internalization of F-1-P versus F-6-P, we included

the otherwise-isogenic mutants of UA159 deficient in either *fruI* (F-1-P) or *levD*

(F-6-P). The results (Fig. 6A) showed that while the wild type and both mutants

formed similar biofilms in BMGS, the *fruI* mutant formed patchy biofilms on

BMFS, resembling those on BMGS. The *levD* mutant behaved like the wild type.

Recent research has suggested that the ability of *S. mutans* to form efficient

biofilms depends on both the synthesis of glucans and the release of

extracellular DNA (eDNA) by bacteria, via both lysis-dependent and -independent

pathways (29). To further study the effects of fructose, the release of eDNA was

quantified using a fluorescent DNA dye in the biofilms of *S. mutans*. The results

were normalized against the viable counts of bacteria (CFU). When grown as

biofilms in BMFS, UA159 released significantly more eDNA per CFU after 24 h

(5-fold) or 48 h (17-fold) of growth compared to BMGS medium (similar to

UA159/pIB184 in Fig. 6C). Significantly lower CFU (5-fold for 24 h, and 20-fold

for 48 h) were recovered from biofilms formed on BMFS than BMGS. As such,

the amounts of eDNA detected per unit volume of supernatant fluid were

comparable for both BMGS and BMFS. Therefore, it appears that the bacteria

incubated in the presence of fructose experienced significantly more lysis, which

resulted in increased release of eDNA and changes in formation of biofilms;

although at this point we cannot rule out the involvement of other routes for

production of eDNA. When treated with a LIVE/DEAD stain and observed under

a confocal laser-scanning microscope, the biofilms formed on BMFS indeed

showed notably more dead cells than those formed on BMGS (Fig. 6B). Similar

impacts of fructose were observed for biofilm development by other genomically

distinct, wild-type isolates of *S. mutans* (30), so the phenomenon is not restricted

to the lab strain UA159 (Fig. S2 in the supplemental material). Furthermore, we

compared the sensitivity of planktonic *S. mutans* cells to the cell wall-degrading

enzymes lysozyme and mutanolysin by extracting chromosomal DNA after

enzymatic treatment, using batch cultures grown in glucose- or fructose-

containing medium. The results showed that 10-fold more DNA could be

harvested from UA159 that was cultured in TV-fructose, compared to cells

cultivated in TV-glucose, after each was treated with lysozyme, or a combination

of lysozyme and mutanolysin, under otherwise-identical conditions (Fig. S3 in the

supplemental material).

As a further test of our hypothesis that F-1-P in particular is responsible for the

fructose effects observed so far, the strains UA159/pIB508 and UA159/pIB184

were compared in their ability to produce eDNA in the 48-h biofilm model. Similar

to UA159, strain UA159/pIB184 yielded significantly lower numbers of CFU, but

much higher levels of eDNA/CFU when growing on BMFS compared BMGS (Fig.

6C). When *sppA* was overexpressed on pIB508, significant reductions in the

release of eDNA/CFU were apparent in comparison to strain UA159/pIB184, an

impact that was especially significant for cells cultured in BMFS. At the same

time, bacterial viable counts also showed significant recovery as an effect of

pIB508, particularly when assayed using BMFS (Fig. 6D). These results suggest

that accumulation of F-1-P in *S. mutans* likely leads to alterations in the cell-

envelope that results in enhanced autolysis and release of eDNA, causing

aberrant biofilm phenotypes.

Other putative phosphatases capable of promoting sugar-phosphate

resistance. The existence in *S. mutans* of a xylitol-inducible phosphatase

capable of dephosphorylating X-5-P has been proposed (31). As xylitol reduced

spp expression (Fig. 3A), we hypothesized that other phosphatase(s) are

produced that may act on xylitol. Based on sequence similarity to SppA and other

bacterial sugar-phosphate phosphatases, a functional genomics approach was

employed to identify genes encoding products that could promote resistance to

xylitol. A total of 9 ORFs were selected and mutated by allelic exchange with an

antibiotic resistance marker, and three (SMU.428, SMU.488 and SMU.718c)

were chosen as potential candidates based on the decreased growth rates (Fig.

S4 in the supplemental material) and yields (Fig. 7A) of the mutants in the

presence of xylitol, compared to the wild type. For confirmation, these genes

were expressed constitutively using pIB184, in both the wild-type and the

respective mutant genetic backgrounds. When assayed for growth on TV-

glucose supplemented with xylitol, mutants that were complemented by each of

these three genes regained resistance to xylitol, and UA159 containing

SMU.718c or SMU.488 expressed from pIB184 also showed enhanced yields,

compared to their respective vector-only controls (Fig. S5 in the supplemental

material). Furthermore, we previously reported that a deletion of the tagatose-

1,6-bisphosphate aldolase (LacD) results in sensitivity to galactose in *S. mutans,*

likely due to accumulation of metabolic intermediates that are normally

catabolized by the tagatose pathway (32). When pIB428 was introduced into the

lacD mutant, there was a significant alleviation of the inhibitory effects of

galactose (Fig. 7B). We found that SMU.428, SMU.488 and SMU.718c are well

conserved in the genomes of a diverse group of *S. mutans* isolates (30)*.* Amino

acid sequence alignments performed using these three proteins, together with

SppA and the AphA phosphatase from *E. coli* (33), showed significant

conservation in those amino acid residues (Fig. S6 in the supplemental material)

that place these proteins in the family of bacterial class B acid phosphatases,

part of the DDDD superfamily of phosphohydrolases (34). Computer analysis

failed to identify any potential signal peptide sequences in the N-terminus of

these proteins, suggesting they are not secreted. Further research is needed to

understand their contribution to xylitol resistance and more broadly, to sugar-

phosphate stress.

Discussion

Carbohydrate fermentation is a highly dynamic process where the relative

activities of various enzymes and resultant intermediates must be maintained at

optimal levels to ensure fitness and survival, and to balance anabolic and

catabolic processes. Perturbations to the systems, such as those that occur in

the oral microbiome during intake of dietary carbohydrates by the host, require

rapid responses by the microbiota. Sugar-related stress responses mediated by

complex regulatory systems have been characterized mostly in Gram-negative

model bacteria, e.g., *E. coli* (8, 35, 36). As a major caries pathogen, *S. mutans*

depends on carbohydrate fermentation for energy generation and virulence. *S.*

mutans is particularly well adapted to metabolizing sucrose, in part through the

activities of three secreted glucosyltransferases (GtfB, GtfC and GtfD) that

convert sucrose into glucan polysaccharides that serve as the scaffolding for

biofilm accumulation and in creation of a diffusion-limiting matrix (37).

Interestingly, Gtfs generate a molecule of free fructose for every glucose moiety

of sucrose that is targeted toward glucan formation. Furthermore, it is now clear

that the majority of the sucrose presented to *S. mutans* is internalized by a

sucrose-specific PTS permease (38). After internalization and phosphorylation,

sucrose is cleaved by a sucrose-6-phosphate hydrolase into G-6-P and fructose.

However, a significant portion of the fructose thus generated is expelled into the

environment and re-internalized by fructose PTS permeases (25, 39). These

observations, when coupled with the fact that fructose itself is a major constituent

of foodstuffs, e.g. with high-fructose corn syrups increasingly used as common

sweeteners, necessitate a better understanding of the metabolism of fructose by

the oral microbiome.

S. mutans maintains at least three PTS permease operons (*fruRKI*, *fruPCD*

and *levDEFG*) plus three phosphofructokinases (FruK, FruP, and

SMU.1191/Pfk). Further, while a number of *S. mutans* PTS permeases can

transport sugars other than their primary cognate carbohydrate, there are no

other instances in *S. mutans* of redundancy in PTS permeases for sugars,

besides those for fructose. This redundancy in PTS permeases, in itself, is

noteworthy and could reflect significant evolutionary benefits to redundant

fructose catabolic systems. Much remains to be clarified regarding the

contributions of these PTS operons to the establishment and persistence of *S.*

mutans in the human oral cavity, and to the organism's cariogenic potential.

However, it is understood that both F-1-P and F-6-P generated by the PTS are

further phosphorylated to create F-1,6-bP, likely via 1-phosphofructokinase FruK

and 6-phosphofructokinase SMU.1191, respectively. F-1,6-bP is then cleaved by

an F-1,6-bP-specfic aldolase into glyceraldehyde 3-phosphate and

dihydroxyacetone phosphate (3). We have demonstrated that removal of FruK

leads to significant growth and fitness defects in *S. mutans*, apparently due to

accumulation of the metabolic intermediate F-1-P (3). Here we present evidence

that *S. mutans* maintains tight control over the levels of F-1-P through an

inducible sugar-phosphate phosphatase (*spp*) operon, consisting of a

transcriptional regulator SppR and a fructose-phosphate hydrolase SppA.

Changes in F-1-P levels, either increase or decrease, significantly alter bacterial

physiology, including cell envelope integrity and biofilm development. We

hypothesize that, aside from being an apparent stress compound in *S. mutans*

(6), that F-1-P may also act as a signaling molecule, the accumulation of which

promotes autolysis and release of extracellular DNA; with the latter being a

critical contributor to formation of biofilms (29).

In support of this hypothesis, our latest RNA deep sequencing analysis in *S.*

mutans has identified a fructose regulon that includes a significant number of

genes that are likely involved in stress responses (25). For example, growth on

fructose up-regulated the *cid* operon (SMU.1700c~1703c) that influences

oxidative stress resistance and modulation of autolysis and biofilm development

(40). Also up-regulated were the *rcrRPQ* and *relPRS* operons (SMU.921 to

SMU.928) which have major impacts on acid and oxidative stress tolerance,

competence development and (p)ppGpp metabolism (41). Thus, fructose and its

intermediates are directly linked to pathways that are critical for persistence,

growth and survival decisions, nutrient acquisition and genetic diversification.

On the other hand, a growing body of evidence suggests that F-1-P is acting

as a regulatory molecule for *S. mutans*. Our recent study on the function of the

fruRKI operon has detailed the transcriptomic impact of accumulation of F-1-P, a

compound that likely serves as an allosteric effector to another DeoR-type

regulator, FruR (3). It was clear that F-1-P influences gene regulation beyond the

fruRKI operon. Specifically, the *fruK* mutant showed altered expression in as

many as 20 known or probable transcriptional regulators, impacting ~20% of the

genome. Key functions ranging from energy metabolism, protein and nucleic acid

biosynthesis, to stress tolerance were affected by fructose. In addition to showing

that overproduction of SppA, which preferentially degrades F-1-P, leads to

phenotypic changes in both xylitol resistance and eDNA release, we have shown

that the ability of fructose to induce *spp* operon expression depends on an intact

FruI protein, which encodes the fructose-PTS permease that creates F-1-P (3).

The lack of interactions between F-1-P and the recombinant SppR protein *in vitro*

could suggest the requirement for other factors in addition to F-1-P that influence

SppR activity or the contribution of other regulatory proteins. Conversely, addition

of the His-tag to SppR altered the F-1-P binding behavior of the recombinant

protein, compared to native SppR. The interaction of SppR with F-6-P that we

observed *in vitro* also points to further complexity in this system. For example, F-

6-P could act as a second allosteric effector for SppR, perhaps when present at

high concentrations. Future research will focus on identifying other regulatory

circuits that respond to the accumulation of F-1-P, both directly and indirectly, in

particular the pathway(s) responsible for cell envelope integrity and biofilm

phenotypes. As stated earlier, the *spp* operon has only been identified in a few

oral streptococcal species, and its specificity for F-1-P could have implications in

understanding niche adaptations and oral microbial ecology.

An intracellular hexose-6-phosphate: phosphohydrolase was purified from

Lactococcus lactis that has a preference for galactose-6-phosphate and F-6-P

(23). This enzyme had an estimated M_r of 60 KDa by SDS-PAGE, was most

active at pH 6.0, and its activity was stimulated by Mn^{2+}, Mg^{2+} and Fe^{2+}, and

inhibited by fluoride. Another bacterial sugar phosphate phosphatase, Spp, was

recently identified in *Pseudomonas fluorescens* that was capable of

dephosphorylating ribose-5-P, F-6-P, and G-6-P (5). The optimal Mg^{2+}

concentration for Spp was determined to be 2.5 mM, whereas addition of Mn^{2+}

reduced the reaction rate. Therefore, SppA appears to share some similarities

with other bacterial phosphohydrolases, yet it stands out with its preference for F-

1-P. Lastly, despite our repeated efforts, we were not able to determine whether

X-5-P is a substrate of SppA, partly due to our inability to obtain or generate

sufficient X-5-P (it is not commercially available). As we continue to resolve this

technical issue, it is important to note the possibility that SppA alleviates xylitol-

related stress through its activities on either F-1-P or F-6-P, as X-5-P has been

suggested to inhibit the activities of phosphofructokinases (9). The same

rationale could apply in interpreting the xylitol phenotypes associated with the

mutants that lacked one of three additional HAD phosphatases; that each

enzyme may have multiple substrates and a stress phenotype could be alleviated

by actions on molecules other than the one triggering the stress. Most notably,

overexpression of SMU.428 abolished galactose-mediated stress in a *lacD*

mutant. SMU.428 is located immediately downstream of the *copYAZ* operon that

is responsible for copper efflux, biofilm formation and stress tolerance, yet its

homologues in a number of other streptococci are closely associated with the

tagatose operon. Clearly though, *S. mutans* possesses multiple

phosphohydrolases and related regulatory proteins capable of mounting a stress

response when triggered by certain sugar-phosphates. Additional research is

needed to understand the function and regulation of these putative sugar-

phosphate phosphatases.

Materials and Methods

Bacterial strains and culture conditions. *Streptococcus mutans* strains (Table

1) were maintained using brain heart infusion (BHI)(Difco Laboratories, Detroit,

MI) agar plates or in liquid BHI medium at 37°C, in an aerobic atmosphere

supplemented with 5% CO_2. To monitor bacterial growth phenotypes and for

most other experiments, a semi-synthetic Tryptone-vitamin (TV) medium (42)

was utilized with various carbohydrates being provided at specified

concentrations. Overnight bacterial cultures were prepared in BHI, then sub-

cultured to exponential phase, and inoculated at 1:100 dilutions into TV media.

Growth was monitored as the change in optical density at 600 nm (OD_{600}) using

a Bioscreen C lab system (Labsystems Oy, Helsinki, Finland). For development

of biofilms in 96-well polystyrene plates or ibidi glass chambers (ibidi GmbH,

Martinsried, Germany), a biofilm medium (BM)(43) was utilized that contained 2

mM sucrose and varying amounts of glucose or fructose, with bacterial inoculum

prepared in BHI. Each of these experiments was repeated at least three times

using independent cultures (biological replicates), with each sample being

represented by at least two wells (technical replicates).

DNA methodology. Standard recombinant DNA techniques were employed in

preparation of genetic materials for applications such as cloning and

mutagenesis (44). To engineer a mutant defective in *sppR*, a set of 4 primers

(Table 2) were designed for PCR amplification of two DNA fragments flanking the

coding sequence of *sppR* (upstream fragment: SMU.507-1 and SMU.507-2GA;

downstream fragment: SMU.507-3GA and SMU.507-4). PCR reactions were

carried out using these primers to generate two DNA fragments that shared >25

bp overlaps with the 5'- and 3'- ends, respectively, of a kanamycin-resistance

cassette. Subsequently, these flanking DNA fragments were used in equimolar

concentrations, together with the kanamycin cassette, in a one-step ligation

reaction that was catalyzed by a Gibson assembly kit (New England Biolabs,

Beverly, MA). The ligated product was then used to transform a naturally

competent UA159 culture, followed by selection on BHI agar plates containing

kanamycin (1 mg/ml). Similar procedures were carried out in the creation of

mutants defective in *sppA*, SMU.428, SMU.488, SMU.718c and other genes. All

strains created in this study have been validated by DNA sequencing to ensure

that no unintended mutations were introduced into the flanking regions of the

gene of interest.

To construct a complementing plasmid for the *sppA* mutant, a pair of primers

(SMU.508-5'Bm and SMU.508-3'SacI) were designed and used for amplification

of the coding sequence of *sppA,* together with its putative ribosome-binding site.

This DNA product was treated with restriction enzymes *Bam*HI and *Sac*I, and

then ligated with an expression vector pIB184 (45) that was linearized with the

same enzymes. After cloning into an *Escherichia coli* host strain BH10B, the

resultant plasmid pIB508 was verified to contain the desired coding and

regulatory sequences and introduced into the *sppA* mutant background, as well

as the wild type UA159. Three additional plasmids, pIB428, pIB488, and

pIB718c, were constructed in the same way, in the pIB184 vector, and used to

complement their respective mutants. To monitor the transcriptional activities of

the *spp* operon, a P*sppR::cat* fusion was constructed by cloning a DNA fragment

containing the intergenic region upstream of *sppR* onto the integration vector

pJL84 that contained a promoterless *cat* gene (46), and the gene fusion was

subsequently delivered into a distal site on the chromosome of *S. mutans* by

double cross-over recombination. The sequence information of all these primers

is included in Table 2.

To assess the resistance of different UA159 cultures to muralytic enzymes, the

bacteria were cultivated overnight in 5 ml TV medium containing 0.5% glucose or

fructose, harvested by centrifugation at 4°C and washed once with a TE buffer

(10 mM Tris-Cl, 1 mM EDTA, pH 8.0). Cells were resuspended in 500 µl of lysis

buffer that contained 50 mM Tris-Cl at pH 8.0, 10 mM EDTA, 10 mg/ml of

lysozyme, and 0.1 mg/ml of RNase A. After addition of 0 or 20 units of

mutanolysin (from *Streptomyces globisporus,* Millipore Sigma, St. Louis, MO),

cells were incubated at 37°C for 1 h, and lysed by adding 40 µl of 20% SDS and

incubating at 80°C for 15 min. Subsequently, 80 µl of 5 M NaCl was added into

each reaction, mixed well, followed by addition of 600 µl of phenol: chloroform:

isoamyl alcohol (25: 24: 1, v/v/v) and 2 min of vortexing. The solution was then

clarified by centrifugation at room temperature at $15000 \times g$ for 5 min, followed by

the removal of 500 µl of the aqueous phase. The genomic DNA was then

precipitated from the aqueous solution by addition of 500 µl 100% isopropanol

and centrifugation as before. After washing once with 75% ethanol, the DNA was

air-dried and dissolved in 50 µl TE buffer and quantified using a Nanodrop

apparatus (Thermo Fisher, Waltham, MA) and agarose-gel electrophoresis.

Engineering and preparation of recombinant proteins of SppR and SppA.

DNA fragments containing the entire coding sequences of SppR and SppA were

each amplified using PCR primers (SMU.507-5'Bm and SMU.507-3'HIII for *sppR*,

and SMU.508-5'Bm and SMU.508-3'HIII for *sppA*)(Table 2) containing

engineered restriction sites, and digested using restriction enzymes *Bam*HI and

*Hin*dIII. After ligating with a pQE30 vector prepared by treating with the same

enzymes to create an N-terminal, 6×His-tagged gene product, the constructs

were introduced into *E. coli* M15. After confirming the fidelity of the PCR and

cloning via DNA sequencing, the expression of each recombinant protein was

induced by adding IPTG to a final concentration of 0.05 mM to cells growing at

mid-exponential phase.

Purification of recombinant proteins, His-SppR and His-SppA, was carried out

using a Ni-NTA agarose according to protocols provided by the supplier

(Qiagen), with some modifications. Briefly, to improve the stability of His-SppR,

10% of glycerol was supplemented to each of the original lysis buffer (50 mM

NaH_2PO_4, 0.3 M NaCl, and 10 mM imidazole, pH 8.0), the wash buffer and

elution buffer. After elution from the column, His-SppR was dialyzed against a

buffer containing 10 mM HEPES pH 7.5, 0.5 M NaCl and 20% glycerol, using a

0.5-mL Slide-A-Lyzer dialysis device with a MWCO of 3.5 KDa (Thermo Fisher).

To avoid carryover of phosphate into the subsequent reactions involving the

recombinant SppA, purification of His-SppA was performed using a phosphate-

free buffer system that was created by replacing the NaH_2PO_4 with 50 mM

HEPES (pH 7.9), and also by supplementing with 10% glycerol. Dialysis of the

purified His-SppA solution was carried out against 10 mM PIPES pH 6.5, 0.5 M

NaCl, and 20% glycerol.

Electrophoretic mobility shift assay (EMSA) and thermal shift. To study the

interaction of His-SppR with the promoter of *sppR*, EMSA was performed using a

biotin-labeled DNA fragment containing the intergenic region upstream of the

sppR coding sequence as previously described (47). When needed, the sugar

phosphates were each used at 5 mM. To study the interaction of SppR with

various sugar phosphates, a 10-µM His-SppR solution was prepared using a

buffer containing 0.1 M HEPES (pH 7.5) and 0.15 M NaCl. After addition of a

SYPRO Orange protein stain (Thermo Fisher), used at a 6000-fold dilution, and 5

mM of each sugar phosphate, the 10-µL reactions were subjected to a melting-

curve treatment on a CFX96 Touch real-time PCR thermocycler (Bio-Rad,

Hercules, CA) by increasing the temperature from 25°C to 75°C in increments of

0.2°C. The RFU measurements were then plotted against the temperature values

to determine the melting temperature (Tm) (28).

Phosphatase assay. The ability of His-SppA to remove the phosphate group

from various sugar-phosphates was measured at 37°C. The basic buffer was

composed of 50 mM PIPES pH 6.5, 5 mM $MgCl_2$, and 1 mM $MnCl_2$. For other pH

values, different buffering reagents were used in place of PIPES at the same

concentration: 2-ethanesulfonic acid (MES) or potassium citrate (K-Cit) for pH

5.5; 1,4-piperazinediethanesulfonic acid (PIPES) for pH 6.0 and pH 6.5; and

HEPES for pH 7.0 and pH 7.5. All initial experiments to determine the effects of

metal ions were carried out in an end-point assay at neutral pH.

   To measure the reaction kinetics of His-SppA on F-1-P, 20 µL of His-SppA that

was diluted to 1 µM using the dialysis buffer was mixed with 40 µL of F-1-P

prepared in the basic buffer, at final concentrations ranging from 0.1 mM to 1.0

mM; for F-1-P ranging from 1.5 mM to 10 mM, 0.25 µM of protein was used. All

components were pre-warmed to 37°C before mixing. With the reaction

maintained at 37°C, a 5-µL aliquot of the reaction was removed at 1 min intervals

and subjected to measurement of free phosphate using a phosphate assay kit

(Malachite Green Reagent, MGR; Sigma). For the kinetics of His-SppA with F-6-

P as a substrate, 20 µL of His-SppA at 2 µM was used in each reaction, and

aliquots of the reaction were taken and measured every 5 min. Each reaction

was repeated twice. The data were analyzed using the Prism from GraphPad

software (La Jolla, CA).

Biofilm development, microscopy and measurement of extracellular DNA

(eDNA). Development of biofilms by *S. mutans* was investigated in 96-well

polystyrene plates (Thermo Fisher) for comparison of overall biofilm mass, or on

glass surfaces using an ibidi 8-well μ-Slide to assess structural changes and

release of eDNA. A semi-defined biofilm medium (BM) was modified to contain

final concentrations of 2 mM sucrose and 18 mM glucose (BMGS) or fructose

(BMFS)(3, 48). Exponential-phase bacterial cultures prepared using BHI were

diluted 1:100 into BMGS or BMFS and incubated at 37°C in an aerobic

environment supplemented with 5% CO_2. Each experiment included at least

three biological replicates. The 96-well plates were incubated for 24 to 48 h,

washed with phosphate-buffered saline (PBS), and stained with 0.1% crystal

violet (CV). After extensive washing with water, the retained CV was eluted with

a solution of 30% acetic acid and quantified by measuring the optical density at

575 nm (OD_{575}).

The ibidi slides were incubated for 48 h, with a replacement of medium after

first 24 h. For observation of biofilm structures, each biofilm was treated using a

LIVE/DEAD BacLight bacterial vitality stain kit (Thermo Fisher) and visualized

using a confocal laser scanning microscopy system (4). To assess the amount of

eDNA released by each biofilm, the entirety of the 48-h biofilm culture was

harvested by scraping off the biofilm material, without washing, and sonicated for

30 sec at 100% power using a water-bath sonicator (model FB120, Thermo

Fisher). A portion of the cell suspension was subjected to serial dilutions and

plated on BHI plates for enumeration of CFU, and the remaining material was

pelleted using a tabletop centrifuge for 2 min at $16,000 \times g$. Subsequently, a

portion of the supernatant fluid was removed and combined 1:1 with a 5 μM

solution of SYTOX Green nucleic acid stain (Thermo Fisher), followed by

measurement of fluorescence (excitation/emission: 504/523 nm) using a

fluorescence-capable Synergy 2 plate reader from BioTek (Winooski, VT). A

standard curve was established using a known DNA solution and the results

were calculated as ng of eDNA released per CFU of the biofilm.

Quantitative reverse transcription PCR (qRT-PCR). Total RNA was extracted

from *S. mutans* cells harvested at $OD_{600} \approx 0.5$ using an RNeasy kit (Qiagen) by

following an established protocol (49). Primers specific to *sppA* were used for

both the reverse transcription and Real-time PCR that followed, with *gyrA* serving

as an internal control (25).

Acknowledgment

This work was supported by DE12236 from the National Institute of Dental and

Craniofacial Research.

References

1.   Postma PW, Lengeler JW, Jacobson GR. 1993.

Phosphoenolpyruvate:carbohydrate phosphotransferase systems of

bacteria. Microbiol Rev 57:543-94.

2.   Irani MH, Maitra PK. 1977. Properties of *Escherichia coli* mutants deficient

in enzymes of glycolysis. J Bacteriol 132:398-410.

3.   Zeng L, Chakraborty B, Farivar T, Burne RA. 2017. Coordinated regulation

of the EII^{Man} and *fruRKI* operons of *Streptococcus mutans* by global and

fructose-specific pathways. Appl Environ Microbiol 83.

4.   Zeng L, Farivar T, Burne RA. 2016. Amino sugars enhance

competitiveness of beneficial commensals with *Streptococcus mutans*

through multiple mechanisms. Appl Environ Microbiol 82:3671-82.

5.   Maleki S, Hrudikova R, Zotchev SB, Ertesvåg H. 2017. Identification of a

new phosphatase enzyme potentially involved in the sugar phosphate

stress response in *Pseudomonas fluorescens*. Applied and Environmental

Microbiology 83.

6.   Richards GR, Patel MV, Lloyd CR, Vanderpool CK. 2013. Depletion of

glycolytic intermediates plays a key role in glucose-phosphate stress in

Escherichia coli. J Bacteriol 195:4816-25.

7.   Korithoski B, Levesque CM, Cvitkovitch DG. 2007. Involvement of the

detoxifying enzyme lactoylglutathione lyase in *Streptococcus mutans*

aciduricity. J Bacteriol 189:7586-92.

8.   Kadner RJ, Murphy GP, Stephens CM. 1992. Two mechanisms for growth

inhibition by elevated transport of sugar phosphates in *Escherichia coli*. J

Gen Microbiol 138:2007-14.

9.   Assev S, Rolla G. 1986. Further studies on the growth inhibition of

Streptococcus mutans OMZ 176 by xylitol. Acta Pathol Microbiol Immunol

Scand B 94:97-102.

10.   Vanderpool CK. 2007. Physiological consequences of small RNA-

751   mediated regulation of glucose-phosphate stress. Curr Opin Microbiol

752   10:146-51.

753   11.   Tanzer JM, Thompson A, Wen ZT, Burne RA. 2006. *Streptococcus*

754   *mutans*: fructose transport, xylitol resistance, and virulence. J Dent Res

755   85:369-73.

756   12.   Trahan L. 1995. Xylitol: a review of its action on mutans streptococci and

757   dental plaque--its clinical significance. Int Dent J 45:77-92.

758   13.   Trahan L, Neron S, Bareil M. 1991. Intracellular xylitol-phosphate

759   hydrolysis and efflux of xylitol in *Streptococcus sobrinus.* Oral Microbiol

760   Immunol 6:41-50.

761   14.   Miyasawa-Hori H, Aizawa S, Takahashi N. 2006. Difference in the xylitol

762   sensitivity of acid production among *Streptococcus mutans* strains and the

763   biochemical mechanism. Oral Microbiol Immunol 21:201-5.

764   15.   Soderling E, Alaraisanen L, Scheinin A, Makinen KK. 1987. Effect of xylitol

765   and sorbitol on polysaccharide production by and adhesive properties of

766   *Streptococcus mutans.* Caries Res 21:109-16.

767   16.   Soderling EM, Hietala-Lenkkeri AM. 2010. Xylitol and erythritol decrease

768   adherence of polysaccharide-producing oral streptococci. Curr Microbiol

769   60:25-9.

770   17.   Trahan L, Mouton C. 1987. Selection for *Streptococcus mutans* with an

771   altered xylitol transport capacity in chronic xylitol consumers. J Dent Res

772   66:982-8.

773   18.   Wen ZT, Browngardt C, Burne RA. 2001. Characterization of two operons

774   that encode components of fructose-specific enzyme II of the

775   sugar:phosphotransferase system of *Streptococcus mutans.* FEMS

776   Microbiol Lett 205:337-42.

777   19.   Dills SS, Seno S. 1983. Regulation of hexitol catabolism in *Streptococcus*

778   *mutans.* J Bacteriol 153:861-6.

779   20.   Cvitkovitch DG, Boyd DA, Hamilton IR. 1995. Regulation of sugar

780   transport via the multiple sugar metabolism operon of *Streptococcus*

781   *mutans* by the phosphoenolpyruvate phosphotransferase system. J

782   Bacteriol 177:5704-6.

783   21.   Coleman JE. 1992. Structure and mechanism of alkaline phosphatase.

784   Annu Rev Biophys Biomol Struct 21:441-83.

785   22.   Dashper SG, Reynolds EC. 1992. pH regulation by *Streptococcus mutans.*

786   J Dent Res 71:1159-65.

787   23.   Thompson J, Chassy BM. 1983. Intracellular hexose-6-

788   phosphate:phosphohydrolase from *Streptococcus lactis*: purification,

789   properties, and function. J Bacteriol 156:70-80.

790   24.   Kuznetsova E, Proudfoot M, Gonzalez CF, Brown G, Omelchenko MV,

791   Borozan I, Carmel L, Wolf YI, Mori H, Savchenko AV, Arrowsmith CH,

792   Koonin EV, Edwards AM, Yakunin AF. 2006. Genome-wide analysis of

793   substrate specificities of the *Escherichia coli* haloacid dehalogenase-like

794   phosphatase family. J Biol Chem 281:36149-61.

795   25.   Zeng L, Burne RA. 2015. Sucrose- and fructose-specific effects on the

796   transcriptome of *Streptococcus mutans*, as determined by RNA

797   sequencing. Appl Environ Microbiol 82:146-56.

798   26.   Abranches J, Chen YY, Burne RA. 2003. Characterization of

799   *Streptococcus mutans* strains deficient in EIIAB^{Man} of the sugar

800   phosphotransferase system. Appl Environ Microbiol 69:4760-9.

801   27.   Abranches J, Nascimento MM, Zeng L, Browngardt CM, Wen ZT, Rivera

802   MF, Burne RA. 2008. CcpA regulates central metabolism and virulence

803   gene expression in *Streptococcus mutans*. J Bacteriol 190:2340-9.

804   28.   Huynh K, Partch CL. 2015. Analysis of protein stability and ligand

805   interactions by thermal shift assay. Curr Protoc Protein Sci 79:28.9.1-14.

806   29.   Liao S, Klein MI, Heim KP, Fan Y, Bitoun JP, Ahn SJ, Burne RA, Koo H,

807   Brady LJ, Wen ZT. 2014. *Streptococcus mutans* extracellular DNA is

808   upregulated during growth in biofilms, actively released via membrane

809   vesicles, and influenced by components of the protein secretion

810   machinery. J Bacteriol 196:2355-66.

811   30.   Cornejo OE, Lefebure T, Pavinski Bitar PD, Lang P, Richards VP,

812   Eilertson K, Do T, Beighton D, Zeng L, Ahn SJ, Burne RA, Siepel A,

813   Bustamante CD, Stanhope MJ. 2013. Evolutionary and population

814   genomics of the cavity causing bacteria *Streptococcus mutans*. Mol Biol

815   Evol 30:881-93.

31.   Pihlanto-Leppala A, Soderling E, Makinen KK. 1990. Expulsion

mechanism of xylitol 5-phosphate in *Streptococcus mutans*. Scand J Dent

Res 98:112-9.

32.   Zeng L, Das S, Burne RA. 2010. Utilization of lactose and galactose by

Streptococcus mutans: transport, toxicity, and carbon catabolite

repression. J Bacteriol 192:2434-44.

33.   Calderone V, Forleo C, Benvenuti M, Thaller MC, Rossolini GM, Mangani

S. 2006. A structure-based proposal for the catalytic mechanism of the

bacterial acid phosphatase AphA belonging to the DDDD superfamily of

phosphohydrolases. Journal of Molecular Biology 355:708-721.

34.   Thaller MC, Schippa S, Rossolini GM. 1998. Conserved sequence motifs

among bacterial, eukaryotic, and archaeal phosphatases that define a new

phosphohydrolase superfamily. Protein Sci 7:1647-52.

35.   Horler RS, Vanderpool CK. 2009. Homologs of the small RNA SgrS are

broadly distributed in enteric bacteria but have diverged in size and

sequence. Nucleic Acids Res 37:5465-76.

36.   Kessler JR, Cobe BL, Richards GR. 2017. Stringent response regulators

contribute to recovery from glucose phosphate stress in *Escherichia coli*.

Applied and Environmental Microbiology 83.

37.   Klein MI, Hwang G, Santos PH, Campanella OH, Koo H. 2015.

Streptococcus mutans-derived extracellular matrix in cariogenic oral

biofilms. Front Cell Infect Microbiol 5:10.

38.   Tanzer JM, Chassy BM, Krichevsky MI. 1971. Sucrose metabolism by

839   *Streptococcus mutans*, SL-I. Biochim Biophys Acta 261:379-87.

840   39.   Zeng L, Burne RA. 2013. Comprehensive mutational analysis of sucrose-

841   metabolizing pathways in *Streptococcus mutans* reveals novel roles for

842   the sucrose phosphotransferase system permease. J Bacteriol 195:833-

843   43.

844   40.   Ahn SJ, Rice KC, Oleas J, Bayles KW, Burne RA. 2010. The

845   *Streptococcus mutans* Cid and Lrg systems modulate virulence traits in

846   response to multiple environmental signals. Microbiology 156:3136-47.

847   41.   Seaton K, Ahn SJ, Sagstetter AM, Burne RA. 2011. A transcriptional

848   regulator and ABC transporters link stress tolerance, (p)ppGpp, and

849   genetic competence in *Streptococcus mutans*. J Bacteriol 193:862-74.

850   42.   Burne RA, Wen ZT, Chen YY, Penders JE. 1999. Regulation of

851   expression of the fructan hydrolase gene of *Streptococcus mutans* GS-5

852   by induction and carbon catabolite repression. J Bacteriol 181:2863-71.

853   43.   Wen ZT, Yates D, Ahn SJ, Burne RA. 2010. Biofilm formation and

854   virulence expression by *Streptococcus mutans* are altered when grown in

855   dual-species model. BMC Microbiol 10:111.

856   44.   Sambrook J, Russell DW. 2001. Molecular cloning: a laboratory manual,

857   3[rd] ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

858   45.   Biswas I, Jha JK, Fromm N. 2008. Shuttle expression plasmids for genetic

859   studies in *Streptococcus mutans*. Microbiology 154:2275-82.

860   46.   Zeng L, Burne RA. 2008. Multiple sugar: phosphotransferase system

861   permeases participate in catabolite modification of gene expression in

862   *Streptococcus mutans*. Mol Microbiol 70:197-208.

863   47.   Zeng L, Burne RA. 2015. NagR differentially regulates the expression of

864   the *glmS* and *nagAB* genes required for amino sugar metabolism by

865   *Streptococcus mutans*. J Bacteriol 197:3533-44.

866   48.   Loo CY, Corliss DA, Ganeshkumar N. 2000. *Streptococcus gordonii*

867   biofilm formation: identification of genes that code for biofilm phenotypes.

868   J Bacteriol 182:1374-82.

869   49.   Ahn SJ, Lemos JA, Burne RA. 2005. Role of HtrA in growth and

870   competence of *Streptococcus mutans* UA159. J Bacteriol 187:3028-38.

871   50.   Ajdic D, McShan WM, McLaughlin RE, Savic G, Chang J, Carson MB,

872   Primeaux C, Tian R, Kenton S, Jia H, Lin S, Qian Y, Li S, Zhu H, Najar F,

873   Lai H, White J, Roe BA, Ferretti JJ. 2002. Genome sequence of

874   *Streptococcus mutans* UA159, a cariogenic dental pathogen. Proc Natl

875   Acad Sci U S A 99:14434-9.

Figure legends

FIGURE 1 Overexpression of *sppA* enhances growth of a *fruK* mutant on

fructose. (A) Relative *sppA* mRNA levels measured in UA159/pIB184,

UA159/pIB508, FruK-13/pIB184, and FruK-13/pIB508. Cultures were prepared

using BHI medium and harvested at the exponential phase. Quantitative reverse

transcription PCR (qRT-PCR) was carried out using the total RNA samples, and

the relative abundance of *sppA* was determined using the $\Delta\Delta$Cq method. The

error bars denote the standard deviations. Student's *t*-test was performed to

determine the statistical significance of the data, with asterisks denoting $P < 0.05$.

(B) FruK-13/pIB508 (red squares) and FruK-13/pIB184 (blue diamonds) were

cultivated in BHI and then diluted into TV medium containing 10 mM fructose.

The cultures were overlaid with mineral oil and OD_{600} was measured every 30

min in a Bioscreen C. The results represent the average of at least three

biological replicates.

FIGURE 2 *In vitro* characterization of SppA as a hexose-phosphate:

phosphohydrolase. The majority of the reactions (except panels B&C) were

performed at 37°C in a basic buffer composed of 50 mM PIPES (pH 6.5), 5 mM

$MgCl_2$, and 1 mM $MnCl_2$. Release of phosphate was measured using a Malachite

Green Reagent (MGR), and the results were divided by time and amounts of

enzyme (A, D), or normalized for relative activities (B, C). (A) 10 mM of the

indicated sugar-phosphates were incubated with His-SppA (0.08 µM for F-1-P,

0.67 µM for the rest), with release of phosphate being monitored over a period of

10 min. (B) 0.1 mM of F-1-P was incubated for 5 min with 0.33 µM of His-SppA

and the specified amounts of metal ions. (C) 0.67 mM F-1-P was incubated with

0.04 µM of His-SppA for 130 sec at the indicated pH values (See Methods for

details). (D) Increasing concentrations of F-1-P and F-6-P were each mixed with

His-SppA and phosphate release was monitored over defined periods of time

(See Methods for details).

FIGURE 3 Transcription of the *spp* operon is regulated by SppR, fructose

derivatives and the PTS. The P*sppR::cat* fusion was integrated into the

chromosome of the wild type (A) or various mutants (B) to monitor the expression

of the operon. All strains were first cultivated to mid-exponential phase (OD_{600} =

0.3 to 0.4) in a TV medium with 10 mM glucose, then (A) 25 or 50 mM of the

specified carbohydrates, or (B) 50 mM of glucose or fructose was added, the

cultures were incubated for 1 h, and cells were harvested for CAT assays. Error

bars denote standard deviations. Asterisks indicate statistically significant

difference (by Student's *t*-test, $P < 0.05$) in comparison (A) with glucose at the

same concentrations, or (B) with the wild type (WT) treated with the same

carbohydrates.

FIGURE 4 *In vitro* interactions of SppR with the *spp* promoter region or F-6-P. (A)

EMSA using recombinant His-SppR and a biotin-labeled probe containing the

promoter region of *sppR.* His-SppR (0.05 to 1 µM) was incubated with 0.4 nM of

DNA probe in a 10-µl reaction for 30 min, before the mixture was resolved on a

non-denaturing polyacrylamide gel. Certain reactions also contained 0.4 µM of

unlabeled probe as a competitor, or 5 mM of the specified sugar phosphate. (B)

The melting temperatures (Tm) of His-SppR in the presence of varying amounts

of F-6-P. A small amount of SYPRO Orange was added to monitor the thermal

denaturation of the protein.

FIGURE 5 Overexpression of SppA enhances resistance to xylitol. Growth

phenotypes of various strains were monitored over time in a Bioscreen C (A, B),

or by measuring the final yield (OD_{600}) after overnight incubation (C). Strains

were cultivated first on BHI to exponential phase, then diluted using TV medium

containing 0.5% glucose in addition to (A, B) 1% xylitol, or (C) the amounts of

xylitol specified on the *x*-axis.

FIGURE 6 Fructose metabolism alters biofilm development by *S. mutans.* BM

containing 2 mM sucrose and 18 mM glucose (BMGS) or 18 mM fructose

(BMFS) was used for biofilm development. (A) Wild type UA159 and its isogenic

mutants *levD* and *fruI* were each incubated for 24 h and the amounts of biofilms

formed were visualized and quantified by crystal violet staining. (B) Biofilms

formed by UA159 were treated with a LIVE/DEAD vitality stain and visualized

under a confocal laser-scanning microscope, with green indicating live cells and

red indicating dead cells. (C) eDNA released from 48-h biofilms formed by strains

UA159/pIB184 (blue) and UA159/pIB508 (orange) were quantified using a DNA

dye (SytoX Green) and a standard curve. The results were normalized using total

CFU from each sample (D). Error bars represent standard deviations. Statistical

analysis was performed using Student's *t* test, with asterisks denoting *P* values: *

<0.05, ** <0.005, and *** <0.0005.

FIGURE 7 HAD family phosphatases contribute to sugar-phosphate tolerance.

(A) *S. mutans* UA159 and its mutants deficient in putative phosphohydrolases.

(B) UA159, a *lacD* mutant, and *lacD* derivatives containing plasmid pIB428 that

overexpresses SMU.428 or the vector pIB184. Bacterial cultures were each

prepared in BHI, then diluted into TV base medium supplemented with (A) 0.5%

glucose and various amounts of xylitol or (B) 0.5% sorbitol and various amounts

of galactose. After 17 to 20 h incubation at 37°C, OD_{600} of the cultures were

recorded. Results are the averages of at least three independent cultures, with

error bars denoting the standard deviations.

Table 1. Bacterial strains and plasmids used in this study.

Strains	Relevant genotypes[a]	Source or reference
Streptococcus mutans		
UA159	Wild type, serotype *c*	(50)
FruK-13	*fruK*M28stop	(3)
MMZ1387	UA159 *sppR*::Km^r	This study
MMZ1388	UA159 *sppR*::Em^r	This study
MMZ1399	UA159 *sppA*::Km^r	This study
MMZ1400	UA159 *sppA*::Em^r	This study
MMZ1416	UA159 P*sppR::cat*	This study
MMZ1518	*levD*::Em^r P*sppR::cat*	This study
MMZ1519	*fruI*::Sp^r P*sppR::cat*	This study
MMZ1520	*manL*::Em^r P*sppR::cat*	This study
MMZ1521	*ccpA*::Sp^r P*sppR::cat*	This study
MMZ1549	UA159 SMU.428::Km^r	This study
MMZ1551	UA159 SMU.718c::Km^r	This study
MMZ1570	UA159 SMU.488::Km^r	This study
MMZ1573	*sppR*::Em^r P*sppR::cat*	This study
MMZ1574	*sppA*::Em^r P*sppR::cat*	This study
Plasmids		
pIB184	Shuttle expression vector	(45)
pIB508	*sppA* cloned into pIB184	This study
pIB428	SMU.428 cloned into pIB184	This study

961   [a] Km^r kanamycin resistance; Sp^r, spectinomycin resistance; Em^r, erythromycin

962   resistance.

963 **Table 2.** Primers used in this study.

Primers	Sequence[a]	Purpose
SMU.507-1	5'- GGT TGG GAT AGT GAT AAA TCG CAT -3'	Deleting *sppR*
SMU.507-2GA	5'- <u>GCC ATT TAT TAT TTC CTT CCT CTT TTA</u> GTT CTT GAT ACA TAA CTT TCC TCT TTT C -3'	Deleting *sppR*
SMU.507-3GA	5'- <u>ATA TTT TAC TGG ATG AAT TGT TTT AGT AGA</u> CGG GCA AAT TTA TCT GAT AAG GT -3'	Deleting *sppR*
SMU.507-4	5'- GCC TGT GCA TTG CAA TT AAG A -3'	Deleting *sppR*
SMU.508-S	5'- ATG GGG CTC TGA TTG TTG AG -3'	qRT-PCR
SMU.508-AS	5'- GCG GAA ACT GCT TCT TGA TT -3'	qRT-PCR
SMU.508-5'Bm	5'- CGA GTA AAA <u>GGA TCC</u> GGG CAA ATT TAT CT -3'	pIB508
SMU.508-3'SacI	5'- AAC CAC TT<u>G AGC TC</u>T GTT TTA AAT TAA AAC A -3'	pIB508
SMU.508-1	5'- CGT CGT GAT TTT GCT CTT TTG A -3'	Deleting *sppA*
SMU.508-2GA	5'- <u>GCC ATT TAT TAT TTC CTT CCT CTT TTA</u> CTT TGC CAT AAC AAA CCC C-3'	Deleting *sppA*
SMU.508-3GA	5'- <u>ATA TTT TAC TGG ATG AAT TGT TTT AGT AGG</u> ACA ACG GTG TTT CCA GGT G-3'	Deleting *sppA*
SMU.508-4	5'- CGA GAC ATA TCA GCA GGA TTG AT -3'	Deleting *sppA*
P507-3'Bm	5'- GTT CTT GAT ACA TA<u>G GAT CCC</u> TCT TTT CA -3'	P*sppR::cat* fusion
P507-5'SacI	5'- AAC TTG TAG <u>GAG CTC</u> TCA TAA GTC ATC AA -3'	P*sppR::cat* fusion
SMU.507-5'Bm	5'- TGA AAA GAG G<u>GG ATC C</u>AT GTA TCA AGA ACA -3'	His-SppR

SMU.507-5'HIII	5'- CAC TTA AAA T<u>AA GCT T</u>AA TTG TCT TTG CCA -3'	His-SppR
SMU.508-5'Bm	5'- AAA G<u>GG ATC C</u>GT TAT GGC AAA GAC AAT TAA -3'	His-SppA
SMU.508-3'HIII	5'- CAA AAA ATC <u>AAG CTT</u> CAA AAA CCA CTT GA -3'	His-SppA
SMU.507-EMSA-5'	5'- GCG TGC AGC GA CCT CCT ATT CCT CTT TTG CTA ATG TTA TCT -3'	DNA probe
SMU.507-EMSA-3'	5'- GTT CTT GAT ACA TAA CTT TCC TCT TTT CA -3'	DNA probe
SMU.428-1	5'- CCA TGG GCT CTG GAA CA -3'	Deleting SMU.428
SMU.428-2GA	5'- <u>GCC ATT TAT TAT TTC CTT CCT CTT TTA</u> GCG TGC CAT CCA TAT CAA TAG CA -3'	Deleting SMU.428
SMU.428-3GA	5'- <u>ATA TTT TAC TGG ATG AAT TGT TTT AGT AGA</u> GCT GGT GTT GCT CAA GCT -3'	Deleting SMU.428
SMU.428-4	5'- GGC CAT ACT AAC GTA CAG CA -3'	Deleting SMU.428
SMU.718c-1	5'- GGG TTA CCC TAG AAG ACT ATC TTA GT -3'	Deleting SMU.718c
SMU.718c-2GA	5'- <u>GCC ATT TAT TAT TTC CTT CCT CTT TTA</u> GCC GTC AAT ATC GAC AGC AAC C -3'	Deleting SMU.718c
SMU.718c-3GA	5'- <u>ATA TTT TAC TGG ATG AAT TGT TTT AGT AGA</u> CAG TGG GGT CGC TTA TGC A -3'	Deleting SMU.718c
SMU.718c-4	5'- GAC TAG CCA AAG TAA TGA AAG AGT GT -3'	Deleting SMU.718c
SMU.488-1	5'- GGG TTG TTA TGG ATA TCC ATG TAC -3'	Deleting SMU.488
SMU.488-2GA	5'- <u>GCC ATT TAT TAT TTC CTT CCT CTT TTA</u> CCC GTA TCT TTG ATG CGT AAG -3'	Deleting SMU.488
SMU.488-3GA	5'- <u>ATA TTT TAC TGG ATG AAT TGT TTT AGT AGA</u> GAG AGC ATC GAG GTG ATT GAT -3'	Deleting SMU.488
SMU.488-4	5'- CCA AAA CAT TCA TTA TGC CTT TAC GCA -3'	Deleting SMU.488

SMU.428-5'Bm	5'- GCT GTG CCA <u>GGA TCC</u> TCG ATT TGT GAT ACA CTA -3'	pIB428
SMU.428-3'RI	5'- GAT AAA AAC <u>GAA TTC</u> CCA GTT CCT TTT TCA TGA -3'	pIB428

964
965   [a] An insertion or mutation is indicated by underlining.

A. Substrate specificity

B. Effects of metal ions

MgCl$_2$	2.5, 5, 10 mM
Others	0.5, 1, 2 mM

C. Effects of pH

D. Kinetics

A

0   0.05   0.2   0.5   1.0   0.5   0.5   0.5   0.5   0.5 µM His-SppR

1000x   cold competitor

F-1-P

F-6-P

F-1,6-bP

G-6-P

Unbound
probe

B

A

B