ESSENTIAL ROLES OF THE *SPPRA* FRUCTOSE-PHOSPHATE

PHOSPHOHYDROLASE OPERON IN CARBOHYDRATE METABOLISM AND

VIRULENCE EXPRESSION BY *STREPTOCOCCUS MUTANS*

Lin Zeng and Robert A. Burne[#]

Department of Oral Biology, University of Florida, Gainesville, Florida, USA

Running title: Adaptation to sugar-phosphate stress

Key words: fructose metabolism, sugar-phosphate stress, xylitol, biofilm, sugar-

phosphate phosphohydrolase

[#]Corresponding Author:

1395 Center Drive, Gainesville, Florida 32610

Phone: 352 273-8850

FAX: 352 273-8829

Email: rburne@dental.ufl.edu

Abstract

The dental caries pathogen *Streptococcus mutans* can ferment a variety of sugars to produce organic acids. Exposure of *S. mutans* to certain non-metabolizable carbohydrates such as xylitol impairs growth and can cause cell death. Recently, the presence of a sugar-phosphate stress in *S. mutans* was demonstrated using a mutant lacking 1-phosphofructokinase (FruK) that accumulates fructose-1-phosphate (F-1-P). Here we studied an operon in *S. mutans, sppRA,* which was highly expressed in the *fruK* mutant. Biochemical characterization of a recombinant SppA protein indicated that it possessed hexose-phosphate phosphohydrolase activity, with preferences for F-1-P and, to a lesser degree, fructose-6-phosphate (F-6-P). SppA activity was stimulated by Mg^{2+} and Mn^{2+}, but inhibited by NaF. SppR, a DeoR-family regulator, repressed the expression of the *sppRA* operon to minimum levels in the absence of the fructose-derived metabolites, F-1-P and likely also F-6-P. Accumulation of F-1-P, as a result of growth on fructose, not only induced *sppA* expression, it significantly altered biofilm maturation through increased cell lysis and enhanced extracellular DNA release. Constitutive expression of *sppA,* via a plasmid or by deleting *sppR*, greatly alleviated fructose-induced stress in a *fruK* mutant, enhanced resistance to xylitol, and reversed effects of fructose on biofilm formation. Finally, by identifying three additional putative phosphatases that are capable of promoting sugar-phosphate tolerance, we show that *S. mutans* is capable of mounting a sugar-phosphate stress response by modulating the levels

41   of certain glycolytic intermediates, functions that are interconnected with the

42   ability of the organism to manifest key virulence behaviors.

Importance

Streptococcus mutans is a major etiologic agent for dental caries, primarily due to its ability to form biofilms on tooth surface and to convert carbohydrates into organic acids. We have discovered a two-gene operon in *S. mutans* that regulates fructose metabolism by controlling the levels of fructose-1-phosphate, a potential signaling compound that affects bacterial behaviors. With fructose becoming increasingly common and abundant in the human diet, we reveal the ways fructose may alter bacterial development, stress tolerance, and microbial ecology in the oral cavity to promote oral diseases.

Introduction

Most bacteria use the sugar:phosphotransferase system (PTS) to internalize mono- and di-saccharides, a process that uses phosphoenolpyruvate (PEP) to energize the transport and concomitant phosphorylation of the carbohydrates (1). While certain components of the PTS are vital to the growth and persistence of many bacteria, the accumulation in cells of phosphorylated carbohydrates can lead to sugar-phosphate stress, a condition that often manifests in growth inhibition or cell death (2-4). A number of mechanisms have been proposed to contribute to sugar-phosphate stress: (i) futile cycling, in which the cells expend energy to internalize and phosphorylate the sugar, but the sugar is then dephosphorylated and expelled (5); (ii) reduction of essential metabolic intermediates arising from an inability to internalize other carbohydrates (6); (iii)

creation of toxic metabolic byproducts, such as methylglyoxal (7, 8), or (iv)

deleterious effects of specific sugar phosphates that act as allosteric modifiers of

enzymes or regulatory proteins, causing dysregulation of metabolism (9). The

majority of research on sugar-phosphate stress has been conducted with Gram-

negative, model organisms, with far fewer studies carried out with known

pathogens or Gram-positive bacteria (10).

Streptococcus mutans, a primary dental caries pathogen, is a Gram-positive,

lactic acid-producing bacterium that depends on, and is particular adept at,

transporting and fermenting carbohydrates for energy production and growth.

The most studied sugar-phosphate stress in *S. mutans* is that induced by xylitol,

a non-metabolizable pentitol that is internalized as xylitol-5-phosphate (X-5-P) by

a fructose-specific PTS, but is subsequently dephosphorylated and released from

the cells (11-13). Although the exact nature of the process remains to be

clarified, including the enzyme(s) responsible for removing the phosphate group

from X-5-P, it is generally believed that both the futile cycling of xylitol and the

presence of X-5-P in cells are detrimental. While futile cycling wastes energy, it

has been proposed that X-5-P is an inhibitor of phosphofructokinase that causes

a block in glycolysis (9). The observed effects of xylitol intake by *S. mutans*

include reductions in high-energy metabolites, such as fructose-1,6-bisphosphate

(F-1,6-bP) and glucose-6-phosphate (G-6-P), as well as impaired glucose-PTS

activity, decreased rates of acid production (14), and inhibition of the formation of

extracellular insoluble polysaccharides from sucrose (15, 16). Given its ability to

negatively affect a caries pathogen, xylitol is widely used as a non- or anti-

cariogenic additive in products that include chewing gum and candies, despite

the fact that spontaneous emergence of xylitol-resistant mutants can occur in

habitual users of xylitol-containing products (17, 18). It has been shown that the

fructose-PTS permease encoded by the *fruI* gene is responsible for transporting

the majority of xylitol, and a *fruI*-deficient mutant lost sensitivity to presence of

xylitol (18). Another example of sugar-phosphate-related stress in *S. mutans* is

one induced by 2-deoxyglucose (2-DG), an analog of glucose that can be

internalized and phosphorylated by the PTS, but cannot be metabolized. Uptake

of 2-DG by *S. mutans* reduces the transport of carbohydrates via the PTS (19)

and the multiple sugar metabolism (*msm*) system, an ATP-binding cassette

(ABC) transporter for melibiose, raffinose, and certain other carbohydrates that

are found in the human diet (20).

Fructose is transported into *S. mutans* via multiple PTS permeases and further

metabolism requires phosphofructokinase enzymes, including a 1-

phosphofructokinase and 6-phosphofructokinase that convert fructose-1-

phosphate (F-1-P) and fructose-6-phosphate (F-6-P), respectively, into F-1,6-bP

that can be further catabolized via glycolysis (3). We recently engineered a point

mutant in *S. mutans* strain UA159 (*fruK*M28stop, strain FruK-13) that prevents

the translation of an intact 1-phosphofructokinase (FruK) enzyme (3). Due to the

accumulation of F-1-P in FruK-13, a severe growth phenotype is evident when

fructose is present. FruK-13 also displayed significant growth defects when

cultured in media containing glucose, sucrose, mannose, galactose, lactose or

certain amino sugars, demonstrated a broad influence of F-1-P on carbohydrate

metabolism and growth of *S. mutans* (3). RNA deep sequencing (RNA-Seq) was

used to define changes in the transcriptome arising from loss of the FruK

enzyme. The expression of 394 genes, including genes encoding ribosomal

proteins, proteins required for biosynthesis of nucleotides and amino acids, and

components of the PTS, were altered in FruK-13 compared to the parental strain

UA159 (3). Among the most up-regulated genes in the *fruK* mutant were two

open reading frames (ORF) in an apparent operonic arrangement, SMU.507 and

SMU.508; designated <u>s</u>ugar-<u>p</u>hosphate <u>p</u>hosphatase <u>R</u>egulator *sppR* and the

phosphatase *sppA*, respectively. SppR is a DeoR-type transcription regulator and

SppA is a predicted hydrolase of the HAD (haloacid dehalogenase) family IIB

that includes enzymes such as trehalose-6-phosphatase, plant and

cyanobacterial sucrose-phosphatases, and a large subfamily of bacterial Cof-like

hydrolases; Cof being an *E. coli* phosphatase (21). While both *sppR* and *sppA*

are well conserved in *S. mutans* strains, homologous systems comprised of both

genes in tandem have only been identified in *Streptococcus gordonii* and

Streptococcus sanguinis (http://www.microbesonline.org). In this communication,

we investigated the regulation and function of the *sppRA* operon and explored

the contribution of its gene products to fitness and cariogenic potential of *S.*

mutans, particularly in the context of regulation of F-1-P levels.

Results

Overexpression of *sppA* enhances growth of FruK-13 in the presence of

fructose. In a previous study with a mutant lacking a 1-phosphofructokinase

(*fruK*), FruK-13, an operon encoding a DeoR-like regulator (SMU.507, *sppR*) and

a predicted hydrolase of the HAD superfamily (SMU.508, *sppA*) was significantly

up-regulated (3). As FruK-13 displays the most severe growth defect on fructose,

likely due to accumulation of F-1-P, it was posited that certain enzymes encoded

by *S. mutans* could eliminate these potentially toxic intermediates. The putative

sugar-phosphate phosphatase encoded by *sppA* was cloned into an expression

vector pIB184 behind a constitutive promoter and the resultant plasmid, pIB508,

was introduced into FruK-13. After confirming enhanced expression of *sppA* by

plasmid pIB508 (Fig. 1A), these resultant strains were compared in their growth

phenotypes in a fructose-based medium. Compared to the vector-only control

strain FruK-13/pIB184, which grew very slowly on a TV-fructose medium, FruK-

13/pIB508 grew at a significantly improved rate (Fig. 1B), providing a strong

support for the hypothesis that SppA could function as a sugar-phosphate

phosphatase that decreased the levels of F-1-P in the *fruK* genetic background.

SppA has hexose-phosphate phosphohydrolase activity. Multiple computer

algorithms predicted that SppA is a hydrolase of the HAD superfamily (e.g.

BLAST, CLUSTAL). To study the function of SppA *in vitro*, the coding sequence

of *sppA* was cloned into pQE30 and expression was induced in *E. coli* to produce

an N-terminally His-tagged recombinant protein, designated His-SppA. His-SppA

has 274 amino acids, excluding the His-tag, and migrated as a single 31-kDa

band in SDS-polyacrylamide gel electrophoresis (SDS-PAGE, Fig. S1). The

protein was relatively stable in a variety of buffers and pH values, although

addition of 0.5 M NaCl and at least 10% of glycerol was required to retain activity

during longer-term storage (>1 month) at 4°C. To test the kinetics of SppA in

removing the phosphate group from presumptive substrates, a set of phosphate-

free buffers were tested to carry out these experiments (see Materials and

Methods for detail), and the phosphate released in each reaction was monitored

using a Malachite green reagent kit.

Substrate specificity. Using 2.5 mM *p*-nitrophenylphosphate (*p*NPP) as a

general colorimetric substrate for phosphatases, phosphatase activity by His-

SppA was detected at 37°C in 50 mM HEPES buffer at pH 7.5. SppA activity on

*p*NPP could be stimulated by the addition of millimolar concentrations of $MgCl_2$

(data not shown). We then conducted a number of preliminary experiments to

gain insights into optimal conditions for activity of His-SppA on a number of sugar

phosphates. Then, as determined by an end-point assay performed under

optimized conditions (Fig. 2A), the preferred substrate for His-SppA was found to

be F-1-P, followed by F-6-P. Very little activity was observed when F-1,6-bP, G-

6-P, or GlcN-6-P were used as substrates, and SppA had no activity on GlcNAc-

6-P under the conditions tested.

Effects of metal ions. Many phosphatases are affected by the presence of

metal ions (22), thus the activity of His-SppA was studied by adding to the

reactions millimolar levels of $CaCl_2$, $CuSO_4$, $MgCl_2$, $MnCl_2$, and $ZnCl_2$. $CaCl_2$,

$CuSO_4$, and $ZnCl_2$ at these levels (0.5 to 2 mM) had no detectable impact to the

activity of His-SppA on F-1-P as a substrate, but the presence of $MgCl_2$ and

$MnCl_2$ significantly enhanced the release of phosphate by the purified,

recombinant enzyme (Fig. 2B). By testing both Mg^{2+} and Mn^{2+} together, it was

determined that optimal enzymatic activities could be elicited when $MgCl_2$ was

present at 5 mM, along with 1 mM $MnCl_2$.

pH. The optimal pH for the activities of His-SppA was determined to be about

6.5, and the rate of phosphate release significantly decreased when the pH

values of the reaction deviated from this point, particularly at values of 6.0 and

below (Fig. 2C). Since *S. mutans* maintains ΔpH values of 0.5 to 1.0 units above

the external environment (23), SppA is probably most active in cells growing in a

moderately acidic environment, such as what would occur when *S. mutans* is

actively metabolizing carbohydrate(s).

Fluoride. Sodium fluoride (NaF) is an inhibitor of many phosphatases. The

presence of 10 mM NaF significantly reduced (by >50%) the rate of phosphate

release from F-1-P by His-SppA (Fig. S2 in the supplemental material).

Reaction kinetics on F-1-P and F-6-P. Finally, using the optimal pH and

metal ion conditions described above, V_{max} and K_m of His-SppA were measured

using F-1-P or F-6-P. The purified enzyme was incubated with increasing

concentrations of each substrate (see Materials and Methods for details) at 37°C,

and release of phosphate over time was monitored by periodic sampling of the

reactions. The release rate of phosphate for each reaction was then plotted

against the substrate levels to estimate rate and substrate affinity (Fig. 2D). For

F-1-P, the V_{max} was 9.50 ± 0.38 µmol/(min x mg protein), and the K_m was 1.90 ±

0.21 mM. For F-6-P, the V_{max} was 0.52 ± 0.03 µmol/(min x mg protein) and the

K_m was 4.83 ± 0.51 mM. Thus, under these conditions, F-1-P is a preferred

substrate for SppA, whereas F-6-P can be dephosphorylated by SppA, albeit less

efficiently. These data are consistent with idea that *S. mutans* may use SppA to

limit accumulation of F-1-P, which can be further phosphorylated (by FruK) into

F-1,6-bP. However, the data may also reflect a preference of the organism to

internalize then further metabolize F-6-P. These parameters, including V_{max} and

$K_{m,}$ are also in line with what has been reported for other members of the HAD

family phosphatases in bacteria (5, 21, 24).

SppR is a negative regulator of *sppA* that responds to fructose-phosphate

signals. SppR is predicted to be a member of the DeoR family, a group of

transcription regulators that often respond allosterically to metabolic

intermediates including phosphorylated carbohydrates. To begin understanding

the mode of regulation by SppR, a deletion mutant of *sppR* was constructed and

the expression of *sppA* was measured using quantitative RT-PCR (qRT-PCR).

The results showed a drastic enhancement in mRNA levels of *sppA* due to the

loss of SppR, when both the wild type and the mutant were growing in the rich

medium BHI (Fig. 1A and data not shown). To ensure that increased expression

of *sppA* was not a result of secondary mutations, plasmid pBGE-*sppR* was

created (25) and introduced into the *sppR* mutant, allowing for stable integration

of the *sppR* gene in the chromosome in the *gtfA* locus. Compared to cells

transformed with the empty vector, complementation of the *sppR* mutant with

sppR in single copy resulted in expression levels of *sppA* that were comparable

to those measured in the wild-type background (Fig.1A).

Since higher expression of both *sppR* and *sppA* was noted using RNA-Seq of

the *fruK* mutant (FruK-13), which should have elevated levels of F-1-P, it was

hypothesized that SppR negatively regulates the *spp* operon, with SppR being

allosterically regulated by fructose phosphates, including F-1-P. To test this

hypothesis, the promoter activity of the operon (P*sppR*) was examined by

utilizing a promoter-reporter gene fusion P*sppR::cat* that was integrated in single

copy at a distant site on the chromosome. Expression of the *cat* fusion was

measured in cells growing on various carbohydrates. A UA159 derivative

carrying the P*sppR::cat* fusion was cultivated in TV medium supplemented with

10 mM glucose to mid-exponential phase, then pulsed for 1 h with 25 mM or 50

mM fructose to increase the levels of fructose-derived metabolites. In addition, 25

mM or 50 mM of glucose, xylitol, or sucrose were also used to treat these cells.

The CAT activities measured (Fig. 3A) on cells pulsed with fructose showed

significant, concentration-dependent increases compared to cells pulsed with

glucose. A slight reduction in CAT activity was noted in cells pulsed with 50 mM

xylitol, and sucrose had little to no impact on the expression of the *spp* promoter

fusion. When compared to glucose, treatment of cells with fructose should result

in accumulation of F-1-P, adding support to the idea that F-1-P is an allosteric

effector capable of relieving the repression by SppR. On the other hand, xylitol

treatment could potentially reduce the capacity of the PTS and carbon flux,

reducing the levels of glycolytic metabolites, including F-1-P and F-6-P. The lack

of response from sucrose treatment could be due to the preference by the

bacterium to utilize the glucose moiety, as our recent study has indicated that

free fructose is expelled by *S. mutans* after sucrose is transported and

phosphorylated on the glucose moiety by ScrA, and the resultant sucrose-6-

phosphate (S-6-P) is cleaved by ScrB (26).

We have previously shown that fructose internalization via FruI likely results in

creation of F-1-P, whereas internalization of fructose by EII^{Lev} (*levDEFG*) yields

F-6-P (3). To test if transport of substrates via different PTS permeases could

affect the regulation of expression of the *spp* operon, the P*sppR::cat* fusion was

placed in the background of mutants deficient in the fructose-PTS permeases,

levD or *fruI*, or in a mutant lacking the primary glucose-PTS permease (*manL*)

(27). A mutant defective in production of catabolite control protein A (CcpA), a

major regulator of genes involved in carbon flow in *S. mutans*, was also included

as CcpA can regulate the expression of certain PTS permease genes (28). The

wild-type control and the mutant strains were grown in TV-glucose to mid-

exponential phase, then pulsed with 50 mM fructose to induce the P*sppR::cat*

fusion. In support of our hypothesis that F-1-P is required for inducing expression

of *spp*, loss of *fruI* abolished fructose-dependent induction of CAT activity,

whereas the *levD* mutant behaved like the wild type (Fig. 3B). Interestingly, a

deletion of the glucose-PTS gene *manL* resulted in significantly increased

expression of the P*sppR::cat* fusion, consistent with our recent study that showed

interconnected regulation of fructose and glucose PTS permeases in *S. mutans*

(3). The *ccpA* mutant showed no difference in response to fructose.

To gauge the overall impact of the regulation by SppR on the *sppRA* operon

and the potential that depletion of F-1-P associated with *sppA* induction could

function in feedback regulation of the operon, we assayed the expression of

P*sppR::cat* fusion in strains deficient in *sppR* and *sppA*, respectively. The results

(Fig. 3B) showed nearly a 2-log increase in CAT activities in the *sppR* mutant

regardless of the sugars tested, and no change in the *sppA* mutant. Together,

these results suggest that SppR exerts a strong, negative regulation over the *spp*

operon, resulting in low levels of expression of *sppA* even when fructose is

present, at least in an otherwise wild-type genetic background.

Measurements of fructose metabolites. To demonstrate that the activities of

the *spp* gene products can regulate the levels of intracellular fructose-derived

metabolites, bacterial cells were grown on TV-glucose to mid-exponential phase,

an additional 50 mM glucose or fructose was added, the cultures were incubated

for 30 min, then cells were harvested. Enzymatic assays were performed on the

deproteinized cell lysates to measure F-1-P levels. The results (Table 2) showed

that treatment of the wild-type strain UA159 by fructose increased F-1-P levels by

~40 fold. In comparison to the wild type, the *sppR* mutant had 3-fold lower levels

of F-1-P when exposed to fructose, consistent with the fact that *sppA* is

overexpressed due to the loss of its negative regulator, SppR. Under the same

condition, the *sppA* mutant had similar levels of F-1-P to that of UA159.

Furthermore, the *fruK* deficient strain FruK-13 produced the highest amounts of

F-1-P when exposed to glucose, providing direct evidence that accumulation of

F-1-P is the cause of the growth defect of FruK-13 on various carbohydrates (3).

However, F-1-P levels in FruK-13 exposed to fructose appeared slightly lower

than the wild type, although not statistically significant; an effect likely caused by

the overexpression of *sppA*. Further, in agreement with the finding that loss of

fruI (but not *levD*) abolished induction by fructose of the P*sppR::cat* fusion (Fig.

3B), here only the *fruI* mutant showed reduced F-1-P levels compared to UA159,

when treated with fructose. When F-6-P was measured using these cell lysates,

similar levels of F-6-P were detected in all four strains tested (Table 2), when

each strain was exposed to either glucose or fructose.

SppR binds to the *spp* promoter, F-6-P and F-1-P. To demonstrate the direct

interaction of SppR with its own promoter region, a recombinant MBP (maltose-

binding protein) -SppR was engineered and overexpressed in *E. coli*. The

purified MBP-SppR was cleaved with a protease to release SppR (Fig. S1B),

which was used in an electrophoretic mobility shift assay (EMSA) that included a

biotin-labeled DNA fragment (*spp* probe) containing the 250-bp intergenic region

immediately upstream of the *sppR* coding sequence. As shown in Fig. 4, the

presence of as little as 62.5 nM SppR resulted in significant shifting of the *spp*

probe. Importantly, when the reaction was repeated with the addition of 5 or 10

mM F-1-P, SppR showed significantly lower DNA-binding activities, as more

probe was left unshifted (Fig. 4). Interestingly, F-6-P also showed a comparable

capacity to reduce the DNA-binding activity of SppR. In comparison, F-1,6-bP

had little effect on the interaction of SppR with the probe. As a negative control, a

purified MBP protein was used in EMSA with the *spp* probe and no shifting of the

probe was observed (data not shown).

As an alternative to EMSA, we also performed a thermal shift assay in which

increasing temperature induces unfolding of SppR, which is monitored using a

protein dye that fluoresces when bound to unfolded polypeptides (29). The

melting temperature (Tm) of proteins usually increases in the presence of

compounds that are bound by the protein. We added to the melting reaction

various amounts of glucose, G-6-P, fructose, F-1-P, F-6-P, F-1,6-bP, GlcN, GlcN-

6-P, GlcNAc and GlcNAc-6-P. While most sugar and sugar phosphates had little

to no impact to the unfolding of SppR, addition of F-6-P increased the Tm in a

dose-responsive manner, indicating a specific interaction between SppR and F-

6-P (Fig. S3 and data not shown). Meantime, addition of F-1-P did not

significantly change the Tm of SppR. Considering the results of our genetic

analysis, we posit that the interactions between SppR and fructose-derived

metabolites modulate the repression of *spp* expression. It is intriguing that F-1-P

showed little affinity for SppR in the thermal shift assay, as it was quite effective

at inhibiting the binding of SppR to the *spp* promoter region. It is possible that

efficient interaction of SppR to F-1-P requires the presence of its cognate DNA

molecules or other factors that were present only in the EMSA reactions. In the

future, we plan to utilize a spectrum of binding assays to further evaluate the

possible allosteric effectors that modulate the DNA-binding affinity of SppR *in*

vivo.

Overexpression of SppA enhances resistance to xylitol. Having

demonstrated that enhanced expression of SppA from pIB508 abolished the

sugar-phosphate stress in strain FruK-13, we tested if overexpression of SppA

could also alleviate stress caused by the presence of xylitol. Strains

UA159/pIB184 and UA159/pIB508 were created and their growth was monitored

in TV containing 0.5% glucose or fructose, with or without 1% xylitol. Without

addition of xylitol, both strains grew similarly well in TV glucose or fructose (data

not shown). When 1% xylitol was added to the TV-glucose medium,

UA159/pIB184 displayed a reduced growth rate compared to when the strain was

grown in TV with glucose alone. Interestingly, UA159/pIB508 showed

significantly faster growth than UA159/pIB184 in the presence of both glucose

and xylitol, indicating enhanced resistance to xylitol as a result of increased

expression of SppA (Fig. 5A; see Fig. 1A for *sppA* mRNA levels). In contrast,

addition of 1% xylitol to TV-fructose had no discernable effect on the growth

phenotypes of either strain (data not shown); a result consistent with the notion

that xylitol is internalized via a fructose-specific PTS (FruI) (18). In this case, our

interpretation is that fructose competes better than xylitol for uptake, so less

xylitol is internalized and there is less of an impact on growth.

Since deletion of *sppR* enhanced the expression of *sppA* (Fig. 3B), we

compared resistance to xylitol by the wild type and its otherwise-isogenic *sppR*

and *sppA* mutants. The *sppR* mutant showed significantly improved growth in

TV-glucose supplemented with xylitol, similar to UA159/pIB508 that

overexpressed *sppA* (Fig. 5B&C). However, loss of *sppA* had no effect on xylitol

resistance, as the *sppA* mutant grew similarly to the wild type. These results

agree with our earlier findings that the expression of *sppA* is likely maintained at

minimal levels, perhaps to avoid futile cycling. We have attempted, but have

failed, to demonstrate the capacity of SppA to dephosphorylate xylitol-5-

phosphate (X-5-P)(see Text S1 in the supplemental material), in large part due to

the difficulty producing sufficient amounts of the compound; it is not commercially

available.

Accumulation of F-1-P enhances eDNA release and affects biofilm

formation. *S. mutans* depends on the development of biofilms for its virulence,

but little information is available on the effects of fructose on *S. mutans* biofilms.

Wild-type cultures in exponential phase were grown in BHI and used to inoculate

a biofilm medium (BM) constituted with 2 mM sucrose and 18 mM glucose

(BMGS) or 18 mM fructose (BMFS), as well as a number of combinations of both

glucose and fructose totaling 18 mM hexose. After 24 h or 48 h of incubation, the

supernates of the culture were removed and development of biofilms was

visualized and quantified by staining with 0.1% crystal violet. As shown in Fig.

6A, growth on fructose (BMFS) changed the accumulation of biomass by *S.*

mutans on a polystyrene surface, resulting in a thinner but more uniform biofilm,

as opposed to a patchy coverage by the biofilms formed on BMGS. Elution of

crystal violet also indicated that greater amounts of dye were retained by the

biofilms formed on fructose than glucose (Fig. 6A). This effect was not likely a

result of reduced growth caused by a lack of glucose, as similar phenotypes were

visible by the presence of as little as 3 mM fructose (data not shown). Also, to

differentiate between effects of internalization of F-1-P versus F-6-P, we included

the otherwise-isogenic mutants of UA159 deficient in either *fruI* (F-1-P) or *levD*

(F-6-P). The results (Fig. 6A) showed that while the wild type and both mutants

formed similar biofilms in BMGS, the *fruI* mutant formed patchy biofilms on

BMFS, resembling those on BMGS. The *levD* mutant behaved like the wild type.

Recent research has suggested that the ability of *S. mutans* to form efficient

biofilms depends on both the synthesis of glucans and the release of

extracellular DNA (eDNA) by bacteria, via both lysis-dependent and -independent

pathways (30). To further study the effects of fructose, the release of eDNA was

quantified using a fluorescent DNA dye in the biofilms of *S. mutans*. The results

were normalized against the viable counts of bacteria (CFU). When grown as

biofilms in BMFS, UA159 released significantly more eDNA per CFU after 24 h

(5-fold) or 48 h (17-fold) of growth compared to BMGS medium (similar to

UA159/pIB184 in Fig. 6C). Significantly lower CFU (5-fold for 24 h, and 20-fold

for 48 h) were recovered from biofilms formed on BMFS than BMGS. As such,

the amounts of eDNA detected per unit volume of supernatant fluid were

comparable for both BMGS and BMFS. Therefore, it appears that the bacteria

incubated in the presence of fructose experienced significantly more lysis, which

resulted in increased release of eDNA and changes in formation of biofilms;

although at this point we cannot rule out the involvement of other routes for

production of eDNA. When treated with a LIVE/DEAD stain and observed under

a confocal laser-scanning microscope, the biofilms formed on BMFS indeed

showed notably more dead cells than those formed on BMGS (Fig. 6B). Similar

impacts of fructose were observed for biofilm development by other genomically

distinct, wild-type isolates of *S. mutans* (31)*,* so the phenomenon is not restricted

to the lab strain UA159 (Fig. S4 in the supplemental material). Furthermore, we

compared the sensitivity of planktonic *S. mutans* cells to the cell wall-degrading

enzymes lysozyme and mutanolysin by extracting chromosomal DNA after

enzymatic treatment, using batch cultures grown in glucose- or fructose-

containing medium. The results showed that 10-fold more DNA could be

harvested from UA159 that was cultured in TV-fructose, compared to cells

cultivated in TV-glucose, after each was treated with lysozyme, or a combination

of lysozyme and mutanolysin, under otherwise-identical conditions (Fig. S5 in the

supplemental material).

As a further test of our hypothesis that F-1-P in particular is responsible for the

fructose effects observed so far, the strains UA159/pIB508 and UA159/pIB184

were compared in their ability to produce eDNA in the 48-h biofilm model. Similar

to UA159, strain UA159/pIB184 yielded significantly lower numbers of CFU, but

much higher levels of eDNA/CFU when growing on BMFS compared BMGS (Fig.

6C). When *sppA* was overexpressed on pIB508, significant reductions in the

release of eDNA/CFU were apparent in comparison to strain UA159/pIB184, an

impact that was especially significant for cells cultured in BMFS. At the same

time, bacterial viable counts also showed significant recovery as an effect of

pIB508, particularly when assayed using BMFS (Fig. 6D). These results suggest

that accumulation of F-1-P in *S. mutans* likely leads to alterations in the cell-

envelope that results in enhanced autolysis and release of eDNA*,* causing

aberrant biofilm phenotypes.

Other putative phosphatases capable of promoting sugar-phosphate

resistance. The existence in *S. mutans* of a xylitol-inducible phosphatase

capable of dephosphorylating X-5-P has been proposed (32). As xylitol reduced

spp expression (Fig. 3A), we hypothesized that other phosphatase(s) are

produced that may act on xylitol. Based on sequence similarity to SppA and other

bacterial sugar-phosphate phosphatases, a functional genomics approach was

employed to identify genes encoding products that could promote resistance to

xylitol. A total of 9 ORFs (SMU.406c, SMU.428, SMU.488, SMU.718c,

SMU.719c, SMU.742, SMU.1108c, SMU.1171c, and SMU.1830c) were selected

and mutated by allelic exchange with an antibiotic resistance marker, and three

(SMU.428, SMU.488 and SMU.718c) were chosen as potential candidates based

on the decreased growth rates (Fig. S6 in the supplemental material) and yields

(Fig. 7A) of the mutants in the presence of xylitol, compared to the wild type. For

confirmation, these genes were expressed constitutively using pIB184, in both

the wild-type and the respective mutant genetic backgrounds. When assayed for

growth on TV-glucose supplemented with xylitol, mutants that were

complemented by each of these three genes regained resistance to xylitol, and

UA159 containing SMU.718c or SMU.488 expressed from pIB184 also showed

enhanced yields, compared to their respective vector-only controls (Fig. S7 in the

supplemental material). Furthermore, we previously reported that a deletion of

the tagatose-1,6-bisphosphate aldolase (LacD) results in sensitivity to galactose

in *S. mutans,* likely due to accumulation of metabolic intermediates that are

normally catabolized by the tagatose pathway (33). When pIB428 was introduced

into the *lacD* mutant, there was a significant alleviation of the inhibitory effects of

galactose (Fig. 7B). We found that SMU.428, SMU.488 and SMU.718c are well

conserved in the genomes of a diverse group of *S. mutans* isolates (31)*.* Amino

acid sequence alignments performed using these three proteins, together with

SppA and the AphA phosphatase from *E. coli* (34), showed significant

conservation in those amino acid residues (Fig. S8 in the supplemental material)

that place these proteins in the family of bacterial class B acid phosphatases,

part of the DDDD superfamily of phosphohydrolases (35). Computer analysis

failed to identify any potential signal peptide sequences in the N-terminus of

these proteins, suggesting they are not secreted. Further research is needed to

understand their contribution to xylitol resistance and more broadly, to sugar-

phosphate stress.

Discussion

Carbohydrate fermentation is a highly dynamic process where the relative

activities of various enzymes and resultant intermediates must be maintained at

optimal levels to ensure fitness and survival, and to balance anabolic and

catabolic processes. Perturbations to the systems, such as those that occur in

the oral microbiome during intake of dietary carbohydrates by the host, require

rapid responses by the microbiota. Sugar-related stress responses mediated by

complex regulatory systems have been characterized mostly in Gram-negative

model bacteria, e.g., *E. coli* (8, 36, 37). As a major caries pathogen, *S. mutans*

depends on carbohydrate fermentation for energy generation and virulence. *S.*

mutans is particularly well adapted to metabolizing sucrose, in part through the

activities of three secreted glucosyltransferases (GtfB, GtfC and GtfD) that

convert sucrose into glucan polysaccharides that serve as the scaffolding for

biofilm accumulation and in creation of a diffusion-limiting matrix (38).

Interestingly, Gtfs generate a molecule of free fructose for every glucose moiety

of sucrose that is targeted toward glucan formation. Furthermore, it is now clear

that the majority of the sucrose presented to *S. mutans* is internalized by a

sucrose-specific PTS permease (39). After internalization and phosphorylation,

sucrose is cleaved by a sucrose-6-phosphate hydrolase into G-6-P and fructose.

However, a significant portion of the fructose thus generated is expelled into the

environment and re-internalized by fructose PTS permeases (26, 40). These

observations, when coupled with the fact that fructose itself is a major constituent

of foodstuffs, e.g. with high-fructose corn syrups increasingly used as common

sweeteners, necessitate a better understanding of the metabolism of fructose by

the oral microbiome.

S. mutans maintains at least three PTS permease operons (*fruRKI*, *fruPCD*

and *levDEFG*) plus three phosphofructokinases (FruK, FruP, and

SMU.1191/Pfk). Further, while a number of *S. mutans* PTS permeases can

transport sugars other than their primary cognate carbohydrate, there are no

other instances in *S. mutans* of redundancy in PTS permeases for sugars,

besides those for fructose. This redundancy in PTS permeases, in itself, is

noteworthy and could reflect significant evolutionary benefits to redundant

fructose catabolic systems. Much remains to be clarified regarding the

contributions of these PTS operons to the establishment and persistence of *S.*

mutans in the human oral cavity, and to the organism's cariogenic potential.

However, it is understood that both F-1-P and F-6-P generated by the PTS are

further phosphorylated to create F-1,6-bP, likely via 1-phosphofructokinase FruK

and 6-phosphofructokinase SMU.1191, respectively. F-1,6-bP is then cleaved by

an F-1,6-bP-specfic aldolase into glyceraldehyde 3-phosphate and

dihydroxyacetone phosphate (3). We have demonstrated that removal of FruK

leads to significant growth and fitness defects in *S. mutans*, apparently due to

accumulation of the metabolic intermediate F-1-P (3). Here we present evidence

that *S. mutans* maintains tight control over the levels of F-1-P through an

inducible s̲ugar-p̲hosphate p̲hosphatase (*spp*) operon, consisting of a

transcriptional regulator SppR and a fructose-phosphate hydrolase SppA.

Changes in F-1-P levels, either increase or decrease, significantly alter bacterial

physiology, including cell envelope integrity and biofilm development. We

hypothesize that, aside from being an apparent stress compound in *S. mutans*

(6), that F-1-P may also act as a signaling molecule, the accumulation of which

promotes autolysis and release of extracellular DNA; with the latter being a

critical contributor to formation of biofilms (30).

In support of this hypothesis, our latest RNA deep sequencing analysis in *S.*

mutans has identified a fructose regulon that includes a significant number of

genes that are likely involved in stress responses (26). For example, growth on

fructose up-regulated the *cid* operon (SMU.1700c~1703c) that influences

oxidative stress resistance and modulation of autolysis and biofilm development

(41). Also up-regulated were the *rcrRPQ* and *relPRS* operons (SMU.921 to

SMU.928) which have major impacts on acid and oxidative stress tolerance,

competence development and (p)ppGpp metabolism (42). Thus, fructose and its

intermediates are directly linked to pathways that are critical for persistence,

growth and survival decisions, nutrient acquisition and genetic diversification

(26).

On the other hand, a growing body of evidence suggests that F-1-P is acting

as a regulatory molecule for *S. mutans*. Our recent study on the function of the

fruRKI operon has detailed the transcriptomic impact of accumulation of F-1-P, a

compound that likely serves as an allosteric effector to another DeoR-type

regulator, FruR (3). It was clear that F-1-P influences gene regulation beyond the

fruRKI operon. Specifically, the *fruK* mutant showed altered expression in as

many as 20 known or probable transcriptional regulators, impacting ~20% of the

genome. Key functions ranging from energy metabolism, protein and nucleic acid

biosynthesis, to stress tolerance were affected by fructose. In addition to showing

that overproduction of SppA, which preferentially degrades F-1-P, leads to

phenotypic changes in both xylitol resistance and eDNA release, we have shown

that the ability of fructose to induce *spp* operon expression depends on an intact

FruI protein, which encodes the fructose-PTS permease that creates F-1-P (3).

The ability of F-1-P to inhibit DNA-binding activity of the purified SppR protein *in*

vitro provided direct evidence for its role as a regulatory effector. At the same

time, the interaction of SppR with F-6-P that we observed also points to further

complexity in this system. Notably, we have performed the same protein:DNA

and protein:metabolite interaction assays using a different, His-tagged version of

the SppR recombinant protein, and observed similar effects of both F-1-P and F-

6-P on the binding of SppR to the promoter region of *spp* operon (data not

shown). At first glance, F-6-P is unlikely the ideal signal molecule for fructose

metabolism as it can also be generated from G-6-P through isomerization.

However, F-6-P could act as a secondary allosteric effector for SppR under

certain conditions, perhaps in a cooperative manner with F-1-P. Future research

will focus on identifying other regulatory circuits that respond to the accumulation

of F-1-P or F-6-P, both directly and indirectly, in particular the pathway(s)

responsible for cell envelope integrity and biofilm phenotypes. As stated earlier,

the *spp* operon has only been identified in a few oral streptococcal species, and

its specificity for F-1-P could have implications in understanding niche

adaptations and oral microbial ecology.

An intracellular hexose-6-phosphate:phosphohydrolase was purified from

Lactococcus lactis that has a preference for galactose-6-phosphate and F-6-P

(24). This enzyme had an estimated M_r of 60 KDa by SDS-PAGE, was most

active at pH 6.0, and its activity was stimulated by Mn^{2+}, Mg^{2+} and Fe^{2+}, and

inhibited by fluoride. Another bacterial sugar phosphate phosphatase, Spp, was

recently identified in *Pseudomonas fluorescens* that was capable of

dephosphorylating ribose-5-P, F-6-P, and G-6-P (5). The optimal Mg^{2+}

concentration for Spp was determined to be 2.5 mM, whereas addition of Mn^{2+}

reduced the reaction rate. Therefore, SppA appears to share some similarities

with other bacterial phosphohydrolases, yet it stands out with its preference for F-

1-P. Lastly, despite our repeated efforts, we were not able to determine whether

X-5-P is a substrate of SppA, partly due to our inability to obtain or generate

sufficient X-5-P (it is not commercially available). As we continue to resolve this

technical issue, it is important to note the possibility that SppA alleviates xylitol-

related stress through its activities on either F-1-P or F-6-P, as X-5-P has been

suggested to inhibit the activities of phosphofructokinases (9). The same

rationale could apply in interpreting the xylitol phenotypes associated with the

mutants that lacked one of three additional HAD phosphatases; that each

enzyme may have multiple substrates and a stress phenotype could be alleviated

by actions on molecules other than the one triggering the stress. Most notably,

overexpression of SMU.428 abolished galactose-mediated stress in a *lacD*

mutant. SMU.428 is located immediately downstream of the *copYAZ* operon that

is responsible for copper efflux, biofilm formation and stress tolerance, yet its

homologues in a number of other streptococci are closely associated with the

tagatose operon. Clearly though, *S. mutans* possesses multiple

phosphohydrolases and related regulatory proteins capable of mounting a stress

response when triggered by certain sugar-phosphates. Additional research is

needed to understand the function and regulation of these putative sugar-

phosphate phosphatases.

Materials and Methods

Bacterial strains and culture conditions. *S. mutans* strains (Table 1) were

maintained using brain heart infusion (BHI)(Difco Laboratories, Detroit, MI) agar

plates or in liquid BHI medium at 37°C, in an ambient atmosphere supplemented

with 5% CO_2. To monitor bacterial growth phenotypes and for most other

experiments, a semi-synthetic tryptone-vitamin (TV) medium (43) was utilized

with various carbohydrates being provided at specified concentrations. Overnight

bacterial cultures were prepared in BHI, then sub-cultured to exponential phase,

and inoculated at 1:100 dilutions into TV media. Growth was monitored as the

change in optical density at 600 nm (OD_{600}) using a Bioscreen C lab system

(Labsystems Oy, Helsinki, Finland). For development of biofilms in 96-well

polystyrene plates or ibidi glass chambers (ibidi GmbH, Martinsried, Germany), a

biofilm medium (BM)(44) was utilized that contained 2 mM sucrose and varying

amounts of glucose or fructose, with bacterial inoculum prepared in BHI. Each of

these experiments was repeated at least three times using independent cultures

(biological replicates), with each sample being represented by at least two wells

(technical replicates).

DNA methodology. Standard recombinant DNA techniques were employed in

preparation of genetic materials for applications such as cloning and

mutagenesis (45). To engineer a mutant defective in *sppR*, a set of 4 primers

(see Table S1 for all primers used in this study) were designed for PCR

amplification of two DNA fragments flanking the coding sequence of *sppR*

(upstream fragment: SMU.507-1 and SMU.507-2GA; downstream fragment:

SMU.507-3GA and SMU.507-4). PCR reactions were carried out using these

primers to generate two DNA fragments that shared >25 bp overlaps with the 5'-

and 3'- ends, respectively, of a kanamycin-resistance cassette. Subsequently,

these flanking DNA fragments were used in equimolar concentrations, together

with the kanamycin cassette, in a one-step ligation reaction that was catalyzed by

a Gibson assembly kit (New England Biolabs, Beverly, MA). The ligated product

was then used to transform a naturally competent UA159 culture, followed by

selection on BHI agar plates containing kanamycin (1 mg/ml). Similar procedures

were carried out in the creation of mutants defective in *sppA*, SMU.428,

SMU.488, SMU.718c and other genes. All strains created in this study have been

validated by DNA sequencing to ensure that no unintended mutations were

introduced into the flanking regions of the gene of interest.

To construct a complementing plasmid for the *sppA* mutant, a pair of primers

(SMU.508-5'Bm and SMU.508-3'SacI) were designed and used for amplification

of the coding sequence of *sppA,* together with its putative ribosome-binding site.

This DNA product was treated with restriction enzymes *Bam*HI and *Sac*I, and

then ligated with an expression vector pIB184 (46) that was linearized with the

same enzymes. After cloning into an *Escherichia coli* host strain BH10B, the

resultant plasmid pIB508 was verified to contain the desired coding and

regulatory sequences and introduced into the *sppA* mutant background, as well

as the wild type UA159. Three additional plasmids, pIB428, pIB488, and

pIB718c, were constructed in the same way, in the pIB184 vector, and used to

complement their respective mutants. To complement the *sppR* mutant, the

coding sequence of *sppR* was PCR-amplified (primers: sppR-5'XbaI and sppR-

3'BsrGI) and cloned into an integration vector, pBGE, that places inserted DNA

fragment under the control of the promoter of the *gtfA* gene (25). To monitor the

transcriptional activities of the *spp* operon, a P*sppR::cat* fusion was constructed

by cloning a DNA fragment containing the intergenic region upstream of *sppR*

onto the integration vector pJL84 that contained a promoterless *cat* gene (47),

and the gene fusion was subsequently delivered into a distal site on the

chromosome of *S. mutans* by double cross-over recombination. The sequence

information of all these primers is included in Table S1.

To assess the resistance of different UA159 cultures to muralytic enzymes, the

bacteria were cultivated overnight in 5 ml TV medium containing 0.5% glucose or

fructose, harvested by centrifugation at 4°C and washed once with a TE buffer

(10 mM Tris-Cl, 1 mM EDTA, pH 8.0). Cells were resuspended in 500 µl of lysis

buffer that contained 50 mM Tris-Cl at pH 8.0, 10 mM EDTA, 10 mg/ml of

lysozyme, and 0.1 mg/ml of RNase A. After addition of 0 or 20 units of

mutanolysin (Millipore Sigma, St. Louis, MO), cells were incubated at 37°C for 1

h, and lysed by adding 40 µl of 20% SDS and incubating at 80°C for 15 min.

Subsequently, 80 µl of 5 M NaCl was added into each reaction, mixed well,

followed by addition of 600 µl of phenol: chloroform: isoamyl alcohol (25: 24: 1,

v/v/v) and 2 min of vortexing. The solution was then clarified by centrifugation at

room temperature at 15000 $\times$ *g* for 5 min, followed by the removal of 500 µl of the

aqueous phase. The genomic DNA was then precipitated from the aqueous

solution by addition of 500 µl 100% isopropanol and centrifugation as before.

After washing once with 75% ethanol, the DNA was air-dried and dissolved in 50

µl TE buffer and quantified using a Nanodrop apparatus (Thermo Fisher,

Waltham, MA) and agarose-gel electrophoresis.

Engineering and preparation of recombinant proteins of SppR and SppA.

DNA fragments containing the entire coding sequences of SppR and SppA were

each amplified using PCR primers (SMU.507-5'Bm and SMU.507-3'HIII for *sppR*,

and SMU.508-5'Bm and SMU.508-3'HIII for *sppA*)(Table S1) containing

engineered restriction sites, and digested using restriction enzymes *Bam*HI and

*Hin*dIII. After ligating with a pQE30 vector prepared by treating with the same

enzymes to create an N-terminal, 6× His-tagged SppA, the construct was

introduced into *E. coli* M15. Similarly, the digested product of *sppR* was cloned

into a plasmid vector pMAL-p2X and expressed as a fusion protein (MBP-SppR)

in *E. coli* NEB Express. In order to produce a recombinant 1-

phosphofructokinase for the F-1-P assay (below), an *E. coli* Fru1PK gene was

also cloned into pQE30, using primers 1-PFKEcoli-5Bm and 1-PFKEcoli-3PstI

(48). After confirming the fidelity of the PCR and cloning via DNA sequencing, the

expression of each recombinant protein was induced by adding IPTG to a final

concentration of 0.05 mM to cells growing at mid-exponential phase.

Purification of recombinant proteins, MBP-SppR and His-SppA, was carried

out using an amylose resin (New England Biolabs) and an Ni-NTA agarose

(Qiagen), respectively, according to protocols provided by the suppliers, with

some modifications. Briefly, to improve the stability of MBP-SppR, 10% of

glycerol and 0.3 M of NaCl were supplemented to the original column buffer (20

mM Tris-Cl, 0.2 M NaCl, and 1 mM EDTA, pH 7.5). After elution from the column,

MBP-SppR was treated with Factor Xa (New England Biolabs) to release the

SppR protein, and the products were dialyzed three times, at a 1:1000 dilution,

against a buffer containing 10 mM HEPES pH 7.5, 0.5 M NaCl and 20% glycerol,

using a Slide-A-Lyzer dialysis device with a MWCO of 3.5 KDa (Thermo Fisher).

To avoid carryover of phosphate into the subsequent reactions involving the

recombinant SppA, purification of His-SppA was performed using a phosphate-

free buffer system that was created by replacing NaH_2PO_4 in the original buffers

with 50 mM HEPES (pH 7.9), and also by supplementing with 10% glycerol.

Dialysis of the purified His-SppA solution was carried out against 10 mM PIPES

pH 6.5, 0.5 M NaCl, and 20% glycerol.

Electrophoretic mobility shift assay (EMSA) and thermal shift. To study the

interaction of SppR with the promoter of *sppR*, EMSA was performed using a

biotin-labeled DNA fragment containing the intergenic region upstream of the

sppR coding sequence as previously described (49). When needed, various

sugar phosphates were each used at 5 or 10 mM. To study the interaction of

SppR with various sugar phosphates, a 10-μM SppR solution was prepared

using a buffer containing 0.1 M HEPES (pH 7.5) and 0.15 M NaCl. After addition

of a SYPRO Orange protein stain (Thermo Fisher), used at a 300-fold dilution,

and various sugar phosphate at specified amounts, the 10-µL reactions were

subjected to a melting-curve treatment on a CFX96 Touch real-time PCR

thermocycler (Bio-Rad, Hercules, CA) by increasing the temperature from 25°C

to 75°C in increments of 0.2°C. The RFU measurements were then plotted

against the temperature values to determine the melting temperature (Tm) (29).

Phosphatase assay. The ability of His-SppA to remove the phosphate group

from various sugar-phosphates was measured at 37°C. The basic buffer was

composed of 50 mM PIPES pH 6.5, 5 mM $MgCl_2$, and 1 mM $MnCl_2$. For other pH

values, different buffering reagents were used in place of PIPES at the same

concentration: 2-ethanesulfonic acid (MES) or potassium citrate (K-Cit) for pH

5.5; 1,4-piperazinediethanesulfonic acid (PIPES) for pH 6.0 and pH 6.5; and

HEPES for pH 7.0 and pH 7.5. All initial experiments to determine the effects of

metal ions were carried out in an end-point assay, using HEPES-based buffers

prepared at neutral pH.

To measure the reaction kinetics of His-SppA on F-1-P, 20 µL of His-SppA that

was diluted to 1 µM using the dialysis buffer was mixed with 40 µL of F-1-P

(Santa Cruz Biotechnology, Dallas, TX) prepared in the basic buffer, at final

concentrations ranging from 0.1 mM to 1.0 mM; for F-1-P ranging from 1.5 mM to

10 mM, 0.25 µM of protein was used. All components were pre-warmed to 37°C

before mixing. With the reaction maintained at 37°C, a 5-µL aliquot of the

reaction was removed at 1 min intervals and subjected to measurement of free

phosphate using a phosphate assay kit (Malachite Green Reagent, MGR;

Sigma). For the kinetics of His-SppA with F-6-P as a substrate, 20 µL of His-

SppA at 2 µM was used in each reaction, and aliquots of the reaction were taken

and measured every 5 min. Each reaction was repeated twice. The data were

analyzed using the Prism from GraphPad software (La Jolla, CA).

F-1-P and F-6-P assays. Bacteria were cultivated overnight in TV with 10 mM

glucose, then diluted 1:20 ratio into 50 mL of the same medium and sub-cultured

until the OD_{600} reached 0.3 ~ 0.4. 50 mM glucose or fructose was then added

into the culture. After another 30 min of incubation, cells were harvested by

centrifugation at 4°C, immediately resuspended in 600 µL PBS, and mixed with

500 µL of glass beads. After two rounds of homogenization at 4°C, cell lysates

were clarified by centrifugation (16,000 × g, 10 min, 4°C). A BCA assay (Thermo

Fisher) was carried out to quantify the protein concentrations. Deproteinized

lysates were generated by mixing 400 µL of cell lysate with 100 µL of 100% (w/v)

trichloroacetic acid and incubating on ice for 10 min. After centrifugation, 420 µL

of the supernate was removed and neutralized by adding 50 µL of 6 N NaOH,

and the pH value of each sample was confirmed using a pH-indicator strip. The

deproteinized cell lysate was used immediately for enzymatic assays or stored at

-80°C.

Each 1 mL master mix of the F-1-P assay included 20 mM HEPES, pH 7.1, 50

mM KCl, 5 mM $MgCl_2$, 1 mM DTT, 2.5 mM ATP, 0.05 mM NADH, 1 µL glycerol-

3-P dehydrogenase (Sigma), 10 µg triose-P isomerase (Sigma), 10 µg of rabbit

muscle aldolase in ammonium sulfate suspension (Sigma), and 5 µL of His-1PFK

(18.6 µg/µl) that was engineered based on the *E. coli* 1-phosphofructokinase

gene. 10 µL of deproteinized cell lysate was mixed with 90 µL of the master mix.

After 15 min of incubation at 37°C, optical density OD_{340} was measured using a

spectrophotometer DU640 and a quartz cuvette, and the F-1-P levels were

determined against a standard curve prepared using F-1-P at concentrations

ranging from 1 to 20 µM. A control reaction was included for each sample by

omitting the His-1PFK protein, and the measurements were subtracted from the

actual results. Each strain was represented by 3 biological replicates, and each

sample was measured at least twice. Levels of F-6-P in the same lysates were

measured using a F-6-P assay kit (Sigma) by following protocols provided by the

supplier.

Biofilm development, microscopy and measurement of extracellular DNA

(eDNA). Development of biofilms by *S. mutans* was investigated in 96-well

polystyrene plates (Thermo Fisher) for comparison of overall biofilm mass, or on

glass surfaces using an ibidi 8-well µ-Slide to assess structural changes and

release of eDNA. A semi-defined biofilm medium (BM) was modified to contain

final concentrations of 2 mM sucrose and 18 mM glucose (BMGS) or fructose

(BMFS)(3, 50). Exponential-phase bacterial cultures prepared using BHI were

diluted 1:100 into BMGS or BMFS and incubated at 37°C in an aerobic

environment supplemented with 5% CO_2. Each experiment included at least

three biological replicates. The 96-well plates were incubated for 24 to 48 h,

washed with phosphate-buffered saline (PBS), and stained with 0.1% crystal

violet (CV). After extensive washing with water, the retained CV was eluted with

a solution of 30% acetic acid and quantified by measuring the optical density at

575 nm (OD_{575}).

The ibidi slides were incubated for 48 h, with a replacement of medium after

first 24 h. For observation of biofilm structures, each biofilm was treated using a

LIVE/DEAD BacLight bacterial vitality stain kit (Thermo Fisher) and visualized

using a confocal laser scanning microscopy system (4). To assess the amount of

eDNA released by each biofilm, the entirety of the 48-h biofilm culture was

harvested by scraping off the biofilm material, without washing, and sonicated for

30 sec at 100% power using a water-bath sonicator (model FB120, Thermo

Fisher). A portion of the cell suspension was subjected to serial dilutions and

plated on BHI plates for enumeration of CFU, and the remaining material was

pelleted using a tabletop centrifuge for 2 min at $16,000 \times g$. Subsequently, a

portion of the supernatant fluid was removed and combined 1:1 with a 5 µM

solution of SYTOX Green nucleic acid stain (Thermo Fisher), followed by

measurement of fluorescence (excitation/emission: 504/523 nm) using a

fluorescence-capable Synergy 2 plate reader from BioTek (Winooski, VT). A

standard curve was established using a known DNA solution and the results

were calculated as ng of eDNA released per CFU of the biofilm.

Quantitative reverse transcription PCR (qRT-PCR). Total RNA was extracted

from *S. mutans* cells harvested at $OD_{600} \approx 0.5$ using an RNeasy kit (Qiagen) by

following an established protocol (51). Primers specific to *sppA* were used for

both the reverse transcription and Real-time PCR that followed, with *gyrA* serving

as an internal control (26).

CAT (*c*hloramphenicol *a*cetyl*t*ransferase) assay. To assess the expression of

P*sppR::cat* fusion in various backgrounds, specific CAT activities from clarified

bacterial lysates were measured according to an established protocol detailed

elsewhere (47).

Acknowledgment

This work was supported by DE12236 from the National Institute of Dental and

Craniofacial Research.

References

1.   Postma PW, Lengeler JW, Jacobson GR. 1993.

Phosphoenolpyruvate:carbohydrate phosphotransferase systems of

bacteria. Microbiol Rev 57:543-94.

2.   Irani MH, Maitra PK. 1977. Properties of *Escherichia coli* mutants deficient

in enzymes of glycolysis. J Bacteriol 132:398-410.

3.   Zeng L, Chakraborty B, Farivar T, Burne RA. 2017. Coordinated regulation

of the EII^{Man} and *fruRKI* operons of *Streptococcus mutans* by global and

fructose-specific pathways. Appl Environ Microbiol 83. e01403-17.

4.   Zeng L, Farivar T, Burne RA. 2016. Amino sugars enhance

824   competitiveness of beneficial commensals with *Streptococcus mutans*

825   through multiple mechanisms. Appl Environ Microbiol 82:3671-82.

826   5.   Maleki S, Hrudikova R, Zotchev SB, Ertesvåg H. 2017. Identification of a

827   new phosphatase enzyme potentially involved in the sugar phosphate

828   stress response in *Pseudomonas fluorescens*. Appl Environ Microbiol 83.

829   e02361-16.

830   6.   Richards GR, Patel MV, Lloyd CR, Vanderpool CK. 2013. Depletion of

831   glycolytic intermediates plays a key role in glucose-phosphate stress in

832   *Escherichia coli*. J Bacteriol 195:4816-25.

833   7.   Korithoski B, Levesque CM, Cvitkovitch DG. 2007. Involvement of the

834   detoxifying enzyme lactoylglutathione lyase in *Streptococcus mutans*

835   aciduricity. J Bacteriol 189:7586-92.

836   8.   Kadner RJ, Murphy GP, Stephens CM. 1992. Two mechanisms for growth

837   inhibition by elevated transport of sugar phosphates in *Escherichia coli*. J

838   Gen Microbiol 138:2007-14.

839   9.   Assev S, Rolla G. 1986. Further studies on the growth inhibition of

840   *Streptococcus mutans* OMZ 176 by xylitol. Acta Pathol Microbiol Immunol

841   Scand B 94:97-102.

842   10.   Vanderpool CK. 2007. Physiological consequences of small RNA-

843   mediated regulation of glucose-phosphate stress. Curr Opin Microbiol

844   10:146-51.

11.   Tanzer JM, Thompson A, Wen ZT, Burne RA. 2006. *Streptococcus*

mutans: fructose transport, xylitol resistance, and virulence. J Dent Res

85:369-73.

12.   Trahan L. 1995. Xylitol: a review of its action on mutans streptococci and

dental plaque--its clinical significance. Int Dent J 45:77-92.

13.   Trahan L, Neron S, Bareil M. 1991. Intracellular xylitol-phosphate

hydrolysis and efflux of xylitol in *Streptococcus sobrinus*. Oral Microbiol

Immunol 6:41-50.

14.   Miyasawa-Hori H, Aizawa S, Takahashi N. 2006. Difference in the xylitol

sensitivity of acid production among *Streptococcus mutans* strains and the

biochemical mechanism. Oral Microbiol Immunol 21:201-5.

15.   Soderling E, Alaraisanen L, Scheinin A, Makinen KK. 1987. Effect of xylitol

and sorbitol on polysaccharide production by and adhesive properties of

Streptococcus mutans. Caries Res 21:109-16.

16.   Soderling EM, Hietala-Lenkkeri AM. 2010. Xylitol and erythritol decrease

adherence of polysaccharide-producing oral streptococci. Curr Microbiol

60:25-9.

17.   Trahan L, Mouton C. 1987. Selection for *Streptococcus mutans* with an

altered xylitol transport capacity in chronic xylitol consumers. J Dent Res

66:982-8.

18.   Wen ZT, Browngardt C, Burne RA. 2001. Characterization of two operons

that encode components of fructose-specific enzyme II of the

sugar:phosphotransferase system of *Streptococcus mutans*. FEMS

Microbiol Lett 205:337-42.

19. Dills SS, Seno S. 1983. Regulation of hexitol catabolism in *Streptococcus*

mutans. J Bacteriol 153:861-6.

20. Cvitkovitch DG, Boyd DA, Hamilton IR. 1995. Regulation of sugar

transport via the multiple sugar metabolism operon of *Streptococcus*

mutans by the phosphoenolpyruvate phosphotransferase system. J

Bacteriol 177:5704-6.

21. Kuznetsova E, Proudfoot M, Gonzalez CF, Brown G, Omelchenko MV,

Borozan I, Carmel L, Wolf YI, Mori H, Savchenko AV, Arrowsmith CH,

Koonin EV, Edwards AM, Yakunin AF. 2006. Genome-wide analysis of

substrate specificities of the *Escherichia coli* haloacid dehalogenase-like

phosphatase family. J Biol Chem 281:36149-61.

22. Coleman JE. 1992. Structure and mechanism of alkaline phosphatase.

Annu Rev Biophys Biomol Struct 21:441-83.

23. Dashper SG, Reynolds EC. 1992. pH regulation by *Streptococcus mutans*.

J Dent Res 71:1159-65.

24. Thompson J, Chassy BM. 1983. Intracellular hexose-6-

phosphate:phosphohydrolase from *Streptococcus lactis*: purification,

properties, and function. J Bacteriol 156:70-80.

25. Zeng L, Burne RA. 2009. Transcriptional regulation of the cellobiose

operon of *Streptococcus mutans*. J Bacteriol 191:2153-62.

26.   Zeng L, Burne RA. 2015. Sucrose- and fructose-specific effects on the

890   transcriptome of *Streptococcus mutans*, as determined by RNA

891   sequencing. Appl Environ Microbiol 82:146-56.

892   27.   Abranches J, Chen YY, Burne RA. 2003. Characterization of

893   *Streptococcus mutans* strains deficient in EIIAB^{Man} of the sugar

894   phosphotransferase system. Appl Environ Microbiol 69:4760-9.

895   28.   Abranches J, Nascimento MM, Zeng L, Browngardt CM, Wen ZT, Rivera

896   MF, Burne RA. 2008. CcpA regulates central metabolism and virulence

897   gene expression in *Streptococcus mutans*. J Bacteriol 190:2340-9.

898   29.   Huynh K, Partch CL. 2015. Analysis of protein stability and ligand

899   interactions by thermal shift assay. Curr Protoc Protein Sci 79:28.9.1-14.

900   30.   Liao S, Klein MI, Heim KP, Fan Y, Bitoun JP, Ahn SJ, Burne RA, Koo H,

901   Brady LJ, Wen ZT. 2014. *Streptococcus mutans* extracellular DNA is

902   upregulated during growth in biofilms, actively released via membrane

903   vesicles, and influenced by components of the protein secretion

904   machinery. J Bacteriol 196:2355-66.

905   31.   Cornejo OE, Lefebure T, Pavinski Bitar PD, Lang P, Richards VP,

906   Eilertson K, Do T, Beighton D, Zeng L, Ahn SJ, Burne RA, Siepel A,

907   Bustamante CD, Stanhope MJ. 2013. Evolutionary and population

908   genomics of the cavity causing bacteria *Streptococcus mutans*. Mol Biol

909   Evol 30:881-93.

32.   Pihlanto-Leppala A, Soderling E, Makinen KK. 1990. Expulsion

mechanism of xylitol 5-phosphate in *Streptococcus mutans*. Scand J Dent

Res 98:112-9.

33.   Zeng L, Das S, Burne RA. 2010. Utilization of lactose and galactose by

Streptococcus mutans: transport, toxicity, and carbon catabolite

repression. J Bacteriol 192:2434-44.

34.   Calderone V, Forleo C, Benvenuti M, Thaller MC, Rossolini GM, Mangani

S. 2006. A structure-based proposal for the catalytic mechanism of the

bacterial acid phosphatase AphA belonging to the DDDD superfamily of

phosphohydrolases. J Mol Biol 355:708-721.

35.   Thaller MC, Schippa S, Rossolini GM. 1998. Conserved sequence motifs

among bacterial, eukaryotic, and archaeal phosphatases that define a new

phosphohydrolase superfamily. Protein Sci 7:1647-52.

36.   Horler RS, Vanderpool CK. 2009. Homologs of the small RNA SgrS are

broadly distributed in enteric bacteria but have diverged in size and

sequence. Nucleic Acids Res 37:5465-76.

37.   Kessler JR, Cobe BL, Richards GR. 2017. Stringent response regulators

contribute to recovery from glucose phosphate stress in *Escherichia coli*.

Appl Environ Microbiol 83. e01636-17.

38.   Klein MI, Hwang G, Santos PH, Campanella OH, Koo H. 2015.

Streptococcus mutans-derived extracellular matrix in cariogenic oral

biofilms. Front Cell Infect Microbiol 5:10.

39.   Tanzer JM, Chassy BM, Krichevsky MI. 1971. Sucrose metabolism by

933   *Streptococcus mutans*, SL-I. Biochim Biophys Acta 261:379-87.

934   40.   Zeng L, Burne RA. 2013. Comprehensive mutational analysis of sucrose-

935   metabolizing pathways in *Streptococcus mutans* reveals novel roles for

936   the sucrose phosphotransferase system permease. J Bacteriol 195:833-

937   43.

938   41.   Ahn SJ, Rice KC, Oleas J, Bayles KW, Burne RA. 2010. The

939   *Streptococcus mutans* Cid and Lrg systems modulate virulence traits in

940   response to multiple environmental signals. Microbiology 156:3136-47.

941   42.   Seaton K, Ahn SJ, Sagstetter AM, Burne RA. 2011. A transcriptional

942   regulator and ABC transporters link stress tolerance, (p)ppGpp, and

943   genetic competence in *Streptococcus mutans*. J Bacteriol 193:862-74.

944   43.   Burne RA, Wen ZT, Chen YY, Penders JE. 1999. Regulation of

945   expression of the fructan hydrolase gene of *Streptococcus mutans* GS-5

946   by induction and carbon catabolite repression. J Bacteriol 181:2863-71.

947   44.   Wen ZT, Yates D, Ahn SJ, Burne RA. 2010. Biofilm formation and

948   virulence expression by *Streptococcus mutans* are altered when grown in

949   dual-species model. BMC Microbiol 10:111.

950   45.   Sambrook J, Russell DW. 2001. Molecular cloning: a laboratory manual,

951   3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

952   46.   Biswas I, Jha JK, Fromm N. 2008. Shuttle expression plasmids for genetic

953   studies in *Streptococcus mutans*. Microbiology 154:2275-82.

47.   Zeng L, Burne RA. 2008. Multiple sugar: phosphotransferase system

permeases participate in catabolite modification of gene expression in

Streptococcus mutans. Mol Microbiol 70:197-208.

48.   Veiga-da-Cunha M, Hoyoux A, Van Schaftingen E. 2000. Overexpression

and purification of fructose-1-phosphate kinase from *Escherichia coli*:

application to the assay of fructose 1-phosphate. Protein Expr Purif 19:48-

52.

49.   Zeng L, Burne RA. 2015. NagR differentially regulates the expression of

the *glmS* and *nagAB* genes required for amino sugar metabolism by

Streptococcus mutans. J Bacteriol 197:3533-44.

50.   Loo CY, Corliss DA, Ganeshkumar N. 2000. *Streptococcus gordonii*

biofilm formation: identification of genes that code for biofilm phenotypes.

J Bacteriol 182:1374-82.

51.   Ahn SJ, Lemos JA, Burne RA. 2005. Role of HtrA in growth and

competence of *Streptococcus mutans* UA159. J Bacteriol 187:3028-38.

52.   Ajdic D, McShan WM, McLaughlin RE, Savic G, Chang J, Carson MB,

Primeaux C, Tian R, Kenton S, Jia H, Lin S, Qian Y, Li S, Zhu H, Najar F,

Lai H, White J, Roe BA, Ferretti JJ. 2002. Genome sequence of

Streptococcus mutans UA159, a cariogenic dental pathogen. Proc Natl

Acad Sci U S A 99:14434-9.

 FIGURE LEGENDS

FIGURE 1. Overexpression of *sppA* enhances growth of a *fruK* mutant on

fructose. (A) Relative *sppA* mRNA levels measured in UA159/pIB184,

UA159/pIB508, FruK-13/pIB184, FruK-13/pIB508, *sppR*/pBGE and *sppR*/pBGE-

sppR. Cultures were prepared using BHI medium and harvested at the

exponential phase. Quantitative reverse transcription PCR (qRT-PCR) was

carried out using the total RNA samples, and the relative abundance of *sppA* was

determined using the $\Delta\Delta$Cq method. The error bars denote the standard

deviations. Student's *t*-test was performed to determine the statistical

significance of the data, with asterisks denoting $P < 0.05$. (B) FruK-13/pIB508 (red

squares) and FruK-13/pIB184 (blue diamonds) were cultivated in BHI and then

diluted into TV medium containing 10 mM fructose. The cultures were overlaid

with mineral oil and OD_{600} was measured every 30 min in a Bioscreen C. The

results represent the average of at least three biological replicates.

FIGURE 2. *In vitro* characterization of SppA as a hexose-phosphate:

phosphohydrolase. The majority of the reactions (except panels B&C) were

performed at 37°C in a basic buffer composed of 50 mM PIPES (pH 6.5), 5 mM

$MgCl_2$, and 1 mM $MnCl_2$. Release of phosphate was measured using a Malachite

Green Reagent (MGR), and the results were divided by time and amounts of

enzyme (A, D), or normalized for relative activities (B, C). (A) 10 mM of the

indicated sugar-phosphates were incubated with His-SppA (0.08 µM for F-1-P,

0.67 µM for the rest), with release of phosphate being monitored over a period of

10 min. (B) 0.1 mM of F-1-P was incubated for 5 min with 0.33 μM of His-SppA

and the specified amounts of metal ions. (C) 0.67 mM F-1-P was incubated with

0.04 μM of His-SppA for 130 sec at the indicated pH values (See Methods for

details). (D) Increasing concentrations of F-1-P and F-6-P were each mixed with

His-SppA and phosphate release was monitored over defined periods of time

(See Methods for details).

1005

FIGURE 3. Transcription of the *spp* operon is regulated by SppR, fructose

derivatives and the PTS. The P*sppR::cat* fusion was integrated into the

chromosome of the wild type (A) or various mutants (B) to monitor the expression

of the operon. All strains were first cultivated to mid-exponential phase (OD_{600} =

0.3 to 0.4) in a TV medium with 10 mM glucose, then (A) 25 or 50 mM of the

specified carbohydrates, or (B) 50 mM of glucose or fructose was added, the

cultures were incubated for 1 h, and cells were harvested for CAT assays. Error

bars denote standard deviations. Asterisks indicate statistically significant

difference (by Student's *t*-test, $P < 0.05$) in comparison (A) with glucose at the

same concentrations, or (B) with the wild type (WT) treated with the same

carbohydrates.

1017

FIGURE 4. *In vitro* interactions of SppR with the *spp* promoter region and

fructose phosphates. EMSA was performed using SppR and a biotin-labeled

probe containing the promoter region of *sppR.* SppR (0 to 125 nM) was

incubated first with various sugar phosphates for 10 min, then with 0.2 nM of

DNA probe for 20 min, before the mixture was resolved on a non-denaturing

polyacrylamide gel.

1024

FIGURE 5. Overexpression of SppA enhances resistance to xylitol. Growth

phenotypes of various strains were monitored over time in a Bioscreen C (A, B),

or by measuring the final yield (OD_{600}) after overnight incubation (C). Strains

were cultivated first on BHI to exponential phase, then diluted using TV medium

containing 0.5% glucose in addition to (A, B) 1% xylitol, or (C) the amounts of

xylitol specified on the *x*-axis.

1031

FIGURE 6. Fructose metabolism alters biofilm development by *S. mutans.* BM

containing 2 mM sucrose and 18 mM glucose (BMGS) or 18 mM fructose

(BMFS) was used for biofilm development. (A) Wild type UA159 and its isogenic

mutants *levD* and *fruI* were each incubated for 24 h and the amounts of biofilms

formed were visualized and quantified by crystal violet staining. (B) Biofilms

formed by UA159 were treated with a LIVE/DEAD vitality stain and visualized

under a confocal laser-scanning microscope, with green indicating live cells and

red indicating dead cells. (C) eDNA released from 48-h biofilms formed by strains

UA159/pIB184 (blue) and UA159/pIB508 (orange) were quantified using a DNA

dye (SytoX Green) and a standard curve. The results were normalized using total

CFU from each sample (D). Error bars represent standard deviations. Statistical

analysis was performed using Student's *t* test, with asterisks denoting *P* values: *

<0.05, ** <0.005, and *** <0.0005.

1045

FIGURE 7. HAD family phosphatases contribute to sugar-phosphate tolerance.

(A) *S. mutans* UA159 and its mutants deficient in putative phosphohydrolases.

(B) UA159, a *lacD* mutant, and *lacD* derivatives containing plasmid pIB428 that

overexpresses SMU.428 or the vector pIB184. Bacterial cultures were each

prepared in BHI, then diluted into TV base medium supplemented with (A) 0.5%

glucose and various amounts of xylitol or (B) 0.5% sorbitol and various amounts

of galactose. After 17 to 20 h incubation at 37°C, OD_{600} of the cultures were

recorded. Results are the averages of at least three independent cultures, with

error bars denoting the standard deviations.

Table 1. Bacterial strains and plasmids used in this study.

Strains or plasmids	Relevant genotypes[a]	Source or reference
UA159	Wild type, serotype *c*	(52)
FruK-13	*fruK*M28stop	(3)
MMZ1387	UA159 *sppR*::Km^r	This study
MMZ1388	UA159 *sppR*::Em^r	This study
MMZ1399	UA159 *sppA*::Km^r	This study
MMZ1400	UA159 *sppA*::Em^r	This study
MMZ1416	UA159 P*sppR::cat*	This study
MMZ1518	*levD*::Em^r P*sppR::cat*	This study
MMZ1519	*fruI*::Sp^r P*sppR::cat*	This study
MMZ1520	*manL*::Em^r P*sppR::cat*	This study
MMZ1521	*ccpA*::Sp^r P*sppR::cat*	This study
MMZ1546	*sppR::*Km^r/pBGE	This study
MMZ1547	*sppR*::Km^r/pBGE-*sppR*	This study
MMZ1549	UA159 SMU.428::Km^r	This study
MMZ1551	UA159 SMU.718c::Km^r	This study
MMZ1570	UA159 SMU.488::Km^r	This study
MMZ1573	*sppR*::Em^r P*sppR::cat*	This study
MMZ1574	*sppA*::Em^r P*sppR::cat*	This study
Plasmids		
pIB184	Shuttle expression vector, Em^r	(46)
pBGE	Integration vector, Em^r	(25)
pIB508	*sppA* cloned into pIB184	This study
pIB428	SMU.428 cloned into pIB184	This study

1056   [a] Km^r kanamycin resistance; Sp^r, spectinomycin resistance; Em^r, erythromycin

1057   resistance.

1058 **Table 2.** Measurements of fructose phosphates (nmole/mg protein).

Strains	F-1-P		F-6-P	
	glucose	fructose	glucose	fructose
UA159	0.46 ± 0.13	18.67 ± 3.30	7.94 ± 0.94	8.85 ± 1.08
sppA	0.92 ± 0.39	18.79 ± 1.43	7.93 ± 3.27	9.64 ± 0.57
sppR	1.30 ± 0.32	6.18 ± 0.56***	9.58 ± 3.26	7.94 ± 2.64
FruK-13	4.17 ± 0.47**	13.19 ± 0.81	8.12 ± 0.26	8.21 ± 1.41
fruI	0.79 ± 0.54	7.31 ± 1.80**	N. A.	N. A.
levD	2.74 ± 1.36	21.95 ± 4.57*	N. A.	N. A.

1059

1060 Glucose-grown cells were treated for 30 min with 50 mM glucose or fructose before the

1061 assay. Results (average ± standard deviation) are each derived from three biological

1062 repeats, with each sample measured twice. Asterisks indicate a statistically significant

1063 difference (by one-way ANOVA, alpha = 0.05) in comparison to UA159 under the same

1064 conditions. N. A., not assayed.

A. Substrate specificity

B. Effects of metal ions

$MgCl_2$	2.5, 5, 10 mM
Others	0.5, 1, 2 mM

C. Effects of pH

D. Kinetics

A

B

	5 mM	10 mM	F-1,6-bP
			F-6-P
	5 mM	10 mM	
			F-1-P
5 mM	10 mM				
0	62.5	125	125	125	125	125	125	125 nM	SppR

← Unbound probe

| 0 | 52.8 | 76.1 | 31.9 | 16.7 | 38.8 | 33.5 | 77.2 | 83.9 | % shifted |

A

B