1

Evolutionary forecasting of phenotypic and genetic

outcomes of experimental evolution in *Pseudomonas*

Peter A. Lind [1]

[1] Department of Molecular Biology, Umeå University, Umeå, Sweden.

Correspondence and requests for materials should be addressed to
Peter A. Lind, Dept. Molecular Biology, Umeå University, SE-901 87 Umeå,
Sweden.

email: peter.lind@umu.se

Keywords: *Pseudomonas*, experimental evolution, genetic architecture, c-di-GMP,
evolutionary predictability, evolutionary forecasting

Short title: Forecasting experimental evolution

Impact statement: Conservation of genotype-to-phenotype maps allows successful
prediction of short-term evolution in *P. protegens* Pf-5 and lays the foundation for
evolutionary forecasting in other *Pseudomonas*.

Abstract

Experimental evolution is often highly repeatable, but the underlying causes are

generally unknown, which prevents extension of evolutionary forecasts to related

species. Data on adaptive phenotypes, mutation rates and targets from the

Pseudomonas fluorescens SBW25 Wrinkly Spreader system combined with

mathematical models of the genotype-to-phenotype map allowed evolutionary

forecasts to be made for several related *Pseudomonas* species. Predicted outcomes of

experimental evolution in terms of phenotype, types of mutations, relative rates of

pathways and mutational targets were then tested in *Pseudomonas protegens* Pf-5. As

predicted, most mutations were found in three specific regulatory pathways resulting

in increased production of Pel exopolysaccharide. Mutations were, as predicted,

mainly found to disrupt negative regulation with a smaller number in upstream

promoter regions. Mutated regions in proteins could also be predicted, but most

mutations were not identical to those previously found. This study demonstrates the

potential of short-term evolutionary forecasting in experimental populations.

Introduction

An increasing number of experimental evolution studies, primarily using microbes, have provided insights into many fundamental questions in evolutionary biology including the repeatability of evolutionary processes (Barrick and Lenski 2013; Jerison and Desai 2015; Long, et al. 2015; Orgogozo 2015; Blount, et al. 2018). Given the ability to control environmental conditions as well as population size and the use of a single asexual organism, such studies could provide an ideal test of our ability to predict evolutionary outcomes in simplified model systems. High repeatability on both phenotypic and genetic level have been observed in a large number of experimental evolution studies (Wichman, et al. 1999; Conrad, et al. 2009; Lee and Marx 2012; Tenaillon, et al. 2012; Barrick and Lenski 2013; Ferguson, et al. 2013; Herron and Doebeli 2013; Blank, et al. 2014; McElroy, et al. 2014; Fraebel, et al. 2017; Kram, et al. 2017; Blount, et al. 2018; Knöppel, et al. 2018), but it has become clear that high repeatability alone is not sufficient for testing evolutionary predictability beyond the prediction that under identical conditions the same evolutionary outcome is likely. The difficulties of moving from repeatability to predictability are largely a result of the lack of knowledge of the genotype-phenotype-fitness map (Figure 1).

Figure 1. Prediction of the adaptive outcomes of experimental evolution requires understanding of how genotypes map to phenotypes and fitness. The differing capacity of genes to translate genotypic variation into phenotypic variation and differences in mutation rates can introduce biases in the production of phenotypic

variation. Natural selection, genetic architecture and mutational biases can both

increase and decrease the predictability of evolution depending on if they can be

recognized beforehand and included into evolutionary forecasting models. (**A**)

Fitness. Mutants that increase to high frequencies in the population are all expected to

have increased fitness as drift is negligible at population sizes typically used for

adaptation experiments with microbes. Much effort has been put into characterizing

the distribution of fitness effects of beneficial mutations, but the shapes of the

distribution and magnitudes of the fittest mutations appear to be highly context

dependent. Many different phenotypes are typically adaptive during experimental

evolution, but in most cases they are not known beforehand and their relative fitness

cannot be predicted. Relative fitness is, in most cases, also expected to be highly

dependent on external environment including the frequency of other adaptive mutants,

which means that even small changes to experimental protocols can lead to

differences in outcomes. (**B**) **Phenotype space**. Each of the adaptive phenotypes can

usually be realized by mutations in different positions and in different genes, but

distinct phenotypes are expected to have similar fitness regardless of genetic

foundations, which can simplify predictions. Depending on the genetic architecture

underlying each trait, which is often unknown, adaptive phenotypes are produced at

different rates. (**C**) **Parameter space**. The adaptive phenotypes are caused by changes

in the molecular networks of cells, which are also influenced by the external

environment. If the wiring of a molecular network underpinning an adaptive

phenotype is well understood, parameterization of the system is possible and

predictive models can be formulated. Mutations can cause functional effects on gene

products, but mutations in some genes are more likely to lead to phenotypic variation.

This can for example be due to differences in mutational robustness of the gene

products themselves or their functions in regulatory networks. (**D**) **Genotype space**.

Mutation rates are not uniform across the genome and mutational hot spots can lead to

bias in the number of mutants producing relevant changes in parameter space and

causing new phenotypes that are presented for natural selection to act upon. The

current understanding of the distribution of mutation rates is limited and

computational predictions have not yet been described. Adding information about

well-characterized mutational hot spots, including indels in homonucleotide tracts or

deletion and duplication between sequence homologies, could possibly improve

prediction compared to a null model with uniform rates. Alternatively experimental

data might be incorporated into models of mutation rates to improve predictions

(Lind, et al. 2019).

There are several problems that need to be solved to develop a model system for true

testing of predictive ability. In some cases adaptive mutations are highly strain

specific, so that for example adaptation of different strains to a specific environment

will produce different results. This is sometime due to the long history of sub-

culturing under laboratory conditions combined with rounds of mutagenesis that has

caused, for example, many *E. coli* and *Salmonella* strains to accumulate diverse

mutations, some of which are rapidly compensated by secondary mutations restoring

fitness (Barrick, et al. 2009; Tenaillon, et al. 2012; Knöppel, et al. 2018). Thus, in

many cases conclusions from one strain cannot be extended to another because of

differences in their genotype-to-phenotype maps (Figure 1C).

Another problem for testing predictability is that in many cases it is not possible to

design an experiment where one specific selective pressure is dominant. For example

experiments with intended adaptation to high temperature (Tenaillon, et al. 2012) or

freeze-thaw-growth cycles (Sleight, et al. 2008) result in similar mutations in *uspA*,

which may indicate adaptation to the medium used (Knöppel, et al. 2018) or generally

stressful conditions. Relatively minor changes in environmental conditions can also

results in divergent mutational patterns (Deatherage, et al. 2017). This means that the

range of possible adaptive phenotypes cannot be defined beforehand (Figure 1B) and

that in many cases the phenotypes that solve the intended selective problem are

outcompeted by other phenotypes with increased fitness (Figure 1A).

A highly specific selective pressure can be applied by selection for antitbiotic

resistance and mutation targets are often highly conserved between different strains

and species and between laboratory and natural populations (O'Neill, et al. 2006;

Schenk, et al. 2012; Brandis, et al. 2015; Jahn, et al. 2017; Lukacisinova and

Bollenbach 2017; Sommer, et al. 2017). However resistance phenotypes are typically

explained solely by the molecular phenotype of a single protein and no alternative

pathways to resistance are known resulting in a relatively simple parameter and

genotype space (Figure 1C, Figure 1D). Thus the prediction will be identical for all

species and it cannot provide a test of prediction from general principles. In many

cases mutants isolated after selection for high-level antibiotic resistance also lacks the

complexity that is inherent to many phenotypic traits where the genotype-to-

phenotype map involves a large number of functional interactions and complex

regulation (Figure 1C). This complexity comes to light in that mutations allowing

adaptation to new environments are commonly found in global regulators of gene

expression such as genes involved in the stringent response, DNA binding proteins,

supercoiling and core genes for RNA and protein synthesis (Barrick, et al. 2009;

Conrad, et al. 2009; Kishimoto, et al. 2010; Tenaillon, et al. 2012; Herron and

Doebeli 2013; Sandberg, et al. 2014; LaCroix, et al. 2015; Deatherage, et al. 2017).

The physiological effects of these mutations are diverse, sometimes affecting the

expression of hundreds of genes making the elucidation of the molecular

underpinnings of the adaptive phenotype (Figure 1C, 1D) extremely complex and thus

difficult to use for predictive modeling.

The wrinkly spreader model in *P. fluorescens* SBW25 (hereafter SBW25) is one of

the best-characterized experimental evolution systems and has several properties that

could make it possible to extend knowledge and principles from this species to related

species (Rainey and Travisano 1998; Spiers, et al. 2002; Spiers, et al. 2003; Spiers

and Rainey 2005; Goymer, et al. 2006; Bantinaki, et al. 2007; McDonald, et al. 2009;

Silby, et al. 2009; Ferguson, et al. 2013; Lind, et al. 2015, 2017b; Lind, et al. 2019).

When the wild type SBW25 is placed into a static growth tube the oxygen in the

medium is rapidly consumed by growing bacteria (Figure 2A). However oxygen

levels at the surface are high and mutants that are able to colonize the air-liquid

interface have a major growth advantage and rapidly increase in frequency (Figure

2A). Several phenotypic solutions to air-liquid interface colonization, all involving

increased cell-cell adhesion, have been described and are distinguishable by their

colony morphology on agar plates (Figure 2A, 2B) (Rainey and Travisano 1998;

Ferguson, et al. 2013; Lind, et al. 2017b). The most successful of these is the Wrinkly

Spreader (WS) (Ferguson, et al. 2013; Lind, et al. 2017b) that overproduces a

cellulosic polymer that is the main structural component of the mat at the air-liquid

interface (Spiers, et al. 2002; Spiers, et al. 2003). The WS phenotype is caused by

mutational activation of c-di-GMP production by a diguanylate cyclase (DGC)

(Figure 2C) (Goymer, et al. 2006). While many different DGCs can be activated to

reach the WS phenotype, some are greatly overrepresented due to larger mutational

target sizes leading to a hierarchy of genetic routes to WS (Figure 2D) (McDonald, et

al. 2009; Lind, et al. 2015). The genotype-to-phenotype map to WS has been

characterized in detail (Goymer, et al. 2006; McDonald, et al. 2009; Lind, et al. 2019)

allowing the development of mathematical models of the three main pathways to WS

(Wsp, Aws and Mws) and the prediction of evolutionary outcomes (Figure 2E) (Lind,

et al. 2019).

Figure 2. The *Pseudomonas fluorescens* SBW25 "wrinkly spreader" model system

has several properties that could allow its extension to other species (**A**) The ancestral

strain that has smooth colony morphology on agar plate is inoculated into static

growth tubes and incubated for several days. Depletion of oxygen in the medium

leads to competition for access to the oxygen-replete surface which is colonized by

mutants with enhanced ability for cell-cell adherence and adherence to the wall of the

tube. The most successful of these mutant types is the wrinkly spreader that has a

distinctive colony morphology due to overproduction of exopolymeric substances

(EPSs) of which a cellulosic polymer is the main structural component (Spiers, et al.

2002; Spiers, et al. 2003; Ferguson, et al. 2013; Lind, et al. 2017b) (**B**) The cell wall

of Gram-negative bacteria, such as *Pseudomonas*, presents several components that

may be used to increase cell-to-cell adhesion and attachment to surfaces. In addition

to different EPSs, capsular polysaccharides (CPSs) and lipopolysaccharides (LPS)

might also be used. A variety of different types of proteins including adhesins,

glycoproteins, fimbriae and pili are also potential phenotypic solutions and

incomplete cleavage of the peptidoglycan (PG) layer can lead to cell-chaining thereby

increasing cell-cell adhesion. In *Pseudomonas fluorescens* at least four alternative

distinct phenotypes are selected for at the surface, involving an alternative EPS, CPS,

LPS and PG, but they all have lower fitness than the WS type (Beaumont, et al. 2009;

Ferguson, et al. 2013; Gallie, et al. 2015; Lind, et al. 2017b). The wrinkly spreader

system is of intermediate complexity in that it allows isolation of a phenotypic subset

of adaptive mutants, those that are selected at the air-liquid interface and produce a

difference in colony morphology, rather than all mutants that increase fitness.

However there is also a large diversity of different genetic and phenotypic solutions to

the dominant adaptive challenge, which makes it of greater complexity than systems

based on single genes, which is often the case for experimental systems using strong

selection for antibiotic resistance. (**C**) The mutational causes of the high fitness WS

phenotype in wild type populations are found in either of three loci, *wspABCDEFR*,

awsXRO and *mwsR,* which all encode diguanylate cyclases that produce the second

messenger c-di-GMP (McDonald, et al. 2009) that is a conserved signal for EPS

production and biofilm formation in many bacterial species. In SBW25 the primary

EPS used is a cellulosic polymer. (**D**) While these three pathways account for >98%

of WS mutants in the wild type, 13 additional rare pathways were found when the

common ones are genetically deleted (Lind, et al. 2015). The large differences in

mutation rates to WS for the different pathways are explained mainly by their

different capacities to translate genotypic variation into phenotypic variation, i.e.

mutational target size, with the three common pathways being subject to negative

regulation, which when disrupted results in overproduction of c-di-GMP and the WS

phenotype (Lind, et al. 2015). The alternative phenotypic solutions are also caused by

inactivating mutations occurring at high rates (Ferguson, et al. 2013; Lind, et al.

2017b). The pathways to WS of intermediate frequency are activated by mutations to

promoter regions, including promoter captures and the most rare are those that require

specific activating mutations in the DGC or double mutations in two negative

regulators. The genetic underpinnings of several adaptive phenotypes have been

elucidated providing a mechanistic understanding of the effects of mutations and why

they are adaptive. (**E**) The three main pathways (Wsp, Aws, Mws) to WS are

particularly well-understood allowing formulation of mathematical models of the

molecular networks and prediction of the relative rates of use of the different

pathways and genes (Lind, et al. 2019).

This study makes initial forecasts of phenotypic and genetic evolutionary outcomes

after static experimental evolution for six *Pseudomonas* species based mainly on their

genome sequence and data from SBW25 (McDonald, et al. 2009; McDonald, et al.

2011; Ferguson, et al. 2013; Lind, et al. 2015, 2017b). Predictions of evolutionary

outcomes were then experimentally tested for the closely related species, *P. protegens*

Pf-5 (hereafter Pf-5) with a highly conserved genetic repertoire of DGCs but that

lacks the main structural component used by WS types in SBW25. Results show that

phenotypes, order of pathways used and types of mutations can be predicted and that

forecasts are robust to changes in environmental conditions.

Results

236 Six *Pseudomonas* species (Figure 3 legend) were chosen based on phylogenetic

237 diversity and their complement of DGCs and EPSs for a first round of predictions.

238 These species encode from none to all three of the main DGCs used in SBW25 and

239 only three species contain genes related to cellulose biosynthesis, the main EPS used

240 in SBW25 (Figure 3). Full details are available in Figure 3 - source data.

Fig 3. Diversity of DGCs and biofilm-related genes for seven *Pseudomonas* species (*P. fluorescens* SBW25, P. *protegens* Pf-5, *P. putida* KT2440, *P. syringae* pv. tomato DC3000, *P. savastanoi* pv. phaseolicola 1448A, *P. aeruginosa* PAO1, *P. stutzeri* ATCC 17588) **(A)** The seven species encode 251 putative DGCs, divided into 87 different homolog groups of which 8 are present in all genomes. WS mutations in SBW25 have been found affecting 13 of these DGCs (marked with *) with an additional nine that have been detected only in combinations with other mutations. SBW25 and Pf-5 share 33 DGCs with 6 unique for each species. It should be noted that not all DGCs are likely to be catalytically active. **(B)** Diversity of biofilm-related genes including putative EPSs, LPS modification, cell chaining, adhesins and known regulators. In SBW25 cellulose-based mats are most successful (encoded by the *wss* operon). A secondary exopolysaccharide (PGA), encoded by *pgaABCD*, can also be used to form a stable mat (Lind, et al. 2017b). Fuzzy spreaders (FS) forms rafts that collapse after becoming too large and the mutational cause is inactivation of *fuzY*, which results in a defect in lipopolysaccharide (LPS) modification (Ferguson, et al. 2013). Cell-chaining (CC) types have loss-of-function mutations in *nlpD* causing a defect in cell division, which leads to segmented chains of cells that can form a weak mat at the surface (Lind, et al. 2017b). The genomes of the *P. aeruginosa* strains PA7, UCBPP-PA14 and LESB58 were also included in the analysis and results were in most cases identical to PAO1 (not shown in Figure 3) with the exception of an absence of homologues for EPS genes *pelA-D* and the DGCs PA2771 in PA14 and PA3343 in PA7.

Ecotype predictions

Given the range of ways that cells can achieve increased adherence and surface colonization by use of different EPSs, LPS modification and cell chaining as demonstrated by the studies with SBW25 (Spiers, et al. 2002; Ferguson, et al. 2013; Lind, et al. 2017b) all species are be expected to colonize the air-liquid interface if access to oxygen is limiting for growth. This could be achieved simply by changes in gene expression in the wild type, but for an experimental evolution study a mutational solution is sought and environmental conditions are chosen so that the wild type strain does not colonize the air-liquid interface. However changing environment presents a further challenge because, as discussed above, it often leads to a different spectrum of adaptive mutations. Thus a foundational requirement for an extended experimental

evolution system to be successful for different species is that the evolutionary

solutions are robust to differences in environmental conditions.

Phenotype predictions

Several different phenotypic solutions can be used to colonize the air-liquid interface

in SBW25 including at least two different EPSs (cellulose (WS) and PGA (PWS),

LPS modification (fuzzy spreaders FS) and cell chaining (CC) (Spiers, et al. 2002;

Ferguson, et al. 2013; Lind, et al. 2017b). However wrinkly spreaders that form

cellulose-based mats are superior and rapidly outcompete all other types (Ferguson, et

al. 2013; Lind, et al. 2017b). Based on the limited data available it is predicted that

cellulose-based biofilms are superior in other species as well and that they will be the

primary structural solution when available as for *P. syringae*, *P. putida* and *P.*

stutzeri. For the three species lacking genes for cellulose biosynthesis, other EPSs are

predicted to be used. Based on studies of *P. aeruginosa* the primary EPS required for

pellicle formation at the air-liquid interface in this species is Pel, encoded by the

pelABCDEFG operon, which is also present in the Pf-5 genome and is predicted to be

the primary phenotypic solution for these species. The genome of *P. savastanoi* lacks

genes for biosynthesis of cellulose and Pel as well as other EPSs that are known to be

able to support mat-formation, such as PGA and there is not sufficient data at this

point to make a prediction of which one is likely to be the primary phenotypic

solution.

Overexpression of EPSs used for mat-formation at the air-liquid interface is in

SBW25 and *P. aeruginosa* linked to mutations increasing c-di-GMP production rather

than mutations in the promoters of, or genes in, the EPS operons themselves. This can

be explained by the role of post-translation regulation by c-di-GMP in the production

of cellulose, Pel, PGA and alginate (Lee, et al. 2007; Römling, et al. 2013; Steiner, et

al. 2013; Morgan, et al. 2014; Liang 2015; Whitney, et al. 2015). In these cases

transcriptional up-regulation alone is not sufficient to cause overproduction because

of lack of a c-di-GMP signal. Possibly there is also an additional benefit to using

activation of the c-di-GMP network in that it reduces motility, which is not needed

when established at the air-liquid interface and which consumes large amount of

energy to sustain and thus is likely to be selected against (Koskiniemi, et al. 2012; Lee

and Marx 2012).

Prediction of types of mutations

Disabling mutations are expected to be more common than enabling mutations and
therefore the prediction is that most mutations will be in genes where loss-of-function
mutations produce an adaptive phenotype (Fig 2D) (Lind, et al. 2015). This is the case
for the large majority of mutations in SBW25 including those activating main DGCs
WspR, AwsR and MwsR, which are all under negative regulation, as well as
disruption of the genes underpinning the FS phenotype (*fuzY*, PFLU0478) and CC
phenotype (*nlpD*, PFLU1301) (McDonald, et al. 2009; Ferguson, et al. 2013; Lind, et
al. 2017b). Next in the hierarchy of mutations are promoter mutations, increasing
transcription, and promoter capture events (Lind, et al. 2015). Less common are
intragenic activating mutations that enable a particular function by for example
increase in catalytic activity or strengthening of interactions to another molecule or
another domain of the same protein (Lind, et al. 2015). Gene duplications occur at a
high rate and clearly have the ability to increase gene expression of DGCs, but they
have not yet been found to cause WS in SBW25, possibly because a two-fold increase
in gene expression is insufficient.

Prediction of pathways used

There are at least 16 different pathways to the WS phenotype in SBW25 with similar
fitness, but they are used at frequencies that vary over several orders of magnitude
based on the differing capacity to translate phenotypic variation into phenotypic
variation (Figure 2D) (Lind, et al. 2015). Mutations in three pathways, Wsp, Aws, and
Mws account for >98% of WS mutations and based on a detailed understanding of the
molecular functions of the genes involved of each pathways mathematical models
predicting at which relative rates the pathways should be used were constructed
(Figure 2E) (Lind, et al. 2019). The prediction results varies depending on the rates of
disabling and enabling mutations, but if it is assumed that disabling mutations are an
order of magnitude more common than enabling mutations the models predict that
Wsp will account for about 54%, Aws 30% and Mws 16% of the WS mutations
(Figure 2E) (Lind, et al. 2019).

When the three common pathways are deleted WS types evolve mainly by mutations
in PFLU0085, which contains an intragenic negative regulator region (Lind, et al.

2015), and this is expected to be the fourth most common pathways when present.

Less common promoter mutations will also appear at rates at least a magnitude lower,

but which DGCs that will be transcriptionally activated cannot be easily predicted

except for assuming it will be homologs of the ones used in SBW25. These DGCs

must be catalytically active and also be localized to the membrane (Farr, et al. 2017).

Possibly the subset of DGCs that are primarily activated by mutations to their

promoters is mainly determined by mutation rate and a higher mutation rate might be

caused by higher transcription and also influenced by gene direction (Sankar, et al.

2016). Most promoter capture deletion events are less than 5 kilobases (Lind, et al.

2015) in size and the lack of an alternative promoter relatively close upstream to the

DGC is likely to rule out these DGCs. The DGCs that can be activated by intragenic

activating mutations cannot now be predicted beyond the simple prediction that these

are the same genes as in SBW25 (Lind, et al. 2015).

Prediction of mutated genes

In addition to predicting the relative rates of the three main pathways, the previously

described mathematical model can also predict which proteins are likely to be

mutated (Lind, et al. 2019). High rates of WS mutations are predicted for WspF,

WspA, WspE, AwsX and AwsR and MwsR (Figure 2E). A significantly lower rate of

enabling mutations is also predicted to occur in WspC, WspR and AwsO (Figure 2E).

Despite the simplicity of null model it closely predicted the mutational targets in

SBW25 with equal rates for WspF, WspA and WspE and rare mutations in WspC and

WspR, suggesting that it is a useful null model also for other species (Lind, et al.

2019).

Prediction of specific mutational targets and effects of mutations

The level of parallelism at the nucleotide level between species is expected to be

dependent both on the number of possible mutations to WS and the degree of

functional conservation of the proteins involved that define the genotype-to-

phenotype map. Mutational hot spots are also expected to contribute to parallelism

when they are conserved, but reduce parallelism when they are not. Based on previous

analysis of patterns of mutations in SBW25 (McDonald, et al. 2009; McDonald, et al.

2011; Lind, et al. 2019) and homology modeling of protein structure using Phyre2

 (Kelley, et al. 2015) regions expected to be mutated were predicted and the likely

molecular consequences of different mutations suggested (Figure 4).

Figure 4. Predicted mutational targets and proposed molecular effects. Black

represents any inactivating mutation including frame shifts, blue represents in frame

inactivating mutations, green represents amino acid substitutions. Numbers in

brackets refers to amino acid residue numbers in SBW25 (**A**) WspA – amino acid

substitutions are expected at the tip of the stalk and in-frame deletion of methylation

sites (**B**) WspF – any inactivating mutation is predicted, amino acid substitutions are

predicted only in areas where they disrupt intermolecular interactions (**C**) WspE –

amino acid substitutions are predicted near the phosphorylation site (**D**) WspR – small

in frame deletion and amino acid substitutions in the linker is predicted to cause

constitutive activation (**E**) AwsO – amino acid substitutions disrupting AwsO

dimerization is predicted to lead to increased binding to AwsX without the presence

of an activating signal (**F**) AwsR - amino acid substitutions in the periplasmic region

or transmembrane helix that disrupt the interaction with AwsX or to the HAMP linker

is predicted (**G**) AwsX – any inactivating mutation that keep the reading frame intact

and do not interfere with expression of downstream AwsR is predicted (**H**) MwsR –

mutations are predicted in the interface between the DGC and phosphodiesterase

domains and in the most C-terminal of the PAS domains resulting in constitutive

activation.

Prediction of fitness effects of WS mutations

While conservation of relative fitness of different phenotypic variants might be

expected it is less clear if the relative fitness of different DCG pathways and

mutations will be conserved between species. Despite this difficulty there might be a

way forward to predict the relative fitness of a large range of mutations with limited

experimental data. The distribution of fitness effects of new mutations have been

found to be bimodal for a large number of genes with different functions with one

mode close to neutrality and one corresponding to a complete loss of a particular

molecular function (Jacquier, et al. 2013; Jimenez, et al. 2013; Firnberg, et al. 2014;

Lind, et al. 2017a; Lundin, et al. 2017). Given that mutations that allow colonization

of the air-liquid interface have large phenotypic effects and are believed to also have

large effects on molecular function, often a complete disruption of an interaction,

adaptive mutations in the same region of a protein are likely to have similar fitness

effects. Thus, an approximation of the distribution of fitness effects could be possible

with relative few mutations for each gene. This is supported by the relatively small

number of WS mutants in SBW25 that have been characterized with sensitive fitness

assays and where mutations in the same gene typically have similar fitness effects

(Lind, et al. 2015; Lind, et al. 2019). If this assumption is true the distribution of

beneficial fitness effects is not continuous and the most advantageous mutations are

not predicted to be equally distributed between pathways or genes. Thus the

prediction would be that mutants isolated after experimental evolution were

concentrated to certain genes even if the mutational rate is similar so that although the

prediction from the null model is equal number of mutations for WspA, WspE and

WspF such distribution is unlikely to be found. While the mutation rates to WS for

the three genes are similar in SBW25, WspA mutants are rarely found after

experimental evolution due to their lower fitness (McDonald, et al. 2009; Lind, et al.

2019). There is however no *a priori* reason to expect that the relative fitness of

mutations in different genes or pathways will be conserved between species.

Inactivating mutations in *fuzY* and *nlpD* producing the alternative adaptive

phenotypes based on LPS modification or cell chaining were also found to have

similar fitness (Ferguson, et al. 2013; Farr 2015). Possibly there are other genes that

can be mutated with similar phenotypes, but that those mutants have lower fitness and

are outcompeted in SBW25. If relative fitness is not conserved between species this

could lead to high convergence on the phenotypic level but with completely different
genetic bases.
436

Experimental test of forecasts in *Pseudomonas protegens* Pf-5
In order to test the predictions presented above, *P. protegens* Pf-5 was used for a
parallel experimental evolution study under static conditions (Figure 1A). Its genome
encodes homologs of all except one of the DGCs used in SBW25 including all the
three common pathways to WS (Wsp, Aws, Mws). Thus the genetic predictions in
terms of types of mutations and mutated genes are nearly identical to SBW25 and the
mathematical null models can be directly applied. However Pf-5 lacks genes for
biosynthesis of cellulose meaning that if c-di-GMP overproduction is the main
pathway used, as predicted, an alternative EPS component must be utilized. The
experimental conditions were modified to test the robustness of predictions to changes
in media composition, temperature and cell wall material compared to those used in
the original SBW25 system (Materials and Methods). After experimental evolution
for five days, dilutions were spread on agar plates and then screened for colonies with
divergent colony morphology, characteristic of many phenotypes that colonize the air-
liquid interface.
452

Genetic pathways
In total 43 independent mutants were isolated and the causal mutations were
identified (Figure 5, Figure 5 – supporting data). As predicted by the null model the
majority (40/43) of mutations were associated with the Wsp, Aws, and Mws pathways
that are subject to negative regulation (Figure 1D). In addition the prediction that
promoter mutations would be the second most common type of mutation was
successful with two mutations found upstream of the *aws* operon, which were
predicted to disrupt the terminator of a high expression ribosomal RNA operon
representing a promoter capture event. Promoter mutations were also found upstream
of PFL_3078, which is the first gene of a putative EPS locus (PFL_3078-3093) that
has not previously been described and that is only present in closely related strains.
The operon encodes genes typical of exopolysaccharide biosynthetic operons making
it highly likely it encodes the main structural component used by these mutants.

Figure 5. A. 43 independent mutants of wild type *Pseudomonas protegens* Pf-5 were isolated after experimental evolution based on their divergent colony morphology and mutations were identified in four operons. Numbers in brackets are the number of independent mutants found. Details are available in Figure 5 – source data. **B.** Experimental evolution with a Δ*wsp* Δ*aws* Δ*mws* triple deletion mutant resulted in WS types with mutations in the DGC PFL_0087.

The mathematical null model (Figure 2E) successfully predicted that of the three common pathways to WS, Wsp would be the most common one (16 mutants) followed by Aws (14) and then Mws (10). Mutations were predominately found in the negative regulators WspF (15 mutants) or AwsX (9), but also in interacting proteins WspE (1) and AwsR (3). Given that the mutational target size is estimated to be smaller for the interacting proteins (Figure 4) this is not surprising. No mutations were found in WspA despite a predicted high rate.

Mutations were predominantly found in predicted regions (Figure 4) for WspF, WspE, AwsX, AwsR and MwsR, but in most cases they were not identical to those in SBW25 (Figure 5 – supporting data). A mutational hot spot was apparent in WspF with 12 out of 15 mutations being identical V271G missense mutations. The previously described mutational hot spots in SBW25 in the *awsX* and *mwsR* genes (Lind, et al. 2019) appeared absent, demonstrating how mutation rate differences can

488 skew evolutionary outcomes even for closely related species with similar genetic
489 architecture.
490

To determine if there were also rare pathways to the WS phenotype the entire *wsp*,
aws, and *mws* operons were deleted and experimental evolution repeated as was
previously done for SBW25 (Lind, et al. 2015). Mutations in the DGC PFL_0087
accounted for six out of the seven WS types found (Figure 5B). This was also the
dominant pathway in the SBW25 Δ*wsp* Δ*aws* Δ*mws* strain where mutations in the
corresponding region of PFLU0085 were responsible for 47% of WS mutants. Thus
the fourth most common pathway is also the same for both species.
498

Mutations in WspA are predicted to be one of the major mutational routes to WS
based on the mathematical model (Lind, et al. 2019), but no mutations were found
either in this study or in SBW25 (McDonald, et al. 2009). However, when the
mutational spectrum of WS mutants was determined in the absence of selection for
growth at the air-liquid interface, *wspA* mutant occurred at rates similar to those of
WspE and WspF, as predicted by the model, and their low frequency after
experimental evolution could be explained by their lower fitness. To investigate if a
WspA mutation could cause WS in *P.protegens* a common deletion found in SBW25
(WspA T293-E299) was introduced and found to cause a WS phenotype and it is
included in the experiments described below.
509

Phenotypic characterization
In total 60 wells were inoculated for the wild type and subjected to experimental
evolution for five days after which air-liquid interface colonization was observed for
the majority of the wells. Mutants with clearly visible changes in colony morphology
were isolated from 43 wells. The experiment was repeated for the Δ*wsp* Δ*aws* Δ*mws*
triple deletion mutant and WS types were detected in 7 wells. Typically a single type
of divergent colonies was observed and one colony for each well was selected for
further characterization at random based on a pre-determined position on the agar
plate. Representative mutations were reconstructed using an allelic exchange protocol
to determine that the mutations are the sole cause of the air-liquid interface
colonization and colony phenotypes and to exclude the influence of secondary
mutations (Figure 6A) before further characterization.

Figure 6. Phenotypic characterization of reconstructed mutants. (**A**) Motility, colony
morphology and air-liquid interface colonization (**B**) Motility assay. As expected if
the c-di-GMP network is activated motility was reduced for most mutants with the
exception of PFL_3078 that is not expected to have increased c-di-GMP levels and
the AwsXRO promoter capture.

The lack of cellulose biosynthetic genes also shows that these ecotypes can evolve by
different phenotypes than in SBW25. Two clearly different phenotypes were observed
with one very similar to the original WS types in SBW25 with a clear motility defect
and mutations in the Wsp, Aws, Mws and PFL_0087 pathways (Figure 6A, 6B). The
other type was less wrinkly, had similar motility as the wild type and promoter
mutations upstream of the PFL_3078-3093 operon (Figure 6A, 6B).

Fitness of adaptive mutants

Two types of fitness assays were performed, similarly as previously described (Lind,
et al. 2015), to measure differences in fitness between the different WS mutants and
the alternative phenotypic solution with the mutation upstream of PFL_3078. The first
assay measures "invasion fitness" where the mutant is allowed to invade a wild type
population from an initial frequency of 1%. This confirms that the mutations are
adaptive and that mutants can colonize the air-liquid interface. The invasion assays
showed that all reconstructed mutants could rapidly invade an ancestral wild type
population (Figure 7A, Figure 7 – source data). Although there were significant
differences between selection coefficients of the mutants (one-way ANOVA $p <$
0.0001), no mutant was significantly different from the most common mutant (WspF
V271G, two-tailed t-test $p > 0.01$).

The second fitness assay measures "competition fitness" and here each mutant is
instead mixed 1:1 with the most common WS type (WspF V271G) at the start of the
competition. The competition assay showed that the ancestral wild type was rapidly
outcompeted by the mutants also at a 1:1 initial ratio (Figure 7B). There was
significant variation in fitness between the WS mutants (one-way ANOVA $p <$
0.0001) and the AwsX had significantly lower selection coefficient (two-tailed t-test p
= 0.009) compared to the reference WspF V271G and one of the MwsR mutants
(E1081K) had significantly higher selection coefficient (two-tailed t-test $p = 0.001$).
The alternative phenotypic solution used by the PFL_3078 promoter mutant resulted
in the lowest fitness (s = -0.1, two-tailed t-test $p < 0.0001$) meaning that it is expected
to be rapidly outcompeted by the WS mutants (Figure 7B). The PFL_0087 mutants
that were only found when the common pathways were deleted had lower fitness
(two-tailed t-tests $p = 0.0005$, $p = 0.001$) and this was also true for the *wspA* mutant
(two-tailed t-test $p = 0.002$), which could explain why these were not found in the
wild type population after experimental evolution.

Figure 7. Fitness of reconstructed *P. protegens* Pf-5 WS mutants was measured in pairwise competitions. **(A)** Invasion fitness was measured relative a dominant ancestral wild type strain with a 1:100 initial ratio to show that mutations were adaptive and that they can increase from rare to colonize the air-liquid interface. Six independent competitions were performed for each pair. **(B)** Competition fitness was measured relative the most common WS mutant (WspF V271G) in a 1:1 initial ratio to compare the fitness of different WS mutants and the alternative phenotypic solution. Four independent competitions were performed for each pair.

Identification of EPS used for air-liquid colonization

SBW25 WS mutants use cellulose as the main structural component, but even though there is high parallelism at the genetic level for Pf-5 WS mutants this cannot be the

case at the phenotypic level as its genome does not encode genes for cellulose biosynthesis. Given that production of Pel exopolysaccharide has been shown to be induced by mutations in *wspF* in *P. aeruginosa* (Hickman, et al. 2005) and that Pel in this species is required for pellicle formation under static growth (Friedman and Kolter 2004) this was predicted to be the main structural component used by Pf-5. To test this prediction the *pel* operon (PFL_2972-PFL_2978) was deleted from Pf-5 and combined with previously characterized WS mutations and fitness was measured. Both invasion fitness (Figure 8A) and competition fitness (Figure 8B) was significantly lower (two-tailed t-tests $p < 0.01$) compared to isogenic strains with an intact *pel* operon (Figure 7A, 7B) except invasion fitness for the AwsX mutant (two-tailed t-tests $p = 0.08$, one outlier). This suggests that Pel polysaccharide serves as an important structural component for colonizing the air-liquid interface and that its production is activated by mutations leading to increased c-di-GMP levels. Although deletion of *pel* in WS mutants resulted in less wrinkly colony morphology it did not result in a smooth ancestral type. Neither did deletion of *pel* abolish the ability to colonize the air liquid interface (Figure 8C) or the ability to invade wild type populations (Figure 8A). This suggests that production of an additional EPS component is induced by increased c-di-GMP levels caused by mutations in Wsp, Aws, Mws and PFL_0087 at least in the absence of *pel*. When the cellulose biosynthetic operon was deleted from SBW25, typical WS mutations in *wsp*, *aws*, and *mws* resulted in air-liquid colonization by use of the alternative structural component encoded by *pgaABCD* and subsequent deletion of the *pgaABCD* operon in these mutants resulted in a wild type colony morphology (Lind, et al. 2017b). Deletion of the *pgaABCD* operon (PFL_0161-PFL_0164) in Pf-5 strains with deletion of the *pel* operon combined with WS mutations in either *wspF*, *awsX* and *mwsR* did not result in a change in colony morphology or loss of ability to colonize the air-liquid interface, which suggests that PGA is not the secondary structural component used or that yet another EPS is also produced in response to increased c-di-GMP levels. As expected if the motility defect observed for WS mutants are primarily caused by high c-di-GMP levels rather than high production of Pel, the motility was also reduced for Wsp, Aws and Mws mutants with the *pel* operon deleted (Figure 8D).

Fig 8. Contribution of *pel* to WS phenotype and fitness (**A**). Deletion of *pel* in WS

mutants reduces invasion fitness. Fitness of reconstructed *P. protegens* Pf-5 WS

mutants without the *pel* operon was measured in pairwise competitions. Invasion

fitness was measured relative a dominant ancestral wild type strain with a 1:100 initial

ratio. Six independent competitions were performed for each pair. (**B**) Deletion of *pel*

in WS mutants reduces competition fitness. Competition fitness was measured

relative the most common WS mutant (WspF V271G) in a 1:1 initial ratio. Four

independent competitions were performed for each pair. (**C**) Deletion of *pel* in WS

mutants did not result in ancestral smooth colony morphology or loss of ability to

colonize the air-liquid interface suggesting a secondary EPS component is produced.

(**D**). Deletion of *pel* did not restore motility showing that Pel overproduction is not the

cause of the motility defect in WS mutants.

Discussion

The extension of the *P. fluorescens* SBW25 experimental evolution system to related species shows promise for true testing of evolutionary forecasting method and models. While there is a diversity of DGCs and EPSs between species leading to differences in forecasts, the conserved role of c-di-GMP and limited number of phenotypes allow the use of previous data to improve predictions and makes the experimental system robust to changes in environmental conditions. The experimental test of initial forecasts for *P. protegens* Pf-5 presented here provides support for the ability to predict some aspects of both genetic and phenotypic evolution while recognizing that the probability of specific mutations cannot in most cases be predicted.

That experimental populations of *Pseudomonas* will colonize the air-liquid interface when incubated under static condition is a prerequisite of extending the model. Given that a range of phenotypic solutions is predicted to be available for all species the evolution of such mutants for *P. protegens* is not surprising. The specific environmental conditions used for experimental evolution often have a major impact on evolutionary outcomes and is also likely to influence relative fitness and possibly mutational biases also in the WS system. However despite major changes in growth medium, temperature and material and physical dimensions of the growth vessel, predictions on both the genetic and phenotypic levels proved successful demonstrating robustness to environmental change and the establishment of a dominant selective pressure, i.e. access to oxygen solved by air-liquid interface colonization.

Phenotypic predictions of the structural basis supporting air-liquid colonization is challenging given the limited previous experimental data. For SBW25 cellulose-based solutions are superior in fitness, but for Pf-5 this solution in not available. The prediction that overproduction of structural exopolysaccharides, rather than fuzzy, cell-chaining or mucoid types, would be the primary solution was successful. One of the two phenotypes found here used the Pel EPS, which could be predicted based on its role in *P. aeruginosa*. However it appears to use a secondary EPS as well, that remains to be identified, given that mutants lacking Pel but with activated DGCs still

colonize the air-liquid interface and have a distinct colony morphology. The second

phenotype used another EPS, encoded by PFL_3078-3093, which had not previously

be described and given that several EPS loci are usually encoded in *Pseudomonas*

genomes its use could not be predicted. However, repeating experimental evolution

using other *Pseudomonas* species is likely to provide more information about which

EPSs can be used to colonize the air-liquid interface and their relative fitness to allow

improved phenotypic predictions. Deletion of the *pel* operon, the unidentified

secondary EPS and PFL_3078-3093 and subsequent experimental evolution could

reveal less fit phenotypic solutions that are expected to exist including fuzzy types

caused by defects in LPS modification, cell-chaining types with defects in cell

division, adhesive proteins or mucoid types using alginate or levan, two EPSs with

lower structural stability.

The general prediction of types of mutations, as described in the hierarchy in Figure

2D, was also successful although the relatively few mutants identified here did not

allow for detection of rare activating mutations or double inactivating mutations. The

majority of mutations were loss-of-function mutations in negative regulators or

interacting proteins followed by less common promoter mutations and promoter

captures. In contrast to SBW25, where all promoter mutations resulted in up-

regulation of DGCs, the mutation upstream of PFL3078-3093 demonstrates the

possibility of direct transcriptional activation of EPS components that are not under

post-translational control of c-di-GMP. Two identical mutations were found over 9 kb

upstream of the *aws* operon, in between a ribosomal RNA operon and the *recCBD*

operon, which encodes key genes for recombination. The molecular effects of these

mutations have not been further investigated, but the resulting WS phenotype is

dependent of the presence of the *aws* operon, deletion of which reversed the

phenotype. This is consistent with an up-regulation of c-di-GMP by AwsR

presumably caused by increased transcription. The mutation is located in the

predicted terminator of the ribosomal RNA operon and increased transcriptional read-

through could put the *aws* operon under control of a very strong *rrn* promoter that is

most highly transcribed during exponential growth. This could explain the relatively

mild colony morphology phenotype as well as high motility of this WS mutant

(Figure 6).

A mathematical null model that incorporates information about the Wsp, Aws and

Mws molecular networks (Null model IV in (Lind, et al. 2019)) successfully predicted

that Wsp would be the most commonly used pathway followed by Aws, and Mws the

most rare. However, the number of mutants isolated here is rather small and the high

frequency of Wsp mutants seems mainly to be caused by a mutational hot spot in

wspF. Still the prediction that the three pathways together would contribute the large

majority of adaptive mutations (40 out of 43) is not trivial given that in SBW25 at

least 13 additional pathways are available to the high fitness WS phenotype (Lind, et

al. 2015). It is also worth noting that direct use of mutation rate data from SBW25

(Lind, et al. 2019) would result in poorer predictions than the mathematical null

model due to a strong mutational hot spot in *aws*X in that species. In addition the

fourth most common pathway to WS, PFL_0087, could be predicted based on data

from SBW25.

For the multi-protein pathways Wsp and Aws, the mathematical model (Lind, et al.

2019) predicted (Figure 2E) that mutations would primarily be found in WspA,

WspE, WspF, AwsX and AwsR. Mutations were detected in all these except in WspA

and the majority was found in the negative regulators WspF and AwsX. WspA

mutations were not found in the original study in SBW25 either (McDonald, et al.

2009), but this was shown to be due to lower fitness relative WspF and WspE mutants

rather than a lower mutation rate to WS (Lind, et al. 2019). This is also a likely

explanation for the absence of WspA mutants here as well (Figure 7B), but it is not

clear if this fitness difference would be conserved in other species or if sometimes

WspA mutants are more fit. Thus the null model prediction of equal rates for WspA,

WspF and WspE is not changed for future experimental tests. Direct comparison

between competitive fitness of mutants in SBW25 and Pf-5 is not possible as it was

measured under different experimental conditions, against different reference mutants

and in most cases the mutations are not identical. However it is interesting to note that

for both species high fitness WS types have mutations in the same proteins (WspF,

WspE, MwsR) and low fitness WS types also appear in the same proteins (WspA,

AwsX, AwsO, PFLU0085/PFL_0087).

The molecular effects of the mutations found here are unknown, but knowledge from

SBW25 and *P. aeruginosa* and their positions in protein structure allowed some

predictions to be made. Inactivating mutations in the negative regulator WspF were

predicted to be either indels or missense mutations in four specific regions. Mutations

were found in two of the predicted regions, one in the vicinity to the methylesterase

active site where mutations are predicted to cause large disruptions in protein

structure and the other one directly disrupting the phosphorylation active site in the

signal receiver domain. No mutations were found in the surface exposed regions

hypothesized to be involved in interactions with WspA and WspE, which could be

due to differences in function between SBW25 and Pf-5 or simply that they appear at

lower frequency and would be detected if additional mutations were isolated. The sole

mutation in WspE is, as predicted, located in the direct vicinity of the phosphorylation

active site. Mutations in AwsX were amino acid substitutions throughout the gene as

well as in frame deletions inactivating the gene as predicted. Mutations in AwsR and

MwsR were also found in predicted regions, but no mutations were found in the small

periplasmic region of AwsR, which is the most commonly targeted region in SBW25.

Known mutational hot spots in *awsX*, *awsR* and *mwsR* in SBW25 (Lind, et al. 2019)

were not conserved in Pf-5 resulting in divergent spectra of mutations, while mutated

regions and predicted functional effects remain conserved between the two species.

Little is known about the molecular function of the putative DGC encoded by

PFL_0087/PFLU0085, but it is clear that a multitude of amino acid substitutions,

deletions and insertions in a more than 40 amino acids long region can lead to WS

(Lind, et al. 2015). Thus it functions as a small intragenic negative regulator region

that might be involved in oligomerization and loss of this interaction results in

constitutive activation of c-di-GMP production.

The diversity of phenotypic solutions observed after experimental evolution is

dependent on fitness differences between the phenotypes, but also on the rate of

which phenotypes are introduced by mutations, which is dependent on the genetic

architecture underlying the trait as well as mutational biases. The Pf-5 strain has at

least three DGC pathways (Wsp, Aws and Mws) that are subject to negative

regulation leading to prediction of a high rate of WS mutants, which are then expected

to outcompete other phenotypic solutions. If instead only one of these pathways were

present, a larger diversity of phenotypes would be expected to be observed with

relative fitness becoming less important as the first mutant that gains a foothold at the

air-liquid interface will have a large advantage and priority effects, i.e. being first,

will increasingly determine which adaptive mutants are observed.

Given that the mutational target upstream of PFL_3078-3093 is likely be relatively

small and that these mutants are rapidly outcompeted by all WS types tested, their

relatively high frequency (3/43) is unexpected. Possibly this is due to a higher

mutation rate at these sites (Sankar, et al. 2016) or that population structure limits

direct competition between these different phenotypes and reduces the importance of

relative fitness. In SBW25 low fitness phenotypes that colonize the air-liquid

interface based on LPS modification or cell-chaining are observed prior to the rise of

WS to high frequencies (Lind, et al. 2017b) due to the presence of mutational hot

spots in these genes which make these mutants appear early during the growth phase

despite their relatively small mutational targets (Ferguson, et al. 2013; Farr 2015;

Lind, et al. 2017b).

In partially predicting evolutionary outcomes in *P. protegens*, this work lays the

foundation for future tests of evolutionary forecasting in related *Pseudomonas* species

by clearly stating predictions on several different levels from phenotype down to

which specific regions of proteins are likely to be mutated. Given what is already

known about the effects of (for now) unpredictable mutational biases and differences

in fitness between different WS types many of the forecasts will inevitably fail.

However hopefully they will fail in interesting ways thereby revealing erroneous

assumptions. The ability to remove common genetic and phenotypic pathways

provides a unique opportunity to also find those pathways that evolution does not

commonly use. This is necessary to determine why forecasts fail and update the

predictive models for another cycle of prediction, experimental evolution and mutant

characterization that make it possible to use this iterative model to define the

information necessary to predict short-term evolutionary processes.

Materials and methods

Strains and media

Pseudomonas protegens Pf-5 (previously known as *P. fluorescens* Pf-5) and

derivatives thereof were used for all experimental evolution and phenotypic

characterization. *E. coli* DH5α was used for cloning PCR fragments for genetic

engineering (Paulsen et al). *P. protegens* Pf-5 was grown in tryptic soy broth

(Tryptone 17g, Soytone 3g, Glucose 2.5g, NaCl 5g, K_2HPO_4 2.5g per liter)

supplemented with 10 mM MgSO4 and 0.2% glycerol (TSBGM) for experimental

evolution and fitness assays. Lysogeny broth (LB) was used during genetic

engineering and LB without NaCl and supplemented with 8% sucrose was used for

counter-selection of *sacB* marker. Solid media were 1.5% agar added to LB or TSB

supplemented with 10 mM MgSO4, 0.2% glycerol and 10 mg/l Congo red. Motility

assays were conducted in 0.3% agar TSB supplemented with 10 mM MgSO4, 0.2%

glycerol. Kanamycin was used at 50 mg/l for *E. coli* or 80 mg/l for *P. protegens* and

gentamicin at 10 mg/l for *E. coli* or 15 mg/L for *P. protegens*. Selection plates for

cloning contained 5-Bromo-4-Chloro-3-Indolyl β-D-Galactopyranoside (X-gal) at 40

mg/l. 100 mg/L nitrofurantoin was used to inhibit growth of *E. coli* donor cells after

conjugation. All strains were stored at -80°C in LB with 10% DMSO.

Experimental evolution

30 central wells of a deep well plate (polypropylene, 1.1 mL, round walls, Axygen

Corning Life Sciences) were inoculated with approximately 10^3 cells each and

incubated at 36°C for 5 days without shaking on two different occasions. Suitable

dilutions were plated on TSBGM plates with Congo red after 5 days and incubated at

36°C for 48 h. Plates were screened for colonies with a visible difference in colony

morphology and one divergent colony per well were randomly selected based only on

its position on the agar plate. In total 43 independent mutants were streaked for single

cells twice before overnight growth in LB and freezing. An identical protocol was

used for the Δ*wsp* Δ*aws* Δ*mws* strain.

Genome sequencing

Seven mutant strains that did not contain mutations in the *wspF* and *awsX* genes were

analyzed by genome resequencing. The strains had mutations in *awsR*, *mwsR*, *wspE,*

upstream PFL_3078 (2 strains) and in the intergenic region between *rrfB* and *recC*

upstream of the awsXRO operon. Genomic DNA was isolated with Genomic DNA

Purification Kit (Thermo Fisher). Sequencing libraries were prepared from 1μg DNA

using the TruSeq PCRfree DNA sample preparation kit (cat# FC- 121-3001/3002,

Illumina Inc.) targeting an insert size of 350bp. The library preparation was

performed according to the manufacturers' instructions (guide#15036187).

Sequencing was performed with MiSeq (Illumina Inc.) paired-end 300bp read length

and v3 sequencing chemistry.

Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The

facility is part of the National Genomics Infrastructure (NGI) Sweden and Science for

Life Laboratory. The SNP&SEQ Platform is also supported by the Swedish Research

Council and the Knut and Alice Wallenberg Foundation. Sequencing data were

analyzed with using Geneious v. 10.2.3 with reads assembled against the *P. protegens*

Pf-5 genome sequence (CP000076.1).

Sanger sequencing

Sanger sequencing were performed by GATC biotech and used to sequence candidate

genes to find adaptive mutations and to confirm reconstructed mutations. Primer

sequences are available in Table S1.

Reconstruction of mutations

Thirteen mutations representing all candidate genes found using Sanger or Illumina

sequencing as well as PFL_0087 and WspA mutations were reconstructed in the wild

type ancestral *P. protegens* Pf-5 to show that they are the cause of the adaptive

phenotype and to be able to assay their fitness effects without the risk of secondary

mutations that might have occurred during experimental evolution. A two-step allelic

replacement protocol was using to transfer the mutation into the ancestor. First a 1-2

kb fragment surrounding the putative adaptive mutations were amplified using PCR

(Phusion High- Fidelity DNA polymerase, Thermo Scientific) and ligated into the

multiple cloning site of the mobilizable pK18mobsac suicide plasmid (FJ437239)

using standard molecular techniques. The ligation mix was then transformed into

competent *E. coli* DH5α using heat shock. After confirmation of correct insert size by

PCR the plasmid was transferred to *P. protegens* Pf-5 by conjugation with the donor

strain and an *E. coli* strain carrying the conjugation helper plasmid pRK2013.

Cultures were grown overnight of the recipient *P. protegens* Pf-5 (20 ml per

conjugation at 30°C in LB) and 2 ml each of the donor and helper *E. coli* strains per

conjugation at 37°C in LB with kanamycin. The culture of *P. protegens* Pf-5 was heat

shocked for 10 minutes at 42°C prior to centrifugation at 4000 rpm for 10 minutes

and resuspension in a small volume of LB. Donor and helper cells were collected by

centrifugation 4000 rpm for 10 minutes, resuspended in LB, and mixed with the

concentrated recipient cells. After another round of centrifugation the conjugation mix

was resuspended in 50 µl LB and spread onto several spots on a LA plate followed by

incubation overnight at 30°C. Each spot of the conjugation mix was scraped of the

plate and resuspended in 200 µl LB each and plated on LA plates with kanamycin, to

select for transfer of the plasmid, and nitrofurantoin that prevents growth of the *E.*

coli donor and helper cells. The pK18mobsac plasmid has a pBR322 type origin and

cannot replicate in *P. protegens* Pf-5 and only cells where the plasmid has integrated

into the chromosome by homologous recombination, with the homology provided by

the cloned fragment, can grow in the presence of kanamycin. After streaking for

single cells on LA plates with kanamycin, the *P. protegens* Pf-5 strains with

integrated plasmids were grown overnight in LB at 30°C without antibiotics to allow

for double crossover homologous recombination resulting in loss of the integrated

plasmid. The plasmid also contains the *sacB* marker conferring sucrose sensitivity,

which allows for counter-selection by plating on LA plates with sucrose. Sucrose

resistant colonies were checked for loss of the kanamycin marker and DNA

sequencing of the cloned region to find strains with the reconstructed mutation and no

other mutations.

Deletion of the *wsp*, *aws*, *mws* and *pelABCDEFG* (PFL_2972-PFL_2978) regions was

accomplished using the same two-step allelic exchange protocol using SOE-PCR to

generated at fragment surrounding the operon as previously described (Ferguson, et

al. 2013; Farr 2015; Lind, et al. 2017b). Gene synthesis (Thermo Fisher) was used to

make DNA fragments used for deletion of PFL_0161-PFL_0164 and WspA T293-

E299. Primer sequences are available in Table S1.

Fitness assays

Two types of competition fitness assays were performed similarly to previously

described (Lind, et al. 2015). The first assay measures invasion fitness, where a

mutant is mixed 1:100 with the wild type ancestor, simulating early stages of air-

liquid interface colonization where a rare mutant establishes and grows at the surface

with no competition from other mutants. The second assay instead measures

competition fitness in a 1:1 competition against a reference mutant strain, which here

was chosen to be the WspF V271G mutant because it was the most commonly found

in the experimental evolution study and thus is highly successful either because of a

high rate of emergence, i.e. a mutational hot spot, or higher fitness than most other

WS mutants. In addition, the WspF V271G mutant has a temperature sensitive colony

morphology phenotype in that it is highly wrinkly at 30°C, but only have a very mild

phenotype when grown at room temperature, thus allowing it to be distinguishable

from both the smooth ancestor and all other wrinkly mutants isolated here.

Fluorescent reference strains of the wild type ancestor and the WspF V271G mutants

were created using a miniTn7 transposon (miniTn7(Gm) PA1/04/03 Gfp.AAV-a)

(Lambertsen, et al. 2004) that allows integration at a defined locus (attTn7) in the

chromosome. This allows the colonies to be distinguished not only by morphology

but also by fluorescence under blue/UV light and gentamicin resistance, which

provides a way to ascertain that secondary adaptive mutants that might occur during

the competition experiment do not bias the results (for example the ancestor could

evolve WS types or a WS mutant can evolve to cheat on the other type by inactivation

of EPS production or reduced c-di-GMP signalling). Introduction of the transposon

into *P. protegens* Pf-5 was performed by tri-parental conjugation from *E. coli* with

helper plasmids pRK2013 (conjugation helper) and pUX-BF13 containing the

transposase genes) using the same conjugation protocol described above.

The invasion assay was performed by mixing shaken overnight cultures of the

competitor 1:100 with the GFP-labeled reference ancestor followed by 1000-fold

dilution and static incubation at 36°C for 48 h in TSBGM medium in deep well plates

(1 ml per well, using only the central 60 wells). For the competition assay, the GFP-

labeled reference strain WspF 271G was mixed 1:1 with the competitor and diluted 6-

fold and grown for 4 h (shaken at 30°C), before plating to determine initial ratios, to

ensure the cells were in a similar physiological state at the start of the competition.

The competition cultures were then diluted 1000-fold in TSBGM medium and grown

in deep well plates (1 ml per well, using only the central 60 wells) static for 24 h at

36°C. Selection coefficients (s) were calculated as previously described (Dykhuizen

1990), where s = 0 is equal fitness, positive is increased fitness and negative is

decreased fitness relative to the reference strain. Briefly s is calculated as the change

in logarithmic ratio over time according to $s = [\ln(R(t)/R(0))]/[t]$, where R is the ratio

of mutant to reference and t is the number of generations of the entire population

during the experiment (estimated from viable counts). The cost of the fluorescent

marker were calculated from control competitions where the GFP-labeled reference

strains (wild type and WspF V271G) were competed against isogenic strains without

the marker and included in each plate under identical conditions during the fitness

assays and used to adjust the selection coefficients to compensate for the cost.

Motility assays

Swimming motility assays were performed in TSBGM plates with 0.3% agar (BD)

and the diameter was measured after 24 h of growth at room temperature. Each strain

was assayed in duplicates on two different plates.

Bioinformatics analysis of DGCs and EPS genes

Homologs for all DGCs in *P. fluorescens* SBW25 were found using the *Pseudomonas*

Ortholog Database at Pseudomonas.com (Winsor, et al. 2016). Blast-p searches for

GGDEF domains were performed to find remaining DGCs in the six *Pseudomonas*

species and their homologs again found using the *Pseudomonas* Ortholog Database

(Whiteside, et al. 2013) and manually inspected. Annotations (Pseudomonas.com. DB

version 17.2) were also searched for diguanylate cyclase and GGDEF. Not all DCCs

found are likely to have diguanylate cyclase activity, but given the difficulties of

predicting which of the partly degenerate active sites are likely to be inactive

combined with the possibilities of mutational activation during experimental

evolution, none were excluded.

There is no simple way to find all genes that can function as structural or regulatory

genes to allow colonization of the air-liquid interface. Thus the selection in Figure 3B

and Figure 3 – source data should not be considered complete. Putative EPS genes

were found using blastp searches with sequences from known exopolysaccharide

biosynthesis proteins including cellulose, PGA, Pel, Psl, Pea, Peb, alginate and levan.

Homologs were then found using the *Pseudomonas* Ortholog Database (Whiteside, et

al. 2013) at Pseudomonas.com (Winsor, et al. 2016). Annotations (Pseudomonas.com.

DB version 17.2) were also searched for glycosyltransferase, glycosyl transferase,

flippase, polysaccharide, lipopolysaccharide, polymerase, biofilm, adhesion and

adhesion. Based on previous work in SBW25 and literature searches a few additional

genes were added.

Acknowledgements

963 This work was supported by grants from Carl Tryggers Foundation for Scientific

964 Research and Magnus Bergvalls Foundation.

Competing interests

967 The author declares no competing interests.

References

Bantinaki E, Kassen R, Knight CG, Robinson Z, Spiers AJ, Rainey PB. 2007.

Adaptive divergence in experimental populations of *Pseudomonas fluorescens*. III.

Mutational origins of wrinkly spreader diversity. Genetics 176:441-453.

http://dx.doi.org/10.1534/genetics.106.069906

Barrick JE, Lenski RE. 2013. Genome dynamics during experimental evolution.

Nature reviews. Genetics 14:827-839. http://dx.doi.org/10.1038/nrg3564

Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, Lenski RE, Kim JF.

2009. Genome evolution and adaptation in a long-term experiment with *Escherichia*

coli. Nature 461:1243-1247. http://dx.doi.org/10.1038/nature08480

Beaumont HJ, Gallie J, Kost C, Ferguson GC, Rainey PB. 2009. Experimental

evolution of bet hedging. Nature 462:90-93. http://dx.doi.org/10.1038/nature08504

Blank D, Wolf L, Ackermann M, Silander OK. 2014. The predictability of molecular

evolution during functional innovation. Proceedings of the National Academy of

Sciences of the United States of America 111:3044-3049.

http://dx.doi.org/10.1073/pnas.1318797111

Blount ZD, Lenski RE, Losos JB. 2018. Contingency and determinism in evolution:

Replaying life's tape. Science 362. http://dx.doi.org/10.1126/science.aam5979

Brandis G, Pietsch F, Alemayehu R, Hughes D. 2015. Comprehensive phenotypic

characterization of rifampicin resistance mutations in *Salmonella* provides insight into

the evolution of resistance in *Mycobacterium tuberculosis*. J Antimicrob Chemother

70:680-685. http://dx.doi.org/10.1093/jac/dku434

Conrad TM, Joyce AR, Applebee MK, Barrett CL, Xie B, Gao Y, Palsson BO. 2009.

Whole-genome resequencing of *Escherichia coli* K-12 MG1655 undergoing short-

term laboratory evolution in lactate minimal media reveals flexible selection of

adaptive mutations. Genome biology 10:R118. http://dx.doi.org/10.1186/gb-2009-10-

10-r118

Deatherage DE, Kepner JL, Bennett AF, Lenski RE, Barrick JE. 2017. Specificity of

genome evolution in experimental populations of Escherichia coli evolved at different

temperatures. Proceedings of the National Academy of Sciences of the United States

of America 114:E1904-E1912. http://dx.doi.org/10.1073/pnas.1616132114

Dykhuizen DE. 1990. Experimental studies of natural selection in bacteria. Annu.

Rev. Ecol. Syst. 21:373-398.

Farr AD. 2015. Formalist features determining the tempo and mode of evolution in

Pseudomonas fluorescens SBW25. [[Auckland, New Zealand]: Massey University.

Farr AD, Remigi P, Rainey PB. 2017. Adaptive evolution by spontaneous domain

fusion and protein relocalization. Nat Ecol Evol 1:1562-1568.

http://dx.doi.org/10.1038/s41559-017-0283-7

Ferguson GC, Bertels F, Rainey PB. 2013. Adaptive Divergence in Experimental

Populations of *Pseudomonas fluorescens*. V. Insight into the Niche Specialist "Fuzzy

Spreader" Compels Revision of the Model Pseudomonas Radiation. Genetics.

http://dx.doi.org/10.1534/genetics.113.154948

Firnberg E, Labonte JW, Gray JJ, Ostermeier M. 2014. A comprehensive, high-

resolution map of a gene's fitness landscape. Molecular biology and evolution

31:1581-1592. http://dx.doi.org/10.1093/molbev/msu081

Fraebel DT, Mickalide H, Schnitkey D, Merritt J, Kuhlman TE, Kuehn S. 2017.

Environment determines evolutionary trajectory in a constrained phenotypic space.

Elife 6. http://dx.doi.org/10.7554/eLife.24669

Friedman L, Kolter R. 2004. Genes involved in matrix formation in *Pseudomonas*

aeruginosa PA14 biofilms. Mol Microbiol 51:675-690.

Gallie J, Libby E, Bertels F, Remigi P, Jendresen CB, Ferguson GC, Desprat N,

Buffing MF, Sauer U, Beaumont HJ, et al. 2015. Bistability in a metabolic network

underpins the de novo evolution of colony switching in Pseudomonas fluorescens.

PLoS biology 13:e1002109. http://dx.doi.org/10.1371/journal.pbio.1002109

Goymer P, Kahn SG, Malone JG, Gehrig SM, Spiers AJ, Rainey PB. 2006. Adaptive

divergence in experimental populations of *Pseudomonas fluorescens*. II. Role of the

GGDEF regulator WspR in evolution and development of the wrinkly spreader

phenotype. Genetics 173:515-526. http://dx.doi.org/10.1534/genetics.106.055863

Herron MD, Doebeli M. 2013. Parallel evolutionary dynamics of adaptive

diversification in *Escherichia coli*. PLoS biology 11:e1001490.

http://dx.doi.org/10.1371/journal.pbio.1001490

Hickman JW, Tifrea DF, Harwood CS. 2005. A chemosensory system that regulates

biofilm formation through modulation of cyclic diguanylate levels. Proceedings of the

National Academy of Sciences of the United States of America 102:14422-14427.

http://dx.doi.org/10.1073/pnas.0507170102

Jacquier H, Birgy A, Le Nagard H, Mechulam Y, Schmitt E, Glodt J, Bercot B, Petit

E, Poulain J, Barnaud G, et al. 2013. Capturing the mutational landscape of the beta-

lactamase TEM-1. Proceedings of the National Academy of Sciences of the United

States of America 110:13067-13072. http://dx.doi.org/10.1073/pnas.1215206110

Jahn LJ, Munck C, Ellabaan MMH, Sommer MOA. 2017. Adaptive Laboratory

Evolution of Antibiotic Resistance Using Different Selection Regimes Lead to

Similar Phenotypes and Genotypes. Front Microbiol 8:816.

http://dx.doi.org/10.3389/fmicb.2017.00816

Jerison ER, Desai MM. 2015. Genomic investigations of evolutionary dynamics and

epistasis in microbial evolution experiments. Curr Opin Genet Dev 35:33-39.

http://dx.doi.org/10.1016/j.gde.2015.08.008

Jimenez JI, Xulvi-Brunet R, Campbell GW, Turk-MacLeod R, Chen IA. 2013.

Comprehensive experimental fitness landscape and evolutionary network for small

RNA. Proceedings of the National Academy of Sciences of the United States of

America 110:14984-14989. http://dx.doi.org/10.1073/pnas.1307604110

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 web

portal for protein modeling, prediction and analysis. Nature protocols 10:845-858.

http://dx.doi.org/10.1038/nprot.2015.053

Kishimoto T, Iijima L, Tatsumi M, Ono N, Oyake A, Hashimoto T, Matsuo M,

Okubo M, Suzuki S, Mori K, et al. 2010. Transition from positive to neutral in

mutation fixation along with continuing rising fitness in thermal adaptive evolution.

PLoS genetics 6:e1001164. http://dx.doi.org/10.1371/journal.pgen.1001164

Knöppel A, Knopp M, Albrecht LM, Lundin E, Lustig U, Näsvall J, Andersson DI.

2018. Genetic Adaptation to Growth Under Laboratory Conditions in *Escherichia coli*

and *Salmonella enterica*. Front Microbiol 9:756.

http://dx.doi.org/10.3389/fmicb.2018.00756

Koskiniemi S, Sun S, Berg OG, Andersson DI. 2012. Selection-driven gene loss in

bacteria. PLoS genetics 8:e1002787. http://dx.doi.org/10.1371/journal.pgen.1002787

Kram KE, Geiger C, Ismail WM, Lee H, Tang H, Foster PL, Finkel SE. 2017.

Adaptation of *Escherichia coli* to Long-Term Serial Passage in Complex Medium:

Evidence of Parallel Evolution. mSystems 2.

http://dx.doi.org/10.1128/mSystems.00192-16

LaCroix RA, Sandberg TE, O'Brien EJ, Utrilla J, Ebrahim A, Guzman GI, Szubin R,

Palsson BO, Feist AM. 2015. Use of adaptive laboratory evolution to discover key

mutations enabling rapid growth of *Escherichia coli* K-12 MG1655 on glucose

minimal medium. Appl Environ Microbiol 81:17-30.

http://dx.doi.org/10.1128/AEM.02246-14

Lambertsen L, Sternberg C, Molin S. 2004. Mini-Tn7 transposons for site-specific

tagging of bacteria with fluorescent proteins. Environmental microbiology 6:726-732.

http://dx.doi.org/10.1111/j.1462-2920.2004.00605.x

Lee MC, Marx CJ. 2012. Repeated, selection-driven genome reduction of accessory

genes in experimental populations. PLoS genetics 8:e1002651.

http://dx.doi.org/10.1371/journal.pgen.1002651

Lee VT, Matewish JM, Kessler JL, Hyodo M, Hayakawa Y, Lory S. 2007. A cyclic-

di-GMP receptor required for bacterial exopolysaccharide production. Mol Microbiol

65:1474-1484. http://dx.doi.org/10.1111/j.1365-2958.2007.05879.x

Liang ZX. 2015. The expanding roles of c-di-GMP in the biosynthesis of

exopolysaccharides and secondary metabolites. Nat Prod Rep 32:663-683.

http://dx.doi.org/10.1039/c4np00086b

Lind PA, Arvidsson L, Berg OG, Andersson DI. 2017a. Variation in mutational

robustness between different proteins and the predictability of fitness effects.

Molecular biology and evolution 34:408-418.

http://dx.doi.org/10.1093/molbev/msw239

Lind PA, Farr AD, Rainey PB. 2017b. Evolutionary convergence in experimental

Pseudomonas populations. ISME J 11:589-600.

http://dx.doi.org/10.1038/ismej.2016.157

Lind PA, Farr AD, Rainey PB. 2015. Experimental evolution reveals hidden diversity

in evolutionary pathways. eLife 4. http://dx.doi.org/10.7554/eLife.07074

Lind PA, Libby E, Herzog J, Rainey PB. 2019. Predicting mutational routes to new

adaptive phenotypes. eLife 8. http://dx.doi.org/10.7554/eLife.38822

Long A, Liti G, Luptak A, Tenaillon O. 2015. Elucidating the molecular architecture

of adaptation via evolve and resequence experiments. Nature reviews. Genetics

16:567-582. http://dx.doi.org/10.1038/nrg3937

Lukacisinova M, Bollenbach T. 2017. Toward a quantitative understanding of

antibiotic resistance evolution. Curr Opin Biotechnol 46:90-97.

http://dx.doi.org/10.1016/j.copbio.2017.02.013

Lundin E, Tang PC, Guy L, Nasvall J, Andersson DI. 2017. Experimental

determination and prediction of the fitness effects of random point mutations in the

biosynthetic enzyme HisA. Molecular biology and evolution.

http://dx.doi.org/10.1093/molbev/msx325

McDonald MJ, Cooper TF, Beaumont HJE, Rainey PB. 2011. The distribution of

fitness effects of new beneficial mutations in *Pseudomonas fluorescens*. Biology

Letters 7:98-100. http://dx.doi.org/Doi 10.1098/Rsbl.2010.0547

McDonald MJ, Gehrig SM, Meintjes PL, Zhang XX, Rainey PB. 2009. Adaptive

divergence in experimental populations of *Pseudomonas fluorescens*. IV. Genetic

constraints guide evolutionary trajectories in a parallel adaptive radiation. Genetics

183:1041-1053. http://dx.doi.org/10.1534/genetics.109.107110

McElroy KE, Hui JG, Woo JK, Luk AW, Webb JS, Kjelleberg S, Rice SA, Thomas

T. 2014. Strain-specific parallel evolution drives short-term diversification during

Pseudomonas aeruginosa biofilm formation. Proceedings of the National Academy of

Sciences of the United States of America 111:E1419-1427.

http://dx.doi.org/10.1073/pnas.1314340111

Morgan JL, McNamara JT, Zimmer J. 2014. Mechanism of activation of bacterial

cellulose synthase by cyclic di-GMP. Nat Struct Mol Biol 21:489-496.

http://dx.doi.org/10.1038/nsmb.2803

O'Neill AJ, Huovinen T, Fishwick CW, Chopra I. 2006. Molecular genetic and

structural modeling studies of *Staphylococcus aureus* RNA polymerase and the

fitness of rifampin resistance genotypes in relation to clinical prevalence. Antimicrob

Agents Chemother 50:298-309. http://dx.doi.org/10.1128/AAC.50.1.298-309.2006

Orgogozo V. 2015. Replaying the tape of life in the twenty-first century. Interface

Focus 5:20150057. http://dx.doi.org/10.1098/rsfs.2015.0057

Rainey PB, Travisano M. 1998. Adaptive radiation in a heterogeneous environment.

Nature 394:69-72. http://dx.doi.org/10.1038/27900

Römling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a

universal bacterial second messenger. Microbiology and molecular biology reviews :

MMBR 77:1-52. http://dx.doi.org/10.1128/MMBR.00043-12

Sandberg TE, Pedersen M, LaCroix RA, Ebrahim A, Bonde M, Herrgard MJ, Palsson

BO, Sommer M, Feist AM. 2014. Evolution of *Escherichia coli* to 42 degrees C and

subsequent genetic engineering reveals adaptive mechanisms and novel mutations.

Molecular biology and evolution 31:2647-2662.

http://dx.doi.org/10.1093/molbev/msu209

Sankar TS, Wastuwidyaningtyas BD, Dong Y, Lewis SA, Wang JD. 2016. The nature

of mutations induced by replication-transcription collisions. Nature.

http://dx.doi.org/10.1038/nature18316

Schenk MF, Szendro IG, Krug J, de Visser JA. 2012. Quantifying the adaptive

potential of an antibiotic resistance enzyme. PLoS genetics 8:e1002783.

http://dx.doi.org/10.1371/journal.pgen.1002783

Silby MW, Cerdeno-Tarraga AM, Vernikos GS, Giddens SR, Jackson RW, Preston

GM, Zhang XX, Moon CD, Gehrig SM, Godfrey SA, et al. 2009. Genomic and

genetic analyses of diversity and plant interactions of *Pseudomonas fluorescens*.

Genome biology 10:R51. http://dx.doi.org/10.1186/gb-2009-10-5-r51

Sleight SC, Orlic C, Schneider D, Lenski RE. 2008. Genetic basis of evolutionary

adaptation by *Escherichia coli* to stressful cycles of freezing, thawing and growth.

Genetics 180:431-443. http://dx.doi.org/10.1534/genetics.108.091330

Sommer MOA, Munck C, Toft-Kehler RV, Andersson DI. 2017. Prediction of

antibiotic resistance: time for a new preclinical paradigm? Nature reviews.

Microbiology 15:689-696. http://dx.doi.org/10.1038/nrmicro.2017.75

Spiers AJ, Bohannon J, Gehrig SM, Rainey PB. 2003. Biofilm formation at the air-

liquid interface by the *Pseudomonas fluorescens* SBW25 wrinkly spreader requires an

acetylated form of cellulose. Molecular microbiology 50:15-27.

Spiers AJ, Kahn SG, Bohannon J, Travisano M, Rainey PB. 2002. Adaptive

divergence in experimental populations of *Pseudomonas fluorescens*. I. Genetic and

phenotypic bases of wrinkly spreader fitness. Genetics 161:33-46.

Spiers AJ, Rainey PB. 2005. The *Pseudomonas fluorescens* SBW25 wrinkly spreader

biofilm requires attachment factor, cellulose fibre and LPS interactions to maintain

strength and integrity. Microbiology 151:2829-2839.

http://dx.doi.org/10.1099/mic.0.27984-0

Steiner S, Lori C, Boehm A, Jenal U. 2013. Allosteric activation of exopolysaccharide

synthesis through cyclic di-GMP-stimulated protein-protein interaction. EMBO J

32:354-368. http://dx.doi.org/10.1038/emboj.2012.315

Tenaillon O, Rodriguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD,

Gaut BS. 2012. The molecular diversity of adaptive convergence. Science 335:457-

461. http://dx.doi.org/10.1126/science.1212986

Whiteside MD, Winsor GL, Laird MR, Brinkman FS. 2013. OrtholugeDB: a bacterial

and archaeal orthology resource for improved comparative genomic analysis. Nucleic

Acids Res 41:D366-376. http://dx.doi.org/10.1093/nar/gks1241

Whitney JC, Whitfield GB, Marmont LS, Yip P, Neculai AM, Lobsanov YD,

Robinson H, Ohman DE, Howell PL. 2015. Dimeric c-di-GMP is required for post-

translational regulation of alginate production in *Pseudomonas aeruginosa*. J Biol

Chem 290:12451-12462. http://dx.doi.org/10.1074/jbc.M115.645051

Wichman HA, Badgett MR, Scott LA, Boulianne CM, Bull JJ. 1999. Different

trajectories of parallel evolution during viral adaptation. Science 285:422-424.

Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. 2016. Enhanced

annotations and features for comparing thousands of *Pseudomonas* genomes in the

Pseudomonas genome database. Nucleic Acids Res 44:D646-653.

http://dx.doi.org/10.1093/nar/gkv1227

1180

Figure 5 - source data. Mutations found after experimental evolution

Genome position CP000076.1	Type	Change	Gene position	Gene locus	Gene symbol	Effect	Mutation found in SBW25
859290	Transition	g->a	-129	PFL_0740	recC	rrfB terminator/recCBD promoter upstream awsXRO	No
859290	Transition	g->a	-129	PFL_0740	recC	rrfB terminator/recCBD promoter upstream awsXRO	No
868744	Transversion	c->a	101	PFL_0743	awsX, yfiR	S34Y	No
868932	Transversion	t->g	289	PFL_0743	awsX, yfiR	Y97D	No
869088	Transversion	t->g	445	PFL_0743	awsX, yfiR	C149G	No
869281	Transversion	t->g	638	PFL_0743	awsX, yfiR	V213G	No
869281	Transversion	t->g	638	PFL_0743	awsX, yfiR	V213G	No
869281	Transversion	t->g	638	PFL_0743	awsX, yfiR	V213G	No
868832-868840	Deletion	Deletion 9 bp	189-197	PFL_0743	awsX, yfiR	Deletion F63-L65	No
868832-868840	Deletion	Deletion 9 bp	189-197	PFL_0743	awsX, yfiR	Deletion F63-L65	No
869040-869072	Deletion	Deletion 33 bp	397-429	PFL_0743	awsX, yfiR	Deletion V133-P143	No
869913	Transition	c->t	581	PFL_0744	awsR, yfiN	A194V	Yes
869994	Transition	a->g	662	PFL_0744	awsR, yfiN	D221G	D->A
870002	Transition	c->t	670	PFL_0744	awsR, yfiN	L226F	No
1301093	Transition	c->t	1982	PFL_1133	wspE	S661L	Yes
1302246	Transition	g->a	808	PFL_1134	wpsF	G270R	Yes
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No

1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302250	Transversion	t->g	812	PFL_1134	wspF	V271G	No
1302327	Transition	c->a	889	PFL_1134	wspF	Q297K	Q->R
1302327	Transition	c->a	889	PFL_1134	wspF	Q297K	Q->R
1301579-1301593	Duplication	Duplication 15 bp	141-155	PFL_1134	wspF	Duplication L51-I55	Del L51-I55
3548383	Transition	c->t	-98	PFL_3078		PFL_3078-PFL_3093 promoter	
3548383	Transition	c->t	-98	PFL_3078		PFL_3078-PFL_3093 promoter	
3548385	Transition	c->t	-96	PFL_3078		PFL_3078-PFL_3093 promoter	
6114402	Transversion	a->t	2237	PFL_5345	mwsR, morA	Q746L	No
6114696	Transition	a->g	2531	PFL_5345	mwsR, morA	H844R	No
6114940	Transition	g->a	2775	PFL_5345	mwsR, morA	M925I	Yes
6115236	Transversion	t->a	3071	PFL_5345	mwsR, morA	L1024Q	No
6115406	Transition	g->a	3241	PFL_5345	mwsR, morA	E1081K	Yes
6115406	Transition	g->a	3241	PFL_5345	mwsR, morA	E1081K	Yes
6115406	Transition	g->a	3241	PFL_5345	mwsR, morA	E1081K	Yes
6115406	Transition	g->a	3241	PFL_5345	mwsR, morA	E1081K	Yes
6115415	Transition	g->a	3250	PFL_5345	mwsR, morA	G1084R	G->S
6115107-6115130	Duplication	Duplication 24 bp	2942-2965	PFL_5345	mwsR, morA	P981-S988	No

Δwsp Δaws Δmws							
86862-86873	Deletion	Deletion 12 bp	1389-1400	PFL_0087		S463R, deletion N464-Q466	
86876-86893	Deletion	Deletion 18 bp	1403-1420	PFL_0087		Deletion D470-S475	
86949-86978	Deletion	Deletion 30 bp	1476-1505	PFL_0087		Deletion Q494-L503	
86949-86978	Deletion	Deletion 30 bp	1476-1505	PFL_0087		Deletion Q494-L503	
86972-87172	Duplication	Duplication 201 bp	1499-1699	PFL_0087		Insertion 67 aa	
87014-87199	Duplication	Duplication 185 bp	1541-1726	PFL_0087		Duplication E544-P571	

Figure 7 - source data. Fitness assays

Invasion - selection coefficients

Wild type	0,003	0,002	-0,005	0,012	0,013	0,010	-0,039	0,021	-0,017
WspF V271G	0,363	0,330	0,303	0,318	0,319	0,340			
WspF Q297K	0,362	0,225	0,347	0,331	0,269	0,282			
WspE S661L	0,334	0,349	0,377	0,362	0,335	0,391			
AwsX del V133-P143	0,300	0,219	0,358	0,223	0,271	0,261			
AwsR A194V	0,308	0,351	0,280	0,262	0,292	0,262			
Promoter AwsXRO	0,412	0,526	0,476	0,348	0,336	0,376			
MwsR M925I	0,392	0,394	0,432	0,346	0,341	0,373			
MwsR E1081K	0,322	0,420	0,357	0,326	0,321	0,283			
PFL_3078 Promoter	0,313	0,351	0,320	0,321	0,334	0,326			
PFL_0087 D470-S475	0,390	0,407	0,433	0,393	0,417	0,429			
PFL_0087 Q494-L503	0,428	0,378	0,383	0,419	0,453	0,433			
WspA del T293-E299	0,435	0,439	0,440	0,444	0,429	0,416			

Competition - selection coefficients

Wild type	-0,230	-0,207	-0,196	-0,175				
WspF V271G	0,033	-0,021	-0,011	-0,029	-0,014	0,027	-0,015	0,010
WspF Q297K	0,015	0,017	-0,009	-0,010				
WspE S661L	-0,008	0,001	0,002	0,050				
AwsX ΔV133-P143	-0,030	-0,082	-0,088	-0,058				
AwsR A194V	-0,113	-0,079	0,008	-0,020				
Promoter AwsXRO	0,041	0,009	0,034	0,039				
MwsR M925I	-0,006	-0,007	-0,011	0,003				
MwsR E1081K	0,085	0,072	0,036	0,050				
PFL_3078 Promoter	-0,066	-0,146	-0,088	-0,102				
PFL_0087 D470-S475	-0,055	-0,056	-0,069	-0,089				
PFL_0087 Q494-L503	-0,056	-0,050	-0,030	-0,026				
WspA del T293-E299	-0,145	-0,068	-0,062	-0,049				

Figure 8 - source data. Fitness assays for del pel mutants

Invasion - selection coefficients

Δ*pel* WspF V271G	0,212	0,253	0,227	0,199	0,197	0,201
Δ*pel* WspF Q297K	0,210	0,162	0,173	0,176	0,212	0,223
Δ*pel* AwsX del V133-P143	0,311	0,222	0,184	0,202	0,203	0,178
Δ*pel* MwsR E1081K	0,225	0,209	0,248	0,204	0,185	0,205
Δ*pel* PFL_0087 D470-S475	0,223	0,244	0,206	0,222	0,259	0,257

Competition - selection coefficients

Δ*pel* WspF V271G	-0,171	-0,177	-0,206	-0,174
Δ*pel* WspF Q297K	-0,041	-0,114	-0,104	-0,069
Δ*pel* AwsX del V133-P143	-0,171	-0,104	-0,200	-0,151
Δ*pel* MwsR E1081K	-0,042	-0,025	-0,112	-0,072
Δ*pel* PFL_0087 D470-S475	-0,179	-0,255	-0,179	-0,237

Table S1. Oligonucleotide primers used for PCR, sequencing and genetic reconst
Cloning

pEX18Gm_MCS_F	tgttgtgtggaattgtgag
pEX18Gm_MCS_R	ctgcaaggcgattaagttg

wsp **operon**

PFL_1133_EcoRI_F	agttgctggcggagaaaac
WspE_1550F	ttcccgctggcccatatcga
WspE_2119F	atcaccgacatcgacatgc
WspE_2213F	gtcctacaaggaccgtga
WspF_154R	caccggcatgatcaggtccat
WspR_70R	gcttcgccgatcatggcctg
WspR_398F	tggtggcgcgcattcgcta
WspR_553R	ccactccagctccaggta
WspR_620R	gtaggtcttgaagtagtcg
PFL_1128_F_EcoRI	tagGAATTCcaagaaggcccgcaagtac
PFL_1128_UR_SOE	cgtatcctgctagattccgaatgagtcagtcctcgaggccat
PFL_1135_DF_SOE	cattcggaatctagcaggatacggagtaaccgcctcagcttc
PFL_1136_F_SalI	catGTCGACgaatcactgattcacggcag

aws **operon**

RecC_F_HindIII	tagaagcttctgataccgcccaagagttc
RecC_R_KpnI	catggtaccccagtcggctcgatatacct
recD_F_EcoRI	acggaattcccactacctgaatgtactg
PFL_0743_R_SOE	cgttagaaatcggctaggcttcagagaaaggccggaaacatg
PFL_0743-SalI_F	tgtgtgctgtcgaccattc
awsR_182R	agctgatggagcgggcgatca
awsR_677F	cgacttcaacgccctgct
PFL_0745_F_SOE	tgaagcctagccgatttctaacgatcgtggtgatcgccgact
awsO_DR	ttacatccgcgaggtgac
PFL_0746_F_HindIII	tgcaagcttcttcttcatcaccccgcaa

mwsR

PFL_5345_UF_EcoRI	actgaattcatggcttccagcttggcct
PFL_5345_UR_SOE	cagtcgattacgtactatagcaggctgtgatcctgcggtttg
mwsR_HindIII_F	gttaagcttagacccagctgttcctgttc
mwsR_2144F	gccgcgacatcagccagca
PFL_5345_SeqF	gaaaaggacctgcgcatg
PFL_5345_SeqR	gaactgcttgaggtagttc
mwsR_3599R	gaaggtgcggtcgatcttca
mwsR_SalI_R	aggccgtcgacgaaggtgc
PFL_5345_DF_SOE	ctgctatagtacgtaatcgactgcatcccctgcttgcgtaat
PFL_5346_R_HindIII	gtcaagctttcagcgatttcgcggaaac

PFL_0087

PFL_0087_KpnI_F	catggtaccgaagccgttgcgacgaatc

PFL_0087_seqF   aagcaaccatttcgcctgt
PFL_0087_seqR   tgccatgttgttgccattg
PFL_0087_HindIII_R   tagaagcttggctcctcaaccagcttc

PFL_3078-3093
PFL_3077_KpnI_F   catggtaccgcgaaagtcccggttgaag
PFL_3078_HindIII_R   tagaagcttgttggccatctcgttcatg

pelABCDEFG
PFL_2971_F_EcoRI   ctggaattcagcgagtactacctggactt
PFL_2972_DelCh   caaggccaatgcggtaaaca
PFL_2972_UR_SOE   attgggccctctatgtcgacatcaactcactcttcagtagatcaatct
PFL_2978_DF_SOE   gatgtcgacatagagggcccaatttgaggagcatcggcaagc
PFL_2979_R_HindIII   agtgcccaagagcagaagc