Classification: Biological Sciences: Evolution

The tempo of linked selection: emergence of a heterogeneous
genomic landscape during a recent radiation of monkeyflowers

Sean Stankowski[1,2,§], Madeline A. Chase[1,3,§], Allison M. Fuiten[1], Peter L. Ralph[1], and
Matthew A. Streisfeld[1*].

[1]Institute of Ecology and Evolution, 335 Pacific Hall 5289, University of Oregon,
Eugene, OR 97405, USA
[2]Present address: Department of Animal and Plant Sciences, University of Sheffield,
Sheffield S10 2TN, UK
[3]Present address: Department of Evolutionary Biology, Evolutionary Biology Centre
(EBC), Uppsala University, Uppsala, Sweden
[§]These authors contributed equally to this work
*Corresponding author: mstreis@uoregon.edu

ORCID ids:
SS: https://orcid.org/0000-0003-0472-9299; MAC: https://orcid.org/0000-0002-7916-
3560; AMF: https://orcid.org/0000-0003-3926-5455 PLR: https://orcid.org/0000-0002-
9459-6866; MAS: https://orcid.org/0000-0002-2660-8642

Keywords: genome assembly | lineage sorting | *Mimulus* | differentiation landscape

Abstract—What are the processes that shape patterns of genome-wide variation
between emerging species? This question is central to our understanding of the origins of
biodiversity and the fundamental principles governing molecular evolution. It is now
clear that indirect selection on linked neutral variation (hereafter 'linked selection') plays
an important role in shaping heterogeneous patterns of genome-wide diversity and
differentiation within and between species. However, we do not know how these
characteristic signatures of linked selection evolve over time. To fill this critical
knowledge gap, we construct the first chromosome-level genome assembly for the bush
monkeyflower, and use it to show that linked selection has been a primary architect of
heterogeneous patterns of lineage sorting, differentiation, and nucleotide diversity across
a recent radiation. By taking advantage of the range of divergence times between the
different pairs of monkeyflower taxa, we also show how the signatures of linked selection
evolve as populations diverge: linked selection occurring within lineages acts to conserve
an ancestral pattern of diversity after a population split, while its joint action in separate
lineages causes a common differentiation landscape to emerge between them. Together,
our study demonstrates linked selection's power to shape patterns of genome-wide
variation within and between taxa, and provides critical insight into how genome-wide
variation evolves during the first 1.5 million years of divergence.

Significance

What are the processes that shape patterns of genome-wide variation between emerging species? Because nucleotides are linked together on chromosomes, even neutral variants are impacted by selection on mutations that arise at neighboring sites. We show that this phenomenon, referred to as linked selection, was important in causing common patterns of differentiation to evolve between taxa during a radiation of monkeyflowers. This signature begins to emerge shortly after divergence begins, but it takes 1.5 million years to become pronounced. This result fills a critical gap in our knowledge about how genomes evolve, it and shows how linked selection shapes patterns of differentiation variation soon after a population split, which is critical to our understanding of divergence and speciation.

Introduction

Since the first discoveries of abundant genetic variation in nature, evolutionary geneticists have sought to understand the processes that shape patterns of polymorphism and divergence within and between species (Lewontin and Hubby 1966; Kimura 1968; Lewontin 1974; Kimura 1986; Gillespie 1994; Casillas and Barbadilla 2017). The neutral theory explained how mutation and drift could shape genetic variation (Kimura 1968; Ohta 1973). Despite work suggesting the importance of non-neutral forces (Langley and Fitch 1974; Gillespie 1994; Kreitman 1996; Hahn 2008), it has remained the default assumption of most molecular genetic analyses, partly because of a lack of concrete, alternative models. However, genome-wide studies have revealed heterogeneous patterns of genetic variation that are inconsistent with purely neutral forces (Begun et al. 2007; Langley et al. 2012; Charlesworth and Charlesworth 2018; Kern and Hahn 2018). These genomic 'landscapes' are important confounders for work in other fields, such as speciation research (Burri 2017a, b; Ravinet et al. 2017), and provide intriguing clues in their own right into the ongoing evolutionary forces shaping our own genomes.

Heterogeneous genomic landscapes are increasingly understood to be formed due to selection on linked neutral variation (Charlesworth and Charlesworth 2018). For example, variable patterns of genetic diversity have now been observed across the genomes of a diverse range of plants and animals and appear to have been shaped by variation in the intensity of linked selection acting across the genome (Corbett-Detig et al. 2015; Kern and Hahn 2018). This occurs because linked selection—acting on deleterious or beneficial mutations—reduces the amount of genetic variation available to future generations at linked sites, similar to a reduction in local effective population size (N_e) (Maynard-Smith and Haigh 1974; Charlesworth et al. 1993; Hudson and Kaplan 1995; Gillespie 2000; Coop and Ralph 2012). In theoretical models, the local intensity of linked selection is determined by the density of targets of selection relative to the recombination rate, such that larger reductions in diversity occur in regions with more frequent selection and less recombination (Maynard-Smith and Haigh 1974; Charlesworth et al. 1993; Hudson and Kaplan 1995; Gillespie 2000).

Compared to patterns of within-species diversity, we know relatively little about how linked selection shapes patterns of genome-wide variation between emerging species (Burri 2017a, b). Although rates of differentiation and lineage sorting should be accelerated in regions of the genome that that have experienced long-term reductions in diversity (Charlesworth 1998; Cutter and Payseur 2013; Pease and Hahn 2013), we do

not know how long it takes for these signatures of linked selection to emerge across the
genome. This gap in our knowledge has important implications for understanding the
genomic basis of speciation, as regions of the genome restricting gene flow may be
mistaken with those experiencing linked selection (Ravinet et al. 2017). Unlike patterns
of diversity, which are inherited from a common ancestor and then maintained by
recurrent linked selection, a heterogeneous pattern of differentiation should emerge
gradually after a population split, as linked selection begins to shape patterns of variation
separately within each lineage. Thus, as the divergence time between populations
increases, recurrent linked selection should strengthen relationships between
differentiation and (*i*) nucleotide diversity, (*ii*) gene density, (*iii*) and recombination rate.
However, these temporal dynamics remain largely unexplored (but see Delmore et al.
2018), which has left a critical gap in our knowledge of the processes that shape patterns
of genome-wide variation between emerging species.
In this paper, we study the temporal signatures of linked selection using taxa from
the bush monkeyflower radiation (Figure 1). This recent radiation of perennial shrubs is
distributed throughout California and consists of seven subsecies of *Mimulus*
aurantiacus, one with two ecotypes (Chase et al. 2017). Together with their sister species
M. clevelandii, they span a variety of divergence times, ranging from locally adapted
ecotypes to species separated by ~1.5 million years (McMinn 1951). The plants inhabit
multiple environments, including temperate coastal regions, mountain ranges, semi-arid
habitats, and offshore islands (Thompson 2005). Most of the taxa are geographically
isolated from one another, though some have parapatric distributions and hybridize in
narrow regions where their distributions overlap (Streisfeld and Kohn 2005, 2007;
Streisfeld et al. 2013; Sobel and Streisfeld 2015; Stankowski et al. 2015; Stankowski and
Streisfeld 2015; Stankowski et al. 2017). Recent phylogenetic studies have confirmed the
monophyly of the radiation and revealed the basic relationships among its taxa
(Stankowski and Streisfeld 2015; Chase et al. 2017), making it an excellent system to
study how the signatures of linked selection evolve over time.
Using whole-genome sequencing and the first chromosome-level reference
assembly for the bush monkeyflower, we reveal heterogeneous patterns of lineage
sorting, diversity, and differentiation that have been shaped by variation in the intensity
of linked selection acting across the genome. By using the different taxon pairs as points
along a divergence continuum, we then show how these signatures of linked selection
evolve over the first 1.5 millions years of divergence. These results have important
implications for our understanding of the origins of biodiversity and the basic principles
governing molecular evolution.
127

Results and Discussion
A chromosome-level genome assembly, map, and annotation for the bush monkeyflower
To facilitate the analysis of genome-wide variation in this group, we constructed
the first chromosome-level reference genome for the bush monkeyflower using a
combination of long-read Single Molecule Real Time (SMRT) sequencing reads
(PacBio), overlapping and mate-pair short-reads (Illumina), and a high-density genetic
map (7,589 segregating markers across 10 linkage groups; Fig. S1; Table S1). Contig
building and scaffolding yielded 1,547 scaffolds, with an N50 size of 1,578 kbp, and a
total length of 207 Mbp. The high-density map allowed us to anchor and orient 94% of

the assembled genome onto 10 linkage groups, which is the number of chromosomes
inferred from karyotypic analyses in all subspecies of *M. aurantiacus* and *M. clevelandii*
(Vickery 1995). Analysis of assembly completeness based on conserved gene space
(Simao et al. 2015) revealed that 93% of 1400 universal single copy orthologous genes
were completely assembled (Table S2). Subsequent annotation yielded 23,018 predicted
genes.
143

144
145
Figure 1. Evolutionary relationships and patterns of genome-wide variation across the
radiation. A) The black tree was constructed from a concatenated alignment of genome-wide
SNPs and is rooted using *M. clevelandii*. The 387 gray trees were constructed from 500 kb
genomic windows. The first number associated with each node or taxon is the bootstrap support
for that clade in the whole genome tree, and the second number is the percentage of window-
based trees in which that clade is present. B) Levels of differentiation (F_{ST}), C) divergence (d_{xy}),
and D) diversity (π) within and among taxa based on the same 500 kb windows. For simplicity,
F_{ST} and d_{xy} are shown only for comparisons with the red ecotype of subspecies *puniceus*.
154

Variation in the extent of lineage sorting across the genome

As a first step toward understanding the processes that have shaped patterns of
genome-wide variation during this radiation, we inferred phylogenetic relationships
among its taxa. Rapid diversification is a hallmark of radiations and can result in
extensive phylogenetic discordance between genomic regions due to incomplete lineage
sorting (ILS) (Tajima 1983; Hudson 1990; Maddison 1997; Pease et al. 2016). To do this,
we sequenced 37 whole genomes from the seven subspecies and two ecotypes of
Mimulus aurantiacus (*n* = 4-5 per taxon) and its sister species *M. clevelandii* (*n* = 3) (Fig.
S2; Table S3). Close sequence similarity allowed us to align reads from all samples to the
reference assembly with high confidence (average 91.7% reads aligned; Table S3). After
mapping, we identified 13.2 million variable sites that were used in subsequent analyses
(average sequencing depth of 21x per individual, Table S3). Relationships were inferred

among the nine taxa using maximum-likelihood (ML) phylogenetic analysis (Stamatakis 2014) based on three different datasets: whole-genome concatenation and 500 kb and 100 kb non-overlapping genomic windows.

The tree topology obtained from the whole genome (concatenated) dataset (Fig. 1) confirmed the same phylogenetic relationships as previous analyses based on reduced-representation sequencing and five methods of phylogenetic reconstruction (Stankowski and Streisfeld 2015; Chase et al. 2017), and were supported by patterns of clustering from principal components analysis (Fig. S3). All seven subspecies formed monophyletic groups with 100% bootstrap support. Relationships within subspecies *puniceus* were more complex, as the red ecotype formed a monophyletic sub-clade within the paraphyletic yellow ecotype, reflecting the recent origin of red flowers from a yellow-flowered ancestor (Stankowski and Streisfeld 2015).

Although the whole genome phylogeny provides a well-supported summary of the relationships among the taxa, window-based analyses revealed extensive phylogenetic discordance at a finer genomic scale (Fig 1A). Despite four of the subspecies forming separate monophyletic groups in nearly all of the 387 window-based trees (500 kb scale; *grandiflorus* 99.7%, *aridus* 99.7%, *parviflorus* 94.1%, and *aurantiacus* 88.1%), only 22 (6%) trees showed the same taxon branching order as the whole-genome tree. While some of this discordance could be generated by gene flow after divergence (Lamichhaney et al. 2015; Stankowski and Streisfeld 2015; Pease et al. 2016; Richards and Martin 2017), our data indicate that the majority is due to incomplete lineage sorting (ILS). Specifically, higher levels of discordance were observed at nodes that were separated by shorter internode lengths ($r^2 = 0.94$, $p < 0.001$; Fig. S4). For example, even though subspecies *puniceus* was monophyletic in the majority of trees (72.6%), individuals from the red and yellow ecotypes only formed monophyletic groups in 27% and 9% of the trees, respectively. Similarly, the closely related subspecies *longiflorus* and *calycinus* were monophyletic in fewer trees than the other subspecies (45.7% and 24.1%, respectively). Thus, it appears that ILS is the primary source of phylogenetic conflict, especially near rapid divergence events, as predicted by theory (Pamilo and Nei 1988; Maddison 1997) and shown in other diverse radiations (Suh et al. 2015; Pease et al. 2016).

Next, we examined how the pattern of lineage sorting varied across the bush monkeyflower genome. Tree discordance due to the stochastic effects of neutral demography should be distributed uniformly across the genome (Hudson 1990). To test this prediction, we computed the correlation between the distance matrix generated from each window-based tree and the whole-genome tree, with a stronger relationship indicating that they are more similar (i.e., less ILS). Plotting this tree concordance score across the 10 linkage groups revealed a striking pattern (Fig. 2A; Fig. S5 for plots along each chromosome and S6 for results for 100 kb windows). Rather than being randomly distributed, trees with low concordance scores tended to cluster together in relatively narrow regions of all 10 chromosomes (Fig. 2A; autocorrelation analysis permutation tests $p = 0.023 - 0.001$; Fig. S7). This non-random pattern suggests that local rates of lineage sorting are determined by differences in the nature and/or strength of selection acting across the bush monkeyflower genome.

Figure 2. Common differentiation and diversity landscapes mirror variation in the local properties of the genome. A) Tree concordance scores for 500 kb non-overlapping genomic windows plotted across the 10 bush monkeyflower chromosomes. B – D) Plots of the *z*-transformed first principal component (PC1) for F_{ST}, d_{xy}, and π in overlapping 500 kb windows (step size = 50 kb). PC1 explains 66%, 70%, and 85% of the variation in F_{ST}, d_{xy} and π, respectively. E – F) Gene count and recombination rate (cM/Mbp) in overlapping 500 kb windows.

Patterns of genome-wide variation and lineage sorting have been shaped by recurrent linked selection

To gain deeper insight into the evolutionary processes that have shaped patterns of genome-wide variation, we used summary statistics in 500 kb windows to quantify patterns of differentiation (F_{ST}), divergence (d_{xy}), and diversity (π) among and within these taxa. The variation in F_{ST} among all 36 pairs of taxa highlights the continuous nature of differentiation across the group (Fig. 1B; Fig. S8), with mean window-based estimates ranging from 0.06 (red vs. yellow ecotypes of *puniceus*) to more than 0.70. Distributions of absolute divergence (d_{xy}) show a similar pattern (Fig. 1C), with mean values ranging from 0.54% (red vs. yellow ecotypes) to 1.6% (yellow ecotype vs. *M. clevelandii*). More strikingly, window-based estimates differ substantially among genomic regions, revealing heterogeneity in levels of differentiation and divergence. In addition, nucleotide diversity (π; Fig. 1D) ranged from 0.09% to 1.26% across genomic windows, but mean estimates were very similar among the ingroup taxa (0.37% to 0.53%) and were only slightly lower in *M. clevelandii* (0.26%).

As with tree concordance, these summary statistics showed non-random patterns of variation across broad regions of the genome ($p < 0.005$; Fig. 2; Fig. S5 Fig. S6; Fig. S7). To summarize variation in each statistic across all 36 pairwise comparisons between taxa (for d_{xy} and F_{ST}) or in the nine taxa (for π), we calculated the first principal component (PC1) for each genomic window. These analyses revealed that variation was highly correlated across genomic regions in all of the monkeyflower taxa. For F_{ST}, PC1

explained 65.9% of the variation across the 36 comparisons. Further, all comparisons
loaded positively onto PC1, indicating that peaks and troughs of F_{ST} tended to occur in
the same genomic regions in all comparisons (Fig. 2B, Table S4). Similarly, patterns of
genome-wide divergence (d_{xy}) and diversity (π) were highly correlated across
comparisons, with PC1 explaining 69.5% and 84.7% of the variation among the window-
based estimates, respectively (Fig 2C-D, Table S4).
Observing similar differentiation, diversity, and divergence 'landscapes' among
these taxa suggests that a common mechanism has been responsible for shaping patterns
of genome-wide variation across the radiation. Recent studies have observed correlated
genomic landscapes among related taxa, concluding that they were generated by a shared
pattern of heterogeneous linked selection (Poelstra et al. 2014; Burri et al. 2015; Van
Doren et al. 2017; Vijay et al. 2017). Indeed, if a region experiences a high level of
linked selection across the phylogenetic tree, then it will have both lower diversity (π)
within species and lower divergence (d_{xy}) between species, because divergence is
determined in part by levels of diversity in the common ancestor (Charlesworth et al.
1993; Charlesworth 1998; Burri 2017a, b). In agreement with this prediction, we
observed a strong positive correlation between PC1 d_{xy} and PC1 π ($r = 0.84$), indicating
that regions of the genome with lower diversity tended to be less diverged in all taxa (Fig.
3, Fig. S9 for scatterplots and Fig. S10 for results at 100 kb scale). Regions with reduced
diversity also tended to show higher differentiation (F_{ST}) ($r = -0.84$) and higher levels of
tree concordance ($r = -0.69$), both of which are predicted by models of linked selection
(Cruickshank and Hahn 2014).

Figure 3. Correlations reveal the impact of heterogeneous linked selection across the genome. Matrix of pairwise correlation coefficients between PC1 F_{ST}, PC1 d_{xy}, PC1 π, tree concordance, gene density, and recombination rate. The heat map and the shape of the ellipse indicate the strength of the correlation and its sign. All correlations are statistically significant at $p < 0.001$. Detailed scatterplots for each relationship can be found in Fig. S9. See Fig. S10 for a correlation matrix for 100 kb windows.

We next identified factors that cause variation in the intensity of linked selection
across the bush monkeyflower genome. Its intensity is determined by the local density of
targets of selection relative to the recombination rate (Maynard-Smith and Haigh 1974;
Charlesworth et al. 1993; Hudson and Kaplan 1995; Gillespie 2000). Specifically, a
higher intensity of linked selection is predicted in regions of the genome that are enriched
for functional elements, because mutations are more likely to have fitness consequences
if they arise in areas that are gene rich. The local recombination rate modulates this
effect, because regions unlinked to functional sites evolve independently of them.

To test these predictions, we used our annotated genome and genetic map to
calculate the number of protein coding genes and the average recombination rate
(cM/Mbp) in each 500 kb window (Fig. 2E-F; Fig. S5; Fig. S6). There was a strong
negative correlation between gene count and recombination rate (r = -0.40), leading to
large variation in the predicted strength of linked selection across the genome. In
addition, we observed strong correlations between PC1 π and both gene count (r = -0.84)
and recombination rate (r = 0.44; Fig. 3; Figs. S9 & S10), both of which indicate that
variation in the intensity of linked selection has shaped common patterns of diversity
across the genome.
Despite only having a direct estimate of gene density and recombination rate
variation from one subspecies (*puniceus*), the presence of a common diversity landscape
implies that the genomic distribution of these features has been conserved in all taxa after
being inherited from their common ancestor. This scenario is consistent with the recent
shared history of the group, and explains how a common pattern of heterogeneous linked
selection has become shared among them.
308
The signature of linked selection becomes stronger with increasing divergence time
Our analyses indicate that linked selection has shaped common patterns of
diversity and differentiation across this radiation. By using the different levels of
divergence between pairs of taxa, we next examined how these signatures evolved
through time. When a population fist splits, levels of diversity (π) and divergence (d_{xy})
are equal, so F_{ST} will be zero across the genome (Fig. S11 for cartoon explanation
assuming a simple model of allopatric divergence, no spatial structure, and large N_e). As
divergence proceeds, differentiation increases, but due to variation in the intensity of
linked selection across the genome, certain regions become differentiated before others.
During the early stages of divergence, lineage specific linked selection has impacted only
a small fraction of the genome, so patterns of differentiation should weakly mirror the
footprint of historical linked selection. However, as the divergence time increases, the
cumulative effects of recurrent linked selection should strengthen the relationships
between F_{ST} and π, gene density, and recombination rate. In contrast, the strength of the
correlations between π and gene density and recombination rate should remain similar
over time, because the shape of the diversity landscape is preserved by recurrent linked
selection despite new mutations arising in each lineage.
To test these predictions, we determined if the strength of the relationship
between the different statistics varied with the level of divergence between taxa. As
expected for a pair of taxa that recently split, the correlation between π and d_{xy} is almost
perfect between the least divergent pairs of taxa ($r \sim 1$), but the correlation decays over
time as ancestral variants fix and new mutations increase d_{xy} (Fig. 4A). Remarkably,
however, the strong correlations between π and gene density ($r \sim 0.8$) and π and
recombination rate ($r \sim 0.4$) barely change with increasing divergence time, as expected if
linked selection continues to act on the same regions in each taxon (Fig. 4B-C). By
contrast, the relationships between F_{ST} and levels of diversity, gene count, and
recombination rate all become stronger with increasing divergence time (Fig. 4D-F),
revealing the build up of heterogeneous differentiation due to the accumulating impact of
recurrent linked selection.
338

Figure 4. Time-course analysis reveals static and dynamic signatures of recurrent linked selection. Correlations between variables (500 kb windows) for all 36 taxonomic comparisons (gray dots) plotted against the average d_{xy} as a measure of divergence time. The top row shows how the relationships between π (each window averaged across a pair of taxa) and (A) d_{xy}, (B) gene count, and (C) recombination rate vary with increasing divergence time. The bottom row (D-E) shows the same relationships, but with F_{ST}. The regressions in each plot are fitted to the eight independent contrasts (colored points) obtained using a phylogenetic correction. The color gradient shows the strength of the correlation.

In addition to showing that the footprint of linked selection is dynamic, the sequence of divergence times provides empirical insight into when linked selection begins to shape patterns of differentiation, and how long it takes for this signature to develop (Fig. 5). Our data indicate that linked selection begins shaping patterns of differentiation at the early stages of population divergence. In population pairs with the most recent divergence times ($d_{xy} = 0.5\% - 0.7\%$), genome-wide patterns of F_{ST} are already correlated with variation in diversity, gene count, and recombination rate ($r \sim 0.4$, Fig. 4D-F). Indeed, these correlations are present between the parapatric red and yellow ecotypes of subspecies *puniceus*, which diverged recently with ongoing gene flow (Stankowski et al. 2015, 2017). This result provides empirical support for earlier theoretical studies indicating that linked selection can shape patterns of differentiation between spatially subdivided populations without a complete barrier to migration (Charlesworth et al. 1997). In contrast to these early time points, where the relationships between π and F_{ST} are moderate (Fig. 5; Fig. S12), the diversity and differentiation landscapes almost perfectly mirror one another in the most divergent comparisons ($r = -0.94$; (Fig. 5; Fig. S12). The build-up of this striking correlation in only 1.5 million years is a testament to linked selection's power to shape genome-wide patterns of variation, both within and between taxa.

Conclusions

Facilitated by the first chromosome-level genome assembly for the bush
monkeyflower, we show that linked selection has been a primary architect of the common
patterns of diversity, differentiation, and lineage sorting across this recent radiation.
Variation in the intensity of linked selection across the genome is conserved among these
taxa and is determined by the distribution of functional elements and variation in the local
recombination rate. By taking advantage of the range of divergence times between the
different pairs of monkeyflower taxa, we have shown how the signatures of linked
selection evolve as populations diverge: linked selection occurring within lineages acts to
preserve an ancestral pattern of diversity after a population split, while its joint action in
separate lineages causes a common differentiation landscape to emerge between them.

In addition to providing a dynamic picture of how the genomic landscape has
evolved over the first 1.5 millions years of divergence, our study has important
implications for the fields of molecular evolution and speciation. For example, even
though the impact of linked selection might be expected to vary across the tree of life due
to factors like differences in genome size, ploidy, mutation rate, and effective population
size (Slotte 2014), our findings support previous studies indicating that little of the
genome is free from its effects (Corbett-Detig et al. 2015; Casillas and Barbadilla 2017;
Kern and Hahn 2018). Second, this work has important implications for interpreting the
causes of heterogeneous genomic landscapes in light of the speciation process.
Specifically, we show that linked selection generates a heterogeneous differentiation
landscape shortly after a population split, which may confound efforts to identify regions

of the genome underlying reproductive isolation between even the youngest incipient
species (Ravinet et al. 2017). However, the common pattern of differentiation provides a
baseline for identifying highly differentiated regions that are unique to a given speciation
event. Thus, characterizing the signatures of linked selection appears to be a necessary
step in obtaining a comprehensive understanding of the processes that shaped patterns of
genome-wide variation between populations and species.
404
Materials and Methods
Genome assembly, high-density linkage map, and annotation
We used a combination of short-read Illumina and long-read Single Molecule,
Real Time (SMRT) sequencing to assemble the genome of a single individual from the
red ecotype of *M. aurantiacus* subspecies *puniceus* (Table S2 for sample collection
location). To assemble resulting scaffolds into pseudomolecules, we generated a high-
density linkage map from an outbred F_2 mapping population (Table S2 for sample
collection locations). Restriction-site associated DNA sequencing (RADseq) was used to
genotype parents and 269 offspring. Map construction was performed using *Lep-MAP2*
(Rastas et al. 2016). After integrating the assembly and genetic map, we made corrections
to the map order using the physical position of markers within the assembled scaffolds.
Genome annotation was conducted using the MAKER pipeline (Holt and Yandell 2011).
See supplementary methods for more details.
418
419 *Genome re-sequencing and variant calling*
420 DNA was extracted from 37 individuals (Table S2), and Illumina sequencing was
421 performed using paired-end 150 bp reads. Raw reads were aligned to the reference
422 assembly, and variant calling was executed with the GATK pipeline (McKenna et al.
423 2010) using the UnifiedGenotyper tool. See supplementary methods for more details.
424
Phylogenetic Analyses
We used *RAxML* v8 to construct genome-wide phylogenies from a concatenated
alignment of all variable sites and from genomic windows. We tested for a relationship
between node concordance (the number of 500 kb window-based trees that recovered a
given node from the genome-wide tree) and internode length using the internode
distances from the genome-wide analysis. Tree concordance scores were generated from
the correlation between the distance matrix from each window-based tree and the whole-
genome tree. Autocorrelation coefficients for tree concordance scores were calculated in
R using custom scripts, and their significance was tested from 1000 random permutations
of the genome-wide data. See supplementary methods for more details.
435
Population genomic analyses
Estimates of nucleotide diversity (π), differentiation (F_{ST}), and divergence (d_{XY})
for 100 kb and 500 kb windows were calculated using Python scripts downloaded from
https://github.com/simonhmartin/genomics_general. Principal components analysis was
used to summarize patterns of variation in these statistics across all taxa (for π) and taxon
comparisons (for F_{ST} and d_{XY}). Average recombination rate (cM/Mbp) for each window
was estimated as the average value across three or more adjacent pairs of mapped
markers in each genomic window. Gene density was estimated as the number of

predicted genes in each window. We used linear regression to test if the strength of the
correlations between different statistics changed with the level of divergence using a set
of eight statistically independent contrasts (Felsenstein 1985; Coyne and Orr 1989). See
supplementary methods for details.
448

Data Accessibility

450 Raw sequencing reads used for the genome assembly, linkage map construction,
451 and genome resequencing and are available on the Short-Read Archive (SRA) under the
452 bioproject ID xxx. The genetic map, annotation, reference genome sequence, and VCF
453 file have been deposited on DRYAD.
454

Acknowledgments

We thank Bill Cresko, Thomas Nelson and Martin Garlovsky for advice and
stimulating discussion. John Willis performed the Illumina mate-pair library preps used
for the genome assembly. Julian Catchen, Clay Small, Susan Bassham, and Janna Fierst
provided technical advice. Doug Turnbull and Maggie Weitzman conducted the Illumina
sequencing at the University of Oregon Genomics Core facility. Funding was provided
by the National Science Foundation grant DEB-1258199 to MS and the Sloan Foundation
to PR.
463

References

465
Begun, D. J., A. K. Holloway, K. Stevens, L. W. Hillier, Y. P. Poh, M. W. Hahn, P. M.
Nista, C. D. Jones, A. D. Kern, C. N. Dewey, L. Pachter, E. Myers, and C. H.
Langley. 2007. Population genomics: Whole-genome analysis of polymorphism
and divergence in Drosophila simulans. Plos Biol 5:2534-2559.
Burri, R. 2017a. Dissecting differentiation landscapes: a linked selection's perspective. J
Evolution Biol 30:1501-1505.
Burri, R. 2017b. Linked selection, demography and the evolution of correlated genomic
landscapes in birds and beyond. Mol Ecol 26:3853-3856.
Burri, R., A. Nater, T. Kawakami, C. F. Mugal, P. I. Olason, L. Smeds, A. Suh, L. Dutoit,
S. Bures, L. Z. Garamszegi, S. Hogner, J. Moreno, A. Qvarnstrom, M. Ruzic, S.
A. Saether, G. P. Saetre, J. Torok, and H. Ellegren. 2015. Linked selection and
recombination rate variation drive the evolution of the genomic landscape of
differentiation across the speciation continuum of Ficedula flycatchers. Genome
Res 25:1656-1665.
Casillas, S., and A. Barbadilla. 2017. Molecular Population Genetics. Genetics 205:1003-
1035.
Charlesworth, B. 1998. Measures of divergence between populations and the effect of
forces that reduce variability. Mol Biol Evol 15:538-543.
Charlesworth, B., and D. Charlesworth. 2018. Neutral variation in the context of
selection. Mol Biol Evol.
Charlesworth, B., M. T. Morgan, and D. Charlesworth. 1993. The effect of deleterious
mutations on neutral molecular variation. Genetics 134:1289-1303.
Charlesworth, B., M. Nordborg, and D. Charlesworth. 1997. The effects of local
selection, balanced polymorphism and background selection on equilibrium

patterns of genetic diversity in subdivided populations. Genetical research 70:155-
174.
Chase, M. A., S. Stankowski, and M. A. Streisfeld. 2017. Genomewide variation provides
insight into evolutionary relationships in a monkeyflower species complex
(Mimulus sect. Diplacus). Am J Bot 104:1510-1521.
Coop, G., and P. Ralph. 2012. Patterns of neutral diversity under general models of
selective sweeps. Genetics 192:205-224.
Corbett-Detig, R. B., D. L. Hartl, and T. B. Sackton. 2015. Natural selection constrains
neutral diversity across a wide range of species. Plos Biol 13:e1002112.
Coyne, J. A., & Orr, H. A. 1989. Patterns of speciation in *Drosophila. Evolution*, 43:362-
381.
Cruickshank, T. E., and M. W. Hahn. 2014. Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. Mol Ecol 23:3133-
3157.
Cutter, A. D., and B. A. Payseur. 2013. Genomic signatures of selection at linked sites:
unifying the disparity among species. Nature reviews. Genetics 14:262-274.
Felsenstein, J. 1985. Phylogenies and the comparative method. *The American*
Naturalist, *125*: 1-15.
Gillespie, J. H. 1994. The Causes of Molecular Evolution. Oxford University Press On
Demand.
Gillespie, J. H. 2000. Genetic drift in an infinite population: The pseudohitchhiking
model. Genetics 155:909-919.
Hahn, M. W. 2008. Toward a selection theory of molecular evolution. Evolution 62:255-
265.
Holt, C., and M. Yandell. 2011. MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. BMC bioinformatics
12:491.
Hudson, R. R. 1990. Gene genealogies and the coalescent process. Oxford Surveys in
Evolutionary Biology 7:44.
Hudson, R. R., and N. L. Kaplan. 1995. Deleterious Background Selection with
Recombination. Genetics 141:1605-1617.
Kern, A. D., and M. W. Hahn. 2018. The neutral theory in light of natural selection. Mol
Biol Evol.
Kimura, M. 1968. Evolutionary rate at the molecular level. Nature 217:624-626.
Kimura, M. 1986. DNA and the neutral theory. Philosophical transactions of the Royal
Society of London. Series B, Biological sciences 312:343-354.
Kreitman, M. 1996. The neutral theory is dead. Long live the neutral theory. BioEssays :
news and reviews in molecular, cellular and developmental biology 18:678-683;
discussion 683.
Lamichhaney, S., J. Berglund, M. S. Almen, K. Maqbool, M. Grabherr, A. Martinez-
Barrio, M. Promerova, C. J. Rubin, C. Wang, N. Zamani, B. R. Grant, P. R.
Grant, M. T. Webster, and L. Andersson. 2015. Evolution of Darwin's finches and
their beaks revealed by genome sequencing. Nature 518:371-375.
Langley, C. H., and W. M. Fitch. 1974. An examination of the constancy of the rate of
molecular evolution. Journal of molecular evolution 3:161-177.

Langley, C. H., K. Stevens, C. Cardeno, Y. C. Lee, D. R. Schrider, J. E. Pool, S. A.
Langley, C. Suarez, R. B. Corbett-Detig, B. Kolaczkowski, S. Fang, P. M. Nista,
A. K. Holloway, A. D. Kern, C. N. Dewey, Y. S. Song, M. W. Hahn, and D. J.
Begun. 2012. Genomic variation in natural populations of Drosophila
melanogaster. Genetics 192:533-598.
Lewontin, R. C. 1974. The Genetic Basis of Evolutionary Change. Columbia University
Press, New York.
Lewontin, R. C., and J. L. Hubby. 1966. A Moleuclar Approach to Study of Genic
Heterozygosity in Natural Populations .2. Amount of Variation and Degree of
Heterozygosity in Natural Populations of Drosophila Pseudoobscura. Genetics
54:595-&.
Maddison, W. P. 1997. Gene trees in species trees. Syst Biol 46:523-536.
Maynard-Smith, J., and J. Haigh. 1974. Hitch-Hiking Effect of a Favorable Gene. Genet
Res 23:23-35.
McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K.
Garimella, D. Altshuler, S. Gabriel, M. Daly, and M. A. DePristo. 2010. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-
generation DNA sequencing data. Genome Res 20:1297-1303.
McMinn, H. E. 1951. Studies in the genus *Diplacus*, Scrophulariaceae. Madrono 11:33-
128.
Ohta, T. 1973. Slightly Deleterious Mutant Substitutions in Evolution. Nature 246:96-98.
Pamilo, P., and M. Nei. 1988. Relationships between Gene Trees and Species Trees. Mol
Biol Evol 5:568-583.
Pease, J. B., D. C. Haak, M. W. Hahn, and L. C. Moyle. 2016. Phylogenomics Reveals
Three Sources of Adaptive Variation during a Rapid Radiation. Plos Biol 14.
Pease, J. B., and M. W. Hahn. 2013. More Accurate Phylogenies Inferred from Low-
Recombination Regions in the Presence of Incomplete Lineage Sorting. Evolution
67:2376-2384.
Poelstra, J. W., N. Vijay, C. M. Bossu, H. Lantz, B. Ryll, I. Muller, V. Baglione, P.
Unneberg, M. Wikelski, M. G. Grabherr, and J. B. W. Wolf. 2014. The genomic
landscape underlying phenotypic integrity in the face of gene flow in crows.
Science 344:1410-1414.
Rastas, P., F. C. F. Calboli, B. C. Guo, T. Shikano, and J. Merila. 2016. Construction of
Ultradense Linkage Maps with Lep-MAP2: Stickleback F-2 Recombinant Crosses
as an Example. Genome Biol Evol 8:78-93.
Ravinet, M., R. Faria, R. K. Butlin, J. Galindo, N. Bierne, M. Rafajlovic, M. A. F. Noor,
B. Mehlig, and A. M. Westram. 2017. Interpreting the genomic landscape of
speciation: a road map for finding barriers to gene flow. J Evolution Biol
30:1450-1477.
Richards, E. J., and C. H. Martin. 2017. Adaptive introgression from distant Caribbean
islands contributed to the diversification of a microendemic adaptive radiation of
trophic specialist pupfishes. Plos Genet 13.
Simao, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov.
2015. BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. Bioinformatics 31:3210-3212.

Slotte, T. 2014. The impact of linked selection on plant genomic variation. Brief Funct
Genomics 13:268-275.
Sobel, J. M., and M. A. Streisfeld. 2015. Strong premating reproductive isolation drives
incipient speciation in Mimulus aurantiacus. Evolution 69:447-461.
Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. Bioinformatics 30:1312-1313.
Stankowski, S., J. M. Sobel, and M. A. Streisfeld. 2015. The geography of divergence
with gene flow facilitates multitrait adaptation and the evolution of pollinator
isolation in Mimulus aurantiacus. Evolution 69:3054-3068.
Stankowski, S., J. M. Sobel, and M. A. Streisfeld. 2017. Geographic cline analysis as a
tool for studying genome-wide variation: a case study of pollinator-mediated
divergence in a monkeyflower. Mol Ecol 26:107-122.
Stankowski, S., and M. A. Streisfeld. 2015. Introgressive hybridization facilitates
adaptive divergence in a recent radiation of monkeyflowers. P Roy Soc B-Biol
Sci 282:154-162.
Streisfeld, M. A., and J. R. Kohn. 2005. Contrasting patterns of floral and molecular
variation across a cline in Mimulus aurantiacus. Evolution 59:2548-2559.
Streisfeld, M. A., and J. R. Kohn. 2007. Environment and pollinator-mediated selection
on parapatric floral races of Mimulus aurantiacus. J Evolution Biol 20:122-132.
Streisfeld, M. A., W. N. Young, and J. M. Sobel. 2013. Divergent Selection Drives
Genetic Differentiation in an R2R3-MYB Transcription Factor That Contributes
to Incipient Speciation in Mimulus aurantiacus. Plos Genet 9.
Suh, A., L. Smeds, and H. Ellegren. 2015. The Dynamics of Incomplete Lineage Sorting
across the Ancient Adaptive Radiation of Neoavian Birds. Plos Biol 13.
Tajima, F. 1983. Evolutionary Relationship of DNA-Sequences in Finite Populations.
Genetics 105:437-460.
Thompson, D. M. 2005. Systematics of Mimulus subgenus Schizoplacus
(Scrophulariaceae). Systematic Botany Monographs 75:1-213.
Van Doren, B. M., L. Campagna, B. Helm, J. C. Illera, I. J. Lovette, and M. Liedvogel.
2017. Correlated patterns of genetic diversity and differentiation across an avian
family. Mol Ecol 26:3982-3997.
Vickery, R. K. 1995. Speciation by Aneuploidy and Polyploidy in Mimulus
(Scrophulariaceae). Great Basin Nat 55:174-176.
Vijay, N., M. Weissensteiner, R. Burri, T. Kawakami, H. Ellegren, and J. B. W. Wolf.
2017. Genomewide patterns of variation in genetic diversity are shared among
populations, species and higher-order taxa. Mol Ecol 26:4284-4295.

Supplementary information for:

The tempo of linked selection: emergence of a heterogeneous genomic landscape during a recent radiation of monkeyflowers

Sean Stankowski[1,2,4], Madeline A. Chase[1,3,4] Allison Fuiten[1], Peter L. Ralph[1], and Matthew A. Streisfeld[1*].

[1]Institute of Ecology and Evolution, 335 Pacific Hall 5289, University of Oregon, Eugene, OR 97405, USA

[2]Present address: Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

[3]Present address: Department of Evolutionary Biology, Evolutionary Biology Centre (EBC), Uppsala University, Uppsala, Sweden

[4]co first authors

*Corresponding author: mstreis@uoregon.edu

Supplementary Materials and Methods

Genome Assembly

We used a combination of short-read Illumina and long-read Single Molecule, Real Time (SMRT) sequencing to assemble the genome of a single individual from the red ecotype of *M. aurantiacus* subspecies *puniceus* (population UCSD; Table S1). Genomic DNA was isolated from leaf tissue using either ZR plant/seed DNA miniprep kits (Zymo Research) or GeneJet Plant Genomic DNA purification kits (Thermo Fisher). Illumina libraries were generated following the *Allpaths-LG* assembly pipeline (Gnerre et al. 2011), which included a single fragment library with average 180 bp insert size and three mate pair libraries (average insert sizes: 3.5-5 kb, 5-7 kb, and 7-13 kb). Libraries were sequenced on the Illumina HiSeq 2500 using paired-end 100 bp reads. An initial scaffold-level assembly was performed with *Allpaths-LG* using default parameters and the *haploidify* function enabled. This assembly yielded 11,123 contigs (N50 = 40.5 kb) and 2,299 scaffolds (N50 = 1,310 kb), for a total assembly size of 193.3 Mbp. Long-read sequencing was performed from the same individual using 12 SMRT cells sequenced on the Pacific Biosystems RS II machine at Duke University. We obtained a total of 6.4 Gbp of sequence, which corresponds to ~21 × coverage of the genome. The PacBio reads were used to re-scaffold the *Allpaths-LG* scaffolds using *Opera-LG* (Gao et al. 2016). This reduced the number of scaffolds to 1,547 (N50 = 1,578 kb).

We then manually improved the scaffold containing the flower color gene *MaMyb2* (Streisfeld et al. 2013). We first aligned this scaffold to a previously published draft sequence assembly from this same individual (Stankowski et al. 2017), which was generated using Illumina short-reads and the *Velvet* assembler (Zerbino and Birney 2008). We used long range PCR and cloning to generate Sanger sequences across three regions within 20 kb of *MaMyb2* that did not assemble well. Genomic DNA was amplified using Phusion high fidelity polymerase (NEB). PCR products were cloned into the pCR2.1 TOPO-TA vector (Life Technologies), and purified plasmids were sequenced with Sanger technology. Resulting sequences were aligned to the scaffold containing *MaMyb2*, and new PCR primers were designed to sequence internal fragments until the entire insert was sequenced. Using this approach, we sequenced a total of 9,824 bp across the three regions. The reference sequence in the assembly was corrected manually to match the Sanger data.

Finally, we gap filled the assembly using the PacBio data and the program *PBJelly* (English et al. 2012). Resulting scaffolds were assembled into pseudomolecules using *Chromonomer* (http://catchenlab.life.illinois.edu/chromonomer/), according to the online manual. This software anchored and oriented scaffolds based on the order of markers in a high-density linkage map (see below) and made corrections to scaffolds when differences occurred between the genetic and physical positions of markers in the map. A final round of gap filling with *PBJelly* was performed to fill any gaps that were created by broken scaffolds in *Chromonomer*. To assess the completeness of the gene space in the assembly, we used both the BUSCO and CEGMA pipelines to estimate the proportion of 956 single copy plant genes (BUSCO) or 248 core eukaryotic genes (CEGMA) that were completely or partially assembled (Parra et al. 2007; Simao et al. 2015). The proportion of these genes present in an assembly has been shown to be correlated with the total proportion of assembled gene space, and thus serves as a good predictor of assembly completeness.

Construction of high-density linkage map

We generated an outbred F_2 mapping population by crossing two F_1 individuals, each the product of crosses between different greenhouse-raised red and yellow ecotype plants collected from one red ecotype and one yellow ecotype population (populations UCSD and LO, respectively; Table S1). We then used restriction-site associated DNA sequencing (RADseq) to genotype F_1 and F_2 individuals. DNA was extracted from leaf material using Zymo ZR plant/seed DNA miniprep kits, and RAD library preparation followed the protocol outlined in Sobel and Streisfeld (Sobel and Streisfeld 2015). Libraries were sequenced on the Illumina HiSeq 2000 platform using single-end 100 bp reads at the Genomics Core Facility, University of Oregon.

Reads were filtered based on quality, and errors in the barcode sequence or RAD site were corrected using the *process_radtags* script in *Stacks v. 1.35* (Catchen et al. 2011; Catchen et al. 2013). Loci were created using the *denovo_map.pl* function of *Stacks*, with three identical raw reads required to create a stack, two mismatches allowed between loci for an individual, and two mismatches allowed when processing the catalog. Single nucleotide polymorphisms (SNPs) were determined and genotypes called using a maximum-likelihood (ML) statistical model implemented in *Stacks* and a stringent χ^2 significance level of 0.01 to distinguish between heterozygotes and homozygotes. We then used the *genotypes* program implemented in *Stacks* to identify a set of 9,029 mappable markers. We specified a 'CP' cross design (F_1 individuals coded as the parents), requiring that a marker was present in at least 85% of progeny at a minimum depth of 12 reads per individual, and we allowed automated corrections to be made to the data.

Linkage map construction was performed using *Lep-MAP2* (Rastas et al. 2016). The data were filtered using the *Filtering* module to include only individuals with less than 15% missing data and excluded markers that showed evidence for extreme segregation distortion (χ^2 test, $P < 0.01$). To assign markers to linkage groups, we used the *SeparateChromosomes* module with a logarithm of odds (LOD) score limit of 20 and no minimum size for linkage groups (LG). This assigned 7,217 markers to 10 linkage groups, which matches the number of chromosomes in *M. aurantiacus*. The *JoinSingles* module was executed again with a LOD limit of 10 to join an additional 877 ungrouped markers to the 10 previously formed LGs. Fifty-seven singles that were not joined at this stage were discarded from the dataset. Initial marker orders were determined using sex-averaged and sex-specific recombination rates using the *OrderMarkers* module. For each LG, we conducted 10 independent runs using the Kosambi mapping function (*useKosambi=1*), with the dataset split into seven pseudofamilies to take advantage of parallel processing. When multiple markers had identical genotypes, only the duplicate marker with the least missing data was used in marker ordering. We retained the marker order from the run with the best likelihood. After removing markers with an error rate > 0.05, the ML order was re-evaluated using the *evaluateOrder* flag. The map contained 8,094 informative loci from 269 F2 individuals, with an average of 3.5% ± SD 3.86 missing data per individual.

After the integration of our assembly and genetic map using the *Chromonomer* software (Amores et al. 2014), we made corrections to the map order based on the physical position of markers within assembled scaffolds. Using the output of

Chromonomer, we identified markers that were out of order in the map compared to their local assembly order and aligned these markers to the assembly from *Chromonomer* using *Bowtie2* v. 2.2.5 (Langmead and Salzberg 2012) with the *very_sensitive* settings to obtain their physical order. We then re-estimated the map using the *evaluateOrder* flag in *Lep-MAP2* as described above, but with the marker order constrained to the physical order (*improveOrder=0*) and with all duplicate markers included in the analysis (*removeDuplicates=0*). After initial map construction, we removed 17 markers with an estimated error rate greater than 5% and estimated the map one last time using the same settings. The final map contained 7,589 markers across the 10 linkage groups.

Genome annotation
Prior to genome annotation, the assembly was soft-masked for repetitive elements and areas of low complexity with *RepeatMasker* (RepeatMasker Open-4.0) using a custom *Mimulus aurantiacus* library created by *RepeatModeler* (RepeatModeler Open-1.0), Repbase repeat libraries (Jurka et al. 2005), and a list of known transposable elements provided by *MAKER* (Holt and Yandell 2011). In total, 30.99% of the genome assembly was masked by *RepeatMasker*. Repetitive elements were annotated with *RepeatModeler*. Hidden Markov Models for gene prediction were generated by *SNAP* (Korf 2004) and *Augustus* (Stanke and Waack 2003)and were trained iteratively to the assembly using *MAKER*, as described by Cantarel et al. (Cantarel et al. 2008). Training was performed on the 14.5 Mbp sequence from LG9. Evidence used by *MAKER* for annotation included protein sequences from *Arabidopsis thaliana*, *Oryza sativa*, *Solanum lycopersicum*, *Solanum tuberosum*, *Daucus carota*, *Vitis vinifera* (all downloaded from EnsemblPlants on 9 August 2016), *Salvia miltiorrhiza* (downloaded from Herbal Medicine Omics Database on 9 August 2016), *Mimulus guttatus v. 2* (downloaded from JGI Genome Portal on 9 August 2016), and all Uniprot/swissprot proteins (downloaded on 18 August 2016) (Goodstein et al. 2012; Nordberg et al. 2013; Kersey et al. 2016)(Herbal Medicine Omics Database; Uniprot). We filtered the annotations with *MAKER* to include: 1) only evidence-based information that also contained assembled protein support, and 2) those *ab initio* gene predictions that did not overlap with the evidence-based annotations and that contained protein family domains, as detected with InterProScan (Quevillon et al. 2005).

Genome re-sequencing and variant calling
We collected leaf tissue from four to five individuals from seven subspecies of *M. aurantiacus*, including both ecotypes of subspecies *puniceus* (Table S3; Fig. S2). In addition, we collected leaf tissue from three individuals of *M. clevelandii*. We extracted DNA from dried tissue using the Zymo Plant/Seed MiniPrep DNA kit following the manufacturer's instructions. We prepared sequencing libraries using the Kapa Biosystems HyperPrep kit, and libraries with an insert size between 400-600 bp were sequenced on the Illumina HiSeq 4000 using paired-end 150 bp reads at the Genomics Core Facility, University of Oregon.

We filtered raw reads using the *process_shortreads* script in *Stacks* v1.46 to remove reads with uncalled bases or poor quality scores. We then aligned the retained reads to the reference assembly using the BWA-MEM algorithm in *BWA* v0.7.15 (Li 2013). An average of 91.7% of reads aligned (range: 82.6-96.0%), and the average

sequencing depth was 21x (range: 15.16 – 30.86x). We then marked PCR duplicates using *Picard* (http://broadinstitute.github.io/picard). We performed an initial run of variant calling using the UnifiedGenotyper tool in *GATK* v3.8 (McKenna et al. 2010) and filtered the data to remove variants with a mapping quality < 50, a quality depth < 4, and a Fisher Strand score > 50. We then used these variants to perform base quality score recalibration for each individual, before performing another run of the UnifiedGenotyper to call final variants. After the second run of variant calling, we removed variants with a mapping quality < 40, a quality depth < 2, and a Fisher Strand score > 60. The final dataset contained 13,233,829 SNPs across the nine taxa. Finally, we ran UnifiedGenotyper with the EMIT_ALL_SITES option to output all variant and invariant genotyped sites.

Phylogenetic analyses
Initially, we used *RAxML* v8 to reconstruct the evolutionary relationships among the nine taxa by concatenating variant sites from across the genome. To investigate patterns of phylogenetic discordance across the genome, we also built trees from windows across the genome. We phased SNPs using *BEAGLE* v4.1 (Browning and Browning 2007), using a window size of 100 kb and an overlap of 10 kb. We incorporated information on recombination rate from the genetic map and did not impute missing genotypes. After phasing, we used *MVFtools* (https://www.github.com/jbpease/mvftools) to run *RAxML* from 100 kb and 500 kb nonoverlapping windows, with the *M. clevelandii* samples set as the outgroups. We then visualized the window trees in *DensiTree* v2.01 (Bouckaert 2010).

To assess concordance between the window-based trees and the whole-genome tree, we converted trees to distance matrices using the *Ape* package in R (Paradis et al. 2004). We then calculated the Pearson's correlation coefficient between the distance matrix from each window and the whole-genome tree, with a stronger correlation indicating higher concordance with the whole-genome tree. We used one-dimensional autocorrelation analysis to determine if variation in tree concordance was randomly distributed across the genome. This involved estimating the autocorrelation between genomic position and tree concordance for each LG with a lag size of 2 Mbp. The significance of the observed value for each LG was determined from a null distribution of autocorrelation coefficients estimated from 1000 random permutations of the genome-wide data.

We also conducted a Principal Components Analysis (PCA) based on all variant sites from across the entire genome using *Plink* v. 1.90 (Chang et al. 2015). Initially, we ran the PCA with all 37 samples, but we consecutively re-ran the analysis by removing different taxa in order to assess clustering patterns among more closely related samples.

Population genomic analyses
To examine how genome-wide patterns of diversity, differentiation, and divergence varied among taxa, we calculated within-taxon nucleotide diversity (π), between-taxon relative differentiation (F_{ST}), and between-taxon absolute divergence (d_{xy}) across non-overlapping 100 kb and 500 kb windows using custom Python scripts downloaded from https://github.com/simonhmartin/genomics_general. We calculated measures of differentiation and divergence across all 36 pairwise comparisons among the nine taxa,

and diversity was estimated separately for each taxon. These scripts estimated π and d_{xy} by dividing the number of sequence differences within a window (either within or between taxa) by the total number of sites in that window. To account for missing data, the script counted the number of differences between each sample, divided by the total number of variant sites that were genotyped within those samples, and then averaged across all pairs of samples. To provide an unbiased estimate of diversity and divergence, we incorporated invariant sites into the calculation by dividing the number of pairwise differences (within and between taxa, respectively) by the total number of genotyped sites (variant and invariant) within a window. F_{ST} was calculated following the measure of K_{ST} (Hudson et al. 1992), equation 9), but was modified to incorporate missing data using the same approach as π and d_{xy}. We filtered the data separately for each taxonomic comparison, so that each site was required to be genotyped in at least three individuals for comparisons within the *M. aurantiacus* complex or at least two individuals for each comparison involving *M. clevelandii.*

We summarized the variation in each statistic across comparisons using a Principal Components Analysis (PCA), with taxon or taxon pair as the variables. Thus, across each window, the first principal component of π, F_{ST}, and d_{xy} provided multivariate measures that explained the greatest covariance in the data. We used a one-dimensional autocorrelation analysis and permutation test to determine if the genome-wide patterns of PC1 π, F_{ST}, and d_{xy} departed from a random expectation, as described above for tree concordance (see section 'phylogenetic analyses').

To examine the relationships among PC1 diversity, differentiation, and divergence, we estimated Pearson's correlation coefficient among all three statistics across genomic windows. Further, we estimated correlations among these three statistics and tree concordance, gene density, and recombination rate. Recombination rate was estimated by comparing the genetic and physical distance (in cM/Mbp) between all pairs of adjacent markers on each LG from the genetic linkage map described above. We removed the top 5% of recombination rates, as these represented unrealistically high values of recombination. A minimum of three estimates was required to obtain an average recombination rate estimate within each window. Gene density was calculated from the number of predicted genes in each window, as determined from the annotation described above.

To determine how the correlations among the statistics (diversity, differentiation, divergence, recombination rate, gene count) changed with increasing divergence time, we examined the correlation coefficient among all pairs of statistics individually for each of the 36 pairwise comparisons. Because diversity was measured within taxa rather than between them, we calculated the mean value of π between each pair of taxa. Also, because many of the pairwise comparisons are non-independent, we applied the phylogenetic correction outlined by (Felsenstein 1985; Coyne and A. 1989) to produce a statistically independent set of data points for this analysis.

As a measure of the divergence time between *M. clevelandii* and *M. aurantiacus*, we estimated the percent sequence divergence (d_{xy}) between individuals of *M. clevelandii* and all subspecies of *M. aurantiacus* combined. We then converted this value into a divergence time T (in generations) using the equation: $T = d_{xy}/(2\mu)$, where μ is the mutation rate, 1.5×10^{-8} (Koch et al. 2001; Brandvain et al. 2014). This value was then converted into years by multiplying by a generation time of two years.

Supplemental References

Amores, A., J. Catchen, I. Nanda, W. Warren, R. Walter, M. Schartl, and J. H. Postlethwait. 2014. A RAD-tag genetic map for the platyfish (Xiphophorus maculatus) reveals mechanisms of karyotype evolution among teleost fish. Genetics 197:625-641.

Bouckaert, R. R. 2010. DensiTree: making sense of sets of phylogenetic trees. Bioinformatics 26:1372-1373.

Brandvain, Y., A. M. Kenney, L. Flagel, G. Coop, and A. L. Sweigart. 2014. Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. Plos Genet 10.

Browning, S. R., and B. L. Browning. 2007. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. Am J Hum Genet 81:1084-1097.

Cantarel, B. L., I. Korf, S. M. C. Robb, G. Parra, E. Ross, B. Moore, C. Holt, A. S. Alvarado, and M. Yandell. 2008. MAKER: An easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res 18:188-196.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. Stacks: an analysis tool set for population genomics. Mol Ecol 22:3124-3140.

Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait. 2011. Stacks: Building and Genotyping Loci De Novo From Short-Read Sequences. G3-Genes Genom Genet 1:171-182.

Chang, C. C., C. C. Chow, L. C. A. M. Tellier, S. Vattikuti, S. M. Purcell, and J. J. Lee. 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4.

Coyne, J. A., and O. H. A. 1989. Patterns of speciation in Drosophila. Evolution 43:362-381.

English, A. C., S. Richards, Y. Han, M. Wang, V. Vee, J. Qu, X. Qin, D. M. Muzny, J. G. Reid, K. C. Worley, and R. A. Gibbs. 2012. Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. PloS one 7:e47768.

Felsenstein, J. 1985. Phylogenies and the comparative method. American Naturalist 125:1-15.

Gao, S., D. Bertrand, B. K. Chia, and N. Nagarajan. 2016. OPERA-LG: efficient and exact scaffolding of large, repeat-rich eukaryotic genomes with performance guarantees. Genome biology 17:102.

Gnerre, S., I. Maccallum, D. Przybylski, F. J. Ribeiro, J. N. Burton, B. J. Walker, T. Sharpe, G. Hall, T. P. Shea, S. Sykes, A. M. Berlin, D. Aird, M. Costello, R. Daza, L. Williams, R. Nicol, A. Gnirke, C. Nusbaum, E. S. Lander, and D. B. Jaffe. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. Proceedings of the National Academy of Sciences of the United States of America 108:1513-1518.

Goodstein, D. M., S. Shu, R. Howson, R. Neupane, R. D. Hayes, J. Fazo, T. Mitros, W. Dirks, U. Hellsten, N. Putnam, and D. S. Rokhsar. 2012. Phytozome: a comparative platform for green plant genomics. Nucleic acids research 40:D1178-1186.

Holt, C., and M. Yandell. 2011. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. BMC bioinformatics 12:491.

Hudson, R. R., D. D. Boos, and N. L. Kaplan. 1992. A Statistical Test for Detecting Geographic Subdivision. Mol Biol Evol 9:138-151.

Jurka, J., V. V. Kapitonov, A. Pavlicek, P. Klonowski, O. Kohany, and J. Walichiewicz. 2005. Repbase Update, a database of eukaryotic repetitive elements. Cytogenetic and genome research 110:462-467.

Kersey, P. J., J. E. Allen, I. Armean, S. Boddu, B. J. Bolt, D. Carvalho-Silva, M. Christensen, P. Davis, L. J. Falin, C. Grabmueller, J. Humphrey, A. Kerhornou, J. Khobova, N. K. Aranganathan, N. Langridge, E. Lowy, M. D. McDowall, U. Maheswari, M. Nuhn, C. K. Ong, B. Overduin, M. Paulini, H. Pedro, E. Perry, G. Spudich, E. Tapanari, B. Walts, G. Williams, M. Tello-Ruiz, J. Stein, S. Wei, D. Ware, D. M. Bolser, K. L. Howe, E. Kulesha, D. Lawson, G. Maslen, and D. M. Staines. 2016. Ensembl Genomes 2016: more genomes, more complexity. Nucleic acids research 44:D574-580.

Koch, M., B. Haubold, and T. Mitchell-Olds. 2001. Molecular systematics of Brassicaceae: evidence from plastidic matK and nuclear Chs sequences. Am J Bot 88:534-544.

Korf, I. 2004. Gene finding in novel genomes. BMC bioinformatics 5:59.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nature methods 9:357-359.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv 1303.3997v1.

Nordberg, H., M. Cantor, S. Dusheyko, S. Hua, A. Poliakov, I. Shabalov, T. Smirnova, I. V. Grigoriev, and I. Dubchak. 2013. The genome portal of the Department of Energy Joint Genome Institute: 2014 updates. Nucleic acids research 42:D26-D31.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of Phylogenetics and Evolution in R language. Bioinformatics 20:289-290.

Parra, G., K. Bradnam, and I. Korf. 2007. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genornes. Bioinformatics 23:1061-1067.

Quevillon, E., V. Silventoinen, S. Pillai, N. Harte, N. Mulder, R. Apweiler, and R. Lopez. 2005. InterProScan: protein domains identifier. Nucleic acids research 33:W116-W120.

Rastas, P., F. C. F. Calboli, B. C. Guo, T. Shikano, and J. Merila. 2016. Construction of Ultradense Linkage Maps with Lep-MAP2: Stickleback F-2 Recombinant Crosses as an Example. Genome Biol Evol 8:78-93.

Simao, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:3210-3212.

Sobel, J. M., and M. A. Streisfeld. 2015. Strong premating reproductive isolation drives incipient speciation in Mimulus aurantiacus. Evolution 69:447-461.

Stanke, M., and S. Waack. 2003. Gene prediction with a hidden Markov model and a new intron submodel. Bioinformatics 19:Ii215-Ii225.

Stankowski, S., J. M. Sobel, and M. A. Streisfeld. 2017. Geographic cline analysis as a tool for studying genome-wide variation: a case study of pollinator-mediated divergence in a monkeyflower. Mol Ecol 26:107-122.

Streisfeld, M. A., W. N. Young, and J. M. Sobel. 2013. Divergent Selection Drives Genetic Differentiation in an R2R3-MYB Transcription Factor That Contributes to Incipient Speciation in Mimulus aurantiacus. Plos Genet 9.

Zerbino, D. R., and E. Birney. 2008. Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. Genome Res 18:821-829.

Supplementary tables

Table S1. Summary of the genetic linkage map constructed using an F2 intercross between the red and yellow ecotypes of subspecies *puniceu*s. The table includes map length in cM for each linkage group (LG), the number of markers associated with each LG, the number of unique map positions, and the average genetic distance in cM between each unique map position. Standard deviations are given in parentheses.

LG	Map length (cM)	Number of markers	Unique map positions	Avg. genetic dist. between unique markers in cM (sd)
1	93.9	969	335	0.28 (0.33)
2	71.37	893	253	0.28 (0.88)
3	76.3	912	256	0.30 (0.65)
4	70.2	851	257	0.28 (0.41)
5	78.7	778	295	0.27 (0.29)
6	69.1	741	247	0.28 (0.88)
7	59.8	738	234	0.26 (0.33)
8	65.6	674	246	0.27 (0.41)
9	68.6	623	182	0.37 (0.58)
10	71.1	410	150	0.48 (0.96)
Avg.	72.74 (8.69)	758.90 (155.75)	245.50 (49.04)	0.31 (0.06)

Table S2. Analysis of gene space completeness in the *M. aurantiacus* genome using CEGMA and BUSCO. The number and percent of core genes found in the final assembly are shown for each analysis (CEGMA, *n* = 248; BUSCO, *n* = 1440).

Analysis	# Genes	% Found in Assembly
CEGMA Complete	233	93.95
CEGMA Partial	244	98.39
BUSCO total complete (duplicated)	1340 (61)	93 (4.2)
BUSCO Fragmented	29	2.0
BUSCO Missing	71	5.0

Table S.3. Sample information for the 37 sequenced samples. Includes their taxon identity, sampling location, percent read alignment, and total sequencing depth.

Sample	Taxon	Latitude	Longitude	% Reads aligned	Seq. Depth
159_83	ssp. *aridus*	32.6630	-116.2230	91.7	21.12
159_84	ssp. *aridus*	32.6630	-116.2230	89.3	21.98
195_1	ssp. *aridus*	32.6300	-116.1429	92.6	20.20
T84	ssp. *aridus*	32.6526	-116.2449	87.2	21.75
T102	ssp. *aurantiacus*	39.0424	-122.7727	94.9	23.74
T104	ssp. *aurantiacus*	39.2045	-123.7646	94.6	25.09
T50	ssp. *aurantiacus*	35.9865	-121.4928	88.3	24.36
T92	ssp. *aurantiacus*	37.8459	-120.6110	94.0	15.16
T144	ssp. *calycinus*	34.1929	-117.2784	93.2	26.00
T150	ssp. *calycinus*	33.8564	-116.8481	94.7	24.02
T90	ssp. *calycinus*	35.5918	-118.5052	91.3	19.97
T91	ssp. *calycinus*	35.3172	-118.5871	95.5	27.91
T101	ssp. *grandiflorus*	39.5536	-121.4301	92.0	16.05
T61	ssp. *grandiflorus*	39.5590	-120.8243	91.6	17.31
T96	ssp. *grandiflorus*	39.0122	-120.7552	92.0	28.21
T99	ssp. *grandiflorus*	39.4376	-121.0599	91.4	23.84
DPR	ssp. *longiflorus*	33.7459	-117.4485	96.0	26.88
SS	ssp. *longiflorus*	34.2722	-118.6100	94.2	30.86
T33	ssp. *longiflorus*	34.3438	-118.5099	94.6	18.87
T8	ssp. *longiflorus*	34.1347	-118.6452	82.6	25.11
KK168	ssp. *parviflorus*	34.0180	-119.6730	91.8	23.66
KK161	ssp. *parviflorus*	34.0180	-119.6730	92.0	19.11
KK180	ssp. *parviflorus*	34.0180	-119.6730	92.4	18.18
KK182	ssp. *parviflorus*	34.0193	-119.6802	91.3	19.46
ELF	ssp. *puniceus*, red	33.0860	-117.1453	93.0	18.20
JMC	ssp. *puniceus*, red	32.7373	-116.9541	93.8	19.06
LH	ssp. *puniceus*, red	33.0609	-117.1188	87.1	19.77
MT	ssp. *puniceus*, red	32.8210	-117.0618	93.7	20.85
UCSD	ssp. *puniceus*, red	32.8894	-117.2362	87.0	18.23
BCRD	ssp. *puniceus*, yellow	32.9496	-116.6380	94.6	20.85
INJ	ssp. *puniceus*, yellow	33.0979	-116.6643	93.1	18.83
LO	ssp. *puniceus*, yellow	32.6767	-116.3312	93.4	18.04
PCT	ssp. *puniceus*, yellow	32.7326	-116.4698	92.3	19.68
POTR	ssp. *puniceus*, yellow	32.6038	-116.6339	90.5	19.27
CLV_GH	*M. clevelandii*	33.1589	-116.8122	92.3	21.31
CLV_11	*M. clevelandii*	33.3391	-116.9325	84.4	15.52
CLV_4	*M. clevelandii*	33.3391	-116.9325	89.3	17.31

Table S.4. Loadings for principal component 1 calculated across all 36 pairwise comparisons (for F_{ST} and d_{xy}) or all nine taxa (for π)

Comparison	F_{ST} PC1	d_{xy} PC1		Taxon	π PC1
AUR_ARI	0.85707	0.95575		ARI	0.9251
BIF_ARI	0.89606	0.86182		AUR	0.86508
CAL_ARI	0.88738	0.95423		AUS	0.97255
CLV_ARI	0.8969	0.39278		BIF	0.77181
FLE_ARI	0.85342	0.94563		CAL	0.96734
LON_ARI	0.9046	0.96057		CLV	0.92017
AUR_BIF	0.81191	0.85563		FLE	0.91044
AUR_CAL	0.32407	0.90376		LON	0.95166
AUR_CLV	0.90725	0.46698		PUN	0.97282
AUR_FLE	0.79434	0.94778			
LON_AUR	0.40962	0.90381			
ARI_AUS	0.90097	0.94262			
AUR_AUS	0.451	0.90512			
BIF_AUS	0.91203	0.90305			
CAL_AUS	0.49255	0.88634			
CLV_AUS	0.90576	0.54264			
FLE_AUS	0.86511	0.94318			
LON_AUS	0.59046	0.88521			
CAL_BIF	0.89043	0.88319			
CLV_BIF	0.91181	0.32045			
FLE_BIF	0.89579	0.86769			
LON_BIF	0.88549	0.89344			
CAL_FLE	0.83543	0.94124			
LON_CAL	0.3786	0.87337			
CAL_CLV	0.90548	0.52607			
FLE_CLV	0.91991	0.45233			
LON_CLV	0.90738	0.51482			
LON_FLE	0.84197	0.94507			
PUN_ARI	0.90783	0.94574			
PUN_AUR	0.51788	0.89861			
PUN_AUS	0.51423	0.84798			
PUN_BIF	0.91936	0.90695			
PUN_CAL	0.62051	0.88316			
PUN_CLV	0.90758	0.55149			
PUN_FLE	0.87615	0.93825			
PUN_LON	0.66912	0.88264			

Supplementary figures

Figure. S1. Map distance (cM/Mbp) vs. physical distance across the 10 linkage groups. Recombination for each marker is estimated relative to the start of the linkage group and plotted at its physical location on each chromosome in the reference assembly.

Figure S2. Geographic distribution of sampling locations for each sample sequenced in this study. Detailed position information for each population can be found in Table S3.

Figure S3. Genome-wide Principal Components Analysis (PCA). Each plot is a separate PCA performed using different sets of taxa. The legend to the right describes the set of taxa included in each analysis, with the capital letter in parentheses and the color representing the specific taxon. A) All taxa; B) all subspecies of *M. aurantiacus*, but excluding *M. clevelandii*; C) only subspecies *aurantiacus*, *longiflorus*, *calycinus*, and the red and yellow ecotypes of subspecies *puniceus*. The percent variation explained by each principal component is given in parentheses.

Figure S4. Incomplete lineage sorting due to rapid diversification. Clades separated by shorter internodes (i.e., separated by less time) are recovered less frequently in window-based trees (500 kb windows). This indicates a strong role for incomplete lineage sorting in areas of the tree where diversification is rapid. The % node concordance is the percentage of window-based of trees that contain a given node in the genome-wide tree, and is plotted against the length of the internode separating each clade. Only clades at and above the level of taxon were included. The red points are the predicted values from a 4-parameter logistic function fitted to the data using an iterative least-squares method.

Figure S5. Patterns of variation plotted across each bush monkeyflower linkage group.
Z-transformed F_{ST}, d_{xy}, and π in overlapping 500 kb windows (step size = 50 kbp). The gray lines are z-transformed scores for each of the 36 pairwise comparisons (F_{ST} and d_{xy}) or nine taxa (π), and the blue line is the z-transformed score for the first principal component (PC1). Estimates of tree concordance, gene count and recombination rate (cM/Mbp) are also shown.

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S5. continued

Figure S6. Common patterns of genome-wide variation mirror variation in the local properties of the genome. Plots are the same as in Fig. 2 of the main text, but for 100 kb windows (step size 10 kb). A) tree concordance; B-D) Z-transformed PC1 for F_{ST}, d_{xy} and π, respectively; E) gene count; and F) recombination rate (cM/Mbp) are ploted across the 10 monkeyflower LGs.

Figure S7. Patterns of variation are non-randomly distributed across the genome.
A) Strong Autocorrelation of tree concordance scores on LG1 over a Mbp scale. B)
Levels of tree concordance, F_{ST}, d_{xy}, and π all show significant autocorrelation at the 2
Mbp scale. The dashed vertical lines show the observed autocorrelation coefficients for
each LG with a 2 Mbp lag. The histogram shows the null distribution of autocorrelation
coefficients (same lag) generated from 1000 random permutations of the genome-wide
values. The observed data are significant at $p = 0.001$ unless stated otherwise.

Figure S8. Patterns of differentiation and divergence for all 36 pairs of taxa. Box plots for each of the 36 pairwise taxonomic comparisons reveal the range of variation in F_{ST} and d_{xy} across the radiation. Moreover, the data show extensive variance among genomic windows within each comparison. Vertical black lines indicate the median, boxes represent the lower and upper quartiles, and whiskers extend to 1.5 times the interquartile range. Taxon abbreviations: *cal, calycinus; lon, longiflorus; aur, aurantiacus; par, parviflorus; ari, aridus; gra, grandiflorus; clv, M. clevelandii.*

Figure S9. Bivariate plots among measures of variation and genomic features across 500 kb genomic windows. Note that this is the same as Figure 3 but with axes units. Also note that the axes are different across rows and columns of the matrix.

Figure S10. Bivariate plots among measures of variation and genomic features across 100 kb genomic windows. The number is the correlation coefficient. Positive correlation coefficients are colored blue and negative coefficients are colored red.

Figure S11. A cartoon depicting the gradual build-up of a heterogeneous differentiation landscape by lineage-specific LS. A) When a population first splits (allopatric divergence with large population sizes), patterns of genome-wide diversity (π pops A and B) are identical due to the complete sharing of ancestral variation among them. Thus, there is no pattern of heterogeneous differentiation between them. B) As time passes, LS begins to act separately within each lineage (LS events are indicated by arrows across the chromosome). This functions to maintain the diversity landscape in the face of new neutral mutations (affected areas in each lineage are shown in color) and also causes ancestral variants to be fixed among them. The result is an increase in F_{ST} in affected areas. C) Long after the split, many LS events have occurred in each lineage. Because the heterogeneous patterns of LS have been conserved since prior to the common ancestor of these taxa (redder areas of the chromosome experience a higher rate of LS), the affected areas are similar between the lineages. As a result, the diversity and differentiation landscapes come to perfectly mirror one another.

Figure S12. Negative correlation between nucleotide diversity and differentiation becomes stronger with increasing divergence time. A) Bivariate plots of the correlation between F_{ST} and π at varying levels of sequence divergence (d_{xy}).

Bonus Haiku!

Peaks and troughs of π,

Static, yet ever-changing.

Their reflection grows.