The meiotic recombination landscape of *Drosophila virilis* is robust to mitotic damage

during hybrid dysgenesis

Lucas W. Hemmer[1†*], Guilherme Dias[2], Brittny Smith[3], Kelley Van Vaerenberghe[1], Ashley

Howard[1], Casey M. Bergman[2] and Justin P. Blumenstiel[1]

[1] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045

[2] Department of Genetics and Institute of Bioinformatics, University of Georgia, Athens, GA

30602

[3] Department of Molecular Biosciences, University of Kansas, Lawrence, KS 66045

[†] Present address: Department of Biology, University of Rochester, Rochester, NY 14627

[*] Corresponding author: lhemmer@ur.rochester.edu

Corresponding author: Lucas W. Hemmer

17   University of Rochester

18   337 Hutchison Hall, Rochester, NY 14627

19   Phone: 402-920-0456

20   Email: lhemmer@ur.rochester.edu

ABSTRACT

Germline DNA damage is a double-edged sword. Programmed double-strand breaks establish the foundation for meiotic recombination and chromosome segregation. However, double-strand breaks also pose a significant challenge for genome stability. Because of this, meiotic double-strand break formation is tightly regulated. However, natural selection can favor selfish behavior in the germline and transposable elements can cause double-strand breaks independent of the carefully regulated meiotic process. To understand how the regulatory mechanisms of meiotic recombination accommodate unregulated transposition, we have characterized the female recombination landscape in a syndrome of hybrid dysgenesis in *Drosophila virilis*. In this system, a cross between two strains of *D. virilis* with divergent transposable element and piRNA profiles results in germline transposition of diverse transposable elements, reduced fertility, and male recombination. We sought to determine how increased transposition during hybrid dysgenesis might perturb the meiotic recombination landscape. Our results show that the overall frequency and distribution of meiotic recombination is extremely robust to germline transposable element activation. However, we also find that hybrid dysgenesis can result in mitotic recombination within the female germline. Overall, these results show that landscape of meiotic recombination may be insensitive to the DNA damage caused by transposition during early development.

INTRODUCTION

Recombination during meiosis plays a critical role in ensuring proper chromosome segregation

(Koehler *et al.* 1996). In addition, meiotic recombination benefits populations since it creates

new genotypic combinations that can facilitate adaptation to changing environments (Weismann

et al. 1904; Burt 2000). While meiosis and sexual reproduction is a conserved process across

many eukaryotes, it can be exploited by selfish genetic elements. Transposable elements (TEs)

are an example of one such class of selfish genetic element. TEs can increase within the genome,

even to the detriment of the host, because fertilization provides a continuous opportunity for TEs

to colonize new genomes (Hickey 1982) and recombination uncouples progenitor TEs from the

harmful effects of descendant insertions. Upon proliferation, TEs can cause mutation, mis-

regulation of gene expression, chromosomal rearrangement, and illegitimate recombination

(Kidwell *et al.* 1977; Kazazian *et al.* 1988; Bennetzen 2000; Slotkin *et al.* 2007; Zhang *et al.*

2011). Moreover, TEs can activate the DNA damage response within developing germline stem

cells and alter stem cell fate (Chen *et al.* 2007; Wylie *et al.* 2014; Tasnim and Kelleher 2018).

The harmful effects of TEs are especially evident in syndromes of hybrid dysgenesis, where

sterility can arise in intraspecific crosses between males carrying TEs and females that lack them

(Bingham *et al.* 1982; Bucheton *et al.* 1984; Yannopoulos *et al.* 1987; Lozovskaya *et al.* 1990).

Hybrid dysgenesis is the result of TE activation in the absence of maternal repression by PIWI-

interacting RNAs (piRNAs) (Aravin *et al.* 2007; Brennecke *et al.* 2008). The piRNA system of

genome defense requires maternal deposition of piRNA to successfully silence TEs across

generations. The combination of unrecognized TEs introduced to a naive genome *via* sperm and

the absence of corresponding piRNAs in the egg results in TE activation and hybrid dysgenesis

(Brennecke *et al.* 2008). A unique syndrome of hybrid dysgenesis in *D. virilis* is observed in

intraspecific crosses between males of an inducing strain (designated strain 160) and reactive

strain females (designated strain 9) (Lozovskaya *et al.* 1990). The primary TE family responsible

for inducing dysgenesis remains unknown and sterility appears to be due to the mass activation

of several TE families abundant in strain 160 but not strain 9. At least four elements are proposed

to contribute significantly to dysgenesis: *Penelope*, *Helena*, *Paris*, and *Polyphemus* (Petrov *et al.*

1995; Evgen'ev *et al.* 1997; Vieira *et al.* 1998; Blumenstiel 2014; Funikov *et al.* 2018). In

addition, activation of TEs during hybrid dysgenesis also leads to transposition of diverse TEs

that are equally abundant between the two strains (Petrov *et al.* 1995).

The impact of hybrid dysgenesis on recombination during meiosis is an open question. TEs may

modulate recombination rates directly since transposition itself can induce illegitimate

recombination at the sites of DNA breaks. As a consequence, in the *P-M* system of hybrid

dysgenesis in *D. melanogaster*, males undergo recombination (Hiraizumi 1971; Kidwell and

Kidwell 1975; Kidwell *et al.* 1977). This is abnormal since meiotic recombination is absent in *D.*

melanogaster males. However, rates of meiotic recombination induced by *P-M* dysgenesis are

typically low, at a rate approximately 2-3% of female meiotic recombination (Preston and Engels

1996). *P-M* induced male recombination is usually attributed to an increased rate of mitotic

exchange induced by DNA damage (Isackson *et al.* 1981). Many of these recombination events

occur near *P* element insertion, require transposase, and are likely the byproduct of *P* element

excision events that are repaired from the homolog (McCarron *et al.* 1994; Sved *et al.* 1995;

Gray *et al.* 1996; Preston and Engels 1996; Preston *et al.* 1996). The effects of hybrid dysgenesis

on female meiotic recombination rates in *D. melanogaster* are less clear. Changes in female

recombination rates were not initially observed during *P-M* hybrid dysgenesis (Hiraizumi 1971;

Chaboissier *et al.* 1995) but later studies found a slight increase (Kidwell 1977; Broadhead *et al.*

1977; Sved *et al.* 1991), often within pericentric heterochromatin. Slightly increased rates of

female recombination were also observed within the *I-R* element system, but these were not

localized near the centromere (Chaboissier *et al.* 1995). However, others have identified no

increase in female recombination rates caused by *P-M* hybrid dysgenesis but, rather, a

redistribution towards regions with low recombination and heterochromatic regions near the

centromere (Slatko 1978; Hiraizumi 1981). There have been no investigations of how hybrid

dysgenesis influences female meiotic recombination in *D. virilis*.

To determine how transposition influences recombination in *Drosophila* females, fine-scale

genetic maps are required. *D. virilis* genetic maps have been obtained only with a limited number

of markers which show that the rate of recombination in *D. virilis* is significantly higher than in

D. melanogaster even though previously estimated rates also differ between studies (Weinstein

1920; Gubenko and Evgen'ev 1984; Huttunen *et al.* 2004) (Table 1). Here, we provide the first

fine-scale genetic map for *D. virilis* using thousands of genotypic markers. Using this map, we

investigate differences in both crossover (CO) frequency and distribution in the hybrid

dysgenesis syndrome of *D. virilis*. Interestingly, we find no significant differences in meiotic

recombination between genetically identical dysgenic and non-dysgenic female progeny. Instead,

we find a strong signature of mitotic recombination that is likely derived from transposition

events in the early germline. Thus, at least in *D. virilis*, meiotic recombination is remarkably

robust to TE activation during hybrid dysgenesis. This suggests that the effects of DNA damage

during early development are not sufficient to trigger changes in the control of recombination

later during meiosis.

110   **MATERIALS AND METHODS**

Fly Stocks and crosses

The hybrid dysgenesis syndrome in *D. virilis* is observed in crosses between reactive strain 9

females and inducer strain 160 males (Lozovskaya *et al.* 1990; Erwin *et al.* 2015). The study was

performed in two stages. A smaller pilot study was performed, followed by a larger second study

that incorporated additional optimization. We observed no differences between these two

experiments, so we combined results for final analysis (see below). For both experiments, each

strain and all subsequent crosses were maintained on standard media at 25°C. Prior to creating

dysgenic and non-dysgenic hybrids, strain 160 and strain 9 were each inbred for 10 generations

by single sib-pair matings. For each direction of the cross, approximately 20 4- to 6-day old

virgin females of one strain and 20 2- to 10-day old males of the other strain were crossed *en*

masse in bottles for six days. Strain 160 males crossed with strain 9 females induce dysgenesis in

the F1 generation while the reciprocal cross yields non-dysgenic F1 progeny. Reciprocal crosses

yield F1 files with identical genetic backgrounds, with the exception of the mitochondrial

genome. By comparing the recombination landscape between F1 females with identical nuclear

genotypes, we obtain a robust analysis of how hybrid dysgenesis influences recombination that

effectively controls for genetic background. Individual virgin F1 females, four days post-

emergence, were backcrossed in single vials to two or three 2- to 10-day old strain 9 males and

maintained in vials for six days. Some non-dysgenic females were only allowed to lay for four or

five days due to their high fertility to prevent vial crowding. Because fertility was low in

dysgenic females, and to increase sample size of progeny within cohorts in the second

experiment, a second brood was obtained from some dysgenic F1 females by transferring to

another vial after ten days. These females were allowed to lay for an additional four days. We

found no difference in recombination between first and second broods (see below), so results

were combined across broods. Female backcross progeny (BC1) were collected once per day and

immediately frozen at -20°C. Between 12 and 20 sisters from each non-dysgenic F1 female was

collected as a sibship. All female progeny of the dysgenic F1 backcrosses were collected.

There is a high amount of variation in fecundity in dysgenic females. Some females are

completely sterile, others have only reduced fecundity and some even have high numbers of

progeny. One approach to analyzing the effects of dysgenesis on recombination would be to

sample only single daughters from each F1 female. However, this approach would not allow for

the discovery of recombination events arising as clusters within the germline. Therefore, we

elected to sequence cohorts of BC1 sisters, balancing our sequencing across cohorts with

different levels of fecundity. To determine if there might be an effect of fecundity on

recombination, all male and female BC1 progeny across the two broods from the second larger

experiment were counted to estimate the fecundity of the F1 mother. In some cases, dysgenic F1

females escape the effects of dysgenesis completely and produce as many progeny as non-

dysgenic females. For these dysgenic F1 females, designated high fecundity, approximately 40

BC1 female progeny were subjected to recombination analysis by sequencing. Progeny produced

by the low fecundity F1 dysgenic females were collected with cohort sizes ranging from one to

40 sisters. By sampling larger cohorts from high fecundity F1 dysgenic females, we sought to

identify clustered recombination events derived within the germline of single females. Power to

detect these events among a small cohort of sisters is low. By examining recombination in both

high fecundity and low fecundity dysgenic females, we also obtained an additional comparison

in the analysis of recombination landscapes between two outcomes of TE activation: TE

activation with strong deleterious effects on fertility versus TE activation with no observable

negative effects on fertility. For a full description of sampling, see Table S1.

DNA extraction, library preparation, and Illumina sequencing

Sequencing libraries were prepared in two batches using different protocols. The first batch

included 192 samples and library preparations were performed following the protocol outlined in

Andolfatto *et al.* (2011) with minor modifications. Single flies were placed into a U-bottom

polypropylene 96-well plate with lysis buffer and 3.5 mm steel grinding balls (BioSpec) then

homogenized with a MiniBeadBeater-96 at 2,100 rpm for 45 seconds. DNA extractions on

homogenized tissue were performed with the Agencourt DNAdvance Genomic DNA Isolation

Kit (Beckman Coulter) following the Insect Tissue Protocol. DNA quantification was spot

checked on some samples and estimated to average 1-2 ng/μl. For each sample, 10 μl of genomic

DNA was digested with 3.3 U of MseI in 20 μls of reaction volume for four hours at 37°C,

followed by heat inactivation at 65°C for 20 min. FC1 and FC2 adaptors (Andolfatto *et al.* 2011)

(Table S2-S3) were ligated to the digested DNA with 1 U of T4 DNA ligase (New England

Biolabs) in 50 μl of reaction volume at 16°C for 5 h and inactivated at 65°C for 10 minutes. The

samples were pooled and concentrated using isopropanol precipitation (1/10 vol NaOAc at pH

5.2, 1 vol of 100% isopropanol, and 1 μl glycogen). The library was resuspended in 125 μl of 1X

Tris-EDTA (pH 8). Adapter dimers were removed with 1.5X vol AMPure XP Beads (Agencourt)

and ligated products were resuspended in 32 μl of 1X Tris-EDTA (pH 8). 200-400 bp DNA

fragments were selected using a BluePippin (Sage Science). Size-selected fragments were

cleaned using 2X vol of AMPure XP beads and resuspended 20 μl of 1X elution buffer (10 μM

Tris, pH 8). Libraries were quantified using a Qubit fluorometer before an 18-cycle PCR

amplification on bar-coded fragments with Phusion high-fidelity PCR Kit (New England

Biolabs). The PCR products were cleaned using 1X vol of AMPure XP Beads.

For the larger second batch (768 samples), sequencing libraries were constructed with an

optimized rapid DNA extraction protocol that uses an in-house Tn5 transposase similar to the

procedure outlined in Picelli *et al.* (2014). DNA was extracted using the *Quick*-DNA 96 kit

(Zymo) and 1-2 ng of DNA was tagmented with Tn5 transposase stored at a concentration of 1.6

mg/ml with pre-annealed oligonucleotides. Tagmentation was performed in 20 μl reaction

volumes containing 2 μl of 5X TAPS-DMF buffer (50 mM TAPS-NaOH, 25 mM $MgCl_2$ (pH

8.5), 50% v/v DMF) and 2 μl of 5x TAPS-PEG buffer (50 mM TAPS-NaOH, 25 mM $MgCl_2$ (pH

8.5), 60% v/v PEG 8000). Samples were incubated at 55°C for 7 min then rapidly lowered to a

holding temperature of 10°C. Reactions were inactivated with 5 μl of 0.2% SDS followed by an

additional incubation at 55°C for 7 min. PCR-based barcoding was performed on 2.5 ul of

sample tagmentation volumes using the KAPA HiFi HotStart ReadyMix PCR Kit (Thermo

Fisher Scientific), 1 μl of 4 μM Index 1 (i7) primers (Table S4), and 1 μl of 4 μM Index 2 (i5)

primers (Table S5) in 9 μl of reaction volume. The PCR thermocycling conditions were: 3 min at

72°C, 2 min 45 sec at 98°C, followed by 14 cycles of 98°C for 15 sec, 62°C for 30 sec, 72°C for

1 min 30 sec. PCR-amplified samples were pooled and the pooled samples were cleaned using

0.8 X vol AMPure XP Beads. We size-selected DNA fragments 250-400 bp from the pooled

sample on a BluePippin and cleaned using 1X vol of AMPure XP Beads.

All libraries were sequenced at the University of Kansas Genomics Core on an Illumina HiSeq

2500 Sequencer with 100 bp single-end sequencing. The first 192 samples were sequenced on

two lanes using the Rapid-Run Mode while the Tn5-produced libraries were sequenced on two

lanes using the High-Output Mode (summary of the output is in File S2).

DNA extraction, library preparation, Pacbio sequencing and assembly

PacBio sequencing was performed on strain 160 after 10 generations of single-sib mating

followed by re-validation for induction of dysgenesis. Females collected for DNA extraction

were allowed to eclose over 10 days, aged for two additional days, starved for 12 hours to

evacuate the gut, then immediately frozen in liquid nitrogen. 500 mg of whole flies were then

used for DNA extraction with Blood Cell and Culture Midi Kit (Qiagen) (Chakraborty *et al.*

2016). The mortar was pre-chilled with liquid nitrogen prior to grinding and the resulting fine

powder was directly transferred into Buffer G2. DNA extraction was performed across 5

columns, using a total of 47.5 mls G2, 190 µl RNAse A (100 mg/ml) and 1.25 ml of Protease

from the Qiagen Kit. 50 mls were placed in a 50°C shaker overnight. After overnight incubation,

debris was spun down and poured onto column. The elution was performed according to

manufacturer's instructions and precipitated with 0.7 volumes of isopropanol, followed by

spooling with a glass rod and resuspending in 100ul EB Buffer. The final DNA concentration

was estimated with a Qubit to be 843 ng/ul, yielding approximately 85 µg. PacBio sequencing

was performed by the University of Michigan DNA Sequencing Core.

Purified strain 160 DNA was used to generate PacBio SMRTbell libraries using the protocol:

Procedure & Checklist 20 kb Template Preparation using BluePippin Size Selection. Briefly,

approximately 10 µg of template was sheared using Covaris g-TUBES to obtain a 20-25 Kb

target length. Sheared DNA was purified using pre-washed AMPureXP beads, analyzed for size

and concentration, then treated with Exo VII to remove single stranded DNA, followed by DNA

damage and end repair. End repaired DNA was then blunt ligated to adaptors to form SMRTbells

and treated with Exo VII plus Exo III to remove any fragments that lack adaptors on both ends.

SMRTbells were purified using pre-washed AMPureXP beads, analyzed for size and

concentration, then run through a Sage Scientific Blue Pippin instrument with 0.75% agarose

dye-free cassette and S1 external marker to remove templates smaller than 10 kb. The PacBio

Binding Calculator was used to determine conditions to anneal primers to SMRTbells and bind

DNA polymerase (P6/C4 chemistry) to form SMRTbell-primer-polymerase complexes.

Complexes were bound to Magbeads and washed to remove unbound template and other

contaminants. Purified complexes with an added Pacific Biosciences internal control were loaded

on a PacBio RS II instrument and run using the MagBead-OCPW protocol. The resulting library

was sequenced on 21 SMRT cells with a movie time of 360 minutes per SMRT cell, totaling

~80-fold coverage of the estimated ~380 Mb *D. virilis* genome (Bosco *et al.* 2007).

Assembly of the PacBio reads was performed using Canu v1.5 with default settings (Koren *et al.*

2017). Canu performs read correction on the 40x longest reads based on the genomeSize

parameter. This parameter was set to 200 Mb after analyzing the read size distribution to avoid

including shorter reads that could result in deterioration of assembly quality. The raw PacBio

reads were aligned back to the Canu assembly with pbmm2 v1.0.0. and the assembly was

polished with Arrow using gcpp v0.01.e12a6d6. PacBio polishing software were downloaded as

part of the pb-assembly metapackage v0.0.6 from bioconda

(https://github.com/PacificBiosciences/pb-assembly). A second round of polishing was

performed after aligning Illumina reads from strain 160 (SRR1200631, Erwin *et al.* 2015) with

BWA-MEM v0.7.17-r1188 (Li 2013) and correcting errors with Pilon v1.22 (Walker *et al.*

2014). Since *D. virilis* strain 160 is largely colinear with the current *D. virilis* reference genome

(strain 15010-1051.87; *Drosophila* 12 Genomes Consortium 2007), we performed reference-

based scaffolding of the strain 160 PacBio assembly using RaGOO v1.1 (Alonge *et al.* 2019).

RaGOO performs scaffolding and correction of potential misassemblies by aligning *de novo*

assemblies to a reference genome and breaking chimeric contigs where necessary. For the

reference, we used an updated version of the SNP-corrected reference genome for strain 160

generated in Erwin *et al.* (2015) that incorporates the chromosome mapping data generated by

Schaeffer *et al.* (2008; see below). General assembly statistics were analyzed with QUAST

v5.0.2 (https://github.com/ablab/quast, commit 67a1136, Gurevich *et al.* 2013). Assembly

completeness was assessed by searching for highly conserved orthologs with BUSCO v3.0.2

(Simão *et al.* 2015) using the Diptera ortholog gene set from OrthoDB v9 (Zdobnov *et al.* 2016).

Assembly statistics are available in Table S6. Coordinates of the mitotic CO clusters (see

methods below) were lifted over to the final version of the PacBio assembly using minimap2

2.16-r922 (Li 2018).

Annotation of genome resources

Illumina-based reference genomes for strains 9 and 160 (Erwin *et al.* 2015) based on the original

Sanger shotgun sequence assembly of *D. virilis* (*Drosophila* 12 Genomes Consortium 2007)

were annotated with the most up-to-date gff file for *D. virilis* (v1.6 Flybase, Gramates *et al.*

2017) using Maker v3.31.9 on default settings (Cantarel *et al.* 2008). Due to errors in the original

reference assembly, genome region 33,464,439-35,498,860 on Chromosome 2 was excluded and

genome region 22,673,797-24,330,628 on Chromosome 5 was placed at position 3,761,384 on

the same chromosome. Thus, previous strain-specific reference genomes (Erwin *et al.* 2015)

were adjusted for two mis-assemblies and updated as '_2' (strain 9, File S3; strain 160, File S4).

TE annotations for Illumina-based reference genomes were obtained using RepeatMasker v4.06

(Tarailo-Graovac and Chen 2009) with the custom TE library from Erwin *et al.* (2015). TE

annotation of the strain 160 PacBio assembly was also obtained using RepeatMasker with the

custom TE library from Erwin *et al.* (2015).

Crossover quantification

Illumina FASTQ files were parsed according to barcode sequences and trimmed by the

University of Kansas Genomics Core facility. The FASTQ files were mapped to the Illumina-

based reference genomes for strains 9 and 160 using the multiplex shotgun genotyping (MSG:

https://github.com/JaneliaSciComp/msg) bioinformatic pipeline for identifying reliable markers

and determining ancestry at those markers using a hidden Markov model (HMM) v0.1

(Andolfatto *et al.* 2011). Briefly, reads were mapped with BWA aln to the strain 9 and 160

parental reference _2 files. Output files were used for HMM determination of ancestry along the

length of the chromosomal segments (see control file, File S5, for settings). The MSG pipeline

provides both ancestry probability calls and CO positions, along with an estimate of the

boundaries for CO positions. Samples with fewer than 10,000 reads were discarded. Double COs

less than 750 kb apart were discarded as these events were considered unlikely. We observed that

reads mapping to regions near the telomere and centromere often predicted the same genotype

across all samples. In principle, this could be driven by segregation distortion. However, these

regions also showed low density for uniquely mapped reads. In addition, segregation distortion

for these regions would drive distortion of linked flanking markers, which we did not see.

Therefore, we considered these regions problematic and removed them from the analysis. COs

located within 500 kb of the telomere of the X and 4th chromosome and COs within 700 kb on

the 2nd, 3rd, and 5th chromosomes were removed. COs near the centromeres were also removed

as follows: within 3.5 Mb on the X chromosome, within 1.1 Mb on the 2nd chromosome, within

1.5 Mb on the 3rd chromosome, within 2.4 Mb on the 4th chromosome, and 2.3 Mb on the 5th

chromosome. The 6th chromosome (corresponding to the 4th in *D. melanogaster*) was also

removed from analysis. In addition, we performed additional manual curation of COs to remove

calls that appeared incorrect. For example, double COs that were spaced closely in samples with

lower numbers of reads and based on ancestry probabilities that were less than 0.9

Data analysis

CO outputs from the MSG pipeline were analyzed with R Version 3.4.2 (R Core Team 2017).

Ancestry probability outputs were used in the R/qtl package (Broman *et al.* 2003) to construct

genetic maps and calculate interference using the Housworth Stahl model (Housworth and Stahl

2003) using the xoi package (Broman and Kwak, Version 0.67-4). We used the lme4 (Bates *et al.*

2015) and lsmeans (Lenth 2016) packages for mixed-model testing of CO events in BC1

progeny. The model construction was performed using the glmer() and glm() functions to test the

random effects of F1 female and fecundity of the F1 female and the fixed effects of dysgenesis

and batches. Fecundity estimates obtained from dysgenic crosses in the second experiment were

first used to determine if fecundity had an effect on total CO count. We found that fecundity had

no effect on CO count ($\chi^2_1 = 2.02$, $p = 0.155$) and excluded it from the final model.

The model for how these effects predict total CO numbers in R was as follows:

glmer(CO.sum ~ batch (fixed) * dys.nondys (fixed) * parent (random),

family=poisson(link=log))

We used likelihood ratio tests to determine the significance of each effect on variance in total CO

number. We used the Biostrings R package (Pagès et al. 2017) to analyze genomic sequences for

correlations between genomic features and recombination. Figures were constructed using

ggplot2 (Wickham 2016).

Analysis of mitotic recombination

Mitotic (or pre-meiotic) recombination is identified by the presence of crossovers that are

common among the progeny of a single parent. These are commonly designated as

recombination clusters and are distinct from hotspots because they are found only within cohorts

of siblings. Clusters indicating germline mitotic recombination were identified as COs in four or

more progeny of a single F1 mother within a 100 kb span. Since the fecundity effects of hybrid

dysgenesis are highly variable, there was imbalance in progeny counts from dysgenic and non-

dysgenic backcrosses. It was therefore necessary to account for this variation in the estimation of

rates of mitotic recombination. This was achieved using a likelihood approach to determine if

rates of mitotic cluster formation (α) within the germline and the frequency of mitotic

recombination within cohorts (β, conditional upon cluster formation) differed between dysgenic

and non-dysgenic parents. Only one mitotic cluster was ever observed per single chromosome so

rate estimate was performed on a per chromosome basis. The probability of not observing a

cluster event (on a given chromosome) is given by the probability that a mitotic recombination

event does not occur in the germline (1-α) plus the probability that a mitotic recombination event

does occur (α) but is not observed among the sampled progeny:

343
$$P_{ClusterNotObs} = (1 - \alpha) + \alpha \times P_{ClusterNotSampled|ClusterOccurred}$$

345   Conditional on mitotic recombination occurring, the probability that it was not observed is equal

346   to the probability that three or fewer progeny within the cohort inherit the recombinant chromatid

347   from the mother. This is given with the binomial probability distribution where β is the

348   frequency of recombinant chromosomes transmitted by the mother with the mitotic event:

350
$$P_{ClusterNotSampled|ClusterOccurred} = \sum_{x=0}^{3} \frac{N!}{x!\,(N-x)!} \beta^x (1 - \beta)^{N-x}$$

352   where N is the total number of progeny in the cohort, β and is the frequency of progeny that

353   inherit the recombinant chromosome. Therefore, parents with three or fewer progeny have

354    $P_{ClusterNotSample|ClusterOccurred} = 1$ under our criteria.

356   When a cluster event is observed, the probability of x progeny with the recombinant

357   chromosome is given by:

$$P(x) = \frac{N!}{x!\,(N-x)!}\beta^x(1-\beta)^{N-x}$$

361 Overall, the probability that a cluster is observed at a given frequency within a cohort is equal to

362 the probability that mitotic recombination happened (α) multiplied by the probability that it is

363 observed at a given frequency, conditional on it having happened:

$$P_{ClusterObserved} = \alpha \times \frac{N!}{x!\,(N-x)!}\beta^x(1-\beta)^{N-x}$$

367 The full likelihood of the data is thus given by:

$$L(Data) = \prod_{i=1}^{m} P_{i.ClusterNotObserved} \prod_{j=1}^{n} P_{j.ClusterObserved}$$

371 where i is index of mothers without an observed mitotic cluster and j as the index of mothers

372 whose progeny indicate a mitotic cluster. Taking the logarithm of our likelihood equation gives

$$\log(L(Data)) = \sum_{i=1}^{m} P_{i.ClusterNotObserved} + \sum_{j=1}^{n} P_{j.ClusterObserved}$$

Mitotic recombination was only ever observed on the X and 3rd chromosomes so a combined

rate was only estimated for these two chromosomes. To estimate rates of mitotic recombination

across dysgenic and non-dysgenic females, we used the Python module Scipy to maximize the

log-likelihood of the data based on α and β. Nested likelihood ratio tests were used to determine

whether there was support for unique values of α or β in dysgenic or non-dysgenic females. Two

three-parameter models were used with distinct cluster formation rates for dysgenic (D) and non-

dysgenic (ND) females (α_D, α_{ND}, β) and, reciprocally, separate frequencies of transmission of the

recombinant chromatid (α, β_D, β_{ND}). We also used as a four-parameter model with individual

estimates for the dysgenic and non-dysgenic mothers (α_D, α_{ND}, β_D, β_{ND}). 95% confidence intervals

(95% CIs) for parameter estimates were obtained by determining parameter values with

likelihood scores two log-likelihood units from the ML estimate with other maximum likelihood

estimated parameters fixed. We tested if models were significantly improved with the inclusion

of additional parameters with a likelihood ratio test (LRT) and a chi-squared distribution with

one degree of freedom for every additional parameter estimated. The Python script for the

maximum likelihood analysis of the mitotic recombination rates is in File S6.

Data availability

All of the de-multiplexed Illumina sequencing reads from BC1 progeny, PacBio reads for strain

160, and the strain 160 PacBio assembly generated in this study are available at the National

Center for Biotechnology Information under accession PRJNA553533. Supplemental files are

available at FigShare.

<div align="center">**RESULTS**</div>

Crossover detection by sequencing

To identify recombination events in reciprocal F1 dysgenic and non-dysgenic progeny, F1

females were backcrossed to reactive strain 9. By sequencing backcross (BC1) progeny, we

identified the recombination events that occurred under the dysgenic and non-dysgenic condition

in the germline of F1 females. F1 dysgenic and non-dysgenic female progeny have identical

nuclear genotypes, which enables a controlled comparison of the effects of TE mobilization on

the recombination landscape. In total, 828 BC1 female progeny were sequenced at sufficient

depth to map recombination breakpoints; 132 samples had fewer than 10,000 reads and were not

included in our analysis. 275 BC1 progeny were sequenced from 20 F1 non-dysgenic females,

311 BC1 progeny were sequenced from 66 low fecundity F1 dysgenic females, and 242 BC1

progeny were sequenced from seven high fecundity F1 dysgenic females. Across all samples, the

MSG pipeline identified a total of 1,150,592 quality-filtered SNPs between the two parental

genomes with an average of distance of 136 bp between SNPs. The MSG HMM uses relative

mapping abundance of reads that are uniquely derived from one of the two parental strains.

Using this information, combined with the crossing scheme, it provides genotype probabilities

for each SNP. For each sample, and at each SNP, the HMM provided a genotype probability of

the BC1 progeny sample being either homozygous for strain 9 (the strain that the F1 progeny

were backcrossed to) or heterozygous. CO breakpoint intervals were then defined based on local

genotype probability calls that switch from greater than 95% to less than 5% along the

chromosome. The median CO breakpoint interval calculated by the MSG HMM was

approximately 18 kb and 84% of all COs localized within a span of 50 kb or less. Seventeen CO

breakpoint intervals were approximately ~ 1 Mb but those were in samples with low read counts

near the 10,000 read cutoff for samples allowed in the analysis.

A high-resolution genetic map of *D. virilis*

Previous studies indicate that the genetic map of *D. virilis* is approximately three times larger

than the genetic map of *D. melanogaster* (Gubenko and Evgen'ev 1984; Huttunen *et al.* 2004).

Critically, the map lengths obtained in those two studies are quite different (Table 1), perhaps

due to a limited number of genetic markers available. Our combined sample has a sufficient

density of markers to provide the first high-resolution recombination map for *D. virilis*.

Combining results from all samples to construct our map was reasonable, since the effects of

dysgenesis were non-significant (see below).

The total genetic map length of *D. virilis* estimated in our combined sample is 732 cM

(centiMorgans) or 2.5 times longer than the genetic map length of *D. melanogaster* (Comeron *et*

al. 2012) (Table 1). The genetic map length estimated in the current study is more than 100 cM

shorter than the first detailed genetic map of *D. virilis* (Gubenko and Evgen'ev 1984) (Table 1).

This may be partly explained by our stringent exclusion of problematic genomic regions.

However, comparing chromosomes that were characterized in all three studies (2, 3, 4, and 5),

our estimate for cM is within or very close to the upper estimate of the two prior studies. In

addition, our cM estimates were more uniform across the chromosomes, which are all fairly

similar in physical length. As expected, the genetic map length of each chromosome in our study

correlates with physical length ($R^2 = 0.851$, $p = 0.025$). There is no significant correlation in the

two prior studies ($R^2 = 0.079$, $p = 0.72$, Huttunen *et al.* 2004; $R^2 = 0.358$, $p = 0.28$, Gubenko and

Evgen'ev 1984, excluding the 6[th] chromosome). The differences in recombination rates between

D. virilis and *D. melanogaster* might appear explained due to their differences in genome size.

The estimated genome size of *D. virilis* is roughly twice the size of the *D. melanogaster* genome,

(404 vs 201 Mb, Bosco *et al.* 2007). Thus, across the entire genome, the average rate of

recombination in *D. virilis* is 1.8 cM/Mb and similar to the average recombination rate of 1.4

cM/Mb in *D. melanogaster*. However, close to half of the *D. virilis* genome is comprised of

satellite DNA, with large expanses of centromeric heterochromatin where little or no crossing

over takes place (Gall and Atherton 1974; Bosco *et al.* 2007). Thus, the *D. virilis* assembly of

euchromatin, where most COs take place, is 206 Mb in length. Accounting for just euchromatic

regions in both species, the average rate of recombination in euchromatin in *D. virilis* is twice as

high as *D. melanogaster* based on euchromatic assembly genome size (3.5 cM/Mb vs. 1.8

cM/Mb). One possible reason for a higher rate of recombination in *D. virilis* may be the fact that

pericentric heterochromatin comprised of satellite DNA may shield the chromosomes arms from

the suppressive centromere effect (Hartmann *et al.* 2019).

Interference reduces the probability of an additional CO in proximity to other COs. We

calculated interference in *D. virilis* using the Housworth-Stahl model to calculate *nu*, a unitless

measure of interference, with a maximum likelihood function based on intercrossover distances

(Housworth and Stahl 2003). If COs are not subject to interference, intercrossover distances are

Poisson distributed and *nu* is equal to one (Broman and Weber 2000). *nu* values greater than one

are observed when COs are spaced more evenly than expected under a Poisson process. Each

chromosome in *D. virilis* has detectable interference between COs with an average *nu* ~ 3 (Table

2). The Houseworth-Stahl model also estimates the percentage of COs produced through an

alternative pathway not subject to interference as the escape parameter *P* (de los Santos *et al.*

2003). Less than 1% of the COs in our study are estimated to be produced through the alternative

CO pathway (Table 2). In contrast to interference, assurance is the recombination control

mechanism that maintains a minimal number of COs on each chromosome to ensure proper

chromosome segregation during meiosis. In the absence of CO assurance and interference, the

distribution of COs should resemble a Poisson distribution with variance in CO number equal to

the mean. We find the CO mean and variance are not equal (7.3 and 4.9, respectively) and the

distribution of CO counts is significantly different from a Poisson distribution (χ^2 = 53.6, p =

5.74E-08). Moreover, *D. virilis* has a higher than expected frequency of individuals with CO

numbers close to the mean (5-8 COs) and a lower than expected frequency of extreme CO

numbers (Figure S1). This indicates the collective action of CO assurance and interference.

In many organisms, the total number of COs between markers in a single tetrad is unobservable

because COs are typically detected using random spore analysis based on only one of the four

chromatids resulting from meiosis. In contrast, tetrad analysis uses the number of COs for each

of the four meiotic products to estimate the frequency of non-exchange tetrads (E_0), single-

exchange tetrads (E_1), or multiple-exchange tetrads (E_n) (Weinstein 1918). We used the

Weinstein method to estimate the frequency of E_0 tetrads for each chromosome in *D. virilis*. The

X chromosome and third chromosome E_0 tetrad frequencies were estimated at 1.2% and 2.1%

respectively. These calculated E_0 tetrad frequencies in *D. virilis* are lower in comparison to *D.*

melanogaster, previously estimated to be 5-10% (Zwick *et al.* 1999; Hughes *et al.* 2018).

However, the Weinstein method indicated negative E_0 tetrad frequencies ranging from -2.6% to -

3.9% for the second, fourth, and fifth chromosomes. Negative values were also sometimes

obtained for other exchange class frequencies (Table S7). Negative tetrad frequencies are a

drawback to using the classic Weinstein method (Zwick *et al.* 1999). Nonetheless, these results

indicate that there are many fewer non-exchange tetrads in *D. virilis* compared to *D.*

melanogaster, consistent with the higher estimated rate of recombination per Mb in *D. virilis*.

Recombination rates are often correlated with certain sequence features, such as GC content,

simple motifs, and nucleotide polymorphism (Begun and Aquadro 1992; Kong *et al.* 2002;

Comeron *et al.* 2012). In *D. virilis*, recombination rates appear to be weakly correlated with GC

content and gene density as not all chromosomes show significant correlations with either

sequence parameter (Table 3). This may be due to unusual patterns of recombination along the

length of the chromosome (discussed later). Simple repeats and SNP density between the two

strains show significant positive correlations with recombination rate on all chromosomes even

after removal of non-recombining regions. Nucleotide diversity is frequently correlated with

recombination rates (Begun and Aquadro 1992; Kong *et al.* 2002) and the strong correlation

between SNP density and recombination in our data is consistent with this pattern (Figure S2).

TE density shows a strong negative correlation with recombination rate but this becomes weaker

when non-recombining regions are removed (Table 3, Figure S2). The similar pattern of weak

correlation between TE density and recombination when regions without recombination are

removed is also seen in *D. melanogaster* (Rizzon *et al.* 2002), where TEs mostly accumulate in

non-recombining pericentromeric heterochromatin (Bartolomé *et al.* 2002; Bergman *et al.* 2006).

No modulation of recombination rates during hybrid dysgenesis

To compare recombination rates in dysgenic and non-dysgenic females, we constructed a full

mixed-effects likelihood model using the lme4 R package (Bates et al. 2015). The direction of

the cross (dysgenic vs. non-dysgenic) and F1 collection batch (pilot vs. second experiment) were

treated as fixed effects; the F1 female of origin was treated as a random effect. In the full model,

we find no difference in the total number of COs between the pilot and second experiment ($\chi^2{}_1 =$

0.10, $p = 0.755$). This suggests that there were no effects of library construction procedure and

justifies combining data sets. Likewise, no effect was found for fecundity in dysgenic flies on

CO numbers ($\chi^2{}_1 = 2.02$, $p = 0.155$). Importantly, dysgenesis did not have a significant effect on

total CO numbers ($\chi^2{}_1 = 0.45$, $p = 0.506$) with nearly identical means in CO number between

dysgenic and non-dysgenic progeny (Figure 1A). There was a marginal interaction between

dysgenesis and batch ($\chi^2{}_1 = 3.17$, $p = 0.075$), but this appears driven by a single high fecundity

dysgenic F1. This F1 female, designated 701, revealed a significantly larger mean CO number

than the other high fecundity dysgenic females (8.3 COs, least squares mean contrast, $p = 0.021$,

Figure 1B). Without the 701 female, the interaction between dysgenesis and batch is non-existent

($\chi^2{}_1 = 0.06$, $p = 0.803$). Overall, full model revealed that the identity of the female had minimal

effect on CO number (variance < 0.0001).

The higher recombination rate per Mb in *D. virilis* in comparison to *D. melanogaster* is due to a

higher number of COs on each chromosome. In *D. melanogaster*, chromosome arms typically

have zero, one, or two COs with three COs on a single chromosome arm being rare (Miller *et al.*

2016). In contrast, a chromosome with three or more COs is common in *D. virilis*, in both

dysgenic and non-dysgenic directions of the cross. Chromosomes with five COs were also

observed (Figure 2). CO counts per chromosome were highly similar between the progeny of

dysgenic and non-dysgenic F1 females ($\chi^2{}_4 = 0.529$, $p = 0.97$). Likewise, there was also no

difference between dysgenic mothers if they were high fecundity ($\chi^2{}_4 = 3.70$, $p = 0.45$) or low

fecundity ($\chi^2{}_4 = 3.45$, $p = 0.49$). Since per chromosome CO counts provide the data for tetrad

analysis, these results indicate that dysgenesis does not modulate tetrad exchange class

frequency.

We also examined the distribution of recombination events along the length of each chromosome

between non-dysgenic flies, high fecundity dysgenic flies, and low fecundity dysgenic flies.

There were no major changes in the distribution of recombination along the length of the

chromosomes (Figure 3). The chromosomal recombination rates between all three groups are

strongly correlated (Table S8). Interference plays a role in determining CO positioning.

Therefore, we determined whether interference was altered by dysgenesis by comparing the

distribution of distances between crossovers identified in the progeny of dysgenic and non-

dysgenic flies. We found no differences in the distribution of crossovers for any chromosome

(Mann-Whitney U test, $p > 0.5$). Overall, we find no differences in the recombination landscape

between dysgenic and non-dysgenic F1 mothers in *D. virilis* at the global level. This suggests

there is little to no feedback between putative activation of the DNA damage response during

dysgenesis in *D. virilis* and the modulation of meiotic recombination.

A signature of early DNA damage and mitotic crossing over in dysgenesis

Despite observing no significant effect of dysgenesis on meiotic recombination rates, we

observed several genomic regions that exhibited a much higher apparent number of COs during

hybrid dysgenesis. For example, within a 500 kb region on the third chromosome, the apparent

recombination rate was 26 cM/Mb, nearly twice as high as any other interval within the genome

(Figure 3C, within the circled region). 32 COs were identified as arising from dysgenic F1

females compared to a single CO identified from non-dysgenic mothers. The COs in this interval

provided evidence for a mitotic recombination cluster because the majority (28/32) were

identified in the progeny of a single highly-fecund dysgenic F1 mother designated 5011. The

mitotic recombination event identified is visible and identifiable in the genotypes of the BC1

progeny of F1 mother 5011 in comparison to the BC1 progeny of another high fecundity

dysgenic female (4029) that lacks a cluster of recombination in this region (Figure 4A-B).

Reciprocal CO products were observed with equal frequency among the BC1 progeny of F1

mother 5011 ($\chi^2_1 = 0.13$, $p = 0.727$, Figure 4B) indicating no transmission bias among

recombinant gametes. Additional unique COs were detected along the entire length of the third

chromosome upstream and downstream of the recombination cluster. The high frequency of

recombinants at the same location identified among this cohort of BC1 progeny likely indicates

an event in the early germline of the F1 female 5011.

Another recombination cluster was identified on the X chromosome, approximately 21.7 Mb

from the telomere. In this region, there was an effective recombination rate of 15.7 cM / Mb

(Figure 3A, within the circled region). Once again, the vast majority of COs within this 500 kb

interval are part of another cluster of recombination attributed to a single highly-fecund dysgenic

F1 female designated 4029 (Figure 4C). The cluster of recombination is revealed in only half of

the progeny of the F1 mother 4029. Interestingly, no additional COs were detected in the portion

of the X chromosome distal from the recombination event and all markers in the distal portion

were heterozygous. The extreme excess of heterozygosity on the X chromosome in the BC1

progeny indicates transmission distortion of the strain 160 genotype distal from the CO from the

4029 mother ($\chi^2_1 = 32$, $p = 0.141E-08$, Figure 4C). Markers proximal to the cluster of

recombination show no transmission distortion ($\chi^2_1 = 0.13$, $p = 0.727$, Figure 4C). Moreover,

crossing over was found to occur in the proximal portion of the X. Thus, it appears that there was

germline loss of heterozygosity (LOH) for the distal region of the chromosome in the germline

of mother 4029.

These two clusters of recombination were identified based on their observed higher rates of

recombination in the dysgenic germline. We infer the clusters are mitotic recombination hotspots

because the chromosomes with the focal recombination event were exclusively found to be

derived from a single F1 mother. Additional mitotic COs may be indistinguishable from meiotic

recombination since such events may be rare and are only evident if the events occur early

enough in development and are associated with depletion of other non-CO germline stem cells.

To uncover additional evidence for other mitotic COs in our data, we screened for clusters of

recombination by identifying CO events within the same 100 kb interval in four or more progeny

of a single F1 mother. Using these criteria, we identified five additional candidate mitotic

recombination clusters in progeny from dysgenic mothers and one additional candidate in

progeny from a non-dysgenic mother (Table 4). Four of these six additional putative clusters of

recombination were also associated with segregation distortion of a single genotype in a

significant portion of the chromosome and no additional COs detected in the distorted region

(Table 4, Figure S3).

To address whether activation of TEs during dysgenesis played a role in causing clustered

mitotic recombination events, we generated a *de novo* PacBio assembly for strain 160 and

determined whether regions of the 160 inducer chromosomes where recombination clusters were

identified contained intact copies of TEs implicated in hybrid dysgenesis (*Penelope*,

Polyphemus, *Helena*, *Skippy*, and *Paris*). Active versions of these TEs are absent in strain 9 and

their activation during hybrid dysgenesis may induce DNA damage on the 160 chromosome for

subsequent repair *via* mitotic recombination. Of these, *Paris* and *Polyphemus* are the most likely

associated with chromosome breaks since they are DNA transposons capable of excision. By

examining the PacBio assembly of strain 160, we found that five clusters of recombination

contained an intact insertion for a TE known to be absent in strain 9 and present in strain 160

within the defined boundaries of the CO (Table 4). Three clusters were associated with

Polyphemus elements in strain 160. One cluster was associated with a *Paris* element and a fifth

cluster on chromosome X contained both elements (Table 4). To determine whether mitotic

recombination events are directly associated with excision during dysgenesis, we performed PCR

and sequencing on original DNA samples used for Illumina genotyping of the BC1 progeny of

F1 mother 5011 with primers that flanked the *Polyphemus* insertion on chromosome 3.

Examination of the one individual that retained the strain 160 haplotype across the CO

breakpoint indicated that even though there was no recombination event for that sample, excision

of *Polyphemus* was identified in the lesion left by the tandem site duplication. In four

recombinant samples, we confirmed that *Polyphemus* was absent. Recombination events initiated

from an excision are expected to be repaired off the non-insertion chromosome and, thus we

were not able to find direct evidence of an excision scar in recombinants lacking the *Polyphemus*

element. Nonetheless, these results support the conclusion that this particular *Polyphemus*

insertion was activated in female 5011 and this was associated with a cluster of mitotic

recombination. Overall, our results suggest clusters of recombination occur more frequently in

dysgenic relative to non-dysgenic females and often occur in regions containing intact copies of

two DNA transposons (*Polyphemus* and *Paris*) associated with hybrid dysgenesis.

Since we identified multiple mitotic clusters of recombination in dysgenic crosses, we sought to

more formally determine whether there was evidence for a statistically significant higher rate of

recombination cluster formation in dysgenic females. Since the criteria for identifying a cluster

was based on observing four or more individuals with a CO within a given span, it was necessary

to account for variation in cohort size. We achieved this by developing a likelihood model where

the probability of observing a set of chromosomes providing evidence of a cluster within a

cohort was a function of the probability that a mitotic event occurs on that chromosome within

an F1 female (α) and the probability of observing that event on a given chromosome (β) among

the sequenced progeny. We considered two, three and four parameter models where either α or β

would be the same under dysgenesis or non-dysgenesis, or there would be a unique value

depending on condition. Using nested likelihood ratio tests, we find support for a three-

parameter model with separate β estimates for dysgenic and non-dysgenic mothers and a shared

α estimate over the two-parameter model ($\alpha = 0.12$, $\beta_{Dys} = 0.78$, $\beta_{NonDys} = 0.11$, LRT, $\chi^2_1 = 51.6$,

$p = 6.92E\text{-}13$, Table S9). Comparing a three-parameter model where α is estimated separately to

the two-parameter model is not supported (LRT, $\chi^2_1 = 3.39$, $p = 0.066$, Table S9). Finally, we do

not find support for a four-parameter model over the three-parameter model with separate β

estimates (LRT, $\chi^2_1 = 1.88$, $p = 0.170$, Table S9). Overall, these results support a similar

baseline rate of occurrence of mitotic events in the dysgenic and non-dysgenic germlines.

However, the frequency of chromosomes that are transmitted with mitotic damage is greater in

the dysgenic germline. Overall, we find evidence that that the total transmission rate of mitotic

damage ($\alpha * \beta$) is more than six times greater in the dysgenic germline (0.096 mitotic COs per

gamete in dysgenic, 0.014 mitotic COs per gamete in non-dysgenic).

653   **DISCUSSION**

A high-resolution genetic map of *D. virilis*

In this study, we obtained the first high-resolution genetic map of *D. virilis*. Combined with

recombination studies in *D. simulans*, *D. mauritiana*, *D. yakuba*, *D. persimilis*, *D. miranda*, *D.*

serrata, *D. mojavensis*, and others (Evgen'ev 1976; True *et al.* 1996; Takano-Shimizu 2001;

Staten *et al.* 2004; Kulathinal *et al.* 2008; Stevison and Noor 2010; Stocker *et al.* 2012;

Smukowski Heil *et al.* 2015; Howie *et al.* 2019), our genetic map of *D. virilis* will aid future

studies of the evolution of recombination in *Drosophila*. Of note is the high rate of

recombination in *D. virilis* in comparison to other species, especially *D. melanogaster*.

Recombination rates in species of *Drosophila* frequently peak in the middle of the chromosome

arm and decrease towards the centromere and telomere (True *et al.* 1996). However, the

distribution of recombination on the second, third, and fourth chromosomes in *D. virilis*

resembles a bimodal distribution (Figure 3). The bimodal distribution may be the result of the

exceptionally high recombination rates in *D. virilis*. When two or more crossovers on a single

chromosome is common, interference preventing CO formation in close proximity would spread

COs more evenly along the length of the chromosome.

Meiotic recombination in light of hybrid dysgenesis in *D. virilis*

This study is one of the few to examine the effects of TE activity on the meiotic recombination

landscape, and the first to do so using the hybrid dysgenesis syndrome in *D. virilis*. Results from

previous studies of hybrid dysgenesis in *D. melanogaster* were conflicting as some found no

effect on female recombination (Hiraizumi 1971; Chaboissier *et al.* 1995) while others found

increases in the recombination rate (Kidwell 1977; Broadhead *et al.* 1977; Sved *et al.* 1991) or

changes in the distribution of recombination (Slatko 1978; Hiraizumi 1981). In addition to

reporting findings using the dysgenic syndrome of a different species, ours is also the first study

to investigate the effects of hybrid dysgenesis on recombination using high-throughput

genotyping rather than phenotypic markers. This allows a greater insight into the fine-scale

changes in recombination rates and distribution that may previously have escaped unnoticed.

Modulation of recombination by hybrid dysgenesis may occur through different mechanisms.

First, mitotic recombination would be directly initiated by double-stranded breaks (DSBs) that

arise from either TE insertion or excision. Second, DSBs caused by TE activity could modulate

global recombination rates through DNA damage signaling. In support of the first hypothesis, we

identified clusters of recombination originating from mitotic CO events in germline stem cells

(GSCs) (Figure 5). During hybrid dysgenesis, the damaging effects of transposons are often

observed in the germline during early development (Engels and Preston 1979; Niki and Chigusa

1986; Sokolova *et al.* 2010). DSBs produced as an outcome of transposition may be repaired by

one of several mechanisms including homologous recombination *via* mitotic crossing over. In

the case of the cluster of COs on the third chromosome in F1 mother 5011, excision of a

Polyphemus DNA transposon may have produced a DSB repaired *via* homologous

recombination in the mitotic germline, in G1 prior to DNA replication, within a developing GSC

(Figure 5A). In this scenario, reciprocal products of the CO would appear in all daughter cells

descended from this GSC and reciprocal products would be observed, on average, in equal

frequency among gametes. Alternatively, a mitotic CO may have occurred after DNA replication

in G2 prior to mitosis in the germline of the 5011 mother (Figure 5B). During mitosis, the

chromatids segregated according to Z segregation (Germani *et al.* 2018; Pimpinelli and Ripoll

1986) such that reciprocal CO products were transmitted to one daughter cell while the other

daughter cell would have received the non-CO chromatids. Other GSCs must have been retained

within the 5011 mother because there are several progeny without the common CO product.

However, the limited number of progeny lacking either of the reciprocal CO chromatids

indicates a depletion of other intact GSCs. We attribute this to an early crisis in GSC survival

due to hybrid dysgenesis, followed by re-population of the GSCs from descendants of a single

cell carrying the CO chromatids. GSCs marked with the mitotic CO were thus able to recover

and rescue fertility after hybrid dysgenesis in the high fecundity female. This is consistent with

the observation that hybrid dysgenesis is associated with an early phase of germline depletion

(Sokolova *et al.* 2013).

A second mitotic CO event on the X, with different consequences, likely occurred in the early

developing germline of the 4029 mother (Figure 5C). In this case, there was apparent loss-of-

heterozygosity (LOH) distal from the CO. This mitotic CO event would have occurred after

DNA replication in G2, with a pattern of X segregation (Germani *et al.* 2018; Pimpinelli and

Ripoll 1986) resulting in each daughter cell receiving one chromatid with the CO and one

without and LOH distal from the CO. This LOH would be responsible for failure to detect

additional meiotic COs derived from the homozygous distal region. The complete transmission

distortion of the distal region suggests severe depletion of GSCs with the reciprocal mitotic CO

products. Again, this is consistent with a severe reduction of GSCs, followed by re-population

from even a single GSC and restoration of fertility in the high fecundity female. Interestingly, the

bounds of the mitotic CO derived from the 4029 mother do not contain intact dysgenesis-

associated TEs nor any other intact TEs in the strain 160 genome. This mitotic CO may therefore

have been the product of a new TE insertion in the genome of the 4029 mother. LOH is also

observed among several clusters of recombination and most of these clusters are associated with

either *Polyphemus* or *Paris* DNA transposons (Table 4, Figure S3). LOH *via* mitotic

recombination is observed after DNA damage or chromosomal breakage in cancer cells (Lasko

1991) and in yeast recombination studies (Symington *et al.* 2014). The greater number of mitotic

recombination events with segregation distortion observed in our data is consistent with previous

observations of non-random segregation of chromatids in clonal analysis; chromatids involved in

mitotic exchange are more likely to segregate into separate daughter cells (X segregation) than

the same daughter cell (Z segregation) in mosaic analyses (Germani *et al.* 2018; Pimpinelli and

Ripoll 1986) Likewise, segregation distortion is frequently observed during hybrid dysgenesis

(Kidwell and Kidwell 1975; Kidwell *et al.* 1977). Our study therefore links segregation

distortion *via* mitotic recombination and subsequent LOH within female germlines as a result of

hybrid dysgenesis.

The observed number of mitotic CO events identified in dysgenic progeny is interesting because

the crossing over pathway is least likely to repair non-programmed DSBs (Preston *et al.* 2006).

Mitotic COs only account for ~1% of all COs detected in our dataset and contribute minimally to

the genetic map length (Figure S4). Interestingly, the mitotic exchange rate is similar to the rates

of male recombination under *P* element hybrid dysgenesis (Kidwell *et al.* 1977; Preston *et al.*

1996). Other pathways of DNA damage repair including non-homologous end joining and

single-strand annealing are probably more common but undetectable in our assay. Rates of

mitotic crossing over may also be higher than what we estimated since many mitotic COs would

not meet our criteria for classifying candidate clusters because many dysgenic mothers produced

fewer than four offspring.

Overall, despite evidence for the mitotic recombination associated with DNA transposon

excision, we found no major differences in the distribution and frequency of meiotic

recombination in *D. virilis* under hybrid dysgenesis, suggesting that DNA damage signaling

during dysgenesis does not modulate meiotic recombination. The DNA damage response has a

critical role in regulating meiotic recombination (Sjöblom and Lähdetie 1996; Lu *et al*. 2010;

Joyce *et al.* 2011). DNA damage response regulators such as *p53* and *chk2* also influence the fate

of GSCs during hybrid dysgenesis (Tasnim and Kelleher 2018). The incomplete penetrance of

hybrid dysgenesis in *D. virilis* may arise from cell to cell variation in the total amount of DNA

damage or in stochastic variation in the DNA damage response. However, we found no

differences in recombination rates between dysgenic flies with minimal germline atrophy (high

fecundity) and severe germline atrophy (low fecundity). This suggests that DNA damage

signaling activated by dysgenesis does not modulate meiotic recombination. This is likely due to

the fact that most TE activity happens in GSCs during early development (Engels and Preston

1979; Sokolova *et al.* 2010). By the onset of meiosis, the harmful effects of TE activity during

dysgenesis may have likely subsided. In *D. virilis*, TE suppression is restored by adulthood in

dysgenic progeny *via* re-establishment of asymmetric piRNAs and the negative impacts of

dysgenesis disappear in subsequent generations (Erwin *et al.* 2015). This suggests that TEs likely

produce few DSBs during meiosis in the *D. virilis* hybrid dysgenesis model. We thus conclude

that the timing of transposition is an important factor that determines how TEs influence

recombination. In the future, it will be worth investigating whether recombination is also robust

to transposition that occurs closer to the initiation of meiotic recombination.

Acknowledgements

We thank John Kelly, Stuart Macdonald, Michael Evgen'ev, Sergei Funikov, members of the

Robert Unckless and Amanda Larracuente labs, and anonymous reviewers for helpful comments

and suggestions. Peter Andolfatto and Audrey Lamb provided guidance on the multiplex shotgun

genotyping protocol and Annemarie Chilton produced the custom Tn5 enzyme. Enzyme

production and development of the library generation protocol was in part supported by NIH R01

OD010974 to Stuart Macdonald (KU) and Tony Long (UC Irvine). We also thank Jennifer

Hackett and the University of Kansas Genome Sequencing Core (NIH COBRE P20GM103638)

and Christina McHenry and the University of Michigan Advanced Genomics Core for library

prep and sequencing. We also thank the KU Center for Research Computing, especially Riley

Epperson, and the Georgia Advanced Computing Resource Center for computational resources.

This work was supported by the University of Kansas, the University of Georgia, support from

COBRE P20GM103638 and NSF Award 1413532 to JB.

REFERENCES

Alonge, M., S. Soyk, S. Ramakrishnan, X. Wang, F. J. Sedlazeck, Z. B. Lippman, and M. C.

Schatz, 2019 Fast and accurate reference-guided scaffolding on draft genomes. bioRxiv

https://doi.org/10.1101/519637

Andolfatto P., D. Davison, D. Erezyilmaz, T. T. Hu, J. Mast, *et al.*, 2011 Multiplexed shotgun

genotyping for rapid and efficient genetic mapping. Genome Res. 21: 610–617.

Aravin A. A., G. J. Hannon, and J. Brennecke, 2007 The Piwi-piRNA pathway provides an

adaptive defense in the transposon arms race. Science 318: 761–764.

Bartolomé C., X. Maside, and B. Charlesworth, 2002 On the abundance and distribution of

transposable elements in the genome of *Drosophila melanogaster*. Mol. Biol. Evol. 19:

926–937.

Bates D., M. Mächler, B. Bolker, and S. Walker, 2015 Fitting linear mixed-effects models using

lme4. J. Stat. Softw. 67. https://doi.org/10.18637/jss.v067.i01

Begun D. J., and C. F. Aquadro, 1992 Levels of naturally occurring DNA polymorphism

correlate with recombination rates in *D. melanogaster*. Nature 356: 519–520.

Bennetzen J. L., 2000 Transposable element contributions to plant gene and genome evolution,

pp. 251–269 in *Plant Molecular Evolution*.

Bergman C. M., H. Quesneville, D. Anxolabéhère, M. Ashburner, 2006 Recurrent insertion and

duplication generate networks of transposable element sequences in the *Drosophila*

melanogaster genome. Genome Biol. 7: R112.

Bingham P. M., M. G. Kidwell, and G. M. Rubin, 1982 The molecular basis of *P-M* hybrid

dysgenesis: the role of the *P* element, a *P*-strain-specific transposon family. Cell 29: 995–

1004.

Blumenstiel J. P., 2014 Whole genome sequencing in *Drosophila virilis* identifies *Polyphemus*, a

recently activated *Tc1*-like transposon with a possible role in hybrid dysgenesis. Mob. DNA

5: 6.

Bosco G., P. Campbell, J. T. Leiva-Neto, and T. A. Markow, 2007 Analysis of *Drosophila*

species genome size and satellite DNA content reveals significant differences among strains

as well as between species. Genetics 177: 1277–1290.

Brennecke J., C. D. Malone, A. A. Aravin, R. Sachidanandam, A. Stark, *et al.*, 2008 An

epigenetic role for maternally inherited piRNAs in transposon silencing. Science 322:

1387–1392.

Broadhead R. S., J. F. Kidwell, and M. G. Kidwell, 1977 Variation of the recombination fraction

in *Drosophila melanogaster* females. J. Hered. 68: 323–326.

Broman K. W., and I. Y. Kwak, R/xoi: tools for analyzing crossover interference. Version 0.67-

4. https://cran.r-project.org/web/packages/xoi/index.html

Broman K. W., and J. L. Weber, 2000 Characterization of human crossover interference. Am. J.

Hum. Genet. 66: 1911–1926.

Broman K. W., H. Wu, S. Sen, and G. A. Churchill, 2003 R/qtl: QTL mapping in experimental

crosses. Bioinformatics 19: 889–890.

Brower-Toland B., S. D. Findley, L. Jiang, L. Liu, H. Yin, *et al.*, 2007 *Drosophila* PIWI

associates with chromatin and interacts directly with HP1a. Genes Dev. 21: 2300–2311.

Bucheton A., R. Paro, H. M. Sang, A. Pelisson, and D. J. Finnegan, 1984 The molecular basis of

I-R hybrid dysgenesis in *Drosophila melanogaster*: identification, cloning, and properties of

the *I* factor. Cell 38: 153–163.

Burt A., 2000 Perspective: sex, recombination, and the efficacy of selection--was Weismann

right? Evolution 54: 337–351.

Cantarel B. L., I. Korf, S. M. C. Robb, G. Parra, E. Ross, *et al.*, 2008 MAKER: an easy-to-use

annotation pipeline designed for emerging model organism genomes. Genome Res. 18:

188–196.

Chaboissier M. C., F. Lemeunier, and A. Bucheton, 1995 *IR* hybrid dysgenesis increases the

frequency of recombination in *Drosophila melanogaster*. Genet. Res. 65: 167–174.

Chakraborty M., J. G. Baldwin-Brown, A. D. Long, and J. J. Emerson, 2016 Contiguous and

accurate de novo assembly of metazoan genomes with modest long read coverage. Nucleic

Acids Res. 44: e147.

Chen Y., A. Pane, and T. Schüpbach, 2007 *cutoff* and *aubergine* mutations result in

retrotransposon upregulation and checkpoint activation in *Drosophila*. Curr. Biol. 17: 637–

642.

Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in

Drosophila melanogaster. PLoS Genet. 8: e1002905.

Drosophila 12 Genomes Consortium, 2007 Evolution of genes and genomes on the *Drosophila*

phylogeny. Nature 450: 203–218.

Engels W. R., and C. R. Preston, 1979 Hybrid dysgenesis in *Drosophila melanogaster*: the

biology of female and male sterility. Genetics 92: 161–174.

Erwin A. A., M. A. Galdos, M. L. Wickersheim, C. C. Harrison, K. D. Marr, *et al.*, 2015 piRNAs

are associated with diverse transgenerational effects on gene and transposon expression in a

hybrid dysgenic syndrome of *D. virilis*. PLoS Genet. 11: e1005332.

Evgen'ev M. B., 1976 The cytogenetic study of crossing over in interspecific hybrids between

Drosophila virilis and *D. texana*. Genetics 83: 355–360.

Evgen'ev M. B., H. Zelentsova, N. Shostak, M. Kozitsina, V. Barskyi, *et al.*, 1997 *Penelope*, a

new family of transposable elements and its possible role in hybrid dysgenesis in

Drosophila virilis. Proc. Natl. Acad. Sci. U. S. A. 94: 196–201.

Funikov S. Y., D. A. Kulikova, G. S. Krasnov, A. P. Rezvykh, L. N. Chuvakova, *et al.*, 2018

Spontaneous gain of susceptibility suggests a novel mechanism of resistance to hybrid

dysgenesis in *Drosophila virilis*. PLoS Genet. 14: e1007400.

Gall J. G., and D. D. Atherton, 1974 Satellite DNA sequences in *Drosophila virilis*. Journal of

Molecular Biology 85: 633–664.

Germani F., C. Bergantinos, and L. A. Johnston, 2018 Mosaic analysis in *Drosophila*. Genetics

208: 473–490.

Gramates L. S., S. J. Marygold, G. D. Santos, J.-M. Urbano, G. Antonazzo, *et al.*, 2017 FlyBase

at 25: looking to the future. Nucleic Acids Res. 45: D663–D671.

Gray Y. H., M. M. Tanaka, and J. A. Sved, 1996 *P*-element-induced recombination in

Drosophila melanogaster: hybrid element insertion. Genetics 144: 1601–1610.

Gubenko I. S., and M. B. Evgen'ev, 1984 Cytological and linkage maps of *Drosophila virilis*

chromosomes. Genetica 65: 127–139.

Gurevich A., V. Saveliev, N. Vyahhi, and G. Tesler, 2013 QUAST: quality assessment tool for

genome assemblies. Bioinformatics 29: 1072–1075.

Hartmann M., J. Umbanhowar, and J. Sekelsky, 2019 Centromere-proximal meiotic crossovers

in *Drosophila melanogaster* are suppressed by both highly-repetitive heterochromatin and

the centromere effect. bioRxiv doi: https://doi.org/10.1101/637561

Hickey D. A., 1982 Selfish DNA: a sexually-transmitted nuclear parasite. Genetics 101: 519–

531.

Hiraizumi Y., 1971 Spontaneous recombination in *Drosophila melanogaster* males. Proc. Natl.

Acad. Sci. U. S. A. 68: 268–270.

Hiraizumi Y., 1981 Heterochromatic recombination in germ cells of *Drosophila melanogaster*

females. Genetics 98: 105–114.

Housworth E. A., and F. W. Stahl, 2003 Crossover interference in humans. Am. J. Hum. Genet.

73: 188–197.

Howie J. M., R. Mazzucco, T. Taus, V. Nolte, and C. Schlotterer, 2018 DNA motifs are not

general predictors of recombination in two *Drosophila* sister species. Genome Biol Evol 11:

1345-1357.

Hughes S. E., D. E. Miller, A. L. Miller, and R. Scott Hawley, 2018 Female meiosis: Synapsis,

recombination, and segregation in *Drosophila melanogaster*. Genetics 208: 875–908.

Huttunen S., J. Aspi, A. Hoikkala, and C. Schlötterer, 2004 QTL analysis of variation in male

courtship song characters in *Drosophila virilis*. Heredity 92: 263–269.

Isackson D. R., T. K. Johnson, and R. E. Denell, 1981 Hybrid dysgenesis in *Drosophila*: the

mechanism of *T-007*-induced male recombination. Mol. Gen. Genet. 184: 539–543.

Jensen P. A., J. R. Stuart, M. P. Goodpaster, J. W. Goodman, and M. J. Simmons, 2008 Cytotype

regulation of *P* transposable elements in *Drosophila melanogaster*: Repressor polypeptides

or piRNAs? Genetics 179: 1785–1793.

Joyce E. F., M. Pedersen, S. Tiong, S. K. White-Brown, A. Paul, *et al.*, 2011 Drosophila ATM

and ATR have distinct activities in the regulation of meiotic DNA damage and repair. J.

Cell Biol. 195: 359–367.

Kazazian H. H. Jr, C. Wong, H. Youssoufian, A. F. Scott, D. G. Phillips, *et al.*, 1988

Haemophilia A resulting from de novo insertion of *L1* sequences represents a novel

mechanism for mutation in man. Nature 332: 164–166.

Kidwell M. G., and J. F. Kidwell, 1975 Spontaneous male recombination and mutation in

isogenic-derived chromosomes of *Drosophila melanogaster*. J. Hered. 66: 367–375.

Kidwell M. G., 1977 Reciprocal differences in female recombination associated with hybrid

dysgenesis in *Drosophila melanogaster*. Genet. Res. 30: 77–88.

Kidwell M. G., J. F. Kidwell, and J. A. Sved, 1977 Hybrid dysgenesis in *Drosophila*

melanogaster: A syndrome of aberrant traits including mutation, sterility and male

recombination. Genetics 86: 813–833.

Koehler K. E., R. S. Hawley, S. Sherman, and T. Hassold, 1996 Recombination and

nondisjunction in humans and flies. Hum. Mol. Genet. 5 Spec No: 1495–1504.

Kong A., D. F. Gudbjartsson, J. Sainz, G. M. Jonsdottir, S. A. Gudjonsson, *et al.*, 2002 A high-

resolution recombination map of the human genome. Nat. Genet. 31: 241–247.

Koren S., B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman, *et al.*, 2017 Canu: scalable and

accurate long-read assembly via adaptive -mer weighting and repeat separation. Genome

Res. 27: 722–736.

Kulathinal R. J., S. M. Bennett, C. L. Fitzpatrick, and M. A. F. Noor, 2008 Fine-scale mapping

of recombination rate in *Drosophila* refines its correlation to diversity and divergence. Proc.

Natl. Acad. Sci. U. S. A. 105: 10051–10056.

Lasko D., 1991 Loss Of Constitutional Heterozygosity In Human Cancer. Annu. Rev. Genet. 25:

281–314.

Lenth R. V., 2016 Least-Squares Means: The R package lsmeans. J. Stat. Softw. 69.

https://doi.org/10.18637/jss.v069.i01

Le Thomas A., A. K. Rogers, A. Webster, G. K. Marinov, S. E. Liao, *et al.*, 2013 Piwi induces

piRNA-guided transcriptional silencing and establishment of a repressive chromatin state.

Genes Dev. 27: 390–399.

Li H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv preprint arXiv.1303.3997.

Li H., 2013 Minimap2: pairwise-alignment for nucleotide sequences. Bioinformatics 34: 3094–

3100.

Lozovskaya E. R., V. S. Scheinker, and M. B. Evgen'ev, 1990 A hybrid dysgenesis syndrome in

Drosophila virilis. Genetics 126: 619–623.

Lu W.-J., J. Chapo, I. Roig, and J. M. Abrams, 2010 Meiotic recombination provokes functional

activation of the p53 regulatory network. Science 328: 1278–1281.

McCarron M., A. Duttaroy, G. Doughty, and A. Chovnick, 1994 Drosophila *P* element

transposase induces male recombination additively and without a requirement for *P* element

excision or insertion. Genetics 136: 1013–1023.

Mézard C., M. T. Jahns, and M. Grelon, 2015 Where to cross? New insights into the location of

meiotic crossovers. Trends Genet. 31: 393–401.

Miller D. E., C. B. Smith, N. Y. Kazemi, A. J. Cockrell, A. V. Arvanitakas, *et al.*, 2016 Whole-

genome analysis of individual meiotic events in *Drosophila melanogaster* reveals That

noncrossover gene conversions are insensitive to interference and the centromere effect.

Genetics 203: 159–171.

Niki Y., and I. Chigusa, 1986 Developmental analysis of the gonadal sterility of P-M hybrid

dysgenesis in *Drosophila melanogaster*. Jpn. J. Genet. 61: 147–156.

Petrov D. A., J. L. Schutzman, D. L. Hartl, and E. R. Lozovskaya, 1995 Diverse transposable

elements are mobilized in hybrid dysgenesis in *Drosophila virilis*. Proc. Natl. Acad. Sci. U.

S. A. 92: 8050–8054.

Picelli S., A. K. Björklund, B. Reinius, S. Sagasser, G. Winberg, *et al.*, 2014 Tn5 transposase

and tagmentation procedures for massively scaled sequencing projects. Genome Res. 24:

2033–2040.

Pimpinelli S., and P. Ripoll., 1986 Nonrandom segregation of centromeres following mitotic

recombination in *Drosophila melanogaster*. Proc. Natl. Acad. Sci. U. S. A. 83: 3900–3903.

Preston C. R., and W. R. Engels, 1996 *P*-element-induced male recombination and gene

conversion in *Drosophila*. Genetics 144: 1611–1622.

Preston C. R., J. A. Sved, and W. R. Engels, 1996 Flanking duplications and deletions associated

with *P*-induced male recombination in *Drosophila*. Genetics 144: 1623–1638.

Preston C. R., C. C. Flores, and W. R. Engels, 2006 Differential usage of alternative pathways of

double-strand break repair in *Drosophila*. Genetics 172: 1055–1068.

Rizzon C., G. Marais, M. Gouy, and C. Biémont, 2002 Recombination rate and the distribution

of transposable elements in the *Drosophila melanogaster* genome. Genome Res. 12: 400–

407.

Santos T. de los, N. Hunter, C. Lee, B. Larkin, J. Loidl, *et al.*, 2003 The Mus81/Mms4

endonuclease acts independently of double-Holliday junction resolution to promote a

distinct subset of crossovers during meiosis in budding yeast. Genetics 164: 81–94.

Schaeffer S. W., A. Bhutkar, B. F. McAllister, M. Matsuda, L. M. Matzkin, *et al.*, 2008 Polytene

chromosomal maps of 11 *Drosophila* species: the order of genomic scaffolds inferred from

genetic and physical maps. Genetics 179: 1601–1655.

Sienski G., D. Dönertas, and J. Brennecke, 2012 Transcriptional silencing of transposons by Piwi

and maelstrom and its impact on chromatin state and gene expression. Cell 151: 964–980.

Simão F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov, 2015

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics 31: 3210–3212.

Sjöblom T., and J. Lähdetie, 1996 Expression of p53 in normal and gamma-irradiated rat testis

suggests a role for p53 in meiotic recombination and repair. Oncogene 12: 2499–2505.

Slatis H. M., 1955 A reconsideration of the Brown-dominant position effect. Genetics 40: 246–

251.

Slatko B. E., 1978 Parameters of male and remale recombination influenced by the *T-007* second

chromosome in *Drosophila melanogaster*. Genetics 90: 257–276.

Slotkin R. K., R. Keith Slotkin, and R. Martienssen, 2007 Transposable elements and the

epigenetic regulation of the genome. Nat. Rev. Genet. 8: 272–285.

Smukowski Heil C. S., C. Ellison, M. Dubin, and M. A. F. Noor, 2015 Recombining without

Hotspots: A Comprehensive Evolutionary Portrait of Recombination in Two Closely

Related Species of *Drosophila*. Genome Biol. Evol. 7: 2829–2842.

Sokolova M. I., E. S. Zelentsova, N. V. Rozhkov, and M. B. Evgeniev, 2010 Morphologic and

molecular manifestations of hybrid dysgenesis in ontogenesis of *Drosophila virilis*. Russ. J.

Dev. Biol. 41: 391–393.

Sokolova M. I., E. S. Zelentsova, N. G. Shostak, N. V. Rozhkov, and M. B. Evgen'ev, 2013

Ontogenetic consequences of dysgenic crosses in *Drosophila virilis*. Int. J. Dev. Biol. 57:

731–739.

Staten R., S. D. Schully, and M. A. F. Noor, 2004 A microsatellite linkage map of *Drosophila*

mojavensis. BMC Genet. 5: 12.

Stevison L. S., and M. A. F. Noor, 2010 Genetic and evolutionary correlates of fine-scale

recombination rate variation in *Drosophila persimilis*. J. Mol. Evol. 71: 332–345.

Stocker A. J., B. B. Rusuwa, M. J. Blacket, F. D. Frentiu, M. Sullivan, *et al.*, 2012 Physical and

linkage maps for *Drosophila serrata*, a model species for studies of clinal adaptation and

sexual selection. G3 2: 287–297.

Sved J. A., L. M. Blackman, A. S. Gilchrist, and W. R. Engels, 1991 High levels of

recombination induced by homologous *P* elements in *Drosophila melanogaster*. Mol. Gen.

Genet. 225: 443–447.

Sved J. A., L. M. Blackman, Y. Svoboda, and R. Colless, 1995 Male recombination with single

and homologous *P* elements in *Drosophila melanogaster*. Mol. Gen. Genet. 246: 381–386.

Symington L. S., R. Rothstein, and M. Lisby, 2014 Mechanisms and regulation of mitotic

recombination in *Saccharomyces cerevisiae*. Genetics 198: 795–835.

Takano-Shimizu T., 2001 Local changes in GC/AT substitution biases and in crossover

frequencies on *Drosophila* chromosomes. Mol. Biol. Evol. 18: 606–619.

Tarailo-Graovac M., and N. Chen, 2009 Using RepeatMasker to identify repetitive elements in

genomic sequences. Curr Protoc Bioinformatics 25: 4.10.1-4.10.14.

Tasnim S., and E. S. Kelleher, 2018 p53 is required for female germline stem cell maintenance in

P-element hybrid dysgenesis. Dev. Biol. 434: 215–220.

True J. R., J. M. Mercer, and C. C. Laurie, 1996 Differences in crossover frequency and

distribution among three sibling species of *Drosophila*. Genetics 142: 507–523.

Vieira J., C. P. Vieira, D. L. Hartl, and E. R. Lozovskaya, 1998 Factors contributing to the hybrid

dysgenesis syndrome in *Drosophila virilis*. Genet. Res. 71: 109–117.

Walker B. J., T. Abeel, T. Shea, M. Priest, A. Abouelliel, *et al.*, 2014 Pilon: an integrated tool for

comprehensive microbial variant detection and genome assembly improvement. PLoS One

9: e112963.

Weinstein A., 1918 Coincidence of crossing over in *Drosophila melanogaster* (*Ampelophila*).

Genetics 3: 135–172.

Weinstein A., 1920 Homologous genes and linear linkage in *Drosophila virilis*. Proc. Natl. Acad.

Sci. U. S. A. 6: 625–639.

Weismann A., J. Arthur Thomson, and M. R. Thomson, 1904 *The evolution theory*.

Westphal T., and G. Reuter, 2002 Recombinogenic effects of suppressors of position-effect

variegation in *Drosophila*. Genetics 160: 609–621.

Wickham H., 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wylie A., W.-J. Lu, A. D'Brot, M. Buszczak, and J. M. Abrams, 2014 p53 activity is selectively

licensed in the *Drosophila* stem cell compartment. Elife 3: e01530.

Yannopoulos G., N. Stamatis, M. Monastirioti, P. Hatzopoulos, and C. Louis, 1987 *hobo* is

responsible for the induction of hybrid dysgenesis by strains of *Drosophila melanogaster*

bearing the male recombination factor 23.5MRF. Cell 49: 487–495.

Zdobnov E. M., F. Tegenfeldt, D. Kuznetsov, R. M. Waterhouse, F. A. Simão, *et al.*, 2016

OrthoDb v9.1: cataloging evolutionary and funcational annotations for animal, fungal,

plant, archaeal, bacterial, and viral orthologs. Nucleic Acids Res 45: D744–D749.

Zhang J., C. Yu, L. Krishnaswamy, and T. Peterson, 2011 Transposable elements as catalysts for

chromosome rearrangements. Methods Mol. Biol. 701: 315–326.

Zwick M. E., D. J. Cutler, and C. H. Langley, 1999 Classic Weinstein: tetrad analysis, genetic

variation and achiasmate segregation in *Drosophila* and humans. Genetics 152: 1615–1629.

TABLES

Table 1: Genetic map lengths of *D. virilis* chromosomes in centiMorgans reported in previous

studies and this study.

	Chromosome					
	X	**2**	**3**	**4**	**5**	**6**
Gubenko & Evgen'ev (1984)	170	257	145	108	203	1
Huttunen *et al.* (2004)	–	118	125	147	60	–
Current study	143.5	160.9	139.6	148.3	140.0	–

1037

1038   **Table 2:** Interference values (*nu*) and frequency of crossovers created in the non-interference

1039   pathway (*P*) for all chromosomes. Both values were estimated with the Housworth-Stahl model

1040   for the entire dataset.

	Chromosome				
	X	2	3	4	5
Nu	3.22	3.17	2.68	3.09	3.37
P	4.81E-02	1.05E-02	2.73E-08	6.91E-03	9.75E-03

1041

1042 **Table 3:** Pearson's correlation coeffecients (R) and p-values between rates of recombination and

1043 sequence parameters calculated in 250 kb intervals.

Sequence Parameter		Chromosome					Total	Total minus Zero CO
		X	2	3	4	5		
GC Content	R	0.08	0.35	0.11	0.18	0.33	0.23	0.15
	p	0.372	<0.001	0.263	0.079	<0.001	<0.001	<0.001
Gene density	R	0.31	0.21	0.12	0.32	0.33	0.19	0.03
	p	<0.001	0.012	0.222	<0.001	<0.001	<0.001	0.506
Simple repeats	R	0.44	0.43	0.31	0.32	0.54	0.39	0.177
	p	<0.001	<0.001	<0.001	<0.001	<0.001	<1E-10	<0.001
SNP Density	R	0.64	0.553	0.60	0.67	0.65	0.62	0.49
	p	<1E-10	<1E-10	<1E-10	<1E-10	<1E-10	<1E-10	<1E-10
TE Density	R	-0.47	-0.47	-0.33	-0.44	-0.49	-0.44	-0.14
	p	<0.001	<0.001	<0.001	<0.001	<0.001	<1E-10	<0.001

1044

1045 **Table 4:** Clusters of recombination identified in the BC1 progeny.

F1 Female Identifier	Chromosome	Position (Mb from Telomere)	CO Progeny / Total Progeny	Distortion Genotype	TEs detected in Strain 160
3*	X	17.7	4 / 29	None	*Polyphemus*
111	3	25.1	4 / 11	Strain 9**	*Polyphemus*
4013	X	11.4	4 / 13	Strain 160	None
4029	X	21.3	14 / 19	Strain 160	None
5011	3	9.8	28 / 32	None	*Polyphemus*
5019	3	20.2	10 / 30	Strain 9**	None
5022	3	23.6	5 / 11	Strain 9**	*Paris*
5089	X	23.4	4 / 11	Strain 160	*Polyphemus* x2, Paris

1046

1047 * The F1 female is non-dysgenic

1048 ** Segregation distortion occurred in some of the progeny. The rest appear normal and probably

1049 did not inherit a mitotic CO chromatid

FIGURES

Figure 1: The distribution of the total crossover (CO) count observed in individual BC1 progeny

with the mean and standard deviation. The mean for each group is represented with a black dot

and the standard deviation is the black line. A) The distribution of the total CO count per BC1

progeny of low fecundity dysgenic, high fecundity and non-dysgenic F1 mothers. B) The

distribution of CO count per BC1 progeny of each high fecundity dysgenic mother with mean

and standard deviation. Asterisks denotes statistical significance by least square means ($p <$

0.05). Progeny from mother 701 had a higher average CO count than progeny from other

mothers while progeny from mother 4029 exhibited a lower average CO count.

1059

1060

1061 **Figure 2:** The proportion of chromosomes grouped by CO count in BC1 progeny of high fecund

1062 dysgenic, low fecund dysgenic, and non-dysgenic F1 mothers. 95% confidence intervals were

1063 calculated by sampling BC1 progeny by bootstrapping 1000 times.

1064

Figure 3: Loess smoothed splines of the recombination rate along the length of each

chromosome in *D. virilis* from the telomere (left) to the centromere (right) with standard error.

The dotted line represents the centromere effect of recombination suppression as recombination

= 0 from the line to the end of the sequence. The rate of recombination was calculated in 500 kb

intervals in F2 progeny of low fecund dysgenic, high fecund and non-dysgenic F1 mothers for

the A) X chromosome, B) 2nd chromosome, C) 3rd chromosome, D) 4th chromosome, and E)

5th chromosome.

1072

1073

Figure 4: Haplotypes of BC1 progeny from a single high-fecund dysgenic mother. A)

Haplotypes of the third chromosome in progeny of the 4029 F1 mother is typical of most

chromosomes with no cluster of recombination. B) Haplotypes of the third chromosome in

progeny of the 5011 F1 mother identify a common recombination breakpoint in most of the

progeny and reciprocal products of recombination in equal frequency (Binomial test, $p > 0.05$).

C) Haplotypes of the X chromosome in progeny of the 4029 F1 mother indicate a common

recombination breakpoint in half of the progeny and extreme segregation distortion of the distal

portion of the chromosome (227 markers 0.5 - 21.4 MB, Binomial test, $p < 1.6E-08$). The

proximal region of the chromosome shows no segregation distortion (86 markers 21.5 - 29.0 Mb

Binomial test, $p > 0.5$).

1084

1085

Figure 5: Models to explain the clusters of recombination on the third and X chromosomes in

the progeny of two high fecundity dysgenic mothers. In the 5011 F1 female, a mitotic crossover

(CO) either occurred A) prior to DNA replication in the early developing germline resulting in

two daughter cells with the CO or B) after DNA replication, followed by a pattern of Z

segregation so that one daughter cell has both CO chromatids. Oocytes produced by these

germline stem cells will transmit the CO as reciprocal products as seen if the CO occurred in G1.

C) A mitotic CO in the 4029 F1 female occurred after DNA replication in the developing

germline and each daughter cell received one CO chromatid and one non-CO chromatid

according to a pattern of X segregation. This results in a loss of heterozygosity (LOH) in the

distal portion of the chromosome to resemble segregation distortion and recombination events

are not detectable.

1097

1098