1   **Title:** Lipidomic and biophysical homeostasis of mammalian membranes in response to dietary lipids is
2   essential for cellular fitness
3   **Short title:** Physical homeostasis of mammalian membranes
4

Authors: Kandice R Levental*, Eric Malmberg, Jessica Symons, Yang-Yi Fan, Robert S. Chapkin,
Robert Ernst, Ilya Levental*
Authors affiliations: Kandice R Levental[a], Eric Malmberg[a], Jessica Symons[a], Yang-Yi Fan[b], Robert S.
Chapkin[b], Robert Ernst[c], Ilya Levental[a]
[a]University of Texas Health Science Center at Houston, Houston, TX, USA
[b]Program in Integrative Nutrition & Complex Diseases and Department of Nutrition & Food Science,
Texas A&M University, College Station, TX, USA.
[c]Department of Medical Biochemistry and Molecular Biology, Medical Faculty, Saarland University,
66421 Homburg, Germany
14

15 ***Co-corresponding authors:** Dr. Ilya Levental and Dr. Kandice Levental
16 **Address:** Suite 4.202A, 6431 Fannin St, Houston, TX, USA 77030
17 **Phone:** +1–713–500–5566
18 **Fax:** +1–713–500–7456
19 **Email:** ilya.levental@uth.tmc.edu ; kandice.r.levental@uth.tmc.edu
20

21 **Keywords:** membrane fluidity, lipidomics, diet, homeostasis, adaptation, saturated, polyunsaturated, fatty
22 acid, cholesterol, SREBP
23

ABSTRACT

Biological membranes form the functional, dynamic interface that hosts a major fraction of all cellular bioactivity. Proper membrane physiology requires maintenance of a narrow range of physicochemical properties, which must be buffered from external perturbations. While homeostatic adaptation of membrane fluidity to temperature variation is a ubiquitous design feature of ectothermic organisms, such responsive membrane adaptation to external inputs has not been directly observed in mammals. Here, we report that challenging mammalian membrane homeostasis by dietary lipids leads to robust lipidomic remodeling to preserve membrane physical properties. Specifically, exogenous polyunsaturated fatty acids (PUFAs) are rapidly and extensively incorporated into membrane lipids, inducing a reduction in membrane packing. These effects are rapidly compensated both in culture and *in vivo* by lipidome-wide remodeling, most notably upregulation of saturated lipids and cholesterol. These lipidomic changes result in recovery of membrane packing and permeability. This lipidomic and biophysical compensation is mediated in part by lipid regulatory machinery, whose pharmacological or genetic abrogation results in cytotoxicity when membrane homeostasis is challenged by dietary lipids. These results reveal an essential mammalian mechanism for membrane homeostasis wherein lipidome remodeling in response to dietary lipid inputs preserves functional membrane phenotypes.

INTRODUCTION

Lipidic membranes are the essential barriers between life and the abiotic world and also mediate most intracellular compartmentalization in eukaryotic cells. However, the functions of membranes are not limited to passive barriers. Approximately one third of all proteins are membrane-embedded [1], and many more are membrane-associated through post-translational modifications, lipid binding, and protein-protein interactions. Thus, a major fraction of cellular bioactivity occurs at membrane interfaces. Importantly, the physicochemical properties of the lipid matrix are key contributors to membrane physiology. A canonical example is membrane viscosity, which determines protein diffusivity, and thus protein-protein interaction frequency. Another is membrane permeability, which governs the diffusion of solutes into and out of the cytosol. Numerous other membrane physical parameters can determine protein behavior, including but not limited to fluidity, permeability, curvature, tension, packing, bilayer thickness, and lateral compartmentalization [2-12].

Because of their central role in protein function, effective maintenance of membrane properties is essential for cell survival in complex and variable environments. In ectothermic (i.e. non-thermoregulating)

organisms, a pervasive challenge to membrane homeostasis comes from temperature variations. Low temperature reduces the motion of lipid acyl chains, causing membranes to laterally contract, stiffen, and become more viscous [13]. Organisms across the tree of life, from prokaryotes to ectothermic animals, respond to such perturbations by tuning membrane lipid composition, down-regulating tightly packing lipids (e.g. containing saturated acyl chains) and up-regulating more loosely packed ones containing unsaturations or methylations in their lipid tails [13-15]. This response was termed 'homeoviscous adaptation', as these lipid changes result in remarkable constancy in membrane fluidity in spite of variable growth conditions [13,14,16]. It is worth noting that while fluidity is maintained at a specific set point, it should not be assumed that it is either the control variable or the physical property being sensed; rather membrane fluidity may correlate with other membrane/lipid control parameters. For example, membrane homeostasis in *Bacillus subtilis* is mediated by the DesK sensor that is believed to sense membrane thickness [17]. In the yeast endoplasmic reticulum (ER), recently discovered sensors are sensitive specifically to lipid packing [5] and bilayer compressibility [18].

Excepting a few specialized instances (e.g. hibernation), mammals and other warm-blooded animals are not subject to large-scale variations in body temperature; thus, there has been relatively little investigation of homeostatic membrane responsiveness in such organisms. However, it is a well-established but under-appreciated fact that mammalian membrane homeostasis is extensively challenged by dietary inputs. Dietary lipids have major impacts on membrane compositions *in vivo* [19,20], and these perturbations must presumably be buffered to maintain cellular functionality. Mammalian lipidomes are much more complex [21-23] than either bacteria [24] or yeast [25,26], suggesting more potential control nodes required to balance the various conflicting demands of mammalian membrane physiology.

The possibility of a homeoviscous response in mammalian cells was suggested by two studies in the 1970s, which used a spontaneously arising mutant of Chinese Hamster Ovary (CHO) cells that is defective in cholesterol regulation [27,28]. These mutants accumulate cholesterol compared to wild-type CHO cells, but maintain normal membrane fluidity, possibly through modulation of their phospholipid profiles. However, the molecular etiology of the defects in these mutants remains unknown, and it was not reported where the cholesterol in these cells was accumulating (possibly storage organelles or lysosomes). Further, limitations of then-available technologies prevented direct demonstration of lipidomic responses to cholesterol modulation. Thus, the relevance of those insights to physiologically relevant perturbations of metabolically normal mammalian cells remains unclear. More recently, homeoviscous adaptation in mammals has been inferred from data-driven modeling approaches, which used the physical properties (melting temperature, intrinsic curvature) of pure lipids to extrapolate those of complex, biological

membranes [29,30]. However, the inherent non-additivity [31] and non-ideality [32] of lipid mixtures suggests that

extrapolation of physical parameters of complex membranes from pure lipids may not be a reliable

approach. Finally, lipid composition and membrane properties have been implicated in the heat shock

response, though usually with a specific focus on signaling of the proteostasis network [33].

Here, we directly evaluate the hypothesis that mammalian membranes homeostatically adapt to dietary

inputs by characterizing the lipidomic and biophysical responses to dietary fatty acids in cultured

mammalian cells and *in vivo*. Using shotgun mass spectrometry, we show that polyunsaturated fatty acids

(PUFAs) are robustly incorporated into membrane phospholipids, introducing significant biophysical

perturbations. This perturbation is counterbalanced by rapid lipidomic remodeling, most notable in the

upregulation of saturated lipids and cholesterol. This remodeling normalizes membrane permeability and

lipid packing, as evaluated by spectral imaging of the solvatochromic dye C-Laurdan. These responses

are mediated in part by transcriptional sterol-regulatory machinery involving the Sterol Regulatory

Element Binding Protein 2 (SREBP2), whose genetic or pharmacological inhibition abrogated lipidomic

and biophysical homeostasis. Finally, we show that the homeostatic membrane response is essential for

cellular fitness, as un-compensated lipidomic perturbations lead to increased membrane permeability and

non-apoptotic cell death when membrane homeostasis is challenged by exogenous fatty acids.

RESULTS

Robust incorporation of PUFAs into membrane lipids

Recent observations revealed that supplementation of cultured mammalian mast cells (rat basophilic

leukemia cells (RBL)) with docosahexaenoic acid (DHA) leads to robust incorporation of this dietary

polyunsaturated fatty acid into membrane lipids [23]. We observed similar effects in isolated human

mesenchymal stem cells [34], cultured Chinese hamster ovary (CHO) cells, and rat primary hippocampal

neurons (Supp Fig S1), confirming that uptake and incorporation of exogenous DHA into membrane lipids

is not cell-type specific. Supplementation designed to recapitulate DHA-enriched diets in mammals (see

Materials and Methods) increased the fraction of DHA-containing glycerophospholipids (GPLs) by nearly

15-fold (from <1 to ~15 mol%) (Fig 1A). Similarly, supplementation with the more common PUFA ω-6

arachidonic acid (AA) increased the fraction of AA-containing lipids by >3-fold (Fig 1B). Surprisingly,

supplementation with monounsaturated (oleic; OA) acid or saturated (palmitic; PA) fatty acid produced

only minimal lipidomic changes (Supp Fig S2). We expect that this disparity in incorporation is associated

with the availability of these fatty acids in cell culture media[35]: cultured cells have access to sufficient

OA and PA such that supplementation at concentrations used here has no effect, whereas PUFA levels are
limited such that supplementation with the physiologically appropriate concentrations used here leads to
robust uptake and incorporation.

To confirm that the membrane incorporation of supplemented PUFAs in cultured cells appropriately
recapitulates *in vivo* conditions, we analyzed membrane lipidomes in mouse cardiac tissue after two weeks
of *ad libitum*-feeding on a semi-purified diet containing either fish oil (FO) or corn oil (CO), as previously
described [36]. The FO diet is highly enriched in long-chain PUFAs, with ω-3 DHA and eicosapentaenoic
acid (EPA; 20:5) comprising ~20% of all fatty acids. In contrast, CO is poor in long-chain PUFAs (1.3%
FAs with 3+ unsaturations), with ω-6 linoleic acid (18:2) being the major PUFA source (>50% of dietary
FAs) (see Supplement Table 1 for full dietary lipid profile). Consistent with previous observations[36-38] and
the *in vitro* measurements here, dietary fish oil supplementation produced a robust incorporation of ω-3
PUFAs into cardiac membrane lipids, with ~18-fold more ω-3 PUFA-containing membrane lipids in the
FO-fed compared to CO-fed tissues (Fig 1D). As a result, FO-fed mice had a significantly higher
proportion of lipids containing long, highly unsaturated acyl chains (5 or 6 double bonds; Fig 1E), which
resulted in an overall increase in unsaturation of FO-fed tissues (Fig 1F).

Fig 1. Supplemented PUFAs are robustly incorporated into membrane phospholipids *in vitro* and *in vivo*. Supplementation of culture media with PUFAs for 3 days (20 μM) leads to a dramatic increase in (A) ω−3 PUFA-containing membrane lipids in RBL cells supplemented with DHA and (B) ω−6 PUFA-containing membrane lipids supplemented with AA. (C) Both treatments result in significantly increased overall unsaturation of membrane lipids. The unsaturation index reflects a concentration-weighted average lipid unsaturation. (D) Mice fed a diet rich in fish oil (FO) have significantly more lipids containing ω−3 PUFAs compared to corn oil (CO) fed mice. (E) Incorporation of dietary ω−3 PUFAs increases cardiac tissue lipids containing very highly unsaturated (5 or 6 double bonds) acyl chains, resulting in an (F) increase in the overall unsaturation of membrane glycerophospholipids. Individual experiments (A-C) or animals (D-F) are shown. Bars represent mean ± SD. **p<0.01, ***p<0.001 for unpaired t-test compared to untreated. Treatment with saturated (PA) or monounsaturated (OA) fatty acids in these conditions had no effect on the lipidome (see Fig S2).

Lipidome remodeling associated with PUFA incorporation

Despite the copious incorporation of exogenous PUFAs into membrane lipids and the resulting increase in overall membrane unsaturation (Fig 1), cells *in vitro* did not show any obvious toxicity or differences in proliferation at these levels of supplementation (Fig S3). This observation was somewhat surprising in light of the central role of lipid unsaturation in membrane physical properties [39,40] and the putatively critical role of those properties in regulating various cellular processes [3,4,7]. Thus, we hypothesized that mammalian cells may compensate for perturbations from exogenous FAs by remodeling their lipidomes. Indeed, while treatments with OA/PA had no effect on overall cell lipidomes (Supp Fig S2; consistent with their lack of incorporation), both AA and DHA supplementation reduced the abundance of other polyunsaturated (i.e. di- and tri-unsaturated) lipid species (Fig 2A-B). This effect could potentially be explained by replacement of these PUFA-containing lipids by AA/DHA-containing ones. Much more surprising was the highly significant increase in fully saturated lipids resulting from PUFA supplementation (Fig 2A), due to the significantly increased abundance of phospholipid-incorporated saturated fatty acids (Fig 2A inset). This analysis reveals approximately 2-fold more fully saturated lipids resulting from DHA and AA supplementation, respectively (Fig 2A). These changes were associated with a significantly reduced overall unsaturation in phospholipids not containing the supplemented FAs (Fig 2B-C, see figure legend for details). These effects were, at least in part, transcriptionally mediated, as evidenced by mRNA levels of the major fatty acid desaturase enzymes SCD1 and SCD2. As previously reported [41], both were substantially down-regulated by DHA treatment (Fig S4), consistent with more saturated and fewer polyunsaturated lipids. The headgroup profile of membrane lipids was not notably affected by any of the treatments (Supp Fig S5).

This lipidomic remodeling in response to PUFA feeding *in vitro* was not limited to a single cell type. The
observations described so far were on a transformed leukocyte cell line (RBL); however, very similar
DHA-induced lipid changes were observed in cultured CHO cells, primary rat hippocampal neurons,
primary human MSCs, and MSCs that were differentiated *in vitro* into osteoblasts or adipocytes (Fig. 2D).
Not only the broad trends of the lipidomic remodeling (more saturated, less unsaturated lipids) were
common between these disparate cell lineages and sources, but also the magnitude of the effects were
quantitatively similar, despite drastic differences in overall lipid composition [23,34,42,43]. The similarity of
the responses between freshly isolated cells, cultured primary cells, and long-term cultured cells lines
derived from two different germ layers and three different organisms suggests that comprehensive
lipidomic remodeling in response to exogenous PUFAs is a general phenomenon for mammalian cells.

Importantly, this response is also evident *in vivo*. The incorporation of highly polyunsaturated PUFAs
from the diet into membrane lipids in mouse cardiac tissue (Fig 1D-E) led to remarkably similar lipidomic
remodeling as was observed in cultured cells. Namely, we observed that FO fed-mice have ~2-fold more
saturated lipids compared to CO-fed mice, consistent with compensatory, homeostatic lipidomic
remodeling *in vivo* (Fig 2E). The striking differences between FO and CO feeding are notable in light of
the fact that CO diets do contain abundant PUFAs in the form of linoleic acid (18:2). These differences
are potentially due to the unique effects of highly polyunsaturated lipids on membrane properties, which
scale with the degree of unsaturation[43,44]. An alternative, non-exclusive possibility is that highly
unsaturated FAs are preferentially taken up into membrane lipids. These details will be resolved by future
studies with various FA feeding protocols.

Figure 2 – Lipidome remodeling induced by PUFA incorporation. (A) Lipidome-wide remodeling of lipid unsaturation induced by DHA or AA supplementation. Both PUFAs induce significantly increased saturated lipids and decrease lipids containing di- and tri-unsaturated lipids. Inset shows the mol% of fully saturated acyl chains in GPLs. (B and C) The unsaturation index (concentration-weighted average lipid unsaturation) of lipids not containing the supplemented FAs and their derivatives is significantly reduced upon PUFA supplementation. For accurate estimates of lipidomic remodeling, the supplemented FAs and their derivatives are removed from the analysis, as the rather extreme over-abundance of those species upon supplementation suppresses the visualization of compensatory effects (e.g. for DHA supplementation, lipids containing 22:6, 20:5, or 24:6 are removed and then the unsaturation index of the "remaining" lipids is calculated). (D) DHA-mediated lipidomic remodeling, indicated by increased saturated lipids and decreased polyunsaturated lipids (2-5 unsaturations), is consistent across multiple cell types, including isolated rat hippocampal neurons, cultured human MSCs, and MSC differentiated into adipogenic or osteogenic lineages. (E) Lipid unsaturation profile in membrane lipids isolated from murine heart tissue after feeding with CO versus FO. Incorporation of ω-3 PUFAs into membrane lipids (see Fig 1D) is associated with other lipidomic remodeling, namely higher levels of saturated lipids. Inset shows the mol% of fully saturated acyl chains in GPLs. All data are average +/- SD for n ≥ 3 biological replicates. A and E are two-way ANOVA with Sidak's multiple comparison test. B-C and insets are student's t-test compared to untreated. *p<0.05, **p<0.01, ***p<0.001.

Lipidomic remodeling is nearly concomitant with exogenous PUFA incorporation

The remodeling associated with PUFA incorporation into membrane lipids suggested the induction of a homeostatic membrane response, wherein saturated lipids are upregulated to compensate for the fluidizing effect of PUFA-containing species. To support this inference, we analyzed the temporal profiles of PUFA

incorporation and associated lipidomic changes *in vitro*. DHA was quickly incorporated into membrane lipids, with significant increases in DHA-containing lipids observed within 1 h of supplementation and half-time of incorporation of ~4 h (Fig 3A-B). The recovery time course was significantly slower, as wash-out of DHA was followed by a return to baseline with a half-time of ~25 h (Fig 3C). This is the approximate doubling time of RBL cells in culture, suggesting that the "wash-out" effect likely results from dilution by new lipid synthesis rather than directed removal of DHA from membrane lipids. Remarkably, the associated lipidome remodeling proceeded with similar temporal profiles to both DHA supplementation and wash-out (Fig 3D-I). The increase in saturated lipids (Fig 3E) and the decrease in di-unsaturated lipids (Fig 3H) were observed shortly after DHA incorporation and similar time courses were also observed for the wash-out (Fig 3F and 3I). These observations reveal unexpectedly abrupt lipidomic perturbations and suggest that such perturbations induce rapid compensatory responses.

The broad compensatory lipidomic remodeling in response to PUFA supplementation in mammalian cells evokes comparisons to classical observations of homeoviscous adaptation in ectothermic (i.e. non-thermoregulating) organisms [13-16,45,46]. There, perturbations of membrane physical properties produced by changes in ambient temperature induce lipid changes whose apparent purpose is to re-normalize membrane physical properties. Our findings suggest that a similar response to fluidizing stimuli occurs in mammalian cells, where it has been co-opted to cope with perturbations from exogenous lipids (e.g. from the diet).

Figure 3 – Time course of DHA incorporation and lipidome remodeling. (A-C) Time course of DHA incorporation into GPLs
following supplementation, then wash-out. Time course of nearly concomitant (D-F) saturated lipid and (G-I) di-unsaturated lipid
changes induced by DHA supplementation. The gray areas in B, E, and H are shown in extended charts to the right of each
panel to illustrate the timing of the remodeling. All data shown are average +/- SD for n ≥ 3 biological replicates. All fits shown
are first-order kinetic models.

Cholesterol upregulation by DHA via SREBP2

The above-described acyl chain remodeling in response to PUFA supplementation was incomplete, as the
unsaturation index of PUFA-supplemented cells remained higher than those from untreated controls (Fig
1C and 1F). If the purpose of this lipidome response is to normalize membrane properties, the acyl chain
remodeling appears insufficient on its own (lipid headgroup remodeling was minimal; Fig S5C). In
contrast to most prokaryotes wherein membrane homeostasis is mediated largely by phospholipids [14],
membrane properties in eukaryotes are regulated to a very significant extent by sterols [47,48]. Thus, we
hypothesized that the adaptive response to PUFA perturbation in mammalian cells was also mediated by
cholesterol. Indeed, DHA supplementation led to significantly increased membrane cholesterol in both
FO-fed murine heart tissue (Fig 4A) and cultured cells (Fig 4B). In cells, the increase in cholesterol

abundance was again quite rapid, with effects observed within 1 h of DHA introduction and reaching a
peak after ~4 h (Fig 4C), essentially concomitant with the acyl chain remodeling (Fig 3). There was also
evidence of a small but notable overshoot, as the cholesterol increase was somewhat attenuated after ~4
hrs. These observations suggest that cholesterol upregulation is an early and potent response to membrane
perturbation.

The machinery for cholesterol production in metazoans is regulated by proteolytic processing of
transcription factors of the sterol regulatory element binding protein (SREBP) family [49]. Specifically,
signals to upregulate cellular cholesterol levels are translated into proteolysis of a membrane-bound
SREBP2 precursor to release a 'mature' cleaved fragment, which translocates to the nucleus to induce
transcription of various target genes, including those for cholesterol synthesis and uptake [49]. Having
observed a robust and rapid increase in cholesterol levels resulting from DHA supplementation, we
evaluated whether SREBP2 processing was associated with this response. Indeed, DHA feeding increased
the production of the 'mature' transcription factor form of SREBP2, with minimal effect on the precursor
form (Fig 4D and Supp Fig S6). In contrast, the proteolytic processing of SREBP1, the SREBP family
member primarily implicated in unsaturated fatty acid metabolism[49], was not affected by DHA under our
conditions. The relationship between these observations and previous reports of PUFA-regulation of
SREBPs are expanded upon in the Discussion.

Our findings suggested that processing of SREBP2 is involved in homeostatic membrane lipid remodeling
in response to perturbation by PUFA incorporation. To test this inference, we measured the response to
PUFA supplementation in cells where SREBP processing was inhibited either genetically or
pharmacologically. SRD-12B cells are clonal variants of CHO cells wherein a genetic defect in the Site 1
Protease (S1P; cleaves SREBPs to produce the transcriptionally active forms) prevents SREBP
processing/activation [50]. In these cells, no notable upregulation of cholesterol (Fig 4F) or saturated lipids
(Fig S7) was observed upon DHA supplementation. Similarly, chemical inhibition of SREBP activation
by the pentacyclic triterpene betulin[51] abrogated DHA-induced upregulation of cholesterol (Fig 4B,
triangles). It is important to point out that S1P also processes other important signaling proteins like ATF6,
responses that would be abrogated in SRD-12B cells (though likely not by betulin treatment). However,
we did not observe induction of ATF6 proteolytic processing by DHA (Fig S8A). We also observed no
significant upregulation of the ATF6 target gene Grp78. Finally, we did not observe induction of HSP90B
(ER stress response gene) or processing of XBP1, suggesting that ER stress is not significantly induced
by DHA feeding (Fig S8B). While these results support a role for SREBP2 processing in PUFA-induced
lipidome remodeling, they explicitly do not rule out other contributors to the homeostatic response. Nor

do they implicate SREBP2 as the sensor for membrane fluidity. Rather, they establish SREBP2 as an important node of the homeostatic response, and one that can be manipulated to evaluate the cellular consequences of disruptions in membrane homeostasis.

One possible explanation for the unexpectedly fast cholesterol response to early stages of DHA accumulation (e.g. 1 h after supplementation) is that DHA-containing lipids are highly fluidizing, even at relatively minor relative abundances [23]. Further, these are likely produced first in the ER and may accumulate at the site of synthesis to much higher levels than in total cell membranes. This local perturbation may induce a rapid and potent response by the cholesterol biosynthetic machinery, which has non-transcriptional as well as transcriptional regulation [52]. It is possible that the early phase of this response is mediated by non-transcriptional regulation of already-available enzymes, whereas the latter phases (including the acyl chain remodeling and increase in saturated lipids) are dependent on transcriptional upregulation of the various machineries.

Fig 4 – Cholesterol upregulation by DHA supplementation. (A) Cholesterol levels are higher in membranes isolated from murine heart tissue after feeding with FO compared to CO. (B) Membrane cholesterol is significantly increased by DHA supplementation of cultured RBLs. Effect is abrogated by simultaneous treatment with 200 nM betulin. (C) Time course suggests cholesterol increase is a rapid response to DHA-mediated membrane perturbation. (D) The 'mature', transcription-competent form of SREBP2 is rapidly produced in response to DHA supplementation (full representative Western blots show in Fig S6). (E) SREBP1 proteolytic processing is not significantly affected by DHA supplementation. (F) Cholesterol is significantly increased in WT CHO cells treated with DHA; this effect is abrogated in cells with a defect in SREBP activation (SRD12B; S1P-negative).

Physical homeostasis following lipidomic perturbation

The above data reveal rapid and comprehensive lipidomic remodeling resulting from the incorporation of

DHA into membrane lipids, wherein lipids that decrease membrane fluidity / increase membrane packing

(saturated lipids and cholesterol) are rapidly upregulated in response to introduction of PUFA-containing

lipids that increase membrane fluidity. These changes are consistent with lipidome remodeling for

homeostatic maintenance of membrane physical properties. To directly evaluate this inference, we

measured membrane packing in live cells using a fluorescence assay that relies on a solvatochromic dye

(C-Laurdan) whose spectral characteristics are dependent on membrane properties [53]. Specifically, the

emission spectrum of C-Laurdan is red-shifted in loosely-packed membranes (due to the enhanced polarity

of the fluorophore nano-environment), and the relative extent of this spectral shift can be quantified by

ratiometric spectroscopy or imaging [54]. The resulting dimensionless parameter called Generalized

Polarization (GP) is a widely used proxy for membrane packing and fluidity [55-59], with higher values

reporting more tightly packed membranes.

The GP of homogenized cell membranes was not significantly affected by 24 h supplementation with

DHA, nor by betulin alone. In contrast, the combination of DHA+betulin significantly reduced lipid

packing (Fig. 5A). Similarly, while wild-type CHO cells did not show a significant effect of DHA on

membrane fluidity after 24 h treatment, DHA supplementation of SRD-12B cells significantly reduced

GP (Fig 5B). These measurements of lipid packing in homogenized whole cell membranes suggested that

inhibition of lipidomic remodeling prevents the homeostatic maintenance of membrane physical

properties.

To gain deeper temporal and spatial insight into DHA-mediated lipidomic remodeling, we used confocal

spectral imaging [60] to image C-Laurdan GP in cells. Figure 5C shows two-dimensional GP maps of DHA

and/or betulin-treated RBL cells with relatively fluid (low GP; blue/green pixels) internal membranes and

relatively packed plasma membranes (high GP; yellow/red pixels) characteristic of mammalian cells [54,56].

The clearly distinct GP regions enabled us to separately quantify the effects of DHA on plasma versus

internal membranes. Figures 5D-E show the GP histograms from the regions within the dotted lines in Fig

5C, representing internal membranes. The packing of these internal membranes was significantly

decreased shortly after introduction of exogenous DHA (Figs 5F purple circles; pooled data for 2 and 4 h

time points are different from 0 with p < 0.01), suggesting reduced packing consistent with the expected
fluidizing effect of PUFA-containing lipids [23,61]. This response eventually reversed, with GP fully
normalizing to baseline by 24 h (Fig. 5F). Thus, DHA incorporation combined with the associated
remodeling produced no net change in overall membrane properties. GP at the plasma membrane (PM)
was not significantly affected by DHA treatment (Fig 5G). However, we point out that all lipidomic
remodeling effects described above for whole cell membranes were also observable in isolated PMs [23].
Namely, supplemented PUFAs accumulate in PM lipids, which are then apparently compensated for by
increases in saturated lipids and cholesterol (Fig S9). One possible explanation for why PMs showed
smaller biophysical changes than internal membranes is that exogenous PUFAs are preferentially
incorporated into lipids at the ER, which is site of most lipid synthesis. If lipidome remodeling occurs
faster than the trafficking of newly synthesized lipids to the PM, then biophysical disruptions to the PM
may be minimal.

The re-normalization of membrane packing was markedly abrograted when lipidomic remodeling was
inhibited by targeting SREBP processing. Namely, betulin treatment suppressed the recovery of internal
membrane packing following DHA treatment, leading to significantly reduced GP in DHA+betulin-
treated RBL cells after 24 h (Fig 5C-F, orange symbols and line). The persistently increased fluidity
resulting from combined betulin+DHA treatment was observed not only in internal membranes, but also
in the PMs (Fig 5G).

To more precisely validate the biophysical perturbation of the PM, we employed a recently characterized
method that relies on a fluorescent reporter of membrane packing (Di-4-ANEPPDHQ, Di4) that integrates
solely into the outer leaflet of the PM (Fig 5H) [62]. Similar to C-Laurdan, the photophysical characteristics
of Di4 are dependent on lipid packing, with fluorescence emission lifetime providing a particularly
sensitive readout, with higher lifetimes indicative of more tightly packed membranes [62,63]. While neither
DHA nor betulin alone significantly affected Di4 lifetime (τ_{Di4}), plasma membrane packing was
significantly reduced in DHA+betulin-treated RBL cells (Fig 5H-I).

Although biophysical membrane perturbations likely have several distinct physiological effects, we
considered the central role of the plasma membrane as a selective barrier. Significant computational and
model membrane literature[64,65] suggests that membrane permeability is dependent on lipid composition,
with more loosely packed, fluid membranes being more permeable to various polar and amphiphilic
compounds. Simulations have specifically analyzed polyunsaturated phospholipids, showing DHA-
containing membranes are 2-3-fold more permeable than monounsaturated counterparts[64]. Thus, we

hypothesized that the permeability of cell membranes would be increased by incorporation of

polyunsaturated lipids. To test this possibility, we developed a method to quantitatively evaluate

membrane permeability to amphiphilic substrates in living cells. Specifically, we adapted a classical cell

viability assay that relies on the passive diffusion of the amphiphilic compound fluorescein diacetate

(FDA) through the PM into the cytoplasm, where cellular hydrolases convert it into fluorescent fluorescein

whose charge inhibits its diffusion out of the cell[66]. Because the catalytic conversion of FDA is highly

efficient[66], the rate of fluorescence increase is a direct readout of FDA flux through the PM. We observed

a linear increase in fluorescence over at least the first 10 mins of measurement, consistent with a constant

flux (Q), as shown in Fig 5J. The permeability can then be calculated from the flux using Fick's Law (see

Methods). The resulting permeability coefficient for FDA was ~2.2×10^{-6} cm/sec, in good agreement with

previously reported values in plant cells[67]. Finally, we tested the effects of DHA and/or betulin on PM

permeability. Consistent with our hypothesis, cells with uncompensated membrane perturbations (i.e.

DHA+betulin) were indeed more susceptible to passive permeation by FDA (Fig 5K).

Fig 5 – Physical membrane homeostasis is disrupted by inhibition of lipidomic response. (A) Spectroscopy of C-Laurdan-labeled homogenized whole cell membranes shows that neither DHA nor betulin alone affected membrane packing (GP) after 24 h, whereas DHA+betulin significantly reduced membrane packing. *p<0.05, **p<0.01 is calculated by students t-test between groups. (B) Spectroscopy of C-Laurdan-labeled membranes from WT CHO cells revealed no effect of DHA supplementation after 24 h, whereas SRD-12B cells (defective in SREBP processing) had significantly reduced membrane packing. **p<0.01 one-sample t-tests compared to untreated. (C) Exemplary GP maps of RBL cells treated with DHA and/or betulin demonstrate temporal changes in lipid packing (GP). Images show the characteristically tighter lipid packing (i.e. greater GP) of the PM compared to internal membranes. The white dotted lines denote the areas from which the GP of the internal membranes was calculated. GP histograms of internal membranes from images in A are shown in D-E. Leftward shift in histograms denotes less tightly packed, more fluid membranes. (D) In control cells, internal membranes initially fluidized by excess DHA-containing lipids (green) recovered to baseline by 24 h (purple). (E) In contrast, internal membrane packing did not recover in betulin-treated cells. (F-G) Experiments in C-E were repeated at least 4 times and the mean ± SD (normalized to GP without DHA) is shown. Internal membranes are initially fluidized by excess DHA-containing lipids (purple), then recover to baseline by 24 h. In contrast, internal membrane packing does not recover in betulin-treated cells (orange). The effect of treatment was significant (p<0.01)

by 2-way ANOVA, and each time point for betulin-treated cells was significantly different from zero (p<0.01). These conclusions

were independent of the areas used to calculate internal membrane GP (see Methods). (G) DHA supplementation had no

significant effect on packing at the plasma membrane (PM); however, when betulin is added to inhibit lipidomic remodeling, the

PM also becomes significantly more fluid upon DHA supplementation. The effect of betulin treatment in PMs was significant

(p<0.001) by 2-way ANOVA. ΔGP represents GP of treated cells normalized to untreated cells within each experiment. *p<0.05,

**p<0.01, ***p<0.001 in D-E of individual time points is calculated by one-sample t-test compared to ΔGP=0. (H) Fluorescence

lifetime imaging of Di4 to probe membrane packing in the PM. Warmer colors are representative of higher lifetimes, indicative

of more tightly packed membranes. RBL cells were treated for 24 h prior to imaging. (I) Only the combined DHA+betulin treatment

significantly reduced Di4 lifetime, consistent with reduction of PM packing. *p<0.05 compared to untreated using paired t-tests.

(J) Exemplary plots of accumulation of fluorescein fluorescence during incubation with FDA. Shown are average +/- SD of

biological repeats and linear regressions. (K) PM permeability to FDA increases by treatment with DHA+betulin. Average +/-

SD for n ≥ 3 biological replicates. **p<0.01 is calculated by one-sample t-tests compared to untreated.

Disruption of membrane homeostasis induces non-apoptotic cell death

In unicellular organisms, the homeoviscous lipidomic response is believed to be necessary for maintaining

membrane physical properties in a range compatible with life [13,15]. Having observed similar lipidomic and

biophysical responses in mammalian membranes, we hypothesized that this adaptation was necessary for

cellular fitness following membrane perturbations. To test this hypothesis, we evaluated the cytotoxic

effects of DHA under pharmacological or genetic inhibition of homeostatic responses produced by

SREBP2. At low doses (up to 20 μM DHA and 500 nM betulin), neither DHA nor betulin alone showed

significant cytotoxicity; however, DHA supplementation in the presence of betulin led to ~50% decrease

in cell number after 3 days of culture (Fig 6A; full dose-response in Supp Fig S10A). We then tested

whether the cytostatic effect shown in Fig 6A was due to reduced proliferation or increased cell death. We

found that the populations of cells at various parts of cell cycle progression were unaffected by any of the

treatments (Fig S11), suggesting that reduced proliferation rates were not the cause of lower cell numbers.

We also did not observe an accumulation of sub-G0 cells, which are reflective of DNA fragmentation,

often associated with apoptosis. However, the combination of DHA and betulin (but neither treatment

alone) significantly increased the proportion of trypan blue positive (TB+) (i.e. dead) cells (Fig 6B). These

observations were confirmed in SRD-12B cells, whose genetic lesion renders them incapable of activating

SREBP [50] and mounting an adaptive lipidomic and biophysical response to DHA supplementation (Figs

4F and 5G). Consistent with the abrogated homeostatic response, DHA had a significant cytotoxic effect

on SRD-12B cells (Fig 6C-D, green). Thus, the reduced cellular viability resulting from uncompensated

membrane perturbations is due to cytotoxicity.

Consistent with increased cell death, we observed a significantly increased fraction of propidium iodide
positive (PI+) cells after 24 hrs of combined DHA+betulin treatment (Fig 6E and I). Despite the lack of
sub-G0 cells, we speculated that apoptosis was possibly involved in inducing cell death, perhaps resulting
from imbalanced signaling at the PM or membrane stress induced by lipidomic perturbations. One
hallmark of apoptosis is the externalization of PS (marked by Annexin V) without loss of membrane
integrity (marked by PI), allowing apoptotic cells to be identified as AnxV+ / PI- by flow cytometry. After
24 h of treatment with DHA and/or betulin, we observed no significant increases in AnxV+ / PI- cells (Fig
6F-I), suggesting that the increased cell death was not apoptotic. Consistently, using a fluorescent marker
of active caspases (CellEvent Caspase-3/7), we observed no induction of apoptosis in cells treated with
DHA+betulin (Fig S12A). Finally, the involvement of apoptosis in a cytotoxic process can be inferred by
using potent caspase inhibitors (e.g. Z-VAD-FMK), which should attenuate apoptosis-induced
cytotoxicity. Treatment with this inhibitor did not inhibit cell death (Fig S12B), supporting a non-apoptotic
cytotoxicity mechanism.

Together, these results reveal that inhibition of lipidome remodeling in mammalian cells markedly
increased non-apoptotic cell death upon membrane perturbation with dietary fatty acids. Because cell
physiology is essentially dependent on maintaining cytoplasmic concentrations of a plethora of small
molecules (water, alcohols, polyamines) and amphiphiles (steroids, fatty acids) that can passively diffuse
through the membrane, it is possible that the significant increases in membrane permeability (Fig 5H-I)
may be responsible for the observed toxicity.

Fig 6 – Inhibition of homeostatic lipidome remodeling induces non-apoptotic cytotoxicity. (A) Neither DHA nor betulin
alone had a significant cytotoxic effect in RBLs, whereas their combination significantly reduced viability. (B) 24hr of DHA+betulin
treatment significantly increased the number of trypan-positive cells. (C) In SRD-12B cells, which fail to upregulate cholesterol
and saturated lipids upon DHA treatment (see Fig 4D and Supp Fig S6), DHA significantly inhibited cell growth, in contrast to
control WT CHO cells. (D) In SRD-12B cells, DHA significantly increased the proportion of TB+ cells, in contrast to WT CHO
cells. (E) DHA+betulin treatment for 24 hr significantly increased the proportion of PI+ cells. (F-I) Flow cytometry showed no
increase in AnxV+/PI- cells (reflective of apoptosis), while the AnxV+/PI+ cells (non-apoptotic cell death) were increased. Flow
cytometry positive controls shown in Fig S13. All data shown as mean +/- SD for ≥ 3 independent experiments. A, B, and D are
one-sample t-tests compared to untreated; E is student's t-test compared to untreated; C is two-way ANOVA for effect of
treatment. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

DISCUSSION

The incorporation of dietary fatty acids into mammalian membrane lipids has been widely observed both
in vitro [23,68,69] and *in vivo* [19,20,37,38,70]. The fact that exogenous fatty acids are so readily used for lipid
synthesis is unsurprising in light of the fact that the enzyme required for *de novo* fatty acid production
(fatty acid synthase) is minimally expressed in most adult human tissues [71]. Indeed, *de novo* lipogenesis
is considered "negligible" [72,73] in adult humans, suggesting that exogenous sources of fatty acids are the
major raw material for maintenance and replenishment of membrane lipids. This reliance on exogenous

inputs for production of components central to cellular architecture and function would seem to present a

major complication for homeostasis.

In ectothermic organisms, homeostatic membrane control has been widely reported in the form of

homeoviscous adaptation (HVA) [13-15]. According to this hypothesis, the relative abundance of saturated

to unsaturated membrane lipids is responsive to temperature changes in order to maintain membrane

fluidity at the level required for the many processes hosted and regulated by cell membranes. Although

such adaptation has been proposed in prokaryotes [14], single-celled eukaryotes [74], and even cold-blooded

animals [75,76], studies on endothermic organisms have been limited [28,29], and there has yet been no direct

observations of HVA in mammals or isolated mammalian cells. Our data directly confirm all major tenets

of cell-autonomous homeoviscous adaptation: (1) lipidomic remodeling resulting from a perturbation of

membrane physical properties (Fig 2-4); (2) recovery of baseline physical properties at a new lipid

composition (Fig 5); and (3) necessity of this response for cellular fitness (Fig 6). These observations

suggest that mammalian cells possess the capacity for homeostatic membrane adaptation analogous to

HVA, and that this response can compensate for perturbations from dietary lipid inputs.

In cold-blooded animals, the homeostatic membrane response also involves modulation of cholesterol

levels [48,76], and we observe a similar response in mammalian cells, mediated at least partially through

activation of SREBP2 (Fig 4). Disruption of SREBP activation resulted in inhibition of the compensatory

response, both with respect to lipid composition (Figs 3 and 4, and Supp Fig S7) and biophysical properties

(Fig 5). The implication of SREBP2 as a node of the sense-and-respond module for membrane adaptation

is consistent with the central role of SREBPs in membrane homeostasis [49]. SREBP transcription factors

have been dubbed the "master regulators of lipid homeostasis", because they direct not only cholesterol

synthesis and uptake, but also proteins associated with membrane lipid metabolism, including those

involved in FA synthesis, elongation, and desaturation [77]. The effects of PUFA supplementation on

SREBP signaling have been extensively studied, and DHA is known to suppress the activation of SREBP1

and its target genes, both in cultured cells and *in vivo* [78-80]. In contrast, SREBP2 is not suppressed by

PUFAs [78,79], revealing that these two complementary regulators of membrane homeostasis have different

functions, despite both being sensitive to membrane cholesterol. In our observations, SREBP2 is

proteolytically activated as part of the homeostatic response. Previous reports [79] have not noted a

significant effect of PUFAs on SREBP2, possibly because those experiments involved acute PUFA

feeding of serum-starved cells, where high levels of activated SREBPs may have suppressed the DHA-

mediated stimulation we observe. We have confirmed the possibility for such an effect in our cells (Fig

S14). It remains to be determined whether SREBP2 is simply an effector of membrane remodeling

downstream of yet-unidentified sensing machinery, or whether this protein (and/or its regulatory
machinery) may itself be capable of sensing perturbations in membrane physical properties. Direct
demonstration of protein responsiveness to membrane packing has recently been described for two yeast
ER proteins [5,18], providing a conceptual and methodological toolbox for identifying other membrane
sensors. Remarkably, despite the ubiquity and importance of membrane homeostasis, the machinery used
for sensing membrane properties remains largely uncharacterized. It is an intriguing observation that Ire1,
a core component of the unfolded protein response (UPR), exhibits a dual sensitivity to unfolded proteins
and aberrant lipid compositions, demonstrating a tight connection between protein-folding and membrane
properties [18]. This connection suggests that the machinery for ameliorating protein-folding stresses may
also be involved in transducing and mitigating membrane stress [81].

With extensive literature on the effects of dietary lipids on membrane compositions [36,37,82-84], it is somewhat
surprising that the remodeling we observe has not been previously reported. We note that in many of the
published datasets, increases in saturated fatty acids have indeed been observed, though rarely remarked
upon. We speculate that the compensatory effects may have been previously overlooked for several
reasons: (a) remodeling effects are obviously less pronounced than the "main" effects (incorporation and
substitution) and may have been considered incidental/irrelevant; (b) lipid remodeling is temporally
regulated and the time point of sample collection may affect its magnitude [84]; (c) most previous reports
have analyzed isolated FAs rather than intact lipids, which may obscure how these are combined into
phospholipids (e.g. it is possible to have more fully saturated lipids without a major increase in saturated
FAs); (d) remodeling of the non-supplemented FAs can be obscured by the high abundance of the
supplemented FAs. For example, if a large fraction of the phospholipidome is converted into DHA-
containing lipids by DHA feeding, then the mol fraction of other lipids obviously decreases, even if the
relative proportion of saturated acyl chains compared to unsaturated is increased; (e) cholesterol is rarely
considered alongside FAs and is a major component of the remodeling we report.

In summary, our observations support the hypothesis that mammalian cells in culture and *in vivo* sense
membrane physical properties and respond to perturbations by comprehensive remodeling of their
lipidomes. In our measurements, the perturbations were induced by supplementation with polyunsaturated
fatty acids (PUFAs); however, it is likely that other fats (e.g. cholesterol) or amphiphiles (bile acids,
anesthetics) may induce similar responses. We show that the compensatory response is necessary for cell
fitness, confirming membrane adaptation as a central requirement for cellular homeostasis.

MATERIALS AND METHODS

Materials: DHA, AA, OA, PA, betulin, GSK2194069, PF-429242, fluorescein diacetate, and 1,10 phenanthroline were obtained from Sigma Aldrich. C-Laurdan was purchased from TPProbes (South Korea). Amplex Red kit to quantify cholesterol was purchased from Invitrogen. Antibodies used: actin (monoclonal clone AC-15, Abcam), SREBP2 (polyclonal, Abcam), SREBP1 (monoclonal, Abcam). Z-DEVD-FMK and myriocin was purchased from Cayman Chemicals). Di-4-ANEPPDHQ (Di4), CellEvent™ Caspase-3/7 Green Detection Reagent, Annexin V-Pacific Blue, and Hoechst 33342 was purchased from ThermoFisher. Fluorescein sodium salt was purchased from Santa Cruz Biotechnology.

Cell culture: Rat basophilic leukemia (RBL) cells were maintained in medium containing 60% modified Eagle's medium (MEM), 30% RPMI, 10% fetal calf serum, 100 units/mL penicillin, and 100 μg/mL streptomycin. Chinese hamster ovary (CHO) cells were maintained in DMEM:F12 (1:1) containing 5% fetal calf serum, 100 units/mL penicillin, and 100 μg/mL streptomycin. SRD12B cells were maintained in DMEM:F12 (1:1) containing 5% fetal calf serum, 50μM sodium mevalonate, 20μM oleic acid, 5μg/mL cholesterol, 100 units/mL penicillin, and 100 μg/mL streptomycin, as previously described[85]. All cells were grown at 37°C in humidified 5% CO_2.

Fatty acid treatments: Fatty acid stock solutions were received as air-purged ampules (Sigma Aldrich) and loaded into BSA immediately upon opening. BSA loading was accompanied by stirring the fatty acid with BSA dissolved in water (2:1 mol/mol FA:BSA), sterile filtering, purging with nitrogen prior to aliquoting, and storing at -80°C. BSA loading, purging, and cold storage were all done to minimize FA oxidation. For all experiments with FA supplementation, cells were incubated with 20 μM FA for 3 days (cells were re-supplemented 24 h prior to analysis). Serum-containing, rather than serum-free, media was used to approximate physiological supplementation, rather than exclusive introduction of specific FAs. These conditions were chosen to approximate *in vivo* settings produced by DHA-enriched diets in mammals: plasma free FA concentrations range from 300 to 750 μM [70,86] and up to 10 mol % of plasma fatty acids are ω-3 DHA in rats fed a high-fish-oil diet [70]. Further, diets rich in ω-3 PUFAs led to significant incorporation of these fats into cell membrane lipids [19,38,70], similar to the levels we observed under our culture feeding conditions (Fig. 1). For these reasons, we believe our culture conditions reasonably approximate physiological dietary membrane perturbations.

Animals and diets: All experimental procedures using laboratory animals were approved by the Public Health Service and the Institutional Animal Care and Use Committee at Texas A&M University. Pathogen-free female C57BL/6 mice (n=8) weighing 16–18 g, were randomly divided into two groups of 4. For 2 weeks, mice had free access to one of the two semi-purified diets, which were adequate in all nutrients [36]. Diets varied only in the oil composition, i.e., either corn oil (CO) or a PUFA-enriched fish-corn oil (FO) mixture (4:1, w/w) at 5 g/100 g diet. The basic diet composition, expressed as g/100 g was: casein, 20; sucrose, 42; cornstarch, 22; cellulose, 6; AIN-76 mineral mix, 3.5; AIN-76 vitamin mix, 1, DL-methionine, 0.3; choline chloride, 0.2; Tenox 20A, 0.1; and oil, 5. The fatty acid composition of the diets was determined by gas chromatography (Supplemental Table 1). Mice were sacrificed by CO2 asphyxiation after two weeks of feeding the CO/FO diet, and the livers and hearts were frozen in liquid nitrogen. 20-50 mg of tissue was mechanically homogenized in Dulbecco's PBS (without Ca^{2+} and Mg^{2+}). Samples were further diluted to 5 mg/mL in Dulbecco's PBS (without Ca^{2+} and Mg^{2+}), and lipidomics analysis was performed as below.

Drug treatments and cell number quantification: RBL cells were treated with or without 20 μM DHA in the presence or absence of betulin (inhibitor of SREBP processing) or GSK2194069 (FAS inhibitor). Cells were treated for 72 h, and then the cell number determined via fluorescein diacetate (FDA) fluorescence. FDA is a cell viability probe which freely diffuses through cell membranes but is trapped in cells following de-acytelation by cytoplasmic esterases in viable cells. The number of viable cells is then directly related to fluorescein fluorescence intensity. For the assay, cells were treated in 96-well plates, gently washed with PBS, and then incubated with fluorescein diacetate (5 μg/mL in PBS) for 2 minutes at 37°C. The plates were then washed again to remove excess FDA and fluorescence was measured at 488 nm excitation and 520 nm emission. For caspase inhibitor studies, 5 and 30 μM Z-DEVD-FMK were added at the same time as DHA and betulin and treated for 72 h prior to FDA analysis.

To count the number of dead cells, cells were treated for 24 h, and the supernatant harvested. The cells were then trypsinized and combined with the supernatant. 0.4% Trypan Blue was added to an equal volume of cells. The cells were then counted using a Countess II cell counter (Thermo Fisher), and the number of Trypan-positive cells recorded. For each experiment the number of cells per well were normalized to untreated, unsupplemented cells.

Active caspase quantification: RBL cells were treated with or without 20 μM DHA in the presence or

absence of 500nM betulin for 24 h. After 24hr, the supernatant was removed, and CellEvent™ Caspase-

3/7 Green Detection Reagent (7.5 μM, diluted in PBS + 0.5% FBS) was added to each well. The cells

were incubated at 37°C for 30 min. At this time Hoechst 33342 (2.5μg/mL final) was added to the solution

and incubated for an addition 5 min at 37°C. Two images using DAPI and GFP filters were randomly

taken for each condition in three different random locations in the well. For a positive control, cells were

treated with 2 mM N-ethylmaleimde in 50 mM HEPES, 2 mM CaCl2, pH 7.4 for 45 min prior to addition

of the caspase detection reagent. Using ImageJ, to determine positive staining for caspase activity, a

threshold was set using the positive controls, and this threshold was maintained across the experiment.

The number of caspase-positive cells was normalized to the number of cells imaged in the same field of

view by counting the Hoechst-positive nuclei.

Lipidomics: Detailed lipidomic analysis was performed by Lipotype, GmbH, as previously described[87].

Briefly, for preparation of crude cell membranes, cells were washed with phosphate buffered saline (PBS),

scraped in 10 mM Tris pH 7.4, and then homogenized with a 27-gauge needle. Nuclei were then pelleted

by centrifugation at 300 xg for 5 min. The supernatant was pelleted by centrifugation at 100,000 xg for 1

h at 4°C. The membrane pellet was then washed and resuspended in 150 mM ammonium bicarbonate.

All lipidomics were performed at Lipotype GmbH (Dresden, Germany) as described previously [21-23,26,34,88]

and detailed below.

Nomenclature: The following lipid names and abbreviations are used: ceramide (Cer), Chol, SM,

diacylglycerol (DAG), lactosyl ceramide (DiHexCer), glucosyl/galactosyl ceramide (HexCer), sterol ester

(SE), and triacylglycerol (TAG), as well as phosphatidic acid (PA), phosphatidylcholine (PC),

phosphatidylethanolamine (PE), phosphatidylglycerol (PG), and phosphatidylinositol (PI),

phosphatidylserine (PS), and their respective lysospecies (lysoPA, lysoPC, lysoPE, lysoPI, and lysoPS)

and ether derivatives (PC O-, PE O-, LPC O-, and LPE O-). Lipid species were annotated according to

their molecular composition as follows: [lipid class]-[sum of carbon atoms in the FAs]:[sum of double

bonds in the FAs];[sum of hydroxyl groups in the long chain base and the FA moiety] (e.g., SM-32:2;1).

Where available, the individual FA composition according to the same rule is given in brackets (e.g.,

18:1;0-24:2;0).

Lipid standards: Synthetic lipid standards were purchased from Sigma-Aldrich (Chol D6), Larodan

(Solna, Sweden) Fine Chemicals (DAG and TAG), and Avanti Polar Lipids (all others).

Lipid extraction for mass spectrometry lipidomics: Lipids were extracted using a two-step

chloroform/methanol procedure [26]. Samples were spiked with internal lipid standard mixture containing:

cardiolipin 16:1/15:0/15:0/15:0 (CL), ceramide 18:1;2/17:0 (Cer), diacylglycerol 17:0/17:0 (DAG),

hexosylceramide 18:1;2/12:0 (HexCer), lyso-phosphatidate 17:0 (LPA), lyso-phosphatidylcholine 12:0

(LPC), lyso-phosphatidylethanolamine 17:1 (LPE), lyso-phosphatidylglycerol 17:1 (LPG), lyso-

phosphatidylinositol 17:1 (LPI), lyso-phosphatidylserine 17:1 (LPS), phosphatidate 17:0/17:0 (PA),

phosphatidylcholine 17:0/17:0 (PC), phosphatidylethanolamine 17:0/17:0 (PE), phosphatidylglycerol

17:0/17:0 (PG), phosphatidylinositol 16:0/16:0 (PI), phosphatidylserine 17:0/17:0 (PS), cholesterol ester

20:0 (CE), sphingomyelin 18:1;2/12:0;0 (SM), triacylglycerol 17:0/17:0/17:0 (TAG) and cholesterol D6

(Chol). After extraction, the organic phase was transferred to an infusion plate and dried in a speed vacuum

concentrator. 1st step dry extract was re-suspended in 7.5 mM ammonium acetate in

chloroform/methanol/propanol (1:2:4, V:V:V) and 2nd step dry extract in 33% ethanol solution of

methylamine in chloroform/methanol (0.003:5:1; V:V:V). All liquid handling steps were performed using

Hamilton Robotics STARlet robotic platform with the Anti Droplet Control feature for organic solvents

pipetting.

MS data acquisition: Samples were analyzed by direct infusion on a QExactive mass spectrometer

(Thermo Scientific) equipped with a TriVersa NanoMate ion source (Advion Biosciences). Samples were

analyzed in both positive and negative ion modes with a resolution of $R_{m/z=200}$=280000 for MS and

$R_{m/z=200}$=17500 for MS/MS experiments, in a single acquisition. MS/MS was triggered by an inclusion list

encompassing corresponding MS mass ranges scanned in 1 Da increments [88]. Both MS and MS/MS data

were combined to monitor CE, DAG and TAG ions as ammonium adducts; PC, PC O-, as acetate adducts;

and CL, PA, PE, PE O-, PG, PI and PS as deprotonated anions. MS only was used to monitor LPA, LPE,

LPE O-, LPI and LPS as deprotonated anions; Cer, HexCer, SM, LPC and LPC O- as acetate adducts and

cholesterol as ammonium adduct of an acetylated derivative [89].

<u>Lipid identification and quantification:</u> Data were analyzed with in-house developed lipid identification

software based on LipidXplorer (20). Data post-processing and normalization were performed using an

in-house developed data management system. Only lipid identifications with a signal-to-noise ratio >5,

and a signal intensity 5-fold higher than in corresponding blank samples were considered for further data

analysis.

<u>Lipidomics data processing:</u> The lipidomic analysis yields a list of >600 individual lipid species and their

picomolar abundances. These were processed by first transforming into mol% of all lipids detected. Next,

the TAG and sterol esters were removed from the analysis, and the remaining data was analyzed as mol%
of membrane lipids. From here, the datasets were broken down further into class composition. In some
cases, the distribution and structural characteristics (e.g. number of carbons or unsaturations in the acyl
chains) of the individual species were analyzed.

Lipidomic results presentation: Both DHA and AA can be converted into other PUFAs. DHA can be
retro-converted to EPA (20:5) which can then be converted to 24:6. AA can be converted into 22:4 and
24:4. In our supplemented cells, we see small, but non-negligible increases in lipids containing these
converted, supplemented fatty acyl chains. With DHA supplementation, the retroconversion represented
<4.8 mol% GPLs. With AA supplementation, lipids containing converted FAs range from 13-15%. Thus,
ω−3 PUFAs are defind here (Fig 1) as lipids containing either 22:6, 20:5, or 24:6 as an acyl chain. ω−6
PUFAs are lipids which contain 20:4, 22:4, or 24:4 as an acyl chain.

C-laurdan spectroscopy: Membrane packing (related to order and fluidity) was determined via C-laurdan
spectroscopy as described [90,91]. Briefly, cells were washed with PBS and stained with 20 μg/mL C-laurdan
for 10 minutes on ice. The emission spectrum from isolated GPMVs was gathered from 400-550 nm with
excitation at 385 nm at 23°C. The GP was calculated according to the following equation:

$$GP = \frac{\sum_{420}^{460} I_x - \sum_{470}^{510} I_x}{\sum_{420}^{460} I_x + \sum_{470}^{510} I_x}$$

C-Laurdan spectral imaging: C-Laurdan imaging was performed as previously described [23,34,60,90,91].
Briefly, cells were washed with PBS and stained with 10 μg/mL C-Laurdan for 10 min on ice, then imaged
via confocal microscopy on a Nikon A1R with spectral imaging at 60x and excitation at 405 nm. The
emission was collected in two bands: 433–463 nm and 473–503 nm. MATLAB (MathWorks, Natick,
MA) was used to calculate the two-dimensional (2D) GP map, where GP for each pixel was calculated
from a ratio of the two fluorescence channels, as previously described [60]. Briefly, each image was binned
(2x2), background subtracted, and thresholded to keep only pixels with intensities greater than 3 standard
deviations of the background value in both channels. Supplemental Fig 15 shows representative images
of each of these image processing steps. The GP image was calculated for each pixel using the above
equation. GP maps (pixels represented by GP value rather than intensity) were imported into ImageJ. To

ensure that the specifics of our quantification method were not biasing the result, we quantified the GP of
internal membranes by three different methods:

1. Line scans drawn across individual cells, as shown in the original figure. PM GP values were taken as peak
   GP values from the periphery of the cell, whereas internal membranes were calculated as the average of all
   values outside the PM peak.
2. The nucleus is visible as a dark spot in C-Laurdan images. We used this as a fiducial marker to draw a
   region-of-interest representing peri-nuclear membranes and calculated the average GP of those pixels.
3. As shown in the images in Figure 5, the internal membranes were masked, and histograms of the pixels
   inside the mask were plotted.

Ultimately, all three methods produced quantitatively similar results, suggesting that this measurement is
robust. We chose to present the results of the third method here because it includes the most information,
allowing us to evaluate the distributions of pixels GPs.

Di4 fluorescence lifetime imaging: Outer leaflet staining of RBL cell PMs was achieved by incubating
cells at 4°C for 8 min with Di4 at 1 µg/ml in phosphate-buffered saline. Cells were quickly washed prior
to immediate imaging in phenol red-free MEM. Lifetime microscopy was performed on a Nikon A1 laser
scanning microscope with an integrated Picoquant time-correlated single photon counting (TC-SPC)
system (Berlin, Germany). The instrument response function (IRF) was determined with a saturated
erythrosine B and KI solution at pH=10 according to the manufacturer's (Picoquant) protocol. Di4
emission was collected at >560 nm, and images were acquired using 20 MHz pulse frequency. The photon
count rate was kept under 10% of the pulse rate by adjusting a manual shutter, and enough frames were
acquired to obtain at least 8×10^3 photons cumulative signal intensity. The fluorescence decay curves were
fitted to a bi-exponential re-convolution function adjusted to the IRF, and the average lifetime was
calculated and represented in the FLIM images as τ_{Di4}. The experiment was repeated 4 times, imaging
15-30 cells per condition each time.

Amplex Red assay: Amplex Red cholesterol assay was performed (according to manufacturer
instructions; Invitrogen) to determine the abundance of cholesterol. Each reading was normalized to
protein concentration (determined by bicinchoninic acid (BCA) assay) in the same samples. Technical
triplicates were measured for each sample. Shown are values normalized to untreated cells. Reported are
the average and standard deviations from at least 3 independent biological replicates.

Western Blot: Cells were washed with ice cold PBS, then scraped into Laemmli lysis buffer (50mM Tris-
HCl, pH 8.0; 2% SDS; 5mM EDTA, pH8.0) supplemented with protease inhibitor cocktail. Protein
concentration was determined using BCA (Pierce), and equal amounts of protein were mixed with
reducing Laemmli sample buffer and loaded onto SDS-PAGE gels. Gels were transferred to PVDF
membranes, which were blocked in 5% BSA. Membranes were incubated with primary antibodies
overnight at 4°C, and detected with either AlexaFluor or HRP-tagged secondary antibodies. Membranes
were imaged using a BioRad ChemiDoc imager. The intensities of the bands were quantified normalized
to actin, and plotted as mean $\pm$ SD of n $\geq$ 3 experiments.

Real Time Quantitative PCR: Total RNA was isolated via Trizol (Sigma) following the manufacturer's
protocol. Reverse transcriptase PCR was performed using the High Capacity cDNA Reverse Transcription
Kit from Applied Biosystems according to manufacturer's protocol. To quantify mRNA expression,
SYBR Fast MasterMix (2x) Universal Dye (#KK4602) from Kapa Biosystems was used in an Eppendorf
Realplex2 Mastercycler. Each primer set for each sample was run in triplicate with 1 ng of cDNA per
well. The primer sets used are:

Gene	Forward Primer	Reverse Primer
GAPDH	GTCTACTGGCGTCTTCACCA	GTGGCAGTGATGGCATGGAC
SCD1	GTGATGTTCCAGAGGAGGTACT	CAGGAACTCAGAAGCCCAGAA
SCD2	GGTGATGTTCCAGAGGAGGTATT	AACTGGAAGACCCCGAACTC
Grp78	CTACGAAGGTGAACGACCCC	ATTTCTTCAGGGGTCAGGCG
HSP90B	TAAGCTCTATGTGCGCCGAG	TCATCGGAATCCACAACACCT
sXBP1	CTGAGTCCGCAGCAGGTG	TTCCAGCTTGGCTGATGAGG

Expression changes were calculated using the delta delta C_T method.

Permeability assay: RBL cells were treated with or without 20 μM DHA in the presence or absence of
500nM betulin for 24 h. At this time, cells were trypsinized, washed 1x with full serum-containing
medium and 2x with phenol red-free MEM, and counted with a Countess II automated cell counter

(ThermoFisher). 250,000 cells were diluted in 2 mL with phenol red-free MEM and added to a quartz 2

mL cuvette. Just prior to reading on the spectrofluorometer, 2.5 ug/mL fluorescein diacetate (stock 1

mg/mL in acetone) was added to the cells and mixed by pipetting. The fluorescence intensity was read at

450 nm excitation and 510 nm emission over a period of 5 minutes (1 read / second). In each of the samples

measured, the slope of the fluorescence increase over time remained constant over the entire 5 minutes,

consistent with a constant flux (Q).

The permeability coefficient (P) of a membrane to a particular substrate is then defined by Fick's Law as:

$$Q = P * A * (C_{out} - C_{in})$$

where A is the area of the membrane and C_{out} and C_{in} are the concentrations of the solute outside and

inside the cells, respectively. In our experiments, $C_{in} = 0$ because FDA is quickly converted into a different

molecule upon entering the cell. We estimate the transport surface area by multiplying a typical

mammalian cell surface area ($\sim$3000 μm^2) by the number of cells in the cuvette (250,000). Using pure

fluorescein for calibration, we measured directly the flux of FDA molecules across the membrane, which

for a typical sample of untreated RBL cells was 6.6 x 10^{-5} nmol/sec. From these parameters, we calculated

a permeability coefficient for FDA to be $\sim$2.2 x 10^{-6} cm/sec, in excellent agreement with values previously

reported in plant cells[67]. We found that the flux scales linearly with cell and FDA concentrations,

consistent with the simple Fick's Law model of passive diffusion followed by rapid active conversion.

Acknowledgements

All fluorescence microscopy was performed at the Center for Advanced Microscopy, Department of

Integrative Biology & Pharmacology at McGovern Medical School, UTHealth. We gratefully

acknowledge Theodore Steck, Yvonne Lange, and members of the Levental lab for their critical feedback

on this manuscript. Funding for this work was provided by the NIH/National Institute of General Medical

Sciences (GM114282, GM124072, GM120351), the Volkswagen Foundation (grant 93091), and the

Human Frontiers Science Program (RGP0059/2019). All authors have no competing interests.

References

1   Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. Predicting transmembrane protein topology
with a hidden Markov model: application to complete genomes. *Journal of molecular biology* **305**, 567-
580, doi:10.1006/jmbi.2000.4315 (2001).

2   Janmey, P. A. & Kinnunen, P. K. Biophysical properties of lipids and dynamic membranes. *Trends in cell*
biology **16**, 538-546, doi:10.1016/j.tcb.2006.08.009 (2006).
3   Los, D. A. & Murata, N. Membrane fluidity and its roles in the perception of environmental signals.
Biochim Biophys Acta **1666**, 142-157, doi:10.1016/j.bbamem.2004.08.002 (2004).
4   Andersen, O. S. & Koeppe, R. E., 2nd. Bilayer thickness and membrane protein function: an energetic
perspective. *Annu Rev Biophys Biomol Struct* **36**, 107-130,
doi:10.1146/annurev.biophys.36.040306.132643 (2007).
5   Covino, R. *et al.* A eukaryotic sensor for membrane lipid saturation. *Molecular cell* **63**, 49-59,
doi:10.1016/j.molcel.2016.05.015 (2016).
6   daCosta, C. J., Dey, L., Therien, J. P. & Baenziger, J. E. A distinct mechanism for activating uncoupled
nicotinic acetylcholine receptors. *Nat Chem Biol* **9**, 701-707, doi:10.1038/nchembio.1338 (2013).
7   Marsh, D. Protein modulation of lipids, and vice-versa, in membranes. *Biochim Biophys Acta* **1778**, 1545-
1575, doi:10.1016/j.bbamem.2008.01.015 (2008).
8   Roux, A. *et al.* Membrane curvature controls dynamin polymerization. *Proc Natl Acad Sci U S A* **107**,
4141-4146, doi:10.1073/pnas.0913734107 (2010).
9   Shi, Z. & Baumgart, T. Membrane tension and peripheral protein density mediate membrane shape
transitions. *Nature communications* **6**, 5974, doi:10.1038/ncomms6974 (2015).
10   Rosholm, K. R. *et al.* Membrane curvature regulates ligand-specific membrane sorting of GPCRs in
living cells. *Nat Chem Biol* **13**, 724-729, doi:10.1038/nchembio.2372 (2017).
11   Sezgin, E., Levental, I., Mayor, S. & Eggeling, C. The mystery of membrane organization: composition,
regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* **18**, 361-374, doi:10.1038/nrm.2017.16 (2017).
12   Lorent, J. H. *et al.* Structural determinants and functional consequences of protein affinity for membrane
rafts. *Nature communications* **8**, 1219, doi:10.1038/s41467-017-01328-3 (2017).
13   Hazel, J. R. Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation?
Annu Rev Physiol **57**, 19-42, doi:10.1146/annurev.ph.57.030195.000315 (1995).
14   Sinensky, M. Homeoviscous adaptation--a homeostatic process that regulates the viscosity of membrane
lipids in Escherichia coli. *Proc Natl Acad Sci U S A* **71**, 522-525 (1974).
15   Ernst, R., Ejsing, C. S. & Antonny, B. Homeoviscous adaptation and the regulation of membrane lipids.
Journal of molecular biology **428**, 4776-4791, doi:10.1016/j.jmb.2016.08.013 (2016).
16   Guschina, I. A. & Harwood, J. L. Mechanisms of temperature adaptation in poikilotherms. *FEBS Lett*
580, 5477-5483, doi:10.1016/j.febslet.2006.06.066 (2006).
17   Aguilar, P. S., Hernandez-Arriaga, A. M., Cybulski, L. E., Erazo, A. C. & de Mendoza, D. Molecular
basis of thermosensing: a two-component signal transduction thermometer in Bacillus subtilis. *EMBO J*
20, 1681-1691, doi:10.1093/emboj/20.7.1681 (2001).
18   Halbleib, K. *et al.* Activation of the unfolded protein response by lipid bilayer stress. *Molecular cell* **67**,
673-684 e678, doi:10.1016/j.molcel.2017.06.012 (2017).
19   Cao, J., Schwichtenberg, K. A., Hanson, N. Q. & Tsai, M. Y. Incorporation and clearance of omega-3
fatty acids in erythrocyte membranes and plasma phospholipids. *Clinical chemistry* **52**, 2265-2272,
doi:10.1373/clinchem.2006.072322 (2006).
20   Oliveira, T. V., Maniero, F., Santos, M. H., Bydlowski, S. P. & Maranhao, R. C. Impact of high
cholesterol intake on tissue cholesterol content and lipid transfers to high-density lipoprotein. *Nutrition*
27, 713-718, doi:10.1016/j.nut.2010.07.018 (2011).
21   Gerl, M. J. *et al.* Quantitative analysis of the lipidomes of the influenza virus envelope and MDCK cell
apical membrane. *J Cell Biol* **196**, 213-221, doi:10.1083/jcb.201108175 (2012).
22   Sampaio, J. L. *et al.* Membrane lipidome of an epithelial cell line. *Proc Natl Acad Sci U S A* **108**, 1903-
1907, doi:1019267108 [pii]
10.1073/pnas.1019267108 (2011).
23   Levental, K. R. *et al.* Polyunsaturated lipids regulate membrane domain stability by tuning membrane
order. *Biophys J* **110(8)**, 1800-1810, doi:10.1016/j.bpj.2016.03.012 (2016).
24   Han, X. in *Lipidomics*   427-442 (John Wiley & Sons, Inc, 2016).
25   Klemm, R. W. *et al.* Segregation of sphingolipids and sterols during formation of secretory vesicles at the
trans-Golgi network. *J Cell Biol* **185**, 601-612, doi:jcb.200901145 [pii]
10.1083/jcb.200901145 (2009).

26   Ejsing, C. S. *et al.* Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proc*
Natl Acad Sci U S A **106**, 2136-2141, doi:0811700106 [pii]
10.1073/pnas.0811700106 (2009).
27   Sinensky, M. Defective regulation of cholesterol biosynthesis and plasma membrane fluidity in a Chinese
hamster ovary cell mutant. *Proc Natl Acad Sci U S A* **75**, 1247-1249 (1978).
28   Sinensky, M. Adaptive alteration in phospholipid composition of plasma membranes from a somatic cell
mutant defective in the regulation of cholesterol biosynthesis. *J Cell Biol* **85**, 166-169 (1980).
29   Dymond, M. K. Mammalian phospholipid homeostasis: homeoviscous adaptation deconstructed by
lipidomic data driven modelling. *Chem Phys Lipids* **191**, 136-146,
doi:10.1016/j.chemphyslip.2015.09.003 (2015).
30   Dymond, M. K., Hague, C. V., Postle, A. D. & Attard, G. S. An in vivo ratio control mechanism for
phospholipid homeostasis: evidence from lipidomic studies. *J R Soc Interface* **10**, 20120854,
doi:10.1098/rsif.2012.0854 (2013).
31   Sodt, A. J., Venable, R. M., Lyman, E. & Pastor, R. W. Nonadditive Compositional Curvature Energetics
of Lipid Bilayers. *Phys Rev Lett* **117**, 138104, doi:10.1103/PhysRevLett.117.138104 (2016).
32   Almeida, P. F. Thermodynamics of lipid interactions in complex bilayers. *Biochim Biophys Acta* **1788**,
72-85, doi:10.1016/j.bbamem.2008.08.007 (2009).
33   Balogh, G. *et al.* Key role of lipids in heat stress management. *FEBS Lett* **587**, 1970-1980,
doi:10.1016/j.febslet.2013.05.016 (2013).
34   Levental, K. R. *et al.* omega-3 polyunsaturated fatty acids direct differentiation of the membrane
phenotype in mesenchymal stem cells to potentiate osteogenesis. *Science advances* **3**, eaao1193,
doi:10.1126/sciadv.aao1193 (2017).
35   Psychogios, N. *et al.* The human serum metabolome. *PloS one* **6**, e16957,
doi:10.1371/journal.pone.0016957 (2011).
36   Fan, Y. Y., McMurray, D. N., Ly, L. H. & Chapkin, R. S. Dietary (n-3) polyunsaturated fatty acids
remodel mouse T-cell lipid rafts. *The Journal of nutrition* **133**, 1913-1920, doi:10.1093/jn/133.6.1913
(2003).
37   Stark, K. D. *et al.* Fatty acid compositions of serum phospholipids of postmenopausal women: a
comparison between Greenland Inuit and Canadians before and after supplementation with fish oil.
Nutrition **18**, 627-630 (2002).
38   Metcalf, R. G. *et al.* Effects of fish-oil supplementation on myocardial fatty acids in humans. *The*
American journal of clinical nutrition **85**, 1222-1228 (2007).
39   van Blitterswijk, W. J., van der Meer, B. W. & Hilkmann, H. Quantitative contributions of cholesterol
and the individual classes of phospholipids and their degree of fatty acyl (un)saturation to membrane
fluidity measured by fluorescence polarization. *Biochemistry* **26**, 1746-1756 (1987).
40   Lande, M. B., Donovan, J. M. & Zeidel, M. L. The relationship between membrane fluidity and
permeabilities to water, solutes, ammonia, and protons. *J Gen Physiol* **106**, 67-84 (1995).
41   Sessler, A. M., Kaur, N., Palta, J. P. & Ntambi, J. M. Regulation of stearoyl-CoA desaturase 1 mRNA
stability by polyunsaturated fatty acids in 3T3-L1 adipocytes. *J Biol Chem* **271**, 29854-29858 (1996).
42   Levental, I. & Veatch, S. L. The continuing mystery of lipid rafts. *J. Mol. Biol.* **428**, 4749-4764 (2016).
43   Lin, X. *et al.* Domain stability in biomimetic membranes driven by lipid polyunsaturation. *J Phys Chem B*
120, 11930-11941, doi:10.1021/acs.jpcb.6b06815 (2016).
44   Filippov, A., Oradd, G. & Lindblom, G. Domain formation in model membranes studied by pulsed-field
gradient-NMR: the role of lipid polyunsaturation. *Biophys J* **93**, 3182-3190,
doi:10.1529/biophysj.107.111534 (2007).
45   Budin, I. *et al.* Viscous control of cellular respiration by membrane lipid composition. *Science*,
doi:10.1126/science.aat7925 (2018).
46   Brankatschk, M. *et al.* A Temperature-Dependent Switch in Feeding Preference Improves Drosophila
Development and Survival in the Cold. *Dev Cell* **46**, 781-793 e784, doi:10.1016/j.devcel.2018.05.028
(2018).
47   Yeagle, P. L. Cholesterol and the cell membrane. *Biochim Biophys Acta* **822**, 267-287 (1985).
48   Crockett, E. Cholesterol Function in Plasma Membranes from Ectotherms: Membrane-Specific Roles in
Adaptation to Temperature. *Integrative and Comparative Biology* **38**, 298-304 (1998).

49 Raghow, R., Yellaturu, C., Deng, X., Park, E. A. & Elam, M. B. SREBPs: the crossroads of physiological
and pathological lipid homeostasis. *Trends Endocrinol Metab* **19**, 65-73, doi:10.1016/j.tem.2007.10.009
(2008).
50 Sakai, J. *et al.* Molecular identification of the sterol-regulated luminal protease that cleaves SREBPs and
controls lipid composition of animal cells. *Molecular cell* **2**, 505-514 (1998).
51 Tang, J. J. *et al.* Inhibition of SREBP by a small molecule, betulin, improves hyperlipidemia and insulin
resistance and reduces atherosclerotic plaques. *Cell Metab* **13**, 44-56, doi:10.1016/j.cmet.2010.12.004
(2011).
52 Ye, J. & DeBose-Boyd, R. A. Regulation of cholesterol and fatty acid synthesis. *Cold Spring Harbor*
perspectives in biology **3**, doi:10.1101/cshperspect.a004754 (2011).
53 Kim, H. M. *et al.* A two-photon fluorescent probe for lipid raft imaging: C-laurdan. *Chembiochem* **8**, 553-
559, doi:10.1002/cbic.200700003 (2007).
54 Owen, D. M., Rentero, C., Magenau, A., Abu-Siniyeh, A. & Gaus, K. Quantitative imaging of membrane
lipid order in cells and organisms. *Nat Protoc* **7**, 24-35, doi:nprot.2011.419 [pii]
10.1038/nprot.2011.419 (2011).
55 Kaiser, H. J. *et al.* Order of lipid phases in model and plasma membranes. *Proc Natl Acad Sci U S A* **106**,
16645-16650, doi:0908987106 [pii]
10.1073/pnas.0908987106 (2009).
56 Gaus, K., Chklovskaia, E., Fazekas de St Groth, B., Jessup, W. & Harder, T. Condensation of the plasma
membrane at the site of T lymphocyte activation. *J Cell Biol* **171**, 121-131, doi:jcb.200505047 [pii]
10.1083/jcb.200505047 (2005).
57 Parasassi, T., De Stasio, G., Ravagnan, G., Rusch, R. M. & Gratton, E. Quantitation of lipid phases in
phospholipid vesicles by the generalized polarization of Laurdan fluorescence. *Biophys J* **60**, 179-189,
doi:10.1016/S0006-3495(91)82041-0 (1991).
58 Sezgin, E. *et al.* Adaptive lipid packing and bioactivity in membrane domains. *PloS one* **10**, e0123930,
doi:10.1371/journal.pone.0123930 (2015).
59 Levental, K. R. & Levental, I. Giant plasma membrane vesicles: models for understanding membrane
organization. *Current topics in membranes* **75**, 25-57, doi:10.1016/bs.ctm.2015.03.009 (2015).
60 Sezgin, E., Waithe, D., Bernardino de la Serna, J. & Eggeling, C. Spectral imaging to measure
heterogeneity in membrane lipid packing. *Chemphyschem* **16**, 1387-1394, doi:10.1002/cphc.201402794
(2015).
61 Feller, S. E., Gawrisch, K. & MacKerell, A. D., Jr. Polyunsaturated fatty acids in lipid bilayers: intrinsic
and environmental contributions to their unique physical properties. *J Am Chem Soc* **124**, 318-326 (2002).
62 Lorent, J. H. *et al.* The molecular and structural asymmetry of the plasma membrane. *bioRxiv*,
https://doi.org/10.1101/698837, doi:10.1101/698837 %J bioRxiv (2019).
63 Owen, D. M. *et al.* Fluorescence lifetime imaging provides enhanced contrast when imaging the phase-
sensitive dye di-4-ANEPPDHQ in model membranes and live cells. *Biophys J* **90**, L80-82,
doi:10.1529/biophysj.106.084673 (2006).
64 Venable, R. M., Kramer, A. & Pastor, R. W. Molecular Dynamics Simulations of Membrane
Permeability. *Chem Rev* **119**, 5954-5997, doi:10.1021/acs.chemrev.8b00486 (2019).
65 Manni, M. M. *et al.* Acyl chain asymmetry and polyunsaturation of brain phospholipids facilitate
membrane vesiculation without leakage. *eLife* **7**, doi:10.7554/eLife.34394 (2018).
66 Rotman, B. & Papermaster, B. W. Membrane properties of living mammalian cells as studied by
enzymatic hydrolysis of fluorogenic esters. *Proc Natl Acad Sci U S A* **55**, 134-141,
doi:10.1073/pnas.55.1.134 (1966).
67 Baron-Epel, O., Hernandez, D., Jiang, L. W., Meiners, S. & Schindler, M. Dynamic continuity of
cytoplasmic and membrane compartments between plant cells. *J Cell Biol* **106**, 715-721,
doi:10.1083/jcb.106.3.715 (1988).
68 Briolay, A., Jaafar, R., Nemoz, G. & Bessueille, L. Myogenic differentiation and lipid-raft composition of
L6 skeletal muscle cells are modulated by PUFAs. *Biochim Biophys Acta* **1828**, 602-613,
doi:10.1016/j.bbamem.2012.10.006 (2013).
69 Seo, J., Barhoumi, R., Johnson, A. E., Lupton, J. R. & Chapkin, R. S. Docosahexaenoic acid selectively
inhibits plasma membrane targeting of lipidated proteins. *FASEB journal : official publication of the*

Federation of American Societies for Experimental Biology **20**, 770-772, doi:10.1096/fj.05-4683fje
(2006).
70 Yaqoob, P. *et al.* Comparison of the effects of a range of dietary lipids upon serum and tissue lipid
composition in the rat. *The international journal of biochemistry & cell biology* **27**, 297-310 (1995).
71 Kusakabe, T. *et al.* Fatty acid synthase is expressed mainly in adult hormone-sensitive cells or cells with
high lipid metabolism and in proliferating fetal cells. *J Histochem Cytochem* **48**, 613-622,
doi:10.1177/002215540004800505 (2000).
72 Weiss, L. *et al.* Fatty-acid biosynthesis in man, a pathway of minor importance. Purification, optimal
assay conditions, and organ distribution of fatty-acid synthase. *Biol Chem Hoppe Seyler* **367**, 905-912
(1986).
73 Murphy, E. J. Stable isotope methods for the in vivo measurement of lipogenesis and triglyceride
metabolism. *J Anim Sci* **84 Suppl**, E94-104 (2006).
74 Nakagawa, Y., Sakumoto, N., Kaneko, Y. & Harashima, S. Mga2p is a putative sensor for low
temperature and oxygen to induce OLE1 transcription in Saccharomyces cerevisiae. *Biochem Biophys Res*
Commun **291**, 707-713, doi:10.1006/bbrc.2002.6507 (2002).
75 Tiku, P. E., Gracey, A. Y., Macartney, A. I., Beynon, R. J. & Cossins, A. R. Cold-induced expression of
delta 9-desaturase in carp by transcriptional and posttranslational mechanisms. *Science* **271**, 815-818
(1996).
76 Burns, M., Wisser, K., Wu, J., Leventel, I. & Veatch, S. L. Miscibility Transition Temperature Scales
with Growth Temperature in a Zebrafish Cell Line. *Biophys J* **113**, 1212-1222,
doi:10.1016/j.bpj.2017.04.052 (2017).
77 Eberle, D., Hegarty, B., Bossard, P., Ferre, P. & Foufelle, F. SREBP transcription factors: master
regulators of lipid homeostasis. *Biochimie* **86**, 839-848, doi:10.1016/j.biochi.2004.09.018 (2004).
78 Yahagi, N. *et al.* A crucial role of sterol regulatory element-binding protein-1 in the regulation of
lipogenic gene expression by polyunsaturated fatty acids. *J Biol Chem* **274**, 35840-35844 (1999).
79 Hannah, V. C., Ou, J., Luong, A., Goldstein, J. L. & Brown, M. S. Unsaturated fatty acids down-regulate
srebp isoforms 1a and 1c by two mechanisms in HEK-293 cells. *J Biol Chem* **276**, 4365-4372,
doi:10.1074/jbc.M007273200 (2001).
80 Xu, J., Teran-Garcia, M., Park, J. H., Nakamura, M. T. & Clarke, S. D. Polyunsaturated fatty acids
suppress hepatic sterol regulatory element-binding protein-1 expression by accelerating transcript decay. *J*
Biol Chem **276**, 9800-9807, doi:10.1074/jbc.M008973200 (2001).
81 Radanovic, T., Reinhard, J., Ballweg, S., Pesek, K. & Ernst, R. An Emerging Group of Membrane
Property Sensors Controls the Physical State of Organellar Membranes to Maintain Their Identity.
BioEssays : news and reviews in molecular, cellular and developmental biology, e1700250,
doi:10.1002/bies.201700250 (2018).
82 Fan, Y. Y., Ly, L. H., Barhoumi, R., McMurray, D. N. & Chapkin, R. S. Dietary docosahexaenoic acid
suppresses T cell protein kinase C theta lipid raft recruitment and IL-2 production. *J Immunol* **173**, 6151-
6160 (2004).
83 Atkinson, T. G., Barker, H. J. & Meckling-Gill, K. A. Incorporation of long-chain n-3 fatty acids in
tissues and enhanced bone marrow cellularity with docosahexaenoic acid feeding in post-weanling
Fischer 344 rats. *Lipids* **32**, 293-302, doi:10.1007/s11745-997-0036-x (1997).
84 Owen, A. J., Peter-Przyborowska, B. A., Hoy, A. J. & McLennan, P. L. Dietary fish oil dose- and time-
response effects on cardiac phospholipid fatty acid composition. *Lipids* **39**, 955-961, doi:10.1007/s11745-
004-1317-0 (2004).
85 Rawson, R. B., DeBose-Boyd, R., Goldstein, J. L. & Brown, M. S. Failure to cleave sterol regulatory
element-binding proteins (SREBPs) causes cholesterol auxotrophy in Chinese hamster ovary cells with
genetic absence of SREBP cleavage-activating protein. *Journal of Biological Chemistry* **274**, 28549-
28556 (1999).
86 Pirro, M. *et al.* Plasma free fatty acid levels and the risk of ischemic heart disease in men: prospective
results from the Quebec Cardiovascular Study. *Atherosclerosis* **160**, 377-384 (2002).
87 Tulodziecka, K. *et al.* Remodeling of the postsynaptic plasma membrane during neural development. *Mol*
Biol Cell **27**, 3480-3489, doi:10.1091/mbc.E16-06-0420 (2016).

88   Surma, M. A. *et al.* An automated shotgun lipidomics platform for high throughput, comprehensive, and
quantitative analysis of blood plasma intact lipids. *European journal of lipid science and technology :*
EJLST **117**, 1540-1549, doi:10.1002/ejlt.201500145 (2015).
89   Liebisch, G. *et al.* High throughput quantification of cholesterol and cholesteryl ester by electrospray
ionization tandem mass spectrometry (ESI-MS/MS). *Biochim Biophys Acta* **1761**, 121-128,
doi:10.1016/j.bbalip.2005.12.007 (2006).
90   Sezgin, E. *et al.* Elucidating membrane structure and protein behavior using giant plasma membrane
vesicles. *Nat Protoc* **7**, 1042-1051, doi:10.1038/nprot.2012.059 (2012).
91   Levental, K. R. & Levental, I. Isolation of giant plasma membrane vesicles for evaluation of plasma
membrane structure and protein partitioning. *Methods in molecular biology* **1232**, 65-77,
doi:10.1007/978-1-4939-1752-5_6 (2015).
981