Title: Homeostatic remodeling of mammalian membranes in response to dietary lipids is essential for cellular

fitness

Short title: Homeostasis of mammalian membranes

Authors: Kandice R Levental, Eric Malmberg, Robert Ernst, Ilya Levental

Authors affiliations: Kandice R Levental[a], Eric Malmberg[a], Robert Ernst[b], Ilya Levental[a]

[a]University of Texas Health Science Center at Houston, Houston, TX, USA

[b]Department of Medical Biochemistry and Molecular Biology, Medical Faculty, Saarland University, 66421

Homburg, Germany

Corresponding author: Dr. Ilya Levental

Address: Suite 4.202A, 6431 Fannin St, Houston, TX, USA 77030

Phone: +1–713–500–5566

Fax: +1–713–500–7456

Email: ilya.levental@uth.tmc.edu

Keywords: membrane fluidity, lipidomics, diet, homeostasis, adaptation, saturated lipid, polyunsaturated fatty

acid, cholesterol, SREBP

Abstract: A major fraction of cellular bioactivity occurs at membranes, with the lipidic matrix constituting a

functional, dynamic interface that regulates protein activity and cell physiology. Proper membrane functionality

requires maintenance of a narrow range of physical properties, which must be buffered from external

perturbations. The most prominent example of such maintenance is homeostatic adaptation of membrane viscosity

to temperature variation, a fundamental and ubiquitous design feature of ectothermic organisms. However, such

responsive membrane adaptation has not been directly observed in mammalian cells. Here, we report that

challenging membrane homeostasis in mammalian cells by dietary lipids leads to robust lipidomic remodeling to

preserve membrane physical properties. Specifically, exogenous polyunsaturated fatty acids (PUFAs) are rapidly

and extensively incorporated into membrane lipids, inducing a reduction in membrane packing. These effects are

rapidly compensated by upregulation of saturated lipids and cholesterol via activation of classical mammalian

sterol regulatory machinery, which results in recovery of membrane packing. Inhibition of this compensatory

response results in decreased cellular fitness when membrane homeostasis is challenged by dietary lipids. These

results reveal an essential mammalian mechanism for membrane homeostasis - analogous to homeoviscous

adaptation - wherein cells remodel their lipidomes in response to dietary lipid inputs in order to preserve

functional membrane phenotypes.

ABSTRACT

A major fraction of cellular bioactivity occurs at membranes, with the lipidic matrix constituting a functional, dynamic interface that regulates protein activity and cell physiology. Proper membrane functionality requires maintenance of a narrow range of physical properties, which must be buffered from external perturbations. The most prominent example of such maintenance is homeostatic adaptation of membrane viscosity to temperature variation, a fundamental and ubiquitous design feature of ectothermic organisms. However, such responsive membrane adaptation has not been directly observed in mammalian cells. Here, we report that challenging membrane homeostasis in mammalian cells by dietary lipids leads to robust lipidomic remodeling to preserve membrane physical properties. Specifically, exogenous polyunsaturated fatty acids (PUFAs) are rapidly and extensively incorporated into membrane lipids, inducing a reduction in membrane packing. These effects are rapidly compensated by upregulation of saturated lipids and cholesterol via activation of classical mammalian sterol regulatory machinery, which results in recovery of membrane packing. Inhibition of this compensatory response results in decreased cellular fitness when membrane homeostasis is challenged by dietary lipids. These results reveal an essential mammalian mechanism for membrane homeostasis - analogous to homeoviscous adaptation - wherein cells remodel their lipidomes in response to dietary lipid inputs in order to preserve functional membrane phenotypes.

INTRODUCTION

Lipidic membranes are the essential barriers between life and the abiotic world and also mediate most intracellular compartmentalization in eukaryotic cells. However, the role of membranes is not limited to passive barriers. Approximately one third of all proteins are membrane-embedded [1], and many more are membrane-associated through post-translational modifications, lipid binding, and protein-protein interactions. Thus, a major fraction of cellular bioactivity occurs at or near membrane interfaces. Importantly, the physicochemical properties of the membrane matrix are key contributors to physiology. A fundamental example is membrane viscosity, which determines protein diffusivity, and thus protein-protein collision/interaction frequency. Another is membrane permeability which governs the diffusion of solutes between the cytosol and the external environment. In addition to these examples, numerous other membrane physical parameters can determine protein behavior, including but not limited to membrane fluidity, permeability, curvature, tension, packing, bilayer thickness, and lateral organization [2-12].

Because of their central role in protein function, the proper maintenance of membrane properties is essential for survival in a complex and variable environment. In ectothermic (i.e. non-thermoregulating) organisms, a pervasive challenge to membrane homeostasis comes in the form of temperature variations. Low temperature

reduces the motion of lipid acyl chains, causing membranes to laterally contract, stiffen, and become more viscous [13]. Organisms across the tree of life, from prokaryotes to ectothermic animals, respond to such perturbations by tuning membrane lipid composition, down-regulating tightly packing lipids (e.g. containing saturated acyl chains) and up-regulating more loosely packed ones containing unsaturations or methylations in their lipid tails [13-15]. This response was termed 'homeoviscous adaptation', as these lipid changes result in remarkable constancy in membrane viscosity in spite of variable growth conditions [13,14,16]. It should be noted that while viscosity is maintained, it must not be assumed that it is either the control variable, or the physical property being sensed; rather viscosity could correlate with other membrane/lipid control parameters. For example, membrane homeostasis in Bacillus subtilis is mediated by the DesK sensor that is thought to respond to membrane thickness [17]. In the yeast endoplasmic reticulum (ER), recently discovered sensors are sensitive specifically to lipid packing [5] and bilayer compressibility [18]. These key insights are exceptions to the rule, as there remains remarkably little known about the molecular mechanisms underlying the sensing of cellular membranes [15,16].

Besides a few specialized instances (e.g. hibernation), mammals and other warm-blooded animals are not subject to large-scale variations in body temperature; thus, there has been relatively little investigation of homeostatic membrane responsiveness in such organisms. However, it is a well-established but under-appreciated fact that mammalian membrane homeostasis is challenged by dietary inputs. Dietary cholesterol and fatty acids have major impacts on membrane compositions *in vivo* [19,20], and these perturbations must presumably be buffered by cellular mechanisms to maintain functionality. Mammalian lipidomes are much more complex [21-23] than either bacteria [24] or yeast [25,26], suggesting many more potential control mechanisms; however, the physiological purpose of mammalian lipidomic complexity remains opaque.

The possibility of a mammalian homeoviscous response was hypothesized from a set of studies in the 1970s, which used a spontaneously arising mutant of Chinese Hamster Ovary (CHO) cells that is defective in cholesterol regulation [27,28]. These mutants accumulate cholesterol compared to wild-type CHO cells, but maintain normal membrane fluidity, possibly through modulation of their phospholipid profiles. However, the molecular etiology of the defects in these mutants remains unknown, and it was not reported where the cholesterol in these cells was accumulating (possibly membranes, but also storage organelles or lysosomes). Further, limitations of then-available technologies prevented direct demonstration of lipidomic responses to cholesterol modulation. Thus, the relevance of those insights to physiologically relevant perturbations of metabolically normal mammalian cells remains unclear. More recently, homeoviscous adaptation in mammals has been inferred from data-driven modeling approaches, which used the physical properties (melting temperature, intrinsic curvature) of pure lipids to extrapolate those of complex, biological membranes [29,30]. However, the inherent non-additivity [31] and non-ideality [32] of lipid mixtures suggests that extrapolation of physical parameters of complex membranes from pure

lipids may not be a reliable approach. Finally, lipid composition and membrane properties have been implicated

in the heat shock response, though usually with a specific focus on signaling in the proteostasis network [33].

Here, we directly evaluate the homeoviscous adaptation hypothesis for mammalian membranes by characterizing

the consequences of membrane perturbations by dietary fatty acids in several mammalian cell types. We show

that polyunsaturated fatty acids (PUFAs) are specifically and robustly incorporated into membrane phospholipids,

introducing significant biophysical perturbations. Mammalian cells respond to this perturbation by nearly

concomitant lipidomic remodeling, most notably upregulation of saturated lipids and cholesterol, yielding

normalization of membrane biophysical properties. This response is centrally mediated by transcriptional sterol-

regulatory machinery involving SREBP2. Finally, we show that the homeostatic membrane response is essential

for cellular fitness, as abrogation of key response nodes leads to cytostatic effects when membrane homeostasis is

challenged by exogenous fatty acids.

RESULTS

Specific and robust incorporation of PUFAs into membrane lipids

Recent observations revealed that docosahexaenoic acid (DHA) supplementation of cultured mammalian mast

cells leads to robust incorporation of this dietary polyunsaturated fatty acid into membrane lipids [23]. We observed

similar effects in isolated human Mesenchymal Stem Cells [34], cultured Chinese hamster ovary (CHO) cells, and

rat primary hippocampal neurons (Supp Fig S1), suggesting that uptake and incorporation of exogenous DHA into

membrane lipids is not cell-type specific. To further characterize the effect, we investigated the extent,

destination, and time-course of exogenous FA incorporation into membrane lipids of a model mammalian cell

type. In rat basophilic leukemia cells (RBL), the fraction of DHA-containing glycerophospholipids (GPLs)

increased by nearly 15-fold (from <1 to ~15 mol%) upon supplementation of cell culture media with 20 µM DHA

(Fig 1A). The supplementation parameters (concentration and delivery vehicle) were selected to recapitulate

physiologically relevant conditions observed *in vivo* under DHA-enriched diets in mammals: ω-3 DHA

comprises up to 10 mol% of plasma fatty acids in rats fed a fish oil-rich diet [35], in which overall plasma free FA

concentrations range from 300 to 750 µM [35,36], suggesting that tens of µM are physiologically appropriate.

Further, ω-3 PUFA-rich diets lead to significant incorporation of these fats into cell membrane lipids *in vivo*

[19,35,37], with DHA-containing lipids reaching levels similar to those observed under our culture supplementation

conditions (Fig. 1A)[23]. Serum-containing, rather than serum-free, media was used to approximate physiological

supplementation, rather than exclusive introduction of specific FAs.

Similar to DHA, supplementation with the more common PUFA ω-6 arachidonic acid (AA) increased the fraction

of AA-containing lipids by >3-fold (Fig 1B). Surprisingly, supplementation with monounsaturated (oleic; OA)

acid or saturated (palmitic; PA) fatty acid produced only minimal lipidomic changes (Fig 1C-D). We speculate

that this disparity in incorporation between PUFAs and more saturated FAs is associated with their availability in

cell culture media. That is, cells have access to sufficient levels of OA and PA such that supplementation has no

effect, whereas PUFA levels are limited such that supplementation with the physiologically appropriate

concentrations used here leads to robust uptake and incorporation.

The detailed lipidomics data provided by MS/MS reveals the destination of the incorporated PUFAs. For DHA,

there was notable channeling into phosphatidylethanolamine (PE) headgroup lipids, particularly those with an

ether-linked hydrophobic tail (PE-O; plasmalogen). In un-supplemented cells, ~70% of all DHA was associated

with a PE headgroup, despite PE lipids comprising only 25% of all GPLs. Supplementation slightly reduced this

effect, although PE lipids still contained the majority of DHA (Fig 1E). These trends were markedly different for

AA (Fig 1F), whose distribution more closely resembled the overall GPL headgroup profile (Supp Fig S2).

Somewhat surprisingly, AA was nearly completely segregated from phosphatidylserine (PS) and

phosphatidylglycerol (PG) headgroups, whereas DHA was not. These trends were also generally evident in

primary human mesenchymal stem cells [34].

Fig 1. PUFAs are robustly and specifically incorporated into membrane phospholipids. Supplementation of culture

media with (A) DHA or (B) AA for 3 days (20 μM) leads to robust enhancement of the levels of these PUFAs in membrane

lipids. In contrast, supplementation with either (C) OA or (D) PA did not have observable effects on the lipidomes. (E) Before

supplementation, DHA is highly non-randomly distributed among membrane phospholipids, enriching in PE-headgroup

classes, most notably PE plasmalogen. Supplementation broadened the distribution of DHA into other major GPLs, though

notable PE and PE-O remained. (F) AA distribution was distributed somewhat more randomly among headgroups, though with

some enrichment in PI and PE. These observations reveal a notably precise targeting of w-3 PUFAs into alkyl-linked PE lipids.

Lipidome remodeling associated with PUFA incorporation

A surprising finding from these experiments was that despite the copious incorporation of exogenous PUFAs into

membrane lipids (Fig 1A-B) and the resulting increase in overall membrane unsaturation (Fig 2B), the cells did

not show any obvious toxicity or differences in proliferation. This observation was unexpected in light of the

central role of lipid unsaturation in membrane physical properties [38,39] and the critical role of those properties in

regulating various cellular processes [3,4,7]. Thus, we hypothesized that cells may be compensating for perturbations

from exogenous FAs by remodeling their lipidomes. Indeed, while treatments with OA/PA had no effect on the

overall cell lipidomes (Supp Fig S3; consistent with their lack of incorporation), both AA and DHA

supplementation led to a reduced abundance of other polyunsaturated (i.e. di- and tri-unsaturated) lipid species

(Fig 2C). This effect could potentially be explained by replacement of one PUFA by another. Much more

surprising was the highly significant increase in fully saturated lipids resulting from PUFA supplementation,

which was due in part to significantly increased abundance of phospholipid-incorporated saturated fatty acids (Fig

2D).

The increase in saturated lipids is particularly striking when lipids containing the supplemented FAs are removed

from the analysis (e.g. ω-3-containing lipids from DHA-supplemented data), as the rather extreme over-

abundance of those species upon supplementation suppresses the visualization of compensatory effects. This

analysis reveals approximately 2-fold and 4-fold more fully saturated lipids resulting from DHA and AA

supplementation, respectively (Fig 2E-F). These changes were associated with a significant net reduction in the

unsaturation index of GPLs not containing the supplemented FAs (Fig 2G-H). These effects were, at least in part,

transcriptionally mediated, as evidenced by mRNA levels of the major fatty acid desaturase enzymes SCD1 and

SCD2. As previously observed [40], both were substantially down-regulated by DHA treatment (Supp Fig S4), fully

consistent with more saturated and fewer polyunsaturated lipids.

A similar compensatory effect was observed for lipid length, i.e. the total number of carbons in lipid acyl chains.

Both AA (20-carbon) and DHA (22-carbon) are relatively long FAs, and thus their incorporation increased the

overall length of GPLs in the membranes (Supp Fig S5 A-B). However, there was also a notable increase in

relatively short non-DHA/AA containing lipids (Supp Fig S5 C-D). Surprisingly, the headgroup profile of GPLs

was not notably affected by any of the treatments (Fig 2A and Supp Fig S3A).

Importantly, the lipidomic remodeling in response to PUFA feeding was not limited to a single cell type. The

observations described so far were made using a transformed leukocyte cell line (RBL); however, very similar

DHA-induced lipid changes were observed in cultured CHO cells, primary rat hippocampal neurons, primary

human MSCs, and MSCs that were differentiated *in vitro* into osteoblasts or adipocytes (Fig. 2I). Remarkably, not

only the broad trends of the lipidomic remodeling (more saturated, less unsaturated lipids) were common between these disparate cell lineages and sources, but also the magnitude of the effects were quantitatively similar, despite drastic differences in overall lipid composition [23,34,41,42]. The similarity of the responses between freshly isolated cells, cultured primary cells, and long-term cultured cells lines derived from two different germ layers and three different organisms suggests that comprehensive lipidomic remodeling in response to exogenous PUFAs is a general phenomenon for mammalian cells.

Figure 2 – Lipidome remodeling induced by PUFA supplementation. (A) Incorporation of PUFAs into membrane lipids has no effect on glycerophospholipids (GPL) headgroup distribution. (B) PUFA supplementation increases membrane unsaturated index. (C) In addition to robust incorporation of AA and DHA, fully saturated lipids are increased while other polyunsaturated lipids are reduced. (D) PUFA supplementation increased abundance of GPL-incorporated saturated fatty acids. (E-F) The compensatory effects in Fig 2C can be more clearly visualized by removing the lipid species containing the supplemented PUFAs from the analysis. (G-H) The unsaturation index of non-DHA/AA-containing lipids is significantly decreased upon PUFA supplementation. (I) DHA-mediated lipidomic remodeling, as indicated by increased saturated lipids and decreased polyunsaturated lipids (2-5 unsaturations), is consistent across multiple cell types, including isolated rat hippocampal neurons,

cultured human MSCs, and MSC differentiated into adipogenic or osteogenic lineages. All data shown are average +/- SD for n ≥ 3 biological replicates. C, E, and F are two-way ANOVA with Sidak's multiple comparison test. B, D, G, and H are student's t-test compared to untreated. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Temporal characteristics of lipidomic remodeling

The remodeling associated with PUFA incorporation into membrane lipids suggested the induction of a homeostatic membrane response, wherein saturated lipids are upregulated to compensate for the fluidizing effect of PUFA-containing species. To support this inference, we analyzed the temporal profiles of PUFA incorporation and associated lipidomic changes. DHA was rapidly incorporated into membrane lipids, with significant increases in DHA-containing lipids observed within 1 h of supplementation and half-time of incorporation of ~4 h (Fig 3A-B). Interestingly, the recovery time course was significantly slower, as wash-out of DHA was followed by a return to baseline with a half-time of ~25 h (Fig 3C). This is the approximate doubling time of RBL cells in culture, suggesting that the "wash-out" effect likely results from dilution by new lipid synthesis rather than directed removal of DHA from membrane lipids. Remarkably, the associated lipidome remodeling showed nearly identical temporal profiles to both DHA supplementation and wash-out (Fig 3D-I). The increase in saturated lipids (Fig 3E) and the decrease in di-unsaturated lipids (Fig 3H) were essentially concomitant with DHA incorporation and similar time courses were also observed for the wash-out (Fig 3F and 3I). These observations reveal unexpectedly rapid lipidomic changes to membrane lipid perturbations and suggest that such perturbations induce nearly simultaneous compensatory responses.

The nature of the broad compensatory lipidomic remodeling in response to PUFA supplementation in mammalian cells evokes classical observations of homeoviscous adaptation in ectothermic (e.g. non-thermoregulating) organisms [13-16], wherein perturbations of membrane physical properties produced by changes in ambient temperature are rapidly compensated for by lipidomic changes designed to re-normalize membrane physical properties. Our findings suggest that a similar response to fluidizing stimuli occurs in mammalian cells, where it has been co-opted to cope with perturbations from exogenous lipids (e.g. from the diet).

Figure 3 – Time course of DHA incorporation and lipidome remodeling. (A-C) Time course of DHA incorporation into GPLs following supplementation, then wash-out. Time course of concomitant (D-F) saturated lipid and (G-I) di-unsaturated lipid changes induced by DHA supplementation. All data shown are average +/- SD for n ≥ 3 biological replicates.

Cholesterol upregulation by DHA via SREBP2

The above-described acyl chain remodeling in response to PUFA supplementation was incomplete, as the unsaturation index of PUFA-supplemented cells remained higher than those from untreated controls (Fig 2B). If the lipidome response we observe were designed to normalize membrane properties, the acyl chain remodeling appears insufficient on its own. In contrast to most prokaryotes where membrane homeostasis is mediated almost exclusively by phospholipids [14], membrane properties in more complex organisms are regulated to a very significant extent by sterols [43,44]. Thus, we hypothesized that the adaptive response to PUFA perturbation in mammalian cells was also mediated by cholesterol. Indeed, DHA supplementation significantly increased cellular cholesterol over the un-supplemented baseline (Fig 4A). This increase in cholesterol abundance was again quite rapid, with effects observed within 1 h of DHA introduction and reaching a peak after ~4 h (Fig 4B), essentially concomitant with the acyl chain remodeling (Fig 3). There was also evidence of a small but notable overshoot, as the cholesterol increase was somewhat attenuated after ~4 hrs. These observations suggest that cholesterol upregulation is an early and potent response to membrane perturbation.

The machinery for cholesterol production in metazoans is regulated by proteolytic processing of transcription
factors of the sterol regulatory element binding protein (SREBP) family [45]. A signal to upregulate cellular
cholesterol level is translated into proteolysis of a membrane-bound SREBP2 precursor to release a 'mature'
cleaved fragment, which translocates to the nucleus to induce transcription of various target genes, including
those for cholesterol synthesis and uptake [45]. Having observed a robust and rapid increase in cholesterol levels
resulting from DHA supplementation, we evaluated whether SREBP2 processing is associated with this response.
Indeed, DHA feeding increased the production of the 'mature' transcription factor form of the SREBP2, with
minimal effects on the precursor form (Fig 4C).

These findings suggested that processing of SREBP2 is involved in homeostatic membrane remodeling in
response to perturbation of mammalian lipid composition by PUFA incorporation. To confirm this inference, we
measured the response to PUFA supplementation in cells lacking functional SREBP processing machinery.
Specifically, we used SRD-12B cells, clonal variants of CHO cells wherein a genetic defect in the Site 1 Protease
(S1P; cleaves SREBPs to produce the transcriptionally active forms) prevents SREBP processing/activation [46]. In
these cells, no notable upregulation of cholesterol (Fig 4D) or saturated lipids (Supp Fig S6) was observed upon
DHA supplementation, confirming the crucial role for SREBP signaling in lipidomic remodeling.

Fig 4 – Cholesterol upregulation by DHA supplementation. (A) Membrane cholesterol is increased by DHA supplementation. (B) Time course suggests cholesterol increase is a rapid response to DHA-mediated membrane perturbation. (C) The 'mature', transcription-competent form of SREBP2 is rapidly produced in response to DHA supplementation. (D) Cholesterol is significantly increased in WT CHO cells treated with DHA; however, in cells with abrogated SREBP activation (SRD12B; S1P-negative), there is no change in cholesterol levels with DHA supplementation. All data shown are average +/- SD for n ≥ 3 biological replicates; significances are one-sample t-test compared to untreated; *p<0.05, **p<0.01.

Order homeostasis following lipidomic perturbation

The above data reveal rapid and comprehensive lipidomic remodeling resulting from the incorporation of DHA into membrane lipids. Lipids that putatively decrease membrane fluidity / increase membrane packing (saturated lipids and cholesterol) are rapidly upregulated in response to introduction of PUFA-containing lipids that should increase membrane fluidity. These observations are consistent with lipidome remodeling for the purpose of homeostatic maintenance of membrane physical properties. To directly evaluate this inference, we measured membrane packing in live cells using a fluorescence assay that relies on a solvatochromic dye (C-Laurdan) whose spectral characteristics are dependent on membrane properties [47]. Specifically, the emission spectrum of C-Laurdan is red-shifted in loosely-packed membranes (due to the enhanced polarity of the fluorophore nano-environment), and the relative extent of this spectral shift can be quantified by ratiometric spectroscopy or imaging [48]. The resulting dimensionless parameter called Generalized Polarization (GP) is a widely used proxy for membrane packing and fluidity [49-52].

The image in Fig 5A shows a map of C-Laurdan GP in live cells generated by confocal spectral imaging [53]. This image reveals the relatively fluid (low GP; blue/green pixels) internal membranes and relatively packed plasma membranes (high GP; yellow/red pixels) characteristic of mammalian cells [48,50] (see trace right). These clearly distinct regions enabled us to separately quantify the effects of DHA on plasma versus internal membranes (Fig 5B-C). In internal membranes, we observed significantly decreased GP shortly after introduction of exogenous DHA (purple circles; pooled data for 2 and 4 h time points are different from 0 with p < 0.01), suggesting reduced packing consistent with the known fluidizing effect of PUFA-containing lipids [23,54]. This response eventually reversed, with fluidity returning fully to baseline at 24 h. These effects were not observed in PMs, which did not significantly diverge from baseline, and actually showed a slight trend toward increased packing with DHA (Fig 5C), possibly suggesting that PM physical properties are regulated more tightly than those of internal membranes. Thus, in both internal and PMs, DHA incorporation combined with the associated remodeling produced no net change in overall membrane properties. These microscopic observations were confirmed by spectroscopic evaluation of membrane packing in crude membrane fractions were isolated from RBL cells (Supp Fig S7).

Consistent with the microscopic measurements, there was no significant difference in C-Laurdan GP between
DHA-treated and untreated membranes after time was allowed for compensation.

Fig 5 – Physical membrane homeostasis. (A) Exemplary GP map of several RBL cells showing the characteristically greater
GP (i.e. tighter lipid packing) of the PM compared to internal membranes. (right) The trace is a line scan of GP values across a
single cell shown by the white line in the image. (B) Temporal changes in lipid packing (GP) after DHA supplementation as a
function of betulin. In control cells, internal membranes are initially fluidized by excess DHA-containing lipids (purple), then
recovered to baseline by 24 hrs. In contrast, internal membrane packing did not recover in betulin-treated cells (orange). Effect
of treatment was highly significant ($p<0.01$) by 2-way ANOVA; each time point for betulin-treated cells was significantly
different from zero ($p<0.01$). (C) The effect of DHA on PMs was minimal compared to internal membranes, though betulin-
treated cells were significantly different from controls ($p<0.001$ by 2-way ANOVA). All data shown are average +/- SD for $n \geq 3$
biological replicates.

Disruption of membrane homeostasis reduces cell fitness

In unicellular organisms, the homeoviscous lipidomic response is necessary for maintaining membrane physical
properties in a range compatible with the biochemical processes necessary for life [13,15]. Having observed similar
lipidomic and biophysical responses in mammalian membranes, we hypothesized that this adaptation was
necessary for cellular fitness under conditions of membrane stress. To test this hypothesis, we evaluated the
cytostatic effects of DHA upon chemical inhibition of key nodes in the membrane response machinery,
specifically by inhibiting SREBP processing using the pentacyclic triterpene betulin [55]. This drug effectively
inhibited DHA-induced upregulation of cholesterol (Fig 6A) and saturated lipids (Supp Fig S8), and therefore

prevented the recovery of membrane packing (Fig 5B, orange squares; and Supp Fig S7), confirming the central
role of SREBPs in the adaptive membrane response to DHA. More importantly, while neither DHA nor betulin
alone showed significant cytostatic effects, together they led to ~50% decrease in cell number after 3 days of
combined treatment (Fig 6C; full dose-response in Supp Fig S9). Similar effects were observed for inhibition of
fatty acid synthase (by 10nM GSK2194069; Fig 6D), which is presumably necessary for production of fatty acids
towards the phospholipid remodeling described in Figures 2-3.

These observations were confirmed in the SRD-12B cells, which have a genetic lesion that renders them
incapable of activating SREBP [46]. Figs 4D and Supp Fig S6 show that these cells failed to upregulate cholesterol
and saturated lipids in response to DHA. Consistently, these cells were also unable to mount an adaptive
membrane response with a significant reduction in membrane packing induced by DHA (Fig 6E). Finally, while
DHA slightly promoted the growth of wild-type CHO cells (Fig 6F, white), it had a marked cytostatic effect on
SRD-12B cells (Fig 6F, green). These results reveal that inhibition of lipidome remodeling in mammalian cells
dramatically reduced cellular fitness upon membrane perturbation with dietary fatty acids.

Fig 6 – Inhibition of homeostatic lipidome remodeling reduces cell fitness under DHA supplementation. (A) 200nM
betulin completely abrogated DHA-induced upregulation of cholesterol and (B) saturated lipids. (C) No significant inhibition of
cell number was observed by treating with either DHA or betulin (0.5 μM) alone, whereas the combination significantly reduced
cell numbers. (D) A similar synergistic effect was observed with DHA and GSK2194069 (FAS inhibitor; 10 nM). (E) In SRD-
12B cells, which fail to upregulate cholesterol and saturated lipids upon DHA treatment (see Fig 4D and Supp Fig S6), DHA
significantly reduced membrane packing, and (F) inhibited cell growth, in contrast to control WT CHO cells, wherein DHA
showed a slightly mitogenic effect. All data shown as average +/- SD for ≥3 independent experiments. A-E are student's t-
tests; F is two-way ANOVA for effect of treatment. *p<0.05, **p<0.01, ***p<0.001.

DISCUSSION

DISCUSSION

The incorporation of dietary fatty acids into mammalian membrane lipids has been extensively observed both *in*
vitro [23,56,57] and *in vivo* [19,20,35,37,58]. The fact that exogenous fatty acids are so readily used for lipid synthesis is
unsurprising in light of the fact that the enzyme required for *de novo* fatty acid production (fatty acid synthase) is
expressed at minimal levels in most adult human tissues [59]. Indeed, *de novo* lipogenesis is considered "negligible"
[60,61] in adult humans, suggesting that exogenous sources of fatty acids are the major raw material for maintenance
and replenishment of membrane lipids. This reliance on exogenous inputs for production of components so
central to cellular architecture and function presents a major complication for mammalian membrane homeostasis.

In ectothermic organisms, such homeostatic membrane control has been widely observed in the form of
homeoviscous adaptation (HVA), which is a ubiquitous response of ectotherm cell membranes to temperature
variations [13-15]. According to this canonical hypothesis, the relative abundance of saturated to unsaturated
membrane lipids is responsive to temperatures changes in order to maintain membrane fluidity at the level
required for the many processes hosted and regulated by cell membranes. Although such adaptation has been
extensively characterized in prokaryotes [14], single-celled eukaryotes [62], and even cold-blooded animals [63,64],
studies on endothermic organisms have been limited [28,29], and there has yet been no direct observations of HVA in
mammalian cells. Our data directly confirm the three major tenets of cell-autonomous homeoviscous adaptation:
(1) lipidomic remodeling resulting from a perturbation of membrane physical properties (Fig 2-4); (2) recovery of
baseline physical properties at a new lipid composition (Fig 5); and (3) necessity of this response for cellular
fitness (Fig 6). These observations suggest that mammalian cells possess the capacity for homeostatic membrane
adaptation analogous to HVA, and that this response can be used to compensate for perturbations from dietary
lipid inputs.

In cold-blooded animals, the homeostatic membrane response also involves modulation of cholesterol levels [44,64],
and we observe a similar response in mammalian cells, mediated at least partially through activation of SREBP2
(Fig 4). It is quite remarkable to note that disruption of SREBP activation either by betulin or in SRD-12B cells
resulted in near-complete abrogation of the compensatory response. This abrogation was notable not only in the
lack of cholesterol upregulation (Fig 4D and 6A), which may have been expected, but also for upregulation of
saturated lipids (Figs S6 and S8), a response not obviously connected to SREBP target genes.

Our implication of SREBP2 as a critical node of the sense-and-respond module for membrane adaptation is fully
consistent with the central role of SREBPs in membrane homeostasis [45]. SREBP transcription factors have been
dubbed the "master regulators of lipid homeostasis", because they affect not only cholesterol synthesis and
uptake, but also proteins associated with membrane lipid metabolism, including those involved in FA synthesis,

elongation, and desaturation [65]. The effect of PUFA supplementation on SREBP function has been extensively

studied, and DHA is known to suppress the activation of SREBP1 and its target genes, both in cultured cells and

in vivo [66-68]. We observed downregulation of two SREBP1 target genes (SCD1 and SCD2, Supp Fig S4)

suggesting a similar effect in our cells. In contrast, SREBP2 is not suppressed by PUFAs [66,67], revealing that these

two complementary regulators of membrane homeostasis have different functions, despite both being sensitive to

membrane cholesterol. In our observations, SREBP2 is induced as a necessary part of the homeostatic response.

Previous measurements have not noted a significant effect of PUFAs on SREBP2, possibly because those

experiments involved acute PUFA feeding of serum-starved cells, where higher levels of activated SREBPs may

have suppressed the DHA-mediated stimulation we observe here. It remains to be determined whether SREBP2 is

simply an effector of membrane remodeling downstream of yet-unidentified sensing machinery, or whether this

protein (and/or its regulatory machinery) may itself be capable of sensing perturbations in membrane physical

properties. Direct demonstration of protein responsiveness to membrane packing has recently been described for

two yeast ER proteins [5,18], providing a conceptual and methodological toolbox for identifying other membrane

sensors. Remarkably, despite the ubiquity and importance of the homeoviscous response, the machinery used for

sensing membrane properties remains almost completely uncharacterized. It is an intriguing observation that Ire1,

a core component of the unfolded protein response (UPR), exhibits a dual sensitivity to unfolded proteins and

aberrant lipid compositions, demonstrating a tight connection between protein-folding and membrane properties

[18]. This connections suggests that the machinery for ameliorating protein-folding stresses may also be involved in

transducing and mitigating membrane stress [69].

In summary, our observations strongly support the hypothesis that mammalian cells generally possess the

machinery for sensing membrane physical properties and responding to perturbations by comprehensive

remodeling of their lipidomes. In our measurements, the major perturbations were induced by physiologically

relevant supplementation with polyunsaturated fatty acids (PUFAs); however, it is likely that other dietary fats

(e.g. cholesterol) and amphiphiles (bile acids, anesthetics) may induce similar responses. We implicate SREBP2

as a central response mediator, consistent with the key role of SREBPs in membrane homeostasis. Finally, we

show that this response is important for cell fitness, confirming membrane adaptation as a central requirement for

cellular homeostasis. Remarkably, detailed analysis of published fatty acid profiles of rodent[70,71] and human[58]

tissues suggests that fully saturated lipids are indeed upregulated *in vivo* following polyunsaturated acid feeding.

Although this observation was not extensively discussed in the above-referenced studies, it directly supports our

conclusions that mammalian cells adapt their lipidomes in response to dietary perturbations, including *in vivo*.

MATERIALS AND METHODS

Materials: Betulin and GSK2194069 were obtained from Sigma Aldrich. C-Laurdan was purchased from
TPProbes (South Korea). Amplex Red kit to quantify cholesterol was purchased from Invitrogen. Antibodies
used: actin (monoclonal clone AC-15, Abcam), SREBP2 (polyclonal, Abcam).

Cell culture: Rat basophilic leukemia (RBL) cells were maintained in medium containing 60% modified Eagle's
medium (MEM), 30% RPMI, 10% fetal calf serum, 100 units/mL penicillin, and 100 μg/mL streptomycin. CHO
cells were maintained in DMEM:F12 (1:1) containing 5% fetal calf serum, 100 units/mL penicillin, and 100
μg/mL streptomycin. SRD12B cells were maintained in DMEM:F12 (1:1) containing 5% fetal calf serum, 50μM
sodium mevalonate, 20μM oleic acid, 5μg/mL cholesterol, 100 units/mL penicillin, and 100 μg/mL streptomycin.
All cells were grown at 37°C in humidified 5% CO2.

Fatty acid treatments: Fatty acid stock solutions were received as air-purged ampules (Sigma Aldrich) and loaded
into BSA immediately upon opening. BSA loading was accompanied by stirring the fatty acid with BSA
dissolved in water (2:1 mol/mol FA:BSA), sterile filtering, purging with nitrogen prior to aliquoting, and storing
at -80°C. BSA loading, purging, and cold storage were all done to minimize FA oxidation. For all experiments
with FA supplementation, cells were incubated with 20 μM FA for 3 days (cells were re-supplemented 24 h prior
to analysis). This concentration was chosen to approximate conditions observed *in vivo* under DHA-enriched diets
in mammals: plasma free FA concentrations range from 300 to 750 μM [35,36] and up to 10 mol % of plasma fatty
acids are ω-3 DHA in rats fed a high-fish-oil diet [35]. Further, diets rich in ω-3 PUFAs led to significant
incorporation of these fats into cell membrane lipids [19,35,37], similar to the levels we observed under our culture
feeding conditions (Fig. 1). For these reasons, we believe our culture conditions reasonably approximate
physiological dietary membrane perturbations.

Drug treatments and cell number quantification: RBL cells were treated with or without 20 μM DHA in the
presence or absence of betulin (inhibitor of SREBP processing) or GSK2194069 (FAS inhibitor). Cells were
treated for 72 h and then the cell number determined via fluorescein diacetate (FDA) fluorescence. FDA is a cell
viability probe which freely diffuses through cell membranes, but is trapped in cells following de-acytelation by
cytoplasmic esterases in viable cells. The number of viable cells is then directly related to fluorescein
fluorescence intensity. For the assay, cells were treated in 96-well plates, gently washed with PBS, and then
incubated with fluorescein diacetate (5 μg/mL in PBS) for 2 minutes at 37°C. The plates were then washed again
to remove excess FDA and fluorescence was measured at 488 nm excitation and 520 nm emission. For each
experiment the number of cells per well were normalized to untreated, unsupplemented cells.

Lipidomics: Detailed lipidomic analysis was performed by Lipotype, GmbH, as previously described[72]. Briefly,
for preparation of crude cell membranes, cells were washed with phosphate buffered saline (PBS), scraped in 10
mM Tris pH 7.4, and then homogenized with a 27-gauge needle. Nuclei were then pelleted by centrifugation at

300 xg for 5 min. The supernatant was pelleted by centrifugation at 100,000 xg for 1 h at 4^{o}C. The membrane

pellet was then washed and resuspended in 150 mM ammonium bicarbonate.

Lipids detected: ceramide (Cer), Chol, SM, diacylglycerol (DAG), lactosyl ceramide (DiHexCer),

glucosyl/galactosyl ceramide (HexCer), sterol ester (SE), and triacylglycerol (TAG), as well as phosphatidic acid

(PA), phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylglycerol (PG), and

phosphatidylinositol (PI), phosphatidylserine (PS), and their respective lysospecies (lysoPA, lysoPC, lysoPE,

lysoPI, and lysoPS) and ether derivatives (PC O-, PE O-, LPC O-, and LPE O-). Lipid species were annotated

according to their molecular composition as follows: [lipid class]-[sum of carbon atoms in the FAs]:[sum of

double bonds in the FAs];[sum of hydroxyl groups in the long chain base and the FA moiety] (e.g., SM-32:2;1).

When available, the individual FA composition according to the same rule is given in brackets (e.g., 18:1;0-

24:2;0)

Lipid extraction: Samples were extracted and analyzed as previously described (39,47), which is a modification

of a previously published method for shotgun lipidomics (48). Briefly, membrane samples were suspended in 150

µL of 150 mM ammonium bicarbonate in water, spiked with 20 µL of internal standard lipid mixture, and then

extracted with 750 µL of a chloroform/methanol 10:1 (v/v) mixture for 2 h at 4_C with 1400 rpm shaking. After

centrifugation (3 min, 3000 **g**) to facilitate phase partitioning, the lower, lipid-containing, organic phase was

collected (first-step extract), and the remaining water phase was further extracted with 750 µL of a

chloroform/methanol 2:1 (v/v) mixture under the same conditions. Again the lower, organic phase was collected

(second-step extract). Extracts were dried in a speed vacuum concentrator, and 120 µL of a dried first-step extract

underwent acetylation with 75 µL of an acetyl chloride/chloroform 1:2 (v/v) mixture for 1 h to derivatize Chol.

After the reaction was completed, the mixture was dried. Then, 120 µL of a dried first-step extract and a

derivatized extract were resuspended in an acquisition mixture with 8 mM ammonium acetate (400 mM

ammonium acetate in methanol/chloroform/methanol/propan-2-ol, 1:7:14:28, v/v/v/v). Next, 120 µL of the

second-step extract was resuspended in an acquisition mixture with 30 µL 33% methylamine in methanol, in 60

mL methanol/chloroform 1:5 (v/v). All liquid-handling steps were performed using a Hamilton (Reno, NV)

STARlet robotic platform with the Anti Droplet Control feature, ensuring the accuracy and reproducibility of

organic solvent pipetting.

Lipid standards: Synthetic lipid standards were purchased from Sigma-Aldrich (Chol D6), Larodan (Solna,

Sweden) Fine Chemicals (DAG and TAG), and Avanti Polar Lipids (all others). The standard lipid mixtures were

chloroform/ methanol 1:1 (v/v) solutions containing Cer 35:1;2, (D18:1;2, 17:0;0); Chol D6; DAG 34:0;0 (17:0;0,

17:0;0); DiHexCer 30:1;2 (D18:1;2.12:0;0); HexCer 30:1;2 (D18:1;2.12:0;0); LPA 17:0;0 (17:0;0); LPC 12:0;0

(12:0;0); LPE 17:1;0 (17:1;0); LPI 17:1;0 (17:1;0); LPS 17:1;0 (17:1;0); PA 34:0;0 (17:0;0, 17:0;0); PC 34:0;0

(17:0;0, 17:0;0); PE 34:0;0 (17:0;0, 17:0;0); PG 34:0;0 (17:0;0, 17:0;0); PI 32:0;0 (16:0;0, 16:0;0); PS 34:0;0

(17:0;0, 17:0;0); SE 20:0;0 (20:0;0); SM 30:1;2 (18:1;2, 12:0;0); and TAG 51:0;0 (17:0;0, 17:0;0, 17:0;0).

Lipid spectrum acquisition: Extracts in the acquisition mixtures were infused with a robotic nanoflow ion source

(TriVersa NanoMate; Advion Biosciences, Ithaca, NY) into a mass spectrometer instrument (Q Exactive, Thermo

Scientific). Cer, DiHexCer, HexCer, lysolipids, and SM were monitored by negative ion mode Fourier transform

mass spectrometry (FT-MS). PA, PC, PE, PI, PS, and ether species were monitored by negative ion mode FT

tandem MS (FT-MS/MS). Acetylated Chol was monitored by positive ion mode FTMS. SE, DAG, TAG, and

species were monitored by positive ion mode FT-MS/MS.

Lipid identification and quantification: Automated processing of acquired mass spectra, and identification and

quantification of detected molecular lipid species were performed with the use of LipidXplorer software (17).

Only lipid identifications with a signal/noise ratio of >5, an absolute abundance of at least 1 pmol, and a signal

intensity fivefold higher than in the corresponding blank samples were considered for further data analysis.

C-laurdan spectroscopy:. Membrane packing (related to order and fluidity) was determined via C-laurdan

spectroscopy as described [73,74]. Briefly, cells were washed with PBS and stained with 20 μg/mL C-laurdan for 10

minutes on ice. The emission spectrum from isolated GPMVs was gathered from 400-550 nm with excitation at

385 nm at 23°C. The GP was calculated according to the following equation:

$$GP = \frac{\sum_{420}^{460} I_x - \sum_{470}^{510} I_x}{\sum_{420}^{460} I_x + \sum_{470}^{510} I_x}$$

C-Laurdan spectral imaging: C-Laurdan imaging was performed as previously described [23,34,53,73,74]. Briefly, cells

were washed with PBS and stained with 10 μg/mL C-Laurdan for 10 min on ice, then imaged via confocal

microscopy on a Nikon A1R with spectral imaging at 60x and excitation at 405 nm. The emission was collected

in two bands: 433–463 nm and 473–503 nm. MATLAB (MathWorks, Natick, MA) was used to calculate the two-

dimensional (2D) GP map, where GP for each pixel was calculated from a ratio of the two fluorescence channels,

as previously described [53]. Briefly, each image was binned (2x2), background subtracted, and thresholded to keep

only pixels with intensities greater than 3 standard deviations of the background value in both channels. The GP

image was calculated for each pixel using the above equation. GP maps (pixels represented by GP value rather

than intensity) were imported into ImageJ. Line scans were drawn across individual cells. PM GP values were

calculated as averages of the peaks as values higher than 0.26. For internal membranes, average GP values were

calculated from line scans around the nucleus (visible as a dark spot in C-Laurdan images). At least 10 cells were

imaged and quantified per experiment with at least 4 individual experiments performed. Reported are the mean

and standard deviations of the average GP values for the individual experiments.

Amplex Red assay: Amplex Red cholesterol assay was performed (according to manufacturer instructions; Invitrogen) to determine the abundance of cholesterol. Each reading was normalized to protein concentration (determined by bicinchoninic acid (BCA) assay) in the same samples. Technical triplicates were measured for each sample. Shown are values normalized to untreated cells. Reported are the average and standard deviations from at least 3 independent biological replicates.

Western Blot: Cells were washed with ice cold PBS, then scraped into Laemmli lysis buffer (50mM Tris-HCl, pH 8.0; 2% SDS; 5mM EDTA, pH8.0) supplemented with protease inhibitor cocktail. Protein concentration was determined using BCA (Pierce), and equal amounts of protein were mixed with reducing Laemmli sample buffer and loaded onto SDS-PAGE gels. Gels were transferred to PVDF membranes, which were blocked in 5% BSA. Membranes were incubated with primary antibodies overnight at 4°C, and detected with either AlexaFluor or HRP-tagged secondary antibodies. Membranes were imaged using a BioRad ChemiDoc imager. The intensities of the bands were quantified normalized to actin, and plotted as mean ± SD of n ≥ 3 experiments.

Real Time Quantitative PCR: Total RNA was isolated via Trizol (Sigma) following the manufacturer's protocol. Reverse transcriptase PCR was performed using the High Capacity cDNA Reverse Transcription Kit from Applied Biosystems according to manufacturer's protocol. To quantify mRNA expression, SYBR Fast MasterMix (2x) Universal Dye (#KK4602) from Kapa Biosystems was used in an Eppendorf Realplex2 Mastercycler. Each primer set for each sample was run in triplicate with 1 ng of cDNA per well. The primer sets used are:

Gene	Forward Primer	Reverse Primer
GAPDH	GTCTACTGGCGTCTTCACCA	GTGGCAGTGATGGCATGGAC
SCD1	GTGATGTTCCAGAGGAGGTACT	CAGGAACTCAGAAGCCCAGAA
SCD2	GGTGATGTTCCAGAGGAGGTATT	AACTGGAAGACCCCGAACTC

Expression changes were calculated using the delta delta C_T method. The data was standardized using a previously published protocol [75] in which the average fold change was log transformed, mean centered, and confidence intervals determined to evaluate statistical significance.

Acknowledgements

523   All confocal microscopy was performed at the Center for Advanced Microscopy, Department of Integrative

524   Biology and Pharmacology McGovern Medical School at the University of Texas Health Science Center at

525   Houston. We acknowledge funding from the Cancer Prevention Research Institute of Texas (CPRIT; R1215),

526   NIH/National Institute of General Medical Sciences (grant nos. GM114282, GM124072, GM120351), and the

527   Volkswagen Foundation (grant no. 93091).

529 References

1   Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. Predicting transmembrane protein topology
with a hidden Markov model: application to complete genomes. *Journal of molecular biology* **305**, 567-
580, (2001).

2   Janmey, P. A. & Kinnunen, P. K. Biophysical properties of lipids and dynamic membranes. *Trends in cell*
biology **16**, 538-546, (2006).

3   Los, D. A. & Murata, N. Membrane fluidity and its roles in the perception of environmental signals.
Biochim Biophys Acta **1666**, 142-157, (2004).

4   Andersen, O. S. & Koeppe, R. E., 2nd. Bilayer thickness and membrane protein function: an energetic
perspective. *Annu Rev Biophys Biomol Struct* **36**, 107-130, (2007).

5   Covino, R. *et al.* A Eukaryotic Sensor for Membrane Lipid Saturation. *Molecular cell* **63**, 49-59, (2016).

6   daCosta, C. J., Dey, L., Therien, J. P. & Baenziger, J. E. A distinct mechanism for activating uncoupled
nicotinic acetylcholine receptors. *Nat Chem Biol* **9**, 701-707, (2013).

7   Marsh, D. Protein modulation of lipids, and vice-versa, in membranes. *Biochim Biophys Acta* **1778**, 1545-
1575, (2008).

8   Roux, A. *et al.* Membrane curvature controls dynamin polymerization. *Proc Natl Acad Sci U S A* **107**,
4141-4146, (2010).

9   Shi, Z. & Baumgart, T. Membrane tension and peripheral protein density mediate membrane shape
transitions. *Nature communications* **6**, 5974, (2015).

10   Rosholm, K. R. *et al.* Membrane curvature regulates ligand-specific membrane sorting of GPCRs in
living cells. *Nat Chem Biol* **13**, 724-729, (2017).

11   Sezgin, E., Levental, I., Mayor, S. & Eggeling, C. The mystery of membrane organization: composition,
regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* **18**, 361-374, (2017).

12   Lorent, J. H. *et al.* Structural determinants and functional consequences of protein affinity for membrane
rafts. *Nature communications* **8**, 1219, (2017).

13   Hazel, J. R. Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation?
Annu Rev Physiol **57**, 19-42, (1995).

14   Sinensky, M. Homeoviscous adaptation--a homeostatic process that regulates the viscosity of membrane
lipids in Escherichia coli. *Proc Natl Acad Sci U S A* **71**, 522-525, (1974).

15   Ernst, R., Ejsing, C. S. & Antonny, B. Homeoviscous adaptation and the regulation of membrane lipids.
Journal of molecular biology **428**, 4776-4791, (2016).

16   Guschina, I. A. & Harwood, J. L. Mechanisms of temperature adaptation in poikilotherms. *FEBS Lett*
580, 5477-5483, (2006).

17   Aguilar, P. S. *et al.* Molecular basis of thermosensing: a two-component signal transduction thermometer
in Bacillus subtilis. *EMBO J* **20**, 1681-1691, (2001).

18   Halbleib, K. *et al.* Activation of the Unfolded Protein Response by Lipid Bilayer Stress. *Molecular cell*
67, 673-684 e678, (2017).

19   Cao, J., Schwichtenberg, K. A., Hanson, N. Q. & Tsai, M. Y. Incorporation and clearance of omega-3
fatty acids in erythrocyte membranes and plasma phospholipids. *Clinical chemistry* **52**, 2265-2272,
(2006).

Oliveira, T. V. *et al.* Impact of high cholesterol intake on tissue cholesterol content and lipid transfers to high-density lipoprotein. *Nutrition* **27**, 713-718, (2011).

Gerl, M. J. *et al.* Quantitative analysis of the lipidomes of the influenza virus envelope and MDCK cell apical membrane. *J Cell Biol* **196**, 213-221, (2012).

Sampaio, J. L. *et al.* Membrane lipidome of an epithelial cell line. *Proc Natl Acad Sci U S A* **108**, 1903-1907, (2011).

Levental, K. R. *et al.* Polyunsaturated lipids regulate membrane domain stability by tuning membrane order. *Biophys J* **110(8)**, 1800-1810, (2016).

Han, X. in *Lipidomics*   427-442 (John Wiley & Sons, Inc, 2016).

Klemm, R. W. *et al.* Segregation of sphingolipids and sterols during formation of secretory vesicles at the trans-Golgi network. *J Cell Biol* **185**, 601-612, (2009).

Ejsing, C. S. *et al.* Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proc Natl Acad Sci U S A* **106**, 2136-2141, (2009).

Sinensky, M. Defective regulation of cholesterol biosynthesis and plasma membrane fluidity in a Chinese hamster ovary cell mutant. *Proc Natl Acad Sci U S A* **75**, 1247-1249, (1978).

Sinensky, M. Adaptive alteration in phospholipid composition of plasma membranes from a somatic cell mutant defective in the regulation of cholesterol biosynthesis. *J Cell Biol* **85**, 166-169, (1980).

Dymond, M. K. Mammalian phospholipid homeostasis: homeoviscous adaptation deconstructed by lipidomic data driven modelling. *Chem Phys Lipids* **191**, 136-146, (2015).

Dymond, M. K., Hague, C. V., Postle, A. D. & Attard, G. S. An in vivo ratio control mechanism for phospholipid homeostasis: evidence from lipidomic studies. *J R Soc Interface* **10**, 20120854, (2013).

Sodt, A. J., Venable, R. M., Lyman, E. & Pastor, R. W. Nonadditive Compositional Curvature Energetics of Lipid Bilayers. *Phys Rev Lett* **117**, 138104, (2016).

Almeida, P. F. Thermodynamics of lipid interactions in complex bilayers. *Biochim Biophys Acta* **1788**, 72-85, (2009).

Balogh, G. *et al.* Key role of lipids in heat stress management. *FEBS Lett* **587**, 1970-1980, (2013).

Levental, K. R. *et al.* omega-3 polyunsaturated fatty acids direct differentiation of the membrane phenotype in mesenchymal stem cells to potentiate osteogenesis. *Science advances* **3**, eaao1193, (2017).

Yaqoob, P. *et al.* Comparison of the effects of a range of dietary lipids upon serum and tissue lipid composition in the rat. *The international journal of biochemistry & cell biology* **27**, 297-310, (1995).

Pirro, M. *et al.* Plasma free fatty acid levels and the risk of ischemic heart disease in men: prospective results from the Quebec Cardiovascular Study. *Atherosclerosis* **160**, 377-384, (2002).

Metcalf, R. G. *et al.* Effects of fish-oil supplementation on myocardial fatty acids in humans. *The American journal of clinical nutrition* **85**, 1222-1228, (2007).

van Blitterswijk, W. J., van der Meer, B. W. & Hilkmann, H. Quantitative contributions of cholesterol and the individual classes of phospholipids and their degree of fatty acyl (un)saturation to membrane fluidity measured by fluorescence polarization. *Biochemistry* **26**, 1746-1756, (1987).

Lande, M. B., Donovan, J. M. & Zeidel, M. L. The relationship between membrane fluidity and permeabilities to water, solutes, ammonia, and protons. *J Gen Physiol* **106**, 67-84, (1995).

Sessler, A. M., Kaur, N., Palta, J. P. & Ntambi, J. M. Regulation of stearoyl-CoA desaturase 1 mRNA stability by polyunsaturated fatty acids in 3T3-L1 adipocytes. *J Biol Chem* **271**, 29854-29858, (1996).

Levental, I. & Veatch, S. L. The continuing mystery of lipid rafts. *J. Mol. Biol.* **428**, 4749-4764, (2016).

Lin, X. *et al.* Domain Stability in Biomimetic Membranes Driven by Lipid Polyunsaturation. *J Phys Chem B* **120**, 11930-11941, (2016).

Yeagle, P. L. Cholesterol and the cell membrane. *Biochim Biophys Acta* **822**, 267-287, (1985).

Crockett, E. Cholesterol Function in Plasma Membranes from Ectotherms: Membrane-Specific Roles in Adaptation to Temperature. *Integrative and Comparative Biology* **38**, 298-304, (1998).

Raghow, R. *et al.* SREBPs: the crossroads of physiological and pathological lipid homeostasis. *Trends Endocrinol Metab* **19**, 65-73, (2008).

Sakai, J. *et al.* Molecular identification of the sterol-regulated luminal protease that cleaves SREBPs and controls lipid composition of animal cells. *Molecular cell* **2**, 505-514, (1998).

47 Kim, H. M. *et al.* A two-photon fluorescent probe for lipid raft imaging: C-laurdan. *Chembiochem* **8**, 553-
559, (2007).
48 Owen, D. M. *et al.* Quantitative imaging of membrane lipid order in cells and organisms. *Nat Protoc* **7**,
24-35, (2011).
49 Kaiser, H. J. *et al.* Order of lipid phases in model and plasma membranes. *Proc Natl Acad Sci U S A* **106**,
16645-16650, (2009).
50 Gaus, K. *et al.* Condensation of the plasma membrane at the site of T lymphocyte activation. *J Cell Biol*
171, 121-131, (2005).
51 Parasassi, T. *et al.* Quantitation of lipid phases in phospholipid vesicles by the generalized polarization of
Laurdan fluorescence. *Biophys J* **60**, 179-189, (1991).
52 Sezgin, E. *et al.* Adaptive lipid packing and bioactivity in membrane domains. *PloS one* **10**, e0123930,
(2015).
53 Sezgin, E., Waithe, D., Bernardino de la Serna, J. & Eggeling, C. Spectral imaging to measure
heterogeneity in membrane lipid packing. *Chemphyschem* **16**, 1387-1394, (2015).
54 Feller, S. E., Gawrisch, K. & MacKerell, A. D., Jr. Polyunsaturated fatty acids in lipid bilayers: intrinsic
and environmental contributions to their unique physical properties. *J Am Chem Soc* **124**, 318-326,
(2002).
55 Tang, J. J. *et al.* Inhibition of SREBP by a small molecule, betulin, improves hyperlipidemia and insulin
resistance and reduces atherosclerotic plaques. *Cell Metab* **13**, 44-56, (2011).
56 Briolay, A., Jaafar, R., Nemoz, G. & Bessueille, L. Myogenic differentiation and lipid-raft composition of
L6 skeletal muscle cells are modulated by PUFAs. *Biochim Biophys Acta* **1828**, 602-613, (2013).
57 Seo, J. *et al.* Docosahexaenoic acid selectively inhibits plasma membrane targeting of lipidated proteins.
FASEB journal : official publication of the Federation of American Societies for Experimental Biology
20, 770-772, (2006).
58 Stark, K. D. *et al.* Fatty acid compositions of serum phospholipids of postmenopausal women: a
comparison between Greenland Inuit and Canadians before and after supplementation with fish oil.
Nutrition **18**, 627-630, (2002).
59 Kusakabe, T. *et al.* Fatty acid synthase is expressed mainly in adult hormone-sensitive cells or cells with
high lipid metabolism and in proliferating fetal cells. *J Histochem Cytochem* **48**, 613-622, (2000).
60 Weiss, L. *et al.* Fatty-acid biosynthesis in man, a pathway of minor importance. Purification, optimal
assay conditions, and organ distribution of fatty-acid synthase. *Biol Chem Hoppe Seyler* **367**, 905-912,
(1986).
61 Murphy, E. J. Stable isotope methods for the in vivo measurement of lipogenesis and triglyceride
metabolism. *J Anim Sci* **84 Suppl**, E94-104, (2006).
62 Nakagawa, Y., Sakumoto, N., Kaneko, Y. & Harashima, S. Mga2p is a putative sensor for low
temperature and oxygen to induce OLE1 transcription in Saccharomyces cerevisiae. *Biochem Biophys Res*
Commun **291**, 707-713, (2002).
63 Tiku, P. E. *et al.* Cold-induced expression of delta 9-desaturase in carp by transcriptional and
posttranslational mechanisms. *Science* **271**, 815-818, (1996).
64 Burns, M. *et al.* Miscibility Transition Temperature Scales with Growth Temperature in a Zebrafish Cell
Line. *Biophys J* **113**, 1212-1222, (2017).
65 Eberle, D. *et al.* SREBP transcription factors: master regulators of lipid homeostasis. *Biochimie* **86**, 839-
848, (2004).
66 Yahagi, N. *et al.* A crucial role of sterol regulatory element-binding protein-1 in the regulation of
lipogenic gene expression by polyunsaturated fatty acids. *J Biol Chem* **274**, 35840-35844, (1999).
67 Hannah, V. C. *et al.* Unsaturated fatty acids down-regulate srebp isoforms 1a and 1c by two mechanisms
in HEK-293 cells. *J Biol Chem* **276**, 4365-4372, (2001).
68 Xu, J. *et al.* Polyunsaturated fatty acids suppress hepatic sterol regulatory element-binding protein-1
expression by accelerating transcript decay. *J Biol Chem* **276**, 9800-9807, (2001).

69   Radanovic, T. *et al.* An Emerging Group of Membrane Property Sensors Controls the Physical State of
670   Organellar Membranes to Maintain Their Identity. *BioEssays : news and reviews in molecular, cellular*
671   *and developmental biology*, e1700250, (2018).
672   70   Fan, Y. Y., McMurray, D. N., Ly, L. H. & Chapkin, R. S. Dietary (n-3) polyunsaturated fatty acids
673   remodel mouse T-cell lipid rafts. *The Journal of nutrition* **133**, 1913-1920, (2003).
674   71   Fan, Y. Y. *et al.* Dietary docosahexaenoic acid suppresses T cell protein kinase C theta lipid raft
675   recruitment and IL-2 production. *J Immunol* **173**, 6151-6160, (2004).
676   72   Tulodziecka, K. *et al.* Remodeling of the postsynaptic plasma membrane during neural development. *Mol*
677   *Biol Cell* **27**, 3480-3489, (2016).
678   73   Sezgin, E. *et al.* Elucidating membrane structure and protein behavior using giant plasma membrane
679   vesicles. *Nat Protoc* **7**, 1042-1051, (2012).
680   74   Levental, K. R. & Levental, I. Isolation of giant plasma membrane vesicles for evaluation of plasma
681   membrane structure and protein partitioning. *Methods in molecular biology* **1232**, 65-77, (2015).
682   75   Willems, E., Leyns, L. & Vandesompele, J. Standardization of real-time PCR gene expression data from
683   independent biological replicates. *Analytical biochemistry* **379**, 127-129, (2008).
684