Gut Microbiota Diversity across Ethnicities in the United States

Authors:

Andrew W. Brooks[1,2], Sambhawa Priya[3,4,5], Ran Blekhman[3,4], Seth R. Bordenstein[1,2,6,7*]

Affiliations:

[1] Vanderbilt Genetics Institute, Vanderbilt University, Nashville, TN, USA.

[2] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3] Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis, MN, USA

[4] Department of Ecology, Evolution, and Behavior, University of Minnesota, Minneapolis, MN, USA

[5] Bioinformatics and Computational Biology Program, University of Minnesota, Minneapolis, MN, USA

[6] Department of Pathology, Microbiology, and Immunology, Vanderbilt University, Nashville, TN, USA.

[7] Vanderbilt Institute for Infection, Immunology and Inflammation, Vanderbilt University, Nashville, TN, USA.

[*] Corresponding author. E-mail: s.bordenstein@vanderbilt.edu (S.R.B)

Abbreviations:

AGP – American Gut Project

ANOSIM – Analysis of Similarity

AUC – Area Under the Curve

BMI – Body Mass Index

F_{ST} – Fixation Index

GWAS - Genome-Wide Association Studies

HMP – Human Microbiome Project

MAF - Minor Allele Frequency

OTU – Operational Taxonomic Unit

PERMANOVA - Permutational Multivariate Analysis of Variance

RF – Random Forest

ROC – Receiver Operating Characteristic

SMOTE – Synthetic Minority Over-sampling Technique

Abstract:

Composed of hundreds of microbial species, the composition of the human gut

microbiota can vary with chronic diseases underlying health disparities that

disproportionally affect ethnic minorities. However, the influence of ethnicity on the gut

microbiota remains largely unexplored and lacks reproducible generalizations across

studies. By distilling associations between ethnicity and differences in two United States

based 16S gut microbiota datasets including 1,673 individuals, we report 12 microbial

genera and families that reproducibly vary by ethnicity. Interestingly, a majority of these

microbial taxa, including the most heritable bacterial family, Christensenellaceae, overlap

with genetically-associated taxa and form co-occurring clusters linked by similar

fermentative and methanogenic metabolic processes. These results demonstrate recurrent

associations between specific taxa in the gut microbiota and ethnicity, providing hypotheses

for examining specific members of the gut microbiota as mediators of health disparities.

Introduction:

The human gut microbiota at fine resolution varies extensively between individuals (1-3), and this variability frequently associates with diet (4-7), age (6, 8, 9), sex (6, 9, 10), body mass index (BMI) (1, 6), and diseases presenting as health disparities (11-14). The overlapping risk factors and burden of many chronic diseases disproportionally affect ethnic minorities in the United States, yet the underlying biological mechanisms mediating these substantial disparities largely remain unexplained. Recent evidence is consistent with the hypothesis that ethnicity associates with variation in microbial abundance, specifically in the oral cavity, gut, and vagina (15-17). To varying degrees, ethnicity can capture many facets of biological variation including social, economic and cultural variation, as well as aspects of human genetic variation and biogeographical ancestry. Ethnicity also serves as a proxy to characterize health disparity incidence in the United States, and while factors such as genetic admixture create ambiguity of modern ethnic identity, self-declared ethnicity has proven a useful proxy for genetic and socioeconomic variation in population scale analyses, including in the Human Microbiome Project (18-20). Microbiota differences have been documented across populations that differ in ethnicity as well as in geography, lifestyle, and sociocultural structure; however, these global examinations cannot disconnect factors such as intercontinental divides and hunter-gatherer versus western lifestyles from ethnically structured differences (21-23). Despite the importance of understanding the interconnections between ethnicity, microbiota, and health disparities, there are no reproducible findings about the influence of ethnicity on differences in the gut microbiota and specific microbial taxa in diverse United States populations, even for healthy individuals (6).

Here, we comprehensively examine connections between self-declared ethnicity and

gut microbiota differences across more than a thousand individuals sampled by the

American Gut Project (AGP, N=1375) (24) and the Human Microbiome Project (HMP, N=298)

(6). Previous studies demonstrated that human genetic diversity in the HMP associates with

differences in microbiota composition(25), and genetic population structure within the HMP

generally delineates self-declared ethnicity (20). Ethnicity was not found to have a

significant association with microbiota composition in a Middle Eastern population,

however factors such as lifestyle and environment that influence microbiota variation across

participants was homogenous compared to the ethnic, sociocultural, economic, and dietary

diversity found within the United States (26). While ethnic diversity is generally

underrepresented in current microbiota studies, evidence supporting an ethnic influence on

microbiota composition among first generation immigrants has been recently demonstrated

in a Dutch population (27). The goal of this examination is to evaluate, for the first time, if

there are reproducible differences in gut microbiota across ethnicities within an overlapping

United States population, as ethnicity is one of the key defining factors for health disparity

incidence in the United States. Lifestyle, dietary, and genetic factors all vary to different

degrees across ethnic groups in the United States, and it will require more even sampling of

ethnic diversity and stricter phenotyping of study populations to disentangle which factors

underlie ethnic microbiota variation in the AGP and HMP.

Results:

Microbiota are subtly demarcated by ethnicity

We first evaluate gut microbiota distinguishability between AGP ethnicities (**Fig 1A**, family taxonomic level, Asians-Pacific Islanders (N=88), Caucasians (N=1237), Hispanics (N=37), and African Americans (N=13)), sexes (female (N=657), male (N=718)), age groups (years grouped by decade), and categorical BMI (underweight (N=70), normal (N=873), overweight (N=318), and obese (N=114)) (Demographic details in **S1A Table**). Age, sex, and BMI were selected as covariates because they are consistent across the AGP and HMP datasets. Additionally, 31 other categorical factors measuring diet, environment, and geography were compared for pairwise differences between two ethnicities using proportions tests, and very few (10 / 894) tests significantly varied (**S1 Table** additional sheets). Interindividual gut microbiota heterogeneity clearly dominates; however, Analyses of Similarity (ANOSIM) reveal subtle but significant degrees of total microbiota distinguishability for ethnicity, BMI, and sex, but not for age (**Fig 1B**, Ethnicity; **Fig 1C**, BMI; **Fig 1D**, Sex; **Fig 1E**, Age) (28). Recognizing that subtle microbiota distinguishability between ethnicities may be spurious, we independently replicate the ANOSIM results from HMP African Americans (N=10), Asians (N=34), Caucasians (N=211) and Hispanics (N=43) (**S2A Table**, R=0.065, p=0.044). We again observe no significant distinguishability for BMI, sex, and age in the HMP. Higher rarefaction depths increase microbiota distinguishability in the AGP across various beta diversity metrics and categorical factors (**S2B Table**), and significance increases when individuals from overrepresented ethnicities are subsampled from the average beta diversity distance matrix (**S2C Table**). Supporting the ANOSIM results, Permutational Multivariate Analysis of Variance (PERMANOVA) models with four different

beta diversity metrics showed that while all factors had subtle but significant associations

with microbiota variation when combined in a single model, effect sizes were highest for

ethnicity in 7 out of 8 comparisons across beta diversity metrics and rarefaction depths in

the AGP and HMP (**S2D Table**). We additionally test microbiota distinguishability by

measuring the correlation between beta diversity and ethnicity, BMI, sex, and age with an

adapted BioEnv test (**S2E Table**) (29). Similar degrees of microbiota structuring occur when

all factors are incorporated (Spearman Rho=0.055, p-values: Ethnicity=0.057, BMI<0.001,

Sex<0.001, Age=0.564). Firmicutes and Bacteroidetes dominated the relative phylum

abundance, with each representing between 35% and 54% of the total microbiota across

ethnicities (**S1 Fig**).

Fig 1. Gut microbiota composition and distinguishability by ethnicity, sex, age and BMI. (A) The average relative abundance of dominant microbial families for each ethnicity. (B-E) Principle coordinates analysis plots of microbiota Bray-Curtis beta diversity and ANOSIM distinguishability for: (B) Ethnicity, (C) Sex, (D) Age, (E) BMI. In B-E, each point represents the microbiota of a single sample, and colors reflect metadata for that sample.

Caucasian points are reduced in size to allow clearer visualization, and p-values are not

corrected across factors which have different underlying population distributions.

We next test for ethnicity signatures in the gut microbiota by analyzing alpha and beta

diversity, abundance and ubiquity distributions, distinguishability, and classification

accuracy (30). Shannon's Alpha Diversity Index (31), which weights both microbial

community richness (Observed OTUs) and evenness (Equitability), significantly varies

across ethnicities in the AGP dataset (Kruskal Wallis, $p=2.8e-8$) with the following ranks:

Hispanics > Caucasians > Asian-Pacific Islanders > African Americans (**Fig 2A**). In the HMP,

there is a significantly lower Shannon diversity for Asian-Pacific Islanders relative to

Caucasians and a trend of lower Shannon diversity for Asian-Pacific Islanders relative to

Hispanics; African Americans change position in diversity relative to other ethnicities,

potentially as a result of undersampling bias. Five alpha diversity metrics, two rarefaction

depths, and separate analyses of Observed OTUs and Equitability generally confirm the

results (**S3A Table**).

Fig 2. Ethnicity associates with diversity and composition of the gut microbiota. (A) Center lines of each boxplot depict the median by which ethnicities were ranked from low (left) to high (right); the lower and upper ends of each box represent the 25th and 75th quartiles respectively; whiskers denote the 1.5 interquartile range, and black dots represent individual samples. Lines in the middle of violin plots depict the mean, and p-values are Bonferroni corrected within each dataset. (B) Left extending violin plots represent intra-ethnic distances for each ethnicity, and right extending violin plots depict all inter-ethnic distances. Center lines depict the mean beta diversity. Significance bars above violin plots

depict Bonferroni corrected pairwise Mann-Whitney-U comparisons of the intra-intra- and intra-inter-ethnic distances. (C) Within each ethnicity, OTUs shared by at least 50% of samples. Colored lines represent a robust ordinary least squares regression within OTUs of each ethnicity, shaded regions represent the 95% confidence interval, R^2 denotes the regression correlation, the OTUs column indicates the number of OTUs with >50% ubiquity for that ethnicity, Mean A/U is the average abundance/ubiquity ratio, and the p_{adj} is the regression significance adjusted and Bonferroni corrected for the number of ethnicities.

If ethnicity impacts microbiota composition, pairwise beta diversity distances (ranging from 0/completely dissimilar to 1/identical) will be greater between ethnicities than within ethnicities. While average gut microbiota beta diversities across all individuals are high (**Fig 2B**, Bray-Curtis=0.808), beta diversities between individuals of the same ethnicity (intra-ethnic, Bray-Curtis=0.806) are subtly, but significantly, lower than those between ethnicities in both the AGP (inter-ethnic, Bray-Curtis=0.814) and HMP datasets (intra-ethnic, Bray-Curtis=0.870 versus inter-ethnic, Bray-Curtis=0.877). We confirm AGP results by subsampling individuals from overrepresented ethnicities across beta metrics and rarefaction depths (**S4A-4B Tables**). Finally, we repeat analyses across beta metrics and rarefaction depths using only the average distance of each individual to all individuals from the ethnicity to which they are compared (**S4C-4D Tables**).

Next, we explore inter-ethnic differences in the number of OTUs shared in at least 50% of individuals within an ethnicity, as the likelihood of detecting a biological signal is improved in more abundant organisms relative to noise that may predominate in lower abundance OTUs. Out of 5,591 OTUs in the total AGP dataset, 101 (1.8%) meet this ubiquity

cutoff in all ethnicities, and 293 (5.2%) OTUs meet the cutoff within at least one ethnicity.

Hispanics share the most ubiquitous OTUs and have the lowest average abundance/ubiquity

(A/U) ratio (**Fig 2C**), indicating stability whereby stability represents a more consistent

appearance of OTUs with lower abundance but higher ubiquity (32). This result potentially

explains their significantly lower intra-ethnic beta diversity distance and thus higher

microbial community overlap relative to the other ethnicities (**Fig 2B**). Comparisons in the

AGP between the higher sampled Hispanic, Caucasian, and Asian-Pacific Islander ethnicities

also reveal a trend wherein higher intra-ethnic community overlap (**Fig 2B**) parallels higher

numbers of ubiquitous OTUs (**Fig 2C**), higher Shannon Alpha diversity (**Fig 2A**), and higher

stability of ubiquitous OTUs as measured by the abundance/ubiquity (A/U) ratio (**Fig 2C**).

We next assess whether a single ethnicity disproportionately impacts total gut

microbiota distinguishability in the AGP by comparing ANOSIM results from the consensus

beta diversity distance matrix when each ethnicity is sequentially removed from the analysis

(**Fig 3A** and **S2E Table**). Distinguishability remains unchanged when the few African

Americans are removed, but is lost upon removal of Asian-Pacific Islanders or Caucasians,

likely reflecting their higher beta diversity distance from other ethnicities (**Fig 3A**). Notably,

removal of Hispanics increases distinguishability among the remaining ethnicities, which

may be due to higher degree of beta diversity overlap observed between Hispanics and other

ethnicities (**S4B Table**). Results conform across rarefaction depths and beta diversity

metrics (**S2F Table**), and pairwise combinations show strong distinguishability between

African Americans and Hispanics (ANOSIM, $R=0.234$, $p=0.005$), and Asian-Pacific Islanders

and Caucasians (ANOSIM, $R=0.157$, $p<0.001$).

Finally, to complement evaluation with ecological alpha and beta diversity we

implement a random forest (RF) supervised learning algorithm to classify gut microbiota

from genus level community profiles into their respective ethnicity. We build four one-

versus-all binary classifiers to classify samples from each ethnicity compared to the rest, and

use two different sampling approaches to train the models, Synthetic Minority Over-

sampling Technique (SMOTE) (33) and down-sampling, for overcoming uneven

representation of ethnicities in both the datasets (see Methods). Given that the area under

the receiver operating characteristic (ROC) curve (or AUC) of a random guessing classifier is

0.5, the models classify each ethnicity fairly well (**Fig 3B**) with average AUCs across sampling

techniques and datasets of 0.78 for Asian-Pacific Islanders, 0.76 for African Americans, 0.69

for Hispanics, and 0.70 for Caucasians. Ethnicity distinguishing RF taxa and out-of-bag error

percentages appear in (**S2 Fig**).

Fig 3. Microbiota distinguishability and classification ability across ethnicities. (A) ANOSIM distinguishability between all combinations of ethnicities. Symbols depict specific ethnicities included in the ANOSIM tests, and boxes denote the R-value as a heatmap, where white indicates increasing and black indicates decreasing distinguishability relative to the R-value with all ethnicities. (B) Average ROC curves (for 10-fold cross-validation) and prediction performance metrics for one-versus-all RF classifiers for each ethnicity, using SMOTE (33) and down subsampling approaches for training.

Recurrent taxon associations with ethnicity

Subtle to moderate ethnicity-associated differences in microbial communities may in

part be driven by differential abundance of certain microbial taxa. 16.2% (130/802) of the

AGP taxa and 20.6% (45/218) of HMP taxa across all classification levels (i.e. phylum to

genus, **S5 Table**) significantly vary in abundance across ethnicities (Kruskal-Wallis,

p_{FDR}<0.05). Between datasets, 19.2% (25/130) of the AGP and 55.6% (25/45) of the HMP

varying taxa replicate in the other dataset, representing a significantly greater degree of

overlap than would be expected by chance (ethnic permutation analysis of overlap, p<0.001

each taxonomic level and all taxonomic levels combined). The highest replication of taxa

varying by abundance occurs with 22.0% of families (9 significant in both datasets / 41

significantly varying families in either dataset), followed by genus with 13.4% (9 significant

in both datasets / 67 significantly varying genera in either dataset).

Among 18 reproducible taxa, we categorize 12 as taxonomically distinct (**Fig 4**) and

exclude 6 where nearly identical abundance profiles between family/genus taxonomy

overlap. Comparing relative abundance differences between pairs of ethnicities for these 12

taxa in the AGP reveals 30 significant differences, of which 20 replicate in the HMP (p<0.05,

Mann-Whitney-U). Intriguingly, all reproducible pairwise differences are a result of

decreases in Asian-Pacific Islanders (**Fig 4**). We also test taxon abundance and

presence/absence associations with ethnicity separately in the AGP using linear and logistic

regression models respectively, and we repeat the analysis while incorporating categorical

sex and continuous age and BMI as covariates (**S6 Table**). Clustering microbial families

based on their abundance correlation reveals two co-occurrence clusters: (i) a distinct

cluster of six Firmicutes and Tenericutes families in the HMP and (ii) an overlapping but

more diverse cluster of 20 families in the AGP (**S3 Fig**). Nine of the 12 taxa found to

recurrently vary in abundance across ethnicities are represented in these clusters (**Fig 4**),

with four appearing in both clusters and the other five appearing either in or closely

correlated with members of both clusters (**S3 Fig**). Furthermore, 90% (18/20) of families in

the AGP cluster and 66% (4/6) of taxa in the HMP cluster significantly vary in abundance

across ethnicities. We also found overlap for AGP and HMP datasets between taxa

significantly varying in abundance across ethnicities (with FDR <0.05) and taxa in RF models

with percentage importance greater than 50% for an ethnicity (**S2B Fig**). Taken together,

these results establish general overlap of the most significant ethnicity-associated taxa

between the these methods, reproducibility of microbial abundances that vary between

ethnicities across datasets, and patterns of co-occurrence among these taxa which could

suggest they are functionally linked.

Fig 4. Ethnicity-associated taxa match between the HMP and AGP. Barplots depict the log10 transformed relative abundance for individuals possessing the respective taxon within each ethnicity, ubiquity appears above (AGP) or below (HMP) bars, and the 25th and 75th percentiles are shown with extending whiskers. Mann-Whitney-U tests evaluate differences in abundance and ubiquity for all individuals between pairs of ethnicities; for example, the direction of change in Victivallaceae is driven by ubiquity while abundance is higher for those possessing the taxon. Significance values are Bonferroni corrected for the six tests within each taxon and dataset, and bold p-values indicate that significance (p<0.05) and direction of change replicate in the AGP and HMP.

Most heritable taxon of bacteria varies by ethnicity

Identified as the most heritable taxon in the human gut (34, 35), the family Christensenellaceae exhibits the second strongest significant difference in abundance across ethnicities in both AGP and HMP datasets (**S5 Table**, Family: AGP, Kruskal-Wallis, p_{FDR}=1.55e-9; HMP, Kruskal-Wallis, p_{FDR}=0.0019). Additionally, Christensenellaceae is variable by sex and BMI (AGP: Sex, Kruskal-Wallis, p_{FDR}=1.22e-12; BMI, Kruskal-Wallis, p_{FDR}=0.0020), and represents some of the strongest pairwise correlations with other taxa in both co-occurrence clusters (**S3 Fig**). There is at least an eight-fold and two-fold reduction in average Christensenellaceae abundance in Asian-Pacific Islanders relative to the other ethnicities in the AGP and HMP respectively (**S5 Table**), and significance of all pairwise comparisons in both datasets show reduced abundance in Asian-Pacific Islanders (**Fig 4**). Christensenellaceae also occur among the top 10 most influential taxa for distinguishing Asian-Pacific Islanders from other ethnicities using RF models for both AGP and HMP

datasets (**S2A Fig**). Abundance in individuals possessing Christensenellaceae and

presence/absence across all individuals significantly associate with ethnicity (**S6 Table**,

Abundance, Linear Regression, $p_{Bonferroni}$=0.006; Presence/Absence, Logistic Regression,

$p_{Bonferroni}$=8.802e-6), but there was only a slight correlation between the taxon's relative

abundance and BMI (**S4 Fig**). Confirming previous associations with lower BMI(36), we

observe that AGP individuals with Christensenellaceae also have a lower BMI (Mean BMI,

23.7±4.3) than individuals without it (Mean BMI, 25.0±5.9; Mann-Whitney-U, p<0.001). This

pattern is separately reflected in African Americans, Asian-Pacific Islanders, and Caucasians

but not Hispanics (**Fig 5**), suggesting that each ethnicity may have different equilibria

between the taxon's abundance and body weight.

Fig 5. Christensenellaceae variably associate with BMI across ethnicities. Boxplots of

BMI for individuals without (unfilled boxplots) and with (filled boxplots)

Christensenellaceae. Significance was determined using one-tailed Mann-Whitney-U tests

for lower continuous BMI values. Black lines indicate the mean relative abundance; the lower

and upper end of each box represent the 25[th] and 75[th] quartiles respectively; and whiskers

denote the 1.5 interquartile range.

Genetic- and ethnicity-associated taxa overlap

Many factors associate with human ethnicity, including a small subset of population

specific genetic variants (estimated ~0.5% genome wide) that vary by biogeographical

ancestry (37, 38); self-declared ethnicity in the HMP is delineated by population genetic

structure (20). Here we investigate whether ethnicity-associated taxa overlap with (i) taxa

that have a significant population genetic heritability in humans (34, 35, 39, 40) and (ii) taxa

linked with human genetic variants in two large Genome-Wide Association Studies (GWAS)-

microbiota analyses (35, 40). All recurrent ethnicity-associated taxa except one were

heritable in at least one study, with seven replicating in three or more studies (**Table 1**).

Likewise, abundance differences in seven recurrent ethnicity-associated taxa demonstrate

significant GWAS associations with at least one variant in the human genome. Therefore, we

assess whether any genetic variants associated with differences in microbial abundance

exhibit significant rates of differentiation (F_{ST}) between 1,000 genomes superpopulations

(38). Out of 49 variants associated with ethnically varying taxa, 21 have higher F_{ST} values

between at least one pair of populations than that of 95% of other variants on the same

chromosome and across the genome; the F_{ST} values of five variants associated with

Clostridiaceae abundance rank above the top 99% (**S7 Table**). Since taxa that vary across

ethnicities exhibit lower abundance in Asian-Pacific Islanders, it is notable that the F_{ST} values

of 18 and 11 variant comparisons for East Asian and South Asian populations, respectively,

are above that of the 95% rate of differentiation threshold from African, American, or

European populations. Cautiously, the microbiota and 1,000 genomes datasets are not

drawn from the same individuals, and disentangling the role of genetic from social and

environmental factors will still require more controlled studies.

Recurrent Ethnicity-Associated Taxa	Heritability	Genetic Associations
Family: Peptococcaceae	0.1213[A], 0.2154[C], 0.26[E]	rs143179968[E]
Family: Dehalobacteriaceae	0.6878[B], 0.3087[C]	
Family: Christensenellaceae	0.3819[A], 0.6170[B], 0.4230[C],	
Order: Clostridiales, Family: Unclassified	0.2914[A], 0.4020[B], 0.1330[C]	*40 Genetic Variants[C]
Genus: *Veillonella*	0.1370[A], 0.2168[D]	rs347941[C]
Order: RF39, Family: Unclassified	0.2341[A], 0.6618[B], 0.3074[C]	rs4883972[C]
Family: Verrucomicrobiaceae	0.1257[A], 0.5973[B], 0.1394[C]	
Family: Victivallaceae		
Family Odoribacteraceae	0.1389[A], 0.1917[D], 0.34[E]	chr7:96414393[E], rs115795847[E]
Genus: *Odoribacter*	0.1916[D]	
Family: Rikenellaceae	0.1299[D], 0.29[E]	rs17098734[C], rs3909540[C], rs147600757[E]
Family: Coriobacteraceae, Genus: Unclassified	0.1364[A], 0.2822[B], 0.1609[C]	rs9357092[E]

Table 1. Most recurrent ethnicity-associated taxa are previously reported heritable

and genetically-associated taxa. The table shows population genetic heritability estimates

and associated genetic variants for the 12 recurrent ethnically varying taxa. The minimum

heritability cutoff was chosen as >0.1, and only exactly overlapping taxonomies were

considered. Studies examined: [A]UKTwins (2014, 'A' measure of additive heritability in ACE

model) (34), [B]Yatsunenko (2014, 'A' measure of additive heritability in ACE model) (34),

[C]UKTwins (2016, 'A' measure of additive heritability in ACE model) (35), [D]Lim (2016, H2r

measure of polygenic heritability in SOLAR (41)) (39), [E]Turpin (2016, H2r measure of

polygenic heritability in SOLAR (41)). *indicates excessive variants were excluded from

table.

Discussion:

Many common diseases associate with microbiota composition and ethnicity, raising

the central hypothesis that microbiota differences between ethnicities can occasionally

serve as a mediator of health disparities. American's self-declared ethnicity can capture

socioeconomic, cultural, geographic, dietary and genetic diversity, and a similarly complex

array of interindividual and environmental factors influence total microbiota composition.

This complexity may result in challenges when attempting to recover consistent trends in

total gut microbiota differences between ethnicities. The challenges in turn emphasize the

importance of reproducibility, both through confirmation across analytical methods and

replication across study populations (15-17, 20, 27, 42). In order to robustly substantiate the

ethnicity-microbiota hypothesis, we evaluated recurrent associations between self-declared

ethnicity and variation in both total gut microbiota and specific taxa in healthy individuals.

Results provide hypotheses for examining specific members of the gut microbiota as

mediators of health disparities.

Our findings from two American datasets demonstrate that: (i) ethnicity consistently

captures gut microbiota with a slightly stronger effect size than other variables such as BMI,

age, and sex, (ii) ethnicity is moderately predictable from total gut microbiota differences,

and (iii) 12 taxa recurrently vary in abundance between the ethnicities, of which the majority

have been previously shown to associate with human genetic variation. Whether shaped

through socioeconomic, dietary, healthcare, genetic, or other ethnicity-related factors,

reproducibly varying taxa represent sources for novel hypotheses addressing health

disparities. For instance, the family Odoribacteraceae and genus *Odoribacter* are primary

butyrate producers in the gut, and they have been negatively associated to severe forms of

Crohn's disease and Ulcerative Colitis in association with reduced butyrate metabolism (43-

45). Asian-Pacific Islanders possess significantly less Odoribacteraceae and *Odoribacter* than

Hispanics and Caucasians in both datasets, and severity of Ulcerative Colitis upon hospital

admission has been shown to be significantly higher in Asian Americans (46). Considering

broader physiological roles, several ethnicity-associated taxa are primary gut anaerobic

fermenters and methanogens (47, 48), and associate with lower BMI and blood triglyceride

levels (36, 49). Indeed, Christensenellaceae, Odoribacteraceae, *Odoribacter*, and the class

Mollicutes containing RF39 negatively associate with metabolic syndrome and demonstrate

significant population genetic heritability in twins (39). Implications for health outcomes

warrant further investigation, but could be reflected by positive correlations of

Odoribacteraceae, *Odoribacter*, Coriobacteriaceae, Christensenellaceae, and the dominant

Verrucomicrobiaceae lineage *Akkermansia* with old age (50, 51). *Akkermansia* associations

with health and ethnicity in western populations may reflect recently arising dietary and

lifestyle effects on community composition, as this mucus consuming taxon is rarely

observed in more traditional cultures globally (23). Moreover, these findings raise the

importance of controlling for ethnicity in studies linking microbiota differences to disease

because associations between specific microbes and a disease could be confounded by

ethnicity of the study participants.

Based on correlations in individual taxon's abundance, a similar pattern of co-

occurrence previously identified as the 'Christensenellaceae Consortium' includes 11 of the

12 recurrent ethnically varying taxa (34), and members of this consortium associate with

genetic variation in the human formate oxidation gene *ALDH1L1*, which is a genetic risk

factor for stroke (35, 52, 53). Formate metabolism is a key step in the pathway reducing

carbon dioxide to methane (54, 55), and increased methane associates with increased

Rikenellaceae, Christensenellaceae, Odoribacteraceae and *Odoribacter* (56). Products of

methanogenic fermentation pathways include short chain fatty acids such as butyrate, which

through reduction of pro-inflammatory cytokines is linked to cancer cell apoptosis and

reduced risk of colorectal cancer (57, 58). Asian Americans are the only ethnic group where

cancer surpasses heart disease as the leading cause of death, and over 70% of Asian

Americans were born overseas, which can affect assimilation into western lifestyles, leading

to reduced access to healthcare and screening, and proper medical education (57, 59-61).

Preliminary results from other groups suggest that the gut microbiome of Southeast Asian

immigrants changes after migration to the United States (Dan Knights, personal

communication). Indeed, as countries in Asia shift toward a more western lifestyle, the

incidence of cancers, particularly gastrointestinal and colorectal cancers, are increasing

rapidly, possibly indicating incompatibilities between traditionally harbored microbiota and

western lifestyles (62-65). Asian Americans have higher rates of type 2 diabetes and

pathogenic infections than Caucasians (66), and two metagenomic functions enriched in

control versus type 2 diabetes cases appear to be largely conferred by cluster-associated

butyrate-producing and motility-inducing Verrucomicrobiaceae and Clostridia taxa reduced

in abundance among AGP and HMP Asian-Pacific Islanders (11). Both induction of cell

motility and butyrate promotion of mucin integrity can protect against pathogenic

colonization and associate with microbial community changes (11, 58, 67). Levels of cell

motility and butyrate are key factors suspected to underlie a range of health disparities

including inflammatory bowel disease, arthritis, and type 2 diabetes (11, 68-70). Patterns of

ethnically varying taxa across ethnicities could result from many factors including varying

diets, environmental exposures, sociocultural influences, human genetic variation and

others. However, regardless of the mechanisms dictating assembly, these results suggest

there is a reproducible, co-occurring group of taxa linked by similar metabolic processes

known to promote homeostasis.

The utility of this work is establishing a framework for studying ethnicity-associated

taxa and hypotheses of how changes in abundance or presence of these taxa may or may not

shape health disparities, many of which also have genetic components. Differing in allele

frequency across three population comparisons and associated with the abundance of

Clostridiales, the genetic variant rs7587067 has a significantly higher frequency in African

(Minor Allele Frequency (MAF)=0.802) versus East Asian (MAF=0.190, F_{ST}=0.54,

Chromosome=98.7%, Genome-Wide=98.9%), admixed American (MAF=0.278, F_{ST}=0.44,

Chromosome=99.0%, Genome-Wide=99.1%), and European populations (MAF=0.267,

F_{ST}=0.45, Chromosome=98.7.3%, Genome-Wide=98.7%). This intronic variant for the gene

HECW2 is a known eQTL (GTeX, eQTL Effect Size=-0.18, p=7.4e-5) (71, 72), and *HECW2*

encodes a ubiquitin ligase linked to enteric gastrointestinal nervous system function through

maintenance of endothelial lining of blood vessels (73, 74). Knockout of *HECW2* in mice

reduced enteric neuron networks and gut motility, and patients with Hirschsprung's disease

have diminished localization of *HECW2* to regions affected by loss of neurons and colon

blockage when compared to other regions of their own colon and healthy individuals (75).

Hirschsprung's disease presenting as full colon blockage is rare and has not undergone

targeted examination as a health disparity, however a possible hypothesis is that lower

penetrance of the disease in individuals with the risk allele at rs7587067 could lead to

subtler effects on gut motility resulting in Clostridiales abundance differences.

Despite the intrigue of connecting the human genome, microbiota and disease

phenotypes, evaluating such hypotheses will require more holistic approaches including

incorporating metagenomics and metabolomics to identify whether enzymes or metabolic

functions reproducibly vary across ethnicities, as well as direct functional studies in model

systems to understand if correlation is truly driven by causation. Further limitations should

also be considered, including recruitment biases for the AGP versus HMP, variation in sample

processing and OTU clustering, and uneven sampling which could only be addressed with

down sampling of over-represented ethnicities. Still, despite these confounders care was

taken to demonstrate the reproducibility of results across statistical methods, ecological

metrics, rarefaction depths, and study populations. Summarily, this work suggests that

abundance differences of specific taxa, rather than whole communities, may represent the

most reliable ethnic signatures in the gut microbiota. A reproducible co-occurring subset of

these taxa link to a variety of overlapping metabolic processes and health disparities, and

contain the most reproducibly heritable taxon, Christensenellaceae. Moreover, a majority of

the microbial taxa associated with ethnicity are also heritable and genetically-associated

taxa, suggesting there is a possible connection between ethnicity and genetic patterns of

biogeographical ancestry that may play a role in shaping these taxa. Our results emphasize

the importance of sampling ethnically diverse populations of healthy individuals in order to

discover and replicate ethnicity signatures in the human gut microbiota, and they highlight

a need to account for ethnic variation as a potential confounding factor in studies linking

microbiota differences to disease. Further reinforcement of these results may lead to

generalizations about microbiota assembly and even consideration of specific taxa as

potential mediators or treatments of health disparities.

Materials and Methods:

Data Acquisition

AGP data was obtained from the project FTP repository located at

ftp://ftp.microbio.me/AmericanGut/. AGP data generation and processing prior to analysis

can be found at: *https://github.com/biocore/American-Gut/tree/master/ipynb/primary-*

processing. All analyses utilized the rounds-1-25 dataset which was released on March 4,

2016. Throughout all analyses, QIIME v1.9.0 was used in an Anaconda environment

[*https://continuum.io*] for all script calls, custom scripts and notebooks were run in the

QIIME 2 Anaconda environment with python version 3.5.2, and plots were post-processed

using Inkscape [*https://inkscape.org/en/*] (76). Ethnicity used in this study was self-declared

by AGP study participants as one of four groups: African American, Asian or Pacific Islander

(Asian-Pacific Islander), Caucasian, or Hispanic. Sex was self-declared as either male, female,

or other. Age was self-declared as a continuous integer of years old, and age categories

defined by the AGP by decade (i.e. 20's, 30's…) were used in this study. BMI was self-declared

as an integer, and BMI categories defined by AGP of underweight, healthy, overweight, and

obese were utilized. A total of 31 categorical metadata factors were assessed for structuring

across ethnicities with a two proportion Z test between pairs of ethnicities using a custom

python script (**S1 Table** additional sheets). The p-values were Bonferroni corrected within

each metadata factor for the number of pairwise ethnic comparisons. 97% Operational

Taxonomic Units (OTUs) generated for each dataset are utilized throughout to maintain

consistency with other published literature, however microbial taxonomy of the HMP is

reassigned using the Greengenes reference database (77). Communities characterized with

16S rDNA sequencing of variable region four followed an identical processing pipeline for all

samples, which was developed and optimized for the Earth Microbiome Project (78). HMP 16S rDNA data processed using QIIME for variable regions 3-5 was obtained from *http://hmpdacc.org/HMQCP/*. Demographic info for individual HMP participants was obtained through dbGaP restricted access to study phs000228.v2.p1, with dbGaP approval granted to SRB and non-human subjects determination IRB161231 granted by Vanderbilt University. Ethnicity and sex were assigned to subjects based on self-declared values, with individuals selecting multiple ethnicities being removed unless they primarily responded as Hispanic, while categorical age and BMI were established from continuous values using the same criteria for assignment as in AGP. The HMP Amerindian population was removed due to severe under-representation. This filtered HMP table was used for community level analyses (ANOSIM, Alpha Diversity, beta intra-inter), however to allow comparison with the AGP dataset, community subset analyses (co-occurrence, abundance correlation, etc...) were performed with taxonomic assignments in QIIME using the UCLUST method with the GreenGenes_13_5 reference.

Quality Control

AGP quality control was performed in Stata v12 (StataCorp, 2011) using available metadata to remove samples (Raw N=9,475): with BMI more than 60 (-988 [8,487]) or less than 10 (-68 [8,419]), missing age (-661 [7,758]), with age greater than 55 years old (-2,777 [4,981]) or less than 18 years old (-582 [4,399]), and blank samples or those not appearing in the mapping file (-482 [3,917]), with unknown ethnicity or declared as other (-131 [3786]), not declared as a fecal origin (-2,002 [1784]), with unknown sex or declared as other (-98 [1686]), or located outside of the United States (-209 [1477]). No HMP individuals were

missing key metadata or had other reasons for exclusion (-0[298]). Final community quality

control for both AGP and HMP was performed by filtering OTUs with less than 10 sequences

and removing samples with less than 1,000 sequences (AGP, -102 [1375]; HMP, -0 [298]). All

analyses used 97% OTUs generated by the AGP or HMP, and unless otherwise noted, results

represent Bray-Curtis beta diversity and Shannon alpha diversity at a rarefaction depth of

1,000 counts per sample.

ANOSIM, PERMANOVA, and BioEnv Distinguishability

The ANOSIM test was performed with 9,999 repetitions on each rarefied table within

a respective rarefaction depth and beta diversity metric (**Fig 1 & S2A-B Table**), with R-

values and p-values averaged across the rarefactions. Consensus beta diversity matrices

were calculated as the average distances across the 100 rarefied matrices for each beta

diversity metric and depth. Consensus distance matrices were randomly subsampled ten

times for subset number of individuals from each ethnic group with more than that subset

number prior to ANOSIM analysis with 9,999 repetitions, and the results were averaged

evaluating the effects of more even representations for each ethnicity (**S2C Table**).

Consensus distance matrices had each ethnicity and pair of ethnicities removed prior to

ANOSIM analysis with 9,999 repetitions, evaluating the distinguishability conferred by

inclusion of each ethnicity (**Fig 3A**, **S2F Table**). Significance was not corrected for the

number of tests to allow comparisons between results of different analyses, metrics, and

depths. PERMANOVA analyses were run using the R language implementation in the Vegan

package (79), with data handled in a custom R script using the Phyloseq package (80).

Categorical variables were used to evaluate the PERMANOVA equation (Beta-Diversity

Distance Matrix ~ Ethnicity + Age + Sex + BMI) using 999 permutations to evaluate

significance, and the R and p values were averaged across 10 rarefactions (**S2D Table**). The

BioEnv test, or BEST test, was adapted to allow evaluation of the correlation and significance

between beta diversity distance matrices and age, sex, BMI, and ethnicity simultaneously

(**S2E Table**) (29). At each rarefaction depth and beta diversity metric the consensus distance

matrix was evaluated for its correlation with the centered and scaled Euclidian distance

matrix of individuals continuous age and BMI, and categorical ethnicity and sex encoded

using patsy (same methodology as original test)[*https://patsy.readthedocs.io/en/latest/#*].

The test was adapted to calculate significance for a variable of interest by comparing how

often the degree of correlation with all metadata variables (age, sex, BMI, ethnicity) was

higher than the correlation when the variable of interest was randomly shuffled between

samples 1,000 times.

Alpha Diversity

527   Alpha diversity metrics (Shannon, Simpson, Equitability, Chao1, Observed OTUs)

were computed for each rarefied table (QIIME: alpha_diversity.py), and results were collated

and averaged for each sample across the tables (QIIME: collate_alpha.py). Pairwise

nonparametric t-tests using Monte Carlo permutations evaluated alpha diversity differences

between the ethnicities with Bonferroni correction for the number of comparisons (**Fig 2A**,

S3 Table, QIIME: compare_alpha_diversity.py). A Kruskal-Wallis test implemented in python

was used to detect significant differences across all ethnicities.

Beta Diversity

Each consensus beta diversity distance matrix had distances organized based on

whether they represented individuals of the same ethnic group, or were between individuals

of different ethnic groups. All values indicate that all pairwise distances between all

individuals were used (**Fig 2B**, **S4A-B Table**), mean values indicate that for each individual

their average distance to all individuals in the comparison group was used as a single point

to assess pseudo-inflation (**S4C-D Table**). A Kruskal-Wallis test was used to calculate

significant differences in intra-ethnic distances across all ethnicities. Pairwise Mann-

Whitney-U tests were calculated between each pair of intra-ethnic distance comparisons,

along with intra-versus-inter ethnic distance comparisons. Significance was Bonferroni

corrected within the number of intra-intra-ethnic and intra-inter-ethnic distance groups

compared, with violin plots of intra- and inter-ethnic beta diversity distances generated for

each comparison.

Random Forest

RF models were implemented using taxa summarized at genus level, which

performed better compared to RF models using OTUs as features, both in terms of

classification accuracy and computational time. We first rarefied OTU tables at sequence

depth of 10,000 (using R v3.3.3 package *vegan's* rrarefy() function) and then summarized

rarefied OTUs at genus-level (or lower characterized level if genus was uncharacterized for

an OTU). We filtered for rare taxa by removing taxa present in fewer than half of the number

of samples in rarest ethnicity (i.e. fewer than 10/2 = 5 samples in HMP and 13/2 = 6 (rounded

down) in AGP), retaining 85 distinct taxa in HMP dataset and 322 distinct taxa in AGP dataset

at genus level. The resulting taxa were normalized to relative abundance and arcsin-sqrt

transformed before being used as features for the RF models. We initially built multi-class

RF model, but since the RF model is highly sensitive to the uneven representation of classes,

all samples were identified as the majority class, i.e. Caucasian. In order to even out the class

imbalance, we considered some sampling approaches, but most existing techniques for

improving classification performance on imbalanced datasets are designed for binary class

imbalanced datasets, and are not effective on datasets with multiple underrepresented

classes. Hence, we adopted the binary classification approach and built four one-versus-all

binary RF classifiers to classify samples from each ethnicity compared to the rest. 10-fold

cross-validation (using R package *caret* (81)) was performed using ROC as the metric for

selecting optimal model. The performance metrics and ROC curves were averaged across the

10 folds (**Fig 3B**). Without any sampling during training the classifiers, most samples were

identified as the majority class, i.e. the Caucasian, by all four one-versus-all RF classifiers. In

order to overcome this imbalance in class representation, we applied two sampling

techniques inside cross-validation: i) down-sampling, and ii) Synthetic Minority Over-

sampling Technique (or SMOTE) (33). In the down-sampling approach, the majority class is

down-sampled by random removal of instances from the majority class. In the SMOTE

approach, the majority class is down-sampled and synthetic samples from the minority class

are generated based on k-nearest neighbors technique (33). Note, the sampling was

performed inside cross-validation on training set, while the test was performed on

unbalanced held-out test set in each fold. In comparison to a no-sampling approach, which

classified most samples as the majority class, i.e., Caucasians, our sampling-based approach

leads to improved sensitivity for classification of minority classes on unbalanced test sets.

Nevertheless, the most accurate prediction remains for the inclusion in the majority class.

The ROC curves and performance metrics table in **Fig 3B** show the sensitivity-specificity

tradeoff and classification performance for one-versus-all classifier for each ethnicity for

both the sampling techniques applied on both the datasets. For both the datasets, down-

sampling shows higher sensitivity and lower specificity and precision for minority classes

(i.e. African Americans, Asian-Pacific Islanders and Hispanics) compared to SMOTE.

However, for the majority class (i.e. Caucasian), down-sampling lowers the sensitivity and

increases the specificity and precision compared to SMOTE. The sensitivity-specificity

tradeoff, denoted by the area under the ROC curve (or AUC) is reduced for Hispanics in both

the datasets. The most important taxa with >50% importance for predicting an ethnicity

using RF model with SMOTE sampling approach are shown in **S2A Fig**. Among the 10 most

important taxa for each ethnicity, there are 9 taxa which overlap between the AGP and HMP

datasets (highlighted by the blue rectangular box); however, which ethnicity they best

distinguish varies between the two datasets. Within each dataset we highlighted taxa which

are distinguishing in RF models and have distinguishing differential abundance in **S2B Fig**,

reporting both the FDR corrected significance for Kruskal-Wallis tests of differential

abundance, and the percent importance for the most distinguished ethnicity of each in RF

models. We also report out-of-bag errors for the final RF classifier that was built using the

optimal model parameters obtained from cross-validation approach corresponding to each

ethnicity and sampling procedure for both AGP and HMP datasets in **S2C Fig**.

Taxon Associations

Taxon differential abundance across categorical metadata groups was performed in

QIIME (QIIME: group_significance.py, **S5 Table**) to examine whether observation counts (i.e.

OTUs and microbial taxon) are significantly different between groups within a metadata category (i.e. ethnicity, sex, BMI, age). The OTU table prior to final community quality control was collapsed at each taxonomic level (i.e. Phylum – Genus; QIIME: collapse_taxonomy.py), with counts representing the relative abundance of each microbial taxon. Differences in the mean abundance of taxa between ethnicities were calculated using Kruskal-Wallis nonparametric statistical tests. P-values are provided alongside false discovery rate and Bonferroni corrected P-values, and taxon were ranked from most to least significant. Results were collated into excel tables by taxonomic level and metadata category being examined, with significant (false discovery rate and Bonferroni P-value < 0.05) highlighted in orange, and taxa that were false discovery rate significant in both datasets were colored red. The Fisher's exact test for the overlap of number of significant taxa between datasets was run at the online portal (http://vassarstats.net/tab2x2.html), with the expected overlap calculated as 5% of the number of significant taxa at all levels within the respective dataset, and the observed 25 taxa that overlapped in our analysis. The permutation analysis was performed by comparing the number of significant taxa (**S5 Table**, p_{FDR}<0.05) overlapping between the AGP and HMP to the number overlapping when the Kruskal-Wallis test was performed 1,000 times with ethnicity randomly permuted. In 1/1000 runs there was one significant taxon overlapping at the family level, and one in 3/1000 permutations at the genus level, with no significant taxa overlapping in any repetitions at higher taxonomic levels. The 12 families and genera that were significantly different were evaluated to not be taxonomically distinct if their abundances across ethnicities at each level represented at least 82-100% (nearly all >95%) of the overlapping taxonomic level, and the genera was used if classified, and family level used if genera was unclassified (g__). Average relative abundances on a log10 scale

among individuals possessing the taxon were extracted for each taxon within each ethnicity,

and the abundance for 12 families and genera were made into barchart figures (**Fig 4**). The

external whisker (AGP above, HMP below) depict the 75th quartile of abundance, and the

internal whisker depicts the 25th quartile. Pairwise Mann-Whitney-U tests were performed

between each pair of ethnicities using microbial abundances among all individuals, and were

Bonferroni corrected for the six comparisons within each taxon and dataset. Bonferroni

significant P-values are shown in the figure, and shown in bold if significance and direction

of change replicate in both datasets. Ubiquity shown above or below each bar was calculated

as the number of individuals in which that taxon was detected within the respective

ethnicity. Additional confirmation of ethnically varying abundance was also performed at

each taxonomic level (**S6 Table**), where the correlation of continuous age and BMI along

with categorically coded sex and ethnicity were simultaneously measured against the log 10

transformed relative abundance of each taxon among individuals possessing it using linear

regression (**S6 Table** - Abundance), and against the presence or absence of the taxon in all

individuals with logistic regression (**S6 Table** - Presence Absence). Significance is presented

for the models each with ethnicity alone, and with all metadata factors included (age, sex,

BMI), alongside Bonferroni corrected p-values, and individual effects of each metadata

factor.

Co-Occurrence Analysis

Bacterial taxonomy was collapsed at the family level, Spearman correlation was

calculated between each pair of families using SciPy (82), and clustermaps were generated

using seaborn (**S3 Fig**), and ethnic associations were drawn from **S5 Table**. Correlations

651 were masked where Bonferroni corrected Spearman p-values were >0.05, and clusters were

652 identified as the most prominent (strongest correlations) and abundance enriched.

653 Enrichment of ethnic association was evaluated by measuring the Mann-Whitney-U of

654 cluster families ethnic associations (p-values, **S5 Table**) compared to the ethnic associations

655 of non-cluster taxa. Cluster associated families were identified as having at least three

656 significant correlations with families within the cluster.

658 Christensenellaceae Analysis

659  The abundance of the family Christensenellaceae was input as relative abundance

660 across all individuals from the family level taxonomic table. Individuals were subset based

661 on the presence/absence of Christensenellaceae and BMIs were compared using a one tailed

662 Mann-Whitney-U test, then each was further subset by ethnicity and BMI compared using

663 one tailed Mann-Whitney-U tests and boxplots within each ethnicity (**Fig 5**).

665 Genetically Associated, Heritable, and Correlated Taxa Analysis

666  Genetically associated taxa from population heritability studies (34, 35, 39, 40) with

667 a minimum heritability (A in ACE models or H2r) >0.1, and from GWAS studies (35, 40) were

668 examined for exact taxonomic overlap with our 12 ethnically-associated taxa. The 42 genetic

669 variants associated with Unclassified Clostridiales are: rs16845116, rs586749, rs7527642,

670 rs10221827, rs5754822, rs4968435, rs17170765, rs1760889, rs6933411, rs2830259,

671 rs7318523, rs17763551, rs2248020, rs1278911, rs185902, rs2505338, rs6999713,

672 rs5997791, rs7236263, rs10484857, rs9938742, rs1125819, rs4699323, rs641527,

673 rs7302174, rs2007084, rs2293702, rs9350764, rs2170226, rs2273623, rs9321334,

rs6542797, rs9397927, rs2269706, rs4717021, rs7499858, rs10148020, rs7524581,

rs11733214, rs7587067 from (35). These 40 variants along with variants in **Table 1** except

for chr7:96414393 (total=49) were then assessed in 1,000 Genomes individuals for

significant differentiation across superpopulations (38). The 1,000 Genomes VCF files were

downloaded (ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/), and variants

with a minor allele frequency less than 0.01 were removed with F_{ST} calculated between each

pair of superpopulations using vcftools (83). The East Asian versus South Asian F_{ST} rates

were not used in the analysis. A custom script was used to examine the F_{ST} for each of the 49

variants and compare to the F_{ST} of all variants on the same chromosome and all variants

genome-wide for that pair of populations, with percentile calculated and the number of

variants with a higher F_{ST} divided by the total number of variants. The eQTL value and

significance for rs7587067 were drawn from the GTEx database (72).

Data and Code Availability

Code, scripts, and data underlying figures are publicly available from the GitHub

repository [https://github.com/awbrooks19/microbiota_and_ethnicity]. Individual

metadata (age, sex, ethnicity…) for the Human Microbiome Project are held under restricted

access available through dbGaP application [NCBI - dbGaP, Human Microbiome Project,

https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi?study_id=phs000228.v3.p1].

Acknowledgements

This work was supported by National Institutes of Health training grants

4T32GM08017810, 5T32GM08017809, and 5T32GM0817808 to AWB, the Vanderbilt Office

of Equity, Diversity and Inclusion to A.W.B. and S.R.B., the Vanderbilt Microbiome Initiative

to S.R.B., and the Alfred P. Sloan Foundation Fellowship to R.B.. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. We thank the American Society of Microbiology for supporting

travel to present this work. We would also like to thank Tony Capra, David Samuels, Patrick

Abbot, Antonis Rokas, and other members of the Vanderbilt Genetics Institute and

Bordenstein Lab for input. The authors acknowledge the Minnesota Supercomputing

Institute (MSI) at the University of Minnesota and the Advanced Computing Center for

Research and Education (ACCRE) at Vanderbilt University for providing resources that

contributed   to   the   research   results   reported   within   this   paper.

Contributions

A.W.B., S.P., R.B., and S.R.B. conceived and designed the research. A.W.B. performed,

analyzed, and interpreted all experiments with the exception of the RF analysis planned and

performed by S.P. and R.B. S.R.B. supervised all experimental designs, data analysis, and data

interpretation. All authors participated in manuscript preparation, editing, and final

approval.

Competing Financial Interests

The authors declare no competing financial interests.

Supplementary Table/Figure Legends:

722 **S1 Fig.** The average relative abundance of dominant microbial phyla for each ethnicity.

a

AGP Taxonomic Lineage	Differential Abundance FDR	RF % Importance for Most Distinguished Ethnicity
Oligella	6.81E-19	100% for Caucasian
Bifidobacterium	1.63E-5	94.49% for Asian-Pacific Islander
Ruminococcaceae	3.92E-6	82.47% for Asian-Pacific Islander
Christensenellaceae	7.35E-9	79.64% for Asian-Pacific Islander
Dialister	0.0358	71.73% for Asian-Pacific Islander
Barnesiellaceae	0.0106	68.41% for Asian-Pacific Islander
Megasphaera	1.57E-5	66.45% for Asian-Pacific Islander

HMP Taxonomic Lineage	Differential Abundance FDR	RF % Importance for Most Distinguished Ethnicity
Odoribacter	0.0006	100% for Asian-Pacific Islander
Succiniclasticum	1.31E-7	100% for African American
Rikenellaceae	0.0042	94.09% for Caucasian
Bacteroides	0.0102	80.69% for Asian-Pacific Islander
Dorea	0.0362	77.50% for African American
Erysipelotrichaceae	0.0362	61.10% for Caucasian
Christensenellaceae	0.0027	57.95% for Caucasian
Veillonella	0.0432	51.32% for Caucasian

c

Sampling Method	African American	Asian or Pacific Islander	Caucasian	Hispanic
SMOTE	8.79%	11.69%	11.70%	8.49%
Down	42.31%	31.82%	29.35%	60.81%

Sampling Method	African American	Asian or Pacific Islander	Caucasian	Hispanic
SMOTE	10.00%	7.56%	10.67%	12.29%
Down	40.00%	33.82%	38.51%	43.02%

S2 Fig. Summary of RF distinguishing taxa and out-of-bag error for each ethnicity. (A)

Importance of taxa for predicting each ethnicity using RF models with SMOTE sampling

approach are shown as percentage contributions, highlighted by color for each ethnicity.

Among the 10 most important taxa for each ethnicity, 9 overlap between the AGP and HMP

datasets (highlighted by the blue rectangular box), however which ethnicity they best

distinguish varies between the two datasets. (B) Taxa which are distinguishing in RF models

and have distinguishing differential abundance in **S5 Table**. The FDR corrected significance

for Kruskal-Wallis tests of differential abundance and the percent importance for the most

distinguished ethnicity of each in RF models are shown. (C) Out-of-bag error percentages for

the final RF classifier that was built using the optimal model parameters obtained from cross-

validation approach corresponding to each ethnicity and sampling procedure for both AGP

and HMP datasets.

S3 Fig. Abundance correlation of microbial families. Spearman correlation clustermaps

of bacterial abundance for families in the AGP and HMP. Numbers within boxes depict the

spearman correlation value with heatmap coloration from blue negative correlation (-1),

white no correlation (0), to red positive correlation (1). Positions have been masked based

on Bonferroni significance <0.05 for the total clustermap of all microbial families. Taxa

within boxes were identified as a highly correlated cluster, and taxa outside the boxes share

multiple correlations with those within the cluster. Blue taxonomic names indicate overlap

of taxa within boxes of both the AGP and HMP, while black indicate multiple correlations

with the clusters in both datasets. The ethnic association column depicts FDR corrected p-

values from Kruskal-Wallis tests in **S5 Table**, which are bolded if <0.05.

S4 Fig. Correlation of BMI with Christensenellaceae abundance. The relationship for

each individual between log10 transformed Christensenellaceae abundance on the y axis and

BMI on the x axis, with statistics slope, R^2, and p fit with a linear regression. Coloration of

each point indicates ethnicity: Yellow – African American; Blue – Asian-Pacific Islander;

Green – Hispanic; Red – Caucasian.

S1 Table. Demographic information for the AGP. Breakdown of age and BMI by sex and

ethnicity. Heatmaps were constructed within each statistic and category (bounded by black

box). The means for all sex and ethnic groups were used as the center (white), with higher

values indicated in red and lower in blue. HMP data is not shown because of data access

restrictions on participant metadata, available through dbGaP application. Additional sheets

depict proportions tests of ethnic structuring for 31 metadata factors, each on their own

sheet.

S2 Table. Microbiota distinguishability by ethnicity, age, sex and BMI. (A) AGP and HMP

ANOSIM distinguishability by ethnicity, age, sex, and BMI at a rarefaction depth of 1,000 and

across four ecological metrics (more details in table). (B) AGP ANOSIM distinguishability by

ethnicity, age, sex, and BMI at rarefaction depths of 1,000 and 10,000. (C) ANOSIM results

for consensus distance matrix while subsampling the maximum number of individuals from

each ethnic group. (D) BioEnv results of correlation between ethnicity, age, sex, and BMI

together with outcome as multivariate beta diversity distance matrices [Distance Matrix =

Ethnicity*x1 + Categorical Age*x2 + Categorical BMI*x3 + Sex*x4 + B]. (E) ANOSIM results

for consensus distance matrix when each ethnicity and group of ethnicities are sequentially

removed from the analysis.

S3 Table. Alpha diversity by ethnicity, age, sex and BMI. Alpha Diversity for Ethnicity,

Age, Sex, and BMI across varying rarefaction depths and beta diversity metrics in AG (4A, 4C-

E), and for ethnicity in the HMP (4B). Results are based on non-parametric permutation

based t-tests, and p-values are Bonferroni corrected within each factor of interest, depth, and

metric.

779 **S4 Table. Comparison of beta diversity distances for within and between ethnicities.**

780 All values depicted are Mann-Whitney-U p-values. (A) All distances between pairs of

781 individuals within each ethnicity were compared between ethnicities across rarefaction

782 depths 1,000 and 10,000, four beta diversity metrics, and with while subsampling over-

783 represented ethnicities. (B) All distances between pairs of individuals within and between

784 each ethnicity were compared between ethnicities. (C) Mean distances between pairs of

785 individuals within each ethnicity were compared between ethnicities. (D) Mean distances

786 between pairs of individuals within and between each ethnicity were compared between

787 ethnicities.

788

789 **S5 Table. Taxa which are differentially abundant by ethnicity, sex, BMI, and age in the**

790 **AGP and HMP.** Kruskal-Wallis results for differential taxa abundance across metadata

791 groupings, including FDR and Bonferroni corrected p-values, and taxa abundance averages

792 within each group. Metadata factors and taxonomic levels are separated by excel tabs.

793

794 **S6 Table. Taxa which are correlated with ethnicity, sex, BMI, and age in the AGP.**

795 Results of linear (Abundance) and logistic (Presence Absence) regression results for

796 differential taxa abundance across metadata factors separated by taxonomic level. Columns

797 in order indicate the taxon name, the number of individuals with non-zero abundance; then

798 the p-value for ethnicity alone, the p-value Bonferroni corrected, the f-test statistic, and R^2;

799 then the same values for the regression with ethnicity, age, sex, and BMI together; then the

800 abundances in each ethnic group, and finally the p-values for each factor broken down.

801

S7 Table. Genetic variants with taxa associations and detailed 1,000 Genomes

population differentiation rates (F_{ST}). Variants in red indicate the variant has at least one

F_{ST} above the 95th percentile for high differentiation between at least one pair of populations.

Columns I-BU represent the values for calculating variant F_{ST} and percentiles. The first two

spaces indicate the two superpopulations being compared. F_{ST} indicates the rate of

differentiation for that variant between that pair of populations. Higher indicates the

number of variants genome-wide with a higher F_{ST}, and total indicates the total genome-wide

variants examined. The columns with chromosome indicate the number of variants with

higher F_{ST} and total variants on the same chromosome as the variant of interest. Percent

indicates the number of variants with a higher F_{ST} divided by the total number of variants.

References:

1.   Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, et al. A core gut
microbiome in obese and lean twins. Nature. 2009;457(7228):480-4.
2.   Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial
gene catalogue established by metagenomic sequencing. Nature. 2010;464(7285):59-65.
3.   Huse SM, Ye Y, Zhou Y, Fodor AA. A core human microbiome as viewed through 16S
rRNA sequence clusters. PLoS One. 2012;7(6):e34242.
4.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, et al. Linking long-term
dietary patterns with gut microbial enterotypes. Science. 2011;334(6052):105-8.
5.   Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al. Diet drives
convergence in gut microbiome functions across mammalian phylogeny and within humans.
Science. 2011;332:970-4.
6.   Human Microbiome Project C. Structure, function and diversity of the healthy human
microbiome. Nature. 2012;486(7402):207-14.
7.   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet
rapidly and reproducibly alters the human gut microbiome. Nature. 2013;505(7484):559-63.
8.   Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al.
Human gut microbiome viewed across age and geography. Nature. 2012;486(7402):222-7.
9.   Davenport ER, Cusanovich DA, Michelini K, Barreiro LB, Ober C, Gilad Y. Genome-Wide
Association Studies of the Human Gut Microbiota. PLoS One. 2015;10(11):e0140301.

10.   Fierera N, Hamadyc M, Lauberb CL, Knight R. The influence of sex, handedness, and
washing on the diversity of hand surface bacteria. Procedings of the National Academy of
Sciences.105(46).
11.   Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of gut
microbiota in type 2 diabetes. Nature. 2012;490(7418):55-60.
12.   Frank DN, Allison AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-
phylogenetic characterization of microbial community imbalances in human inflammatory
bowel diseases. Procedings of the National Academy of Sciences. 2007(104):13780–5.
13.   Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes associated with
obesity and IBD. FEBS Lett. 2014;588(22):4223-33.
14.   Zackular JP, Baxter NT, Iverson KD, Sadler WD, Petrosino JF, Chen GY, et al. The gut
microbiome modulates colon tumorigenesis. MBio. 2013;4(6):e00692-13.
15.   Mason MR, Nagaraja HN, Camerlengo T, Joshi V, Kumar PS. Deep sequencing identifies
ethnicity-specific bacterial signatures in the oral microbiome. PLoS One. 2013;8(10):e77287.
16.   Ravela J, Gajera P, Abdob ZG, Schneiderc M, Koeniga SSK, McCullea SL, et al. Vaginal
microbiome of reproductive-age women. Procedings of the National Academy of Sciences.
2011;108:4680-7.
17.   Fettweis JM, Brooks JP, Serrano MG, Sheth NU, Girerd PH, Edwards DJ, et al. Differences
in vaginal microbiome in African American women versus women of European ancestry.
Microbiology. 2014;160(Pt 10):2272-82.
18.   Williams DR, Priest N, Anderson NB. Understanding associations among race,
socioeconomic status, and health: Patterns and prospects. Health Psychol. 2016;35(4):407-11.
19.   Mersha TB, Abebe T. Self-reported race/ethnicity in the age of genomic research: its
potential impact on understanding health disparities. Human Genomics. 2015;9(1):1.
20.   Kolde R, Franzosa EA, Rahnavard G, Hall AB, Vlamakis H, Stevens C, et al. Host genetic
variation and its microbiome interactions within the Human Microbiome Project. Genome Med.
2018;10(1):6.
21.   Clemente JC PE, Blaser MJ, Sandhu K, Gao K, Wang B, Magda M, Hidalgo G, et al. The
microbiome of uncontacted Amerindians. Science Advances. 2015;3.
22.   Rampelli S, Schnorr SL, Consolandi C, Turroni S, Severgnini M, Peano C, et al.
Metagenome Sequencing of the Hadza Hunter-Gatherer Gut Microbiota. Curr Biol.
2015;25(13):1682-93.
23.   Smits SA, Leach J, Sonnenburg ED, Gonzalez CG, Lichtman JS, Reid G, et al. Seasonal
cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania. Science.
2017;357:802-6.
24.   McDonald D, Birmingham A, Knight R. Context and the human microbiome.
Microbiome. 2015;3:52.
25.   Blekhman R, Goodrich JK, Huang K, Sun Q, Bukowski R, Bell JT, et al. Host genetic
variation impacts microbiome composition across human body sites. Genome Biol.
2015;16:191.
26.   Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, et al. Environment
dominates over host genetics in shaping human gut microbiota. Nature. 2018;555(7695):210-5.

27.   Deschasaux M, Bouter KE, Prodan A, Levin E, Groen AK, Herrema H, et al. Depicting the
composition of gut microbiota in a population with varied ethnic origins but shared geography.
Nat Med. 2018.

28.   Clarke KR. Non-parametric multivariate analyses of changes in community structure.
Australian Journal of Ecology. 1993;18:117-43.

29.   Clarke KR, Ainsworth M. A method of linking multivariate community structure to
environmental variables. Marine Ecology. 1993;92:205-19
.

30.   Knights D, Costello EK, Knight R. Supervised classification of human microbiota. FEMS
Microbiol Rev. 2011;35(2):343-59.

31.   Shannon CE. A mathematical theory of communication. Bell Syst Tech J. 1948;27:379-
423.

32.   Hester ER, Barott KL, Nulton J, Vermeij MJ, Rohwer FL. Stable and sporadic symbiotic
communities of coral and algal holobionts. ISME J. 2016;10(5):1157-69.

33.   Chawla NV, Bowyer KW, Hall LO and Kegelmeyer WP. SMOTE: Synthetic Minority Over-
sampling Technique. Journal of Artificial Intelligence Research. 2002;16:321-57.

34.   Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, et al. Human genetics
shape the gut microbiome. Cell. 2014;159(4):789-99.

35.   Goodrich JK, Davenport ER, Beaumont M, Jackson MA, Knight R, Ober C, et al. Genetic
Determinants of the Gut Microbiome in UK Twins. Cell Host Microbe. 2016;19(5):731-43.

36.   Fu J, Bonder MJ, Cenit MC, Tigchelaar EF, Maatman A, Dekens JA, et al. The Gut
Microbiome Contributes to a Substantial Proportion of the Variation in Blood Lipids. Circ Res.
2015;117(9):817-24.

37.   Pennisi E. Human Genetic Variation. Science. 2007;318(5858):1842-3.

38.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A
global reference for human genetic variation. Nature. 2015;526(7571):68-74.

39.   Lim MY, You HJ, Yoon HS, Kwon B, Lee JY, Lee S, et al. The effect of heritability and host
genetics on the gut microbiota and metabolic syndrome. Gut. 2016.

40.   Turpin W, Espin-Garcia O, Xu W, Silverberg MS, Kevans D, Smith MI, et al. Association of
host genome with intestinal microbial composition in a large healthy cohort. Nature Genetics.
2016;48(11):1413-7.

41.   Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees.
Am J Hum Genet. 1998;62(5):1198-211.

42.   Rothschild D, Weissbrod O, Barkan E, Korem T, Zeevi D, Costea PI, et al. Environmental
factors dominate over host genetics in shaping human gut microbiota composition. BioRxiv.
2017.

43.   Morgan XC, Tickle TL, Sokol H, Gevers D, Huttenhower C. Dysfunction of the intestinal
microbiome in inflammatory bowel disease and treatment. Genome Biology. 2012;7(979).

44.   Lewis JD, Chen EZ, Baldassano RN, Otley AR, Griffiths AM, Lee D, et al. Inflammation,
Antibiotics, and Diet as Environmental Stressors of the Gut Microbiome in Pediatric Crohn's
Disease. Cell Host Microbe. 2015;18(4):489-500.

45.   Goker M, Gronow S, Zeytun A, Nolan M, Lucas S, Lapidus A, et al. Complete genome
sequence of Odoribacter splanchnicus type strain (1651/6). Stand Genomic Sci. 2011;4(2):200-
9.

46.   Castaneda G, Liu B, Torres S, Bhuket T, Wong RJ. Race/Ethnicity-Specific Disparities in
the Severity of Disease at Presentation in Adults with Ulcerative Colitis: A Cross-Sectional Study.
Dig Dis Sci. 2017.
47.   Boucias DG, Cai Y, Sun Y, Lietze VU, Sen R, Raychoudhury R, et al. The hindgut lumen
prokaryotic microbiota of the termite Reticulitermes flavipes and its responses to dietary
lignocellulose composition. Mol Ecol. 2013;22(7):1836-53.
48.   LATHAM MJ, WOLIN MJ. Fermentation of Cellulose by Ruminococcus flavefaciens in the
Presence and Absence ofMethanobacterium ruminantium. Appl Environ Microbiol.
1977;34(3):297-301.
49.   Falony G, Raes J. Population-level analysis of gut microbiome variation. Science.
2016;352(6285):560-4.
50.   Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, et al. Gut Microbiota
and Extreme Longevity. Curr Biol. 2016;26(11):1480-5.
51.   Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, et al. Age-
Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic Inflammation, and
Macrophage Dysfunction. Cell Host Microbe. 2017;21(4):455-66 e4.
52.   Xie W, Wood AR, Lyssenko V, al. e. Genetic Variants Associated With Glycine
Metabolism and Their Role in Insulin Sensitivity and Type 2 Diabetes. Diabetes. 2013;62.
53.   Williams SR, Yang Q, Chen F, Liu X, Keene KL, Jacques P, et al. Genome-wide meta-
analysis of homocysteine and methionine metabolism identifies five one carbon metabolism
loci and a novel association of ALDH1L1 with ischemic stroke. PLoS Genet.
2014;10(3):e1004214.
54.   Petersen LM, Bautista EJ, Nguyen H, Hanson BM, Chen L, Lek SH, et al. Community
characteristics of the gut microbiomes of competitive cyclists. Microbiome. 2017;5(1):98.
55.   Nakamura N, Lin HC, McSweeney CS, Mackie RI, Gaskins HR. Mechanisms of microbial
hydrogen disposal in the human colon and implications for health and disease. Annu Rev Food
Sci Technol. 2010;1:363-95.
56.   Parthasarathy G, Chen J, Chen X, Chia N, O'Connor HM, Wolf PG, et al. Relationship
Between Microbiota of the Colonic Mucosa vs Feces and Symptoms, Colonic Transit, and
Methane Production in Female Patients With Chronic Constipation. Gastroenterology.
2016;150(2):367-79 e1.
57.   Jackson CS, Oman M, Patel AM, Vega KJ. Health disparities in colorectal cancer among
racial and ethnic minorities in the United States. J Gastrointest Oncol. 2016;7(Suppl 1):S32-43.
58.   Lopetuso LR, Scaldaferri F, Petito V, Gasbarrini A. Commensal Clostridia: leading players
in the maintenance of gut homeostasis. Gut Pathogens. 2013.
59.   Sy DF. The Center for Asian Health Engages Communities in Research to Reduce Asian
American Health Disparities. US Department of Health & Human Services, National Institute on
Minority Health and Health Disparities.
60.   Hwang H. Colorectal Cancer Screening among Asian Americans. Asian Pacific Journal of
Cancer Prevention. 2013;14(7):4025-32.
61.   Oh KM, Kreps GL, Jun J. Colorectal Cancer Screening Knowledge, Beliefs, and Practices of
Korean Americans. American Journal of Health Behavior. 2013;37(3):381-94.
62.   Sankaranarayanan R, Ramadas K, Qiao Y-l. Managing the changing burden of cancer in
Asia. BMC Medicine. 2014;12(3).

63.   Pourhoseingholi MA. Increased burden of colorectal cancer in Asia. World J Gastrointest
Oncol. 2012;4(4):68-70.
64.   Pourhoseingholi MA, Vahedi M, Baghestani AR. Burden of gastrointestinal cancer in
Asia; an overview. Gastroenterology and Hepatology. 2015.
65.   Pourhoseingholi MA. Epidemiology and burden of colorectal cancer in Asia-Pacific
region: what shall we do now? Translational Gastrointestinal Cancer. 2014;3(4):169-73.
66.   Report CHDaI. 2013.
67.   Cao H, Liu X, An Y, Zhou G, Liu Y, Xu M, et al. Dysbiosis contributes to chronic
constipation development via regulation of serotonin transporter in the intestine. Sci Rep.
2017;7(1):10322.
68.   Mosca A, Leclerc M, Hugot JP. Gut Microbiota Diversity and Human Diseases: Should We
Reintroduce Key Predators in Our Ecosystem? Front Microbiol. 2016;7:455.
69.   Zhang X, Zhang D, Jia H, Feng Q, Wang D, Liang D, et al. The oral and gut microbiomes
are perturbed in rheumatoid arthritis and partly normalized after treatment. Nat Med.
2015;21(8):895-905.
70.   Singh VP, Proctor SD, Willing BP. Koch's postulates, microbial dysbiosis and
inflammatory bowel disease. Clin Microbiol Infect. 2016;22(7):594-9.
71.   Sherry ST WM, Kholodov M, Baker J, Phan L, Smigielski EM, Sirotkin K. dbSNP: the NCBI
database of genetic variation. Nucleic Acids Research. 2001;29(308).
72.   Consortium GT. The Genotype-Tissue Expression (GTEx) project. Nat Genet.
2013;45(6):580-5.
73.   Qiu X, Wei R, Li Y, Zhu Q, Xiong C, Chen Y, et al. NEDL2 regulates enteric nervous system
and kidney development in its Nedd8 ligase activity-dependent manner. Oncotarget.
2016;7(21).
74.   Wei R, Qiu X, Wang S, Li Y, Wang Y, Lu K, et al. NEDL2 is an essential regulator of enteric
neural development and GDNF/Ret signaling. Cell Signal. 2015;27(3):578-86.
75.   O'Donnell AM, Coyle D, Puri P. Decreased expression of NEDL2 in Hirschsprung's
disease. J Pediatr Surg. 2016;51(11):1839-42.
76.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al.
QIIME allows analysis of high-throughput community sequencing data. Nat Method.
2010;7:335-6.
77.   DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a
chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl Environ
Microbiol. 2006;72(7):5069-72.
78.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J.
2012;6(8):1621-4.
79.   Anderson MJ. A new method for non-parametric multivariate analysis of variance.
Australian Journal of Ecology. 2001;26:32-46.
80.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. PLoS One. 2013;8:e61217.
81.   Kuhn M. A short introduction to the caret package. 2017.
82.   Jones E, Oliphant T, Peterson P. Open Source Scientific Tools for Python. 2001.

83.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
format and VCFtools. Bioinformatics. 2011;27(15):2156-8.
1009