Gut Microbiota Diversity across Ethnicities in the United States

Authors:

Andrew W. Brooks[1,2], Sambhawa Priya[3,4,5], Ran Blekhman[3,4], Seth R. Bordenstein[1,2,6,7*]

Affiliations:

[1] Vanderbilt Genetics Institute, Vanderbilt University, Nashville, TN, USA.

[2] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3]Department of Genetics, Cell Biology, and Development, University of Minnesota,

Minneapolis, MN, USA

[4]Department of Ecology, Evolution, and Behavior, University of Minnesota, Minneapolis,

MN, USA

[5] Bioinformatics and Computational Biology Program, University of Minnesota,

Minneapolis, MN, USA

[6] Department of Pathology, Microbiology, and Immunology, Vanderbilt University,

Nashville, TN, USA.

[7]Vanderbilt Institute for Infection, Immunology and Inflammation, Vanderbilt University,

Nashville, TN, USA.

[*] Corresponding author. E-mail: s.bordenstein@vanderbilt.edu (S.R.B)

Abbreviations:

AGP – American Gut Project

ANOSIM – Analysis of Similarity

AUC – Area Under the Curve

BMI – Body Mass Index

F_{ST} – Fixation Index

GWAS - Genome-Wide Association Studies

HMP – Human Microbiome Project

MAF - Minor Allele Frequency

OTU – Operational Taxonomic Unit

PERMANOVA - Permutational Multivariate Analysis of Variance

RF – Random Forest

ROC – Receiver Operating Characteristic

SMOTE – Synthetic Minority Over-sampling Technique

Abstract:

Composed of hundreds of microbial species, the composition of the human gut

microbiota can vary with chronic diseases underlying health disparities that

disproportionally affect ethnic minorities. However, the influence of ethnicity on the gut

microbiota remains largely unexplored and lacks reproducible generalizations across

studies. By distilling associations between ethnicity and gut microbiota variation in two

American datasets including 1,673 individuals, we report 12 microbial genera and families

that reproducibly vary by ethnicity. Interestingly, a majority of these microbial taxa,

including the most heritable bacterial family, Christensenellaceae, overlap with genetically-

associated lineages and form co-occurring clusters of taxa linked by similar fermentative and

methanogenic metabolic processes. These results demonstrate recurrent associations

between specific taxa in the gut microbiota and ethnicity, providing hypotheses for

examining specific members of the gut microbiota as mediators of health disparities.

Introduction:

The human gut microbiota at fine resolution varies extensively between individuals (1-3), and this variability frequently associates with diet(4-7), age(6, 8, 9), sex(6, 9, 10), body mass index (BMI) (1, 6), and diseases presenting as health disparities (11-14). The overlapping risk factors and burden of many chronic diseases disproportionally affect ethnic minorities in the United States, yet the underlying biological mechanisms mediating these substantial disparities largely remain unexplained. Recent evidence is consistent with the hypothesis that ethnicity associates with microbial abundance, specifically in the oral cavity, gut, and vagina (15-17). Ethnicity can capture many facets of biological variation including social, economic and cultural variation as well as aspects of human genetic variation and biogeographical ancestry (18, 19). Despite the importance of understanding the interconnections between ethnicity, microbiota, and health disparities, there are no replicated generalizations about the influence of ethnicity on variation in the gut microbiota and specific microbial taxa in diverse American populations, even for healthy individuals (6).

Here, we comprehensively examine connections between self-declared ethnicity and gut microbiota variation in more than a thousand individuals sampled by the American Gut Project (AGP, N=1375) (20) and the Human Microbiome Project (HMP, N=298) (6). Human genetic diversity in the HMP has been shown to associate with differences in microbiota composition, and it has been demonstrated that genetic population structure within the HMP partially delineates self-declared ethnicity (21). Ethnicity was not found to have a significant association with microbiota composition in a middle-eastern population, however microbiota influencing factors such as lifestyle and environment across participants was

homogenous compared to the ethnic, sociocultural, economic, and dietary diversity found

within the United States (22).

Results:

Ethnicity subtly demarcates microbiota

We first evaluate gut microbiota distinguishability between AGP ethnicities (**Fig 1A**,

family taxonomic level, Asians-Pacific Islanders (N=88), Caucasians (N=1237), Hispanics

(N=37), and African Americans (N=13)), sexes (female (N=657), male (N=718)), age groups

(years grouped by decade), and categorical BMI (underweight (N=70), normal (N=873),

overweight (N=318), and obese (N=114)) (Demographic details in **S1 Table**). 97%

Operational Taxonomic Units (OTUs) generated for each dataset are utilized throughout to

maintain consistency with other published literature, however microbial taxonomy of the

HMP is reassigned using the Greengenes reference database (23). While interindividual

microbiota heterogeneity dominates, Analyses of Similarity (ANOSIM) reveal subtle but

significant degrees of total microbiota distinguishability for ethnicity, BMI, and sex, but not

for age (**Fig 1B**, Ethnicity; **Fig 1C**, BMI; **Fig 1D**, Sex; **Fig 1E**, Age) (24). Recognizing that subtle

microbiota distinguishability between ethnicities may be spurious, we independently

replicate the ANOSIM results from HMP African Americans (N=10), Asians (N=34),

Caucasians (N=211) and Hispanics (N=43) (**S2A Table**, R=0.065, p=0.044), and observe no

significant distinguishability for BMI, sex, and age. Higher rarefaction depths increase

microbiota distinguishability in the AGP across various beta diversity metrics and

categorical factors (**S2B Table**), and significance increases when individuals from

overrepresented ethnicities are subsampled from the average beta diversity distance matrix

(**S2C Table**). Supporting the ANOSIM results, Permutational Multivariate Analysis of

Variance (PERMANOVA) models with four different beta diversity metrics showed that while

all factors had subtle but significant associations with microbiota variation when combined

in a single model, effect sizes were highest for ethnicity in 7 out of 8 comparisons across beta

diversity metrics and rarefaction depths in the AGP and HMP (**S2D Table**). We additionally

test microbiota distinguishability by measuring the correlation between beta diversity and

ethnicity, BMI, sex, and age with an adapted BioEnv test (**S2E Table**) (25). Similar degrees

of microbiota structuring occur when all factors are incorporated (Spearman Rho=0.055, p-

values: Ethnicity=0.057, BMI<0.001, Sex<0.001, Age=0.564). Firmicutes and Bacteroidetes

dominated the relative phylum abundance, with each representing between 35% and 54%

of the total microbiota across ethnicities (**S1 Fig**).

Fig 1. Gut microbiota composition and distinguishability by ethnicity, sex, age and

BMI. (A) The average relative abundance of dominant microbial families for each ethnicity.

(B-E) Principle coordinates analysis plots of microbiota Bray-Curtis beta diversity and

ANOSIM distinguishability for: (B) Ethnicity, (C) Sex, (D) Age, (E) BMI. In B-E, each point

represents the microbiota of a single sample, and colors reflect metadata of that individual.

Caucasian points are reduced in size to allow clearer visualization.

We next test for ethnicity signatures in the gut microbiota by analyzing alpha and beta

diversity, abundance and ubiquity distributions, distinguishability, and classification

accuracy (26). Shannon's Alpha Diversity Index (27), which weights both microbial

community richness (Observed OTUs) and evenness (Equitability), significantly varies

across ethnicities in the AGP dataset (Kruskal Wallis, p=2.8e-8) with the following ranks:

Hispanics > Caucasians > Asian-Pacific Islanders > African Americans (**Fig 2A**). Some of these

results replicate in the HMP dataset, where we find a significantly lower Shannon diversity

for Asian-Pacific Islanders relative to Caucasians, and a trend of lower Shannon diversity for

Asian-Pacific Islanders relative to Hispanics. Five alpha diversity metrics, two rarefaction

depths, and separate analyses of Observed OTUs and Equitability generally confirm the

results (**S3A Table**).

Fig 2. Ethnicity associates with diversity and composition of the gut microbiota. (A) Center lines of each boxplot depict the median by which ethnicities were ranked from low (left) to high (right); the lower and upper end of each box represent the 25th and 75th quartiles respectively; whiskers denote the 1.5 interquartile range, and black dots represent individual samples. Lines in the middle of violin plots depict the mean, and p-values are Bonferroni corrected within each dataset. (B) Left extending violin plots represent intra-ethnic distances for each ethnicity, and right extending violin plots depict all inter-ethnic distances. Center lines depict the mean beta diversity. Significance bars above violin plots

depict Bonferroni corrected pairwise Mann-Whitney-U comparisons of the intra-intra- and intra-inter-ethnic distances. (C) Within each ethnicity, OTUs shared by at least 50% of samples. Colored lines represent a robust ordinary least squares regression within OTUs of each ethnicity, shaded regions represent the 95% confidence interval, R^2 denotes the regression correlation, the OTUs column indicates the number of OTUs with >50% ubiquity for that ethnicity, Mean A/U is the average abundance/ubiquity ratio, and the p_{adj} is the regression significance adjusted and Bonferroni corrected for the number of ethnicities.

If ethnicity impacts microbiota composition, pairwise beta diversity distances (ranging from 0/completely dissimilar to 1/identical) will be greater between ethnicities than within ethnicities. While average gut microbiota beta diversities across all individuals are high (**Fig 2B**, Bray-Curtis=0.808), beta diversities between individuals of the same ethnicity (intra-ethnic, Bray-Curtis=0.806) are subtly, but significantly, lower than those between ethnicities in both the AGP (inter-ethnic, Bray-Curtis=0.814) and HMP datasets (intra-ethnic, Bray-Curtis=0.870 versus inter-ethnic, Bray-Curtis=0.877). We confirm AGP results by subsampling individuals from overrepresented ethnicities across beta metrics and rarefaction depths (**S4A-4B Tables**). Finally, we repeat analyses across beta metrics and rarefaction depths using only the average distance of each individual to all individuals from the ethnicity to which they are compared (**S4C-4D Tables**).

Next, we explore inter-ethnic variation in the number of OTUs shared in at least 50% of individuals within an ethnicity. Out of 5,591 OTUs in the total AGP dataset, 101 (1.8%) meet this ubiquity cutoff in all ethnicities, and 293 (5.2%) unique OTUs meet the cutoff within at least one ethnicity. Hispanics share the most ubiquitous OTUs and have the lowest

average abundance/ubiquity (A/U) ratio (**Fig 2C**), indicating higher stability of lower

abundance shared OTUs (28). This result potentially explains their significantly lower intra-

ethnic beta diversity distance and thus higher microbial community overlap relative to the

other ethnicities (**Fig 2B**). Comparisons in the AGP between the higher sampled Hispanic,

Caucasian, and Asian-Pacific Islander ethnicities also reveal a trend wherein higher intra-

ethnic community overlap (**Fig 2B**) parallels higher numbers of ubiquitous OTUs (**Fig 2C**),

higher Shannon Alpha diversity (**Fig 2A**), and higher stability of ubiquitous OTUs as

measured by the abundance/ubiquity (A/U) ratio (**Fig 2C**).

We next assess whether a single ethnicity disproportionately impacts total gut

microbiota distinguishability in the AGP by comparing ANOSIM results from the consensus

beta diversity distance matrix when each ethnicity is sequentially removed from the analysis

(**Fig 3A** and **S2E Table**). Distinguishability remains unchanged when the few African

Americans are removed, but is lost upon removal of Asian-Pacific Islanders or Caucasians

(**Fig 3A**). Notably, removal of Hispanics increases distinguishability among the remaining

ethnicities, which may be due to higher degree of beta diversity overlap observed between

Hispanics and other ethnicities (**S4B Table**). Results conform across rarefaction depths and

beta   diversity   metrics   (**S2F   Table**),   and   pairwise   combinations   show   strong

distinguishability between African Americans and Hispanics (ANOSIM, R=0.234, p=0.005),

and Asian-Pacific Islanders and Caucasians (ANOSIM, R=0.157, p<0.001).

Finally, to complement evaluation with ecological alpha and beta diversity we

implement a random forest supervised learning algorithm to classify gut microbiota from

genus level community profiles into their respective ethnicity. We build four one-versus-all

binary classifiers to classify samples from each ethnicity compared to the rest, and use two

different sampling approaches to train the models, Synthetic Minority Over-sampling Technique (SMOTE) (29) and down-sampling, for overcoming uneven representation of ethnicities in both the datasets (see Methods). Given that the area under the receiver operating characteristic (ROC) curve (or AUC) of a random guessing classifier is 0.5, the models classify each ethnicity fairly well (**Fig 3B**) with average AUCs across sampling techniques and datasets of 0.78 for Asian-Pacific Islanders, 0.76 for African Americans, 0.69 for Hispanics, and 0.70 for Caucasians.

Fig 3. Microbiota distinguishability and classification ability across ethnicities. (A) ANOSIM distinguishability between all combinations of ethnicities. Symbols depict specific ethnicities included in the ANOSIM tests, and boxes denote the R-value as a heatmap, where

white indicates increasing and black indicates decreasing distinguishability relative to the R-

value with all ethnicities. (B) Average ROC curves (for 10-fold cross-validation) and

prediction performance metrics for one-versus-all random forest classifiers for each

ethnicity, using SMOTE (29) and down subsampling approaches for training.

Recurrent taxa associations with ethnicity

Subtle to moderate ethnicity-associated variation in microbial communities may in

part be strongly driven by differential abundance of certain microbial lineages. We find that

16.2% (130/802) of the AGP taxa and 20.6% (45/218) of HMP taxa across all classification

levels (i.e. phylum to genus, **S5 Table**) significantly vary in abundance across ethnicities

(Kruskal-Wallis, $p_{FDR}<0.05$). Between datasets, 19.2% (25/130) of the AGP and 55.6%

(25/45) of the HMP varying lineages replicate in the other dataset, representing a

significantly greater degree of overlap than would be expected by chance (AGP replicated,

Fisher's exact one-tailed test, expected 5% overlap (7 overlapping vs. 123 not overlapping)

and observed (25 overlapping vs. 105 not overlapping), p=5.26e-4; HMP replicated, Fisher's

exact test, expected 5% overlap (2 overlapping vs. 43 not overlapping) and observed (25

overlapping vs. 20 not overlapping), p=4.72e-8; ethnic permutation analysis of overlap,

p<0.001 each taxonomic level and all taxonomic levels combined). The highest replication of

taxonomic lineages varying by abundance occurs with 22.0% (9/41) of families across

datasets, followed by genus with 13.4% (9/67).

Among 18 reproducible lineages, we categorize 12 as unique (**Fig 4**) and exclude 6

where nearly identical abundance profiles between family/genus taxonomy overlap.

Comparing relative abundance differences between pairs of ethnicities for these 12 taxa in

AGP and HMP reveals 20 out of 30 significant (p<0.05, Mann-Whitney-U) differences

replicated. Intriguingly, all reproducible pairwise differences are a result of decreases in

Asian-Pacific Islanders (**Fig 4**).

We also test taxon abundance and presence/absence associations with ethnicity

separately in the AGP using linear and logistic regression models respectively, and we repeat

the analysis while incorporating categorical sex and continuous age and BMI as covariates

(**S6 Table**). Clustering microbial families based on their abundance correlation reveals two

co-occurrence clusters: (i) a distinct cluster of six Firmicutes and Tenericutes families in the

HMP and (ii) an overlapping but more diverse cluster of 20 families in the AGP (**S2 Fig**). Nine

of the 12 taxa found to recurrently vary in abundance across ethnicities are represented in

these clusters (**Fig 4**), with four appearing within both clusters, and the other five appearing

either within or closely correlated with members of both clusters (**S2 Fig**). Further, 90%

(18/20) of families in the AGP cluster and 66% (4/6) of taxa in the HMP cluster significantly

vary in abundance across ethnicities. Taken together, these results establish general overlap

of the most significantly ethnically-associated taxa between the three methods,

reproducibility of microbial abundances that vary between ethnicities across datasets, and

patterns of co-occurrence among these taxa which could suggest they are functionally linked.

Fig 4. Ethnicity-associated taxa match between the HMP and AGP. Barplots depict the

log10 transformed relative abundance for individuals possessing the respective taxon within

each ethnicity, ubiquity appears above (AGP) or below (HMP) bars, and the 25th and 75th

percentiles are shown with extending whiskers. Mann-Whitney-U tests evaluate differences

in abundance and ubiquity for all individuals between pairs of ethnicities; for example, the

direction of change in Victivallaceae is driven by ubiquity while abundance is higher for

those possessing the taxon. Significance values are Bonferroni corrected for the six tests

within each taxon and dataset, and bold p-values indicate that significance (p<0.05) and

direction of change replicate in the AGP and HMP.

Most heritable taxon varies by ethnicity

Identified as the most heritable human gut taxon (30, 31), the family

Christensenellaceae exhibits the second strongest significant differences in abundance

across ethnicities in both AGP and HMP datasets (**S5 Table**, Family: AGP, Kruskal-Wallis,

p_{FDR}=1.55e-9; HMP, Kruskal-Wallis, p_{FDR}=0.0019). Additionally, Christensenellaceae is

variable by sex and BMI (AGP: Sex, Kruskal-Wallis, p_{FDR}=1.22e-12; BMI, Kruskal-Wallis,

p_{FDR}=0.0020), and represents some of the strongest pairwise correlations with other taxa in

both co-occurrence clusters (**S2 Fig**). There is at least an eight-fold and two-fold reduction

in average Christensenellaceae abundance in Asian-Pacific Islanders relative to the other

ethnicities in the AGP and HMP respectively (**S5 Table**), and significance of all pairwise

comparisons in both datasets show reduced abundance in Asian-Pacific Islanders (**Fig 4**).

Abundance in individuals possessing Christensenellaceae and presence/absence across all

individuals significantly associate with ethnicity (**S6 Table**, Abundance, Linear Regression,

$p_{Bonferroni}$=0.006; Presence/Absence, Logistic Regression, $p_{Bonferroni}$=8.802e-6), but there was

only a slight correlation between the taxon's relative abundance and BMI (**S3 Fig**).

Confirming previous associations with lower BMI(32), we observe that AGP individuals with

Christensenellaceae also have a significantly lower BMI (Mean BMI, 23.7±4.3) than

individuals without it (Mean BMI, 25.0±5.9; Mann-Whitney-U, p<0.001). This pattern is

separately reflected in African Americans, Asian-Pacific Islanders, and Caucasians but not

Hispanics (**Fig 5**), suggesting that each ethnicity may have different equilibria between the

taxon's abundance and body weight.

Fig 5. Christensenellaceae variably associates with BMI across ethnicities. Boxplots of
BMI for individuals without (unfilled boxplots) and with (filled boxplots)
Christensenellaceae. Significance was determined using one-tailed Mann-Whitney-U tests
for lower continuous BMI values. Black lines indicate the mean relative abundance; the lower
and upper end of each box represent the 25th and 75th quartiles respectively; and whiskers
denote the 1.5 interquartile range.

Genetic- and ethnicity-associated taxa overlap

Many factors associate with human ethnicity, including a small subset of population
specific genetic variants (estimated ~0.5% genome wide) that vary by biogeographical
ancestry (33, 34), and self-declared ethnicity in the HMP is delineated by population genetic
structure (21). Here we investigate whether ethnicity-associated taxa overlap with (i) taxa
that have a significant population genetic heritability in humans (30, 31, 35, 36) and (ii) taxa
linked with human genetic variants in two large Genome-Wide Association Studies (GWAS)-

microbiota analyses (31, 36). All except one recurrent ethnicity-associated taxa are heritable

in at least one study, with seven replicating in three or more studies (**Table 1**). Likewise,

abundance variation in seven recurrent ethnicity-associated taxa demonstrate significant

GWAS associations with at least one variant in the human genome, therefore we assess

whether any genetic variants associated with differences in microbial abundance show

significant rates of differentiation (F_{ST}) between 1,000 genomes superpopulations (34). Out

of 49 variants associated with ethnically varying taxa, 21 have higher F_{ST} values between at

least one pair of populations than that of 95% of other variants on the same chromosome

and across the genome, and the F_{ST} values of five variants associated with Clostridiaceae

abundance rank above the top 99% (**S7 Table**). Since taxa that vary across ethnicities exhibit

lower abundance in Asian-Pacific Islanders, it is notable that the F_{ST} values of 18 and 11

variant comparisons for East Asian and South Asian populations, respectively, are above that

of the 95% rate of differentiation threshold from African, American, or European

populations. Critically, the microbiota and 1,000 genomes datasets are not drawn from the

same individuals, and disentangling the role of genetic from social and environmental factors

will still require more controlled studies.

Recurrent Ethnicity-Associated Taxa	Heritability	Genetic Associations
Family: Peptococcaceae	0.1213 [A], 0.2154 [C], 0.26 [E]	rs143179968 [E]
Family: Dehalobacteriaceae	0.6878 [B], 0.3087 [C]	
Family: Christensenellaceae	0.3819 [A], 0.6170 [B], 0.4230 [C],	
Order: Clostridiales, Family: Unclassified	0.2914 [A], 0.4020 [B], 0.1330 [C]	*40 Genetic Variants [C]
Genus: *Veillonella*	0.1370 [A], 0.2168 [D]	rs347941 [C]
Order: RF39, Family: Unclassified	0.2341 [A], 0.6618 [B], 0.3074 [C]	rs4883972 [C]
Family: Verrucomicrobiaceae	0.1257 [A], 0.5973 [B], 0.1394 [C]	
Family: Victivallaceae		
Family Odoribacteraceae	0.1389 [A], 0.1917 [D], 0.34 [E]	chr7:96414393 [E], rs115795847 [E]
Genus: *Odoribacter*	0.1916 [D]	
Family: Rikenellaceae	0.1299 [D], 0.29 [E]	rs17098734 [C], rs3909540 [C], rs147600757 [E],
Family: Coriobacteraceae, Genus:	0.1364 [A], 0.2822 [B], 0.1609 [C]	rs9357092 [E]

Table 1. Recurrent ethnicity-associated taxa overlap heritable and genetically-associated taxa. The table shows population genetic heritability estimates and associated genetic variants for the 12 recurrent ethnically varying taxa. The minimum heritability cutoff was chosen as >0.1, and only exactly overlapping taxonomies were considered. Studies examined: [A]UKTwins (2014, A in ACE model) (30), [B]Yatsunenko (2014, A in ACE model) (30), [C]UKTwins (2016, A in ACE model) (31), [D]Lim (2016, H2r in SOLAR (37)) (35), [E]Turpin (2016, H2r in SOLAR (37)). *indicates excessive variants were excluded from table.

Discussion:

Many common diseases associate with microbiota composition and ethnicity, raising the central hypothesis that microbiota variation between ethnicities can occasionally serve as a mediator of health disparities. American's self-declared ethnicity can capture socioeconomic, cultural, geographic, dietary and genetic diversity, and a similarly complex array of interindividual and environmental factors influence total microbiota composition, resulting in challenges when trying to consistently recover variation in total gut microbiota between ethnicities. These challenges inform the importance of reproducibility, both through confirmation across analytical methods and replication across study populations. In order to more fully evaluate this hypothesis, baseline generalizations are drawn here about the impact of ethnicity on gut microbiota variation in healthy individuals, and is concordant with recent literature in single populations suggesting that ethnicity plays a subtle but reproducible role in microbiota assembly (15-17, 21, 22).

Whether shaped through socioeconomic, dietary, healthcare, genetic, or other ethnicity-related factors, the replicated, varying taxa represent sources for novel hypotheses

addressing health disparities. For instance, the family Odoribacteraceae and genus

Odoribacter are primary butyrate producers in the gut, and have been negatively linked to

severe forms of Crohn's disease and Ulcerative Colitis in association with reduced butyrate

metabolism (38-40). Asian-Pacific Islanders possess significantly less Odoribacteraceae and

Odoribacter than Hispanics and Caucasians in both datasets, and severity of Ulcerative Colitis

upon hospital admission has been shown to be significantly higher in Asian Americans (41).

Considering broader physiological roles, several ethnicity-associated taxa are primary gut

anaerobic fermenters and methanogens (42, 43), and associate with lower BMI and blood

triglyceride levels (32, 44). Indeed, Christensenellaceae, Odoribacteraceae, *Odoribacter*, and

the class Mollicutes containing RF39 negatively associate with metabolic syndrome and

demonstrate significant population heritability in twins (35). Implications for health

outcomes warrant further investigation, but could be reflected by positive correlations of

Odoribacteraceae, *Odoribacter*, Coriobacteriaceae, Christensenellaceae, and the dominant

Verrucomicrobiaceae lineage *Akkermansia* with old age (45, 46). Moreover, these findings

raise the importance of controlling for ethnicity in studies linking microbiota variation to

disease because associations between specific microbes and a disease could be confounded

by ethnicity of the study subjects.

Based on correlations in individual taxon's abundance, a similar pattern of co-

occurrence previously identified as the Christensenellaceae 'consortium' includes 11 of the

12 recurrent ethnically varying taxa (30), and members of this consortium associate with

genetic variation in the human formate oxidation gene *ALDH1L1* which is a genetic risk factor

for stroke (31, 47, 48). Formate metabolism is a key step in the pathway reducing carbon

dioxide to methane (49, 50), and increased methane associates with increased Rikenellaceae,

Christensenellaceae, Odoribacteraceae and *Odoribacter* (51). Products of methanogenic

fermentation pathways include short chain fatty acids such as butyrate, which through

reduction of pro-inflammatory cytokines has been linked to cancer cell apoptosis and

reduced risk of colorectal cancer (52, 53). Asian Americans are the only ethnic group where

cancer surpasses heart disease as the leading cause of death, and over 70% of Asian

Americans were born overseas, which can affect assimilation into western lifestyles, leading

to reduced access to healthcare and screening, and proper medical education (52, 54-56).

Indeed, as countries in Asia shift toward a more western lifestyle, the incidence of cancers,

particularly gastrointestinal and colorectal cancers, are increasing rapidly, possibly

indicating incompatibilities between traditionally harbored microbiota and western

lifestyles (57-60). Asian Americans have higher rates of type 2 diabetes and pathogenic

infections than Caucasians (61), and two metagenomic functions enriched in control versus

type 2 diabetes cases appear to be largely conferred by cluster-associated butyrate-

producing and motility-inducing Verrucomicrobiaceae and Clostridia lineages reduced in

abundance among AGP and HMP Asian-Pacific Islanders (11). Both induction of cell motility

and butyrate promotion of mucin integrity can protect against pathogenic colonization and

associate with microbial community changes (11, 53, 62). Levels of cell motility and butyrate

are key factors suspected to underlie a range of health disparities including inflammatory

bowel disease, arthritis, and type 2 diabetes (11, 63-65). Patterns of ethnically varying taxa

across ethnicities could result from many factors including varying diets, environmental

exposures, sociocultural influences, human genetic variation and others. However,

regardless of the mechanisms dictating assembly, these results suggest there is a

reproducible, co-occurring group of taxa linked by similar metabolic processes known to

promote homeostasis.

The utility of this work is establishing a framework for studying ethnicity-associated

taxa and hypotheses of how changes in abundance or presence of these taxa may or may not

shape health disparities, many of which also have genetic components. Differing in allele

frequency across three population comparisons and associated with the abundance of

Clostridiales, the genetic variant rs7587067 has a significantly higher frequency in African

(Minor Allele Frequency (MAF)=0.802) versus East Asian (MAF=0.190, F_{ST}=0.54,

Chromosome=98.7%, Genome-Wide=98.9%), admixed American (MAF=0.278, F_{ST}=0.44,

Chromosome=99.0%, Genome-Wide=99.1%), and European populations (MAF=0.267,

F_{ST}=0.45, Chromosome=98.7.3%, Genome-Wide=98.7%). This intronic variant for the gene

HECW2 is a known eQTL (GTeX, eQTL Effect Size=-0.18, p=7.4e-5) (66, 67), and *HECW2*

encodes a ubiquitin ligase linked to enteric gastrointestinal nervous system function through

maintenance of endothelial lining of blood vessels (68, 69). Knockout of *HECW2* in mice

reduced enteric neuron networks and gut motility, and patients with Hirschsprung's disease

have diminished localization of *HECW2* to regions affected by loss of neurons and colon

blockage when compared to other regions of their own colon and healthy individuals (70).

Hirschsprung's disease presenting as full colon blockage is rare and has not undergone

targeted examination as a health disparity, however a possible hypothesis is that lower

penetrance of the disease in individuals with the risk allele at rs7587067 could lead to

subtler effects on gut motility resulting in Clostridiales abundance variation.

Another example is that the abundance of Rikenellaceae in the gut is strongly and

reproducibly associated with variant rs62171178, which was identified as an eQTL for gene

PHOSPHO2 within human gut tissue (36, 67). The primary substrate of the protein encoded

by *PHOSPHO2* is vitamin B6 (71), which shows increased deficiencies in germ free compared

to conventionally reared rats (72). Interestingly, microbial vitamin B6 biosynthesis and

salvage was the best predictor of chronic fatigue and irritable bowel syndromes (73, 74).

Despite the intrigue of connecting the human genome, microbiota and disease phenotypes,

evaluating these hypotheses will require more holistic approaches such as incorporating

metagenomics and metabolomics to identify whether enzymes or metabolic functions

reproducibly vary across ethnicities, as well as direct functional studies in model systems to

understand if correlation is truly driven by causation.

Further limitations should also be considered, including recruitment biases for the

AGP versus HMP, variation in sample processing and OTU clustering, and uneven sampling

which could only be addressed with down sampling of over-represented ethnicities. Still,

despite these confounders care was taken to demonstrate the reproducibility of results

across statistical methods, ecological metrics, rarefaction depths, and study populations.

Summarily, this work suggests that abundance variation of specific taxa, rather than whole

communities, may represent the most reliable ethnic signatures in the gut microbiota. A

reproducible co-occurring subset of these taxa link to a variety of overlapping metabolic

processes and health disparities, and contain the most heritable bacterial family,

Christensenellaceae. Moreover, a majority of the microbial taxa associated with ethnicity are

also heritable and genetically-associated lineages, suggesting there is a possible connection

between ethnicity and genetic patterns of biogeographical ancestry that may play a role in

shaping these taxa. Our results emphasize the importance of sampling ethnically diverse

populations of healthy individuals in order to discover and replicate ethnicity signatures in

the human gut microbiota, and they highlight a need to account for ethnic variation as a

potential confounding factor in studies linking microbiota variation to disease. Further

reinforcement of these results may lead to generalizations about microbiota assembly and

even consideration of specific taxa as potential mediators or treatments of health disparities.

Materials and Methods:

Data Acquisition

AGP data was obtained from the project FTP repository located at *ftp://ftp.microbio.me/AmericanGut/*. AGP data generation and processing prior to analysis can be found at: *https://github.com/biocore/American-Gut/tree/master/ipynb/primary-processing*. All analyses utilized the rounds-1-25 dataset which was released on March 4, 2016. Throughout all analyses, QIIME v1.9.0 was used in an Anaconda environment [*https://continuum.io*] for all script calls, custom scripts and notebooks were run in the QIIME 2 Anaconda environment with python version 3.5.2, and plots were post-processed using Inkscape [*https://inkscape.org/en/*] (75). Ethnicity used in this study was self-declared by AGP study participants as one of four groups: African American, Asian or Pacific Islander (Asian-Pacific Islander), Caucasian, or Hispanic. Sex was self-declared as either male, female, or other. Age was self-declared as a continuous integer of years old, and age categories defined by the AGP by decade (i.e. 20's, 30's…) were used in this study. BMI was self-declared as an integer, and BMI categories defined by AGP of underweight, healthy, overweight, and obese were utilized. Microbiota communities were characterized using 16S rDNA sequencing of variable region four and OTU clustering at 97% similarity, following an identical processing pipeline for all samples developed and optimized for the Earth Microbiome Project (76). HMP 16S rDNA data processed using QIIME for variable regions 3-5 was obtained from *http://hmpdacc.org/HMQCP/*. Demographic info for individual HMP participants was obtained through dbGaP restricted access to study phs000228.v2.p1, with dbGaP approval granted to SRB and non-human subjects determination IRB161231 granted by Vanderbilt University. Ethnicity and sex were assigned to subjects based on self-declared

values, with individuals selecting multiple ethnicities being removed unless they primarily

responded as Hispanic, while categorical age and BMI were established from continuous

values using the same criteria for assignment as in AGP. The HMP Amerindian population

was removed due to severe under-representation. This filtered HMP table was used for

community level analyses (ANOSIM, Alpha Diversity, beta intra-inter), however to allow

comparison with the AGP dataset, community subset analyses (co-occurrence, abundance

correlation, etc…) were performed with taxonomic assignments in QIIME using the UCLUST

method with the GreenGenes_13_5 reference.

Quality Control

AGP quality control was performed in Stata v12 (StataCorp, 2011) using available

metadata to remove samples (Raw N=9,475): with BMI more than 60 (-988 [8,487]) or less

than 10 (-68 [8,419]), missing age (-661 [7,758]), with age greater than 55 years old (-2,777

[4,981]) or less than 18 years old (-582 [4,399]), and blank samples or those not appearing

in the mapping file (-482 [3,917]), with unknown ethnicity or declared as other (-131

[3786]), not declared as a fecal origin (-2,002 [1784]), with unknown sex or declared as other

(-98 [1686]), or located outside of the United States (-209 [1477]). No HMP individuals were

missing key metadata or had other reasons for exclusion (-0[298]). Final community quality

control for both AGP and HMP was performed by filtering OTUs with less than 10 sequences

and removing samples with less than 1,000 sequences (AGP, -102 [1375]; HMP, -0 [298]). All

analyses used 97% OTUs generated by the AGP or HMP, and unless otherwise noted, results

represent Bray-Curtis beta diversity and Shannon alpha diversity at a rarefaction depth of

1,000 counts per sample.

ANOSIM, PERMANOVA, and BioEnv Distinguishability

The ANOSIM test was performed with 9,999 repetitions on each rarefied table within a respective rarefaction depth and beta diversity metric (**Fig 1 & S2A-B Table**), with R-values and p-values averaged across the rarefactions. Consensus beta diversity matrices were calculated as the average distances across the 100 rarefied matrices for each beta diversity metric and depth. Consensus distance matrices were randomly subsampled ten times for subset number of individuals from each ethnic group with more than that subset number prior to ANOSIM analysis with 9,999 repetitions, and the results were averaged evaluating the effects of more even representations for each ethnicity (**S2C Table**). Consensus distance matrices had each ethnicity and pair of ethnicities removed prior to ANOSIM analysis with 9,999 repetitions, evaluating the distinguishability conferred by inclusion of each ethnicity (**Fig 3A**, **S2F Table**). Significance was not corrected for the number of tests to allow comparisons between results of different analyses, metrics, and depths. PERMANOVA analyses were run using the R language implementation in the Vegan package (77), with data handled in a custom R script using the Phyloseq package (78). Categorical variables were used to evaluate the PERMANOVA equation (Beta-Diversity Distance Matrix ~ Ethnicity + Age + Sex + BMI) using 999 permutations to evaluate significance, and the R and p values were averaged across 10 rarefactions (**S2D Table**). The BioEnv test, or BEST test, was adapted to allow evaluation of the correlation and significance between beta diversity distance matrices and age, sex, BMI, and ethnicity simultaneously (**S2E Table**) (25). At each rarefaction depth and beta diversity metric the consensus distance matrix was evaluated for its correlation with the centered and scaled Euclidian distance

matrix of individuals continuous age and BMI, and categorical ethnicity and sex encoded

using patsy (same methodology as original test)[*https://patsy.readthedocs.io/en/latest/#*].

The test was adapted to calculate significance for a variable of interest by comparing how

often the degree of correlation with all metadata variables (age, sex, BMI, ethnicity) was

higher than the correlation when the variable of interest was randomly shuffled between

samples 1,000 times.

Alpha Diversity

Alpha diversity metrics (Shannon, Simpson, Equitability, Chao1, Observed OTUs)

were computed for each rarefied table (QIIME: alpha_diversity.py), and results were collated

and averaged for each sample across the tables (QIIME: collate_alpha.py). Pairwise

nonparametric t-tests using Monte Carlo permutations evaluated alpha diversity differences

between the ethnicities with Bonferroni correction for the number of comparisons (**Fig 2A**,

S3 Table, QIIME: compare_alpha_diversity.py). A Kruskal-Wallis test implemented in python

was used to detect significant differences across all ethnicities.

Beta Diversity

Each consensus beta diversity distance matrix had distances organized based on

whether they represented individuals of the same ethnic group, or were between individuals

of different ethnic groups. All values indicate that all pairwise distances between all

individuals were used (**Fig 2B**, **S4A-B Table**), mean values indicate that for each individual

their average distance to all individuals in the comparison group was used as a single point

to assess pseudo-inflation (**S4C-D Table**). A Kruskal-Wallis test was used to calculate

significant differences in intra-ethnic distances across all ethnicities. Pairwise Mann-

Whitney-U tests were calculated between each pair of intra-ethnic distance comparisons,

along with intra-versus-inter ethnic distance comparisons. Significance was Bonferroni

corrected within the number of intra-intra-ethnic and intra-inter-ethnic distance groups

compared, with violin plots of intra- and inter-ethnic beta diversity distances generated for

each comparison.

Random Forest

RF models were implemented using taxa summarized at genus level, which

performed better compared to RF models using OTUs as features, both in terms of

classification accuracy and computational time. We first rarefied OTU tables at sequence

depth of 10,000 (using R v3.3.3 package *vegan's* rrarefy() function) and then summarized

rarefied OTUs at genus-level (or lower characterized level if genus was uncharacterized for

an OTU). We filtered for rare taxa by removing taxa present in fewer than half of the number

of samples in rarest ethnicity (i.e. fewer than 10/2 = 5 samples in HMP and 13/2 = 6 (rounded

down) in AGP), retaining 85 distinct taxa in HMP dataset and 322 distinct taxa in AGP dataset

at genus level. The resulting taxa were normalized to relative abundance and arcsin-sqrt

transformed before being used as features for the RF models. We initially built multi-class

RF model, but since the RF model is highly sensitive to the uneven representation of classes,

all samples were identified as the majority class, i.e. Caucasian. In order to even out the class

imbalance, we considered some sampling approaches, but most existing techniques for

improving classification performance on imbalanced datasets are designed for binary class

imbalanced datasets, and are not effective on datasets with multiple underrepresented

classes. Hence, we adopted the binary classification approach and built four one-versus-all

(OVA) binary RF classifiers to classify samples from each ethnicity compared to the rest. 10-

fold cross-validation (using R package *caret* (79)) was performed using ROC as the metric

for selecting optimal model. The performance metrics and ROC curves were averaged across

the 10 folds (**Fig 3B**). Without any sampling during training the classifiers, most samples

were identified as the majority class, i.e. the Caucasian, by all four OVA RF classifiers. In order

to overcome this imbalance in class representation, we applied two sampling techniques

inside cross-validation: i) down-sampling, and ii) Synthetic Minority Over-sampling

Technique (or SMOTE) (29). In the down-sampling approach, the majority class is down-

sampled by random removal of instances from the majority class. In the SMOTE approach,

the majority class is down-sampled and synthetic samples from the minority class are

generated based on k-nearest neighbors technique (29). Note, the sampling was performed

inside cross-validation on training set, while the test was performed on unbalanced held-out

test set in each fold. The ROC curves and performance metrics table in **Fig 3B** show the

sensitivity-specificity tradeoff and classification performance for OVA classifier for each

ethnicity for both the sampling techniques applied on both the datasets. For both the

datasets, down-sampling shows higher sensitivity and lower specificity and precision for

minority classes (i.e. African Americans, Asian-Pacific Islanders and Hispanics) compared to

SMOTE. However, for the majority class (i.e. Caucasian), down-sampling lowers the

sensitivity and increases the specificity and precision compared to SMOTE. The sensitivity-

specificity tradeoff, denoted by the area under the ROC curve (or AUC) is reduced for

Hispanics in both the datasets.

Taxon Associations

Taxon differential abundance across categorical metadata groups was performed in

QIIME (QIIME: group_significance.py, **S5 Table**) to examine whether observation counts (i.e.

OTUs and microbial taxon) are significantly different between groups within a metadata

category (i.e. ethnicity, sex, BMI, age). The OTU table prior to final community quality control

was collapsed at each taxonomic level (i.e. Phylum – Genus; QIIME: collapse_taxonomy.py),

with counts representing the relative abundance of each microbial taxon. Differences in the

mean abundance of taxa between ethnicities were calculated using Kruskal-Wallis

nonparametric statistical tests. P-values are provided alongside false discovery rate and

Bonferroni corrected P-values, and taxon were ranked from most to least significant. Results

were collated into excel tables by taxonomic level and metadata category being examined,

with significant (false discovery rate and Bonferroni P-value < 0.05) highlighted in orange,

and taxa that were false discovery rate significant in both datasets were colored red. The

Fisher's exact test for the overlap of number of significant taxa between datasets was run at

the online portal (http://vassarstats.net/tab2x2.html), with the expected overlap calculated

as 5% of the number of significant lineages at all taxonomic level within the respective

dataset, and the observed 25 taxa that overlapped in our analysis. The permutation analysis

was performed by comparing the number of significant taxa (**S5 Table**, $p_{FDR}<0.05$)

overlapping between the AGP and HMP to the number overlapping when the Kruskal-Wallis

test was performed 1,000 times with ethnicity randomly permuted. In 1/1000 runs there

was one significant taxon overlapping at the family level, and one in 3/1000 permutations at

the genus level, with no significant taxa overlapping in any repetitions at higher taxonomic

levels. The 12 families and genera that were significantly different were evaluated to not be

'unique' if their abundances across ethnicities at each level represented at least 82-100%

(nearly all >95%) of the overlapping taxonomic level, and the genera was used if classified,

and family level used if genera was unclassified (g__). Average relative abundances on a log10

scale among individuals possessing the taxon were extracted for each taxon within each

ethnicity, and the abundance for 12 families and genera were made into barchart figures (**Fig**

4). The external whisker (AGP above, HMP below) depict the 75th quartile of abundance, and

the internal whisker depicts the 25th quartile. Pairwise Mann-Whitney-U tests were

performed between each pair of ethnicities using microbial abundances among all

individuals, and were Bonferroni corrected for the six comparisons within each taxon and

dataset. Bonferroni significant P-values are shown in the figure, and shown in bold if

significance and direction of change replicate in both datasets. Ubiquity shown above or

below each bar was calculated as the number of individuals in which that taxon was detected

within the respective ethnicity. Additional confirmation of ethnically varying abundance was

also performed at each taxonomic level (**S6 Table**), where the correlation of continuous age

and BMI along with categorically coded sex and ethnicity were simultaneously measured

against the log 10 transformed relative abundance of each taxon among individuals

possessing it using linear regression (**S6 Table** - Abundance), and against the presence or

absence of the taxon in all individuals with logistic regression (**S6 Table** - Presence Absence).

Significance is presented for the models each with ethnicity alone, and with all metadata

factors included (age, sex, BMI), alongside Bonferroni corrected p-values, and individual

effects of each metadata factor.

Co-Occurrence Analysis

Bacterial taxonomy was collapsed at the family level, Spearman correlation was

calculated between each pair of families using SciPy (80), and clustermaps were generated

using seaborn (**S2 Fig**), and ethnic associations were drawn from **S5 Table**. Correlations

were masked where Bonferroni corrected Spearman p-values were >0.05, and clusters were

identified as the most prominent (strongest correlations) and abundance enriched.

Enrichment of ethnic association was evaluated by measuring the Mann-Whitney-U of

cluster families ethnic associations (p-values, **S5 Table**) compared to the ethnic associations

of non-cluster taxa. Cluster associated families were identified as having at least three

significant correlations with families within the cluster.

Christensenellaceae Analysis

The abundance of the family Christensenellaceae was input as relative abundance

across all individuals from the family level taxonomic table. Individuals were subset based

on the presence/absence of Christensenellaceae and BMIs were compared using a one tailed

Mann-Whitney-U test, then each was further subset by ethnicity and BMI compared using

one tailed Mann-Whitney-U tests and boxplots within each ethnicity (**Fig 5**).

Genetically Associated, Heritable, and Correlated Taxa Analysis

Genetically associated taxa from population heritability studies (30, 31, 35, 36) with

a minimum heritability (A in ACE models or H2r) >0.1, and from GWAS studies (31, 36) were

examined for exact taxonomic overlap with our 12 ethnically-associated taxa. The 42 genetic

variants associated with Unclassified Clostridiales are: rs16845116, rs586749, rs7527642,

rs10221827, rs5754822, rs4968435, rs17170765, rs1760889, rs6933411, rs2830259,

rs7318523, rs17763551, rs2248020, rs1278911, rs185902, rs2505338, rs6999713,

rs5997791, rs7236263, rs10484857, rs9938742, rs1125819, rs4699323, rs641527,

rs7302174, rs2007084, rs2293702, rs9350764, rs2170226, rs2273623, rs9321334,

rs6542797, rs9397927, rs2269706, rs4717021, rs7499858, rs10148020, rs7524581,

rs11733214, rs7587067 from (31). These 40 variants along with variants in **Table 1** except

for chr7:96414393 (total=49) were then assessed in 1,000 Genomes individuals for

significant differentiation across superpopulations (34). The 1,000 Genomes VCF files were

downloaded (ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/), and variants

with a minor allele frequency less than 0.01 were removed with F_{ST} calculated between each

pair of superpopulations using vcftools (81). The East Asian versus South Asian F_{ST} rates

were not used in the analysis. A custom script was used to examine the F_{ST} for each of the 49

variants and compare to the F_{ST} of all variants on the same chromosome and all variants

genome-wide for that pair of populations, with percentile calculated and the number of

variants with a higher F_{ST} divided by the total number of variants. The eQTL value and

significance for rs7587067 were drawn from the GTEx database (67).

Data and Code Availability

Code, scripts, and data underlying figures are publicly available from the GitHub

repository   [https://github.com/awbrooks19/microbiota_and_ethnicity].   Individual

metadata (age, sex, ethnicity…) for the Human Microbiome Project are held under restricted

access available through dbGaP application [NCBI - dbGaP, Human Microbiome Project,

https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi?study_id=phs000228.v3.p1].

Acknowledgements

641   This work was supported by National Institutes of Health training grants 4T32GM08017810, 5T32GM08017809, and 5T32GM0817808 to AWB, the Vanderbilt Office of Equity, Diversity and Inclusion to A.W.B. and S.R.B., the Vanderbilt Microbiome Initiative to S.R.B., and the Alfred P. Sloan Foundation Fellowship to R.B.. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank the American Society of Microbiology for supporting travel to present this work. We would also like to thank Tony Capra, David Samuels, Patrick Abbot, Antonis Rokas, and other members of the Vanderbilt Genetics Institute and Bordenstein Lab for input. The authors acknowledge the Minnesota Supercomputing Institute (MSI) at the University of Minnesota and the Advanced Computing Center for Research and Education (ACCRE) at Vanderbilt University for providing resources that contributed to the research results reported within this paper.

Contributions

655   A.W.B., S.P., R.B., and S.R.B. conceived and designed the research. A.W.B. performed, analyzed, and interpreted all experiments with the exception of the Random Forest analysis planned and performed by S.P. and R.B. S.R.B. supervised all experimental designs, data analysis, and data interpretation. All authors participated in manuscript preparation, editing, and final approval.

Competing Financial Interests

662   The authors declare no competing financial interests.

Supplementary Table/Figure Legends:

S1 Fig. The average relative abundance of dominant microbial phyla for each ethnicity.

S2 Fig. Abundance correlation of microbial families. Spearman correlation clustermaps of bacterial abundance for families in the AGP and HMP. Numbers within boxes depict the spearman correlation value with heatmap coloration from blue negative correlation (-1), white no correlation (0), to red positive correlation (1). Positions have been masked based on Bonferroni significance <0.05 for the total clustermap of all microbial families. Lineages within boxes were identified as a highly correlated cluster, and lineages outside the boxes share multiple correlations with those within the cluster. Blue taxonomic names indicate overlap of taxa within boxes of both the AGP and HMP, while black indicate multiple correlations with the clusters in both datasets. The ethnic association column depicts FDR corrected p-values from Kruskal-Wallis tests in **S5 Table**, which are bolded if <0.05.

S3 Fig. Correlation of BMI with Christensenellaceae abundance. The relationship for each individual between log10 transformed Christensenellaceae abundance on the y axis and BMI on the x axis, with statistics slope, R^2, and p fit with a linear regression. Coloration of each point indicates ethnicity: Yellow – African American; Blue – Asian-Pacific Islander; Green – Hispanic; Red – Caucasian.

S1 Table. Demographic information for the AGP. Breakdown of age and BMI by sex and ethnicity. Heatmaps were constructed within each statistic and category (bounded by black box). The means for all sex and ethnic groups were used as the center (white), with higher

values indicated in red and lower in blue. HMP data is not shown because of data access

restrictions on participant metadata, available through dbGaP application.

S2 Table. Microbiota distinguishability by ethnicity, age, sex and BMI. (A) AGP and HMP

ANOSIM distinguishability by ethnicity, age, sex, and BMI at a rarefaction depth of 1,000 and

across four ecological metrics (more details in table). (B) AGP ANOSIM distinguishability by

ethnicity, age, sex, and BMI at rarefaction depths of 1,000 and 10,000. (C) ANOSIM results

for consensus distance matrix while subsampling the maximum number of individuals from

each ethnic group. (D) BioEnv results of correlation between ethnicity, age, sex, and BMI

together with outcome as multivariate beta diversity distance matrices [Distance Matrix =

Ethnicity*x1 + Categorical Age*x2 + Categorical BMI*x3 + Sex*x4 + B]. (E) ANOSIM results

for consensus distance matrix when each ethnicity and group of ethnicities are sequentially

removed from the analysis.

S3 Table. Alpha diversity by ethnicity, age, sex and BMI. Alpha Diversity for Ethnicity,

Age, Sex, and BMI across varying rarefaction depths and beta diversity metrics in AG (4A, 4C-

E), and for ethnicity in the HMP (4B). Results are based on non-parametric permutation

based t-tests, and p-values are Bonferroni corrected within each factor of interest, depth, and

metric.

S4 Table. Comparison of beta diversity distances for within and between ethnicities.

All values depicted are Mann-Whitney-U p-values. (A) All distances between pairs of

individuals within each ethnicity were compared between ethnicities across rarefaction

depths 1,000 and 10,000, four beta diversity metrics, and with while subsampling over-

represented ethnicities. (B) All distances between pairs of individuals within and between

each ethnicity were compared between ethnicities. (C) Mean distances between pairs of

individuals within each ethnicity were compared between ethnicities. (D) Mean distances

between pairs of individuals within and between each ethnicity were compared between

ethnicities.

S5 Table. Taxa which are differentially abundant by ethnicity, sex, BMI, and age in the

AGP and HMP. Kruskal-Wallis results for differential taxa abundance across metadata

groupings, including FDR and Bonferroni corrected p-values, and taxa abundance averages

within each group. Metadata factors and taxonomic levels are separated by excel tabs.

S6 Table. Taxa which are correlated with ethnicity, sex, BMI, and age in the AGP.

Results of linear (Abundance) and logistic (Presence Absence) regression results for

differential taxa abundance across metadata factors separated by taxonomic level. Columns

in order indicate the taxon name, the number of individuals with non-zero abundance; then

the p-value for ethnicity alone, the p-value Bonferroni corrected, the f-test statistic, and R^2;

then the same values for the regression with ethnicity, age, sex, and BMI together; then the

abundances in each ethnic group, and finally the p-values for each factor broken down.

S7 Table. Genetic variants with taxa associations and detailed 1,000 Genomes

population differentiation rates (F_{ST}). Variants in red indicate the variant has at least one

F_{ST} above the 95th percentile for high differentiation between at least one pair of populations.

Columns I-BU represent the values for calculating variant F_{ST} and percentiles. The first two

spaces indicate the two superpopulations being compared. F_{ST} indicates the rate of

differentiation for that variant between that pair of populations. Higher indicates the

number of variants genome-wide with a higher F_{ST}, and total indicates the total genome-wide

variants examined. The columns with chromosome indicate the number of variants with

higher F_{ST} and total variants on the same chromosome as the variant of interest. Percent

indicates the number of variants with a higher F_{ST} divided by the total number of variants.

References:

1.   Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, et al. A core
gut microbiome in obese and lean twins. Nature. 2009;457(7228):480-4.
2.   Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut
microbial gene catalogue established by metagenomic sequencing. Nature.
2010;464(7285):59-65.
3.   Huse SM, Ye Y, Zhou Y, Fodor AA. A core human microbiome as viewed through 16S
rRNA sequence clusters. PLoS One. 2012;7(6):e34242.
4.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, et al. Linking long-
term dietary patterns with gut microbial enterotypes. Science. 2011;334(6052):105-8.
5.   Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al. Diet
drives convergence in gut microbiome functions across mammalian phylogeny and within
humans. Science. 2011;332:970-4.
6.   Human Microbiome Project C. Structure, function and diversity of the healthy
human microbiome. Nature. 2012;486(7402):207-14.
7.   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet
rapidly and reproducibly alters the human gut microbiome. Nature. 2013;505(7484):559-
63.
8.   Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al.
Human gut microbiome viewed across age and geography. Nature. 2012;486(7402):222-7.
9.   Davenport ER, Cusanovich DA, Michelini K, Barreiro LB, Ober C, Gilad Y. Genome-
Wide Association Studies of the Human Gut Microbiota. PLoS One. 2015;10(11):e0140301.
10.   Fierera N, Hamadyc M, Lauberb CL, Knight R. The influence of sex, handedness, and
washing on the diversity of hand surface bacteria. Procedings of the National Academy of
Sciences.105(46).
11.   Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of
gut microbiota in type 2 diabetes. Nature. 2012;490(7418):55-60.

12.   Frank DN, Allison AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-
phylogenetic characterization of microbial community imbalances in human inflammatory
bowel diseases. Procedings of the National Academy of Sciences. 2007(104):13780–5.
13.   Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes associated with
obesity and IBD. FEBS Lett. 2014;588(22):4223-33.
14.   Zackular JP, Baxter NT, Iverson KD, Sadler WD, Petrosino JF, Chen GY, et al. The gut
microbiome modulates colon tumorigenesis. MBio. 2013;4(6):e00692-13.
15.   Mason MR, Nagaraja HN, Camerlengo T, Joshi V, Kumar PS. Deep sequencing
identifies ethnicity-specific bacterial signatures in the oral microbiome. PLoS One.
2013;8(10):e77287.
16.   Ravela J, Gajera P, Abdob ZG, Schneiderc M, Koeniga SSK, McCullea SL, et al. Vaginal
microbiome of reproductive-age women. Procedings of the National Academy of Sciences.
2011;108:4680-7.
17.   Fettweis JM, Brooks JP, Serrano MG, Sheth NU, Girerd PH, Edwards DJ, et al.
Differences in vaginal microbiome in African American women versus women of European
ancestry. Microbiology. 2014;160(Pt 10):2272-82.
18.   Williams DR, Priest N, Anderson NB. Understanding associations among race,
socioeconomic status, and health: Patterns and prospects. Health Psychol. 2016;35(4):407-
11.
19.   Mersha TB, Abebe T. Self-reported race/ethnicity in the age of genomic research: its
potential impact on understanding health disparities. Human Genomics. 2015;9(1):1.
20.   McDonald D, Birmingham A, Knight R. Context and the human microbiome.
Microbiome. 2015;3:52.
21.   Kolde R, Franzosa EA, Rahnavard G, Hall AB, Vlamakis H, Stevens C, et al. Host
genetic variation and its microbiome interactions within the Human Microbiome Project.
Genome Med. 2018;10(1):6.
22.   Rothschild D, Weissbrod O, Barkan E, Korem T, Zeevi D, Costea PI, et al.
Environmental factors dominate over host genetics in shaping human gut microbiota
composition. BioRxiv. 2017.
23.   DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes,
a chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl
Environ Microbiol. 2006;72(7):5069-72.
24.   Clarke KR. Non-parametric multivariate analyses of changes in community
structure. Australian Journal of Ecology. 1993;18:117-43.
25.   Clarke KR, Ainsworth M. A method of linking multivariate community structure to
environmental variables. Marine Ecology. 1993;92:205-19
.
26.   Knights D, Costello EK, Knight R. Supervised classification of human microbiota.
FEMS Microbiol Rev. 2011;35(2):343-59.
27.   Shannon CE. A mathematical theory of communication. Bell Syst Tech J.
1948;27:379-423.
28.   Hester ER, Barott KL, Nulton J, Vermeij MJ, Rohwer FL. Stable and sporadic
symbiotic communities of coral and algal holobionts. ISME J. 2016;10(5):1157-69.
29.   N.V. C, K.W. B, L.O. H, W.P. K. SMOTE: Synthetic Minority Over-sampling Technique.
Journal of Artificial Intelligence Research. 2002;16:321-57.

30.   Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, et al. Human
genetics shape the gut microbiome. Cell. 2014;159(4):789-99.
31.   Goodrich JK, Davenport ER, Beaumont M, Jackson MA, Knight R, Ober C, et al. Genetic
Determinants of the Gut Microbiome in UK Twins. Cell Host Microbe. 2016;19(5):731-43.
32.   Fu J, Bonder MJ, Cenit MC, Tigchelaar EF, Maatman A, Dekens JA, et al. The Gut
Microbiome Contributes to a Substantial Proportion of the Variation in Blood Lipids. Circ
Res. 2015;117(9):817-24.
33.   Pennisi E. Human Genetic Variation. Science. 2007;318(5858):1842-3.
34.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A
global reference for human genetic variation. Nature. 2015;526(7571):68-74.
35.   Lim MY, You HJ, Yoon HS, Kwon B, Lee JY, Lee S, et al. The effect of heritability and
host genetics on the gut microbiota and metabolic syndrome. Gut. 2016.
36.   Turpin W, Espin-Garcia O, Xu W, Silverberg MS, Kevans D, Smith MI, et al.
Association of host genome with intestinal microbial composition in a large healthy cohort.
Nature Genetics. 2016;48(11):1413-7.
37.   Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general
pedigrees. Am J Hum Genet. 1998;62(5):1198-211.
38.   Morgan XC, Tickle TL, Sokol H, Gevers D, Huttenhower C. Dysfunction of the
intestinal microbiome in inflammatory bowel disease and treatment. Genome Biology.
2012;7(979).
39.   Lewis JD, Chen EZ, Baldassano RN, Otley AR, Griffiths AM, Lee D, et al. Inflammation,
Antibiotics, and Diet as Environmental Stressors of the Gut Microbiome in Pediatric Crohn's
Disease. Cell Host Microbe. 2015;18(4):489-500.
40.   Goker M, Gronow S, Zeytun A, Nolan M, Lucas S, Lapidus A, et al. Complete genome
sequence of Odoribacter splanchnicus type strain (1651/6). Stand Genomic Sci.
2011;4(2):200-9.
41.   Castaneda G, Liu B, Torres S, Bhuket T, Wong RJ. Race/Ethnicity-Specific Disparities
in the Severity of Disease at Presentation in Adults with Ulcerative Colitis: A Cross-
Sectional Study. Dig Dis Sci. 2017.
42.   Boucias DG, Cai Y, Sun Y, Lietze VU, Sen R, Raychoudhury R, et al. The hindgut lumen
prokaryotic microbiota of the termite Reticulitermes flavipes and its responses to dietary
lignocellulose composition. Mol Ecol. 2013;22(7):1836-53.
43.   LATHAM MJ, WOLIN MJ. Fermentation of Cellulose by Ruminococcus flavefaciens in
the Presence and Absence ofMethanobacterium ruminantium. Appl Environ Microbiol.
1977;34(3):297-301.
44.   Falony G, Raes J. Population-level analysis of gut microbiome variation. Science.
2016;352(6285):560-4.
45.   Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, et al. Gut
Microbiota and Extreme Longevity. Curr Biol. 2016;26(11):1480-5.
46.   Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, et al. Age-
Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic Inflammation,
and Macrophage Dysfunction. Cell Host Microbe. 2017;21(4):455-66 e4.
47.   Xie W, Wood AR, Lyssenko V, al. e. Genetic Variants Associated With Glycine
Metabolism and Their Role in Insulin Sensitivity and Type 2 Diabetes. Diabetes. 2013;62.
48.   Williams SR, Yang Q, Chen F, Liu X, Keene KL, Jacques P, et al. Genome-wide meta-
analysis of homocysteine and methionine metabolism identifies five one carbon

metabolism loci and a novel association of ALDH1L1 with ischemic stroke. PLoS Genet.
2014;10(3):e1004214.
49. Petersen LM, Bautista EJ, Nguyen H, Hanson BM, Chen L, Lek SH, et al. Community
characteristics of the gut microbiomes of competitive cyclists. Microbiome. 2017;5(1):98.
50. Nakamura N, Lin HC, McSweeney CS, Mackie RI, Gaskins HR. Mechanisms of
microbial hydrogen disposal in the human colon and implications for health and disease.
Annu Rev Food Sci Technol. 2010;1:363-95.
51. Parthasarathy G, Chen J, Chen X, Chia N, O'Connor HM, Wolf PG, et al. Relationship
Between Microbiota of the Colonic Mucosa vs Feces and Symptoms, Colonic Transit, and
Methane Production in Female Patients With Chronic Constipation. Gastroenterology.
2016;150(2):367-79 e1.
52. Jackson CS, Oman M, Patel AM, Vega KJ. Health disparities in colorectal cancer
among racial and ethnic minorities in the United States. J Gastrointest Oncol. 2016;7:S32-
43.
53. Lopetuso LR, Scaldaferri F, Petito V, Gasbarrini A. Commensal Clostridia: leading
players in the maintenance of gut homeostasis. Gut Pathogens. 2013.
54. Sy DF. The Center for Asian Health Engages Communities in Research to Reduce
Asian American Health Disparities. US Department of Health & Human Services, National
Institute on Minority Health and Health Disparities.
55. Hwang H. Colorectal Cancer Screening among Asian Americans. Asian Pacific Journal
of Cancer Prevention. 2013;14(7):4025-32.
56. Oh KM, Kreps GL, Jun J. Colorectal Cancer Screening Knowledge, Beliefs, and
Practices of Korean Americans. American Journal of Health Behavior. 2013;37(3):381-94.
57. Sankaranarayanan R, Ramadas K, Qiao Y-l. Managing the changing burden of cancer
in Asia. BMC Medicine. 2014;12(3).
58. Pourhoseingholi MA. Increased burden of colorectal cancer in Asia. World J
Gastrointest Oncol. 2012;4(4):68-70.
59. Pourhoseingholi MA, Vahedi M, Baghestani AR. Burden of gastrointestinal cancer in
Asia; an overview. Gastroenterology and Hepatology. 2015.
60. Pourhoseingholi MA. Epidemiology and burden of colorectal cancer in Asia-Pacific
region: what shall we do now? Translational Gastrointestinal Cancer. 2014;3(4):169-73.
61. Report CHDaI. 2013.
62. Cao H, Liu X, An Y, Zhou G, Liu Y, Xu M, et al. Dysbiosis contributes to chronic
constipation development via regulation of serotonin transporter in the intestine. Sci Rep.
2017;7(1):10322.
63. Mosca A, Leclerc M, Hugot JP. Gut Microbiota Diversity and Human Diseases: Should
We Reintroduce Key Predators in Our Ecosystem? Front Microbiol. 2016;7:455.
64. Zhang X, Zhang D, Jia H, Feng Q, Wang D, Liang D, et al. The oral and gut
microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment.
Nat Med. 2015;21(8):895-905.
65. Singh VP, Proctor SD, Willing BP. Koch's postulates, microbial dysbiosis and
inflammatory bowel disease. Clin Microbiol Infect. 2016;22(7):594-9.
66. Sherry ST WM, Kholodov M, Baker J, Phan L, Smigielski EM, Sirotkin K. dbSNP: the
NCBI database of genetic variation. Nucleic Acids Research. 2001;29(308).
67. Consortium GT. The Genotype-Tissue Expression (GTEx) project. Nat Genet.
2013;45(6):580-5.

68.   Qiu X, Wei R, Li Y, Zhu Q, Xiong C, Chen Y, et al. NEDL2 regulates enteric nervous
system and kidney development in its Nedd8 ligase activity-dependent manner.
Oncotarget. 2016;7(21).
69.   Wei R, Qiu X, Wang S, Li Y, Wang Y, Lu K, et al. NEDL2 is an essential regulator of
enteric neural development and GDNF/Ret signaling. Cell Signal. 2015;27(3):578-86.
70.   O'Donnell AM, Coyle D, Puri P. Decreased expression of NEDL2 in Hirschsprung's
disease. J Pediatr Surg. 2016;51(11):1839-42.
71.   Roberts SJ, Stewart AJ, Schmid R, Blindauer CA, Bond SR, Sadler PJ, et al. Probing the
substrate specificities of human PHOSPHO1 and PHOSPHO2. Biochim Biophys Acta.
2005;1752(1):73-82.
72.   Ikeda M, Hosotani T, Kurimoto K, Mori T, Ueda T, Kotake Y, et al. The differences of
the metabolism related to vitamin B6-dependent enzymes among vitamin B6-deficient
germ-free and conventional rats. Nutritional Science Vitaminology. 1979;131(9).
73.   Nagy-Szakal D, Williams BL, Mishra N, Che X, Lee B, Bateman L, et al. Fecal
metagenomic profiles in subgroups of patients with myalgic encephalomyelitis/chronic
fatigue syndrome. Microbiome. 2017;5(1):44.
74.   Bhui KS, Dinos S, Ashby D, Nazroo J, Wessely S, White PD. Chronic fatigue syndrome
in an ethnically diverse population: the influence of psychosocial adversity and physical
inactivity. BMC Medicine. 2011;9(26).
75.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al.
QIIME allows analysis of high-throughput community sequencing data. Nat Method.
2010;7:335-6.
76.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.
ISME J. 2012;6(8):1621-4.
77.   Anderson MJ. A new method for non-parametric multivariate analysis of variance.
Australian Journal of Ecology. 2001;26:32-46.
78.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. PLoS One. 2013;8:e61217.
79.   Kuhn M. A short introduction to the caret package. 2017.
80.   Jones E, Oliphant T, Peterson P. Open Source Scientific Tools for Python. 2001.
81.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011;27(15):2156-8.

Taxonomic Lineage
(Blue: Overlap Within Cluster | Bold: Overlap Correlated with Cluster)

Ethnic Association
(Bold: FDR Significant p-value)

American Gut Project

Taxonomic Lineage	Ethnic Association
k__Bacteria; p__Proteobacteria; c__Deltaproteobacteria; o__Desulfovibrionales; f__Desulfovibrionaceae	**0.00067**
k__Bacteria; p__Firmicutes; c__Clostridiales; o__Clostridiales; f__Peptococcaceae	**0.00138**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Ruminococcaceae	**0.00085**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__[Mogibacteriaceae]	**0.00171**
k__Bacteria; p__Cyanobacteria; c__4C0d-2; o__YS2; f__	**3.38e-6**
k__Bacteria; p__Proteobacteria; c__Alphaproteobacteria; o__RF32; f__	**4.10e-6**
k__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__[Barnesiellaceae]	**0.00606**
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Rikenellaceae	**7.36e-6**
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__[Odoribacteraceae]	**0.00087**
k__Bacteria; p__Verrucomicrobia; c__Verrucomicrobiae; o__Verrucomicrobiales; f__Verrucomicrobiaceae	**3.93e-6**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__EtOH8	0.60348
k__Bacteria; p__Synergistetes; c__Synergistia; o__Synergistales; f__Synergistaceae	0.24561
k__Bacteria; p__Proteobacteria; c__Betaproteobacteria; o__Burkholderiales; f__Oxalobacteraceae	**0.01552**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__	**9.29e-9**
k__Bacteria; p__Tenericutes; c__Mollicutes; o__RF39; f__	**0.00281**
k__Archaea; p__Euryarchaeota; c__Methanobacteria; o__Methanobacteriales; f__Methanobacteriaceae	**0.00142**
k__Bacteria; p__Tenericutes; c__RF3; o__ML615J-28; f__	**0.00517**
k__Bacteria; p__; c__; o__SHA-98; f__	**0.00342**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Dehalobacteriaceae	**0.00336**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Christensenellaceae	**1.55e-9**
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Bacteroidaceae	0.65799
k__Bacteria; p__Proteobacteria; c__Betaproteobacteria; o__Burkholderiales; f__Alcaligenaceae	0.38689
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Porphyromonadaceae	**0.00517**
k__Bacteria; p__Proteobacteria; c__Gammaproteobacteria; o__Enterobacteriales; f__Enterobacteriaceae	0.24790
k__Bacteria; p__Actinobacteria; c__Actinobacteria; o__Bifidobacteriales; f__Bifidobacteriaceae	**2.11e-5**
k__Bacteria; p__Firmicutes; c__Bacilli; o__Lactobacillales; f__Streptococcaceae	0.64088
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Clostridiaceae	0.64362
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Lachnospiraceae	0.35217
k__Bacteria; p__Firmicutes; c__Erysipelotrichi; o__Erysipelotrichales; f__Erysipelotrichaceae	0.99041
k__Bacteria; p__Actinobacteria; c__Coriobacteria; o__Coriobacteriales; f__Coriobacteriaceae	0.70082
k__Bacteria; p__Verrucomicrobia; c__Opitutae; o__[Cerasicoccales]; f__[Cerasicoccaceae]	0.99041
k__Bacteria; p__Lentisphaerae; c__[Lentisphaeria]; o__Victivallales; f__Victivallaceae	**0.00606**
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__S24-7	0.38689
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Prevotellaceae	0.21918
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__[Paraprevotellaceae]	0.96330
k__Archaea; p__Euryarchaeota; c__Thermoplasmata; o__E2; f__[Methanomassiliicoccaceae]	0.54235

Human Microbiome Project

Spearman Correlation

Color scale: 1.0 / 0.8 / 0.6 / 0.4 / 0.2 / 0.0 / −0.2 / −0.4 / −0.6 / −0.8 / −1.0

Taxonomic Lineage	Ethnic Association
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Ruminococcaceae	0.18789
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__	**0.00463**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Christensenellaceae	**0.00193**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Dehalobacteriaceae	**0.02336**
k__Bacteria; p__Tenericutes; c__Mollicutes; o__RF39; f__	**0.00718**
k__Bacteria; p__Tenericutes; c__RF3; o__ML615J-28; f__	0.08836
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Prevotellaceae	**0.03608**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Peptococcaceae	**0.02300**
k__Bacteria; p__Firmicutes; c__Erysipelotrichi; o__Erysipelotrichales; f__Erysipelotrichaceae	0.23592
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Clostridiaceae	0.23592
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__Lachnospiraceae	0.31072
k__Bacteria; p__Verrucomicrobia; c__Verrucomicrobiae; o__Verrucomicrobiales; f__Verrucomicrobiaceae	**0.04723**
k__Bacteria; p__Firmicutes; c__Bacilli; o__Lactobacillales; f__Streptococcaceae	0.57953
k__Bacteria; p__Actinobacteria; c__Coriobacteriia; o__Coriobacteriales; f__Coriobacteriaceae	**0.00660**
k__Bacteria; p__Firmicutes; c__Clostridia; o__Clostridiales; f__[Mogibacteriaceae]	0.23592
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Bacteroidaceae	**0.00629**
k__Bacteria; p__Lentisphaerae; c__[Lentisphaeria]; o__Victivallales; f__Victivallaceae	**0.02558**
k__Bacteria; p__Proteobacteria; c__Betaproteobacteria; o__Burkholderiales; f__Oxalobacteraceae	0.18294
k__Bacteria; p__Firmicutes; c__Clostridia; o__SHA-98; f__	0.59964
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__S24-7	0.11834
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__Rikenellaceae	**0.00376**
k__Bacteria; p__Bacteroidetes; c__Bacteroidia; o__Bacteroidales; f__[Odoribacteraceae]	**0.00629**