AlleleAnalyzer: a tool for personalized and allele-specific sgRNA design

Kathleen C. Keough[1,2], Svetlana Lyalina[2,3], Michael P. Olvera[2], Sean Whalen[2], Bruce R. Conklin[2,4],

Katherine S. Pollard[2,5,6]

[1]*Pharmaceutical Sciences and Pharmacogenomics Graduate Program at the University of California, San*

Francisco, California, USA

[2]*Gladstone Institutes, San Francisco, California, USA*

[3]*Bioinformatics Graduate Program at the University of California, San Francisco, California, USA*

[4]*Departments of Biostatistics, Medicine, Ophthalmology and Pharmacology, University of California, San*

Francisco, California, USA

[5]*Department of Epidemiology & Biostatistics, Institute for Human Genetics, Quantitative Biology Institute,*

and Institute for Computational Health Sciences, University of California, San Francisco, California, USA

[6]*Chan Zuckerberg Biohub, California, USA*

Abstract

The CRISPR/Cas system is a highly specific genome editing tool capable of distinguishing alleles differing

by even a single base pair. However, current tools only design sgRNAs for a reference genome, not taking

into account individual variants which may generate, remove, or modify CRISPR/Cas sgRNA sites. This

may cause mismatches between designed sgRNAs and the individual genome they are intended to target,

leading to decreased experimental performance. Here we describe AlleleAnalyzer, a tool for designing

personalized and allele-specific sgRNAs for genome editing. We leverage >2,500 human genomes to

identify optimized pairs of sgRNAs that can be used for human therapeutic editing in large populations in

the future.

Keywords

CRISPR, sgRNA design, genomics, genome surgery, genome editing, computational biology

Background

The CRISPR/Cas genome-editing system is highly specific, with the ability to discriminate between similar

genomic sites, even alleles, based on a single nucleotide difference[1]. In order to target a genomic region

with the CRISPR system, a single-guide RNA (sgRNA) must be designed that is specific to the region of

interest. While current sgRNA design tools incorporate various data relating to predicted efficiency and

specificity such as epigenetic marks and chromatin accessibility[2–4], in the vast majority of cases, sgRNAs

are designed using reference genomes, such as the hg38 assembly for human or the GRCm38 assembly for

mouse. Since sgRNAs are often used on cell lines or organisms with many nucleotide differences from the

reference (e.g., on average 0.1% of a human genome[5]). Despite the finding that sgRNAs can sometimes

tolerate a single basepair mismatch, these mismatches frequently negatively impact sgRNA efficiency and

render imprecise the results of specificity prediction[2, 6, 7]. Furthermore, the use of CRISPR to research

areas such as haploinsufficiency, genomic imprinting, and dominant negative diseases require allele-

specific sgRNA design. To address these challenges, we developed AlleleAnalyzer, a software tool that

designs personalized and allele-specific sgRNAs for individual genomes, identifies pairs of sgRNAs to

generate excisions likely to block expression of a gene, and leverages patterns of shared variation from

>2,500 human genomes to design sgRNA pairs for that will have the greatest utility in a target population.

Results and Discussion

Incorporating genetic variation into sgRNA design enables personalized and allele-specific CRISPR

experiments. Personalized design involves accounting for variants that disrupt, generate or modify sgRNA

sites in a given genome. A genetic variant can impact sgRNA sites by being located in or near a protospacer

Fig. 1. *Analysis of allele specific sgRNA sites*

A) In a sample genome, tools designing sgRNAs for the reference genome are imperfect matches due to genetic variants, exemplified by guide 1. This tool designs personalized sgRNAs, as demonstrated by guides 2 and 3, which incorporate homozygous and avoid heterozygous variants. It also designs allele-specific sgRNAs based on incorporation of heterozygous variants, shown by guides 4-6. B) Most variants annotated by the 1000 Genomes Project (1KGP) are in or near a PAM site, demonstrating both a need as well as an opportunity for sgRNA personalization. C) Analysis of common variants, and variants in an individual cell line within the Brunello sgRNA library .

adjacent motif (PAM site), potentially generating or eliminating sgRNA sites in an individual in a

heterozygous or homozygous manner. Rather than being an impediment, these variants can be incorporated

into sgRNA design, yielding personalized or allele-specific sgRNAs, depending on variant zygosity (Figure

1a). Because Cas nucleases have different PAM sequences, a variant may impact an sgRNA site for one

Cas but not another. We analyzed 11 Cas types (Supplementary Table 1) and ~81 million genome-wide

variants annotated by the 1000 Genomes Project[8] (1KGP), finding that most variants impact sgRNA sites

for at least one Cas type, even when considering only variants in PAMs, which are putatively more allele-

specific[1] (Figure 1b). The likelihood that a variant impacts an sgRNA site differs across Cas nucleases

(range: 19-98%), is positively correlated with PAM frequency in the reference genome (Pearson rho=0.9,

p=0.04), and is negatively correlated with PAM size (Pearson rho=-0.9, p=0.05). In fact, 3.6% of sgRNAs

in the widely used Brunello genome-wide CRISPR screening sgRNA library[9] contain at least one

common genetic variant (AF > 5% in the 1KGP cohort), and 2.1% of these sgRNAs contain a variant in the

individual human genome of an induced pluripotent stem cell (iPSC) line WTC, commonly used for disease

modeling [10] (Figure 1c), impacting ~13% of protein-coding genes in both cases. Failing to account for

variants can reduce the efficacy of sgRNAs and also generate unexpected off-target effects. These results

emphasize the importance of designing sgRNAs using the personal genome of the patient or cell line where

they will be deployed, or at least accounting for both heterozygous and homozygous genetic variants when

interpreting results using generic sgRNA libraries.

Genetic variants are not just an impediment to sgRNA design; they can be leveraged to establish new

therapeutic and research possibilities. Questions that allele-specific editing could help address include

haploinsufficiency, imprinting, and allele-specific gene regulation, as well as discovery and correction of

heterozygous disease variants. One promising example is genome surgery to treat dominant negative

disease by excising only the disease causing copy of a gene, an approach which rescues healthy phenotypes

in cell and animal models of dominant negative diseases including Huntington's disease[11] and retinitis

pigmentosa[12, 13]. We assessed this strategy genome-wide by attempting to design a pair of allele-specific

sgRNAs for each human protein-coding gene that could generate a genomic excision and eliminate protein

production from just one allele. Given a Cas nuclease, an estimated maximum distance between the two

sgRNAs on the haplotype to be excised, and allele-specific sgRNA sites, it is possible to classify genes–or

other genomic elements, such as enhancers–as putatively targetable or not (Supplementary Figure 1). We

use the term putatively targetable when a pair of allele-specific sgRNAs exists but has not yet been tested,

because it will not always be possible to cut specifically at a site and coding exon excision will not always

stop expression[14]. If we choose a maximum distance of 10 kilobases (kb) between sgRNAs, require the

sgRNAs to be within the gene including introns, and consider 11 Cas varieties, the average individual from

1KGP is putatively targetable for allele-specific excision at 77% of protein-coding genes. This rate is evenly

distributed across chromosomes but varies by Cas nuclease and gene (Supplementary Figure 2). For genes

that are not putatively targetable, additional allele-specific sgRNA sites may be found by leveraging non-

coding variants up- and down-stream of the gene, or even in distal enhancers for the gene. Genome-wide,

we found that by simply including the 5 kb flanking regions of each gene, we can increase the expected

proportion of targetable protein-coding genes per individual from 77% to 85%. We conclude that allele-

specific excision is applicable to the vast majority of genes in most human genomes.

Since some genes in a given individual do not have a pair of allele-specific sgRNAs, we asked if gene

silencing with a single allele-specific sgRNA within the coding sequence (single-guide strategy) makes

more genes excisable. We compared paired-guide and single-guide strategies for allele-specific gene

knockout in the individual human genome of the WTC iPSC line [10] and found that more than twice as

Fig. 2: Targeting pairs of allele specific polymorphisms
A) 82 pairs of allele-specific sgRNA sites for SpCas9 are shared by at least 10% of 1KGP in the gene RHO including 1 kb flanking the gene. B) Development of a set cover optimization algorithm allows targeting of the largest population possible with the fewest allele-specific sgRNA pairs. C) Most targetable individuals have at least one of five SpCas9 sgRNA pairs optimized to be highly shared among the 1KGP cohort in the gene RHO plus the 1 kb regions flanking the gene. D) All possible allele-specific sgRNAs for SpCas9 in RHO plus the flanking 1 kb sequences were designed and scored for predicted specificity using CRISPOR. In the violinplot showing the score distribution for sgRNAs designed against the reference and alternate alleles per each heterozygous variant, the inner boxplot denotes the quartiles of each dataset.

many genes are putatively targetable with paired guides (Supplementary Figure 3), because one or both

sgRNAs can fall in introns or untranslated regions whereas single sgRNAs are limited to coding regions.

Genes that are putatively targetable with a single- and not paired-guide approach tend to have less than two

heterozygous variants in the gene, indicating lack of multiple variants as the primary reason a paired-guide

strategy fails. These genes likely could be putatively targetable with a paired-guide strategy by

incorporating flanking, promoter, or other regulatory regions. We therefore recommend paired-guides for

allele-specific gene excision.

Genome editing sgRNAs do not need to be designed one genome at a time. Variants that impact sgRNA

sites are often shared among large proportions of the individuals within and sometimes between populations

due to haplotype structure. Allele sharing varies by population and locus, as individuals with common

ancestry will share haplotypes that harbor specific sets of variants. We therefore developed an algorithm to

identify allele-specific sgRNA guide pairs for a given gene that cover the maximum number of individuals

in a population; these have the broadest therapeutic potential, similar to designing a drug to treat as many

people as possible. Specifically, our method seeks to cover the most people with the fewest sgRNA pairs

using their shared heterozygous variants; this is similar to the set cover problem in that the algorithm

identifies an optimal combination rather than simply selecting most shared sgRNA pairs, which could

disproportionately favor one group over another [15]. Our algorithm generates optimized pairs of sgRNAs

that can be used to study or treat genetic diseases in large groups, potentially eliminating the need to develop

new sgRNA pairs for each patient or cell line, with practical implications for the development of genome

surgery as a field. Our algorithm can also be used to identify sgRNA pair combinations applicable to a

custom cohort, enabling researchers to design guides that are maximally shared among multiple cell lines,

for example, which would improve experimental efficiency.

As a case study, we investigated the feasibility of excising one allele of exon 1 of *RHO,* which can cause

dominant negative macular dystrophy[13]. Considering the gene plus 1 kb of flanking sequence on either

side, there are 82 pairs of allele-specific sgRNA sites for SpCas9 that are shared by >10% of all 1KGP

individuals, with the number and composition of these pairs varying across 1KGP populations (Figure 2a,

Supplementary Figure 4). We sought to identify an optimal combination of five allele-specific sgRNA pairs

to target the majority of the 1KGP cohort (Figure 2b). We found that five allele-specific sgRNA pairs could

putatively excise one allele of *RHO* while leaving the other allele intact in ~88% of 1KGP individuals with

at least two variants, or 57% of the overall 1KGP population (Figure 2c). We also demonstrated how

avoiding heterozygous variants and incorporating homozygous variants enables personalized sgRNA

design in the *RHO* locus for the WTC genome for many Cas varieties, including SpCas9, SaCas9 and cpf1

(Cas12a) (Supplementary Figure 5, Supplementary Tables 2 and 3). The dominant negative disease gene

RHO clearly demonstrates the power of using genetic variation in sgRNA design.

We incorporated these methods into AlleleAnalyzer, an open-source software tool (Supplementary Figure

6). This tool designs personalized and allele-specific sgRNAs for unique individuals and cohorts, given

their genetic variants, and optimizes sgRNA pairs to cover many individuals based on shared variants. To

our knowledge, this is the first computational resource that designs personalized and allele-specific CRISPR

sgRNAs, thus expanding and building upon the existing repertoire of sgRNA design tools (Supplementary

Table 4). We integrated the specificity scoring capabilities of CRISPOR[4] to enable users to stratify guides

by that metric as desired (Figure 2d). The AlleleAnalyzer toolkit and tutorials are available along with the

database   of   annotated   1KGP   variants   (Supplementary   Table   5)   at

https://github.com/keoughkath/AlleleAnalyzer.

Conclusions

The genetic variation aware sgRNA design tool AlleleAnalyzer is an important step towards effective

deployment of CRISPR-based technologies in diverse genomes, including but not limited to research and

therapeutic development for once incurable dominant negative diseases.

Methods

PAM occurrence in the human reference genome

PAM frequency

The AlleleAnalyzer tool includes a script enabling scanning of a reference genome fasta file for existing

PAM sites. We used this to identify PAM sites for 11 Cas types (Supplementary Table 1) in the reference

human genomes hg19 and hg38.

PAM size

PAM sizes were equated as the sum of non-N (A, C, G or T) bases in a PAM site. Thus "NGG" for SpCas9

would have size 2, and "NNGRRT" for SaCas9 would have size 4.

AlleleAnalyzer analysis of the 1000 Genomes cohort

Annotation of variants

Genetic variants were determined to generate or destroy an allele-specific sgRNA site if they were proximal

to or in a PAM site (Figure 1a). Sufficient proximity to a PAM site was defined for this study as 20 base

pairs based on the common length of sgRNA recognition sequences. For all Cas varieties this was the 20

base pairs 5' of the PAM, except for cpf1 (Cas12a) for which it was 3' of the PAM. The sgRNA design

tools that are part of AlleleAnalyzer allow different user-defined sgRNA lengths and addition of Cas

enzymes and PAMs. There is evidence to suggest that genetic variants that generate or destroy a PAM are

more likely to lead to allele-specific Cas activity compared to those in the seed sequence[1]; AlleleAnalyzer

thus provides options to differentiate between CRISPR sites in a PAM site versus the sgRNA recognition

sequence. All variants genome-wide were annotated for the 1KGP cohort for reference genomes hg19 nd

hg38 and are available for querying; an example subset of these data for the first 100 variants annotated by

1KGP on chromosome 1 in reference genome hg19 is available in Supplementary Table 5.

Generation of gene set

The gene set analyzed was compiled using the canonical transcripts for RefSeq gene annotations for human reference genome hg19 and hg38 downloaded using the UCSC table browser[16], and filtered for genes with at least one coding exon. When non-protein-coding genes were excluded, 15,199 genes were evaluated for hg19, and 16,143 for hg38. Values reported in the text are for hg19 unless stated otherwise, but analyses were conducted for both reference genomes with similar results.

Allele-specific putative gene targetability genome-wide

Putative allele-specific targetability of a gene is defined here as whether a gene contains a pair of allele-specific sgRNA sites for at least one of the 11 Cas enzymes evaluated that are less than 10 kb apart on the same haplotype in an individual that will disrupt a coding exon (Supplementary Figure 1). This metric was calculated for each protein-coding gene for all 2,504 1KGP individuals.

Set cover analysis

In order to determine optimal pairs of sgRNAs to cover large groups of people in a particular gene, we applied set cover optimization which we implemented using the Python package PuLP[17]. The aim was to maximize the number of individuals from the 1KGP for whom a user-supplied maximum number of sgRNA pairs would putatively target a given gene. This script can also be used to determine a minimum percentage of people to be covered by a set of sgRNA pairs.

WTC sequencing

The genome for the iPSC line WTC[10] was sequenced by the Allen Cell Science Institute. Analysis and variant calls in the reference genome hg19 were done according to GATK version 3.7 best practices[18] and phased using Beagle version 4.1 with default settings[19].

WTC targetability analysis

Variant annotation procedures were the same as in the 1KGP analysis. The same genes lists used in the

1KGP analysis were analyzed in WTC, except when specified in the text, for the cases of 1 kb flanking the

gene *RHO*, or when analyzing targetability for all genes + 5 kb flanking vs. genic region only.

Packages used

Python

Docopt was used for handling of command-line arguments. Pandas[20] version 0.21.0 and NumPy[21]

version 1.13.3 and elements of the standard Python distribution sys, os, and regex were used for multiple

aspects of data analysis. PuLP[17] version 1.6.8 was used for set cover analysis. PyTables[22] was used for

data management. Biopython[23] and pyfaidx[24] were used for Fasta processing. Scripts from

CRISPOR[4] were integrated into AlleleAnalyzer to facilitate specificity scoring of sgRNAs.

R

Packages used to generate arcplots included viridis version 0.5.1, viridisLite version 0.3.0, igraph version

1.1.2, ggraph version 1.0.0, ggplot2 version 2.2.1, reshape2 version 1.4.3, dplyr version 0.7.4, tidyr version

0.7.2, and readr version 1.1.1.

Bioinformatics

Bcftools versions 1.5 and 1.6 were used to manipulate VCF and BCF files.

Scripts

Scripts were written in Python version 3.6.1, R version 3.3.2 and Bash version 3.2.57.

Data Availability

1KGP phase 3 data were downloaded from the 1KGP website (http://www.internationalgenome.org/). The

reference hg19 and hg38 genome data were downloaded from the UCSC genome browser. The 1KGP

analysis dataset has been made available for public access online at

(http://lighthouse.ucsf.edu/public_files_no_password/excisionFinderData_public/1kgp_dat/).

Code Availability

All data processing and analysis scripts as well as the sgRNA design tool are located at

github.com/keoughkath/AlleleAnalyzer.

List of Abbreviations

sgRNA: single-guide RNA

PAM site: protospacer adjacent motif site

1KGP: 1000 Genomes Project

kb: kilobases (1000 genomic basepairs)

iPSC: induced pluripotent stem cell

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of Data and Material

All data processing and analysis scripts as well as the sgRNA design tool are located at

github.com/keoughkath/AlleleAnalyzer. 1KGP phase 3 data were downloaded from the 1KGP website

(http://www.internationalgenome.org/). The reference hg19 and hg38 genome data were downloaded from

the UCSC genome browser. The 1KGP analysis dataset has been made available for public access online

at   (http://lighthouse.ucsf.edu/public_files_no_password/excisionFinderData_public/1kgp_dat/).   WTC

whole-genome   sequencing   data   is   made   available   by   the   Allen   Institute   at

(https://www.allencell.org/genomics.html).

Competing Interests

B.R.C. is a founder of Tenaya Therapeutics, a company focused on finding treatments for heart failure,

including the use of CRISPR interference to interrogate genetic cardiomyopathies. B.R.C. and K.S.P. hold

equity in Tenaya, and Tenaya provides research support for heart failure related research to B.RC and K.S.P.

Funding

B.R.C and K.S.P. were supported by the Gladstone Institutes. K.S.P. received support from the National

Institutes of Health (UM1HL098179, P01HL089707, R01MH109907). B.R.C received support from the

National Institutes of Health (U01HL100406, P01HL089707, R01HL130533, 1R01EY028249). K.C.K

received support from the Claire Giannini Fund and the UCSF Discovery Fellows Program.

Authors' Contributions

K.C.K, S.W., B.R.C., and K.S.P. conceived the project. K.C.K, S.W., S.L., B.R.C., and K.S.P. designed the

experiments, K.C.K, S.L., M.P.O., B.R.C., and K.S.P. analyzed data. K.C.K, B.R.C., and K.S.P. wrote the

paper with editing from all other authors.

Acknowledgements

We are grateful for valuable scientific input from Anthony Moore, and members of the Pollard and Conklin

labs. We thank Anders Riutta from the Gladstone Bioinformatics Core for his Github troubleshooting

expertise. We are grateful to Maximilian Haeussler for his assistance and feedback with the integration of

CRISPOR.

References

1. Kathleen A. Christie, David G. Courtney, Larry A. DeDionisio CCS, Shyamasree De Majumdar, Laura

C. Mairs MAN& CBTM. Towards personalised allele-specific CRISPR gene editing to treat autosomal

dominant disorders. Sci Rep. 2017; November:1–11. doi:10.1038/s41598-017-16279-4.

2. Doench JG, Hartenian E, Graham DB, Tothova Z, Hegde M, Smith I, et al. Rational design of highly

active sgRNAs for CRISPR-Cas9–mediated gene inactivation. Nat Biotechnol. 2014;32:1262–7.

doi:10.1038/nbt.3026.

3. Horlbeck MA, Gilbert LA, Villalta JE. Compact and highly active next-generation libraries for

CRISPR-mediated gene repression and activation. 2016;9:1–20.

4. Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, et al. Evaluation of off-target

and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. Genome

Biol. 2016;17:148.

5. National Institutes of Health. NIH Curriculum Supplement Series. Biological Sciences Curriculum

Study. 2007. doi:10.1371/journal.pone.0075601.

6. Scott DA, Zhang F. Implications of human genetic variation in CRISPR-based therapeutic genome

editing. Nat Med. 2017;23:1095–101.

7. Lessard S, Francioli L, Alfoldi J, Tardif J-C, Ellinor PT, MacArthur DG, et al. Human genetic variation

alters CRISPR-Cas9 on- and off-targeting specificity at therapeutically implicated loci. Proc Natl Acad

Sci U S A. 2017;114:E11257–66. doi:10.1073/pnas.1714640114.

8. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A

global reference for human genetic variation. Nature. 2015;526:68–74.

9. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, et al. Optimized sgRNA

design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol. 2016;34

November 2015:1–12. doi:10.1038/nbt.3437.

10. Drubin DG, Hyman AA. Stem cells: the new "model organism." Mol Biol Cell. 2017;28:1409–11.

doi:10.1091/mbc.E17-03-0183.

11. Shin JW, Kim K-H, Chao MJ, Atwal RS, Gillis T, MacDonald ME, et al. Permanent inactivation of

Huntington's disease mutation by personalized allele-specific CRISPR/Cas9. Hum Mol Genet.

2016;25:4566–76.

12. Gao X, Tao Y, Lamas V, Huang M, Yeh W-H, Pan B, et al. Treatment of autosomal dominant hearing

loss by in vivo delivery of genome editing agents. Nature. 2018;553:217–21.

13. Bakondi B, Lv W, Lu B, Jones MK, Tsai Y, Kim KJ, et al. In Vivo CRISPR/Cas9 Gene Editing

Corrects Retinal Dystrophy in the S334ter-3 Rat Model of Autosomal Dominant Retinitis Pigmentosa.

Mol Ther. 2015;24 September:556–63. doi:10.1038/mt.2015.220.

14. Chen X, Xu F, Zhu C, Ji J, Zhou X, Feng X, et al. Dual sgRNA-directed gene knockout using

CRISPR/Cas9 technology in Caenorhabditis elegans. Sci Rep. 2014;4:7581.

15. Clarkson KL. Algorithms for polytope covering and approximation. In: Lecture Notes in Computer

Science. 1993. p. 246–52.

16. Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, et al. The UCSC Table

Browser data retrieval tool. Nucleic Acids Res. 2004;32 Database issue:D493-6.

17. Mitchell S, OSullivan M, others. PuLP: a linear programming toolkit for python. Univ Auckl. 2011.

18. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et al. From

fastQ data to high-confidence variant calls: The genome analysis toolkit best practices pipeline. 2013.

19. Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for

whole-genome association studies by use of localized haplotype clustering. Am J Hum Genet.

2007;81:1084–97.

20. McKinney W. Data Structures for Statistical Computing in Python. In: van der Walt S, Millman J,

editors. Proceedings of the 9th Python in Science Conference. 2010. p. 51–6.

21. Stéfan van der Walt SCC and GV. The NumPy Array: A Structure for Efficient Numerical

Computation. Comput Sci Eng. 2011;13:22–30.

22. Francesc Alted IV and others. PyTables: Hierarchical Datasets in Python.

23. Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: Freely available

Python tools for computational molecular biology and bioinformatics. Bioinformatics. 2009;25:1422–3.

24. Shirley MD, Ma Z, Pedersen BS, Wheelan SJ. Efficient "pythonic" access to FASTA files using

pyfaidx. PeerJ Inc.; 2015.

Supplementary Figure 1

A pair of allele-specific sgRNA sites is defined at putatively targetable if their predicted excision will

disrupt at least one protein-coding exon.

Supplementary Figure 2

This faceted density plot shows the percentage of putatively targetable 1KGP individuals (2,504 total individuals) per

protein-coding gene for 11 types of Cas nuclease.

Supplementary Figure 3

Many more genes are targetable in the genome of WTC with a paired (dual)- as opposed to single-guide strategy. The number of variants in a gene is influential in determining targetability. Many genes that are not dual- or single-guide targetable have very few variants, and the genes that are only targetable with a single-guide approach compared to a dual-guide approach also tend to have fewer variants. All 11 Cas varieties are considered in this analysis.

Rhodopsin + 1 kb flanking

356 **Supplementary Figure 4**

357 Shared pairs of sgRNAs per locus vary by population. We show allele-specific sgRNA site pairs shared by

358 at least 10% of each population for SpCas9 in the gene *RHO* plus the 1 kb flanking regions in the five super-

359 populations in the 1KGP as well as the overall 1KGP cohort.

361   *Nucleases evaluated: SpCas9, SaCas9, cpf1 (Cas12a)*

Supplementary Figure 5

Integrative Genomics Viewer track view of allele-specific (lavender, upper track) and personalized (multi-colored, middle track) sgRNAs for SpCas9, SaCas9 and cpf1 (Cas12a) in the gene *RHO* plus 1 kb flanking in WTC. Allele-specific guides are shaded according to position of the variant in the guide, with variants closer to the PAM being darker based on their putative greater specificity. The track labeled "WTC genetic variants" (top) denotes genetic variants in WTC in this locus, of which there are only heterozygous variants. The bottom track shows the RefSeq annotation for the first exon of this gene in hg38.

372 **Supplementary Figure 6**

373 Flowchart for the AlleleAnalyzer software tool.

Common name(s)	Abbreviation	PAM	Properties
SpCas9	SpCas9	NGG	*Streptococcus pyogenes* (Sp) Cas9., most widely used version with dozens of variants using same PAM, e.g. eSpCas9, SpCas9-HF1, eSpCas9 1.1 and more (Jinek et al. 2012)
SpCas9 VRER Variant	SpCas9-V1	NGCG	Version of SpCas9 with alternative targeting range (Kleinstiver et al. 2015)
SpCas9 EQR Variant	SpCas9-V2	NGAG	Version of SpCas9 with alternative targeting range (Kleinstiver et al. 2015)
SpCas9 VQR Variant	SpCas9-V3	NGAN or NGNG	Version of SpCas9 with wider targeting range (Kleinstiver et al. 2015)
SaCas9	SaCas9	NNGRRT	*Staphylococcus aureus* (Sa) Cas9. Small relative to SpCas9, (Horvath et al. 2008, Jiang et al. 2013)
SaCas9 KKH Variant	SaCas9-V1	NNNRT	Version of SaCas9 with 2 to 4-fold increased targeting range relative of SaCas9 (Kleinstiver et al. 2015)
nmCas9	nmCas9	NNNNGATT	*Neisseria meningitidis* (Nm) Cas9, with different PAM site (Hou et al. 2013)
cpf1	cpf1	TTTN	Multiple variations, notably opposite orientation system and sticky-end cut rather than blunt. Multiple species exist, including from *Acidaminucoccus* and *Lachnospiraceae.* (Zetsche et al. 2015)
StCas9 1	StCas9-V1	NNAGAA	*Streptococcus thermophilus* (St) Cas9. Smaller relative of SpCas9. Increased specificity. (Kleinstiver et al. 2015, Muller et al. 2016)
StCas9 2	StCas9-V2	NGGNG	*Streptococcus thermophilus* (St) Cas9. Smaller relative of SpCas9. Increased specificity. (Muller et al 2016)
cjCas9	cjCas9	NNNNACA	*Campylobacter jejuni* Cas9. Smallest Cas9 ortholog to date, easy to package (Kim et al. 2017)

Supplementary Table 1

11 types of Cas enzyme were evaluated, each of which has a distinct PAM site.

Supplementary Table 2

All possible allele-specific sgRNAs for SpCas9, SaCas9 and cpf1 (Cas12a) in the region surrounding the

first exon of *RHO* WTC (Supplementary Figure 6).

Supplementary Table 3

All possible personalized sgRNAs for SpCas9, SaCas9 and cpf1 (Cas12a) in the region surrounding the

first exon of *RHO* WTC (Supplementary Figure 6). WTC has no homozygous variants in this region, thus

allele frequency and variant-related columns are blank. However, the sgRNAs are designed to avoid the 9

heterozygous variants that WTC has in this region.

	Personalized sgRNA design	Allele-specific sgRNA design	Supports any genome	Paired sgRNA design	Optimizes pairs of sgRNAs to maximize coverage in a cohort	Single or multi-locus design	Multiple Cas varieties	Ability to add novel Cas enzymes	Off-target scoring
AlleleAnalyzer	yes	yes	yes	yes	yes	yes	yes	yes	yes
CRISPOR (Haeussler et al. 2016)	no	no	yes	no	no	yes	yes	no	yes
GuideScan (Perez et al. 2017)	no	no	no	yes	no	yes	yes	no	yes
E-CRISP (Heigwer et al. 2014)	no	no	no	yes	no	yes	no	no	yes
MIT design tool (Hsu et al. 2013)	no	no	no	no	no	yes	no	no	yes
CRISPRscan (Moreno-Mateos et al. 2015)	no	no	no	no	no	no	yes	no	yes
FlashFry (McKenna & Shendure, 2018)	no	no	yes	no	no	yes	yes	no	yes

409   **Supplementary Table 4**

410   Comparison of AlleleAnalyzer features with other commonly used CRISPR sgRNA design tools.

412   **Supplementary Table 5**

413   An example subset of variant annotations for the first 100 variants on chromosome 1 from 1KGP.