Title

Choanoflagellate transfection illuminates their cell biology and the ancestry of animal septins

Short Title

Robust transfection in choanoflagellates

Authors

David S. Booth[1], Heather Szmidt-Middleton, and Nicole King[2]*

Howard Hughes Medical Institute and Department of Molecular and Cell Biology

University of California, Berkeley

Berkeley, CA 94720

[1] https://orcid.org/0000-0002-4724-4702

[2] https://orcid.org/0000-0002-6409-1111

*Corresponding author

Email: nking@berkeley.edu

Phone: (510) 643-9395

ABSTRACT

As the closest living relatives of animals, choanoflagellates offer unique insights into animal

origins and core mechanisms underlying animal cell biology. However, unlike traditional model

organisms, such as yeast, flies and worms, choanoflagellates have been refractory to DNA

delivery methods for expressing foreign genes. Here we report the establishment of a robust

method for expressing transgenes in the choanoflagellate *Salpingoeca rosetta*, overcoming

barriers that have previously hampered DNA delivery and expression. To demonstrate how this

method accelerates the study of *S. rosetta* cell biology, we engineered a panel of fluorescent

protein markers that illuminate key features of choanoflagellate cells. We then investigated the

localization of choanoflagellate septins, a family of GTP-binding cytoskeletal proteins that are

hypothesized to regulate the multicellular rosette development in *S. rosetta.* Fluorescently

tagged septins localized to the basal pole of *S. rosetta* single cells and rosettes in a pattern

resembling septin localization in animal epithelia. The establishment of transfection in *S. rosetta*

and its application to the study of septins represent critical advances in the growth of *S. rosetta*

as an experimental model for investigating choanoflagellate cell biology, core mechanisms

underlying animal cell biology, and the origin of animals.

INTRODUCTION

First described in the mid-19th century, choanoflagellates inspired great debate regarding animal taxonomy (James-Clark, 1868; Kent, 1871; Leadbeater, 2015). The most diagnostic morphological feature of choanoflagellates, a "collar complex" composed of a single apical flagellum surrounded by a collar of actin-filled microvilli (Fig. 1), was interpreted as evidence of a special relationship between choanoflagellates and sponges, whose choanocytes (or "collar cells") each bear a collar complex. Subsequent phylogenetic analyses and the discovery of cells with a collar complex in nearly all animal phyla have revealed that sponges and all other animals are monophyletic, with choanoflagellates as their closest living relatives (Fig. 1) (Lang *et al.*, 2002; King *et al.*, 2008; Ruiz-Trillo *et al.*, 2008; Burger *et al.*, 2003; Fairclough *et al.*, 2013; Brunet and King, 2017). Moreover, comparative genomic analyses have revealed that choanoflagellates, animals, and other holozoans express genes required for animal multicellularity and embryogenesis (Richter *et al.*, 2018; Sebe-Pedros *et al.*, 2017), including cadherins (Abedin and King, 2008), tyrosine kinases (Suga *et al.*, 2012; Manning *et al.*, 2008), and Myc (Young *et al.*, 2011). Thus, comparisons among animals and choanoflagellates have the potential to provide unique insights into animal origins and core features of animal cell biology that are not conserved in other experimental models, such as yeast (King, 2004).

The choanoflagellate *Salpingoeca rosetta* (previously named *Proterospongia* sp. (King *et al.*, 2003)) has recently emerged as an experimentally tractable model. *S. rosetta* develops from a single founding cell into a spherical, multicellular "rosette" (Fig. 1C) through serial rounds of cell division in a process that evokes the earliest stages of animal embryogenesis (Fairclough *et al.*, 2010). Since the establishment of the first *S. rosetta* cultures almost twenty years ago, *S. rosetta* has become increasingly amenable to cell and molecular biological approaches due to the sequencing of its genome (Fairclough *et al.*, 2013), the establishment of forward genetic screens (Levin and King, 2013; Levin *et al.*, 2014), the ability to experimentally control key events in its life history (Dayel *et al.*, 2011), and the discovery that environmental bacteria induce multicellular rosette development and mating (Alegado *et al.*, 2012; Dayel *et al.*, 2011; Woznica *et al.*, 2016; Woznica *et al.*, 2017).

An important remaining barrier to the study of molecular and cellular mechanisms in *S. rosetta* has been the inability to perform transfection and transgene expression. Furthermore, the absence of the RNA interference pathway in *S. rosetta* has precluded gene knockdowns (Richter *et al.*, 2018; Fairclough *et al.*, 2013). Here we report the establishment of a robust

nucleofection-based method to transfect and express transgenes in *S. rosetta.* By engineering

plasmids with *S. rosetta* regulatory sequences driving the expression of fluorescently-tagged *S.*

rosetta proteins, we have developed a broad panel of markers for the study of choanoflagellate

cell biology *in vivo*. As a first application, we used transgene expression to characterize septins,

genes with conserved roles in fungal (Helfer and Gladfelter, 2006; Berepiki and Read, 2013)

and animal development (Fairclough *et al.*, 2013; Adam *et al.*, 2000; Neufeld and Rubin, 1994;

O'Neill and Clark, 2013; Kim, S. K. *et al.*, 2010), that have been hypothesized to regulate rosette

development (Fairclough *et al.*, 2013). By imaging fluorescently-tagged septins in live cells, we

show that their localization in *S. rosetta* resembles that in animal epithelia, providing a potential

evolutionary link between the mechanisms underlying animal and choanoflagellate

multicellularity.

RESULTS

A robust method to transfect *S. rosetta*

To detect successful transfection, we started by engineering four different DNA plasmid

constructs, each with different *S. rosetta* regulatory sequences fused to a gene, *nanoluc* (Hall *et*

al., 2012), encoding a highly sensitive luciferase (Fig. S1B). Because no choanoflagellate

promoters had previously been mapped, we increased the likelihood of cloning sequences that

would drive robust gene expression by fusing *nanoluc* to non-coding sequences flanking a set of

genes – *elongation factor L (efl)*, *α-tubulin (tub)*, *non-muscle actin (act)*, and *histone H3 (H3)* –

that each exhibit high expression, lack introns in their open reading frames and have well-

annotated 5'- and 3'-untranslated regions (Fig. S1A) (Fairclough *et al.*, 2013).

Next, we set out to deliver these DNA plasmid constructs into *S. rosetta* cells using

nucleofection, an electroporation-based technique that has proven particularly effective for

transfection of diverse eukaryotes (Janse, Ramesar *et al.*, 2006; Caro *et al.*, 2012; Vinayak *et*

al., 2015), including mammalian primary cells that are resistant to transfection (Gresch *et al.*,

2004; Hamm *et al.*, 2002). To quantify transfection efficiency, we performed luciferase assays

on cell lysates. Through many trials, we eventually achieved a low level of transfection with

nucleofection by improving conditions for culturing *S. rosetta* cells (Fig. S2), modifying

approaches for handling cells throughout the nucleofection procedure (Supplementary

Information), and screening thirty unique combinations of electrical pulses and buffers (Fig. S3).

Optimization around these initial conditions culminated in a procedure that provided robust

and reproducible transfection of *S. rosetta* (Fig. 2A; Methods and

http://www.protocols.io/groups/king-lab). When used in the optimized transfection procedure, all

four transfection reporters drove strong expression of nanoluc protein, producing luminescence

signals that were over three orders of magnitude above the detection limit (Fig. 2B).

Because this was the first example, to our knowledge, of successful transgene expression

in any choanoflagellate, we sought to identify which steps in the optimized protocol were most

essential. Using the p*H3-nanoluc* transfection reporter, we quantified how the omission of each

step impacted transfection efficiency (Fig. 2C). In addition to the use of an optimal electrical

pulse during nucleofection, the two most important steps were priming the cells through the

enzymatic removal of the extracellular matrix prior to nucleofection (Fig. 2A, step 1; Fig. S4) and

the inclusion of carrier DNA during nucleofection (Fig. 2A, step 2; Fig. S5); eliminating either of

these steps resulted in a nearly complete loss of signal.

Priming the cells for nucleofection was a novel step motivated by our observation that *S.*

rosetta cells are surrounded by a potentially protective extracellular coat (Dayel *et al.*, 2011;

Levin *et al.*, 2014; Leadbeater, 2015), and the inclusion of carrier DNA (pUC19) in nucleofection

reactions eliminated the need to include large quantities of reporter construct plasmid (Fig. S5).

Another improvement was the development of a recovery buffer that enhanced transfection ten-

fold (Fig. 2A, step3), presumably by promoting membrane resealing after nucleofection (Rols

and Teissie, 1990; Rols and Teissie, 1989).

Luciferase assays performed on cell lysates gave a sensitive read-out of population-wide

nanoluc expression but did not allow the examination of live, transfected cells nor reveal the

proportion of cells that were successfully transfected. Therefore, we next engineered eight

reporters with different fluorescent proteins placed under the control of regulatory sequences

from the *S. rosetta actin* homolog. Fluorescence was readily detected from cells transfected with

reporters encoding mTFP1 (Ai *et al.*, 2006), mWasabi (Ai *et al.*, 2008), sfGFP (Pedelacq *et al.*,

2006), mNeonGreen (Shaner *et al.*, 2013), mPapaya (Hoi *et al.*, 2013), TagRFP-T(Shaner *et al.*,

2008), mCherry (Shaner *et al.*, 2004), and tdTomato (Shaner *et al.*, 2004). In contrast, an eGFP

(Yang *et al.*, 1996) reporter failed to yield fluorescent cells, likely due to protein misfolding, as

cells transfected with the 'super-folder' variant of GFP (sfGFP) did fluoresce properly. In

transfected cells, the fluorescent signal was distributed throughout the nucleus and cytosol yet

excluded from membrane bound compartments (Fig. 3A).

We observed that transfected cells resembled untransfected cells in their shape, motility,

and ability to propagate, indicating that transfection did not irreparably harm *S. rosetta*.

Fluorescence persisted through multiple cell divisions, yet the diminishing signal in daughter

cells indicated that transfection was transient (Fig. S6A). Importantly, using flow cytometry one

to two days after transfection, we found that ~1% of the population was reproducibly

transfected, and fluorescence-activated cell sorting enriched this transfected cell population

(Fig. S6B). This transfection frequency is comparable to high frequency episomal transformation

of the model yeast *Saccharomyces cerevisiae* that ranges from 1 – 10% (Schiestl and Gietz,

1989; Kawai *et al.*, 2010), and similar transfection frequencies are achieved in model

apicocomplexans (Janse, Franke-Fayard *et al.*, 2006; Caro *et al.*, 2012).

Fluorescent markers illuminate the cell architecture of *S. rosetta*

To demonstrate the versatility of the new method for transfection and simultaneously

explore the cell biology of *S. rosetta in vivo*, we designed a set of fluorescent reporters to mark

key features of *S. rosetta* cells: the nucleus, cytoplasm, collar, filopodia, flagellum, membrane,

mitochondria and endoplasmic reticulum (ER). For each fluorescent reporter, the *mCherry* gene

was fused in-frame to *S. rosetta* DNA sequences encoding conserved proteins or peptides that

localize to specific organelles or subcellular regions in yeast and mammalian cells. To

benchmark each fluorescent marker, we compared its localization in transfected cells to cellular

landmarks known from electron and immunofluorescence micrographs (Fig. S7) (Abedin and

King, 2008; King *et al.*, 2009; King *et al.*, 2008; Sebe-Pedros *et al.*, 2013; Leadbeater, 2015).

Electron micrographs have revealed two distinct regions in the nucleus: the darkly-stained

nucleolus positioned in the center and the surrounding, more lightly-stained nucleoplasm

(Leadbeater, 2015; Burkhardt *et al.*, 2014). As predicted, mCherry fused to either the carboxy

terminus of H3 or the amino terminus of the simian virus 40 nuclear localization signal localized

primarily to the *S. rosetta* nucleoplasm and was excluded from the cytoplasm (Fig. 3B, C)

(Kanda *et al.*, 1998; Kalderon *et al.*, 1984). In contrast, the cytoplasmic marker EFL-mCherry

(Huh *et al.*, 2003) localized to the cytosol and was excluded from the nucleus (Fig. 3D).

Two of the most diagnostic features of the choanoflagellate cell are the actin-filled collar

and the flagellum, which is comprised of microtubules (Karpov and Leadbeater, 1998). A fusion

of mCherry to the filamentous actin-binding peptide Lifeact (Riedl *et al.*, 2008) highlighted the

parallel arrangement of straight microvilli in the collar (Fig. 3G and 3J), as well as filopodia

extending from the basal pole of the cell (Fig. 3G, lower arrow)(Karpov and Leadbeater, 1998;

Sebe-Pedros *et al.*, 2013). In live cells, Lifeact-mCherry revealed the native structure of the

collar, which can be distorted in cells fixed for staining with fluorescent phalloidin or actin

antibodies (Sebe-Pedros *et al.*, 2013). Lifeact-mCherry also showed details of actin filament

organization that have not previously been evident, such as the existence of actin filaments that

originate in the cell body and coalesce at the base of the collar to form each microvillus (Fig.

3G', upper arrow; improved immunofluorescence techniques also preserve these cortical actin

filaments, Fig. 1B). A fusion of α-tubulin to mCherry (Straight *et al.*, 1997) illuminated individual

cortical microtubules emanating from the base of the flagellum to the basal pole of the cell (Fig.

3K', arrow) and allowed visualization of the rapidly beating flagellum in live cells (Fig. 3H and

3K).

A cell membrane marker, with a geranyl-geranylation sequence fused to mCherry (Wang

and Casey, 2016; Reid *et al.*, 2004), outlined the entire cell, including the flagellum, collar, and

cell body (Fig. 3I and 3L), and faintly marked the Golgi apparatus (Fig. 3I', arrow). In live cells,

the cell membrane marker captured the formation of a phagocytic cup engulfing bacterial prey

(Fig. S8)(Dayel and King, 2014). The ER marker (Friedman *et al.*, 2011), which included the

amino terminal signal sequence from the secreted protein Rosetteless (Levin *et al.*, 2014) and a

carboxy terminal ER retention sequence from the ER resident chaperone BiP (PTSG_07223),

highlighted the continuity of the ER with the nuclear envelope and the distribution of ER

throughout the cell, including around vacuoles (Fig. 3F). A mitochondrial marker (Friedman *et*

al., 2011) with an amino terminal targeting sequence from *S. cerevisiae* Cytochrome C Oxidase,

Subunit IV revealed a network of mitochondria (Nunnari *et al.*, 1997) that is enriched around the

nucleus and extends throughout the cell (Fig. 3G). Taken together, these fluorescent markers

demonstrate new experimental capabilities to rapidly tag proteins and to monitor their

localization in distinct cellular compartments and locales.

Transgenesis reveals septin localization in live *S. rosetta* single cells and rosettes

A major motivation for establishing transgenics in *S. rosetta* was to rapidly characterize

candidate genes for multicellularity. Therefore, we investigated the localization of septins, a

family of paralogous genes hypothesized to contribute to multicellular development in *S. rosetta*

(Fairclough *et al.*, 2013). The assembly of septin monomers into higher order structures is

important for septin localization (McMurray *et al.*, 2011) and for the conserved roles of septins in

regulating cytokinesis and cell polarity in fungi and animals. In animals, septins also function in

phagocytosis (Huang *et al.*, 2008), ciliogenesis (Hu *et al.*, 2010), and planar cell polarity (Kim *et*

al., 2010). Septin proteins have a characteristic domain architecture with a diagnostic amino

terminal guanosine triphosphate binding domain (G-domain), and most septins also have a

carboxy terminal coiled-coil domain (Pan *et al.*, 2007; Nishihama *et al.*, 2011) (Fig. 4A). Septin

paralogs interact directly through their G-domains to form heteromeric filaments, and these

heteromeric filaments interact with each other through the septin coiled-coil domains to form

higher order assemblies (Bertin *et al.*, 2008; Garcia *et al.,* 2011; Sirajuddin *et al.*, 2007).

S. rosetta expresses four septin paralogs, three of which have well-annotated genome

models in the current draft genome (Fairclough *et al.*, 2013). We started by examining the

localization of the *S. rosetta* septin protein *Sr*Septin2 (PTSG_07215), a septin with domains

conserved in human and fungal septins (Fairclough *et al.*, 2013) that are necessary for septin

filament formation (Sirajuddin *et al.*, 2007; Bertin *et al.*, 2008). Strikingly, mTFP1-*Sr*Septin2 was

enriched at the basal poles of single and rosettes cells (Fig. 4B and 4E) and at points of contact

between adjacent cells in rosettes (Fig. 4E). This mTFP1-*Sr*Septin2 fusion likely revealed the

native localization of *Sr*Septin2 in *S. rosetta* because septins visualized by immunofluorescence

microscopy in yeast (Ford and Pringle, 1991; Kim, H. B. *et al.,* 1991; Cid *et al.*, 1998; Haarer

and Pringle, 1987), *Drosophila* (Adam *et al.*, 2000; Silverman-Gavrila *et al.*, 2008; Neufeld and

Rubin, 1994), and mammalian cells (Spiliotis *et al.*, 2008) display the same localization as

septins tagged with fluorescent proteins.

To investigate whether *Sr*Septin2 localized to the basal pole of cells might be part of

heteromeric septin filaments, we examined the localization of another septin paralog, *Sr*Septin6

(PTSG_06009) and found that *Sr*Septin6 displays the same basal localization as *Sr*Septin2 (Fig.

4C). Such colocalization (Fig. S9) and the sequence homology with septins that have previously

been shown to form heteromeric filaments (Sirajuddin *et al.*, 2007) strongly suggests that

*Sr*Septin2 and *Sr*Septin6 assemble together at the basal pole. We further found that the basal

localization of *Sr*Septin2 requires the coiled-coil domain, as a complete deletion of the coiled-

coil domain (SrSeptin2ΔCC; Fig. 4A, D, F) eliminated *Sr*Septin2 enrichment at the basal pole

when expressed in wild-type cells. Unexpectedly, mTFP1-*Sr*Septin2ΔCC formed ectopic rings

around vesicles in the cytosol in wild-type cells (Fig. 4D and 4F). The localization of mTFP1-

*Sr*Septin2ΔCC resembled the formation of ectopic septin filaments at convex membranes and

the depletion of septins at the concave membranes in the filamentous fungi *Ashbya gosypii*

(Meseroll *et al.*, 2012) upon the deletion of the coiled-coil domain of the fungal septin Shs1p.

Similarly, vesicles to which mTFP1-*Sr*Septin2ΔCC localized have convex membranes as

opposed to the concave membrane at the basal end of *S. rosetta* where wild-type *Sr*Septin2
localized.
The basal and lateral localization of *Sr*Septin2 and *Sr*Septin6 in rosettes is reminiscent of
septin localization in polarized epithelial cells (Fares *et al.*, 1995; Spiliotis *et al.*, 2008), in which
septins interact with the positive ends of microtubules that are growing toward the basal pole
(Bowen *et al.*, 2011). In choanoflagellates, microtubules radiate down from the apical
microtubule organizing centers (Karpov and Leadbeater, 1998), with the plus ends meeting at
the basal pole of each cell, similar to the orientation of microtubule plus ends toward the basal
pole in animal epithelia (Meads and Schroer, 1995). To examine if septins also interact with the
plus ends of microtubules in *S. rosetta*, we co-transfected cells with mTFP1-*Sr*Septin2 and the
tubulin marker α-tubulin-mCherry (Fig. 4G). Fluorescence microscopy showed that septin
filaments intercalate between cortical microtubules at the basal pole of the cell (Fig. 4G). These
data are consistent with conserved interactions between septins and microtubules from yeast to
animals (Kusch *et al.*, 2002; Kremer *et al.*, 2005; Sellin *et al.*, 2011; Spiliotis *et al.*, 2008),
including at the plus-ends of microtubules in choanoflagellates and animal epithelia (Bowen *et*
al., 2011).
247
DISCUSSION
By synthesizing our growing knowledge of *S. rosetta* biology with a rigorous
characterization and optimization of each step in the transfection procedure, we have developed
a robust method for transgenesis in *S. rosetta* that can be easily implemented by other
laboratories. This method overcomes numerous barriers that prevented efficient DNA delivery in
our prior attempts using diverse methods, including standard electroporation, lipofection,
bombardment, and cell-penetrating peptides. A key breakthrough for this study was the
discovery that the extracellular coat surrounding *S. rosetta* might present a barrier for
transfection, which motivated the development of a method to gently remove the extracellular
material surrounding *S. rosetta,* thereby sensitizing cells for transfection. Additional
improvements to the transfection procedure, such as a step for promoting the closure of the
plasma membrane after electrical pulsation, were designed to address the unique challenges
that arise from culturing *S. rosetta* in sea water. Just as our method was informed by
approaches developed in model microeukaryotes (*Chlamydomonas* and yeast), the methods we
have established in *S. rosetta* will likely extend to aid gene delivery in diverse non-model marine

microeukaryotes. Overall, the gestalt of continually improving choanoflagellate husbandry (Levin

and King, 2013), developing protocols for priming and recovering cells during nucleofection, and

extensively optimizing transfection based on a quantitative assay produced a robust method for

gene delivery in *S. rosetta*.

This work also provides a foundational set of vectors for expressing transgenes in *S.*

rosetta (Dataset S1). In these vectors, the expression of luciferase or fluorescent proteins was

placed under the control of native regulatory elements. From these vectors, we constructed a

panel of fluorescently tagged subcellular markers that serve as references for monitoring the

localization of other proteins in *S. rosetta*. For example, through our pilot study of *Sr*Septin2 and

*Sr*Septin6, the use of these new transgenic tools revealed that septins localize to the basal pole

of choanoflagellates, mirroring their localization in animal epithelial cells (Spiliotis *et al.*, 2008;

Fares *et al.*, 1995).

Observing septin localization in *S. rosetta* contributes to our understanding of how septin

functions evolved prior to the evolution of an epithelium in stem animals. A hallmark of animal

epithelia is tightly connected cells that each have an apical and basal pole comprised of distinct

lipids, proteins, and cytoskeletal structures (Rodriguez-Boulan and Macara, 2014). Septins help

shape epithelia and other types of cells by serving as buttresses (Tanaka-Takiguchi *et al.*, 2009;

Tooley *et al.*, 2009) and diffusion barriers (Barral *et al.*, 2000; Takizawa *et al.*, 2000; Hu *et al.*,

2010) for membranes that have distinct geometries (Bridges *et al.*, 2016; Cannon *et al.*, 2018)

and lipid compositions (Casamayor and Snyder, 2003; Tanaka-Takiguchi *et al.*, 2009; Bertin *et*

al., 2010; Bridges *et al.*, 2014) and by interacting with microtubules (Spiliotis, 2010) and

filamentous actin (Mavrakis *et al.*, 2014). While comparisons between animals and fungi have

revealed conserved functions of septins in cell organization (Spiliotis and Gladfelter, 2012), the

divergence of the lineages that gave rise to fungi and animals over one billion years ago

(Parfrey *et al.*, 2011) resulted in important differences in fungal and animal cell biology (Stajich

et al., 2009). In fungi, septins facilitate polarized cell growth toward the new daughter cell (Kim

et al., 1991; Berepiki and Read, 2013), compartmentalize connected daughter cells (Barral *et*

al., 2000; Takizawa *et al.*, 2000; Helfer and Gladfelter, 2006) , and mediate cytokinesis

(Hartwell, 1971). In addition to their conserved roles in cell division (Neufeld and Rubin, 1994),

animal septins have specific functions in epithelia that maintain apical-basal polarity (Spiliotis *et*

al., 2008), planar cell polarity (Kim *et al.*, 2010), intercellular adhesion (Kim, J. and Cooper,

2018; Park *et al.*, 2015) , and ciliogenesis (Hu *et al.*, 2010; Kim *et al.*, 2010). The basal

localization of septins in *S. rosetta* suggests that septin filaments organize a distinct region at

the basal end of the cell, perhaps supporting intercellular contacts at the basal ends of cells in

rosettes (Fig. 4E). Consistent with this hypothesis is the observation that the Rosetteless

protein, which is necessary for rosette development, localizes to the basal end of cells prior to

secretion into the interior of rosettes where the basal ends of cells meet (Levin *et al.*, 2014).

Continued study of septin function in *S. rosetta* will establish the mechanisms by which septins

facilitate multicellular development, and further comparisons with non-metazoan holozoans with

recently established transgenic methods, such as *Capsaspora owczarzaki* (Parra-Acero *et al.*,

2018) and *Creolimax fragrantissima* (Suga and Ruiz-Trillo, 2013), will provide further insights

into the ancestral functions of septins.

Previous analyses of gene function in choanoflagellates relied on custom antibodies

(Abedin and King, 2008; Young *et al.*, 2011; Burkhardt *et al.*, 2014; Levin *et al.*, 2014), laborious

forward genetic screens (Levin *et al.*, 2014), and *in vitro* biochemistry (Burkhardt *et al.*, 2014).

The ability to express transgenes in *S. rosetta* described here will accelerate studies of the

ancestral functions of animal genes that are conserved in choanoflagellates. We anticipate that

future work will build on this approach, eventually leading to the development of methods for

stable transgenesis and genome editing in *S. rosetta*. Combining an expanded repertoire of

approaches for investigating gene function in-depth in *S. rosetta* with comparisons to other

experimentally-tractable choanoflagellates (Li *et al.,* 2018; Richter *et al.*, 2018) and non-

choanozoans (Suga and Ruiz-Trillo, 2013; Parra-Acero *et al.*, 2018) promises to yield

increasingly mechanistic insights into the ancestry of animal cell biology.

MATERIALS AND METHODS

Cell culture and media preparation

S. rosetta was cultured with a single bacterial species, *E. pacifica*, that serves as a food source (Levin and King, 2013). Media recipes are provided in Table S1. Cultures were established from frozen aliquots by adding 1 ml of thawed cells to 10 ml of 0.2x High Nutrient Media (Table S1). After the cells reached a density of 10^4 cells/ml, the culture was split 1:2 into 1x High Nutrient Media with a constant volume of 0.24 ml/cm^2. After this initial split (denoted as day 0), cells were passaged in 1x High Nutrient Media according to the following schedule: 1:4 dilution on day 1, 1:8 dilution on day 2, 1:16 on day 3. Subsequently cells were passaged every day at a 1:24 dilution or every other day as a 1:48 dilution of cells.

Based on the recommendation from Lonza to use a medium with a low calcium concentration for transfecting mammalian cells, we searched for a seawater recipe with a lower concentration of calcium than the routinely-used artificial seawater made from Tropic of Marin Salts (Levin and King, 2013), which has a calcium concentration of 9.1 mM at a salinity of 35 g/kg (Atkinson and Bingman, 1998). The AK seawater formulation (Table S1) that has been used to culture marine algae (Hallegraeff *et al.*, 2004) and dinoflagellates (Skelton *et al.*, 2009) and has a calcium concentration of 2.7 mM. We found that *S. rosetta* grows more rapidly in 1x High Nutrient Media prepared in AK seawater rather than seawater prepared with Tropic of Marin Salts (Fig. S2A). Therefore, we switched to a growth medium based on AK seawater for routine culturing. After optimizing the nucleofection protocol, we demonstrated that growing *S. rosetta* in AK seawater also resulted in higher transfection efficiencies (Fig. S2B) than growing *S. rosetta* in seawater prepared with Tropic of Marin Salts.

Reporter plasmid design and molecular cloning

Dataset S1 lists the complete inventory of engineered plasmids with a summary of primers, cloning methods, and annotations for constructing each plasmid. Complete plasmid sequences and plasmids have also been deposited at Addgene (http://www.addgene.org/Nicole_King). Below is a brief summary of considerations for designing plasmids, and a more detailed description of standard molecular cloning methods for engineering plasmids can be found in the Supplementary Information.

Cloning regulatory regions from *S. rosetta* genes. Because we had no previous knowledge about the architecture of choanoflagellate regulatory regions, we aimed to clone as

much as 1000 bp upstream and downstream of targeted open reading frames as these

fragments are slightly larger than the mean intergenic distance of 885 bp (Sebe-Pedros *et al.*,

2017). Of necessity, the cloned intergenic sequences reported here were shorter to avoid

repetitive CA and GT sequences that were present before the putative promoter and after the

3'-UTR, respectively. To increase the specificity of primers, we designed the primers to anneal

to regions with a GC content ≤ 50%, as the *S. rosetta* genome is 56% GC. Ultimately, the

cloned regions that encompass the promoter and the 5'-UTR ranged in size from 550 bp to

1095 bp and those encompassing the 3'UTR ranged from 200 bp to 807 bp.

Synthetic gene design. Synthetic reporter genes (*nanoluc* and the genes encoding

diverse fluorescent proteins listed in Supplementary File 1) were codon optimized to match the

codon usage of the set of highly expressed intron-less genes listed in Fig. S1, as codon usage

can be biased for highly expressed genes (Hiraoka *et al.*, 2009). A codon usage table (Dataset

S2) was generated from the coding sequences of highly expressed intronless genes (Dataset

S2A) and from all coding sequences (Dataset S2B) using the 'cusp' tool in Emboss (Rice *et al.*,

2000). The codon usage table was then used to generate a codon optimized DNA sequence for

each target protein sequence with the 'backtranseq' tool on Emboss. The DNA sequences were

further edited by making synonymous substitutions with less frequently used codons to change

restriction enzyme sites and to remove repetitive sequences. Finally, sequences were added to

the ends of these designed genes for cloning with restriction enzymes or Gibson assembly. The

engineered reporter gene sequences are available through Addgene (Dataset S1;

http://www.addgene.org/Nicole_King).

Subcellular marker design. Dataset S3 provides the amino acid sequences for all of the

subcellular markers reported in Fig. 3. To ensure that the fluorescent protein tag for each

marker would not interfere with the functions of proteins or peptides that determine localization,

some of the constructs were engineered to have a flexible linker sequence (SGGSGGS)

separating the fluorescent protein and the localization signals.

376
Optimized transfection protocol

The protocol is summarized in Fig. 2 and detailed protocols for reagent preparation and

transfection are available at protocols.io at the following link:

http://www.protocols.io/groups/king-lab

Culture. Two days prior to transfection, a culture flask (Corning, Cat. No. 353144) was

seeded with *S. rosetta* at a density of 5,000 cells/ml in 200 ml of 1x High Nutrient Media. The

culture was supplemented with 2 mg of frozen *E. pacifica* by resuspending a 10 mg pellet of

flash-frozen *E. pacifica* in 1 ml of media and then adding 200 μl of the resuspended pellet to the

culture of *S. rosetta*.

Wash. After 36-48 hours of growth, bacteria were washed away from *S. rosetta* cells

through three consecutive rounds of centrifugation and resuspension in sterile AK seawater.

The culture flask was vigorously shaken for 30 sec to homogenize the 200 ml that was seeded

two days prior (see above) and then transferred to 50 ml conical tubes and spun for 5 min at

2000 x g and 22°C. The supernatant was removed with a serological pipette, and residual

media was removed with a fine tip transfer pipette. The cell pellets were resuspended in a total

volume of 100 ml of AK seawater, vigorously shaken in their conical tubes for 30 sec, and then

centrifuged for 5 min at 2200 x g and 22°C. The supernatant was removed as before. Each cell

pellet was resuspended in 50 ml of AK seawater, vigorously shaken for 30 sec, and centrifuged

for 5 min at 2400 x g and 22°C. After the supernatant was removed, the cells were resuspended

in a total volume of 100 μl of AK seawater. A 100-fold dilution of cells was counted on a Luna-FL

automated cell counter (Logos Biosystems) and the remaining cells were diluted to a final

concentration of $5x10^7$ choanoflagellate cells/ml. The resuspended cells were divided into 100 μl

aliquots with $5x10^6$ cells per aliquot to immediately prime cells in the next step. A 200 ml culture

typically yields 6-8 aliquots of cells.

Prime. After washing away bacteria, each aliquot of *S. rosetta* cells was incubated in

priming buffer to remove the extracellular material coating the cell. The 100 μl aliquots that

contained $5x10^6$ cells were centrifuged for 5 min at 800 x g and at room temperature. The

supernatant was removed with a fine tip micropipette. Cells were resuspended in 100 μl of

priming buffer (40 mM HEPES-KOH, pH 7.5; 34 mM Lithium Citrate; 50 mM L-Cysteine; 15%

(w/v) PEG 8000; and 1 μM papain) and then incubated for 30 min. Priming was quenched by

adding 2 μl of 50 mg/ml bovine serum albumin-fraction V (Sigma) and then centrifuged for 5 min

at 1250 x g and 22°C with the centrifuge brake set to a 'soft' setting. The supernatant was

removed with a fine-tip micropipette, and the cells were resuspended in 25 μl of SF Buffer

(Lonza).

Nucleofect. Each transfection reaction was prepared by adding 2 μl of 'primed' cells

resuspended in SF buffer to a mixture of 14 μl of SF Buffer; 2 μl of 20 μg/μl pUC19; 1 μl of 250

mM ATP, pH 7.5; 1 μl of 100 mg/ml Sodium Heparin; and ≤7 μl of reporter DNA. (Note that

higher volumes of nucleofection lead to lower transfection frequencies; thus, reporter DNA

should be as concentrated as possible, not exceeding 7 μl. Also, see below for "Note about

titrating reporter plasmids.") The transfection reaction was transferred to one well of a 96-well

nucleofection plate (Lonza). The nucleofection plate was placed in a Nucleofector 4d 96-well

Nucleofection unit (Lonza), and the CM156 pulse was applied to each well.

Rest and recover. Immediately after pulsation, 100 μl of ice-cold recovery buffer (10 mM

HEPES-KOH, pH 7.5; 0.9 M Sorbitol; 8% (w/v) PEG 8000) was added to the cells, Recovery

buffer was gently mixed with the transfected cells by firmly tapping the side of the plate and then

incubating the samples for 5 min. The whole volume of the transfection reaction plus the

recovery buffer was transferred to 1 ml of 1x High Nutrient Media in a 12-well plate. After the

cells recovered for 1 hour, 5 μl of a 10 mg frozen *E. pacifica* pellet resuspended in media (see

above), was added to each well. The cells were grown for 24 to 48 hours before assaying for

luminescence or fluorescence.

Note about establishing transfection in non-model microeukaryotes. Establishing a

transfection protocol for *S. rosetta* required adapting several different transfection procedures

for a variety of eukaryotic cells to meet the unique requirements of *S. rosetta*. While the specific

details for transfecting *S. rosetta* may not be readily applicable to other organisms, the general

considerations and the process for optimization that led to the development of the transfection

protocol described here could inform efforts to transfect other microeukaryotes. Therefore, we

have included a summary in the supplementary information (See text and Figs. S10 and S11) of

the initial development and optimization of the aforementioned protocol.

Nanoluc reporter assay

To measure relative transfection efficiency resulting from different transfection protocols

and promoters, we performed luciferase assays on lysates of transfected cells. Cells transfected

with 2.5 μg of *nanoluc* reporter plasmids were pelleted by centrifuging for 10 min at 4200 x g

and 4°C. The supernatant was removed and the cells were resuspended in 50 μl of NanoGlo

buffer (Promega) and then transferred to a well of a white, opaque 96-well plate (Greiner Bio-

one Cat No.655083). Luminescence was immediately recorded on a Spectramax L Microplate

Reader (Molecular Devices) with a 1 min dark adaption and 10 sec dwell time with the

photomultiplier gain set to photon counting mode.

Based on standard definitions from analytical chemistry (Harris, 2007) the detection limit

was set to three standard deviations above the background signal such that any signal above

the detection limit has less than a 1% chance of arising from random error. The limit of detection

was calculated in two different ways. First, the y-axis intercept and standard deviation were

calculated from a standard curve (Harris, 2007) fit to a serial dilution of nanoluc versus

luciferase activity (Fig. S3A). To decrease the bias toward higher luciferase values, the standard

curve was fit with the objective *(O)*: $O \equiv \min \sum_i \frac{|m_i - c_i|}{m_i}$, where m is the measured luciferase value

for a given data point i and c is the calculated luciferase value. Second, the detection limit was

also determined as three standard deviations above the mean of eight replicate luciferase

measurements of cells transfected without any reporter plasmid, which resulted in the same

calculated detection limit.

Reproducibility in luciferase assays was assessed by performing at least two independent

experiments on separate days with different preparations of 'primed' cells; data presented in

Fig. 2, S2, S3, S10, and S11 represent one of the independent experiments. Within each

experiment from the same preparation of 'primed' cells, replicate measurements were

performed by setting up three to five independent transfections for each condition (shown as

black dots); bar graphs in Fig. 2, S2, S3, S10, and S11 show the mean values of the five

independent transfections with error bars showing the standard deviation. Before performing

statistical tests that rely on a normal distribution, luciferase data were transformed to a log-

normal distribution by taking the base-10 logarithm of luciferase values as gene expression data

from luciferase assays display a log-normal distribution (Muranaka *et al.*, 2013).

466
Flow Cytometry

To measure the percentage of cells expressing each of the different transgenes under

different transfection conditions, we used flow cytometry. Cells were transfected with 10 μg of

mWasabi or 10 μg of TagRFP-T reporter plasmids for flow cytometry because these

fluorophores produced the highest fluorescence signal upon illumination with the 488 or 561 nm

lasers, respectively. To prepare cells for flow cytometry, cultures from 10-12 transfections were

pooled 24 hours after transfection and centrifuged for 15 min at 3600 x g and 4°C. The

supernatant was removed with a fine-tip transfer pipette to avoid disturbing the pellet. The

pelleted cells were resuspended in 500 μl of 0.22 μm filtered AK seawater and then filtered

through a 40 μm filter.

Because a large number of bacteria were present in the cultures, *S. rosetta* cells were

gated based on the area of forward scattering signal versus the area of the side scattering

signal and the area of the forward scattering signal versus the height of the forward scattering

signals. To differentiate transfected cells from untransfected cells, fluorescence signal was

measured using lasers and filters for the fluorophores FITC (Green Fluorescence) and PE (Red

Fluorescence); untransfected cells form a population along the y=x line of these plots, and the

population of transfected cells are skewed along one axis that corresponds to the fluorophore.

The transfected cells were gated to exclude >99.99% of untransfected cells as determined from

a negative control reaction that was transfected without a fluorescent reporter (Fig. 2E, left

panel).

Live cell imaging

An important benefit of transgenics is the ability to visualize protein localization and cell

architecture in living cells. To this end, we have established improved protocols for live cell

imaging in *S. rosetta*. Glass-bottomed dishes were prepared for live cell microscopy by corona-

treating the glass for 10 s. Afterwards, 300 μl of 0.1 mg/ml poly-D-lysine was applied to the

glass cover (18 μl/cm^2), incubated for 10 min at room temperature, and then removed. Excess

poly-D-lysine was washed away from the glass surface with three rinses of 500 μl artificial

seawater.

Cells transfected with 5 μg of each fluorescent reporter were prepared for microscopy by

centrifuging 1-2 ml of transfected cells for 10 min at 3,600 x g and 4°C. After centrifugation, the

supernatant was removed and the cell pellet was resuspended in 200 μl of 4/5 Tropic of Marin

artificial seawater with 100 mM LiCl. Lithium chloride slows flagellar beating, as in spermatozoa

(Brokaw, 1987; Gibbons and Gibbons, 2013), to decrease the movement of cells during

imaging. The resuspended cells were pipetted on top of the poly-D-lysine coated glass-bottom

dish and adsorbed on the surface for 10 min. Lastly, 200 μl of 20% (w/v) Ficoll 400 dissolved in

4/5 Tropic of Marin artificial seawater with 100 mM LiCl was pipetted drop-by-drop on top of the

cells. The addition of Ficoll decreases flagellar movement by increasing the viscosity of the

media (Wilson *et al.*, 2015; Pate and Brokaw, 1980) without significantly changing the

osmolarity or refractive index of the sample (GE Lifesciences).

Confocal microscopy was performed on a Zeiss Axio Observer LSM 880 with an Airyscan

detector and a 63x/NA1.40 Plan-Apochromatic oil immersion objective. The mTFP1 and

mCherry fluorophores were selected for two-color imaging due to their high photostability and

minimal spectral overlap. Confocal stacks were acquired in superresolution mode using ILEX

line scanning and two-fold averaging and the following settings: 40 nm x 40 nm pixel size, 93

nm z-step, 0.9-1.0 μsec/pixel dwell time, 850 gain, 458 nm laser operating at 5% laser power,

561 nm laser operating at 3% laser power, 458/561 nm multiple beam splitter, and 495-550 nm

band-pass/570 nm long-pass filter. Images were initially processed using the automated
Airyscan algorithm (Zeiss) and then reprocessed by setting the Airyscan threshold 0.5 units
higher than the value reported from automated Airyscan processing. The stacks were further
processed by correcting for signal decay, background, and flickr in Zen Blue (Zeiss). Last, FIJI
(Schindelin *et al.*, 2012) was used to apply a gamma factor to each channel and subtract the
background using a 100 pixel radius.

Epifluorescence and differential interference contrast images were recorded using a Zeiss
Axio Observer.Z1/7 Widefield microscope with a Hamamatsu Orca-Flash 4.0 LT CMOS Digital
Camera and 40x/NA 1.1 LD C-Apochromatic water immersion, 63x/NA1.40 Plan-Apochromatic
oil immersion, or 100x NA 1.40 Plan-Apochromatic oil immersion objectives. Green fluorescent
proteins were imaged with a 38 HE filter set and red fluorescent proteins with a 43 HE filter set.
Images were processed by applying a gamma factor and background subtracting fluorescence
channels in FIJI.

Note about titrating reporter plasmids. A titration of fluorescent reporter plasmids
showed that 10 μg of total reporter plasmid(s) best balanced transfection efficiency, brightness,
and a faithful indication of subcellular architecture. We caution that high plasmid concentrations
can result in the overexpression of fluorescent markers, leading to aberrant localization of the
marker and gross changes in cell morphology. Such artefacts can be avoided by performing a
titration to determine the best concentration of plasmid and recording images from cells with a
range of fluorescence intensities that result from any transfection. One of the best markers to
assess optimal reporter plasmid concentrations is the tubulin marker because of its distinct
localization that can be benchmarked with immunofluorescence.

Immunofluorescence staining and imaging
Immunfluorescence was performed as previously described (Woznica *et al.*, 2016) with
modifications to better preserve features of the cytoskeleton. Two milliliters of cells were
concentrated by centrifugation for 10 min at 2750 x g and 4°C. The cells were resuspended in
400 μl of artificial seawater and applied to poly-L-lysine coated coverslips (BD Biosciences)
placed in the bottom of each well of a 24-well cell culture dish. After allowing the cells to settle
on the coverslip for 30 min, 150 μl of the cell solution was gently removed from the side of the
dish. It is crucial to leave a small layer of buffer on top of cells to preserve the cell morphology,
hence the 250 μl of liquid left in the well. All of the subsequent washes and incubations during
the staining procedure were performed by adding and removing 200 μl of the indicated buffer.

Cells were fixed in two stages. First, the coverslip was washed once with 6% acetone in

cytoskeleton buffer (10 mM MES, pH 6.1; 138 KCl, 3 mM $MgCl_2$; 2 mM EGTA; 675 mM

Sucrose), which better preserves the actin cytoskeleton (Cramer and Mitchison, 1995; Symons

and Mitchison, 1991), and then incubated for 10 min at room temperature after a second

application of the acetone solution. Subsequently, the coverslip was washed once with 4%

formaldehyde diluted in cytoskeleton buffer and then incubated for 15 min at room temperature

after a second application of the formaldehyde solution. Last, the coverslip was gently washed

three times with cytoskeleton buffer.

Cells were permeabilized by washing the coverslip once with permeabilization buffer (100

mM PIPES, pH 6.95; 2 mM EGTA; 1 mM $MgCl_2$; 1% (w/v) bovine serum albumin-fraction V;

0.3% (v/v Triton X-100) and then incubated for 30 min upon a second addition of

permeabilization buffer. After the permeabilization buffer was removed, the coverslip was

washed once with primary antibody, 50 ng/ml mouse E7 anti-tubulin antibody (Developmental

Studies Hybridoma Bank) diluted in permeabilization buffer, and then incubated for 1 h in a

second application of primary antibody. The coverslip was gently washed twice in

permeabilization buffer. Next, the coverslip was washed once with secondary antibody, 8 ng/ml

Donkey anti-mouse IgG–AlexaFluor568 (ThermoFisher) diluted in permeabilization buffer, and

then incubated for 1 h after a second application of secondary antibody. Afterwards, the

coverslip was washed once in permeabilization buffer and then three times with PEM (100 mM

PIPES-KOH, pH 6.95; 2 mM EGTA; 1 mM $MgCl_2$). The coverslip was washed once with 10

μg/ml Hoechst 33342 and 4 U/ml Phalloidin-AlexaFluor488 in PEM and then incubated for 30

min with a second application of Hoechst33342/Phalloidin. Finally, the coverslip was washed

once in PEM.

To prepare a slide for mounting, 10 μl of Pro-Long Diamond (Invitrogen) was added to a

slide. The coverslip was gently removed from the well with forceps, excess buffer was blotted

from the side with a piece of filter paper, and the coverslip was gently placed on the drop of Pro-

Long diamond. The mounting media was allowed to cure overnight before visualization.

Images were acquired on a Zeiss LSM 880 Airyscan confocal microscope with a 63x

objective (as described for live cell imaging) by frame scanning in the superresolution mode with

the following settings: 35 nm x 35 nm pixel size; 80 nm z-step; 0.64 μs/pixel dwell time; 561 nm

laser operating at 1.5% power with a 488/561 nm beam splitter, a 420-480 nm/495-620 nm

band pass filter, and a gain of 750; 488 nm laser operating at 1.5% power with a 488/561 nm

beam splitter, a 420-480 nm/495-550 nm band pass filter, and a gain of 750; and 405 nm laser

operating at 1.5% power with a 405 nm beam splitter, a 420-480 nm/495-550 nm band pass

filter, and a gain of 775.

Acknowledgements

Laura Wetzel, Monika Sigg, Hannah Elzinga, Lily Helfrich, and Reef Aldayafleh helped with experiments and reagent preparation. Corey Allard, as part of the Marine Biological Laboratory's Physiology Course, helped with early tests of priming conditions. Kent McDonald generously provided a transmission electron micrograph from samples prepared by Pawel Burkhardt. We thank these people for providing access and support for scientific instruments: Russell Vance and Lab for use of their luminometer, Hector Nolla and Alma Valeros in the Flow Cytometry Facility, and the UC Berkeley DNA Sequencing Facility. The following individuals generously donated reagents and provided technical support: Brad Hook and Dee Czarniecki from Promega, Ethan Brooks from Lonza, and Colleen Manning from Zeiss. We appreciate scientific discussions and advice from these individuals: David Schaffer, Sabrina Sun, Jorge Ortiz, Niren Murthy, Tara DeBoer, Fyodor Urnov, Matt Welch and Lab, Rebecca Heald and Lab, Abby Dernberg and Lab, and Amy Gladfelter. We thank members of the King lab for helpful discussions, research support, and comments on the manuscript, especially Arielle Woznica, Kayley Hake, and Ben Larson. An additional thanks to the following people for providing comments on the manuscript: Candace Britton, Pawel Burkhardt, Matt Daugherty, Galo Garcia, Tera Levin, Kristin Patrick, and Dan Richter. DSB is supported as a Simons Foundation Postdoctoral Fellow of the Jane Coffin Childs Memorial Fund for Biomedical Research. This work was funded in part by a grant from the Gordon and Betty Moore Foundation's Marine Microbiology Initiative for establishing Emerging Model Systems.

Author Contributions

DSB and NK conceived of the project and wrote the manuscript. DSB, HSM, and NK designed experiments and interpreted data. DSB and HSM collected data.

REFERENCES

Abedin, M. and King, N. (2008). The premetazoan ancestry of cadherins. Science *5865*, 946-
948.

Adam, J. C., Pringle, J. R. and Peifer, M. (2000). Evidence for functional differentiation among
Drosophila septins in cytokinesis and cellularization. Mol. Biol. Cell *9*, 3123-3135.

Ai, H. W., Henderson, J. N., Remington, S. J. and Campbell, R. E. (2006). Directed evolution of
a monomeric, bright and photostable version of *Clavularia* cyan fluorescent protein: structural
characterization and applications in fluorescence imaging. Biochem. J. *3*, 531-540.

Ai, H. W., Olenych, S. G., Wong, P., Davidson, M. W. and Campbell, R. E. (2008). Hue-shifted
monomeric variants of *Clavularia* cyan fluorescent protein: identification of the molecular
determinants of color and applications in fluorescence imaging. BMC Biol., 13.

Alegado, R. A., Brown, L. W., Cao, S., Dermenjian, R. K., Zuzow, R., Fairclough, S. R., Clardy,
J. and King, N. (2012). A bacterial sulfonolipid triggers multicellular development in the closest
living relatives of animals. eLife, e00013.

Atkinson, M. J. and Bingman, C. (1998). Elemental composition of commercial seasalts. Journal
of Aquariculture and Aquatic Sciences *2*, 39-43.

Barral, Y., Mermall, V., Mooseker, M. S. and Snyder, M. (2000). Compartmentalization of the
cell cortex by septins is required for maintenance of cell polarity in yeast. Mol. Cell *5*, 841-851.

Berepiki, A. and Read, N. D. (2013). Septins are important for cell polarity, septation and
asexual spore formation in *Neurospora crassa* and show different patterns of localisation at
germ tube tips. PLoS One *5*, e63843.

Bertin, A., McMurray, M. A., Grob, P., Park, S. S., Garcia, G., Patanwala, I., Ng, H. L., Alber, T.,
Thorner, J. and Nogales, E. (2008). *Saccharomyces cerevisiae* septins: supramolecular
organization of heterooligomers and the mechanism of filament assembly. Proc. Natl. Acad. Sci.
U. S. A. *24*, 8274-8279.

Bertin, A., McMurray, M. A., Thai, L., Garcia, G., Votin, V., Grob, P., Allyn, T., Thorner, J. and
Nogales, E. (2010). Phosphatidylinositol-4,5-bisphosphate promotes budding yeast septin
filament assembly and organization. J. Mol. Biol. *4*, 711-731.

Bowen, J. R., Hwang, D., Bai, X., Roy, D. and Spiliotis, E. T. (2011). Septin GTPases spatially
guide microtubule organization and plus end dynamics in polarizing epithelia. J. Cell Biol. *2*,
187-197.

Bridges, A. A., Jentzsch, M. S., Oakes, P. W., Occhipinti, P. and Gladfelter, A. S. (2016).
Micron-scale plasma membrane curvature is recognized by the septin cytoskeleton. J. Cell Biol.
1, 23-32.

Bridges, A. A., Zhang, H., Mehta, S. B., Occhipinti, P., Tani, T. and Gladfelter, A. S. (2014).
Septin assemblies form by diffusion-driven annealing on membranes. Proc. Natl. Acad. Sci. U.
S. A. *6*, 2146-2151.

Brokaw, C. J. (1987). A lithium-sensitive regulator of sperm flagellar oscillation is activated by
cAMP-dependent phosphorylation. J. Cell Biol. *4*, 1789-1798.

Brunet, T. and King, N. (2017). The Origin of Animal Multicellularity and Cell Differentiation.
Dev. Cell. *2*, 124-140.

Burger, G., Forget, L., Zhu, Y., Gray, M. W. and Lang, B. F. (2003). Unique mitochondrial
genome architecture in unicellular relatives of animals. Proc. Natl. Acad. Sci. U. S. A. *3*, 892-
897.

Burkhardt, P., Gronborg, M., McDonald, K., Sulur, T., Wang, Q. and King, N. (2014).
Evolutionary insights into premetazoan functions of the neuronal protein homer. Mol. Biol. Evol.
9, 2342-2355.

Cannon, K. S., Woods, B. L., Crutchley, J. M. and Gladfelter, A. S. (2018). An amphipathic helix
enables septins to sense micron-scale membrane curvature. bioRxiv.

Caro, F., Miller, M. G. and DeRisi, J. L. (2012). Plate-based transfection and culturing technique
for genetic manipulation of *Plasmodium falciparum*. Malar J., 22.

Casamayor, A. and Snyder, M. (2003). Molecular dissection of a yeast septin: distinct domains
are required for septin interaction, localization, and function. Mol. Cell. Biol. *8*, 2762-2777.

Cid, V. J., Adamikova, L., Cenamor, R., Molina, M., Sanchez, M. and Nombela, C. (1998). Cell
integrity and morphogenesis in a budding yeast septin mutant. Microbiology *Pt 12*, 3463-3474.

Cramer, L. P. and Mitchison, T. J. (1995). Myosin is involved in postmitotic cell spreading. J.
Cell Biol. *1*, 179-189.

Dayel, M. J., Alegado, R. A., Fairclough, S. R., Levin, T. C., Nichols, S. A., McDonald, K. and
King, N. (2011). Cell differentiation and morphogenesis in the colony-forming choanoflagellate
Salpingoeca rosetta. Dev. Biol. *1*, 73-82.

Dayel, M. J. and King, N. (2014). Prey capture and phagocytosis in the choanoflagellate
Salpingoeca rosetta. PLoS One *5*, e95577.

Fairclough, S. R. *et al.* (2013). Premetazoan genome evolution and the regulation of cell
differentiation in the choanoflagellate *Salpingoeca rosetta*. Genome Biol. *2*, r15.

Fairclough, S. R., Dayel, M. J. and King, N. (2010). Multicellular development in a
choanoflagellate. Curr. Biol. *20*, 875.

Fares, H., Peifer, M. and Pringle, J. R. (1995). Localization and possible functions of *Drosophila*
septins. Mol. Biol. Cell *12*, 1843-1859.

Ford, S. K. and Pringle, J. R. (1991). Cellular morphogenesis in the *Saccharomyces cerevisiae*
cell cycle: localization of the CDC11 gene product and the timing of events at the budding site.
Dev. Genet. *4*, 281-292.

Friedman, J. R., Lackner, L. L., West, M., DiBenedetto, J. R., Nunnari, J. and Voeltz, G. K.
(2011). ER tubules mark sites of mitochondrial division. Science *6054*, 358-362.

Garcia, G., Bertin, A., Li, Z., Song, Y., McMurray, M. A., Thorner, J. and Nogales, E. (2011).
Subunit-dependent modulation of septin assembly: budding yeast septin Shs1 promotes ring
and gauze formation. J. Cell Biol. *6*, 993-1004.

Gibbons, B. H. and Gibbons, I. R. (2013). Lithium reversibly inhibits microtubule-based motility
in sperm flagella. Nature *5968*, 560-562.

Gresch, O. *et al.* (2004). New non-viral method for gene transfer into primary cells. Methods *2*,
151-163.

Haarer, B. K. and Pringle, J. R. (1987). Immunofluorescence localization of the *Saccharomyces*
cerevisiae CDC12 gene product to the vicinity of the 10-nm filaments in the mother-bud neck.
Mol. Cell. Biol. *10*, 3678-3687.

Hall, M. P. *et al.* (2012). Engineered luciferase reporter from a deep sea shrimp utilizing a novel
imidazopyrazinone substrate. ACS Chem. Biol. *11*, 1848-1857.

Hallegraeff, G. M., Anderson, D. M. and Cembella, A. D. (2004). Manual on Harmful
Marine Microalgae, Paris, France: United Nations Educational, Scientific and Cultural
Organization.

Hamm, A., Krott, N., Breibach, I., Blindt, R. and Bosserhoff, A. K. (2002). Efficient transfection
method for primary cells. Tissue Eng. *2*, 235-245.

Harris, D. C. (2007). Quantitative chemical analysis, New York, NY: W.H. Freeman and Co.

Hartwell, L. H. (1971). Genetic control of the cell division cycle in yeast. IV. Genes controlling
bud emergence and cytokinesis. Exp. Cell Res. *2*, 265-276.

Helfer, H. and Gladfelter, A. S. (2006). AgSwe1p regulates mitosis in response to
morphogenesis and nutrients in multinucleated *Ashbya gossypii* cells. Mol. Biol. Cell *10*, 4494-
4512.

Hiraoka, Y., Kawamata, K., Haraguchi, T. and Chikashige, Y. (2009). Codon usage bias is
correlated with gene expression levels in the fission yeast *Schizosaccharomyces pombe*. Genes
Cells *4*, 499-509.

Hoi, H., Howe, E. S., Ding, Y., Zhang, W., Baird, M. A., Sell, B. R., Allen, J. R., Davidson, M. W.
and Campbell, R. E. (2013). An engineered monomeric *Zoanthus* sp. yellow fluorescent protein.
Chem. Biol. *10*, 1296-1304.

Hu, Q., Milenkovic, L., Jin, H., Scott, M. P., Nachury, M. V., Spiliotis, E. T. and Nelson, W. J.
(2010). A septin diffusion barrier at the base of the primary cilium maintains ciliary membrane
protein distribution. Science *5990*, 436-439.

Huang, Y. W., Yan, M., Collins, R. F., Diciccio, J. E., Grinstein, S. and Trimble, W. S. (2008).
Mammalian septins are required for phagosome formation. Mol. Biol. Cell *4*, 1717-1726.

Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S. and
O'Shea, E. K. (2003). Global analysis of protein localization in budding yeast. Nature *6959*, 686-
691.

James-Clark, H. (1868). On the Spongiae Ciliatae as Infusoria Flagellata: or observations on the
structure, animality and relationship of *Leucosolenia botryoides* Bowerbank. The Annals and
Magazine of Natural History, 250–64.

Janse, C. J., Franke-Fayard, B., Mair, G. R., Ramesar, J., Thiel, C., Engelmann, S.,
Matuschewski, K., van Gemert, G. J., Sauerwein, R. W. and Waters, A. P. (2006). High
efficiency transfection of *Plasmodium berghei* facilitates novel selection procedures. Mol.
Biochem. Parasitol. *1*, 60-70.

Janse, C. J., Ramesar, J. and Waters, A. P. (2006). High-efficiency transfection and drug
selection of genetically transformed blood stages of the rodent malaria parasite *Plasmodium*
berghei. Nat. Protoc. *1*, 346-356.

Kalderon, D., Roberts, B. L., Richardson, W. D. and Smith, A. E. (1984). A short amino acid
sequence able to specify nuclear location. Cell *3 Pt 2*, 499-509.

Kanda, T., Sullivan, K. F. and Wahl, G. M. (1998). Histone-GFP fusion protein enables sensitive
analysis of chromosome dynamics in living mammalian cells. Curr. Biol. *7*, 377-385.

Karpov, S. A. and Leadbeater, B. S. C. (1998). Cytoskeleton Structure and Composition in
Choanoflagellates. J. Eukaryot. Microbiol. *3*, 361-367.

Kawai, S., Hashimoto, W. and Murata, K. (2010). Transformation of *Saccharomyces cerevisiae*
and other fungi: methods and possible underlying mechanism. Bioeng. Bugs *6*, 395-403.

Kent, W. S. (1871). Affinities of the sponges. Nature. *4*, 184.

Kim, H. B., Haarer, B. K. and Pringle, J. R. (1991). Cellular morphogenesis in the
Saccharomyces cerevisiae cell cycle: localization of the CDC3 gene product and the timing of
events at the budding site. J. Cell Biol. *4*, 535-544.

Kim, J. and Cooper, J. A. (2018). Septins regulate junctional integrity of endothelial monolayers.
Mol. Biol. Cell *13*, 1693-1703.

Kim, S. K. *et al.* (2010). Planar cell polarity acts through septins to control collective cell
movement and ciliogenesis. Science *5997*, 1337-1340.

King, N. (2004). The unicellular ancestry of animal development. Dev. Cell. *3*, 313-325.

King, N., Hittinger, C. T. and Carroll, S. B. (2003). Evolution of key cell signaling and adhesion
protein families predates animal origins. Science *5631*, 361-363.

King, N. *et al.* (2008). The genome of the choanoflagellate *Monosiga brevicollis* and the origin of
metazoans. Nature *7180*, 783-788.

King, N., Young, S. L., Abedin, M., Carr, M. and Leadbeater, B. S. (2009). Visualizing the
subcellular localization of actin, beta-tubulin, and DNA in *Monosiga brevicollis*. Cold Spring Harb
Protoc. *2*, pdb.prot5150.

Kremer, B. E., Haystead, T. and Macara, I. G. (2005). Mammalian septins regulate microtubule
stability through interaction with the microtubule-binding protein MAP4. Mol. Biol. Cell *10*, 4648-
4659.

Kusch, J., Meyer, A., Snyder, M. P. and Barral, Y. (2002). Microtubule capture by the cleavage
apparatus is required for proper spindle positioning in yeast. Genes Dev. *13*, 1627-1639.

Lang, B. F., O'Kelly, C., Nerad, T., Gray, M. W. and Burger, G. (2002). The closest unicellular
relatives of animals. Curr. Biol. *20*, 1773-1778.

Leadbeater, B. S. C. (2015). The Choanoflagellates: Evolution, Biology and Ecology,
Cambridge, United Kingdom: Cambridge University Press.

Levin, T. C., Greaney, A. J., Wetzel, L. and King, N. (2014). The rosetteless gene controls
development in the choanoflagellate *S. rosetta*. eLife, e04070.

Levin, T. C. and King, N. (2013). Evidence for sex and recombination in the choanoflagellate
Salpingoeca rosetta. Curr. Biol. *21*, 2176-2180.

Li, R., Neundorf, I. and Nitsche, F. (2018). First Efficient Transfection in Choanoflagellates using
Cell-Penetrating Peptides. bioRxiv.

Manning, G., Young, S. L., Miller, W. T. and Zhai, Y. (2008). The protist, *Monosiga brevicollis*,
has a tyrosine kinase signaling network more elaborate and diverse than found in any known
metazoan. Proc. Natl. Acad. Sci. U. S. A. *28*, 9674-9679.

Mavrakis, M., Azou-Gros, Y., Tsai, F. C., Alvarado, J., Bertin, A., Iv, F., Kress, A., Brasselet, S.,
Koenderink, G. H. and Lecuit, T. (2014). Septins promote F-actin ring formation by crosslinking
actin filaments into curved bundles. Nat. Cell Biol. *4*, 322-334.

McMurray, M. A., Bertin, A., Garcia, G., Lam, L., Nogales, E. and Thorner, J. (2011). Septin
filament formation is essential in budding yeast. Dev. Cell. *4*, 540-549.

Meads, T. and Schroer, T. A. (1995). Polarity and nucleation of microtubules in polarized
epithelial cells. Cell Motil. Cytoskeleton *4*, 273-288.

Meseroll, R. A., Howard, L. and Gladfelter, A. S. (2012). Septin ring size scaling and dynamics
require the coiled-coil region of Shs1p. Mol. Biol. Cell *17*, 3391-3406.

Muranaka, T., Kubota, S. and Oyama, T. (2013). A single-cell bioluminescence imaging system
for monitoring cellular gene expression in a plant body. Plant Cell Physiol. *12*, 2085-2093.

Neufeld, T. P. and Rubin, G. M. (1994). The *Drosophila* peanut gene is required for cytokinesis
and encodes a protein similar to yeast putative bud neck filament proteins. Cell *3*, 371-379.

Nishihama, R., Onishi, M. and Pringle, J. R. (2011). New insights into the phylogenetic
distribution and evolutionary origins of the septins. Biol. Chem. *8-9*, 681-687.

Nunnari, J., Marshall, W. F., Straight, A., Murray, A., Sedat, J. W. and Walter, P. (1997).
Mitochondrial transmission during mating in *Saccharomyces cerevisiae* is determined by
mitochondrial fusion and fission and the intramitochondrial segregation of mitochondrial DNA.
Mol. Biol. Cell *7*, 1233-1242.

O'Neill, R. S. and Clark, D. V. (2013). The *Drosophila melanogaster* septin gene Sep2 has a
redundant function with the retrogene Sep5 in imaginal cell proliferation but is essential for
oogenesis. Genome *12*, 753-758.

Pan, F., Malmberg, R. L. and Momany, M. (2007). Analysis of septins across kingdoms reveals
orthology and new motifs. BMC Evol. Biol., 103.

Parfrey, L. W., Lahr, D. J., Knoll, A. H. and Katz, L. A. (2011). Estimating the timing of early
eukaryotic diversification with multigene molecular clocks. Proc. Natl. Acad. Sci. U. S. A. *33*,
13624-13629.

Park, T. J., Kim, S. K. and Wallingford, J. B. (2015). The planar cell polarity effector protein
Wdpcp (Fritz) controls epithelial cell cortex dynamics via septins and actomyosin. Biochem.
Biophys. Res. Commun. *2*, 562-566.

Parra-Acero, H., Ros-Rocher, N., Perez-Posada, A., Kożyczkowska, A., Sánchez-Pons, N.,
Nakata, A., Suga, H., Najle, S. R. and Ruiz-Trillo, I. (2018). Transfection of *Capsaspora*
owczarzaki, a close unicellular relative of animals. Development.

Pate, E. F. and Brokaw, C. J. (1980). Movement of spermatozoa in viscous environments. J.
Exp. Biol., 395-397.

Pedelacq, J. D., Cabantous, S., Tran, T., Terwilliger, T. C. and Waldo, G. S. (2006). Engineering
and characterization of a superfolder green fluorescent protein. Nat. Biotechnol. *1*, 79-88.

Reid, T. S., Terry, K. L., Casey, P. J. and Beese, L. S. (2004). Crystallographic analysis of CaaX
prenyltransferases complexed with substrates defines rules of protein substrate selectivity. J.
Mol. Biol. *2*, 417-433.

Rice, P., Longden, I. and Bleasby, A. (2000). EMBOSS: the European Molecular Biology Open
Software Suite. Trends Genet. *6*, 276-277.

Richter, D., Fozouni, P., Eisen, M. and King, N. (2018). Gene family innovation, conservation
and loss on the animal stem lineage. eLife, e34226.

Riedl, J. *et al.* (2008). Lifeact: a versatile marker to visualize F-actin. Nat. Methods *7*, 605-607.

Rodriguez-Boulan, E. and Macara, I. G. (2014). Organization and execution of the epithelial
polarity programme. Nat. Rev. Mol. Cell Biol. *4*, 225-242.

Rols, M. P. and Teissie, J. (1990). Modulation of electrically induced permeabilization and fusion
of Chinese hamster ovary cells by osmotic pressure. Biochemistry *19*, 4561-4567.

Rols, M. P. and Teissie, J. (1989). Ionic-strength modulation of electrically induced
permeabilization and associated fusion of mammalian cells. Eur. J. Biochem. *1*, 109-115.

Ruiz-Trillo, I., Roger, A. J., Burger, G., Gray, M. W. and Lang, B. F. (2008). A phylogenomic
investigation into the origin of metazoa. Mol. Biol. Evol. *4*, 664-672.

Schiestl, R. H. and Gietz, R. D. (1989). High efficiency transformation of intact yeast cells using
single stranded nucleic acids as a carrier. Curr. Genet. *5-6*, 339-346.

Schindelin, J. *et al.* (2012). Fiji: an open-source platform for biological-image analysis. Nat.
Methods *7*, 676-682.

Sebe-Pedros, A., Burkhardt, P., Sanchez-Pons, N., Fairclough, S. R., Lang, B. F., King, N. and
Ruiz-Trillo, I. (2013). Insights into the origin of metazoan filopodia and microvilli. Mol. Biol. Evol.
9, 2013-2023.

Sebe-Pedros, A., Degnan, B. M. and Ruiz-Trillo, I. (2017). The origin of Metazoa: a unicellular
perspective. Nat. Rev. Genet. *8*, 498-512.

Sellin, M. E., Holmfeldt, P., Stenmark, S. and Gullberg, M. (2011). Microtubules support a disk-
like septin arrangement at the plasma membrane of mammalian cells. Mol. Biol. Cell *23*, 4588-
4601.

Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N., Palmer, A. E. and Tsien, R.
Y. (2004). Improved monomeric red, orange and yellow fluorescent proteins derived from
Discosoma sp. red fluorescent protein. Nat. Biotechnol. *12*, 1567-1572.

Shaner, N. C. *et al.* (2013). A bright monomeric green fluorescent protein derived from
Branchiostoma lanceolatum. Nat. Methods *5*, 407-409.

Shaner, N. C., Lin, M. Z., McKeown, M. R., Steinbach, P. A., Hazelwood, K. L., Davidson, M. W.
and Tsien, R. Y. (2008). Improving the photostability of bright monomeric orange and red
fluorescent proteins. Nat. Methods *6*, 545-551.

Silverman-Gavrila, R. V., Hales, K. G. and Wilde, A. (2008). Anillin-mediated targeting of peanut
to pseudocleavage furrows is regulated by the GTPase Ran. Mol. Biol. Cell *9*, 3735-3744.

Sirajuddin, M., Farkasovsky, M., Hauer, F., Kuhlmann, D., Macara, I. G., Weyand, M., Stark, H.
and Wittinghofer, A. (2007). Structural insight into filament formation by mammalian septins.
Nature *7160*, 311-315.

Skelton, H. M., Burkholder, J. M. and Parrow, M. W. (2009). Axenic culture of the heterotrophic
dinoflagellate *Pfiesteria shumwayae* in a semi-defined medium. J. Eukaryot. Microbiol. *1*, 73-82.

Spiliotis, E. T. (2010). Regulation of microtubule organization and functions by septin GTPases.
Cytoskeleton (Hoboken) *6*, 339-345.

Spiliotis, E. T. and Gladfelter, A. S. (2012). Spatial guidance of cell asymmetry: septin GTPases
show the way. Traffic *2*, 195-203.

Spiliotis, E. T., Hunt, S. J., Hu, Q., Kinoshita, M. and Nelson, W. J. (2008). Epithelial polarity
requires septin coupling of vesicle transport to polyglutamylated microtubules. J. Cell Biol. *2*,
295-303.

Stajich, J. E., Berbee, M. L., Blackwell, M., Hibbett, D. S., James, T. Y., Spatafora, J. W. and
Taylor, J. W. (2009). The fungi. Curr. Biol. *18*, 840.

Straight, A. F., Marshall, W. F., Sedat, J. W. and Murray, A. W. (1997). Mitosis in living budding
yeast: anaphase A but no metaphase plate. Science *5325*, 574-578.

Suga, H., Dacre, M., de Mendoza, A., Shalchian-Tabrizi, K., Manning, G. and Ruiz-Trillo, I.
(2012). Genomic survey of premetazoans shows deep conservation of cytoplasmic tyrosine
kinases and multiple radiations of receptor tyrosine kinases. Sci. Signal. *222*, ra35.

Suga, H. and Ruiz-Trillo, I. (2013). Development of ichthyosporeans sheds light on the origin of
metazoan multicellularity. Dev. Biol. *1*, 284-292.

Symons, M. H. and Mitchison, T. J. (1991). Control of actin polymerization in live and
permeabilized fibroblasts. J. Cell Biol. *3*, 503-513.

Takizawa, P. A., DeRisi, J. L., Wilhelm, J. E. and Vale, R. D. (2000). Plasma membrane
compartmentalization in yeast by messenger RNA transport and a septin diffusion barrier.
Science *5490*, 341-344.

Tanaka-Takiguchi, Y., Kinoshita, M. and Takiguchi, K. (2009). Septin-mediated uniform bracing
of phospholipid membranes. Curr. Biol. *2*, 140-145.

Tooley, A. J., Gilden, J., Jacobelli, J., Beemiller, P., Trimble, W. S., Kinoshita, M. and Krummel,
M. F. (2009). Amoeboid T lymphocytes require the septin cytoskeleton for cortical integrity and
persistent motility. Nat. Cell Biol. *1*, 17-26.

Vinayak, S., Pawlowic, M. C., Sateriale, A., Brooks, C. F., Studstill, C. J., Bar-Peled, Y.,
Cipriano, M. J. and Striepen, B. (2015). Genetic modification of the diarrhoeal pathogen
Cryptosporidium parvum. Nature *7561*, 477-480.

Wang, M. and Casey, P. J. (2016). Protein prenylation: unique fats make their mark on biology.
Nat. Rev. Mol. Cell Biol. *2*, 110-122.

Wilson, K. S., Gonzalez, O., Dutcher, S. K. and Bayly, P. V. (2015). Dynein-deficient flagella
respond to increased viscosity with contrasting changes in power and recovery strokes.
Cytoskeleton (Hoboken) *9*, 477-490.

Woznica, A., Cantley, A. M., Beemelmanns, C., Freinkman, E., Clardy, J. and King, N. (2016).
Bacterial lipids activate, synergize, and inhibit a developmental switch in choanoflagellates.
Proc. Natl. Acad. Sci. U. S. A. *28*, 7894-7899.

Woznica, A., Gerdt, J. P., Hulett, R. E., Clardy, J. and King, N. (2017). Mating in the Closest
Living Relatives of Animals Is Induced by a Bacterial Chondroitinase. Cell *6*, 1183.e11.

Yang, T. T., Cheng, L. and Kain, S. R. (1996). Optimized codon usage and chromophore
mutations provide enhanced sensitivity with the green fluorescent protein. Nucleic Acids Res.
22, 4592-4593.

Young, S. L., Diolaiti, D., Conacci-Sorrell, M., Ruiz-Trillo, I., Eisenman, R. N. and King, N.
(2011). Premetazoan ancestry of the Myc-Max network. Mol. Biol. Evol. *10*, 2961-2971.

 FIGURE LEGENDS

Figure 1: Introduction to *Salpingoeca rosetta*, an experimentally-tractable model

choanoflagellate

(A) *S. rosetta* and other choanoflagellates (abbreviated as 'choano') are the closest living

relatives of animals. **(B, C)** *S. rosetta* has a complex life history that includes single cells **(B)** and

multicellular rosettes **(C)**. Immunofluorescence in fixed, permeabilized *S. rosetta* single cells **(B)**

highlights the diagnostic cellular architecture of choanoflagellates, including a single apical

flagellum (f) made of microtubules (white) surrounded by a collar (co) filled with F-actin (red) of

microvilli. Staining for tubulin also illuminates cortical microtubules (cm) that run in parallel

tracks along the cell periphery from the apical to the basal poles of each cell. DNA staining

(blue) highlights the choanoflagellate nucleus (n) and the nucleoids of bacterial prey (b) present

in choanoflagellate cultures. **(C)** In multicellular rosettes (stained as in panel **B**), the basal poles

of cells are oriented toward the interior of the rosette and the apical flagella point outward.

Figure 2: A robust procedure for transfecting *S. rosetta*

(A) A summary of the step-wise procedure to transfect *S. rosetta* with DNA plasmids. To

prepare *S. rosetta* for transfection, cells were harvested at mid-log phase and then washed to

remove bacteria (depicted as grey ovals). *S. rosetta* cells (depicted with an apical collar,

flagellum, and nucleus; n) were primed for nucleofection (step 1) through washing with a buffer

that degrades extracellular material. A DNA plasmid encoding a highly sensitive luciferase,

nanoluc, or a fluorescent protein was then transfected into the nucleus with a nucleofector (step

2). Immediately after transfection, the cells rested in a buffer that promotes membrane closure

(step 3). Finally, the cells were transferred into 1x High Nutrient Media prepared with AK

seawater for two days (step 4) before we assayed the expression of nanoluc or fluorescent

proteins from the transfected DNA. **(B)** Non-coding DNA sequences flanking the coding

sequences for *S. rosetta elongation factor L* (pEFL), *α-tubulin* (pTub), *β-actin* (pAct), and

histone H3 (pH3) genes drive the expression of a codon-optimized *nanoluc* reporter gene. 2.5

µg of pEFL-*nanoluc,* pTub-*nanoluc,* pAct-*nanoluc*, and pH3-*nanoluc* reporter plasmids were

each transfected into *S. rosetta* and the cells were subsequently assayed for luciferase

expression. Each reporter produced a luminescence signal that was at least three orders of

magnitude greater than the detection limit (dotted line) and significantly greater (one-way

ANOVA, $p < 0.001$) than the background from a negative control, in which cells were
transfected with an empty pUC19 vector (None). See Materials and Methods for details on
replicates and statistical tests. **(C)** Systematically omitting each step of the transfection
procedure revealed critical steps for the delivery and expression of plasmid DNA in *S. rosetta*
cells. As a baseline for comparison, cells with 2.5 *µ*g of pH3-*nanoluc* reporter (row b) produced
a luciferase signal that was three orders of magnitude greater than the background detected
from cells transfected without the reporter plasmid (row a). Omitting the priming step by
incubating cells in artificial seawater instead of priming buffer (row c), decreased luciferase
signal by over two orders of magnitude. Nucleofection without carrier DNA (row d) or the
application of the CM156 electrical pulse (row e) resulted in a complete loss of luciferase signal,
indicating that both were essential for successful transfection. Directly transferring cells to sea
water after nucleofection instead of a buffer that promotes membrane resealing during the rest
step (row f) decreased the luciferase signal almost ten-fold. Finally, despite the fact that most
prey bacteria were washed out prior to nucleofection, addition of fresh prey bacteria did not
appear to be necessary. Supplementing transfected cells with fresh prey bacteria at the start of
the recovery step had seemingly little effect on transfection success (row g), probably due to the
persistence of a small number of live bacteria throughout the nucleofection procedure. **(D and**
E) Fluorescent reporters mark transfected cells. Live cells transfected with a pAct-m*Wasabi*
reporter construct could be observed by fluorescence microscopy **(D)** and quantified by flow
cytometry **(E)**. Untransfected cells were used to draw a gate that includes 99.99% of cells, or
four-standard deviations above the mean fluorescence value (left). The same gate was applied
to a population of transfected cells (right) to categorize the mWasabi- population. Cells with
higher values of green fluorescence that laid outside of the mWasabi- gate are categorized as
mWasabi+. The efficiency of transformation, as quantified by three independent flow cytometry
experiments, was ~1% in a population of 1 million cells.
950
Figure 3: Fluorescent markers illuminate the cell biology of *S. rosetta* in live cells
Fluorescent subcellular markers expressed from reporter plasmids in live *S. rosetta* cells were
constructed by fusing *mCherry* in frame to genes encoding localization peptides and proteins
(Datasets S1 and S3). Twenty-four hours after co-transfecting cells with 5 *µ*g of a plasmid
encoding a subcellular marker fused to the mCherry protein and 5 *µ*g of a plasmid encoding
untagged mTFP1 that served as a whole cell marker, live cells were visualized by
superresolution microscopy with a Zeiss LSM 880 Airyscan. The variation in localization of the

whole cell mTFP1 marker stems from cell-to-cell differences in the number and localization of

vacuoles, which exclude mTFP1. In panels **A - I**, the cells are oriented with the apical flagellum

at the top and the nucleus, when included in the plane of focus (**A'' - F''),** is indicated with a

dotted white line. **(A)** Without localization signals (None), fluorescent proteins (mCherry, **A'**, and

mTFP1, **A''**) were distributed throughout the cell with a slight enrichment in the nucleus and

complete exclusion from other membrane bound compartments. **(B and C)** A fusion of mCherry

to the carboxy terminus of Histone H3 **(B')** or the amino terminus of a simian virus 40 nuclear

localization signal **(**NLS; **C')** was confined to the nucleus, whereas mCherry fused to the

carboxy terminus of elongation factor L **(**EFL; **D)** was excluded from the nucleus and restricted

to the cytosol. **(E)** The endoplasmic reticulum (ER) was highlighted by fusing the signal

sequence from Rosetteless (PTSG_03555) and an ER retention sequence (HDEL from

PTSG_07223) to the amino and carboxy termini of mCherry, respectively. **(F)** The mitochondrial

network was highlighted by fusing a targeting sequence from *S. cerevisiae* CoxIV to the amino

terminus of mCherry. **(G)** A Lifeact peptide fused to the amino terminus of mCherry marked

filamentous actin (F-actin) that forms filipodia (arrowhead) and actin filaments in the cell body

that coalesce to form the collar (arrow). **(H)** Fusing mCherry to the amino terminus of α-Tubulin

highlighted parallel tracks of microtubules (arrowhead) that extended subcortically from the

apical pole to the basal pole of cells and microtubules that emerged from the apical pole of the

cell body to form the flagellum. The flagellum undulates rapidly in live cells and can be difficult to

image in total; in this cell the most distal tip of the flagellum is captured in the plane of focus

(arrow). **(I)** A plasma membrane marker constructed by fusing a geranyl-geranylation sequence

(PTSG_00306) to the carboxy terminus of mCherry outlined the entire cell shape, including the

collar, flagellum, and cell body. The membrane marker also weakly highlighted the Golgi

(arrowhead). The food vacuole (asterisk) was often visualized due to autofluorescence from

ingested bacteria or through accumulation of the fluorescent markers in the food vacuole,

perhaps through autophagy. **(J - L)** Orthogonal views along the xy and xz axes from confocal

micrographs showed fine details of cell architecture that were highlighted by transfecting cells

with F-actin, microtubule, and plasma membrane markers fused to mCherry (magenta). In xz

views, each cell is oriented with the flagellum facing toward the top of the micrograph; the

flagella appeared shorter and blurred because of the sigmoidal shape of the flagellar beat.

Lifeact **(J)** and the plasma membrane **(L)** markers fused to mCherry showed the microvilli

(arrowheads). **(K)** The α-tubulin-mCherry showed the subcortical tracks of microtubules at the

cell periphery (arrowhead) and the microtubule organizing center (arrow).

Figure 4: Septins assemble at the basal pole of *S. rosetta* cells

(A) *Sr*Septin2 has a prototypical protein domain architecture of septins, with an amino-terminal Septin G-domain that mediates filament formation and a carboxy terminal coiled-coil domain that mediates higher order assemblies of septin filaments. To investigate the localization of *Sr*Septin2, we engineered fusions with mTFP1 at the amino terminus and created a truncation of the coiled-coil domain (ΔCC). **(B)** A mTFP1-*Sr*Septin2 fusion protein localized to the basal pole of unicellular cells (**B'**, arrowhead). Co-transfecting cells with mTFP1-*Sr*Septin2 and a plasma membrane marker revealed *Sr*Septin2 distributed throughout the cytosol and enriched at the basal pole in confocal slices through the center of the cell. **(C)** mTFP1-*Sr*Septin6 mirrored the enrichment of mTFP1-*Sr*Septin2 at the basal pole (**C'**, arrowhead). The overlapping localization of *Sr*Septin2 and *Sr*Septin6 was compatible with these proteins forming heteromeric filaments with each other and other septin paralogs. **(D)** Consistent with the coiled-coil domain mediating the localization of septins through the formation of higher-order structures, *Sr*Septin2ΔCC localized throughout the cytoplasm, with no visible enrichment at the basal pole. Surprisingly, the deletion also caused ectopic filaments (**D'**; arrowheads) to form around membrane-bound vesicles that were, based on their size and position in the cell, presumably food vacuoles. **(E)** In rosettes, mTFP1-*Sr*Septin2 localized to points of cell-cell contact corresponding to the basal poles of cells (**E'**; arrowhead). **(F)** As in single cells, mTFP1-*Sr*Septin2ΔCC in rosettes was distributed throughout the cytosol and formed ectopic filaments (**F'**; arrowheads) around vacuoles. In panels **E** and **F**, *S. rosetta* single cells were transfected as in panels **B** and **C**, immediately induced to develop into rosettes (Woznica *et al.*, 2016), and imaged the next day. **(G)** *Sr*Septin2 intercalated between microtubules at the basal pole of the cell. Co-transfecting cells with mTFP1-*Sr*Septin2 and the α-tubulin marker showed *Sr*Septin2 filaments intercalated between microtubules at the basal pole in confocal slices that capture the cell cortex to easily visualize microtubule tracks. (**G'**, **G''**, **G'''**; box). **G''''** shows a 4x magnification of the basal pole of a representative cell (boxed region from **G'**, **G''**, **G'''**). In panels B-F, autofluorescence from ingested bacteria or through accumulation of the fluorescent markers highlights the food vacuole (asterisk).

Figure 1

Introduction to *Salpingoeca rosetta*, an experimentally-tractable model choanoflagellate

Figure 2

A robust procedure for transfecting *S. rosetta*

Figure 3

Fluorescent markers illuminate the cell biology of *S. rosetta* in live cells

Marker | **Whole Cell** | **Merge**

None — A′, A″, A‴
H3 — B′, B″, B‴
NLS — C′, C″, C‴
EFL — D′, D″, D‴
Endoplasmic Reticulum — E′, E″, E‴
Mitochondria — F′, F″, F‴

Marker | **Whole Cell** | **Merge**

Lifeact — G′, G″, G‴
Tubulin — H′, H″, H‴
Plasma Membrane — I′, I″, I‴

2 μm

Lifeact | **Tubulin** | **Plasma Membrane**

xy — J′, K′, L′

xz — J″, K″, L″

2 μm

Figure 4

Septins assemble at the basal pole of *S. rosetta* cells