**Title:** Natural *Wolbachia* infections are common in the major malaria vectors in

Central Africa

**Running title:** *Wolbachia* diversity in African *Anopheles*

**Authors**

Diego Ayala[1,2,*], Ousman Akone-Ella[2], Nil Rahola[1,2], Pierre Kengne[1], Marc F.

Ngangue[2,3], Fabrice Mezeme[2], Boris K. Makanga[2], Carlo Costantini[1], Frédéric

Simard[1], Franck Prugnolle[1], Benjamin Roche[1,4], Olivier Duron[1] & Christophe Paupy[1].

**Affiliations**

[1] MIVEGEC, IRD, CNRS, Univ. Montpellier, Montpellier, France.

[2] CIRMF, Franceville, Gabon.

[3] ANPN, Libreville, Gabon

[4] UMMISCO, IRD, Montpellier, France.

[*] **Corresponding author:**

Diego Ayala, MIVEGEC, IRD, CNRS, Univ. Montpellier, 911 av Agropolis, BP
64501, 34394 Montpellier, France; phone: +33(0)4 67 41 61 47; email:
diego.ayala@ird.fr
21

**Abstract**

During the last decade, the endosymbiont bacterium *Wolbachia* has emerged as a biological tool for vector disease control. However, for long time, it was believed that *Wolbachia* was absent in natural populations of *Anopheles*. The recent discovery that species within the *Anopheles gambiae* complex hosts *Wolbachia* in natural conditions has opened new opportunities for malaria control research in Africa. Here, we investigated the prevalence and diversity of *Wolbachia* infection in 25 African *Anopheles* species in Gabon (Central Africa). Our results revealed the presence of *Wolbachia* in 16 of these species, including the major malaria vectors in this area. The infection prevalence varied greatly among species, confirming that sample size is a key factor to detect the infection. Moreover, our sequencing and phylogenetic analyses showed the important diversity of *Wolbachia* strains that infect *Anopheles*. Co-evolutionary analysis unveiled patterns of *Wolbachia* transmission within *Anopheles* species, suggesting that past independent acquisition events were followed by co-cladogenesis. The large diversity of *Wolbachia* strains that infect natural populations of *Anopheles* offers a promising opportunity to select suitable phenotypes for suppressing *Plasmodium* transmission and/or manipulating *Anopheles* reproduction, which in turn could be used to reduce the malaria burden in Africa.

**Key-words**: *Anopheles*, *Wolbachia*, diversity, co-evolution, disease control.

## Introduction

Malaria still affects millions of people and is the cause of thousands of deaths worldwide, although sub-Saharan Africa pays the highest tribute [1]. Currently, vector-control measures (e.g., insecticide-treated bed nets or indoor residual spray) are the largest contributors to malaria eradication [2]. If these interventions are maintained or increased, malaria burden should be drastically reduced in Africa before 2030 [3]. These predictions are based on the constant effectiveness of these methods. However, the spread of insecticide resistance [4] and vector behavioural changes related to the massive use of bed nets [5] might challenge malaria eradication in the coming decades. Therefore, it is vital to develop alternative and environmentally friendly control strategies for the millennium development goal of malaria eradication [6].

Several methods have been proposed to accompany or replace the use of synthetic insecticides [7]. Among them, the use of the maternally inherited *Wolbachia* bacteria (α-proteobacteria, Anaplasmataceae family) has emerged as a promising alternative biological tool for fighting malaria and other vector-borne diseases [7-11]. This bacterium exhibits a large spectrum of interactions with its hosts: from mutualism and commensalism to parasitism [12]. Moreover, *Wolbachia* can invade mosquito populations and/or prevent vector-borne infections in some of the most important mosquito vectors [8,11,13]. Indeed, *Aedes aegypti* populations that were artificially infected with *Wolbachia* have been successfully used to suppress dengue transmission in laboratory conditions, and have been released in natural populations of this mosquito [14,15]. Similarly, laboratory studies showed that infection of *Anopheles* (the vector of human malaria) with *Wolbachia* strains has a negative impact on the transmission of *Plasmodium* parasites [16-18], providing a relevant alternative for

malaria control. Unfortunately, only one stable transfected *Wolbachia* colony has

been described in *Anopheles stephensi* [18]. Therefore, data on the use *Wolbachia* for

*Anopheles* control remain scarce and mainly concern experimental studies in

laboratory conditions [18,19], due to technical (i.e., egg microinjection) and biological

(i.e., competitive exclusion with the bacterium *Asaia*) difficulties to carry out

transinfections in *Anopheles*, despite multiple assays [20-22]. For long time, it was

believed that *Wolbachia* was absent in natural populations of *Anopheles* [20]. However,

in the last few years, three studies reported that *An. gambiae, An. coluzzii* and *An.*

*arabiensis* (three major malaria vectors) populations from Burkina Faso and Mali

(West Africa) are naturally infected by *Wolbachia* [23-25]. Noteworthy, they showed a

negative correlation between *Wolbachia* infection and *Plasmodium* development [24,25].

Moreover, a very recent report suggests that other *Anopheles* species also are infected

with *Wolbachia*[22]. These findings support the development of novel vector control

strategies based on *Wolbachia-Anopheles* interactions. However, although *Wolbachia*

naturally infects 40% - 60% of arthropods [26,27], infection of *Anopheles* species is still

not well documented. Moreover, during the last decade, screens in many other malaria

mosquito species worldwide (n=38) did not bring any evidence of *Wolbachia*

infection [9,20,28].

In this study, we investigated the presence of *Wolbachia* in 25 *Anopheles* species in

Gabon, Central Africa. We sampled mosquitoes across the country and in a variety of

ecological settings, from deep rainforest to urban habitats. By using a molecular

approach, we confirmed *Wolbachia* presence in 16 species, including all the major

malaria vectors in Central Africa (*An. gambiae, An. coluzzii, An. funestus, Anopheles*

*nili* and *Anopheles moucheti*). The prevalence of *Wolbachia* infection was particularly

high in *An. nili* and *An. moucheti*. Phylogenetic analysis revealed that all the infected

mosquito species hosted *Wolbachia* bacteria belonging to the supergroup A or B (both

exhibit high genetic diversity). Finally, we explored the co-evolution between

*Wolbachia* and *Anopheles*. The results have direct implications for the development of

new and environmentally friendly vector control strategies and open new directions

for research on pathogen transmission and reproductive manipulation.

**Results**

***Wolbachia naturally infects a large number of Anopheles species from Gabon.***

In this study, we screened 648 mosquitoes from eight sites in Gabon (Fig. 1, Table 1,

Table S1). On the basis of their morphological traits [29] and molecular analysis results

[30-34], we identified 25 *Anopheles* species (Text S1). Our sample included all the

species in which the presence of *Wolbachia* was previously investigated in Africa

(*An. gambiae, An. coluzzii, An. funestus* and *An. coustani)*, with the exception of *An.*

*arabiensis* that is absent in Gabon (Table 1)[35]. By PCR amplification of a *16S* rRNA

fragment [24], we found 70 *Wolbachia*-positive specimens that belonged to 16 different

*Anopheles* species, distributed throughout the country (Fig. 1, Table S1). With few

exceptions, *Wolbachia* infection rate was lower than 15% in most species (n=11), as

observed in other arthropods [26,27]. On the other hand, for *An. moucheti, An. "GAB-2",*

*An. "GAB-3", An. jebudensis* and *An. nili,* more than 50% of sampled mosquitoes

were infected (Table 1), as previously reported in other mosquito species where

prevalence can be very high [36,37]. To our knowledge, none of these species was

previously screened for *Wolbachia* infection.

***Wolbachia is maternally inherited in An. moucheti****.*

Although *Wolbachia* is mainly maternally transmitted [12], horizontal transmission may

occasionally occur in natural conditions [38-40]. To confirm the maternal transmission in

the infected mosquito species, we focused on *An. moucheti* for logistic reasons (i.e.,

highest *Wolbachia* prevalence and ease of sampling). Although no laboratory *An.*

*moucheti* strain is currently available, we obtained eggs from six *Wolbachia*-infected

females. In total, we analysed the infectious status of 79 progeny by PCR

amplification of the same *16S* rRNA fragment [24] (Table S2), and found that 70 were

infected, with an average maternal transmission frequency of 97.54% (range: 90% -

100%).

**Naturally occurring *Wolbachia* strains in *Anopheles* reveal high genetic diversity**.

By sequence analysis of the *16S* rRNA fragment PCR-amplified from each *Anopheles*

sample (Table 1), we could assign the *Wolbachia* strains to three pre-existing

supergroups: A (n=5), B (n=64) and C (n=1) (Fig. 2). Specifically, we detected

supergroup B *Wolbachia* in 64 mosquitoes belonging to all 16 infected *Anopheles*

species. We found supergroup A *Wolbachia* in five individuals from four species (*An.*

*funestus, An. coluzzii, An. vinckei* and *An. carnevalei*), thus providing examples of

multiple infections, as previously observed in *Ae. albopictus* [41] (Fig. 2). None of the

mosquitoes examined was co-infected by *Wolbachia* strains belonging, for instance,

to the supergroups A and B. Moreover, we confirmed that the *Wolbachia* strains

previously identified in *An. gambiae s.l.* from Burkina Faso and Mali are included in

the supergroups A and B [23,25]. Finally, we found that one *An. coustani* individual was

infected by a *Wolbachia* strain from supergroup C that is known to infect only filarial

worms. Therefore, we investigated the presence of filarial nematode DNA in the

mosquito by PCR amplification and sequencing of a fragment of the *COI* filarial gene

[42], followed by phylogenetic analysis with RAxML. Our results confirmed the

presence of *Dirofilaria immitis* in this specimen (Fig. S1). This canine filarial parasite

hosts *Wolbachia* and is transmitted by many mosquitoes, including *Anopheles* [43].

Therefore, it is not surprising to find an *An. coustani* specimen infected by this filarial

nematode. This specimen was excluded from further investigations.

To expand our knowledge on the *Wolbachia* strains that infect natural *Anopheles*

populations, we PCR amplified, sequenced and analysed a *16S* rRNA fragment and

also fragments from three other conserved *Wolbachia* genes (*coxA, fbpA* and *ftsZ*) that

are commonly used for strain typing and evolutionary studies [44] (Fig. 2). We used a

new nested PCR protocol (see *Methods*) for samples that could not be genotyped

using the classical Multi-Locus Sequence Typing (MLST) primers (Table S1). Our

phylogenetic analyses confirmed the *16S* results, assigning most of the species to

supergroups A and B. Few samples (asterisks in Fig. 2, gene *coxA*) showed some

incongruence relative to the *16S* results. They suggest signals of recent recombination

between the supergroups A and B, as previously demonstrated [44]. Detailed sequence

analysis revealed that mosquito species belonging to the same group or complex (i.e.,

*An. moucheti* and *An. gambiae*) displayed a common *Wolbachia* haplotype (defined

here as a unique allelic profile) (Fig. 2 and Fig. 3). Conversely, some species with

lower prevalence (i.e., *An. coluzzii, An. marshallii, An. vinckei* or *An. funestus*)

displayed a variety of haplotypes. The case of *An. vinckei* was particularly interesting

because the three infected specimens displayed different haplotypes for the analysed

*Wolbachia* genes. Moreover, one specimen (*An. vinckei* M002, Fig. 2) was infected

by a completely different *Wolbachia* strain. Overall, the *Wolbachia* haplotypes

identified in this study were different from the allelic profiles of the previously

annotated *Wolbachia* strains or of the strain that infects *An. gambiae* in Burkina Faso

and Mali [23,25] (Fig. 2 and Fig. 3). Within supergroup B, we could easily distinguish at

least two strains. The strain infecting *An. moucheti* (*wANMO*) was similar to the one

identified in *An. gambiae* (in our study) or *An. marshallii*, while the strain infecting

*An. nili* (*wANNI*) was more closely related to those found in other mosquito species,

such as *Ae. albopictus* or *Cx. quinquefasciatus* (Fig. 2 and Fig. 3). Conversely, the

other haplotypes were associated with one specific host.

***Wolbachia* independently evolves in malaria-transmitting mosquitoes.**

As *Wolbachia* is mainly a maternally inherited bacterium, the host mitochondrial

DNA (mtDNA) is a suitable marker to study its evolutionary history in *Anopheles* [45].

Analysis of *COII* sequences from 176 specimens belonging to the 25 *Anopheles*

species collected in Gabon provided the most exhaustive phylogenetic tree of

*Anopheles* in Central Africa (Fig. 3). This analysis highlighted the independent

acquisition and loss of *Wolbachia* across the different *Anopheles* species clades.

Moreover, the genetic distances of *Wolbachia* strains and their *Anopheles* host were

not correlated (Mantel test, p >0.05) (Fig. S2). Nevertheless, mosquitoes from the

same complex, and therefore genetically very close, shared the same *Wolbachia*

supergroup and haplotypes (Fig. 3 and Fig. S2). Finally, we investigated how

*Wolbachia* evolved within each *Anopheles* species [46]. Our results revealed that

*Wolbachia*-infected and non-infected mosquitoes shared the same mtDNA haplotype

(Fig. 2), indicating that infection status and host haplotypes are not associated.

**Discussion**

The present study provides three key conclusions. First, the genus *Anopheles* includes

a large number of species that are naturally infected by *Wolbachia* (16/25), with high

infection prevalence among major malaria vectors. Second, *Anopheles*-infecting

*Wolbachia* bacteria show high genetic diversity, with similar haplotypes detected in

different *Anopheles* species. Third, the independent evolution of *Wolbachia* and

*Anopheles* might be interpreted as multiple acquisition events with horizontal

transmission. The large diversity of *Wolbachia* strains that infect many natural

*Anopheles* populations could represent a major opportunity for reducing pathogen

transmission and/or for reproductive manipulation in *Anopheles* with the aim of

decreasing malaria burden in Africa.

For long time, the scientific community has been interested in finding new ways to

use *Wolbachia* for fighting vector-borne diseases [7-9,11]. In arthropods, *Wolbachia* is

broadly spread, including among *Culex* and *Aedes* mosquitoes. Conversely, it was

believed that in the genus *Anopheles*, *Wolbachia* infections were mainly limited to

species within the gambiae complex [23,25], and few other species [22]. Several hypotheses

can be put forward to explain this. First, low infection prevalence or local variations

could have hindered the discovery of *Wolbachia* infections. In our study, most

*Anopheles* species exhibited a prevalence lower than 15% (Table 1). This pattern is

common in many other arthropods [26,27], and it is usually associated with a weak

manipulation of the host reproduction and/or imperfect maternal transmission [47].

However, high *Wolbachia* prevalence rates have been reported for some mosquitoes

species, such as *Culex pipiens*[36] or *Aedes albopictus*[48]. In general, our sampling effort

was higher than in previous studies (n <30) [9,28], and this could explain why we found

more infected species. Our statistical analysis showed that a sample size of 60

individuals per species is needed to quantify correctly prevalence rates lower than

15%, with a probability of 95% (Fig. S3). Moreover, local frequency variations

among populations could also hinder the detection of *Wolbachia* infections [36]. For

instance, we sampled *An. coluzzii* in three different sites, but we only found

*Wolbachia*-infected mosquitoes at La Lopé (Fig 1, Table S1). Therefore, sampling in

different localities and in different seasons might improve detection rates. Second, it

could be difficult to detect low-density *Wolbachia* infections in *Anopheles* with the

routinely used molecular tools, as previously reported for other arthropods [49,50] and

recently in *Anopheles gambiae* [25]. Our results indicate that conventional PCR

amplification (*wsp*-targeting primers [44]) analysis allowed detecting *Wolbachia*

infection only in 6 of the 16 species (*An. moucheti, An m. nigeriensis, An. "GAB-3",*

*An nili, An. jebudensis* and *An. vinckei*) under study, presumably because of the high

*Wolbachia* density. Moreover, some *Anopheles* species with high *Wolbachia* infection

rates, such as *An. moucheti* or *An. nili*, were never screened before.

Unexpectedly, our work revealed that *Anopheles* species are infected by different

*Wolbachia* strains. Although three previous studies reported *Wolbachia* infection in

the *An. gambiae* germline [23-25], we do not know whether *Wolbachia* naturally invades

and is maternally inherited in the infected mosquito species from Gabon. Indeed,

horizontal gene transfer (resulting in the insertion of *Wolbachia* genes within the

mosquito genome), or parasitism (e.g., by filarial nematodes) could explain the

detection of *Wolbachia* genes in an organism without maternal transmission.

However, the *Wolbachia* sequences we identified were genetically close to those

found in other Diptera, and we did not observe any signal of extensive divergence,

which would be expected in the case of horizontal gene transfer [51,52] (Figs 2 and 3).

243 Alternatively, mosquitoes could have been parasitized by *Wolbachia*-hosting

244 nematodes [53,54] or mites [55]. However, these *Wolbachia* strains belong to other, easily

245 distinguishable supergroups (Fig. 2 and Fig. S1). Moreover, the analysis of *An.*

246 *moucheti* F1 progeny confirms, at least in this species, that no other biological

247 *Wolbachia* contamination was present in our analysis. In conclusion, our data suggest

248 that *Wolbachia* is naturally present in the *Anopheles* species of Central Africa

249 analysed in our study, and that is maternally inherited in *An. moucheti* (Table S2).

251 In Central African *Anopheles*, *Wolbachia* acquisition seems to be independent of the

252 host phylogeny (Figs 2 and 3). Our results revealed that the genetic distances between

253 *Wolbachia* and *Anopheles* are not positively correlated (Mantel test, $p > 0.05$) (Fig.

254 S2). Therefore, *Wolbachia* and the host lineage seem to evolve independently. The

255 different larval ecology of these species suggests other ways of lateral transfer (e.g.,

256 during nectar feeding [39]). On the other hand, we found that all the *Anopheles* species

257 belonging to the same complex shared related *Wolbachia* strains (Fig. 3). Permeable

258 reproductive barriers among members of the same complex could facilitate the

259 intermittent movement of the bacterium [56]. Interestingly, although they share similar

260 *Wolbachia* strains, sibling species showed different infection prevalence. Indeed, *An.*

261 *carnevalei* and *An. m. nigeriensis* exhibited frequencies lower than 15%, whereas *An.*

262 *nili* and *An. moucheti*, their respective counterparts and the most important malaria

263 vectors in their complex, displayed frequencies higher than 60% (Table 1). Moreover,

264 our *An. gambiae* and *An. coluzzii* populations were infected by different *Wolbachia*

265 strains than those detected in Burkina Faso and Mali. Similarly, in mosquitoes [36] and

266 ants [57], the same species is infected by different *Wolbachia* strains according to the

267 region. The availability of whole-genome sequences for *Wolbachia* strains [58] will

enlighten the intricate phylogenetic relationships among the different strains in

*Anopheles*.

**Conclusions**

*Wolbachia* has emerged as a biological tool for controlling vector-borne diseases [14,15].

In this study, we demonstrated the natural presence of this endosymbiont bacterium in

a large number of *Anopheles* species, including the five major malaria vectors in

Central Africa. It has been shown that *Wolbachia* ability to interfere with pathogen

transmission depends on the bacterium strain. Therefore, our results offer the

opportunity to determine whether the different *Anopheles*-infecting *Wolbachia* strains

affect *Plasmodium* transmission and/or *Anopheles* reproduction. Indeed, the three

most infected species (*An. moucheti*, *An. nili* and *An. vinckei*) play an important role

in human (the first two) and non-human malaria transmission in the deep forest of

Gabon [35,59]. Therefore, we could investigate *Wolbachia* positive [24] and negative [60,61]

effects on the susceptibility to *Plasmodium* infection in their natural hosts. Moreover,

the strongest effect on suppression of pathogen transmission or reproductive

manipulation has been observed in *Wolbachia* transinfections [16,18,62-66]. Therefore, the

availability of *Wolbachia* strains that infect natural *Anopheles* populations offers

promising opportunities for experimental and theoretical studies in *Anopheles*, and

also in other mosquito families that are vectors of other diseases, including *Ae.*

*aegypti* and *Ae. albopictus.* In conclusion, our findings are merely the "tip of the

iceberg" in the study of *Wolbachia* use to control vector-borne diseases, particularly

in malaria.

**Methods**

**Research and ethics statements**

Mosquitoes were collected in Gabon under the research authorization

AR0013/16/MESRS/CENAREST/CG/CST/CSAR and the national park entry

authorization AE16008/PR/ANPN/SE/CS/AEPN. Mosquito sampling using the

human-landing catch (HLC) method was performed under the protocol

0031/2014/SG/CNE approved by the National Research Ethics Committee of Gabon.

**Mosquito sampling and DNA extraction**

Mosquitoes were collected in eight sites across Gabon, Central Africa, from 2012 to

2016 (Fig. 1, Table 1, Fig. S1). These sites included sylvatic (national parks) and

anthropic habitats (villages and cities). Adult females were collected using CDC light

traps, BG traps and HLC. Collected specimens were taxonomically identified

according to standard morphological features [29,67]. Then, they were individually stored

in 1.5 mL tubes at -20°C and sent to CIRMF for molecular analysis. When possible,

at least 30 mosquitoes (from 1 to 58) for each *Anopheles* species from different sites

were selected for genomic analysis. Total genomic DNA was extracted from the

whole body using the DNeasy Blood and Tissue Kit (Qiagen), according to the

manufacturer's instructions. Genomic DNA was eluted in 100 μL of TE buffer.

Specimens belonging to the *An. gambiae* complex, *An. funestus* group, *An. moucheti*

complex and *An. nili* complex were molecularly identified using PCR-based

diagnostic protocols [30-33,68].

*Wolbachia* **screening and Multi Locus Sequence Typing (MLST) analysis**

*Wolbachia* infection in adult females was detected by nested PCR amplification of a

*Wolbachia*-specific *16S* rDNA fragment (~400 bp) using 2 μL of host genomic DNA,

according to the protocol developed in Catteruccia's laboratory [24]. Amplification of

this *16S* rDNA fragment in infected *Aedes albopictus* and *Culex pipiens* genomic

DNA (data not shown) confirmed the performance of this nested PCR protocol to

detect *Wolbachia* in many different mosquito species[24]. To detect potential

contaminations, *Ae. albopictus* and *Cx quinquefasciatus* from Gabon were used as

positive controls, and water and *Ae. Aegypti* as negative controls. Moreover, PCR

amplifications for each species were carried out independently and in different days.

The amplicon size was checked on 1.5% agarose gels, and amplified *16S* rDNA

fragments were sent to Genewiz (UK) for sequencing (forward and reverse) to

confirm the presence of *Wolbachia*-specific sequences. The DNA quality of all

samples was confirmed by the successful amplification of a fragment (~800 bp) of the

mitochondrial gene *COII* in all the *Anopheles* species under study [34,69]. PCR products

were run on 1.5% agarose gels, and *COII* fragments from 176 mosquito specimens of

the 25 species were sequenced (forward and reverse) by Genewiz (UK) for the

*Anopheles* phylogenetic studies. *Wolbachia*-positive genomic DNA samples (2

μL/sample) were then genotyped by MLST using three loci, *coxA (~450 bp) ftsZ*

*(~500 bp)* and *fbpA (~460 bp)* [70], and according to standard conditions [44]. If the three

fragments could not be amplified, a newly-developed nested PCR protocol was used.

Specifically, after the first run with the standard primers, 2 μL of the obtained product

was amplified again using internal primers specific for each gene: *coxA* (coxA_NF-2:

5'-TTTAACATGCGCGCAAAAGG-3';                    coxA_NR-2:                    5'-

TAAGCCCAACAGTGAACATATG-3'),          *ftsZ*          (ftsZ_NF-2:          5'-

ATGGGCGGTGGTACTGGAAC-3';                    ftsZ_NR-2:                    5'-

AGCACTAATTGCCCTATCTTCT-3'),          and          *fbpA*          (fbpA_NF-1:          5'-

AGCTTAACTTCTGATCAAGCA-3';                    fbpA_NR-1:                    5'-

TTCTTTTTCCTGCAAAGCAAG-3'). Cycling conditions for *coxA* and *ftsZ* were:

94°C for 5min, followed by 36 cycles at 94°C for 15s, 55°C for 15s and 72°C for 30s,

and a final extension step at 72°C for 10min. For *fbpA*, they were: 94°C for 5min

followed by 36 cycles at 94°C for 30s, 59°C for 45s and 72°C for 90s, and a final

extension step at 72°C for 10 min. The resulting fragments (*coxA*, 357bp; *fbpA*, 358-

bp; and *ftsZ*, 424-bp) were sequenced bidirectionally by Genewiz (UK). The new

sequences obtained in this study were submitted to GenBank (accession numbers XX

to XX, Supplementary Table 1). Unfortunately, the other three MLST genes (*gatB*,

*wsp* and *hcpA*) could not be amplified, due to technical problems (i.e. multiple bands)

*Phylogenetic and statistical analysis*

All *Wolbachia* sequences for the *16S, coxA, fbpA* and *ftsZ* gene fragments and for

*Anopheles* COII were corrected using *Geneious* R10 [71]. The resulting consensus

sequences for each gene were aligned with sequences that represent the main known

*Wolbachia* supergroups obtained from GenBank (see Table S1). Only unique

haplotypes for each species were included in the analysis (haplotype was defined as a

unique allelic profile for each examined locus). Inference of phylogenetic trees was

performed using the maximum likelihood (ML) method and RAxML [72] with a

substitution model GTR + CAT [73] and 1000 bootstrapping replicates. Finally, all

MLST *Wolbachi*a sequences were used to build phylogenetic trees using RAxML

(GTR+CAT model, 1000 bootstrapping replicates). Trees were visualized with iTOL

v.3.4.3 [74].

To quantify the accuracy of the observed *Wolbachia* infection prevalence, the

influence of sample size on its estimation was assessed. For this, it was assumed that

*Wolbachia* prevalence within a host species followed a beta binomial distribution [27]

yielding many species with a low or a high *Wolbachia* prevalence but few with an

intermediate one. This allowed quantifying, for each sample size, the proportion of

samples (over 1,000 realizations) that could yield an estimate that was not

significantly different from the prevalence over the whole population with a z-test and

a significance threshold at 95%. As expected, sample size could be small for very low

or very high prevalence (60 individuals are enough in 95% of cases for these extreme

prevalence values), while it was much higher for intermediate prevalence values (up

to 150 individuals for a prevalence value close to 50%).

All statistical analyses were performed using "R" v3.2.5 (R Development Core Team,

http://cran.r-project.org/), with the addition of the "ggplot2" library [75].

**Author contributions**

D.A., O.D. and C.P. designed the experiments. D.A., O.A., N.R. and P.K., performed

the experiments. D.A., F.S., C.C., O.D and C.P. analysed the data. D.A. performed the

sequencing analysis. D.A. and B.R. performed the statistical analysis. D.A., N.R.,

M.N., F.M., B.M., C.P. and F.P. provided samples for the analysis. D.A., O.D. and

C.P. wrote the manuscript.

**Competing interests**

The authors declare no competing (financial and non-financial) interests.

**Acknowledgements**

We thank the "Agence Nationale de la Preservation de la Nature" (ANPN) and the

"Centre National de la Recherche Scientifique et Technologique of Gabon"

(CENAREST) that authorized this study and facilitated the access to the national

parks of La Lopé, Moukalaba-Doudou and Plateaux Batékés. We specially thank

Sonia Lekana for helping with the preliminary sequencing analysis.

**References**

1 WHO. World Malaria Report 2015. (World Health Organization, 2015).
2 Bhatt, S. *et al.* The effect of malaria control on Plasmodium falciparum in
Africa between 2000 and 2015. *Nature* **526**, 207-+,
doi:10.1038/nature15535 (2015).
3 Griffi, J. T. *et al.* Potential for reduction of burden and local elimination of
malaria by reducing Plasmodium falciparum malaria transmission: a
mathematical modelling study. *Lancet Infect. Dis.* **16**, 465-472,
doi:10.1016/s1473-3099(15)00423-5 (2016).
4 Ranson, H. & Lissenden, N. Insecticide Resistance in African Anopheles
Mosquitoes: A Worsening Situation that Needs Urgent Action to Maintain
Malaria Control. *Trends in Parasitology* **32**, 187-196,
doi:10.1016/j.pt.2015.11.010 (2016).
5 Pates, H. & Curtis, C. in *Annual Review of Entomology* Vol. 50 *Annual*
*Review of Entomology* 53-70 (Annual Reviews, 2005).
6 Newby, G. *et al.* The path to eradication: a progress report on the malaria-
eliminating countries. *Lancet* **387**, 1775-1784 (2016).
7 McGraw, E. A. & O'Neill, S. L. Beyond insecticides: new thinking on an
ancient problem. *Nat Rev Micro* **11**, 181-193 (2013).
8 Hoffmann, A. A., Ross, P. A. & Rasic, G. Wolbachia strains for disease
control: ecological and evolutionary considerations. *Evolutionary*
*Applications* **8**, 751-768, doi:10.1111/eva.12286 (2015).
9 Bourtzis, K. *et al.* Harnessing mosquito-Wolbachia symbiosis for vector
and disease control. *Acta Tropica* **132**, S150-S163,
doi:10.1016/j.actatropica.2013.11.004 (2014).
10 Kambris, Z., Cook, P. E., Phuc, H. K. & Sinkins, S. P. Immune Activation by
Life-Shortening Wolbachia and Reduced Filarial Competence in
Mosquitoes. *Science* **326**, 134-136, doi:10.1126/science.1177531 (2009).
11 Iturbe-Ormaetxe, I., Walker, T. & Neill, S. L. O. Wolbachia and the
biological control of mosquito-borne disease. *Embo Reports* **12**, 508-518,
doi:10.1038/embor.2011.84 (2011).
12 Werren, J. H., Baldo, L. & Clark, M. E. Wolbachia: master manipulators of
invertebrate biology. *Nature Reviews Microbiology* **6**, 741-751,
doi:10.1038/nrmicro1969 (2008).
13 Dodson, B. L. *et al.* Wolbachia Enhances West Nile Virus (WNV) Infection
in the Mosquito Culex tarsalis. *Plos Neglect Trop D* **8**,
doi:10.1371/journal.pntd.0002965 (2014).
14 Hoffmann, A. A. *et al.* Successful establishment of Wolbachia in Aedes
populations to suppress dengue transmission. *Nature* **476**, 454-U107,
doi:10.1038/nature10356 (2011).

15 Schmidt, T. L. *et al.* Local introduction and heterogeneous spatial spread
of dengue-suppressing Wolbachia through an urban population of Aedes
aegypti. *PLOS Biology* **15**, e2001894, doi:10.1371/journal.pbio.2001894
(2017).
16 Hughes, G. L., Koga, R., Xue, P., Fukatsu, T. & Rasgon, J. L. Wolbachia
Infections Are Virulent and Inhibit the Human Malaria Parasite
Plasmodium Falciparum in Anopheles Gambiae. *Plos Pathogens* **7**,
doi:10.1371/journal.ppat.1002043 (2011).
17 Kambris, Z. *et al.* Wolbachia Stimulates Immune Gene Expression and
Inhibits Plasmodium Development in Anopheles gambiae. *Plos Pathogens*
**6**, doi:10.1371/journal.ppat.1001143 (2010).
18 Bian, G. *et al.* Wolbachia Invades Anopheles stephensi Populations and
Induces Refractoriness to Plasmodium Infection. *Science* **340**, 748-751,
doi:10.1126/science.1236192 (2013).
19 Hughes, G. L., Vega-Rodriguez, J., Xue, P. & Rasgon, J. L. Wolbachia Strain
wAlbB Enhances Infection by the Rodent Malaria Parasite Plasmodium
berghei in Anopheles gambiae Mosquitoes. *Applied and Environmental*
*Microbiology* **78**, 1491-1495, doi:10.1128/aem.06751-11 (2012).
20 Hughes, G. L. *et al.* Native microbiome impedes vertical transmission of
Wolbachia in Anopheles mosquitoes. *Proceedings of the National Academy*
*of Sciences* **111**, 12498-12503, doi:10.1073/pnas.1408888111 (2014).
21 Rossi, P. *et al.* Mutual exclusion of Asaia and Wolbachia in the
reproductive organs of mosquito vectors. *Parasites & Vectors* **8**,
doi:10.1186/s13071-015-0888-0 (2015).
22 Jeffries, C. L. *et al.* Novel Wolbachia strains in Anopheles malaria vectors
from Sub-Saharan Africa. *bioRxiv*, doi:10.1101/338434 (2018).
23 Baldini, F. *et al.* Evidence of natural Wolbachia infections in field
populations of Anopheles gambiae. *Nature Communications* **5**,
doi:10.1038/ncomms4985 (2014).
24 Shaw, W. R. *et al.* Wolbachia infections in natural Anopheles populations
affect egg laying and negatively correlate with Plasmodium development.
*Nature Communications* **7**, doi:10.1038/ncomms11772 (2016).
25 Gomes, F. M. *et al.* Effect of naturally occurring *Wolbachia* in *Anopheles*
*gambiae s.l.* mosquitoes from Mali on *Plasmodium falciparum* malaria
transmission. *PNAS* **114**, 12566-12571, doi:10.1073/pnas.1716181114
(2017).
26 Duron, O. *et al.* The diversity of reproductive parasites among arthropods:
Wolbachia do not walk alone. *Bmc Biology* **6**, doi:10.1186/1741-7007-6-
27 (2008).
27 Zug, R. & Hammerstein, P. Still a Host of Hosts for Wolbachia: Analysis of
Recent Data Suggests That 40% of Terrestrial Arthropod Species Are
Infected. *Plos One* **7**, doi:10.1371/journal.pone.0038544 (2012).
28 Osei-Poku, J., Han, C., Mbogo, C. M. & Jiggins, F. M. Identification of
Wolbachia Strains in Mosquito Disease Vectors. *PLOS ONE* **7**, e49922,
doi:10.1371/journal.pone.0049922 (2012).
29 Gillies, M. T. & Coetzee, M. C. *A Supplement to the Anophelinae of Africa*
*South of the Sahara (Afrotropical region).* (The South African Institute for
Medical Research, 1987).

30 Santolamazza, F. *et al.* Insertion polymorphisms of SINE200
retrotransposons within speciation islands of Anopheles gambiae
molecular forms. *Malar J* **7**, 163, doi:1475-2875-7-163 [pii]
10.1186/1475-2875-7-163 (2008).
31 Cohuet, A. *et al.* Species identification within the *Anopheles funestus* group
of malaria vectors in Cameroon and evidence for a new species. *American*
*Journal of Tropical Medicine and Hygiene* **69**, 200-205 (2003).
32 Kengne, P. *et al.* Molecular differentiation of three closely related
members of the mosquito species complex, *Anopheles moucheti*, by
mitochondrial and ribosomal DNA polymorphism. *Medical and Veterinary*
*Entomology* **21**, 177-182 (2007).
33 Kengne, P., Awono-Ambene, P., Nkondjio, C. A., Simard, F. & Fontenille, D.
Molecular identification of the *Anopheles nili* group of African malaria
vectors. *Medical    and    Veterinary    Entomology*    **17**,    67-74,
doi:doi:10.1046/j.1365-2915.2003.00411.x (2003).
34 Rahola, N. *et al.* Description of Anopheles gabonensis, a new species
potentially involved in rodent malaria transmission in Gabon, Central
Africa.    *Infection    Genetics    and    Evolution*    **28**,    628-634,
doi:10.1016/j.meegid.2014.05.012 (2014).
35 Makanga, B. *et al.* Ape malaria transmission and potential for ape-to-
human    transfers    in    Africa.    *PNAS*    **113**,    5329-5334,
doi:10.1073/pnas.1603008113 (2016).
36 Dumas, E. *et al.* Population structure of Wolbachia and cytoplasmic
introgression in a complex of mosquito species. *Bmc Evolutionary Biology*
**13**, doi:10.1186/1471-2148-13-181 (2013).
37 Duron, O. *et al.* Transposable element polymorphism of Wolbachia in the
mosquito Culex pipiens: evidence of genetic diversity, superinfection
andrecombination. *Mol Ecol* **14**, doi:10.1111/j.1365-294X.2005.02495.x
(2005).
38 Ahmed, M. Z., De Barro, P. J., Ren, S. X., Greeff, J. M. & Qiu, B. L. Evidence for
horizontal transmission of secondary endosymbionts in the Bemisia
tabaci    cryptic    species    complex.    *PLoS    One*    **8**,    e53084,
doi:10.1371/journal.pone.0053084 (2013).
39 Li, S. J. *et al.* Plant-mediated horizontal transmission of Wolbachia
between    whiteflies.    *The    ISME    journal*    **11**,    1019-1028,
doi:10.1038/ismej.2016.164 (2017).
40 Werren, J. H., Zhang, W. & Guo, L. R. Evolution and phylogeny of
Wolbachia: reproductive parasites of arthropods. *Proc Biol Sci* **261**, 55-63,
doi:10.1098/rspb.1995.0117 (1995).
41 Sinkins, S. P., Braig, H. R. & Oneill, S. L. WOLBACHIA SUPERINFECTIONS
AND THE EXPRESSION OF CYTOPLASMIC INCOMPATIBILITY. *Proceedings*
*of    the    Royal    Society    B-Biological    Sciences*    **261**,    325-330,
doi:10.1098/rspb.1995.0154 (1995).
42 Casiraghi, M., Anderson, T. J. C., Bandi, C., Bazzocchi, C. & Genchi, C. A
phylogenetic    analysis    of    filarial    nematodes:    comparison    with    the
phylogeny of Wolbachia endosymbionts. *Parasitology* **122**, 93-103,
doi:10.1017/s0031182000007149 (2001).

43 Simon, F. *et al.* Human and Animal Dirofilariasis: the Emergence of a
Zoonotic Mosaic. *Clinical Microbiology Reviews* **25**, 507-+,
doi:10.1128/cmr.00012-12 (2012).
44 Baldo, L. *et al.* Multilocus sequence typing system for the endosymbiont
Wolbachia pipientis. *Appl Environ Microbiol* **72**, doi:10.1128/aem.00731-
06 (2006).
45 Richardson, M. F. *et al.* Population Genomics of the Wolbachia
Endosymbiont in Drosophila melanogaster. *PLOS Genetics* **8**, e1003129,
doi:10.1371/journal.pgen.1003129 (2012).
46 Charlat, S. *et al.* The joint evolutionary histories of Wolbachia and
mitochondria in Hypolimnas bolina. *BMC Evolutionary Biology* **9**, 64,
doi:10.1186/1471-2148-9-64 (2009).
47 Engelstadter, J. & Hurst, G. D. D. The Ecology and Evolution of Microbes
that Manipulate Host Reproduction. *Annual Review of Ecology Evolution*
*and Systematics* **40**, 127-149,
doi:10.1146/annurev.ecolsys.110308.120206 (2009).
48 Kittayapong, P., Baimai, V. & O'Neill, S. L. Field prevalence of Wolbachia in
the mosquito vector Aedes albopictus. *American Journal of Tropical*
*Medicine and Hygiene* **66**, 108-111 (2002).
49 Arthofer, W., Riegler, M., Avtzis, D. N. & Stauffer, C. Evidence for low-titre
infections in insect symbiosis: Wolbachia in the bark beetle Pityogenes
chalcographus (Coleoptera, Scolytinae). *Environmental Microbiology* **11**,
1923-1933, doi:10.1111/j.1462-2920.2009.01914.x (2009).
50 Augustinos, A. A. *et al.* Detection and Characterization of Wolbachia
Infections in Natural Populations of Aphids: Is the Hidden Diversity Fully
Unraveled? *Plos One* **6**, doi:10.1371/journal.pone.0028695 (2011).
51 Woolfit, M., Iturbe-Ormaetxe, I., McGraw, E. A. & O'Neill, S. L. An Ancient
Horizontal Gene Transfer between Mosquito and the Endosymbiotic
Bacterium Wolbachia pipientis. *Mol. Biol. Evol.* **26**, 367-374,
doi:10.1093/molbev/msn253 (2009).
52 Dunning Hotopp, J. C. *et al.* Widespread lateral gene transfer from
intracellular bacteria to multicellular eukaryotes. *Science* **317**, 1753-
1756, doi:10.1126/science.1142490 (2007).
53 Robert, V., Ayala, D. & Simard, F. in *Entomologie médicale et vétérinaire*
(eds G. Duvallet, D. Fontenille, & V. Robert) (IRD Editions, 2017).
54 Clements, A. N. *The Biology of Mosquitoes: Development, Nutrition and*
*Reproduction v. 1.* (CABI Publishing, 1999).
55 Simmons, T. W. & Hutchinson, M. L. A Critical Review of All Known
Published Records for Water Mite (Acari: Hydrachnidiae) and Mosquito
(Diptera: Culicidae) Parasitic Associations From 1975 to Present. *Journal*
*of Medical Entomology* **53**, 737-752, doi:10.1093/jme/tjw052 (2016).
56 Pombi, M. *et al.* Dissecting functional components of reproductive
isolation among closely related sympatric species of the Anopheles
gambiae complex. *Evolutionary Applications* **10**, 1102-1120,
doi:10.1111/eva.12517 (2017).
57 Tsutsui, N. D., Kauppinen, S. N., Oyafuso, A. F. & Grosberg, R. K. The
distribution and evolutionary history of Wolbachia infection in native and
introduced populations of the invasive argentine ant (Linepithema

humile). *Molecular Ecology* **12**, 3057-3068, doi:10.1046/j.1365-
294X.2003.01979.x (2003).

58   Gerth, M., Gansauge, M. T., Weigert, A. & Bleidorn, C. Phylogenomic
analyses uncover origin and spread of the Wolbachia pandemic. *Nature*
*Communications* **5**, doi:10.1038/ncomms6117 (2014).

59   Paupy, C. *et al.* Anopheles moucheti and Anopheles vinckei Are Candidate
Vectors of Ape Plasmodium Parasites, Including Plasmodium
praefalciparum in Gabon. *PLoS ONE* **8**, doi:e57294
10.1371/journal.pone.0057294 (2013).

60   Hughes, G. L., Rivero, A. & Rasgon, J. L. Wolbachia Can Enhance
Plasmodium Infection in Mosquitoes: Implications for Malaria Control?
*Plos Pathogens* **10**, doi:10.1371/journal.ppat.1004182 (2014).

61   Zele, F. *et al.* Wolbachia increases susceptibility to Plasmodium infection
in a natural system. *Proceedings of the Royal Society B-Biological Sciences*
**281**, doi:10.1098/rspb.2013.2837 (2014).

62   Blagrove, M. S. C., Arias-Goeta, C., Failloux, A. B. & Sinkins, S. P. Wolbachia
strain wMel induces cytoplasmic incompatibility and blocks dengue
transmission in Aedes albopictus. *Proceedings of the National Academy of*
*Sciences of the United States of America* **109**, 255-260,
doi:10.1073/pnas.1112021108 (2012).

63   Moreira, L. A. *et al.* A Wolbachia Symbiont in Aedes aegypti Limits
Infection with Dengue, Chikungunya, and Plasmodium. *Cell* **139**, 1268-
1278, doi:10.1016/j.cell.2009.11.042 (2009).

64   Bian, G., Xu, Y., Lu, P., Xie, Y. & Xi, Z. The endosymbiotic bacterium
Wolbachia induces resistance to dengue virus in Aedes aegypti. *PLoS*
*Pathog* **6**, e1000833, doi:10.1371/journal.ppat.1000833 (2010).

65   Walker, T. *et al.* The wMel Wolbachia strain blocks dengue and invades
caged Aedes aegypti populations. *Nature* **476**, 450-453,
doi:10.1038/nature10355 (2011).

66   Joubert, D. A. *et al.* Establishment of a Wolbachia Superinfection in Aedes
aegypti Mosquitoes as a Potential Approach for Future Resistance
Management. *PLoS Pathog* **12**, e1005434,
doi:10.1371/journal.ppat.1005434 (2016).

67   Gillies, M. T. & de Meillon, B. *The anophelinae of Africa, south of the Sahara.*
Vol. 54 (The South African Institute for Medical Research, 1968).

68   Fanello, C., Santolamazza, F. & della Torre, A. Simultaneous identification
of species and molecular forms of the *Anopheles gambiae* complex by PCR-
RFLP. *Medical and Veterinary Entomology* **16**, 461-464 (2002).

69   Ndo, C. *et al.* Population genetic structure of the malaria vector *Anopheles*
*nili* in sub-Saharan Africa. *Malaria Journal* **9**, doi:161
10.1186/1475-2875-9-161 (2010).

70   Baldo, L. *et al.* Multilocus sequence typing system for the endosymbiont
Wolbachia pipientis. *Applied and Environmental Microbiology* **72**, 7098-
7110, doi:10.1128/aem.00731-06 (2006).

71   Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop
software platform for the organization and analysis of sequence data.
*Bioinformatics* **28**, 1647-1649, doi:10.1093/bioinformatics/bts199
(2012).

72    Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313,
doi:10.1093/bioinformatics/btu033 (2014).
73    Stamatakis, A. in *International Parallel and Distributed Processing*
*Symposium.* ( IEEE/ACM).
74    Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL): an online tool for
phylogenetic tree display and annotation. *Bioinformatics* **23**, 127-128,
doi:10.1093/bioinformatics/btl529 (2007).
75    Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer
Publishing Company, Incorporated, 2009).
76    Boundenga, L. *et al.* Haemosporidian Parasites of Antelopes and Other
Vertebrates from Gabon, Central Africa. *PLoS One* **11**, e0148958,
doi:10.1371/journal.pone.0148958 (2016).
77    Hamon, J. & Mouchet, J. Secondary vectors of human malaria in Africa.
*Medecine tropicale : revue du Corps de sante colonial* **21**, 643-660 (1961).
78    Olson, D. M. *et al.* Terrestrial ecoregions of the worlds: A new map of life
on Earth. *Bioscience* **51**, 933-938, doi:10.1641/0006-
3568(2001)051[0933:teotwa]2.0.co;2 (2001).
79    Neafsey, D. E. *et al.* Highly evolvable malaria vectors: The genomes of 16
Anopheles mosquitoes. *Science* **347**, 43-+, doi:10.1126/science.1258522
(2015).
649

651 **Figures and Tables**

653 **Table 1**. Summary of the *Anopheles* species screened in this study.

| Group/complex | Species | Malaria role | Infected | Tested | Infection (%) |
|---|---|---|---|---|---|
| gambiae | *An. gambiae* | H | 5 | 44 | 11 |
| | *An. coluzzii* | H | 2 | 58 | 3 |
| | *An. brunnipes* | | 0 | 1 | 0 |
| | *An. cinctus* | | 0 | 2 | 0 |
| moucheti | *An. moucheti* | H, P, A | 30 | 42 | 71 |
| | *An. nigeriensis* | h | 1 | 27 | 4 |
| | *An. "GAB-2"* | | 5 | 8 | 63 |
| | *An. "GAB-3"* | | 1 | 1 | 100 |
| | *An. gabonensis* | A | 0 | 29 | 0 |
| funestus | *An. funestus* | H | 2 | 37 | 5 |
| | *An. implexus* | | 1 | 26 | 4 |
| | *An. jebudensis* | | 1 | 2 | 50 |
| | *An. maculipalpis* | | 0 | 29 | 0 |
| nili | *An. nili* | H, A | 11 | 19 | 58 |
| | *An. carnevalei* | h, A | 2 | 29 | 7 |
| | *An. "GAB-1"* | | 0 | 19 | 0 |
| | *An. hancocki* | h | 1 | 41 | 2 |
| | *An. theileri* | h | 0 | 24 | 0 |
| | *An. rodhesiensis* | | 0 | 4 | 0 |
| coustani | *An. coustani* | h, A | 2 | 35 | 6 |
| | *An. paludis* | h, A | 1 | 16 | 6 |
| | *An. gr coustani* | h | 0 | 51 | 0 |
| | *An. squamosus* | | 0 | 32 | 0 |
| | *An. marshallii* | h, P, A | 2 | 42 | 5 |
| | *An. vinckei* | P, A | 3 | 30 | 10 |
| | | | **70** | **648** | |

Malaria role: known role for each species in malaria transmission [35,53,76,77] in humans

(H: major, h: secondary), primates (P), other animals (A), or unknown (blank).

**Figure 1**. Sampling sites and *Wolbachia* infection prevalence.

Map of Gabon showing the main African habitat types ([78], freely available at

http://maps.tnc.org/gis_data.html) and the villages were sampling took place (black

dots). The map was drawn using ArcGIS Basic v.10. The prevalence of *Wolbachia*

infection (number of infected *Anopheles* species and individuals) per site is presented

in bar charts. The pink colour indicates positive species/individuals and blue the total

number of species/individuals screened for *Wolbachia* infection at that site. CCB:

Cocobeach; LOP: Lopé; MKG: Mikongo; BTK: National Park of Plateaux Batékés;

FCV: Franceville; LBV: Libreville; MKB: National Park of Moukalaba-Doudou;

BKB: Bakoumba.

**Figure 2.** Circular phylograms of the *Wolbachia* strains isolated in the 16 *Anopheles*

species.

The phylogenetic trees were built with RAxML [72]. The names of the *Anopheles*

species from which the *Wolbachia*-specific sequences were isolated in this study are

shown in blue (positive for *Wolbachia* supergroup B), red (positive for supergroup A)

and brown (positive for supergroup C), while the names of mosquitoes species

(*Diptera*) from which the previously published *Wolbachia* sequences were isolated

are in green. Red dots show branches supporting a bootstrap >70% from 1000

replicates. (A) Circular phylogenetic tree using the *Wolbachia*-specific *16S* rRNA

fragment and *Anaplasma marginale* as outgroup. Different *Wolbachia* strains found in

the same *Anopheles* species are connected by pink lines. The pink bar charts indicate

the number of identical *Wolbachia* haplotypes found in each species. Scale bar

corresponds to nucleotide substitutions per site. (B) Circular phylogenetic trees based

on the *coxA, fbpA and ftsZ* fragment sequences using *Dirofilaria immitis* (supergroup

C) as outgroup. Specimens with a different supergroup assignation than *16S* are

marked with asterisks.

**Figure 3**. Maximum likelihood phylogeny of the 25 *Anopheles* species under study

and *Wolbachia* haplotypes.

The tree was inferred with RAxML [72] using the sequences of the *COII* fragment from

176 *Anopheles* specimens belonging to the 25 species under study and rooted with

*Anopheles darlingi* as outgroup (New World mosquito, diverged 100 Myr ago [79]). Red

dots in branches represent bootstrap values >70% from 1000 replicates. The shape of

each field column represents the *16S* (rectangle), *coxA* (rhombus), *fbpA* (triangle) and

*ftsZ* (hexagon) genes. The different *Wolbachia* gene haplotypes (i.e., unique allelic

profiles) are indicated with colour codes (all pink = the newly identified wANMO

strain). The bar chart size indicates the number of individuals of the same species with

the same haplotype, and the colour represents their infection status: grey, non-

infected; blue, infected by the *Wolbachia* supergroup B; red, infected by supergroup

A; brown, infected by supergroup C.

**Supplementary Materials**

**Text S1.** Mosquito taxonomic and molecular identification.

**Table S1.** Mosquitoes screened in this study and their accession numbers.

ID: Specimen identification name; Sites: Collection locations; Species: Morphological

and molecular identification; COII_hap: COII haplotype for each species; COII:

accession number for the cytochrome oxidase subunit II gene; 16S: accession number

for the 16S rRNA gene; ftsZ: accession number for the filamenting temperature-

sensitive mutant Z protein; fbpA: accession number the fructose-bisphosphate

putative aldolase protein; coxA: accession number for the cytochrome c oxidase

subunit I.

**Table S2.** Summary of the mosquitoes collected and screened in this study

**Table S3**. *Anopheles moucheti* F1 used to estimate vertical transmission.

**Figure S1**. Rooted maximum likelihood phylogeny of the filarial *Wolbachia* sequence

isolated from one *An. coustani* specimen.

The tree was inferred with RAxML [72] using the sequence of the filarial *COII* fragment

amplified from the *An. coustani* specimen BNG78 (in blue) and  public sequences

(NCBI) and rooted with *Brugia malayi* as outgroup. The black dot on the branch

indicates a bootstrap value >70% from 1000 replicates.

**Figure S2**. Scatterplot showing the genetic distances between *Wolbachia* strains (*16S*)

and infected *Anopheles* species *(COII).*

Genetic distances were estimated as the number of different bases between the

sequences of each pair of infected *Anopheles* specimens. The smoothed conditional

mean (blue line) and the 95% confidence intervals (grey area) were plotted using the

smoothing "gam" function of the ggplo2 library [75].

**Figure S3**. Probability of detecting *Wolbachia* infection.

The probability was estimated for each sample size and infection prevalence value.

The probability of correct estimation follows a black-blue gradient. The dashed line

indicates the 90% probability of infection detection.

# Figure 1

# Figure 2

# Figure 3

Tree scale: 0.01