Title:

Identification and characterization of conserved and divergent genes encoding the

nuclear envelope LINC complex in maize (*Zea mays* L.).

Running Title:

Maize LINC complex

Authors and Affiliations:

Gumber, Hardeep K.[1]; McKenna, Joseph F.[2]; Estrada, Amado L.[1]; Tolmie, Andrea F.[2]

Graumann, Katja[2]; Bass, Hank W.[1]

1. Department of Biological Science, Florida State University, Tallahassee, FL,
   USA, 32306-4295
2. Department of Biological and Medical Sciences, Faculty of Health and Life
   Sciences, Oxford Brookes University, Oxford, UK, OX30BP.

Corresponding Author:

bass@bio.fsu.edu

Key words:

LINC, SUN, KASH, nuclear envelope, maize

SUMMARY STATEMENT

Genes encoding maize candidates for the core LINC and associated complex proteins have been comprehensively identified with functional validation by one or more assays for several of the KASH genes.

ABSTRACT

The LINC (Linker of Nucleoskeleton to Cytoskeleton) complex serves as an essential multi-protein structure spanning the nuclear envelope. It connects the cytoplasm to the nucleoplasm and functions to maintain nuclear shape and architecture, as well as regulates chromosome dynamics during mitosis and meiosis. Knowledge of LINC complex composition and function in the plant kingdom is relatively limited, especially in the monocots which include the cereal grains and other grass species. We identified and classified 22 genes encoding candidate LINC complex and associated proteins in maize through bioinformatic and biochemical approaches. Representative KASH candidates were functionally validated in one or more assays including nuclear envelope localization, native or heterologous co-immunoprecipitation with antisera for ZmSUN2, and fluorescence recovery after photobleaching. These findings support a summary working model of the entire maize LINC and associated proteins complex with components found to be conserved across eukaryotes, unique to plants, or highly divergent and grass-specific. This model contributes a new experimental system for the cell biology of the nuclear envelope and new opportunities for future studies of the LINC complex in a model crop species.

List of Symbols and Abbreviations:

CDS   coding sequence

CRWN1   CROWDED NUCLEI 1

FRAP   Fluorescence Recovery After Photobleaching

INM   inner nuclear membrane

KASH   Klarsicht/ANC-1/Syne-1 homology

LINC   linker of nucleoskeleton and cytoskeleton

MLKP   Maize LINC KASH AtWIP-like

MLKG   Maize LINC KASH Grass-specific

MLKS   Maize LINC KASH AtSINE-like

MLKT   Maize LINC KASH AtWIT-like

MKAKU4   Maize KAKU4-like

NE   nuclear envelope

NET   nuclear envelope transmembrane protein

NCH   NMCP/CRWN-Homologous

NEAP   nuclear envelope-associated protein

ONM   outer nuclear membrane

SUN   Sad1/UNC-84

WIP   WPP domain-interacting protein

WIT   WPP domain-interacting tail-anchored protein

INTRODUCTION

In eukaryotic cells, the nuclear envelope (NE) is a structural hallmark that encapsulates the biparentally-inherited genetic material, the chromosomes. The NE is comprised of inner and outer nuclear membranes, with embedded nuclear pore complexes (NPC) and a variety of nuclear envelope transmembrane proteins (NETs). Two NETs, Sad1/UNC-84 (SUN), and Klarsicht/ANC-1/Syne-1 homology (KASH) domain proteins, interact in the perinuclear space to form the core of an evolutionarily conserved Linker of Nucleoskeleton and Cytoskeleton (LINC) complex which spans the NE (Crisp et al., 2006). LINC complexes carry out many functions ranging from organizing the shape and position of the nucleus as a mobile and pliable organelle to direct mechanotransduction to effects on chromatin structure and dynamics in somatic and meiotic cells (Kim et al., 2015; Kracklauer et al., 2013; Meier, 2016; Meier et al., 2017; Razafsky and Hodzic, 2015; Starr and Fridolfsson, 2010; Tamura et al., 2015; Tapley and Starr, 2013). The biological importance of NE functions is reflected in part through the wide range of NE-defective developmental disorders such as laminopathies or envelopathies (Burke and Stewart, 2014; Fridkin et al., 2009; Janin et al., 2017; Razafsky and Hodzic, 2015).

The core components of the LINC complex are the inner nuclear membrane (INM) SUN-domain proteins (Hagan and Yanagida, 1995; Malone et al., 1999) and the outer nuclear membrane (ONM) KASH-domain proteins (Starr and Han, 2002). The SUN-domain proteins are known to interact with chromatin, nuclear lamins, or lamin-like proteins (Haque et al., 2006; Hodzic et al., 2004; Zhou et al., 2015b). The KASH-domain proteins are known to interact directly or indirectly with cytoskeletal structures

including microfilaments, microtubules, intermediate filaments, microtubule organizing

centers, and organelles (Luxton and Starr, 2014; Starr and Fridolfsson, 2010). LINC

complexes are highly conserved in all eukaryotes yet many of the components have

evolved rapidly or recently, limiting the power of sequence conservation to identify

functional homologs.

The body of knowledge of the LINC complex comes primarily from studies in

opisthokonts, especially metazoans and yeast, with some but limited information for

other eukaryotes, including plants. Knowledge of plant LINC complex components,

especially in Arabidopsis, has recently began to expand (Meier, 2016; Meier et al.,

2017; Poulet et al., 2017a; Tiang et al., 2012; Zhou et al., 2015a). The first SUN

domain protein recognized in plants, OsSad1, was found as part of nuclear proteomic

study in rice (Moriguchi et al., 2005). Its relationship to the *Saccharomyces pombe* Sad1

was confirmed by localization of OsSad1-GFP fusion protein to the nuclear periphery in

onion epidermal cells (Moriguchi et al., 2005). Subsequently, two classes of SUN

domain proteins were found to be present in plants; first, the CCSD/C-terminal SUN

group, in which the SUN domain is near the C-terminus and second, the PM3/Mid-SUN

group, in which the SUN domain is centrally located (Graumann and Evans, 2010;

Graumann et al., 2014; Murphy et al., 2010; Oda and Fukuda, 2011). More recently, a

large number of SUN domain proteins have been identified in many plant species by

homology searches (reviewed by Meier, 2016; Meier et al., 2017; Poulet et al., 2017a).

The C-terminal SUN proteins are similar to those in animals, but the mid-SUN

proteins are less well studied as a group despite their known occurrence in plants

(Poulet et al., 2017a), yeast (Friederichs et al., 2011), Dictyostelium (Shimada et al.,

2010) and mice (Sohaskey et al., 2010). Among the conserved roles that SUN proteins

play in plants are those involving the maintenance of the shape and size of the nucleus

(Graumann et al., 2010; Graumann et al., 2014), plant growth (Graumann et al., 2014),

and meiotic chromosome behavior (Murphy et al., 2014; Varas et al., 2015).

The first plant KASH proteins to be described were the WPP-domain-interacting

proteins (WIPs). The WIPs are plant-specific, outer nuclear envelope associated

proteins originally discovered as RanGAP-NE anchoring proteins (Xu et al., 2007) and

later shown to bind to SUN and possess KASH-like features (Meier et al., 2010; Zhou et

al., 2012). A related group of ONM LINC-associated plant proteins are the WPP-

interacting tail-anchored (WIT) proteins (Zhao et al,. 2008), which bind directly to WIPs,

indirectly to SUN, and also have WPP-interacting domains. WITs possess the LINC-like

activity of connecting the NE to the actin cytoskeleton through their association with

plant specific Myosin XI-i (Tamura et al., 2013).   In addition to the WIPs and WITs, two

other categories of ONM KASH proteins have been described in plants, the SUN-

Interacting Nuclear Envelope proteins (SINEs) and the TIR-KASH (TIKs), both of which

are NETs (Graumann et al., 2014; Zhou et al., 2014).

There are currently recognized three plant-specific categories of genes encoding

proteins in the INM or at the intranuclear periphery that interact with SUN or each other

to associate with the LINC complex. These encode the INM Nuclear Envelope

Associated Proteins (NEAPs) and nucleoplasmic proteins, including Nuclear Matrix

Constituent Proteins (NMCPs, also called CRWNs) and the AtCRWN-binding

nucleoplasmic protein AtKAKU4s (Ciska et al., 2013; Dittmer and Richards, 2008; Goto

et al., 2014; Masuda et al., 1997; Pawar et al., 2016; Wang et al., 2013). While NEAPs

and CRWNs can be directly associated with LINC complexes by binding to C-terminal

SUN proteins, KAKU4 is indirectly associated with LINC complexes via interaction with

CRWNs (Goto et al., 2014). These interconnected proteins are thought have roles in

chromatin architecture and nuclear morphology, placing the LINC complex squarely in

genetic and genomic response pathways in plant cells (Poulet et al., 2017b).

In this study, we have identified, named and tested some of the core LINC

complex components in maize, an important agricultural and economic species with

historic contributions to genetics, cytogenetics, epigenetics, and meiosis (reviewed by

Nannas and Dawe, 2015). We include comparisons across phyla ranging from single

cellular chlorophyte to advanced land plants to highlight both conserved and divergent

maize LINC complex components.

RESULTS

Identification of maize genes encoding LINC or LINC-associated components

We developed a working model, shown in Figure 1, for the maize LINC complex

and associated proteins as identified through bioinformatic or experimental evidence.

The proteins are depicted as residing in one of the four cellular locations, the cytoplasm,

the ONM of the NE, the INM of the NE, or the nucleoplasm. Despite the considerable

difficulty in assigning certain orthology for all of the maize LINC components, we have

established the following nomenclature for these candidate maize LINC genes. The

genes encoding KASH-like and associated ONM proteins are named with the common

prefix of *MLK* for *Maize LINC KASH* followed by a one letter designation to delineate the

four KASH-related subgroups in maize: *G* for Grass-specific, *S* for *AtSINE*-like, *P* for

AtWIP-like, or *T* for *AtWIT*-like. The genes encoding the INM LINC proteins fall into two

groups, the previously published core LINC SUN domain proteins (Murphy et al., 2010),

and the genes designated *MNEAP* for Maize *AtNEAP*-like. Finally, the genes encoding

the nucleoplasmic LINC-associated proteins are designated as *NCH* for NMCP/CRWN

homologs or *MKAKU4* for *Maize KAKU4-like*. In summary the 22 known or candidate

core LINC and associated proteins in maize are comprised of 10 MLKs, 5 SUNs, 3

NEAPs, 2 NCHs, and 2 MKAKU4s. For all of these, the gene IDs in maizeGDB, PLAZA

gene, and UniPROT are summarized in Table 1 and diagrammed with their protein

domain structures in Figure 2. Of these, the KASH and SUN proteins are the most

similar to their animal counterparts, whereas the MLKGs, MLKTs, MNEAPs, MKAKU4s

appear to be more characteristic of plants.

The MLK genes: ten candidates for maize ONM core LINC genes

A hallmark of the KASH domain proteins is that they have limited sequence

conservation and are not reliably recognized by sequence similarity searches. Instead,

they contain a conserved structure consisting of a C-terminal transmembrane domain,

and a short periplasmic tail with characteristic terminal residues including a penultimate

proline followed by a threonine or serine. Only a few plant genes with similarity to

animal KASH genes have been identified so far (Zhou et al., 2012; Zhou et al., 2014).

To find all of the potential maize LINC ONM components, we first searched the maize

proteome (B73 AGPv4), from ensembl.gramene.org (Jiao et al., 2017) for KASH

proteins with the following characteristic features; (1) a TM domain at the C-terminus

followed by 9-40 amino acids that could extend into the perinuclear space, and (2)

terminal four residues of [DTVAMPLIFY], [VAPIL], P, and T (Poulet et al., 2017a; Zhou

et al., 2014). This search led to the identification of a total of eight genes encoding

maize KASH-like candidates, four with homology to AtWIPs (*MLKP1 – MLKP4*), two

with homology to AtSINEs (*MLKS1* and *MLKS2*) and two found only in members of the

grass family, Poaceae (*MLKG1* and *MLKG2*). We next searched the maize proteome

for WIT homologs using AtWIT1 and AtWIT2 protein sequences as queries (Zhao et al.,

2008) and found two maize genes, *MLKT1* and *MLKT2*, both encoding proteins with the

AtWIT-like TM domain at the C-terminus and multiple coiled-coil (CC) domains. Given

the rapid divergence of LINC gene sequences and the lack of distinct and diagnostic

domains, and gene content differences between maize cultivars (Hirsch et al., 2016), it

is possible that maize has even more MLKT genes not readily detected using these

criteria.

The four MLKP (AtWIP-like) genes belong to the PLAZA gene family designated

HOMO4M002181, and encode proteins with multiple coiled-coil domains (MLKP1-

MLKP4, Fig 2). The previously identified AtWIP proteins interact with AtWITs and

RanGAP1 and exhibit coiled-coil domain-dependent homo- and hetero-multimerization

(Zhao et al., 2008; Zhou et al., 2012). The MLKP proteins are therefore predicted to

interact with themselves, each other, ZmSUN proteins, and RanGAP proteins with WPP

domains. The two MLKS (AtSINE-like) genes belong to PLAZA gene family designated

HOMO4M002474, and encode proteins with armadillo (ARM) repeats (MLKS1- MLKS2,

Fig. 2). ARM domains have been reported in a variety of proteins across plant and

animal kingdoms as actin-binding and protein-protein interaction domain (Coates,

2003). The MLKS proteins are predicted, therefore, to interact and colocalize with actin,

as has been demonstrated for AtSINE1 (Zhou et al., 2014). The two MLKG (grass-

specific) genes lack both CC and armadillo domains and have been assigned to

different PLAZA gene families - HOMO4M009187 for *MLKG1* and HOMO4M008346 for

MLKG2.

Beyond the eight candidate MLK genes that we predict to encode maize ONM

core LINC complex proteins, we also found four other maize genes that met some of

our criteria. For instance, some encoded proteins with the terminal PT residues, but did

not satisfy the single C-terminal TM criterion. It remains unknown whether these genes

(B73v4 gene IDs Zm00001d024832, Zm00001d036395, Zm00001d005590 and

Zm00001d017144) encode actual variant LINC proteins or simply resemble LINC

proteins.

The eight maize INM LINC genes; five SUN and three NEAP genes

The five genes encoding maize SUN-domain proteins belong to two subfamilies

distinguished by the location of the SUN domain (Murphy et al., 2010). They are not

newly identified in this study, but are included in Table 1 for the sake of completion,

along with their most recent B73v4 gene IDs. ZmSUN1 and ZmSUN2 proteins have the

canonical C-terminal SUN domain, whereas the ZmSUN3, ZmSUN4,and ZmSUN5 have

a central SUN domain that is characteristic of, but not limited to plants (Graumann et al.,

2014). Despite the conservation of SUN-domain proteins as a group, and their

assignment to one of two subfamilies, direct assignment of orthology for individual SUN

genes across taxonomic groups such as eudicots (Arabidopsis) versus monocots

(maize) has yet to be established.

Another major family of INM LINC proteins are the higher plant-specific proteins

called Nuclear Envelope-Associated Proteins, the NEAPs (reviewed by Groves et al.,

2018; Meier et al., 2017; Poulet et al., 2017a). NEAPs exist as a multigene family, first

discovered in Arabidopsis and shown to interact with SUN-domain proteins (Pawar et

al., 2016). Genetic analysis of AtNEAPs showed that they affect several phenotypes,

including primary root growth, nuclear morphology, and chromatin organization (Pawar

et al., 2016) . NEAPs have three protein structural features, the presence of extensive

coiled coil domains, nuclear localization sequences, and C-terminal TM domains. We

used these criteria to identify three maize NEAP genes (Table 1, Fig. 2), *MNEAP1*,

MNEAP2, and *MNEAP3*. Interestingly, we observed partial homology to N-terminus of

structural maintenance of chromosomes domain (SMC_N, PFAMID PF02463) proteins

in the MNEAPs, further implicating them in DNA and chromosomal functions.

The MNEAP1 and MNEAP2 proteins both end with a characteristic C-terminal

hydrophobic patch and two basic terminal residues KR (Pawar et al., 2016). These two

proteins share 92% sequence identity and are very similar to the protein encoded by

one of the two sorghum NEAPs genes (sobic.009G033800). The *MNEAP1* and

MNEAP2 most likely arose from the genome duplication event that occurred following

the divergence of maize from sorghum, ~ 11 MYA (Gaut et al., 2000). *MNEAP3* fits our

initial criteria but is otherwise quite different in many respects including a unusually

large number of splice variant transcript models and the lack of KR terminal residues.

Consequently, *MNEAP3* may be a complex locus or alternatively, pseudogene no

longer under evolutionary pressure to maintain its structure.

The four maize nucleoplasmic LINC-associated genes; two *NCH* and two *KAKU4*

genes

In mammals, the SUN-domain proteins of the LINC complex interact with the

nuclear lamina which is located at the periphery of the nucleoplasm and made up of

lamin proteins. Plant nuclei lack laminA/B orthologs, but they do have a lamin-like

meshwork that is thought to consist of lamin-like proteins known as NMCP/CRWN as

well as CRWN-binding nuclear proteins, KAKU4, (Ciska et al., 2013; Goto et al., 2014).

These proteins are plant-specific, collectively thought to comprise plant nuclear

architecture proteins, and are associated with the LINC complex through CRWN-SUN

interactions (Goto et al., 2014).

Using AtCRWN and DcNMCP protein sequences as queries, we identified two

maize genes, *NCH1* and *NCH2*, representing the two major clades of NMCP/CRWN

genes (Ciska et al., 2013; Wang et al., 2013). The maize *NHC1* is most similar to

DcNMCP1, *DcNMCP3*, and *AtCRWN1*. The genes in this group are members of

PLAZA gene family HOM04M004640 (Table 1). The maize *NCH2* gene is most similar

to *DcNMCP2*, *AtCRWN2*, *AtCRWN3*, and *AtCRWN4*. The genes in this group are

members of the PLAZA gene family HOM04M003564. Both NCH1 and NCH2 proteins

share several features with their plant homologs. These features include large proteins

(~100 kDa or greater), multiple coiled-coil domains over the majority of the protein,

multiple CDK1 phosphorylation sites, and a nuclear localization sequence near the C-

terminus (Fig. 2).

Using AtKAKU4 protein sequence similarity searches, we identified two highly

similar maize KAKU4-like genes, *MKAKU41* and *MKAKU42*. These two genes and

AtKAKU4 are all members of the PLAZA gene family HOM04M004991. KAKU4 and

KAKU4-like genes appear to be limited to flowering plants and encode proteins that

have been described as candidate lamina components (Goto et al., 2014). As a group,

they may be remotely associated with the LINC complexes but are included here

because of the known interaction of AtKAKU4 with AtCRWN proteins as well as the

genetic evidence that implicates them in LINC functions and nuclear architecture (Goto

et al., 2014). Aside from a NLS near the N-terminus, the MKAKU4 proteins do not

contain any notable structural protein domains or motifs (Fig. 2).

A mix of ancient and derived genes encode components of the maize LINC

complex

Given that the time of appearance and the evolutionary conservation of genes

often reflects functional properties, we examined the maize LINC candidate genes and

their counterparts in a variety of plant species as summarized in Figure 3. Recent

phylogenetic analyses of plant LINC genes or sub-families therein have been published

(Ciska et al., 2013; Poulet et al., 2017a; Zhou et al., 2014), with emphasis on

Arabidopsis and other eudicot species, but did not include the latest PLAZA gene

families or maize B73v4 annotations (Jiao et al., 2017; Van Bel et al., 2018). The

species listed (Fig. 3) are arranged in increasing evolutionary distance from maize

starting with sorghum and extending to the unicellular green alga, *Chlamydomonas*

reinhardtii.

In plants, the most highly conserved members of the core LINC and associated

complex belong to gene families that encode the C-terminal and mid-SUNs, NCH2 and

the AtSINE-like KASH proteins MLKS1 and MLKS2). These gene families are present

in all the plant species listed (shaded blue or green, Fig. 3). Curiously, of all the genes

listed, only the mid-SUN genes are listed as present in Chlamydomonas PLAZA gene

families. The next most highly conserved plant LINC gene families encode the MNEAP

and NCH1 proteins. They are present in seed-bearing plants (shaded tan, Fig. 3),

which includes pine, a conifer species. The third most highly conserved plant LINC

gene families encode MLKP, MLKT and MKAKU4 proteins, all of which are found in

flowering plants, the angiosperms (shaded pink, Fig. 3). In contrast, the most recent to

appear in evolutionary time are the gene families that encode the MLKG proteins,

present in the grass family within the monocots (shaded yellow, Fig. 3). The biological

significance of the timing of evolutionary appearance remains to be determined, but we

generally expect that the more widely distributed gene families have more ancient or

basal functions, whereas the more recent gene families could have resulted from

duplication followed by either regulatory divergence, such as tissue-specific expression,

or gain of new biological functions.

Maize LINC genes exhibit tissue-specific and developmental co-expression

patterns

We analyzed the expression patterns of the maize LINC candidate genes as

shown in Figure 4 in order to examine them for evidence of expression and potential

functional groupings that should be reflected in their differential or co-expression

patterns. For this, we use the expanded maize gene expression atlas which contains

transcriptome data from numerous tissues broadly sampled across most of the

developmental stages of the maize life cycle (Stelpflug et al., 2016). For comparison,

we included three housekeeping genes (HKG, Fig. 4A), *CDK1*, *RPN*, and *SGT1* (Lin et

al., 2014).   We excluded *MNEAP3*, which lacked a gene model when the atlas was

produced, and *MLKG1*, which had zero reads in all fifteen tissues selected for this

study. Despite their exclusion, these two are presumed to be real genes and not

pseudogenes because of the expression of their corresponding transcripts detected in

one or more tissues (Chettoor et al., 2014; Walley et al., 2016; Wang et al., 2016; Xu et

al., 2012)*.*

Most of the candidate maize core LINC and LINC-associated genes showed a

pattern of broad expression across multiple tissues, with co-expression patterns

(dendrogram, Fig. 4A) revealing several distinct groupings.   The "root" co-expression

group (orange box, Fig. 4A) includes three root samples and 12 DAP developing seed.

The "young inflorescence" co-expression group (blue box, Fig. 4A) includes male

(immature tassel) and female (immature cob) tissues. The "meristem" co-expression

group (green box, Fig. 4A) includes the developing embryo and growing shoot tip

tissues. These last two groups together form a larger co-expression cluster from

tissues that share the property of having actively dividing cell populations. Beyond

these three groups, several tissues show distinctive patterns of LINC gene expression,

not clustering with other tissues, including anthers (male pollen-bearing organs),

topmost seedling leaf, meiotic tassel, and silks (female, pollen-receiving organs).

To characterize the general expression levels of the candidate maize LINC

genes, we plotted the median FPKM values across different tissues (Fig. 4B). Maize

LINC genes generally exhibit low to moderate expression levels, similar to that of the

house-keeping *CDK1* gene (Fig. 4B). The more highly expressed LINC-related genes

had transcript levels that ranged from 80-140 median FPKM and included the

MKAKU4s, the C-terminal SUNs, the NCHs, and some of the KASH genes, *MLKP3* and

MLKP4. The lower expressed genes had transcript levels that ranged from 3-50

median FPKM and included the mid-SUNs, the MNEAPs, and most of the other KASH

genes. As a group, the maize AtWIP-like genes (*MLKP1-4*) have higher expression

levels than the AtSINE-like genes (*MLKS1*, *MLKS2*), with *MLKP3* and *MLKP4* showing

the most uniform and broad expression. In contrast, the mid-SUN *SUN5* gene exhibited

a most striking pattern of expression in only pollen. The *Poaceae*-specific gene,

MLKG2, appears to be expressed in tissues characterized by meristem and active cell

division, but absent from others such as endosperm, root, leaf, and silks.

Among the highly similar gene pairs that are likely to have arisen as part of a

recent genome duplication (Gaut et al., 2000), some of them (*MLKP1 and MKLP2;*

MLKP3 and MLKP4; and MKAKU41 and MKAKU42) show similar expression levels

whereas other gene pairs, such as *MNEAP1* and *MNEAP2* exhibit differential

expression levels. Overall, this expression analysis provides evidence for LINC and

associated proteins genes being transcribed and regulated in a tissue-specific manner.

The co-expression groupings predict possible but untested genetic or physical

interactions that may reflect shared functions.

Identification of LINC and associated proteins from native co-IP using anti-

ZmSUN2 antibodies.

In order to find native cellular interactors of SUN proteins, we carried out co-

immunoprecipitation (co-IP) optimized for maize NE proteins from purified earshoot

nuclei as summarized Table 2. Maize earshoots of ~3-5 cm in length yield high quality

cellular extracts from fast-growing somatic tissues several days prior to silking and

fertilization. The antibody used for native co-IP was the affinity-purified sera designated

anti-SUN2-NPAP as previously characterized (Murphy et al., 2014). Control co-IP

experiments were conducted using either no antibodies or irrelevant but similarly

affinity-purified rabbit sera, anti-ZmNDPK1 (Kopylov et al., 2015).   A key technical

advance was the use of the short-chain crosslinker dithiobis(succinimidyl propionate),

DSP. We adapted the membrane protein cross-link immunoprecipitation (MCLIP)

method for use with maize because it was shown to effectively recover SUN-interacting

proteins from U2OS cells (Jafferali et al., 2014). We found that the MCLIP method

using DSP together with light formaldehyde fixation to preserve the nuclei resulted in

substantially increased final recovery and detection of known or predicted maize

homologs of plant LINC, NE, or SUN-interacting proteins (Tamura et al., 2013; Wang et

al., 2013; Zhou et al., 2012).

The proteins recovered from the co-IP assays were subjected to tryptic digestion

and identified by liquid chromatography tandem mass spectrometry (LC-MS/MS).   Over

140 proteins were identified (Supplemental Table 2) using software thresholds of 95%

protein and 95% peptide. As expected, SUN2 was the most abundant protein, but we

also repeatedly detected SUN1, which could result from crossreactivity of ZmSUN1 with

our anti-SUN2 sera, or from actual co-IP of ZmSUN1 bound to SUN2, as expected from

documented hetero multimerization of SUN proteins in plants and animals (Sosa et al.,

2012).

The co-precipitated proteins identified were considered high confidence if they

were detected in multiple replicates but not in the negative controls. The specificity of

the modified MCLIP method is evidenced by recovery of core LINC components and

predicted cytoplasmic and nucleoplasmic proteins. Among the predicted core LINC

components repeatedly found in the top co-IP hits were three ONM proteins (MLKG1,

MLKS2, and MLKT2) two INM proteins (ZmSUN1 and ZmSUN2), two NMCP/CRWN

homologs (NCH1 and NCH2), two RAN family proteins (RAN GTPase and RAN-GAP2)

and five nuclear pore proteins (NUP133, NUP96, NUP88, GP210, and NUA).

We also found proteins that were co-precipitated but likely via indirect interaction

with SUN. These include the cytoskeletal proteins TUB6, TUB4 and Actin-1/ARP7. The

Actin-1/ARP7 protein could be cytoplasmic or nucleoplasmic or both because its

annotation includes the designation of nuclear actin, an intriguing possibility in light of

recent findings showing cross-species binding of Arabidopsis AtARP7 to the carrot

nuclear matrix protein DcNMCP1 (Mochizuki et al., 2017). We recovered several

nuclear proteins with DNA-binding activities including transcription factors bZIP, ARF10,

and the linker histone H1a.   Overall, the co-IP resulted in confirmatory identification of

native SUN2-interacting proteins with known or suspected LINC proteins, plus several

additional proteins that represent new targets for future investigation. To further verify

some of the LINC candidates from either the Co-IP or bioinformatic analyses, we

selected three genes for further localization and interaction assays *in vivo* using

transient heterologous expression assays.

In planta evidence of NE targeting and ZmSUN2 interaction for three candidate

maize LINC proteins

In order to investigate the subcellular localization of the maize KASH proteins, we

transiently expressed N-terminal GFP fusions of MLKG1, MLKP1 or MLKP2 in _Nicotiana_

benthamiana leaf tissue as summarized in Figure 5. Expression of a positive control,

mCherry-ZmSUN2, shows that maize NE proteins can be properly expressed from the

plasmids and localized to the NE in this leaf cell heterologous expression system. All

three of the previously untested ONM maize LINC candidate proteins, MLKG1, MLKP1

and MLKP2, also localized to the nuclear periphery (Fig. 5A). These localization

patterns are comparable to those previously reported for AtKASH proteins in similarly

designed expression assays (Zhou et al., 2012; Zhou et al., 2014).   For GFP-MLKG1, a

network-like pattern at the cell periphery was also seen (Fig. 5B). In order to further

examine this pattern, we co-expressed GFP-MLKG1 with an ER marker and an actin

marker and found that GFP-MLKG1 colocalizes with both the ER and the actin network

(Fig. 5B). Similar colocalization patterns have previously been demonstrated for

AtSINE1 (Zhou et al., 2014) but, unlike AtSINE1, MLKG1 does not possess an ARM

domain or other known actin binding motifs.

The C-termini of the KASH proteins are known to be required for NE localization

which is mediated through interaction with SUN proteins (Sosa et al., 2012; Zhou et al.,

2012; Zhou et al., 2014). To test whether this is also true of our maize KASH

candidates, we created and tested three classes of C-terminal deletions or replacement

constructs. ΔKASH represents constructs in which KASH domain, that is, all the amino

acid residues following the TMD are deleted (Fig. S1A). Δ4r, ΔKASHr and ΔTMr

represent constructs in which terminal four residues, KASH domain, or everything

beginning with TMD have been replaced with a non-genic 38 aa terminus from the

vector backbone, respectively (Fig. 5C, S1A).

We observed that each of these C-terminal deletion and replacement constructs

led to the loss of NE localization (Fig. 5C; Sup 1B). The modified proteins showed

localization patterns that resembled soluble proteins, visible throughout the cytoplasm

and nucleoplasm (Fig. 5C, supplemental figure 1). For comparison, we expressed GFP-

only as a control (Supplementary Figure 1B), which illustrates the distribution of a

soluble protein without a TM domain in the cytoplasm and nucleoplasm. Interestingly, in

every case, the Δ4 and Δ4r modifications were sufficient to abolish NE localization

despite the fact that the fusion protein still has a TMD. These results are consistent with

the idea that the terminal residues of these proteins may confer several properties

including stable post-translational membrane insertion and targeting to the ONM. In

order to investigate whether the NE localization seen for these maize KASH proteins

involves binding to maize SUN proteins, *in vivo*, we co-expressed three different maize

GFP-KASH with or without ZmSUN2 or deletion derivatives of ZmSUN2 (Figure 6A). If

they do interact, we would expect to see a reduction in the mobility of GFP-KASH as

measured by fluorescence recovery after photobleaching and indicative of interacting

partners, as previously described (Zhou et al., 2014). For two of the three, MLKG1 and

MLKP2, we did indeed observe reduced mobility when co-expressed with ZmSUN2 as

summarized in Fig. 6B. When GFP-MLKG1 is transiently expressed by itself, 63±8% of

the protein population is mobile while 36% remain immobilized. However, upon co-

expression with mCherry-ZmSUN2, only 49±12% of the protein population was mobile

indicating that the co-expressed ZmSUN2 substantially increased the immobile fraction

of GFP-MLKG1 from 36% to 51% (Fig. 6 A,B). We interpret this shift as indirect

evidence for ZmSUN2 interaction with GFP-MLKG1. To test this idea, GFP-MLKG1

was co-expressed with a domain deletion mutant of ZmSUN2 lacking the SUN domain

(mCherry-ZmSUN2ΔSUN) and found that the mobility of MLKG1 was similar to that of

the MLKG1 alone (68±7%; Fig. 6 B,C). This finding indicates that the SUN domain is

required for mediating the change in GFP-MLKG1 mobility, providing strong indirect

evidence that ZmSUN2 interacts with MLKG1. To further test if the coiled-coil domain of

the SUN protein is required for reduced mobility, we created a mCherry-ZmSUN2

lacking the coiled-coil domain, mCherry-ZmSUN2ΔCC.   Co-expression of this mCherry-

ZmSUN2ΔCC with GFP-MLKG1 had a similar effect on KASH mobility to that of the full

length SUN2 (Fig. 6 B, C). This result establishes that the ZmSUN2-co-expression-

dependent increase in the immobile fraction of MLKG1 occurs to a similar degree with

or without the CC domain.

Similar changes in protein mobility were observed for GFP-MLKP2 (Fig. 6 F,G)

when co-expressed with full-length ZmSUN2, ZmSUN2ΔSUN or ZmSUN2ΔCC. Co-

expression with ZmSUN2 or ZmSUNΔCC resulted in significant increases in the

immobile fraction of GFP-KASH from 51% to 69% or 71%, respectively. to GFP-

MLKG1, co-expression of GFP-MLKP2 with ZmSUN2ΔSUN did not result in a

significant change in the mobile fraction of the maize KASH. In contrast, we did not

observe statistically significant changes in GFP-MLKP1 mobility when co-expressed

with full-length or deletion constructs of ZmSUN2 (Fig. 6 D, E), even though the

changes observed were all in the same directions as those for GFP-MLKG1 and GFP-

MLKP2. Representative FRAP images, individual repeat curve data and halftime

recovery values are shown in Supplementary Figures 2, 3, and 4.

Of the 3 candidates tested, MLKG1 was already found by native co-IP to interact

with SUN2, but the other two, MLKP1 and MLKP2 were not, presumably because of

their low expression levels in earshoot (Fig. 4). In order to check for their ability to

interact with SUN2 *in planta*, we assayed them by co-IP in tobacco tissues where they

were co-expressed as heterologous fluorescent fusion proteins (Supp. Fig 5A). Using

anti-GFP antisera, we observed co-precipitation of mCherry-SUN2, but not in control

extracts from non-infiltrated or mCherry-SUN2 only samples. These additional tests

confirmed that all three of the proteins tested, MLKG1, MLKP1, and MLKP2, showed

SUN2 interaction in one or more direct or indirect interaction assays (summary table in

Supp. 5B).

In summary, the fusion protein transient expression showed that all three

candidates tested showed localization to the NE, and deletion analysis established that

the last four amino acid residues were required for that localization. Furthermore,

these showed evidence of ZmSUN2 interaction *in planta* from one or more experiments,

inferred from co-expression FRAP or detected in native or heterologous co-IP

experiments. These findings provide the first experimental evidence for the

identification of multiple LINC complex components in monocot plant species,

evolutionarily distant from Arabidopsis and relevant to establishing the conservation of

the LINC complex across all eukaryotes.

DISCUSSION

Here we set out to systematically identify all of the maize LINC core and associated components as a way to advance nuclear envelope research. Despite the wide variety of functionalities attributed to LINC complex in plants, animals, and yeast, there remains a considerable number of unknowns regarding the mechanisms and functions of individual components. Our findings represent a valuable model grass crop species for future studies of the plant nuclear envelope and important biological processes such as meiotic chromosome behavior, development, signaling, and pathology (Kim et al., 2015; Kracklauer et al., 2013; Meier et al., 2017; Razafsky and Hodzic, 2015; Tapley and Starr, 2013). Maize provides a valuable plant complement to Arabidopsis, where SUN1 and SUN2 double knockouts result in disrupted meiotic progression, excessive chromosomal interlocks, and reduced fertility (Varas et al., 2015). However, Arabidopsis does not exhibit a canonical early prophase telomere bouquet cluster on the nuclear envelope, nor has it been shown to have a meiotic SUN belt (Murphy et al., 2014). These differences highlight one example of an important principle that not all findings from one plant species can be generalized across the kingdom. That, together with the difficulty in identifying orthologs illustrate the importance of defining the composition of the LINC complex in multiple species where genetic and cytological resources afford efficient and comparative analysis of the associated biological functions.

Searching for maize KASH and associated ONM proteins, we found homologs of other plant genes encoding AtWIP and AtSINE1/2-like proteins, but not those encoding AtSINE3/4, SINE5 or AtTIK (Graumann et al., 2014; Zhou et al., 2014). Genes not found

in maize show limited evolutionary distributions. *AtSINE3/4* are found only in

brassicaceae, *SINE5* only in medicago, and *AtTIK* only to Arabidopsis (Poulet et al.,

2017a). Similarly, *MLKG1* and *MLKG2* genes are found only in maize or other grasses

but not eudicot species such as Arabidopsis or Medicago. In contrast, the *MLKT* genes

of maize are clearly homologous *AtWITs*, a rare case in which sequence conservation is

sufficient for identification of an ONM LINC component. It is worth noting, however, that

other maize genes resembling ONM LINC components were found but fell short of the

criteria we applied in this study. It remains an intriguing but recurring observation that

ONM LINC proteins contribute to ancient structure but often exist as highly divergent in

extant species. In this regard, plant KASH genes are similar to those from opisthokonts,

with characteristic variation in protein sequence and structure (Meier et al., 2017).

536   Searching for additional maize genes encoding INM proteins, we found three

homologs of the plant-specific *NEAP gene*s. The *NEAP* genes are limited to seed

plants (Poulet et al., 2017a) and not found in any of the representatives of the more

ancient moss and algal species included in this study (Fig. 3). The *MNEAP1* and

MNEAP2 genes encode protein sequences that end in two basic residues, like their

eudicot homologs, but the functional significance of these terminal residues remains

unknown. Interestingly, the maize NEAPs have SMC_N domains which are currently

uncharacterized but may point to a chromosome tethering or nuclear structural role.

The maize NEAPs named in this study are three of the eleven maize genes assigned to

the PLAZA gene family HOM04M002962. Consequently, the actual number of the

maize NEAP genes may grow as revised bioinformatic and functional criteria for their

recognition emerges. In contrast, we did not find evidence for additional SUN genes of

maize beyond the five previously described (Murphy et al., 2010; Murphy et al., 2014).

Searching for nucleoplasmic proteins associated with plant LINC complexes, we

found two maize genes with homology to *NMCP/CRWN* genes, *NCH1* and *NCH2*, both

of which are deeply conserved in plants. Each of these are associated with multiple

transcript models (Wang et al., 2016), suggestive of splice variants, an intriguing

observation in light of the tendency for animal lamin genes to also have many splice

variants (Broers et al., 2006). Identification of plant nuclear matrix components that

interact with the nuclear envelope provides new opportunities to investigate the

mechanisms that underlie how signals are transduced via the LINC complex.

Cellular localization experiments confirmed the NE localization of fluorescent

fusion protein constructs for ZmSUN2 and all three of the maize KASH homologues

tested, MLKG1, MLKP1 and MLKP2 (Fig. 5). Interestingly, GFP-MLKG1 colocalizes

with the ER and the actin network, implicating it in cytoskeletal signal transduction to the

nucleus. Relatedly, Arabidopsis actin is tethered to the LINC complex through direct

interactions with AtSINE1 and through myosin XI-i - WIT-WIP interactions (Tamura et

al., 2013; Xu et al., 2007; Zhou et al., 2014). The presence of *SINE1* (*MLKS1* and

MLKS2), *WIP* (*MLKP1 - MLKP4*) and *WIT* (*MLKT1* and *MLKT2*) homologs in maize,

together with the poaceae-specific *MLKG1* suggests that plant genomes may encode a

diverse repertoire of of proteins to connect and integrate NE with cytoskeletal

processes.

To provide additional experimental evidence on the composition of the maize

LINC complex, we used a ZmSUN2 Co-IP method optimized for membrane proteins

together with isolated nuclei. We found a large number of different proteins that were

generally consistent with predictions from our working model (Fig. 1) and previous

reports (Dittmer and Richards, 2008; Goto et al., 2014; Pawar et al., 2016; Tamura et

al., 2010; Tamura et al., 2013; Wang et al., 2013; Zhou et al., 2012; Zhou et al., 2015b;

Zhou et al., 2015c). Among the expected and reported LINC and LINC-associated

proteins, our native Co-IP detected SUN, WIPs, WIT, NCH, and RAN cycle proteins.

In relation to the cytoplasmic and ONM side of the LINC complex, we detected

proteins predicted to bind directly to ZmSUN2 including MLKG1 and MLKS2. We also

co-precipitated proteins predicted to bind indirectly to ZmSUN2 including MLKT2, two

RAN cycle proteins, RAN GTPase, RAN-GAP2, Tub4, Tub6, and the nuclear pore

proteins NUP133, NUP96, NUP88, GP210, and NUA. Notably, we did not detect all of

the nuclear pore proteins, consistent with previous studies showing interactions of a

specific and limited subset of NPC proteins with SUN-domain proteins in association

with transcriptional activation and mRNA export processes (Gerber et al., 2004; Li and

Noegel, 2015). In relation to the nucleoplasmic and INM side of the LINC complex, we

detected two INM proteins, ZmSUN1 and ZmSUN2; and two NMCP/CRWN proteins,

NCH1 and NCH2.

The Co-IP experiments also yielded an interesting list of additional proteins

beyond the core LINC and associated proteins summarized in our working model.

Some of these proteins appear to belong to non-specific protein categories, but in fact

have been found in various NE studies including nuclear actin (ARP7/Actin-1),

transcription factors (bZIP5, ARF10, MYBR35, bHLH72/bHLH041), Histone H1 (H1a),

and ribosomal proteins (L19, L6, and L22-2) (Horigome et al., 2011; Hotta et al., 2007;

Mochizuki et al., 2017; Nakayama et al., 2008). Nuclear actin is increasingly recognized

as playing key roles in chromosome dynamics and gene expression (reviewed by

(Spichal and Fabre, 2017) whereas a plant linker H1 histone was recently shown to

have MTOC activity and reside at the nuclear periphery (Boruc et al., 2012; Nakayama

et al., 2008).

Given the increasing recognition that the NE and nucleoskeletal proteins are

implicated in chromatin remodeling and gene regulation, these additional, non-LINC

proteins discovered through native co-IP represent ideal targets and experimental

inroads for mechanistic studies of LINC-mediated dynamic processes coupled to

chromatin structure and nuclear architecture.   For instance, the NE cooperates with

nucleoskeletal proteins to modulate euchromatin and heterochromatin organization

(Poulet et al., 2017b; Wang et al., 2013). Emerging breakthroughs in HiC mapping of

nuclear architecture are likely to uncover additional principles of genome organization

that involve LINC-binding nuclear proteins (Grob et al., 2014). Given the importance of

lamins in human health, it is possible that plant NMCP/CRWN/NCH defects may even

model animal disease processes.

Both Co-IP and FRAP provide complementary assays for investigating *in planta*

binding of maize KASH candidates to ZmSUN2 (Fig. 6). For example, MLKG1 was co-

purified from maize extracts using SUN2 antisera (Table 2) while FRAP assays

demonstrated a ZmSUN2-dependent change in the mobility of GFP-MLKG1 and GFP-

MLKP2. The heterologous expression system described in this and previous studies

(Zhou et al., 2012; Zhou et al., 2014) demonstrates an efficient means to detect,

isolate, and dissect protein-protein interactions for distantly related grass species such

as maize, which is not easily transformed. They also provide an efficient means to carry

out *in vivo* tests of candidate interactors (e.g. Supp Fig 5) for which biochemical

amounts of native tissue may be limiting.

Overall, this study provides a comprehensive list of maize LINC candidates (Fig.

1) with multiple lines supporting experimental evidence. Through our efforts to identify

the the full ensembl of LINC components in a model eukaryote, we have contributed to

valuable new avenues of research to address unanswered questions regarding the role

of the nuclear envelope as an integrative structure coordinating nuclear and cellular

functions.

METHODS

Bioinformatic identification of MLKs

The maize B73 proteome v4

(ftp://ftp.gramene.org/pub/gramene/CURRENT_RELEASE/fasta/zea_mays/pep/) was

searched for sequences ending in motif [DTVAMPLIFY][VAPIL]PT (brackets indicate

alternative amino acid residues at the respective position). The selected proteins were

further searched for the presence of TMD using TMHMM (Krogh et al., 2001) and

Phobius softwares (Käll et al., 2004). Proteins lacking TMD or specific ending were

discarded. The selected maize MLKs were then searched on PLAZA monocot v4

(https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v4_monocots/) and those

belonging to same homolog family were grouped together.

Protein feature calculations and domain searches

Theoretical Molecular weight and isoelectric point (pI) of candidate proteins was

calculated using pI/Mw tool of ExPASy (https://web.expasy.org/compute_pi/). Coiled coil

domains were identified by Jpred (http://www.compbio.dundee.ac.uk/jpred/). For the

identification of conserved domain sequences, the candidate protein sequences were

searched on Uniprot (http://www.uniprot.org/) , NCBI Conserved Domain Database

(https://www.ncbi.nlm.nih.gov/cdd/), PFAM (https://pfam.xfam.org/), SMART

(http://smart.embl-heidelberg.de/). The domains identified were then plotted using IBS

1.0.3 (Illustrator for Biological sequences) software (Liu et al., 2015).

Identification of LINC orthologs in other organisms and phylogenetic

reconstruction

Sequences of orthologs of Maize LINC candidates belonging to matching HOMO gene

families were retrieved from PLAZA v4

(https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v4_monocots/). Sequences of

all species selected for this study are present in PLAZA except *Medicago truncatula*,

which were retrieved from Gramene. The protein sequences were further screened for

the presence of characteristic protein features. The MLKS/MLKG/MLKP orthologs were

searched for the presence of TMD towards C-terminus and for terminal four residues of

the proteins to be VIPT, VVPT, AVPT, PLPT, TVPT, LVPT or PPPS. For MLKT

orthologs, proteins having high sequence similarity and C-terminal TMD were retained.

For NCHs, KAKUs and NEAPs, sequences having high sequence similarity to

Arabidopsis CRWNS, KAKU4 and NEAPs were selected. For phylogenetic

reconstruction of each protein category, sequences were first aligned using MUSCLE

and Neighbour end joining trees were generated in MEGA 7.0.21 (Kumar et al., 2016)

using default parameters.

Transcriptome

The expanded maize gene expression atlas (Stelpflug et al., 2016) was

downloaded for analysis. Log_{10} Fragments Per Kilobase of exon per Million fragments

mapped (FPKM) values for 15 tissues were used to construct heatmap using Heatmap2

package of R. To group tissues by similar expression profiles, hierarchical clustering

was done using Hclust function of R.

Co-IP protocol

Maize (Zea mays L) B73 seeds were obtained from WF Thompson (NC State

University, Raleigh, NC, USA, accession "27 FARMS", bulked for NCSU, denoted

B73NC27). Seeds were grown at the Florida State University Mission Road Research

Facility (Tallahassee, FL). Young earshoots (3-5 cm long) were harvested from 7-8-

week-old plants, flash frozen in liquid nitrogen and stored at -80 °C until use. Ten

grams tissue, were ground under liquid nitrogen with a mortar and pestle and cross-

linked with 0.1% formaldehyde by stirring for 10 min in 100 mL ice-cold Buffer A (15

mM PIPES-NaOH, pH 6.8, 0.32 mM sorbitol, 80 mM KCl, 20 mM NaCl, 0.5 mM EGTA,

2 mM EDTA, 1 mM DTT, 0.15 mM spermine, and 0.5 mM spermidine; (Belmont et al.,

1987). The nuclei were isolated as described in (Vera et al., 2014) and resuspended in

5 ml PBS. The isolated nuclei were treated with 2 mM DSP (dithiobis(succinimidyl

propionate) (Cat. No. 22585, Thermo Scientific) crosslinker for 2 h at 4 °C. To lyse the

nuclei, 7 M urea and 1% Triton X-114 and Plant protease inhibitor cocktail (Cat. No.

P9599, Sigma) were added and incubated for 20 min on ice. The ruptured nuclei were

sonicated three times in Bioruptor (Diagenode) for 5 minutes each at medium setting.

Urea was diluted with 8 volumes of PBS. Homogenates were centrifuged at 10,000g

for 10 min at 0 °C to remove cellular debris. The supernatant was collected and pre-

cleared by adding 0.5 mg Protein A dynabeads (Cat No.10000D, Thermo Fisher) for 30

min with rotation at 4 °C. The beads were removed by placing tubes on magnetic rack

(Cat. No.12321D, Thermo Fisher). The cleared lysate was split into 3 equal aliquots. 3

µg α- SUN2 antibody was added to first aliquot, 3 µg NDPK control antibody to the

second tube and no antibody added to the third tube. The tubes were incubated at 4 °C

with rotation for overnight followed by the addition of 0.5 mg Protein A magnetic beads

to each tube for one-hour incubation at 4 °C. The antigen-antibody bound beads were

captured with the help of magnetic rack and washed thrice with PBS supplemented

with 0.02% Tween-20. The protein complexes bound to the beads were digested on

the using ProteoExtract All-in-One Trypsin Digestion Kit (Cat. No. 650212,

Calbiochem) according to manufacturer's instructions. Briefly, beads were

resuspended in Extraction buffer and digestion buffer was added, reduced using the

provided reducing agent for 10 minutes at 37°C. Samples were cooled to RT and then

blocked using blocking reagent for 10 minutes at room temperature. Trypsin at a final

concentration of 8 ng/µl was added and incubated for 2 hours at 37°C with shaking.

The reaction was stopped by acidification. Digests were vacuum dried and submitted

for LC-MS/MS to Translational Lab at Florida State University.

Mass spectrometry and data analysis

Digested peptides were resuspended in 0.1% Formic acid for loading onto LC-MS at

Translational Lab, Florida State University. An externally calibrated Thermo LTQ

Orbitrap Velos nLC-ESI-LTQ-Orbitrap (high-resolution electrospray tandem mass

spectrometer) was used with the following parameters. A 2 cm, 100 μm i.d. (internal

diameter) trap column (SC001 Easy Column from Thermo-scientific) was followed by a

10 cm analytical column of 75 μm i.d. (SC200 Easy Column from Thermo-scientific).

Both trap column and analytical column had C18-AQ packaging. Separation was carried

out using Easy nanoLC II (Thermo-Scientific) with a continuous, vented column

configuration. A 5 μL (~500 ng) sample was aspirated into a 20 μL sample loop and

loaded onto the trap. The flow rate was set to 300 nL/min for separation on the

analytical column. Mobile phase A was composed of 99.9 H_2O (EMD Omni Solvent),

and 0.1% formic acid and mobile phase B was composed of 99.9% ACN, and 0.1%

formic acid. A 1 h linear gradient from 0% to 45% B was performed. The LC eluent was

directly nano-sprayed into an LTQ Orbitrap Velos mass spectrometer (Thermo

Scientific). During the chromatographic separation, the LTQ Orbitrap Velos was

operated in a data-dependent mode and under direct control of the Xcalibur software

(Thermo Scientific). The MS data were acquired using the following parameters: 10

data-dependent collisional-induced-dissociation (CID) MS/MS scans per full scan. All

measurements were performed at room temperature and three technical replicates. The

raw files were analyzed using Proteome Discoverer (version 1.4) software package with

SequestHT and Mascot search nodes using maize proteome database available at

Gramene (AGVP3.0). The resulting msf files were further analyzed by the 'Scaffold

version 4.4' the proteome validator software at 90% peptide and protein threshold with a

minimum of 1 peptide.

Constructs and gene cloning

Coding sequences of MLKP1 and MLKP2 genes fused to C-terminus of eGFP-FLAG-

HA sequence were custom synthesized by Genscript as pHWBF08 and pHWBF06

respectively. eGFP-FLAG-HA-MLKP2 and eGFP-FLAG-HA-MLKP2 sequences were

PCR amplified with att flanking primers and cloned individually into pDONR221 vector

by BP cloning (Cat. No. 1235019, Invitrogen) to generate pHWBF08EC and

pHWBF06EC entry clones respectively. These were moved to destination vector

pH7WG2 (Karimi et al., 2002) by LR recombination reaction of Gateway cloning system

(Cat. No. 1235019, Invitrogen) to obtain expression vectors pPK1iexp and pPK1Cexp

respectively. For MLKG1, coding region was PCR amplified from maize B73 using gene

specific primers (ST3) and cloned into pCRII vector by TOPO cloning (Cat. No.

K461020, Thermo Fisher). After sequence verification, the coding sequence was

amplified with forward primer flanked with BamHI restriction site sequence and reverse

primer flanked by SbfI restriction site sequence. The PCR amplified product was double

digested with BamHI and SbfI (NEB) and ligated in pECGFP (Bass lab) donor vector in

frame and downstream of eGFP-FLAG-HA sequence. The fusion sequence was then

moved to pH7WG2 destination vector by LR recombination to generate expression

vector called pPK1Dexp. Deletion constructs for MLKP1 and MLKP2 were generated

using corresponding full length entry clones as template to PCR amplify required

regions with internal primers containing flanking att sequences (ST3) without a stop

codon and cloned into pDONR211 by BP cloning. These deletion constructs were

subcloned in pH7WG2 destination vector by LR recombination and the final expression

cassettes contains additional 38 aa residues

"NPAFLYKVVISRGHARVRKNHQSLSTNLSLSIFLQNNV" from the non-genic region of

the vector backbone at their C-termini.

MLKG1ΔKASH was generated using Q5 DNA mutagenesis kit (Cat. No. E0554S,

NEB). Briefly, two primers were designed in opposite directions, going outwards from

KASH domain. The entire entry vector (pECPK1D) minus the KASH domain was

amplified using high fidelity Q5 polymerase provided with the kit. The template

contamination was removed and vector re-ligated using manufacturer provided enzyme

mix containing a kinase, a ligase and DpnI to create entry clone pPK1DdelKASH. The

entry clone were then moved to pH7WG2 destination vector by LR recombination.

Agrobacterium tumefaciens **transformation**

A. tumefaciens was transformed with expression vectors using established heat shock

protocols (Graumann and Evans, 2010). Briefly, competent cells were thawed on ice

and 4ul plasmid DNA was added. Cells were incubated for 5 min on ice, 5 min in liquid

nitrogen and 5 min at 37 °C 1 ml LB medium was added and cells were incubated for 2h

at 28 °C before being plated out and left to grow for 48 h at 28 °C.

Nicotiana benthamiana **transformation and microscopy**

Transient expression of fluorescent fusion proteins was carried out as previously

described (Sparkes et al., 2006). Briefly, Agrobacteria were suspended in infiltration

buffer at an OD of 0.05 and pressed into the lower leaf epidermis with a syringe. 3 days

post infiltration, leaf sections were imaged with a Zeiss LSM 800 confocal microscope

using a 100X oil immersion objective, NA 1.46 and 1.5X digital zoom. GFP was excited

at 488 nm and emission collected between 500-550 nm, mCherry was excited with a

561 nm laser and emission collected from 565-620 nm. For FRAP, 5 pre scans and then

a 160x160 pixel ROI is bleached with the 488 nm laser at 100% for 10 iterations.

Recovery data was collected over 36 s (240 scans, 0.13 s framerate). Mean ROI

intensity data was extracted using Zen Blue (version 2.3) and normalised as described

in (Martinière et al., 2012). One phase association recovery curves, plateau values and

halftimes were produced in Graphpad (version 6). ANOVA statistical test with multiple

comparisons was performed. For DAPI staining, leaf sections were first incubated in 1

ml water containing 5 ul 20% Triton X-100 and 1ul DAPI. DAPI excitation was

performed with the 405 nm laser and emission between 400-450 nm imaged. DAPI/GFP

or DAPI/mCherry imaging was performed using multitrack line scanning.

Direct Co-IP and western blot

1 g of *N. benthamiana* leaves were harvested and ground in liquid nitrogen.

The powdered tissue was homogenized in 5 mL of radioimmunoprecipitation assay

buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% SDS, 0.5% Na-deoxycholate, 1%

NP-40, 1 mM PMSF, and 1% protease inhibitor cocktail (Cat. No. P9599, Sigma) in a

dounce homogenizer kept on ice. The homogenate was centrifuged at 10,000 xg and

the supernatant was collected for co-IP. One tenth of the protein extracts was used as

the input sample and concentrated using acetone precipitation. The rest of the lysate

was subjected to co-IP using dynabeads coated with rabbit anti-GFP antibody (Cat. No.

ab290, Abcam). The lysate was incubated with beads for 4 hours after which the beads

were collected using magnetic rack and washed with PBST three times. The antibody-

protein complex bound to beads was eluted using 50 mM Glycine (pH 2.8). The

immunoprecipitates and the input samples were boiled in 4X Laemelli's gel loading

buffer, fractionated on 15% SDS-PAGE and transferred to polyvinylidene difluoride

membranes (Cat. No. 88518. Thermo Scientific), and detected with a mouse anti-GFP

(1:3,000; Cat No. MA5-15256,Thermo Scientific) or rabbit anti-SUN2-NPAP (1:1,000;

Murphy et al 2014) antibody at room temperature for 1 h. After three 15-min washes in

PBS-T buffer at room temperature, the membranes were incubated with a 1:3,000

dilution of anti-rabbit or anti-mouse IgG horseradish peroxidase–linked antibody (Santa

Cruz Biotechnology, Santa Cruz, CA) for 1 h at room temperature, then subjected to 3 ×

15 min washes in PBST buffer at room temperature. The PVDF membrane was

incubated with chemiluminescent reaction kit for 3-5 min at room temperature (Millipore,

Immobilon detection kit, WBKL50100, Billerica, MA) and exposed to X-ray film in dark

room to visualize the protein bands.

FIGURE LEGENDS

Figure 1. **Working model of the plant LINC complex; core and associated**

proteins. Core LINC and associated proteins are depicted as residing in one of the four

cellular locations, each represented by different background shaded colors: green,

cytoplasm (CYTO); yellow, outer nuclear membrane (ONM); pink inner nuclear

membrane (INM); and blue NUC (nucleoplasm). The core LINC proteins (dashed box

enclosed) shown include SUN-domain proteins labeled "SUN", KASH-related proteins

"KASH/MLK", and AtWIT-related proteins labeled "WIT/MLKT". The LINC-associated

proteins shown include cytoplasmic proteins labeled "RanGAP" and "motor proteins",

INM proteins labeled "NEAP/MNEAP", and nucleoplasmic proteins labeled

"NMCP/CRWN/NCH" and "KAKU4/MKAKU4". Maize gene names and their known

Arabidopsis homolog types are listed. This composite model places proteins together on

the basis of previously documented pair-wise interactions representing a generalized

consensus assembly but not meant to depict a single complex.

Figure 2. Domain organization of candidates for the maize core LINC and

associated proteins complex. Schematic representations of predicted proteins from

canonical transcript models are diagrammed to show the location of domains (color

coded and listed at bottom). The proteins are drawn to scale, using a separate scale for

the last two larger proteins, NCH1 and NCH2. The four C-terminal residues and stop

codon (*) location of the KASH and MNEAP group proteins are indicated. The

previously published SUN-domain proteins (Murphy et al., 2010) are not included. The

partial SMC N-terminal domains in the MNEAPs are not diagrammed but are located at

AA 24-308 for MNEAP1, AA 18-309 for MNEAP2, and at AA 3-109 for MNEAP3.

Potential CDK1 phosphorylation sites are shown (blue circles) for NCH1 and NCH2.

Figure 3. Evolutionarily conservation of plant LINC complex components.

Candidate maize LINC gene names, cellular location predictions, gene family group,

and PLAZA gene family ID (each starting with shared prefix of HOMO4M) are listed in

the first four columns.   The other columns provide counts of the numbers of genes for

each species (top row) in each Plaza gene family (4th column). The general grouping

color key is indicated on the top left of the figure. Species included are Zema, *Zea*

mays; Sobi*, Sorghum bicolor,* Seit, *Setaria viridis;* Hovu, *Hordeum vulgare;*

Trae,*Triticum aestivum;* Brdi, *Brachypodium distachyon;* Orsa*, Oryza sativa spp.*

Japonica; Zoma*, Zostera marina;* Arth*, Arabidopsis thaliana;* Potr*, Populus trichocarpa;*

Vivi*, Vitis vinifera;* Metr*, Medicago truncatula;* Piab*, Picea abies;* Phpa*, Physcomitrella*

patens and Chre, *Chlamydomonas reinhardtii.* Broader grouping names are provided

along the top (labeled grey boxes).

Figure 4: Gene expression levels and co-expression clustering of 20 maize LINC

genes. Expression levels of 20 maize candidate LINC genes were analyzed using

published transcriptome data (Stelpflug et al., 2016) and including the genes

designated as housekeeping genes (HKG) in a previous study (Lin et al., 2014). (A)

Expression profiles of LINC genes in several tissues developmental stages. Data are

presented as log10 Fragments Per Kilobase of exon per Million fragments mapped

(FPKM) values from published transcriptome. Hierarchical clustering grouped tissues

according to similar expression profiles. The root, young inflorescence, and meristem

groups are denoted by orange, blue, and green boxes, respectively. The growth stage

of maize plant at the time of tissue harvesting are shown in parentheses; V, vegetative

stage; R, reproductive stage; DAP, days after pollination. (B) The Median FPKM values

of the tissues shown in panel A are plotted, ranked by value for individual genes.

Housekeeping genes are shown in red.

Figure 5. Subcellular localisation of transiently expressed GFP-MLKP fusions in

N. benthamiana. Agrobacterium-mediated leaf transformation with expression

constructs was carried out followed by imaging three days later using 100X lense with

confocal microscopy to detect DAPI (405 nm excitation laser), GFP (488 nm excitation

laser), or mCherry (561 nm excitation laser). Representative focal sections are shown

for GFP (green) or mCherry (purple). (A) GFP-MLKG1, GFP-MLKP1, GFP-MLKP2 and

mCherry-ZmSUN2 (green) are localised at the nuclear periphery, surrounding the

nucleus (n) stained with DAPI (magenta). (B**)** GFP-MLKG1 (green) is also present in a

network-like pattern at the cell periphery, where it partially co-localises with the ER

marker HDEL-mRFP (magenta, top row) and the actin marker lifeact-mRFP (magenta,

bottom row). (C**)** Schematic overview depicting the terminal deletions that were replaced

with 38 aa extension, $(N)_{38}$. (D) GFP-MLKP2Δ4r, GFP-MLKP2ΔKASHr and GFP-

MLKP2Δ4TMr (green) localise in the cytoplasm and the DAPI-stained nucleoplasm

(magenta). All scale bars 10μm.

Figure 6. Fluorescence recovery after photobleaching (FRAP) co-expression

interaction assays of MLK LINC candidates with ZmSUN2 in *N. benthamiana*. (A)

Schematic overview depicting the full length mCherry-ZmSUN2 and internal deletion

constructs that removed the coiled coil domain (ΔCC) or the SUN domain (ΔSUN).

Normalized averaged (n=26-37) intensity FRAP curves for (B) GFP-MLKG1, (D) GFP-

MKLP1, or (F) GFP-MLKP2 expressed alone (green), with full-length mCherry-SUN2

(black), with mCherry-SUN2-ΔCC (blue), or with mCherry-SUN2-ΔSUN (pink). Plateau

values are plotted to the right for assays with GFP-MLKG1 (C), GFP-MLKP1 (E), and

GFP-MLKP2 (G). Box plot error bars represent SD (blue) and mean (red) values.

ANOVA statistical analyses with multiple comparisons are noted (ns = not significant =

p≥0.05, **** = significant with P ≤0.0001).

Supplementary Figure 1. Subcellular localisation of transiently expressed GFP-

MLKp domain deletion fusions in *N. benthamiana*. Full-length, domain deletions or

replacements, and GFP only controls were assayed as described in Figure 5.   (A)

Schematic overview of domain composition of the constructs.   Constructs are listed at

the left of each row for images take through the middle of the nuclei (n in DAPI,

magenta). The right-most images (Cell periphery) were taken at a different focal plane

in order to image the cytoplasm near the cell cortex. Free GFP (bottom row, green) is

observed in the cytoplasm and, to a lesser extent, in the DAPI-stained nucleoplasm (

magenta). All scale bars 10μm.

Supplementary Figure 2. FRAP images from the co-expression interaction

assays. Representative images of cells during the FRAP assays for the experiments

summarized in Figure 6. The FRAP ROIs (white boxes) are shown along with images

taken at the times indicated across the top for (A) GFP-MLKG1, (B) GFP-MLKP1, and

(C) GFP-MLKP2. Scale bar denotes 2µm. FRAP ROI shown by white box. All scale

bars 2µm.

Supplementary Figure 3. Normalised intensity FRAP curves co-plotted from all

experiments performed in *N. benthamiana* leaves. The GFP fusion protein

constructs (MLKG1, MKLP1, and MLKP2) are grouped by columns as labeled at top.

The co-expression construct are grouped by row as labeled at left. Individual data

points (blue dots), single experiment recovery curves (black lines) and global average

recovery curves (red lines) and replicate numbers (n) are shown in each panel.

Supplementary Figure 4. FRAP recovery halftime values. The recovery half times

(t1/2(s)) are plotted for the full-length and co-expression experiments summarized in

Figure 6. The SD (blue lines) and mean (red lines) are shown for (A) GFP-MLKG1, (B)

GFP-MLKP1, and (C) GFP-MLKP2. ANOVA statistical analyses with multiple

comparisons are noted (ns = not significant = p≥0.05, ** = significant with P ≤0.01, ****

= significant with P ≤0.0001). Replicate numbers range from n=26 to n=35, as listed in

Supplementary Figure 3.

Supplementary Figure 5. Co-immunoprecipitation assays of MLKP1 and MLKP2

with ZmSUN2 in *N. benthamiana*. Total protein extracts from *N. benthamiana* leaves

harvested 3 days post-infection were subjected to co-IP using dynabeads coated with

anti-GFP antibody.   (A) Western Blot of the co-precipitates and inputs with primary

antibodies listed on the right side of the blot. The FP construct combinations used for

infiltration are listed on the top of the blot. The numbers on the left are protein molecular

weight standards. (B) A summary of KASH-SUN interaction evidence tabulated from

this experiment (Heterologous co-IP) or other experiments in this study (Native co-IP

from Table 2, Heterologous FRAP form Figure 6 and Supplemental Figures 2-4).

ACKNOWLEDGEMENTS

Thank Eric Richards and Iris Meier for helpful and insightful comments on the

manuscript, Allison Jevitt for artwork on Figure 1, Dr. Rakesh K Singh and staff of the

Translational lab (College of Medicine, FSU) for assistance and advice on Co-IP and

Mass spectrometry. The work was supported by grants from the National Science

Foundation to HWB (NSF IOS 1444532) and Oxford Brookes University to KG, and

from a Dissertation Research Grant to HKG (Florida State University Graduate School).

Competing interests

949   The authors declare no competing or financial interests

REFERENCES

Belmont, A. S., Sedat, J. W. and Agard, D. A. (1987). A three-dimensional approach

to mitotic chromosome structure: evidence for a complex hierarchical organization.

J. Cell Biol. **105**, 77–92.

Boruc, J., Zhou, X. and Meier, I. (2012). Dynamics of the plant nuclear envelope and
nuclear pore. *Plant Physiol.* **158**, 78–86.

Broers, J. L. V., Ramaekers, F. C. S., Bonne, G., Yaou, R. B. and Hutchison, C. J.
(2006). Nuclear lamins: laminopathies and their role in premature ageing. *Physiol.*
Rev. **86**, 967–1008.

Burke, B. and Stewart, C. L. (2014). Chapter One - Functional Architecture of the
Cell's Nucleus in Development, Aging, and Disease. In *Current Topics in*
Developmental Biology (ed. Stewart, C. L.), pp. 1–52. Academic Press.

Chettoor, A. M., Givan, S. A., Cole, R. A., Coker, C. T., Unger-Wallace, E.,
Vejlupkova, Z., Vollbrecht, E., Fowler, J. E. and Evans, M. M. (2014). Discovery
of novel transcripts and gametophytic functions via RNA-seq analysis of maize
gametophytic transcriptomes. *Genome Biol.* **15**, 414.

Ciska, M., Masuda, K. and Moreno Díaz de la Espina, S. (2013). Lamin-like
analogues in plants: the characterization of NMCP1 in Allium cepa. *J. Exp. Bot.* **64**,
1553–1564.

Coates, J. C. (2003). Armadillo repeat proteins: beyond the animal kingdom. *Trends*
Cell Biol. **13**, 463–471.

Crisp, M., Liu, Q., Roux, K., Rattner, J. B., Shanahan, C., Burke, B., Stahl, P. D. and
Hodzic, D. (2006). Coupling of the nucleus and cytoplasm: role of the LINC
complex. *J. Cell Biol.* **172**, 41–53.

Dittmer, T. A. and Richards, E. J. (2008). Role of LINC proteins in plant nuclear
morphology. *Plant Signal. Behav.* **3**, 485–487.

Fridkin, A., Penkner, A., Jantsch, V. and Gruenbaum, Y. (2009). SUN-domain and
KASH-domain proteins during development, meiosis and disease. *Cell. Mol. Life*
Sci. **66**, 1518–1533.

Friederichs, J. M., Ghosh, S., Smoyer, C. J., McCroskey, S., Miller, B. D., Weaver,
K. J., Delventhal, K. M., Unruh, J., Slaughter, B. D. and Jaspersen, S. L. (2011).

The SUN protein Mps3 is required for spindle pole body insertion into the nuclear
membrane and nuclear envelope homeostasis. *PLoS Genet.* **7**, e1002365.

Gaut, B. S., Le Thierry d'Ennequin, M., Peek, A. S. and Sawkins, M. C. (2000).
Maize as a model for the evolution of plant nuclear genomes. *Proc. Natl. Acad. Sci.*
U. S. A. **97**, 7008–7015.

Gerber, A. P., Herschlag, D. and Brown, P. O. (2004). Extensive association of
functionally and cytotopically related mRNAs with Puf family RNA-binding proteins in
yeast. *PLoS Biol.* **2**, E79.

Goto, C., Tamura, K., Fukao, Y., Shimada, T. and Hara-Nishimura, I. (2014). The
Novel Nuclear Envelope Protein KAKU4 Modulates Nuclear Morphology in
Arabidopsis. *Plant Cell* **26**, 2143–2155.

Graumann, K. and Evans, D. E. (2010). Plant SUN domain proteins: components of
putative plant LINC complexes? *Plant Signal. Behav.* **5**, 154–156.

Graumann, K., Runions, J. and Evans, D. E. (2010). Characterization of SUN-domain
proteins at the higher plant nuclear envelope. *Plant J.* **61**, 134–144.

Graumann, K., Vanrobays, E., Tutois, S., Probst, A. V., Evans, D. E. and Tatout, C.
(2014). Characterization of two distinct subfamilies of SUN-domain proteins in
Arabidopsis and their interactions with the novel KASH-domain protein AtTIK. *J.*
Exp. Bot. **65**, 6499–6512.

Grob, S., Schmid, M. W. and Grossniklaus, U. (2014). Hi-C analysis in Arabidopsis
identifies the KNOT, a structure with similarities to the flamenco locus of Drosophila.
Mol. Cell **55**, 678–693.

Groves, N. R., Biel, A. M., Newman-Griffis, A. H. and Meier, I. (2018). Dynamic
Changes in Plant Nuclear Organization in Response to Environmental and
Developmental Signals. *Plant Physiol.* **176**, 230–241.

Hagan, I. and Yanagida, M. (1995). The product of the spindle formation gene sad1+
associates with the fission yeast spindle pole body and is essential for viability. *J.*

Cell Biol. **129**, 1033–1047.

Haque, F., Lloyd, D. J., Smallwood, D. T., Dent, C. L., Shanahan, C. M., Fry, A. M.,
Trembath, R. C. and Shackleton, S. (2006). SUN1 interacts with nuclear lamin A
and cytoplasmic nesprins to provide a physical connection between the nuclear
lamina and the cytoskeleton. *Mol. Cell. Biol.* **26**, 3738–3751.

Hirsch, C. N., Hirsch, C. D., Brohammer, A. B., Bowman, M. J., Soifer, I., Barad, O.,
Shem-Tov, D., Baruch, K., Lu, F., Hernandez, A. G., et al. (2016). Draft Assembly
of Elite Inbred Line PH207 Provides Insights into Genomic and Transcriptome
Diversity in Maize. *Plant Cell* **28**, 2700–2714.

Hodzic, D. M., Yeater, D. B., Bengtsson, L., Otto, H. and Stahl, P. D. (2004). Sun2 is
a novel mammalian inner nuclear membrane protein. *J. Biol. Chem.* **279**, 25805–
25812.

Horigome, C., Okada, T., Shimazu, K., Gasser, S. M. and Mizuta, K. (2011).
Ribosome biogenesis factors bind a nuclear envelope SUN domain protein to
cluster yeast telomeres. *EMBO J.* **30**, 3799–3811.

Hotta, T., Haraguchi, T. and Mizuno, K. (2007). A novel function of plant histone H1:
microtubule nucleation and continuous plus end association. *Cell Struct. Funct.* **32**,
79–87.

Jafferali, M. H., Vijayaraghavan, B., Figueroa, R. A., Crafoord, E., Gudise, S.,
Larsson, V. J. and Hallberg, E. (2014). MCLIP, an effective method to detect
interactions of transmembrane proteins of the nuclear envelope in live cells.
Biochim. Biophys. Acta **1838**, 2399–2403.

Janin, A., Bauer, D., Ratti, F., Millat, G. and Méjat, A. (2017). Nuclear envelopathies:
a complex LINC between nuclear envelope and pathology. *Orphanet J. Rare Dis.*
12, 147.

Jiao, Y., Peluso, P., Shi, J., Liang, T., Stitzer, M. C., Wang, B., Campbell, M. S.,
Stein, J. C., Wei, X., Chin, C.-S., et al. (2017). Improved maize reference genome

with single-molecule technologies. *Nature* **546**, 524–527.

Käll, L., Krogh, A. and Sonnhammer, E. L. L. (2004). A combined transmembrane
topology and signal peptide prediction method. *J. Mol. Biol.* **338**, 1027–1036.

Karimi, M., Inzé, D. and Depicker, A. (2002). GATEWAY^TM vectors for Agrobacterium-
mediated plant transformation. *Trends Plant Sci.* **7**, 193–195.

Kim, D. I., Birendra, K. C. and Roux, K. J. (2015). Making the LINC: SUN and KASH
protein interactions. *Biol. Chem.* **396**, 295–310.

Kopylov, M., Bass, H. W. and Stroupe, M. E. (2015). The maize (Zea mays L.)
nucleoside diphosphate kinase1 (ZmNDPK1) gene encodes a human NM23-H2
homologue that binds and stabilizes G-quadruplex DNA. *Biochemistry* **54**, 1743–
1757.

Kracklauer, M. P., Link, J. and Alsheimer, M. (2013). LINCing the nuclear envelope to
gametogenesis. *Curr. Top. Dev. Biol.* **102**, 127–157.

Krogh, A., Larsson, B., Von Heijne, G. and Sonnhammer, E. L. L. (2001). Predicting
transmembrane protein topology with a hidden markov model: application to
complete genomes1. *J. Mol. Biol.* **305**, 567–580.

Kumar, S., Stecher, G. and Tamura, K. (2016). MEGA7: Molecular Evolutionary
Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874.

Li, P. and Noegel, A. A. (2015). Inner nuclear envelope protein SUN1 plays a
prominent role in mammalian mRNA export. *Nucleic Acids Res.* **43**, 9874–9888.

Lin, F., Jiang, L., Liu, Y., Lv, Y., Dai, H. and Zhao, H. (2014). Genome-wide
identification of housekeeping genes in maize. *Plant Mol. Biol.* **86**, 543–554.

Liu, W., Xie, Y., Ma, J., Luo, X., Nie, P., Zuo, Z., Lahrmann, U., Zhao, Q., Zheng, Y.,
Zhao, Y., et al. (2015). IBS: an illustrator for the presentation and visualization of
biological sequences. *Bioinformatics* **31**, 3359–3361.

Luxton, G. W. G. and Starr, D. A. (2014). KASHing up with the nucleus: novel

functional roles of KASH proteins at the cytoplasmic surface of the nucleus. *Curr.*
Opin. Cell Biol. **28**, 69–75.

Malone, C. J., Fixsen, W. D., Horvitz, H. R. and Han, M. (1999). UNC-84 localizes to
the nuclear envelope and is required for nuclear migration and anchoring during C.
elegans development. *Development* **126**, 3171–3181.

Martinière, A., Lavagi, I., Nageswaran, G., Rolfe, D. J., Maneta-Peyret, L., Luu, D.-
T., Botchway, S. W., Webb, S. E. D., Mongrand, S., Maurel, C., et al. (2012). Cell
wall constrains lateral diffusion of plant plasma-membrane proteins. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 12805–12810.

Masuda, K., Xu, Z. J., Takahashi, S., Ito, A., Ono, M., Nomura, K. and Inoue, M.
(1997). Peripheral framework of carrot cell nucleus contains a novel protein
predicted to exhibit a long alpha-helical domain. *Exp. Cell Res.* **232**, 173–181.

Meier, I. (2016). LINCing the eukaryotic tree of life - towards a broad evolutionary
comparison of nucleocytoplasmic bridging complexes. *J. Cell Sci.* **129**, 3523–3531.

Meier, I., Zhou, X., Brkljacić, J., Rose, A., Zhao, Q. and Xu, X. M. (2010). Targeting
proteins to the plant nuclear envelope. *Biochem. Soc. Trans.* **38**, 733–740.

Meier, I., Richards, E. J. and Evans, D. E. (2017). Cell Biology of the Plant Nucleus.
Annu. Rev. Plant Biol. **68**, 139–172.

Mochizuki, R., Tsugama, D., Yamazaki, M., Fujino, K. and Masuda, K. (2017).
Identification of candidates for interacting partners of the tail domain of DcNMCP1, a
major component of the Daucus carota nuclear lamina-like structure. *Nucleus* **8**,
312–322.

Moriguchi, K., Suzuki, T., Ito, Y., Yamazaki, Y., Niwa, Y. and Kurata, N. (2005).
Functional isolation of novel nuclear proteins showing a variety of subnuclear
localizations. *Plant Cell* **17**, 389–403.

Murphy, S. P., Simmons, C. R. and Bass, H. W. (2010). Structure and expression of
the maize (Zea mays L.) SUN-domain protein gene family: evidence for the

existence of two divergent classes of SUN proteins in plants. *BMC Plant Biol.* **10**,
269.

Murphy, S. P., Gumber, H. K., Mao, Y. and Bass, H. W. (2014). A dynamic meiotic
SUN belt includes the zygotene-stage telomere bouquet and is disrupted in
chromosome segregation mutants of maize (Zea mays L.). *Front. Plant Sci.* **5**, 314.

Nakayama, T., Ishii, T., Hotta, T. and Mizuno, K. (2008). Radial microtubule
organization by histone H1 on nuclei of cultured tobacco BY-2 cells. *J. Biol. Chem.*
283, 16632–16640.

Nannas, N. J. and Dawe, R. K. (2015). Genetic and genomic toolbox of Zea mays.
Genetics **199**, 655–669.

Oda, Y. and Fukuda, H. (2011). Dynamics of Arabidopsis SUN proteins during mitosis
and their involvement in nuclear shaping. *Plant J.* **66**, 629–641.

Pawar, V., Poulet, A., Détourné, G., Tatout, C., Vanrobays, E., Evans, D. E. and
Graumann, K. (2016). A novel family of plant nuclear envelope-associated proteins.
J. Exp. Bot. **67**, 5699–5710.

Poulet, A., Probst, A. V., Graumann, K., Tatout, C. and Evans, D. (2017a). Exploring
the evolution of the proteins of the plant nuclear envelope. *Nucleus* **8**, 46–59.

Poulet, A., Duc, C., Voisin, M., Desset, S., Tutois, S., Vanrobays, E., Benoit, M.,
Evans, D. E., Probst, A. V. and Tatout, C. (2017b). The LINC complex contributes
to heterochromatin organisation and transcriptional gene silencing in plants. *J. Cell*
Sci. **130**, 590–601.

Razafsky, D. and Hodzic, D. (2015). Nuclear envelope: positioning nuclei and
organizing synapses. *Curr. Opin. Cell Biol.* **34**, 84–93.

Shimada, N., Inouye, K., Sawai, S. and Kawata, T. (2010). SunB, a novel Sad1 and
UNC-84 domain-containing protein required for development of Dictyostelium
discoideum. *Dev. Growth Differ.* **52**, 577–590.

Sohaskey, M. L., Jiang, Y., Zhao, J. J., Mohr, A., Roemer, F. and Harland, R. M.
(2010). Osteopotentia regulates osteoblast maturation, bone formation, and skeletal
integrity in mice. *J. Cell Biol.* **189**, 511–525.

Sosa, B. A., Rothballer, A., Kutay, U. and Schwartz, T. U. (2012). LINC complexes
form by binding of three KASH peptides to domain interfaces of trimeric SUN
proteins. *Cell* **149**, 1035–1047.

Sparkes, I. A., Runions, J., Kearns, A. and Hawes, C. (2006). Rapid, transient
expression of fluorescent fusion proteins in tobacco plants and generation of stably
transformed plants. *Nat. Protoc.* **1**, 2019–2025.

Spichal, M. and Fabre, E. (2017). The Emerging Role of the Cytoskeleton in
Chromosome Dynamics. *Front. Genet.* **8**, 60.

Starr, D. A. and Fridolfsson, H. N. (2010). Interactions between nuclei and the
cytoskeleton are mediated by SUN-KASH nuclear-envelope bridges. *Annu. Rev.*
Cell Dev. Biol. **26**, 421–444.

Starr, D. A. and Han, M. (2002). Role of ANC-1 in tethering nuclei to the actin
cytoskeleton. *Science* **298**, 406–409.

Stelpflug, S. C., Sekhon, R. S., Vaillancourt, B., Hirsch, C. N., Buell, C. R., de Leon,
N. and Kaeppler, S. M. (2016). An Expanded Maize Gene Expression Atlas based
on RNA Sequencing and its Use to Explore Root Development. *Plant Genome* **9**,.

Tamura, K., Fukao, Y., Iwamoto, M., Haraguchi, T. and Hara-Nishimura, I. (2010).
Identification and characterization of nuclear pore complex components in
Arabidopsis thaliana. *Plant Cell* **22**, 4084–4097.

Tamura, K., Iwabuchi, K., Fukao, Y., Kondo, M., Okamoto, K., Ueda, H., Nishimura,
M. and Hara-Nishimura, I. (2013). Myosin XI-i links the nuclear membrane to the
cytoskeleton to control nuclear movement and shape in Arabidopsis. *Curr. Biol.* **23**,
1776–1781.

Tamura, K., Goto, C. and Hara-Nishimura, I. (2015). Recent advances in

understanding plant nuclear envelope proteins involved in nuclear morphology. *J.*

Exp. Bot. **66**, 1641–1647.

Tapley, E. C. and Starr, D. A. (2013). Connecting the nucleus to the cytoskeleton by

SUN-KASH bridges across the nuclear envelope. *Curr. Opin. Cell Biol.* **25**, 57–62.

Tiang, C.-L., He, Y. and Pawlowski, W. P. (2012). Chromosome organization and

dynamics during interphase, mitosis, and meiosis in plants. *Plant Physiol.* **158**, 26–

34.

Van Bel, M., Diels, T., Vancaester, E., Kreft, L., Botzki, A., Van de Peer, Y.,

Coppens, F. and Vandepoele, K. (2018). PLAZA 4.0: an integrative resource for

functional, evolutionary and comparative plant genomics. *Nucleic Acids Res.* **46**,

D1190–D1196.

Varas, J., Graumann, K., Osman, K., Pradillo, M., Evans, D. E., Santos, J. L. and

Armstrong, S. J. (2015). Absence of SUN1 and SUN2 proteins in Arabidopsis

thaliana leads to a delay in meiotic progression and defects in synapsis and

recombination. *Plant J.* **81**, 329–346.

Vera, D. L., Madzima, T. F., Labonne, J. D., Alam, M. P., Hoffman, G. G.,

Girimurugan, S. B., Zhang, J., McGinnis, K. M., Dennis, J. H. and Bass, H. W.

(2014). Differential nuclease sensitivity profiling of chromatin reveals biochemical

footprints coupled to gene expression and functional DNA elements in maize. *Plant*

Cell **26**, 3883–3893.

Walley, J. W., Sartor, R. C., Shen, Z., Schmitz, R. J., Wu, K. J., Urich, M. A., Nery, J.

R., Smith, L. G., Schnable, J. C., Ecker, J. R., et al. (2016). Integration of omic

networks in a developmental atlas of maize. *Science* **353**, 814–818.

Wang, H., Dittmer, T. A. and Richards, E. J. (2013). Arabidopsis CROWDED NUCLEI

(CRWN) proteins are required for nuclear size control and heterochromatin

organization. *BMC Plant Biol.* **13**, 200.

Wang, B., Tseng, E., Regulski, M., Clark, T. A., Hon, T., Jiao, Y., Lu, Z., Olson, A.,

Stein, J. C. and Ware, D. (2016). Unveiling the complexity of the maize

transcriptome by single-molecule long-read sequencing. *Nat. Commun.* **7**, 11708.

Xu, X. M., Meulia, T. and Meier, I. (2007). Anchorage of plant RanGAP to the nuclear

envelope involves novel nuclear-pore-associated proteins. *Curr. Biol.* **17**, 1157–

1163.

Xu, X. H., Chen, H., Sang, Y. L., Wang, F., Ma, J. P., Gao, X.-Q. and Zhang, X. S.

(2012). Identification of genes specifically or preferentially expressed in maize silk

reveals similarity and diversity in transcript abundance of different dry stigmas. *BMC*

Genomics **13**, 294.

Zhao, Q., Brkljacic, J. and Meier, I. (2008). Two distinct interacting classes of nuclear

envelope-associated coiled-coil proteins are required for the tissue-specific nuclear

envelope targeting of Arabidopsis RanGAP. *Plant Cell* **20**, 1639–1651.

Zhou, X., Graumann, K., Evans, D. E. and Meier, I. (2012). Novel plant SUN–KASH

bridges are involved in RanGAP anchoring and nuclear shape determination. *J. Cell*

Biol. **196**, 203–211.

Zhou, X., Graumann, K., Wirthmueller, L., Jones, J. D. G. and Meier, I. (2014).

Identification of unique SUN-interacting nuclear envelope proteins with diverse

functions in plants. *J. Cell Biol.* **205**, 677–692.

Zhou, X., Graumann, K. and Meier, I. (2015a). The plant nuclear envelope as a

multifunctional platform LINCed by SUN and KASH. *J. Exp. Bot.* **66**, 1649–1659.

Zhou, X., Groves, N. R. and Meier, I. (2015b). Plant nuclear shape is independently

determined by the SUN-WIP-WIT2-myosin XI-i complex and CRWN1. *Nucleus* **6**,

144–153.

Zhou, X., Groves, N. R. and Meier, I. (2015c). SUN anchors pollen WIP-WIT

complexes at the vegetative nuclear envelope and is necessary for pollen tube

targeting and fertility. *J. Exp. Bot.* **66**, 7299–7307.

1195

1196

Table 1: Maize LINC Genes

Maize Gene name[B]	Gene Identifier[A]					Protein Sequence Features					Domains		
	UniProt[C]	B73v3[D]	B73v4[D]	PLAZA gene[E]	Plaza gene family[E]	N-terminal Sequence	C-terminal Sequence	Total length (AA)	Molecular weight (kDa)	pI[G]	CC	TMD	Other (location)
MLKG1	A0A096TG70	GRMZM2G370707	Zm00001d002723	ZM02G25900	HOM04M009187	MDMPP	VVPT*	258	28	4.82	74- 94	224 - 246	KASH (247 - 258)
MLKG2	A0A1D6M739	GRMZM2G117392	Zm00001d038539	ZM06G27180	HOM04M008346	MTKPT	VMPPT*	227	23.74	5.02	-	191 - 214	KASH (215 - 227)
MLKS1	A0A1D6KGN6	GRMZM2G094850	Zm00001d031134	ZM01G30030	HOM04M002474	MATSL	VVPT*	617	67.2	8.59	-	579 - 601	KASH (602-617)
MLKS2	K7UCH8	GRMZM2G000608	Zm00001d052955	ZM04G32760	HOM04M002474	MGRSL	LVPT*	637	68.49	6.55	-	605 - 623	KASH (624 - 637), ARM (3 - 517)
MLKP1	B4FDR2	GRMZM2G151418	Zm00001d003334	ZM02G12590	HOM04M002181	MSMNY	AVPT*	294	33.73	4.6	68 - 91, 109 -136, 149 - 183, 222 - 256	264 - 286	KASH (289 - 294)
MLKP2	K7TL67	GRMZM2G034882	Zm00001d025667	ZM10G18600	HOM04M002181	MLMHY	VVPT*	290	33.57	4.63	68 - 91, 109 -136, 152 -179	258 - 280	KASH (281 - 290)
MLKP3	B7ZXA8	GRMZM2G003992	Zm00001d005997	ZM02G32490	HOM04M002181	MDSGA	IVPT*	538	57.9	5.47	319 - 343, 358 - 378, 452 - 500	506 - 529	KASH (530 - 538)
MLKP4	C0PH11	GRMZM2G176407	Zm00001d020879	ZM07G17460	HOM04M002181	MDSGA	IVPT*	532	57.6	4.7	313 - 337, 352 - 372, 401 - 421, 446 - 480	502 - 523	KASH (524 - 532)
MLKT1	C0P4P9	GRMZM2G148723	Zm00001d039272	ZM03G00380	HOM04M000144	MDFHN	VNDC*	653	72.35	4.86	94-114, 135-155, 201-228, 374-415, 455-517	620-640	KASH (641 - 653)
MLKT2	K7UYZ4	GRMZM2G006661	Zm00001d010047	ZM08G14110	HOM04M000144	MDTVI	GSPA*	667	74.51	4.72	96-116, 130-157, 218-238, 316-392, 411-431, 453-473, 481-508, 527-547	643-661	KASH (662 - 667)
ZmSUN1	B6THU8	GRMZM2G109818	Zm00001d015450	ZM05G20890	HOM04M002694	MTMSA	PSTGG*	462	50.96	9.07	169 - 189, 195 - 229	116 - 141	SUN (277-455)
ZmSUN2	B6TY16	GRMZM2G440614	Zm00001d040331	ZM03G08780	HOM04M002694	MASPS	AGSQA*	439	47.5	7.78	168-188	84 - 109	SUN (264-478)
ZmSUN3	A0A1D6N5R4	GRMZM2G120262	Zm00001d042643	ZM03G25290	HOM04M001731	MQRSR	ILIIQ*	613	68.04	4.86	484 - 511	33 - 55, 554 - 574, 595 -612	SUN (194 - 358)
ZmSUN4	K7VAL9	GRMZM2G005483	Zm00001d012240	ZM08G30590	HOM04M001731	MSLSC	ILIIQ*	639	70.54	5.22	510 -537	57 - 79, 580 -600, 621 -638	SUN (218 - 382)
ZmSUN5	A0A1D6FYJ8	AC194341.4_FG003	Zm00001d011277	ZM08G23230	HOM04M001731	MSRKR	VLLYN*	607	65.86	5.46	414 - 434, 495 - 523	46 - 66, 525 -544, 572 - 588	SUN (152 - 322)
MNEAP1	B6TAT2	GRMZM2G042593	Zm00001d010918	ZM08G20470	HOM04M002962	MSVSE	VILAKR*	348	39.75	8.94	221 - 248	330-346	NA
MNEAP2	K7TRE8	GRMZM2G061728	Zm00001d024756	ZM10G11720	HOM04M002962	MQCLS	VILAKR*	347	39.73	8.97	149 - 83 220 - 247	329-345	NA
MNEAP3	B6TCY4	(absent)	Zm00001d002875	(absent)	HOM04M002962	MSLLK	FIHV*	170	19.89	8.94	14 – 34, 84-104	125-150	NA
NCH1	K7U7Q6	GRMZM2G015875	Zm00001d051600	ZM04G22450	HOM04M004640	MFTPQ	SFFTT*	1156	132.82	5.13	160 – 278, 312 – 350, 362 – 522, 537 – 579, 592 – 701	NA	
NCH2	A0A1D6NAM9	GRMZM2G320013	Zm00001d043335	ZM03G30810	HOM04M003564	MASPR	KMFKE*	896	105.04	5.25	131 – 165, 250 – 281, 309 – 336, 386 – 532, 635 – 721	NA	
MKAKU41	A0A1D6JGB5	GRMZM2G104199	Zm00001d026487	ZM10G25400	HOM04M004991	MASLF	RGRAT*	554	59.59	5.29	NA	NA	
MKAKU42	A0A1D6DVM2	GRMZM2G521862	Zm00001d002012	ZM02G01940	HOM04M004991	MASLF	EGRAT*	591	64.87	5.66	NA	NA	

FOOTNOTES

A	Gene IDs are listed as of 2017
B	New gene names MKLP, MLKS, Maize LINC KASH AtSINE-like; MLKG, Maize LINC KASH Grass-specific; Maize LINC KASH AtWIP-like: MLKT, Maize LINC KASH AtWIT-like
C	UniProt ID from the UnitProt Consortium as of December 2017 (Nucleic Acids Res. 45: D158-D169 (2017))
D	B73v3 and B73v4 gene models from MaizeGDB.org as of December 2017
E	Plaza gene and family IDs from version 4 of Plaza Gene (https://doi.org/10.1093/nar/gkx1002)
F	KASH domain refers to amino acids post transmembrane domain
G	pI, Isoelectric point calculated from the full-length protein using ExPASy Compute "pI/Mw"

Table 2: Proteins co-precipitated with ZmSUN2 pull down on maize ear shoot nuclei

	B73v4 Maize gene id	Maize gene name	Gene Description'	Exclusive unique peptide count	SUN IP (Normalized total spectra)	Irrelevant antibody control IP	No Antibody control IP
LINC components	Zm00001d040331	ZmSUN2	SUN domain protein2	11	384.45	0	0
	Zm00001d015450	ZmSUN1	SUN domain protein1	3	78.398	0	0
	Zm00001d051600	NCH1	Protein CROWDED NUCLEI 1	2	82.803	0	0
	Zm00001d043335	NCH2	Nuclear lamina component	3	28.871	0	0
	Zm00001d002723	MLKG1	OSJNBa0010H02.21-like protein	2	14.459	0	0
	Zm00001d052955	MLKS2	ARM repeat superfamily protein	2	1.2886	0	0
	Zm00001d010047	MLKT2	WPP domain-interacting tail-anchored protein 1	1	26.117	0	0
Nucleoporins	Zm00001d052282	NUA	Nuclear-pore anchor	7	87.161	0	0
	Zm00001d048023	NUP133	Nuclear pore complex protein NUP133	1	18.911	0	0
	Zm00001d046566	NUP96	Nuclear pore complex protein NUP96	1	7.253	0	0
	Zm00001d026263	NUP88	Nuclear pore complex protein NUP88	1	7.206	0	0
	Zm00001d005355	GP210	Nuclear pore complex protein GP210	1	7.206	0	0
Cytoskeletal elements	Zm00001d013410	Actin-1	ARP7 Nuclear actin	3	82.756	0	0
	Zm00001d012556	Tub6	beta- Tubulin 6	5	40.576	0	0
	Zm00001d015348	Tub4	beta- Tubulin 4	1	7.253	0	0
Ribosomal proteins	Zm00001d044287	L19	Ribosomal protein L19	2	40.623	0	0
	Zm00001d017047	L6	60S ribosomal protein L6	1	30.569	0	0
	Zm00001d022463	L22-2	60S ribosomal protein L22-2	1	11.658	0	0
RAN	Zm00001d017239	RanGAP2	RAN GTPase-activating protein 2	1	14.459	0	0
	Zm00001d048223	RABA6a	Ras-related protein RABA6a	2	14.412	0	0
DNA/RNA binding	Zm00001d052820	RNA binding protein	RNA-binding (RRM/RBD/RNP motifs) family protein	2	21.665	0	0
	Zm00001d025638	bZIP5	Basic-leucine zipper domain	1	11.658	0	0
	Zm00001d042267	ARF10	Auxin response factor 10	1	11.658	0	0
	Zm00001d011297	MYBR35	Putative MYB DNA-binding domain superfamily protein	1	7.206	0	0
	Zm00001d016354	bHLH72/bHLH041	Putative transcription factor bHLH041	1	7.206	0	0
Histone	Zm00001d034479	H1a	Histone 1 (H1)	1	26.117	0	0

Figure 1

Figure 2

Figure 3

Gene names	Location	Group	Plaza gene family (HOMO4M)	Zema	Sobi	Seit	Hovu	Trae	Brdi	Orsa	Zoma	Arth	Potr	Vivi	Soly	Metr	Piab	Phpa	Chre
MLKG1	ONM	KASH, grass	009187	1	1	1	1	3	1	1	0	0	0	0	0	0	0	0	0
MLKG2	ONM	KASH, grass	008346	1	1	1	1	6	1	0	0	0	0	0	0	0	0	0	0
MLKS1- MLKS2	ONM	KASH, SINE	002474	2	2	2	1	6	2	2	1	2	2	1	1	1	2	7	0
MLKP1 – MLKP4	ONM	KASH, WIP	002181	4	3	3	2	7	3	3	1	3	2	1	1	2	0	0	0
MLKT1-MLKT2	ONM	WIT	000144	2	1	2	2	5	1	2	2	2	1	2	2	1	0	0	0
ZmSUN1- ZmSUN2	INM	C-term SUN	002694	2	2	2	2	6	2	2	2	2	1	1	1	1	1	2	0
ZmSUN3- ZmSUN5	INM	mid-SUN	001731	3	2	2	2	6	2	2	3	3	3	2	1	3	2	2	1
MNEAP1 – MNEAP3	INM	NEAP	002962	3	2	2	1	2	1	2	2	4	2	2	1	1	1	0	0
NCH1	NUC	NMCP/CRWN	004640	1	1	1	2	3	1	1	4	1	2	2	2	2	2	0	0
NCH2	NUC	NMCP/CRWN	003564	1	1	1	1	3	1	1	4	3	3	1	1	1	2	2	0
MKAKU41- MKAKU42	NUC	KAKU4	004991	2	1	1	1	3	1	1	2	1	2	1	1	1	0	0	0

Column groupings: Viridiplantae (Green plants) spans all species. Land plants (Zema–Phpa). Seed plants (Zema–Piab). Angiosperms (Flowering plants) = Monocots (Zema–Zoma) + Eudicots (Arth–Metr). Poaceae (Grasses) = Zema–Orsa. Gymno = Piab. Bryophyte = Phpa. Algae = Chre.

Figure 4

A

Figure 5

Figure 6