Title:

Identification and characterization of genes encoding the nuclear envelope LINC

complex in the monocot species *Zea mays*.

Running Title:

Maize LINC complex

Authors and Affiliations:

Gumber, Hardeep K.[1]; McKenna, Joseph F.[2]; Estrada, Amado L.[1]; Tolmie, Andrea F.[2]

Graumann, Katja[2]; Bass, Hank W.[1]

1. Department of Biological Science, Florida State University, Tallahassee, FL,
   USA, 32306-4295
2. Department of Biological and Medical Sciences, Faculty of Health and Life
   Sciences, Oxford Brookes University, Oxford, UK, OX30BP.

Corresponding Author:

bass@bio.fsu.edu

Key words:

KASH, LINC, maize, nuclear envelope, SUN

SUMMARY STATEMENT

Genes encoding maize candidates for the core LINC and associated complex proteins have been comprehensively identified with functional validation by one or more assays for several of the KASH genes.

ABSTRACT

The LINC (Linker of Nucleoskeleton to Cytoskeleton) complex is an essential multi-protein structure spanning the nuclear envelope. It connects the cytoplasm to the nucleoplasm, functions to maintain nuclear shape and architecture, and regulates chromosome dynamics during cell division. Knowledge of LINC complex composition and function in the plant kingdom is primarily limited to Arabidopsis, but critically missing from the evolutionarily distant monocots which include grasses, the most important agronomic crops worldwide. To fill this knowledge gap, we identified and characterized 22 maize genes, including a new grass-specific KASH gene family. Using bioinformatic, biochemical, and cell biological approaches, we provide evidence that representative KASH candidates localize to the nuclear periphery and interact with ZmSUN2 *in vivo*. FRAP experiments using domain-deletion constructs verified that this SUN-KASH interaction was dependent on the SUN but not the coiled-coil domain of ZmSUN2. A summary working model is proposed for the entire maize LINC complex encoded by conserved and divergent gene families. These findings expand our knowledge of the plant nuclear envelope in a model grass species, with implications for both basic and applied cellular research.

List of Symbols and Abbreviations:

CDS   coding sequence

CRWN1   CROWDED NUCLEI 1

FRAP   Fluorescence Recovery After Photobleaching

INM   inner nuclear membrane

KASH   Klarsicht/ANC-1/Syne-1 homology

LINC   linker of nucleoskeleton and cytoskeleton

MLKP   Maize LINC KASH AtWIP-like

MLKG   Maize LINC KASH Grass-specific

MLKS   Maize LINC KASH AtSINE-like

MLKT   Maize LINC KASH AtWIT-like

MKAKU4   Maize KAKU4-like

NE   nuclear envelope

NET   nuclear envelope transmembrane protein

NCH   NMCP/CRWN-Homologous

NEAP   nuclear envelope-associated protein

ONM   outer nuclear membrane

SUN   Sad1/UNC-84

TM   transmembrane

WIP   WPP domain-interacting protein

WIT   WPP domain-interacting tail-anchored protein

INTRODUCTION

In eukaryotic cells, the nuclear envelope (NE) is a structural hallmark that encapsulates the biparentally-inherited genetic material, the chromosomes. The NE is comprised of inner and outer nuclear membranes, with embedded nuclear pore complexes (NPC) and a variety of nuclear envelope transmembrane proteins (NETs). Two NETs, Sad1/UNC-84 (SUN), and Klarsicht/ANC-1/Syne-1 homology (KASH) domain proteins, interact in the perinuclear space to form the core of an evolutionarily conserved Linker of Nucleoskeleton and Cytoskeleton (LINC) complex which spans the NE (Crisp et al., 2006). LINC complexes carry out many functions ranging from organizing the shape and position of the nucleus as a mobile and pliable organelle to direct mechanotransduction to effects on chromatin structure and dynamics in somatic and meiotic cells (Kim et al., 2015; Kracklauer et al., 2013; Meier, 2016; Meier et al., 2017; Razafsky and Hodzic, 2015; Starr and Fridolfsson, 2010; Tamura et al., 2015; Tapley and Starr, 2013). The biological importance of NE functions comes primarily from studies of opisthokonts, and in humans is reflected through the wide range of NE-defective developmental disorders such as laminopathies or envelopathies (Burke and Stewart, 2014; Fridkin et al., 2009; Janin et al., 2017; Razafsky and Hodzic, 2015).

The core components of the LINC complex are the inner nuclear membrane (INM) SUN-domain proteins (Hagan and Yanagida, 1995; Malone et al., 1999) and the outer nuclear membrane (ONM) KASH-domain proteins (Starr and Han, 2002). The SUN-domain proteins are known to interact with chromatin, nuclear lamins, or lamin-like proteins (Haque et al., 2006; Hodzic et al., 2004; Zhou et al., 2015b). The KASH-domain proteins are known to interact directly or indirectly with cytoskeletal structures

including microfilaments, microtubules, intermediate filaments, microtubule organizing

centers, and organelles (Luxton and Starr, 2014; Starr and Fridolfsson, 2010). LINC

complexes are highly conserved in all eukaryotes, yet many of the components have

evolved rapidly or recently, limiting the power of sequence conservation to identify

functional homologs.

The body of knowledge of the LINC complex comes primarily from studies in

opisthokonts, especially metazoans and yeast. However, there exists a severe gap in

our knowledge of plant LINC functions, limiting our ability to understand, predict, and

modify plants to meet global challenges in agriculture and biorenewable resources

(Godfray et al., 2010). The first SUN domain protein recognized in plants, OsSad1, was

found as part of nuclear proteomic study in rice (Moriguchi et al., 2005). Its relationship

to the *Saccharomyces pombe* Sad1 was confirmed by localization of OsSad1-GFP

fusion protein to the nuclear periphery in onion epidermal cells (Moriguchi et al., 2005).

Subsequently, two classes of SUN domain proteins were found to be present in plants;

first, the CCSD/C-terminal SUN group, in which the SUN domain is near the C-terminus

and second, the PM3/Mid-SUN group, in which the SUN domain is centrally located

(Graumann and Evans, 2010; Graumann et al., 2014; Murphy et al., 2010; Oda and

Fukuda, 2011). More recently, a large number of SUN domain proteins have been

identified in many plant species by homology searches (reviewed by Meier, 2016; Meier

et al., 2017; Poulet et al., 2017a).

The C-terminal SUN proteins are similar to those in animals, but the mid-SUN

proteins are less well studied as a group despite their known occurrence in plants

(Poulet et al., 2017a), yeast (Friederichs et al., 2011), Dictyostelium (Shimada et al.,

2010) and mice (Sohaskey et al., 2010). Among the conserved roles that SUN proteins

play in plants are those involving the maintenance of the shape and size of the nucleus

(Graumann et al., 2010; Graumann et al., 2014), plant growth (Graumann et al., 2014),

and meiotic chromosome behavior (Murphy et al., 2014; Varas et al., 2015).

The first plant KASH proteins to be described were the WPP-domain-interacting

Other components of the plant LINC complex have been described primarily in a plant

model system, Arabidopsis, a eudicot species that diverged from the economically

important monocot species ~200 MYA (Meier, 2016; Meier et al., 2017; Poulet et al.,

2017a; Tiang et al., 2012; Zeng et al., 2014; Zhou et al., 2015a). proteins (WIPs). The

WIPs are plant-specific, outer nuclear envelope associated proteins originally

discovered as RanGAP-NE anchoring proteins (Xu et al., 2007) and later shown to bind

to SUN and possess KASH-like features (Meier et al., 2010; Zhou et al., 2012). A

related group of ONM LINC-associated plant proteins are the WPP-interacting tail-

anchored (WIT) proteins (Zhao et al,. 2008), which bind directly to WIPs, indirectly to

SUN, and also have WPP-interacting domains. WITs possess the LINC-like activity of

connecting the NE to the actin cytoskeleton through their association with plant specific

Myosin XI-i (Tamura et al., 2013).   In addition to the WIPs and WITs, two other

categories of ONM KASH proteins have been described in plants, the SUN-Interacting

Nuclear Envelope proteins (SINEs) and the TIR-KASH (TIKs), both of which are NETs

(Graumann et al., 2014; Zhou et al., 2014).

There are currently three recognized plant-specific categories of genes encoding

proteins in the INM or at the intranuclear periphery that interact with SUN or each other

to associate with the LINC complex. These encode the INM Nuclear Envelope

Associated Proteins (NEAPs) and nucleoplasmic proteins, including Nuclear Matrix

Constituent Proteins (NMCPs, also called CRWNs) and the AtCRWN-binding

nucleoplasmic protein AtKAKU4s (Ciska et al., 2013; Dittmer and Richards, 2008; Goto

et al., 2014; Masuda et al., 1997; Pawar et al., 2016; Wang et al., 2013). While NEAPs

and CRWNs can be directly associated with LINC complexes by binding to C-terminal

SUN proteins, KAKU4 is indirectly associated with LINC complexes via interaction with

CRWNs (Goto et al., 2014). These interconnected proteins are thought to have roles in

chromatin architecture and nuclear morphology, placing the LINC complex squarely in

genetic and genomic response pathways in plant cells (Poulet et al., 2017b).

There exist profound biological differences and vast evolutionary distances

between monocots (grasses) and eudicots. Given this fact, together with the

importance of the nucleus and the LINC complex in basic developmental processes, we

set out to define and characterize for the first time the entire LINC complex in the crop

grass model species, maize (reviewed by Nannas and Dawe, 2015). We include

comparisons across phyla ranging from single cellular chlorophyte to advanced land

plants, to highlight both conserved and divergent maize LINC complex components.

RESULTS

Identification of maize genes encoding LINC or LINC-associated components

We developed a working model, shown in Figure 1, for the maize LINC complex

and associated proteins as identified through bioinformatic or experimental evidence.

The proteins are depicted as residing in one of the four cellular locations, the cytoplasm,

the ONM of the NE, the INM of the NE, or the nucleoplasm. Despite the considerable

difficulty in assigning certain orthology for all of the maize LINC components, we have

established the following nomenclature for these candidate maize LINC genes. The

genes encoding KASH-like and associated ONM proteins are named with the common

prefix of *MLK* for *Maize LINC KASH* followed by a one letter designation to delineate the

four KASH-related subgroups in maize: *G* for Grass-specific, *S* for *AtSINE*-like, *P* for

AtWIP-like, or *T* for *AtWIT*-like. The genes encoding the INM LINC proteins fall into two

groups, the previously published core LINC SUN domain proteins (Murphy et al., 2010),

and the genes designated *MNEAP* for Maize *AtNEAP*-like. Finally, the genes encoding

the nucleoplasmic LINC-associated proteins are designated *NCH* for NMCP/CRWN

homologs or *MKAKU4* for *Maize KAKU4-like*. In summary, the 22 known or candidate

core LINC and associated proteins in maize are 10 MLKs, 5 SUNs, 3 NEAPs, 2 NCHs,

and 2 MKAKU4s. For all of these, the gene IDs in maizeGDB, PLAZA gene, and

UniPROT are summarized in Table 1, Table S1, and diagrammed with their protein

domain structures in Figure 2. Of these, the KASH and SUN proteins are the most

similar to their animal counterparts, whereas the MLKGs, MLKTs, MNEAPs, MKAKU4s

appear to be more characteristic of plants.

The MLK genes: ten candidates for maize ONM core LINC genes

A hallmark of the KASH domain proteins is that they have limited sequence

conservation and are not reliably recognized by sequence similarity searches. Instead,

they contain a conserved structure consisting of a C-terminal transmembrane domain

and a short periplasmic tail with characteristic terminal residues including a penultimate

proline followed by a threonine or serine. Only a few plant genes with similarity to

animal KASH genes have been identified so far (Zhou et al., 2012; Zhou et al., 2014).

To find all of the potential maize LINC ONM components, we first searched the maize

proteome (B73 AGPv4) from ensembl.gramene.org (Jiao et al., 2017) for KASH proteins

with the following characteristic features: (1) a TM domain at the C-terminus followed

by 9-40 amino acids that could extend into the perinuclear space, and (2) terminal four

residues of [DTVAMPLIFY], [VAPIL], P, and T (Poulet et al., 2017a; Zhou et al., 2014).

This search led to the identification of a total of eight genes encoding maize KASH-like

candidates, four with homology to AtWIPs (*MLKP1 – MLKP4*), two with homology to

AtSINEs (*MLKS1* and *MLKS2*) and two found only in members of the grass family,

Poaceae (*MLKG1* and *MLKG2*).   We next searched the maize proteome for WIT

homologs using AtWIT1 and AtWIT2 protein sequences as queries (Zhao et al., 2008)

and found two maize genes, *MLKT1* and *MLKT2*, both encoding proteins with the

AtWIT-like TM domains at their C-termini and multiple coiled-coil (CC) domains. Given

the rapid divergence of LINC gene sequences and the lack of distinct and diagnostic

domains, and gene content differences between maize cultivars (Hirsch et al., 2016), it

is possible that maize has even more MLKT genes not readily detected using these

criteria.

The four MLKP (AtWIP-like) genes belong to the PLAZA gene family designated

HOMO4M002181, and encode proteins with multiple coiled-coil domains (MLKP1-

MLKP4, Fig. 2). The previously identified AtWIP proteins interact with AtWITs and

RanGAP1 and exhibit coiled-coil domain-dependent homo- and hetero-multimerization

(Zhao et al., 2008; Zhou et al., 2012). The MLKP proteins are therefore predicted to

interact with themselves, each other, ZmSUN proteins, and RanGAP proteins with WPP

domains. The two MLKS (AtSINE-like) genes belong to PLAZA gene family designated

HOMO4M002474, and encode proteins with armadillo (ARM) repeats (MLKS1- MLKS2,

Fig. 2). ARM domains have been reported in a variety of proteins across plant and

animal kingdoms as actin-binding and protein-protein interaction domain (Coates,

2003). The MLKS proteins are predicted, therefore, to interact and co-localize with

actin, as has been demonstrated for AtSINE1 (Zhou et al., 2014). The two MLKG

(grass-specific) genes lack both CC and armadillo domains and have been assigned to

different PLAZA gene families - HOMO4M009187 for *MLKG1* and HOMO4M008346 for

MLKG2.

Beyond the eight candidate MLK genes that we predict to encode maize ONM

core LINC complex proteins, we also found four other maize genes that met some of

our criteria. For instance, some encoded proteins with the terminal PT residues, but did

not satisfy the single C-terminal TM domain criterion. It remains unknown whether

these genes (B73v4 gene IDs Zm00001d024832, Zm00001d036395, Zm00001d005590

and Zm00001d017144) encode actual variant LINC proteins or simply resemble LINC

proteins.

The eight maize INM LINC genes; five SUN and three NEAP genes

The five genes encoding maize SUN-domain proteins belong to two subfamilies

distinguished by the location of the SUN domain (Murphy et al., 2010). They are not

newly identified in this study, but are included in Table 1 for the sake of completion,

along with their most recent B73v4 gene IDs. ZmSUN1 and ZmSUN2 proteins have the

canonical C-terminal SUN domain, whereas the ZmSUN3, ZmSUN4,and ZmSUN5 have

a central SUN domain that is characteristic of, but not limited to plants (Graumann et al.,

2014). Despite the conservation of SUN-domain proteins as a group, and their

assignment to one of two subfamilies, direct assignment of orthology for individual SUN

genes across taxonomic groups such as eudicots (Arabidopsis) versus monocots

(maize) has yet to be established.

Another major family of INM LINC proteins are the higher plant-specific proteins

called Nuclear Envelope-Associated Proteins, the NEAPs (reviewed by Groves et al.,

2018; Meier et al., 2017; Poulet et al., 2017a). NEAPs exist as a multigene family, first

discovered in Arabidopsis and shown to interact with SUN-domain proteins (Pawar et

al., 2016). Genetic analysis of AtNEAPs showed that they affect several phenotypes,

including primary root growth, nuclear morphology, and chromatin organization (Pawar

et al., 2016) . NEAPs have three protein structural features, the presence of extensive

coiled coil domains, nuclear localization sequences, and C-terminal TM domains. We

used these criteria to identify three maize NEAP genes (Table 1, Fig. 2), *MNEAP1*,

MNEAP2, and *MNEAP3*. Interestingly, we observed partial homology to N-terminus of

structural maintenance of chromosomes domain (SMC_N, PFAMID PF02463) proteins

in the MNEAPs, further implicating them in DNA and chromosomal functions.

The MNEAP1 and MNEAP2 proteins both end with a characteristic C-terminal

hydrophobic patch and two basic terminal residues KR (Pawar et al., 2016). These two

proteins share 92% sequence identity and are very similar to the protein encoded by

one of the two sorghum NEAPs genes (sobic.009G033800). The *MNEAP1* and

MNEAP2 most likely arose from the genome duplication event that occurred following

the divergence of maize from sorghum, ~ 11 MYA (Gaut et al., 2000). *MNEAP3* fits our

initial criteria but is otherwise quite different in many respects including an unusually

large number of splice variant transcript models and the lack of KR terminal residues.

Consequently, *MNEAP3* may be a complex locus or alternatively, pseudogene no

longer under evolutionary pressure to maintain its structure.

The four maize nucleoplasmic LINC-associated genes; two *NCH* and two *KAKU4*

genes

In mammals, the SUN-domain proteins of the LINC complex interact with the

nuclear lamina, which is located at the periphery of the nucleoplasm and made up of

lamin proteins. Plant nuclei lack laminA/B orthologs, but they do have a lamin-like

meshwork that is thought to consist of lamin-like proteins known as NMCP/CRWN as

well as CRWN-binding nuclear proteins, KAKU4, (Ciska et al., 2013; Goto et al., 2014).

These proteins collectively are thought to comprise a significant portion of the plant

nucleoskeleton and they are coupled to the LINC complex via interactions with SUN

domain proteins (Goto et al., 2014).

Using AtCRWN and DcNMCP protein sequences as queries , we identified two

maize genes, *NCH1* and *NCH2*, representing the two major clades of NMCP/CRWN

genes (Ciska et al., 2013; Wang et al., 2013). The maize *NHC1* is most similar to

DcNMCP1, *DcNMCP3*, and *AtCRWN1*. The genes in this group are members of

PLAZA gene family HOM04M004640 (Table S1). The maize *NCH2* gene is most

similar to *DcNMCP2*, *AtCRWN2*, *AtCRWN3*, and *AtCRWN4*. The genes in this group

are members of the PLAZA gene family HOM04M003564. Both NCH1 and NCH2

proteins share several features with their plant homologs. These features include large

proteins (~100 kDa or greater), multiple coiled-coil domains over the majority of the

protein, multiple CDK1 phosphorylation sites, and a nuclear localization sequence near

the C-terminus (Fig. 2).

Using AtKAKU4 protein sequence similarity searches, we identified two highly

similar maize KAKU4-like genes, *MKAKU41* and *MKAKU42*. These two genes and

AtKAKU4 are all members of the PLAZA gene family HOM04M004991. KAKU4 and

KAKU4-like genes appear to be limited to flowering plants and encode proteins that

have been described as candidate lamina components (Goto et al., 2014). As a group,

they may be remotely associated with the LINC complexes but are included here

because of the known interaction of AtKAKU4 with AtCRWN proteins as well as the

genetic evidence that implicates them in LINC functions and nuclear architecture (Goto

et al., 2014). Aside from a NLS near the N-terminus, the MKAKU4 proteins do not

contain any notable structural protein domains or motifs (Fig. 2).

A mix of ancient and derived genes encode components of the maize LINC

complex

Given that the time of appearance and the evolutionary conservation of genes

often reflects functional properties, we examined the maize LINC candidate genes and

their counterparts in a variety of plant species as summarized in Figure 3. Recent

phylogenetic analyses of plant LINC genes or sub-families therein have been published

(Ciska et al., 2013; Poulet et al., 2017a; Zhou et al., 2014), with emphasis on

Arabidopsis and other eudicot species, but did not include the latest PLAZA gene

families or maize B73v4 annotations (Jiao et al., 2017; Van Bel et al., 2018). The

species listed (Fig. 3) are arranged in increasing evolutionary distance from maize

starting with sorghum and extending to the unicellular green alga, *Chlamydomonas*

reinhardtii.

In plants, the most highly conserved members of the core LINC and associated

complex belong to gene families that encode the C-terminal and mid-SUNs, NCH2 and

the AtSINE-like KASH proteins MLKS1 and MLKS2). These gene families are present

in all the plant species listed (shaded blue or green, Fig. 3). Curiously, of all the genes

listed, only the mid-SUN genes are listed as present in Chlamydomonas PLAZA gene

families. The next most highly conserved plant LINC gene families encode the MNEAP

and NCH1 proteins. They are present in seed-bearing plants (shaded tan, Fig. 3),

which includes pine, a conifer species. The third most highly conserved plant LINC

gene families encode MLKP, MLKT and MKAKU4 proteins, all of which are found in

flowering plants, the angiosperms (shaded pink, Fig. 3). In contrast, the most recent to

appear in evolutionary time are the gene families that encode the MLKG proteins,

present in the grass family within the monocots (shaded yellow, Fig. 3). The biological

significance of the timing of evolutionary appearance remains to be determined, but we

generally expect that the more widely distributed gene families have more ancient or

basal functions, whereas the more recent gene families could have resulted from

duplication followed by either regulatory divergence, such as tissue-specific expression,

or gain of new biological functions.

Maize LINC genes exhibit tissue-specific and developmental co-expression

patterns

We analyzed the expression patterns of the maize LINC candidate genes as

shown in Figure 4 in order to examine them for evidence of expression and potential

functional groupings that should be reflected in their differential or co-expression

patterns. For this, we used the expanded maize gene expression atlas which contains

transcriptome data from numerous tissues broadly sampled across most of the

developmental stages of the maize life cycle (Stelpflug et al., 2016). For comparison,

we included three housekeeping genes (HKG, Fig. 4A), *CDK1*, *RPN*, and *SGT1* (Lin et

al., 2014).   We excluded *MNEAP3*, which lacked a gene model when the atlas was

produced, and *MLKG1*, which had zero reads in all fifteen tissues selected for this

study. Despite their exclusion, these two are presumed to be real genes and not

pseudogenes because of the expression of their corresponding transcripts detected in

one or more tissues (Chettoor et al., 2014; Walley et al., 2016; Wang et al., 2016; Xu et

al., 2012)*.

Most of the candidate maize core LINC and LINC-associated genes showed a

pattern of broad expression across multiple tissues, with co-expression patterns

(dendrogram, Fig. 4A) revealing several distinct groupings.   The "root" co-expression

group (orange box, Fig. 4A) includes three root samples and 12 DAP developing seed.

The "young inflorescence" co-expression group (blue box, Fig. 4A) includes male

(immature tassel) and female (immature cob) tissues. The "meristem" co-expression

group (green box, Fig. 4A) includes the developing embryo and growing shoot tip

tissues. These last two groups together form a larger co-expression cluster from

tissues that share the property of having actively dividing cell populations. Beyond

these three groups, several tissues show distinctive patterns of LINC gene expression,

not clustering with other tissues, including anthers (male pollen-bearing organs),

topmost seedling leaf, meiotic tassel, and silks (female, pollen-receiving organs).

To characterize the general expression levels of the candidate maize LINC

genes, we plotted the median FPKM values across different tissues (Fig. 4B). Maize

LINC genes generally exhibit low to moderate expression levels, similar to those of the

house-keeping *CDK1* gene (Fig. 4B). The more highly expressed LINC-related genes

had transcript levels that ranged from 80-140 median FPKM; these genes included the

MKAKU4s, the C-terminal SUNs, the NCHs, and some of the KASH genes, *MLKP3* and

MLKP4. The less-expressed genes had transcript levels that ranged from 3-50 median

FPKM; these included the mid-SUNs, the MNEAPs, and most of the other KASH genes.

As a group, the maize AtWIP-like genes (*MLKP1-4*) have higher expression levels than

the AtSINE-like genes (*MLKS1*, *MLKS2*), with *MLKP3* and *MLKP4* showing the most

uniform and broad expression. In contrast, the mid-SUN *SUN5* gene exhibited a most

striking pattern of expression in only pollen. The *Poaceae*-specific gene, *MLKG2,*

appears to be expressed in tissues characterized by meristem and active cell division,

but absent from others such as endosperm, root, leaf, and silks.

Among the highly similar gene pairs that are likely to have arisen as part of a

recent genome duplication (Gaut et al., 2000), some of them (*MLKP1 and MKLP2;*

MLKP3 and MLKP4; and MKAKU41 and MKAKU42) show similar expression levels

whereas other gene pairs, such as *MNEAP1* and *MNEAP2,* exhibit differential

expression levels. Overall, this expression analysis provides evidence that genes

encoding LINC and associated proteins are transcribed and regulated in a tissue-

specific manner. The co-expression groupings predict possible but untested genetic or

physical interactions that may reflect shared functions.

Identification of LINC and associated proteins from endogenous maize co-IP

using anti-ZmSUN2 antibodies.

In order to find naturally occuring interactors of maize SUN proteins, we carried

out co-immunoprecipitation (co-IP) optimized for NE proteins from purified earshoot

nuclei as summarized Table 2. Maize earshoots of ~3-5 cm in length yield high quality

cellular extracts from fast-growing somatic tissues several days prior to silking and

fertilization. The antibodies used for co-IP were the affinity-purified serum designated

anti-SUN2-NPAP as previously characterized (Murphy et al., 2014). Control co-IP

experiments were conducted using either no antibodies or irrelevant but similarly

affinity-purified rabbit sera, anti-ZmNDPK1 (Kopylov et al., 2015).   A key technical

advance was the use of the short-chain crosslinker dithiobis(succinimidyl propionate),

DSP. We adapted the membrane protein cross-link immunoprecipitation (MCLIP)

method for use with maize because it was shown to effectively recover SUN-interacting

proteins from U2OS cells (Jafferali et al., 2014). We found that the MCLIP method,

using DSP together with light formaldehyde fixation to preserve the nuclei, resulted in

substantially increased final recovery and detection of known or predicted maize

homologs of plant LINC, NE, or SUN-interacting proteins (Tamura et al., 2013; Wang et

al., 2013; Zhou et al., 2012).

The proteins recovered from the co-IP assays were subjected to tryptic digestion

and identified by liquid chromatography tandem mass spectrometry (LC-MS/MS).   Over

140 proteins were identified (Table S3) using software thresholds of 95% protein and

95% peptide. As expected, SUN2 was the most abundant protein, but we also

repeatedly detected SUN1, which could result from cross-reactivity of ZmSUN1 with our

anti-SUN2 serum, or from actual co-IP of ZmSUN1 bound to SUN2, as expected from

documented hetero multimerization of SUN proteins in plants and animals (Sosa et al.,

2012).

The identified co-precipitated proteins were considered high confidence if they

were detected in multiple replicates but not in the negative controls. The specificity of

the modified MCLIP method is evidenced by recovery of core LINC components and

predicted cytoplasmic and nucleoplasmic proteins. Among the predicted core LINC

components repeatedly found in the top co-IP hits were three ONM proteins (MLKG1,

MLKS2, and MLKT2) two INM proteins (ZmSUN1 and ZmSUN2), two NMCP/CRWN

homologs (NCH1 and NCH2), two RAN family proteins (RAN GTPase and RAN-GAP2)

and five nuclear pore proteins (NUP133, NUP96, NUP88, GP210, and NUA).

We also found proteins that were co-precipitated but likely via indirect interaction

with SUN. These include the cytoskeletal proteins TUB6, TUB4 and Actin-1/ARP7. The

Actin-1/ARP7 protein could be cytoplasmic or nucleoplasmic or both because its

annotation includes the designation of nuclear actin, an intriguing possibility in light of

recent findings showing cross-species binding of Arabidopsis AtARP7 to the carrot

nuclear matrix protein DcNMCP1 (Mochizuki et al., 2017). We recovered several

nuclear proteins with DNA-binding activities including transcription factors bZIP, ARF10,

and the linker histone H1a.   Overall, the co-IP resulted in confirmatory identification of

naturally occuring SUN2-interacting proteins with known or suspected LINC proteins,

plus several additional proteins that represent new targets for future investigation. To

further verify some of the LINC candidates from either the Co-IP or bioinformatic

analyses, we selected three genes for further localization and interaction assays *in vivo*

using transient heterologous expression assays.

***In planta* evidence of NE targeting and ZmSUN2 interaction for three candidate**

maize LINC proteins

In order to investigate the subcellular localization of the maize KASH proteins, we

transiently expressed N-terminal GFP fusions of MLKG1, MLKP1 or MLKP2 in *Nicotiana*

benthamiana leaf tissue as summarized in Figure 5. Expression of a positive control,

mCherry-ZmSUN2, shows that maize NE proteins can be properly expressed from the

plasmids and localized to the NE in this leaf cell heterologous expression system. All

three of the previously untested ONM maize LINC candidate proteins, MLKG1, MLKP1

and MLKP2, also localized to the nuclear periphery (Fig. 5A). These localization

patterns are comparable to those previously reported for AtKASH proteins in similarly

designed expression assays (Zhou et al., 2012; Zhou et al., 2014).   For GFP-MLKG1, a

network-like pattern at the cell periphery was also seen (Fig. 5B; Fig. S2C). In order to

further examine this pattern, we co-expressed GFP-MLKG1 with an ER marker and an

actin marker and found that GFP-MLKG1 co-localizes with both the ER and the actin

network (Fig. 5B). Similar co-localization patterns have previously been demonstrated

for AtSINE1 (Zhou et al., 2014) but, unlike AtSINE1, MLKG1 does not possess an ARM

domain or other known actin binding motifs.

The C-termini of the KASH proteins are known to play a key role in its NE localization

which is mediated through interaction with SUN proteins (Sosa et al., 2012; Zhou et al.,

2012; Zhou et al., 2014). To test for a role of the C-terminal residues in our maize

KASH candidates, we created and tested several classes of C-terminal deletion

constructs (Fig. 5C). The Δ4 deletion constructs are lacking the last 4 C-terminal amino

acids. The ΔKASH deletion constructs are lacking all of the amino acid residues

following the TM domain, made with (MLKP1/2 ΔKASH) or without (MLKG1ΔKASH) a

minor three alanine stabilizing spacer. The ΔTM deletion constructs are lacking the

entire C-terminus starting from the beginning of the predicted TM domain (Fig. 5C,

S1,S2).   For MLKP1, both the ΔKASH or ΔTM protein truncations changed the

localization pattern to that resembling soluble proteins (Fig. 5D, S2) whereas the Δ4

truncation showed a small increase in apparent solubility, but retained considerable

targeting to the NE. For MLKP2Δ4, MLKP2ΔKASH, and MKLG1ΔKASH, the terminal

truncations still resembled that of the full length proteins with localization to the NE. For

comparison, we expressed GFP-only as a control (Fig. S2B), which illustrates the

distribution of a soluble protein without a TM domain in the cytoplasm and nucleoplasm.

Interestingly, the removal of the KASH domains from either MLKP2 or MLKG1 did not

abolish its ability to localize to the nuclear periphery, suggestive of other or additional

determinants that enable MLKP2 and MLKG1 associations with the NE membranes

(e.g. Fig. 5B, S2C).

In order to investigate whether maize KASH binds to maize SUN *in vivo*, we co-

expressed three different maize GFP-KASH with or without ZmSUN2. The GFP-KASH

proteins co-localized to the NE with mCherry-ZmSUN2 full length (Fig. S7). If GFP-

KASH and ZmSUN2 do interact, we would expect to see a reduction in the mobility of

GFP-KASH as measured by fluorescence recovery after photobleaching and indicative

of interacting partners, as previously described (Zhou et al., 2014). For MLKG1 and

MLKP2, we did indeed observe reduced mobility when co-expressed with ZmSUN2

(Fig. 6B,F; S4-5). When GFP-MLKG1 is transiently expressed by itself, 63±8% of the

protein population is mobile while 36% remains immobilized. However, upon co-

expression with mCherry-ZmSUN2, only 49±12% of the protein population was mobile,

indicating that the co-expressed ZmSUN2 substantially increased the immobile fraction

of GFP-MLKG1 from 36% to 51% (Fig. 6A,B). We interpret this shift as indirect

evidence for ZmSUN2 interaction with GFP-MLKG1. To test this idea, GFP-MLKG1

was co-expressed with a domain deletion mutant of ZmSUN2 lacking the SUN domain

(mCherry-ZmSUN2ΔSUN); we found that the mobility of MLKG1 was similar to that of

the MLKG1 alone (68±7%; Fig. 6 B,C). This finding indicates that the SUN domain is

required for mediating the change in GFP-MLKG1 mobility, providing strong indirect

evidence that ZmSUN2 interacts with MLKG1. To further test if the coiled-coil domain of

the SUN protein is required for reduced mobility, we created a version of mCherry-

ZmSUN2 lacking the coiled-coil domain, mCherry-ZmSUN2ΔCC. Co-expression of this

mCherry-ZmSUN2ΔCC with GFP-MLKG1 had a similar effect on KASH mobility to that

of the full length SUN2 (Fig. 6 B, C; S4-5). This result establishes that the ZmSUN2-co-

expression-dependent increase in the immobile fraction of MLKG1 occurs to a similar

degree when co-expressed with ZmSUN2 with or without the CC domain.

Similar changes in protein mobility were observed for GFP-MLKP2 (Figs 6 F, G

and S4-5) when co-expressed with full-length ZmSUN2, ZmSUN2ΔSUN or

ZmSUN2ΔCC. Co-expression with ZmSUN2 or ZmSUNΔCC resulted in significant

increases in the immobile fraction of GFP-KASH from 51% to 69% or 71%, respectively.

In contrast, we did not observe statistically significant changes in GFP-MLKP1 mobility

when co-expressed with full-length or deletion constructs of ZmSUN2 (Figs 6D, E and

S4), even though the changes observed were all in the same directions as those for

GFP-MLKG1 and GFP-MLKP2 (Figs 6, S3-5).

Of the 3 candidates tested, MLKG1 was already found by endogenous maize

co-IP to interact with SUN2, but the other two, MLKP1 and MLKP2 were not,

presumably because of their low expression levels in earshoot (Fig. 4). In order to

check for their ability to interact with SUN2 *in planta*, we assayed them by co-IP in

tobacco tissues where they were co-expressed as heterologous fluorescent fusion

proteins (Fig. S6A). Using anti-GFP antisera, we observed co-precipitation of mCherry-

SUN2, but not in control extracts from non-infiltrated or mCherry-SUN2 only samples.

These additional tests confirmed that all three of the proteins tested, MLKG1, MLKP1,

and MLKP2, showed SUN2 interaction in one or more direct or indirect interaction

assays (Table in Supplemental Figure 6B).

In summary, all three of the KASH candidates tested showed NE localization in

the transient heterologous expression system. They also showed evidence of in planta

interaction with ZmSUN2, supported by one or more lines of evidence, co-expression

FRAP, endogenous co-IP, or heterologous co-IP. These findings, together with our

bioinformatic, transcriptomic, and phylogenetic analyses provide the first experimental

evidence for an entire core LINC complex and associated proteins in a model monocot

species.

DISCUSSION

The human population growth rate imposes a great challenge in agriculture and

food production. It has been estimated that in order to provide food for a projected

population of 9.15 billion in 2050, agricultural productivity must increase by 60% (from

2013 Summit Report at https://plantsummit.wordpress.com/). For both biological and

social reasons, it is necessary to discover and characterize new plant genes that inform

our knowledge how the cell nucleus impacts organismal traits. This is especially true for

model crop species such as maize with annual production at over 400 million bushels

per year (FAOSTAT 2016, http://www.fao.org/faostat/en/). This study, the first to define

the composition of the maize LINC complex and associated proteins, represents a

major advance that leverages knowledge from non-plant eukaryotes regarding the role

of the LINC complex in gene regulation, chromatin dynamics, and nuclear architecture

(reviewed by Alam et al., 2016; Hieda, 2017; Kim et al., 2015; Poulet et al., 2017b).

Here we set out to systematically identify all of the maize LINC core and

associated components, adding a valuable new and different model species for

investigation of the plant nuclear envelope and important biological processes such as

meiotic chromosome behavior, development, signaling, and pathology (Kracklauer et

al., 2013; Meier et al., 2017; Razafsky and Hodzic, 2015; Tapley and Starr, 2013).

Maize provides a valuable plant complement to Arabidopsis. Both maize, a monocot,

and Arabidopsis, a eudicot, have made important contributions to biology, yet they have

quite different body plans, seed development pathways, genome sizes, and gene

expression profiles, having diverged 200 MYA (Brendel et al., 2002; Conklin et al.,

2018; Dukowic-Schulze et al., 2014; Wang et al., 2012; Zeng et al., 2014).

In Arabidopsis, SUN1 and SUN2 double knockouts are known to disrupt

conserved functions such as meiotic progression, excessive chromosomal interlocks,

and reduced fertility (Varas et al., 2015). However, Arabidopsis does not exhibit a

canonical early prophase telomere bouquet cluster on the nuclear envelope, nor has it

been shown to have a meiotic SUN belt (Armstrong et al., 2001; Hurel et al., 2018;

Murphy et al., 2014). These differences may highlight the limitations of extrapolating

across divergent species within the plant kingdom. Furthermore, the difficulty in

identifying certain orthologs for some LINC components illustrates the importance of

defining the composition of the LINC complex in multiple species, especially in species

with well-developed genetic and cytological resources that enable comparative analysis

of the associated biological functions.

Searching for maize KASH and associated ONM proteins, we found homologs of

plant genes encoding AtWIP and AtSINE1/2-like proteins, but not those encoding

AtSINE3/4, SINE5 or AtTIK (Graumann et al., 2014; Zhou et al., 2014). Curiously, the

AtSINE3/4 genes are found only in brassicaceae, the *SINE5* genes are found only in

medicago, and the *AtTIK* genes are found only in Arabidopsis (Poulet et al., 2017a).

Similarly, *MLKG1* and *MLKG2* genes are found only in maize or other grasses and

appear to be completely absent from eudicot species such as Arabidopsis or Medicago.

In contrast, the *MLKT* genes of maize are clearly homologous to the *AtWITs* genes, a

rare case in which sequence conservation is sufficient to identify candidate ONM LINC

components. Notably, additional maize genes resembling ONM LINC components were

found and may have NE functions, but fell short of the criteria we applied in this study.

It remains an intriguing but recurring observation that ONM LINC proteins are part of an

ancient structure, but they can be highly divergent. In this regard, plant KASH genes are

similar to those from opisthokonts, with characteristic variation in protein sequence and

structure (Meier et al., 2017).

Searching for additional maize genes encoding INM proteins, we found three

homologs of the plant-specific *NEAP gene*s. The *NEAP* genes are limited to seed

plants (Poulet et al., 2017a) and not found in any of the representatives of the more

ancient moss and algal species included in this study (Fig. 3). The *MNEAP1* and

MNEAP2 genes encode protein sequences that end in two basic residues, like their

eudicot homologs, but the functional significance of these terminal residues remains

unknown. Interestingly, the maize NEAPs have SMC_N domains which are currently

uncharacterized but may point to a chromosome tethering or nuclear structural role.

The maize NEAPs named in this study are three of the eleven maize genes assigned to

the PLAZA gene family HOM04M002962. Consequently, the actual number of the

maize NEAP genes may grow as revised bioinformatic and functional criteria for their

recognition emerges. In contrast, we did not find evidence for additional SUN genes of

maize beyond the five previously described (Murphy et al., 2010; Murphy et al., 2014).

Searching for nucleoplasmic proteins associated with plant LINC complexes, we

found two maize genes with homology to *NMCP*/*CRWN* genes, *NCH1* and *NCH2*, both

of which are deeply conserved in plants. Each of these are associated with multiple

transcript models (Wang et al., 2016) suggestive of splice variants, an intriguing

observation in light of the tendency for animal lamin genes to also have many splice

variants (Broers et al., 2006). Identification of plant nuclear matrix components that

interact with the nuclear envelope provides a more complete view of the physical bridge

across the NE. As such, this emerging model provides new opportunities to investigate

the mechanisms that underlie how signals are transduced via the LINC complex.

Cellular localization experiments confirmed the NE localization of fluorescent

fusion protein constructs for ZmSUN2 and all three of the maize KASH homologues

tested, MLKG1, MLKP1 and MLKP2 (Fig. 5). Interestingly, GFP-MLKG1 co-localizes

with the ER and the actin network, implicating it in cytoskeletal signal transduction to the

nucleus. Relatedly, Arabidopsis actin is tethered to the LINC complex through direct

interactions with AtSINE1 and through the interaction of myosin XI-i with WIT-WIP

(Tamura et al., 2013; Xu et al., 2007; Zhou et al., 2014). The presence of *SINE1*, *WIP*

and *WIT* homologs in maize, together with the poaceae-specific *MLKG1,* suggests that

plant genomes may encode a diverse repertoire of ONM proteins to connect and

integrate NE with cytoskeletal processes.

To provide additional experimental evidence on the composition of the maize

LINC complex, we used a ZmSUN2 co-IP method optimized for membrane proteins

together with isolated nuclei. We found a large number of different proteins that were

generally consistent with predictions from our working model (Fig. 1) and previous

reports (Dittmer and Richards, 2008; Goto et al., 2014; Pawar et al., 2016; Tamura et

al., 2010; Tamura et al., 2013; Wang et al., 2013; Zhou et al., 2012; Zhou et al., 2015b;

Zhou et al., 2015c). Among the expected and reported LINC and LINC-associated

proteins, our endogenous co-IP detected SUN, WIPs, WIT, NCH, and RAN cycle

proteins.

In relation to the cytoplasmic and ONM side of the LINC complex, we detected

proteins predicted to bind directly to ZmSUN2 including MLKG1 and MLKS2. We also

co-precipitated proteins predicted to bind indirectly to ZmSUN2 including MLKT2, two

RAN cycle proteins, RAN GTPase, RAN-GAP2, Tub4, Tub6, and the nuclear pore

proteins NUP133, NUP96, NUP88, GP210, and NUA. Notably, we did not detect all of

the nuclear pore proteins, consistent with previous studies showing interactions of a

specific and limited subset of NPC proteins with SUN-domain proteins in association

with transcriptional activation and mRNA export processes (Gerber et al., 2004; Li and

Noegel, 2015). In relation to the nucleoplasmic and INM side of the LINC complex, we

detected two INM proteins, ZmSUN1 and ZmSUN2; and two NMCP/CRWN proteins,

NCH1 and NCH2.

The co-IP experiments yielded an interesting list of additional proteins beyond

the core LINC and associated proteins summarized in our working model. Some of

these proteins appear to belong to non-specific protein categories but in fact have been

found in various NE studies involving nuclear actin (ARP7/Actin-1), transcription factors

(bZIP5, ARF10, MYBR35, bHLH72/bHLH041), Histone H1 (H1a), and ribosomal

proteins (L19, L6, and L22-2) (Horigome et al., 2011; Hotta et al., 2007; Mochizuki et al.,

2017; Nakayama et al., 2008). Nuclear actin is increasingly recognized as playing key

roles in chromosome dynamics and gene expression (reviewed by Spichal and Fabre,

2017) whereas a plant linker H1 histone was recently shown to have MTOC activity and

reside at the nuclear periphery (Boruc et al., 2012; Nakayama et al., 2008).

Given the increasing recognition that the NE and nucleoskeletal proteins are

implicated in chromatin remodeling and gene regulation, these additional non-LINC

proteins discovered through co-IP represent ideal experimental targets for investigating

regulatory mechanisms of chromatin structure and nuclear architecture. For instance,

the NE cooperates with nucleoskeletal proteins to modulate euchromatin and

heterochromatin organization (Poulet et al., 2017b; Wang et al., 2013). Emerging

breakthroughs in HiC mapping of nuclear architecture are likely to uncover additional

principles of genome organization that involve LINC-binding nuclear proteins (Grob et

al., 2014). Given the importance of lamins in human health, it is possible that plant

NMCP/CRWN/NCH defects may shed light on NE-associated diseases.

Both co-IP and FRAP provide complementary assays for investigating *in planta*

binding of maize KASH candidates to ZmSUN2 (Fig. 6). For example, MLKG1 was co-

purified from maize extracts using SUN2 antisera (Table 2) while FRAP assays

demonstrated a ZmSUN2-dependent change in the mobility of GFP-MLKG1 and GFP-

MLKP2. The heterologous expression system described in this and previous studies

(Zhou et al., 2012; Zhou et al., 2014) represents an efficient means to detect, isolate,

and dissect protein-protein interactions for distantly related grass species such as

maize, which is not easily transformed. It also provides an efficient means to carry out

in vivo tests of candidate interactors (e.g. Fig. S6) for which biochemical amounts of

native tissue may be limiting.

Overall, this study provides the first comprehensive list of maize LINC candidates

(Fig. 1), several of which were validated with multiple lines of evidence. Through the

identification of an ensemble of LINC components in a model eukaryote, we have

established valuable new avenues of research to address unanswered questions

regarding the role of the nuclear envelope as an integrative structure coordinating

nuclear and cellular functions.

METHODS

Bioinformatic identification of MLKs

The maize B73 proteome v4

(ftp://ftp.gramene.org/pub/gramene/CURRENT_RELEASE/fasta/zea_mays/pep/) was

searched for sequences ending in motif [DTVAMPLIFY][VAPIL]PT (brackets indicate

alternative amino acid residues at the respective position). The selected proteins were

further searched for the presence of TM domain using TMHMM (Krogh et al., 2001) and

Phobius software (Käll et al., 2004). Proteins lacking TM domain or specific ending were

discarded. The selected maize MLKs were then searched on PLAZA monocot v4

(https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v4_monocots/) and those

belonging to same homolog family were grouped together.

Protein feature calculations and domain searches

Theoretical molecular weight and isoelectric point (pI) of candidate proteins was

calculated using the pI/Mw tool of ExPASy (https://web.expasy.org/compute_pi/). Coiled

coil domains were identified by Jpred (http://www.compbio.dundee.ac.uk/jpred/). For the

identification of conserved domain sequences, the candidate protein sequences were

searched on Uniprot (http://www.uniprot.org/), NCBI Conserved Domain Database

(https://www.ncbi.nlm.nih.gov/cdd/), PFAM (https://pfam.xfam.org/), and SMART

(http://smart.embl-heidelberg.de/). The domains identified were then plotted using IBS

1.0.3 (Illustrator for Biological sequences) software (Liu et al., 2015).

Identification of LINC orthologs in other organisms and phylogenetic

reconstruction

Sequences of orthologs of Maize LINC candidates belonging to matching HOMO gene

families were retrieved from PLAZA v4

(https://bioinformatics.psb.ugent.be/plaza/versions/plaza_v4_monocots/). Sequences of

all species selected for this study are present in PLAZA except *Medicago truncatula*,

which were retrieved from Gramene. The protein sequences were further screened for

the presence of characteristic protein features. The MLKS/MLKG/MLKP orthologs were

searched for the presence of TM domains towards C-terminus and for terminal four

residues of the proteins to be VIPT, VVPT, AVPT, PLPT, TVPT, LVPT or PPPS. For

MLKT orthologs, proteins having high sequence similarity and C-terminal TM domains

were retained. For NCHs, KAKUs and NEAPs, sequences having high sequence

similarity to Arabidopsis CRWNS, KAKU4 and NEAPs were selected. For phylogenetic

reconstruction of each protein category, sequences were first aligned using MUSCLE

and Neighbour end joining trees were generated in MEGA 7.0.21 (Kumar et al., 2016)

using default parameters.

Transcriptome

685   The expanded maize gene expression atlas (Stelpflug et al., 2016) was

686 downloaded for analysis. Log$_{10}$ Fragments Per Kilobase of exon per Million fragments

687 mapped (FPKM) values for 15 tissues were used to construct a heatmap using the

688 Heatmap2 package of R. To group tissues by similar expression profiles, hierarchical

689 clustering was done using the Hclust function of R.

Co-IP protocol

Maize (Zea mays L) B73 seeds were obtained from WF Thompson (NC State

University, Raleigh, NC, USA, accession "27 FARMS", bulked for NCSU, denoted

B73NC27). Seeds were grown at the Florida State University Mission Road Research

Facility (Tallahassee, FL). Young earshoots (3-5 cm long) were harvested from 7-8-

week-old plants, flash frozen in liquid nitrogen and stored at -80 °C until use. Ten

grams tissue was ground under liquid nitrogen with a mortar and pestle, and cross-

linked with 0.1% formaldehyde by stirring for 10 min in 100 mL ice-cold Buffer A (15

mM PIPES-NaOH, pH 6.8, 0.32 mM sorbitol, 80 mM KCl, 20 mM NaCl, 0.5 mM EGTA,

2 mM EDTA, 1 mM DTT, 0.15 mM spermine, and 0.5 mM spermidine; (Belmont et al.,

1987). The nuclei were isolated as described in (Vera et al., 2014) and resuspended in

5 ml PBS. The isolated nuclei were treated with 2 mM DSP (dithiobis(succinimidyl

propionate) (Cat. No. 22585, Thermo Scientific) crosslinker for 2 h at 4 °C. To lyse the

nuclei, 7 M urea, and 1% Triton X-114 and plant protease inhibitor cocktail (Cat. No.

P9599, Sigma) were added and incubated for 20 min on ice. The ruptured nuclei were

sonicated three times in Bioruptor (Diagenode) for 5 minutes each at medium setting.

Urea was diluted with 8 volumes of PBS. Homogenates were centrifuged at 10,000 xg

for 10 min at 0 °C to remove cellular debris. The supernatant was collected and pre-

cleared by adding 0.5 mg Protein A dynabeads (Cat No.10000D, Thermo Fisher) for 30

min with rotation at 4 °C. The beads were removed by placing tubes on magnetic rack

(Cat. No.12321D, Thermo Fisher). The cleared lysate was split into 3 equal aliquots,

and 3 μg α- SUN2 antibody was added to first aliquot, 3 μg NDPK control antibody to

the second tube and no antibody added to the third tube. The tubes were incubated at

4 °C with rotation overnight, then 0.5 mg Protein A magnetic beads were added to

each tube for one-hour incubation at 4 °C. The antigen-antibody bound beads were

captured with the help of a magnetic rack and washed thrice with PBS supplemented

with 0.02% Tween-20. The protein complexes bound to the beads were digested using

ProteoExtract All-in-One Trypsin Digestion Kit (Cat. No. 650212, Calbiochem)

according to manufacturer's instructions. Briefly, beads were resuspended in extraction

buffer and digestion buffer was added, then reduced using the provided reducing agent

for 10 minutes at 37°C. Samples were cooled to RT and then blocked using blocking

reagent for 10 minutes at room temperature. Trypsin at a final concentration of 8 ng/μl

was added and incubated for 2 hours at 37°C with shaking. The reaction was stopped

by acidification. Digests were vacuum dried and submitted for LC-MS/MS to the

Translational Lab at Florida State University.

Mass spectrometry and data analysis

Digested peptides were resuspended in 0.1% Formic acid for loading onto LC-MS at the

Translational Lab, Florida State University. An externally calibrated Thermo LTQ

Orbitrap Velos nLC-ESI-LTQ-Orbitrap (high-resolution electrospray tandem mass

spectrometer) was used with the following parameters. A 2 cm, 100 µm i.d. (internal

diameter) trap column (SC001 Easy Column from Thermo-scientific) was followed by a

10 cm analytical column of 75 µm i.d. (SC200 Easy Column from Thermo-scientific).

Both trap column and analytical column had C18-AQ packaging. Separation was carried

out using Easy nanoLC II (Thermo-Scientific) with a continuous, vented column

configuration. A 5 µL (~500 ng) sample was aspirated into a 20 µL sample loop and

loaded onto the trap. The flow rate was set to 300 nL/min for separation on the

analytical column. Mobile phase A was composed of 99.9 H_2O (EMD Omni Solvent) and

0.1% formic acid, and mobile phase B was composed of 99.9% ACN and 0.1% formic

acid. A 1 h linear gradient from 0% to 45% B was performed. The LC eluent was directly

nano-sprayed into an LTQ Orbitrap Velos mass spectrometer (Thermo Scientific).

During the chromatographic separation, the LTQ Orbitrap Velos was operated in a data-

dependent mode and under direct control of the Xcalibur software (Thermo Scientific).

The MS data were acquired using the following parameters: 10 data-dependent

collisional-induced-dissociation (CID) MS/MS scans per full scan. All measurements

were performed at room temperature and three technical replicates. The raw files were

analyzed using Proteome Discoverer (version 1.4) software package with SequestHT

and Mascot search nodes using the maize proteome database available at Gramene

(AGVP3.0). The resulting msf files were further analyzed by the 'Scaffold version 4.4'

proteome validator software at 90% peptide and protein threshold with a minimum of 1

peptide.

Constructs and gene cloning

Coding sequences of MLKP1 and MLKP2 genes fused to the C-terminus of eGFP-

FLAG-HA sequence were custom synthesized by Genscript as pHWBF08 and

pHWBF06 respectively. Sequences eGFP-FLAG-HA-MLKP1 and eGFP-FLAG-HA-

MLKP2 were PCR amplified with att flanking primers and cloned individually into

pDONR221 vector by BP cloning (Cat. No. 1235019, Invitrogen) to generate

pHWBF08EC and pHWBF06EC entry clones respectively. These were moved to

destination vector pH7WG2 (Karimi et al., 2002) by LR recombination reaction of

Gateway cloning system (Cat. No. 1235019, Invitrogen) to obtain expression vectors

pPK1iexp and pPK1Cexp respectively. For MLKG1, the coding region was PCR

amplified from maize B73 using gene specific primers (Table S4) and cloned into pCRII

vector by TOPO cloning (Cat. No. K461020, Thermo Fisher). After sequence

verification, the coding sequence was amplified with forward primer flanked with BamHI

restriction site sequence and reverse primer flanked by SbfI restriction site sequence.

The PCR amplified product was double digested with BamHI and SbfI (NEB) and ligated

in pECGFP (Bass lab) donor vector in frame and downstream of the eGFP-FLAG-HA

sequence. The fusion sequence was then moved to pH7WG2 destination vector by LR

recombination to generate an expression vector called pPK1Dexp. Deletion constructs

for MLKP1 and MLKP2 were generated using corresponding full length entry clones as

template to PCR amplify required regions with internal primers containing flanking att

sequences (Table S4) <mark>with stop codon and cloned into pDONR211 by BP cloning.</mark>

These deletion constructs were subcloned in pH7WG2 destination vector by LR

recombination. MLKG1ΔKASH was generated using the Q5 DNA mutagenesis kit (Cat.

No. E0554S, NEB). Briefly, two primers were designed in opposite directions, going

outwards from the KASH domain. The entire entry vector (pECPK1D) minus the KASH

domain was amplified using high fidelity Q5 polymerase provided with the kit. The

template contamination was removed and vector re-ligated using manufacturer-provided

enzyme mix containing a kinase, a ligase and DpnI to create entry clone

MLKP1delKASH. The entry clone was then moved to pH7WG2 destination vector by

LR recombination. Sequences of all the plasmids generated in this study are given in

Table S5.

***Agrobacterium tumefaciens* transformation**

Using established heat shock protocols, *A. tumefaciens* was transformed with

expression vectors (Graumann and Evans, 2010). Briefly, competent cells were thawed

on ice and 4ul plasmid DNA was added. Cells were incubated for 5 min on ice, 5 min in

liquid nitrogen, and 5 min at 37 °C, then 1 ml LB medium was added and cells were

incubated for 2h at 28 °C before being plated out and left to grow for 48 h at 28 °C.

***Nicotiana benthamiana* transformation and microscopy**

Transient expression of fluorescent fusion proteins was carried out as previously

described (Sparkes et al., 2006). Briefly, Agrobacterium were suspended in infiltration

buffer at an OD of 0.05 and pressed into the lower leaf epidermis with a syringe. Three

days post infiltration, leaf sections were imaged with a Zeiss LSM 800 confocal

microscope using a 100X oil immersion objective, NA 1.46 and 1.5X digital zoom. GFP

was excited at 488 nm and emission collected between 500-550 nm; mCherry was

excited with a 561 nm laser and emission collected from 565-620 nm. For FRAP, we

performed 5 pre scans and then a 160x160 pixel ROI was bleached with the 488 nm

laser at 100% for 10 iterations. Recovery data was collected over 36 s (240 scans, 0.13

s framerate). Mean ROI intensity data was extracted using Zen Blue (version 2.3) and

normalised as described in (Martinière et al., 2012). One phase association recovery

curves, plateau values and halftimes were produced in Graphpad (version 6). The

ANOVA statistical test was performed with multiple comparisons. For DAPI staining, leaf

sections were first incubated in 1 ml water containing 5 ul 20% Triton X-100 and 1ul

DAPI. DAPI excitation was performed with the 405 nm laser and emission between 400-

450 nm imaged. DAPI/GFP or DAPI/mCherry imaging was performed using multitrack

line scanning.

Direct Co-IP and western blot

Leaves of *N. benthamiana* (1 g) were harvested and ground in liquid nitrogen. The

powdered tissue was homogenized in 5 mL of radioimmunoprecipitation assay buffer

(50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% SDS, 0.5% Na-deoxycholate, 1% NP-40,

1 mM PMSF, and 1% protease inhibitor cocktail (Cat. No. P9599, Sigma) in a dounce

homogenizer kept on ice. The homogenate was centrifuged at 10,000 xg and the

supernatant was collected for co-IP. One tenth of the protein extract was used as the

input sample and concentrated using acetone precipitation. The rest of the lysate was

subjected to co-IP using dynabeads coated with rabbit anti-GFP antibody (Cat. No.

ab290, Abcam). The lysate was incubated with beads for 4 hours, after which the beads

were collected using magnetic rack and washed with PBST three times. The antibody-

protein complex bound to beads was eluted using 50 mM Glycine (pH 2.8). The

immunoprecipitates and the input samples were boiled in 4X Laemelli's gel loading

buffer, fractionated on 15% SDS-PAGE and transferred to polyvinylidene difluoride

membranes (Cat. No. 88518. Thermo Scientific), and detected with a mouse anti-GFP

(1:3,000; Cat No. MA5-15256,Thermo Scientific) or rabbit anti-SUN2-NPAP (1:1,000;

Murphy et al 2014) antibody at room temperature for 1 h. After three 15-min washes in

PBS-T buffer at room temperature, the membranes were incubated with a 1:3,000

dilution of anti-rabbit or anti-mouse IgG horseradish peroxidase–linked antibody (Santa

Cruz Biotechnology, Santa Cruz, CA) for 1 h at room temperature, then subjected to 3 ×

15 min washes in PBST buffer at room temperature. The PVDF membrane was

incubated with chemiluminescent reaction kit for 3-5 min at room temperature (Millipore,

Immobilon detection kit, WBKL50100, Billerica, MA) and exposed to X-ray film in dark

room to visualize the protein bands.

FIGURE LEGENDS

Figure 1. **Working model of the plant LINC complex; core and associated**

proteins. Core LINC and associated proteins are depicted as residing in one of the four

cellular locations, each represented by different background shaded colors: green,

cytoplasm (CYTO); yellow, outer nuclear membrane (ONM); pink inner nuclear

membrane (INM); and blue NUC (nucleoplasm). The core LINC proteins (dashed box

enclosed) shown include SUN-domain proteins labeled "SUN", KASH-related proteins

"KASH/MLK", and AtWIT-related proteins labeled "WIT/MLKT". The LINC-associated

proteins shown include cytoplasmic proteins labeled "RanGAP" and "motor proteins,"

and "cytoskeleton"; the INM proteins labeled "NEAP/MNEAP"; and the nucleoplasmic

proteins labeled "NMCP/CRWN/NCH" and "KAKU4/MKAKU4". Maize gene names and

their known Arabidopsis homolog types are listed. This composite model places

proteins together on the basis of previously documented pair-wise interactions

representing a generalized consensus assembly but not meant to depict a single

complex.

Figure 2. Domain organization of candidates for the maize core LINC and

associated proteins complex. Schematic representations of predicted proteins from

canonical transcript models are diagrammed to show the location of domains (color

coded and listed at bottom). The proteins are drawn to scale, using a separate scale for

the last two larger proteins, NCH1 and NCH2. The four C-terminal residues and stop

codon (*) location of the KASH and MNEAP group proteins are indicated. The

previously published SUN-domain proteins (Murphy et al., 2010) are not included. The

partial SMC N-terminal domains in the MNEAPs are not diagrammed but are located at

AA 24-308 for MNEAP1, AA 18-309 for MNEAP2, and at AA 3-109 for MNEAP3.

Potential CDK1 phosphorylation sites are shown (blue circles) for NCH1 and NCH2.

Figure 3. Evolutionary conservation of plant LINC complex components.

Candidate maize LINC gene names, cellular location predictions, gene family group,

and PLAZA gene family ID (each starting with shared prefix of HOMO4M) are listed in

the first four columns.   The other columns provide counts of the numbers of genes for

each species (top row) in each Plaza gene family (4th column). The general grouping

color key is indicated on the top left of the figure. Species included are Zema, *Zea*

mays; Sobi*, Sorghum bicolor,* Seit, *Setaria viridis;* Hovu, *Hordeum vulgare;*

Trae,*Triticum aestivum;* Brdi, *Brachypodium distachyon;* Orsa*, Oryza sativa spp.*

Japonica; Zoma*, Zostera marina;* Arth*, Arabidopsis thaliana;* Potr*, Populus trichocarpa;*

Vivi*, Vitis vinifera;* Metr*, Medicago truncatula;* Piab*, Picea abies;* Phpa*, Physcomitrella*

patens and Chre, *Chlamydomonas reinhardtii.* Broader grouping names are provided

along the top (labeled grey boxes).

Figure 4: Gene expression levels and co-expression clustering of 20 maize LINC

genes. Expression levels of 20 maize candidate LINC genes were analyzed using

published transcriptome data (Stelpflug et al., 2016) and including the genes

designated as housekeeping genes (HKG) in a previous study (Lin et al., 2014). (A)

Expression profiles of LINC genes in several tissues developmental stages. Data are

presented as log10 Fragments Per Kilobase of exon per Million fragments mapped

(FPKM) values from published transcriptome. Hierarchical clustering grouped tissues

according to similar expression profiles. The root, young inflorescence, and meristem

groups are denoted by orange, blue, and green boxes, respectively. The growth stage

of maize plant at the time of tissue harvesting are shown in parentheses; V, vegetative

stage; R, reproductive stage; DAP, days after pollination. (B) The Median FPKM values

of the tissues shown in panel A are plotted, ranked by value for individual genes.

Housekeeping genes are shown in red.

Figure 5. Subcellular localization of transiently expressed GFP-MLKP fusions in

N. benthamiana. Agrobacterium-mediated leaf transformation with expression

constructs was carried out followed by imaging three days later using 100X lens with

confocal microscopy to detect DAPI (405 nm excitation laser), GFP (488 nm excitation

laser), or mCherry/mRFP (561 nm excitation laser). Representative focal sections are

shown for GFP (green), mCherry (green in panel A, bottom row) mRFP (majenta in

panel B), or DAPI (majenta in panels A and D). (A) GFP-MLKG1, GFP-MLKP1, GFP-

MLKP2 and mCherry-ZmSUN2 (green) are localized at the nuclear periphery,

surrounding the nucleus (n) stained with DAPI (magenta). (B) GFP-MLKG1 (green) is

also present in a network-like pattern at the cell periphery, where it partially co-localizes

with the ER marker HDEL-mRFP (magenta, top row) and the actin marker lifeact-mRFP

(magenta, bottom row). (C) Schematic overview depicting the terminal deletion

constructs. (D) Images from deletion constructs (listed at left) are shown as GFP only

(1st column), DAPI only (2nd column), or overlay (3rd column). GFP-MLKP1ΔKASH

shows additional localization to the cytoplasm and the nucleoplasm (magenta), whereas

the others retain a tendency to localize to the nuclear periphery. All scale bars 10μm.

Figure 6. Fluorescence recovery after photobleaching (FRAP) co-expression interaction assays of MLK LINC candidates with ZmSUN2 in *N. benthamiana*. (A) Schematic overview depicting the full length mCherry-ZmSUN2 and internal deletion constructs that removed the coiled coil domain (ΔCC) or the SUN domain (ΔSUN). Normalized averaged (n=26-37) intensity FRAP curves for (B) GFP-MLKG1, (D) GFP-MKLP1, or (F) GFP-MLKP2 expressed alone (green), with full-length mCherry-SUN2 (black), with mCherry-SUN2-ΔCC (blue), or with mCherry-SUN2-ΔSUN (pink). Plateau values are plotted to the right for assays with GFP-MLKG1 (C), GFP-MLKP1 (E), and GFP-MLKP2 (G). Box plot error bars represent SD (blue) and mean (red) values. ANOVA statistical analyses with multiple comparisons are noted (ns = not significant = p≥0.05, **** = significant with P ≤0.0001).

ACKNOWLEDGEMENTS

Thank Eric Richards and Iris Meier for helpful and insightful comments on the manuscript, Allison Jevitt for artwork on Figure 1, Dr. Rakesh K Singh and staff of the Translational Lab (College of Medicine, FSU) for assistance and advice on co-IP and Mass spectrometry. The work was supported by grants from the National Science Foundation to HWB (NSF IOS 1444532) and Oxford Brookes University to KG, and from a Dissertation Research Grant to HKG (Florida State University Graduate School).

Competing interests

The authors declare no competing or financial interests.

REFERENCES

Alam, S. G., Zhang, Q., Prasad, N., Li, Y., Chamala, S., Kuchibhotla, R., Kc, B., Aggarwal, V., Shrestha, S., Jones, A. L., et al. (2016). The mammalian LINC complex regulates genome transcriptional responses to substrate rigidity. *Sci. Rep.* **6**, 38063.

Armstrong, S. J., Franklin, F. C. and Jones, G. H. (2001). Nucleolus-associated telomere clustering and pairing precede meiotic chromosome synapsis in Arabidopsis thaliana. *J. Cell Sci.* **114**, 4207–4217.

Belmont, A. S., Sedat, J. W. and Agard, D. A. (1987). A three-dimensional approach to mitotic chromosome structure: evidence for a complex hierarchical organization. *J. Cell Biol.* **105**, 77–92.

Boruc, J., Zhou, X. and Meier, I. (2012). Dynamics of the plant nuclear envelope and nuclear pore. *Plant Physiol.* **158**, 78–86.

Brendel, V., Kurtz, S. and Walbot, V. (2002). Comparative genomics of Arabidopsis and maize: prospects and limitations. *Genome Biol.* **3**, REVIEWS1005.

Broers, J. L. V., Ramaekers, F. C. S., Bonne, G., Yaou, R. B. and Hutchison, C. J. (2006). Nuclear lamins: laminopathies and their role in premature ageing. *Physiol. Rev.* **86**, 967–1008.

Burke, B. and Stewart, C. L. (2014). Chapter One - Functional Architecture of the Cell's Nucleus in Development, Aging, and Disease. In *Current Topics in Developmental Biology* (ed. Stewart, C. L.), pp. 1–52. Academic Press.

Chettoor, A. M., Givan, S. A., Cole, R. A., Coker, C. T., Unger-Wallace, E., Vejlupkova, Z., Vollbrecht, E., Fowler, J. E. and Evans, M. M. (2014). Discovery of novel transcripts and gametophytic functions via RNA-seq analysis of maize gametophytic transcriptomes. *Genome Biol.* **15**, 414.

Ciska, M., Masuda, K. and Moreno Díaz de la Espina, S. (2013). Lamin-like analogues in plants: the characterization of NMCP1 in Allium cepa. *J. Exp. Bot.* **64**, 1553–1564.

Coates, J. C. (2003). Armadillo repeat proteins: beyond the animal kingdom. *Trends Cell Biol.* **13**, 463–471.

Conklin, P. A., Strable, J., Li, S. and Scanlon, M. J. (2018). On the mechanisms of development in monocot and eudicot leaves. *New Phytol.*

Crisp, M., Liu, Q., Roux, K., Rattner, J. B., Shanahan, C., Burke, B., Stahl, P. D. and Hodzic, D. (2006). Coupling of the nucleus and cytoplasm: role of the LINC complex. *J. Cell Biol.* **172**, 41–53.

Dittmer, T. A. and Richards, E. J. (2008). Role of LINC proteins in plant nuclear
morphology. *Plant Signal. Behav.* **3**, 485–487.

Dukowic-Schulze, S., Harris, A., Li, J., Sundararajan, A., Mudge, J., Retzel, E. F.,
Pawlowski, W. P. and Chen, C. (2014). Comparative transcriptomics of early
meiosis in Arabidopsis and maize. *J. Genet. Genomics* **41**, 139–152.

Fridkin, A., Penkner, A., Jantsch, V. and Gruenbaum, Y. (2009). SUN-domain and
KASH-domain proteins during development, meiosis and disease. *Cell. Mol. Life*
Sci. **66**, 1518–1533.

Friederichs, J. M., Ghosh, S., Smoyer, C. J., McCroskey, S., Miller, B. D., Weaver,
K. J., Delventhal, K. M., Unruh, J., Slaughter, B. D. and Jaspersen, S. L. (2011).
The SUN protein Mps3 is required for spindle pole body insertion into the nuclear
membrane and nuclear envelope homeostasis. *PLoS Genet.* **7**, e1002365.

Gaut, B. S., Le Thierry d'Ennequin, M., Peek, A. S. and Sawkins, M. C. (2000).
Maize as a model for the evolution of plant nuclear genomes. *Proc. Natl. Acad. Sci.*
U. S. A. **97**, 7008–7015.

Gerber, A. P., Herschlag, D. and Brown, P. O. (2004). Extensive association of
functionally and cytotopically related mRNAs with Puf family RNA-binding proteins
in yeast. *PLoS Biol.* **2**, E79.

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J.
F., Pretty, J., Robinson, S., Thomas, S. M. and Toulmin, C. (2010). Food
security: the challenge of feeding 9 billion people. *Science* **327**, 812–818.

Goto, C., Tamura, K., Fukao, Y., Shimada, T. and Hara-Nishimura, I. (2014). The
Novel Nuclear Envelope Protein KAKU4 Modulates Nuclear Morphology in
Arabidopsis. *Plant Cell* **26**, 2143–2155.

Graumann, K. and Evans, D. E. (2010). Plant SUN domain proteins: components of
putative plant LINC complexes? *Plant Signal. Behav.* **5**, 154–156.

Graumann, K., Runions, J. and Evans, D. E. (2010). Characterization of SUN-domain
proteins at the higher plant nuclear envelope. *Plant J.* **61**, 134–144.

Graumann, K., Vanrobays, E., Tutois, S., Probst, A. V., Evans, D. E. and Tatout, C.
(2014). Characterization of two distinct subfamilies of SUN-domain proteins in
Arabidopsis and their interactions with the novel KASH-domain protein AtTIK. *J.*
Exp. Bot. **65**, 6499–6512.

Grob, S., Schmid, M. W. and Grossniklaus, U. (2014). Hi-C analysis in Arabidopsis
identifies the KNOT, a structure with similarities to the flamenco locus of
Drosophila. *Mol. Cell* **55**, 678–693.

Groves, N. R., Biel, A. M., Newman-Griffis, A. H. and Meier, I. (2018). Dynamic

Changes in Plant Nuclear Organization in Response to Environmental and
Developmental Signals. *Plant Physiol.* **176**, 230–241.

Hagan, I. and Yanagida, M. (1995). The product of the spindle formation gene sad1+
associates with the fission yeast spindle pole body and is essential for viability. *J.*
Cell Biol. **129**, 1033–1047.

Haque, F., Lloyd, D. J., Smallwood, D. T., Dent, C. L., Shanahan, C. M., Fry, A. M.,
Trembath, R. C. and Shackleton, S. (2006). SUN1 interacts with nuclear lamin A
and cytoplasmic nesprins to provide a physical connection between the nuclear
lamina and the cytoskeleton. *Mol. Cell. Biol.* **26**, 3738–3751.

Hieda, M. (2017). Implications for Diverse Functions of the LINC Complexes Based on
the Structure. *Cells* **6**,.

Hirsch, C. N., Hirsch, C. D., Brohammer, A. B., Bowman, M. J., Soifer, I., Barad, O.,
Shem-Tov, D., Baruch, K., Lu, F., Hernandez, A. G., et al. (2016). Draft
Assembly of Elite Inbred Line PH207 Provides Insights into Genomic and
Transcriptome Diversity in Maize. *Plant Cell* **28**, 2700–2714.

Hodzic, D. M., Yeater, D. B., Bengtsson, L., Otto, H. and Stahl, P. D. (2004). Sun2 is
a novel mammalian inner nuclear membrane protein. *J. Biol. Chem.* **279**, 25805–
25812.

Horigome, C., Okada, T., Shimazu, K., Gasser, S. M. and Mizuta, K. (2011).
Ribosome biogenesis factors bind a nuclear envelope SUN domain protein to
cluster yeast telomeres. *EMBO J.* **30**, 3799–3811.

Hotta, T., Haraguchi, T. and Mizuno, K. (2007). A novel function of plant histone H1:
microtubule nucleation and continuous plus end association. *Cell Struct. Funct.* **32**,
79–87.

Hurel, A., Phillips, D., Vrielynck, N., Mézard, C., Grelon, M. and Christophorou, N.
(2018). A cytological approach to studying meiotic recombination and chromosome
dynamics in Arabidopsis thaliana male meiocytes in three dimensions. *Plant J.* **95**,
385–396.

Jafferali, M. H., Vijayaraghavan, B., Figueroa, R. A., Crafoord, E., Gudise, S.,
Larsson, V. J. and Hallberg, E. (2014). MCLIP, an effective method to detect
interactions of transmembrane proteins of the nuclear envelope in live cells.
Biochim. Biophys. Acta **1838**, 2399–2403.

Janin, A., Bauer, D., Ratti, F., Millat, G. and Méjat, A. (2017). Nuclear envelopathies:
a complex LINC between nuclear envelope and pathology. *Orphanet J. Rare Dis.*
12, 147.

Jiao, Y., Peluso, P., Shi, J., Liang, T., Stitzer, M. C., Wang, B., Campbell, M. S.,
Stein, J. C., Wei, X., Chin, C.-S., et al. (2017). Improved maize reference genome

with single-molecule technologies. *Nature* **546**, 524–527.

Käll, L., Krogh, A. and Sonnhammer, E. L. L. (2004). A combined transmembrane
topology and signal peptide prediction method. *J. Mol. Biol.* **338**, 1027–1036.

Karimi, M., Inzé, D. and Depicker, A. (2002). GATEWAY^{TM} vectors for Agrobacterium-
mediated plant transformation. *Trends Plant Sci.* **7**, 193–195.

Kim, D. I., Birendra, K. C. and Roux, K. J. (2015). Making the LINC: SUN and KASH
protein interactions. *Biol. Chem.* **396**, 295–310.

Kopylov, M., Bass, H. W. and Stroupe, M. E. (2015). The maize (Zea mays L.)
nucleoside diphosphate kinase1 (ZmNDPK1) gene encodes a human NM23-H2
homologue that binds and stabilizes G-quadruplex DNA. *Biochemistry* **54**, 1743–
1757.

Kracklauer, M. P., Link, J. and Alsheimer, M. (2013). LINCing the nuclear envelope to
gametogenesis. *Curr. Top. Dev. Biol.* **102**, 127–157.

Krogh, A., Larsson, B., Von Heijne, G. and Sonnhammer, E. L. L. (2001). Predicting
transmembrane protein topology with a hidden Markov model: application to
complete genomes. *J. Mol. Biol.* **305**, 567–580.

Kumar, S., Stecher, G. and Tamura, K. (2016). MEGA7: Molecular Evolutionary
Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874.

Li, P. and Noegel, A. A. (2015). Inner nuclear envelope protein SUN1 plays a
prominent role in mammalian mRNA export. *Nucleic Acids Res.* **43**, 9874–9888.

Lin, F., Jiang, L., Liu, Y., Lv, Y., Dai, H. and Zhao, H. (2014). Genome-wide
identification of housekeeping genes in maize. *Plant Mol. Biol.* **86**, 543–554.

Liu, W., Xie, Y., Ma, J., Luo, X., Nie, P., Zuo, Z., Lahrmann, U., Zhao, Q., Zheng, Y.,
Zhao, Y., et al. (2015). IBS: an illustrator for the presentation and visualization of
biological sequences. *Bioinformatics* **31**, 3359–3361.

Luxton, G. W. G. and Starr, D. A. (2014). KASHing up with the nucleus: novel
functional roles of KASH proteins at the cytoplasmic surface of the nucleus. *Curr.*
Opin. Cell Biol. **28**, 69–75.

Malone, C. J., Fixsen, W. D., Horvitz, H. R. and Han, M. (1999). UNC-84 localizes to
the nuclear envelope and is required for nuclear migration and anchoring during C.
elegans development. *Development* **126**, 3171–3181.

Martinière, A., Lavagi, I., Nageswaran, G., Rolfe, D. J., Maneta-Peyret, L., Luu, D.-
T., Botchway, S. W., Webb, S. E. D., Mongrand, S., Maurel, C., et al. (2012). Cell
wall constrains lateral diffusion of plant plasma-membrane proteins. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 12805–12810.

Masuda, K., Xu, Z. J., Takahashi, S., Ito, A., Ono, M., Nomura, K. and Inoue, M.
(1997). Peripheral framework of carrot cell nucleus contains a novel protein
predicted to exhibit a long alpha-helical domain. *Exp. Cell Res.* **232**, 173–181.

Meier, I. (2016). LINCing the eukaryotic tree of life - towards a broad evolutionary
comparison of nucleocytoplasmic bridging complexes. *J. Cell Sci.* **129**, 3523–3531.

Meier, I., Zhou, X., Brkljacić, J., Rose, A., Zhao, Q. and Xu, X. M. (2010). Targeting
proteins to the plant nuclear envelope. *Biochem. Soc. Trans.* **38**, 733–740.

Meier, I., Richards, E. J. and Evans, D. E. (2017). Cell Biology of the Plant Nucleus.
Annu. Rev. Plant Biol. **68**, 139–172.

Mochizuki, R., Tsugama, D., Yamazaki, M., Fujino, K. and Masuda, K. (2017).
Identification of candidates for interacting partners of the tail domain of DcNMCP1,
a major component of the Daucus carota nuclear lamina-like structure. *Nucleus* **8**,
312–322.

Moriguchi, K., Suzuki, T., Ito, Y., Yamazaki, Y., Niwa, Y. and Kurata, N. (2005).
Functional isolation of novel nuclear proteins showing a variety of subnuclear
localizations. *Plant Cell* **17**, 389–403.

Murphy, S. P., Simmons, C. R. and Bass, H. W. (2010). Structure and expression of
the maize (Zea mays L.) SUN-domain protein gene family: evidence for the
existence of two divergent classes of SUN proteins in plants. *BMC Plant Biol.* **10**,
269.

Murphy, S. P., Gumber, H. K., Mao, Y. and Bass, H. W. (2014). A dynamic meiotic
SUN belt includes the zygotene-stage telomere bouquet and is disrupted in
chromosome segregation mutants of maize (Zea mays L.). *Front. Plant Sci.* **5**, 314.

Nakayama, T., Ishii, T., Hotta, T. and Mizuno, K. (2008). Radial microtubule
organization by histone H1 on nuclei of cultured tobacco BY-2 cells. *J. Biol. Chem.*
283, 16632–16640.

Nannas, N. J. and Dawe, R. K. (2015). Genetic and genomic toolbox of Zea mays.
Genetics **199**, 655–669.

Oda, Y. and Fukuda, H. (2011). Dynamics of Arabidopsis SUN proteins during mitosis
and their involvement in nuclear shaping. *Plant J.* **66**, 629–641.

Pawar, V., Poulet, A., Détourné, G., Tatout, C., Vanrobays, E., Evans, D. E. and
Graumann, K. (2016). A novel family of plant nuclear envelope-associated
proteins. *J. Exp. Bot.* **67**, 5699–5710.

Poulet, A., Probst, A. V., Graumann, K., Tatout, C. and Evans, D. (2017a). Exploring
the evolution of the proteins of the plant nuclear envelope. *Nucleus* **8**, 46–59.

Poulet, A., Duc, C., Voisin, M., Desset, S., Tutois, S., Vanrobays, E., Benoit, M.,
Evans, D. E., Probst, A. V. and Tatout, C. (2017b). The LINC complex contributes
to heterochromatin organisation and transcriptional gene silencing in plants. *J. Cell*
Sci. **130**, 590–601.

Razafsky, D. and Hodzic, D. (2015). Nuclear envelope: positioning nuclei and
organizing synapses. *Curr. Opin. Cell Biol.* **34**, 84–93.

Shimada, N., Inouye, K., Sawai, S. and Kawata, T. (2010). SunB, a novel Sad1 and
UNC-84 domain-containing protein required for development of Dictyostelium
discoideum. *Dev. Growth Differ.* **52**, 577–590.

Sohaskey, M. L., Jiang, Y., Zhao, J. J., Mohr, A., Roemer, F. and Harland, R. M.
(2010). Osteopotentia regulates osteoblast maturation, bone formation, and
skeletal integrity in mice. *J. Cell Biol.* **189**, 511–525.

Sosa, B. A., Rothballer, A., Kutay, U. and Schwartz, T. U. (2012). LINC complexes
form by binding of three KASH peptides to domain interfaces of trimeric SUN
proteins. *Cell* **149**, 1035–1047.

Sparkes, I. A., Runions, J., Kearns, A. and Hawes, C. (2006). Rapid, transient
expression of fluorescent fusion proteins in tobacco plants and generation of stably
transformed plants. *Nat. Protoc.* **1**, 2019–2025.

Spichal, M. and Fabre, E. (2017). The Emerging Role of the Cytoskeleton in
Chromosome Dynamics. *Front. Genet.* **8**, 60.

Starr, D. A. and Fridolfsson, H. N. (2010). Interactions between nuclei and the
cytoskeleton are mediated by SUN-KASH nuclear-envelope bridges. *Annu. Rev.*
Cell Dev. Biol. **26**, 421–444.

Starr, D. A. and Han, M. (2002). Role of ANC-1 in tethering nuclei to the actin
cytoskeleton. *Science* **298**, 406–409.

Stelpflug, S. C., Sekhon, R. S., Vaillancourt, B., Hirsch, C. N., Buell, C. R., de Leon,
N. and Kaeppler, S. M. (2016). An Expanded Maize Gene Expression Atlas based
on RNA Sequencing and its Use to Explore Root Development. *Plant Genome* **9**,.

Tamura, K., Fukao, Y., Iwamoto, M., Haraguchi, T. and Hara-Nishimura, I. (2010).
Identification and characterization of nuclear pore complex components in
Arabidopsis thaliana. *Plant Cell* **22**, 4084–4097.

Tamura, K., Iwabuchi, K., Fukao, Y., Kondo, M., Okamoto, K., Ueda, H., Nishimura,
M. and Hara-Nishimura, I. (2013). Myosin XI-i links the nuclear membrane to the
cytoskeleton to control nuclear movement and shape in Arabidopsis. *Curr. Biol.* **23**,
1776–1781.

Tamura, K., Goto, C. and Hara-Nishimura, I. (2015). Recent advances in

understanding plant nuclear envelope proteins involved in nuclear morphology. *J.*
Exp. Bot. **66**, 1641–1647.

Tapley, E. C. and Starr, D. A. (2013). Connecting the nucleus to the cytoskeleton by
SUN-KASH bridges across the nuclear envelope. *Curr. Opin. Cell Biol.* **25**, 57–62.

Tiang, C.-L., He, Y. and Pawlowski, W. P. (2012). Chromosome organization and
dynamics during interphase, mitosis, and meiosis in plants. *Plant Physiol.* **158**, 26–
34.

Van Bel, M., Diels, T., Vancaester, E., Kreft, L., Botzki, A., Van de Peer, Y.,
Coppens, F. and Vandepoele, K. (2018). PLAZA 4.0: an integrative resource for
functional, evolutionary and comparative plant genomics. *Nucleic Acids Res.* **46**,
D1190–D1196.

Varas, J., Graumann, K., Osman, K., Pradillo, M., Evans, D. E., Santos, J. L. and
Armstrong, S. J. (2015). Absence of SUN1 and SUN2 proteins in Arabidopsis
thaliana leads to a delay in meiotic progression and defects in synapsis and
recombination. *Plant J.* **81**, 329–346.

Vera, D. L., Madzima, T. F., Labonne, J. D., Alam, M. P., Hoffman, G. G.,
Girimurugan, S. B., Zhang, J., McGinnis, K. M., Dennis, J. H. and Bass, H. W.
(2014). Differential nuclease sensitivity profiling of chromatin reveals biochemical
footprints coupled to gene expression and functional DNA elements in maize. *Plant*
Cell **26**, 3883–3893.

Walley, J. W., Sartor, R. C., Shen, Z., Schmitz, R. J., Wu, K. J., Urich, M. A., Nery, J.
R., Smith, L. G., Schnable, J. C., Ecker, J. R., et al. (2016). Integration of omic
networks in a developmental atlas of maize. *Science* **353**, 814–818.

Wang, G., Wang, F., Wang, G., Wang, F., Zhang, X., Zhong, M., Zhang, J., Lin, D.,
Tang, Y., Xu, Z., et al. (2012). Opaque1 encodes a myosin XI motor protein that is
required for endoplasmic reticulum motility and protein body formation in maize
endosperm. *Plant Cell* **24**, 3447–3462.

Wang, H., Dittmer, T. A. and Richards, E. J. (2013). Arabidopsis CROWDED NUCLEI
(CRWN) proteins are required for nuclear size control and heterochromatin
organization. *BMC Plant Biol.* **13**, 200.

Wang, B., Tseng, E., Regulski, M., Clark, T. A., Hon, T., Jiao, Y., Lu, Z., Olson, A.,
Stein, J. C. and Ware, D. (2016). Unveiling the complexity of the maize
transcriptome by single-molecule long-read sequencing. *Nat. Commun.* **7**, 11708.

Xu, X. M., Meulia, T. and Meier, I. (2007). Anchorage of plant RanGAP to the nuclear
envelope involves novel nuclear-pore-associated proteins. *Curr. Biol.* **17**, 1157–
1163.

Xu, X. H., Chen, H., Sang, Y. L., Wang, F., Ma, J. P., Gao, X.-Q. and Zhang, X. S.

(2012). Identification of genes specifically or preferentially expressed in maize silk
reveals similarity and diversity in transcript abundance of different dry stigmas.
BMC Genomics **13**, 294.

Zeng, L., Zhang, Q., Sun, R., Kong, H., Zhang, N. and Ma, H. (2014). Resolution of
deep angiosperm phylogeny using conserved nuclear genes and estimates of early
divergence times. *Nat. Commun.* **5**, 4956.

Zhao, Q., Brkljacic, J. and Meier, I. (2008). Two distinct interacting classes of nuclear
envelope-associated coiled-coil proteins are required for the tissue-specific nuclear
envelope targeting of Arabidopsis RanGAP. *Plant Cell* **20**, 1639–1651.

Zhou, X., Graumann, K., Evans, D. E. and Meier, I. (2012). Novel plant SUN–KASH
bridges are involved in RanGAP anchoring and nuclear shape determination. *J.
Cell Biol.* **196**, 203–211.

Zhou, X., Graumann, K., Wirthmueller, L., Jones, J. D. G. and Meier, I. (2014).
Identification of unique SUN-interacting nuclear envelope proteins with diverse
functions in plants. *J. Cell Biol.* **205**, 677–692.

Zhou, X., Graumann, K. and Meier, I. (2015a). The plant nuclear envelope as a
multifunctional platform LINCed by SUN and KASH. *J. Exp. Bot.* **66**, 1649–1659.

Zhou, X., Groves, N. R. and Meier, I. (2015b). Plant nuclear shape is independently
determined by the SUN-WIP-WIT2-myosin XI-i complex and CRWN1. *Nucleus* **6**,
144–153.

Zhou, X., Groves, N. R. and Meier, I. (2015c). SUN anchors pollen WIP-WIT
complexes at the vegetative nuclear envelope and is necessary for pollen tube
targeting and fertility. *J. Exp. Bot.* **66**, 7299–7307.

1204

Table 1. Maize LINC genes

Short Name[A]	Full Name[A]	Gene id[B]	Length (AA)	Domains		
				Coiled coil	TMD	Other (location)
MLKG1	Maize LINC KASH Grass-specific1	Zm00001d002723	258	74-94	224-246	KASH (247-258)
MLKG2	Maize LINC KASH Grass-specific2	Zm00001d038539	227	-	19-214	KASH (215-227)
MLKS1	Maize LINC KASH AtSINE-like1	Zm00001d031134	617	-	579-601	KASH (602-617)
MLKS2	Maize LINC KASH AtSINE-like2	Zm00001d052955	637	-	605-623	KASH (624-637), ARM (3-517)
MLKP1	Maize LINC KASH AtWIP-like1	Zm00001d003334	308	82-105, 123-150,163-197, 236-270	278-300	KASH (301-308)
MLKP2	Maize LINC KASH AtWIP-like2	Zm00001d025667	290	68-91, 109-136, 152-179	258-280	KASH (281-290)
MLKP3	Maize LINC KASH AtWIP-like3	Zm00001d005997	538	319-343, 358-378, 452-500	506-529	KASH (530-538)
MLKP4	Maize LINC KASH AtWIP-like4	Zm00001d020879	532	313-337, 352-372, 401-421, 446-480	502-523	KASH (524- 532)
MLKT1	Maize LINC KASH AtWIT-like1	Zm00001d039272	653	94-114, 135-155, 201-228, 374-415, 455-517	620-640	KASH (641-653)
MLKT2	Maize LINC KASH AtWIT-like2	Zm00001d010047	667	96-116, 130-157, 218-238, 316-392, 411-431, 453-473,481-508, 527-547	643-661	KASH (662-667)
ZmSUN1	SUN domain protein1	Zm00001d015450	462	169-189, 195-229	116-141	SUN (277-455)
ZmSUN2	SUN domain protein2	Zm00001d040331	439	168-188	84-109	SUN (264-478)
ZmSUN3	SUN domain protein3	Zm00001d042643	613	484-511	33-55, 554-574, 595-612	SUN (194 -358)
ZmSUN4	SUN domain protein4	Zm00001d012240	639	510 -537	57-79, 580-600, 621-638	SUN (218-382)
ZmSUN5	SUN domain protein5	Zm00001d011277	607	414-434, 495-523	46-66, 525-544, 572-588	SUN (152-322)
MNEAP1	Maize Nuclear Envelope-Associated Protein1	Zm00001d010918	348	221-248	330-346	NA
MNEAP2	Maize Nuclear Envelope-Associated Protein2	Zm00001d024756	347	149-83, 220-247	329-345	NA
MNEAP3	Maize Nuclear Envelope-Associated Protein3	Zm00001d002875	170	14-34, 84-104	125-150	NA
NCH1	NMCP/CRWN-Homologous1	Zm00001d051600	1156	160-278, 312-350, 362-522, 537-579, 592-701	NA	NA
NCH2	NMCP/CRWN-Homologous2	Zm00001d043335	896	131-165, 250-281, 309-336, 386-532, 635-721	NA	NA
MKAKU41	Maize KAKU4-Like1	Zm00001d026487	554	NA	NA	NA
MKAKU42	Maize KAKU4-Like2	Zm00001d002012	591	NA	NA	NA

1206

1207   [A]Gene names approved by and listed at MaizeGDB (maizegdb.org)

1208   [B]B73v4 gene models from MaizeGDB.org as of December 2017. Detailed version of this table with B73v3,

1209   PLAZA and UniPROT gene ids is provided as Table S1.

1210

1211

 Table 2. Proteins co-precipitated with ZmSUN2 pull down on maize ear shoot nuclei

	Maize gene id (B73v4)	Maize gene name	Gene Description	Exclusive unique peptide count (IP46)	SUN IP (Normalized total spectra)	Irrelevant antibody control IP	No Antibody control IP
Core LINC components	Zm00001d040331	ZmSUN2	SUN domain protein2	11	384.45	0	0
	Zm00001d015450	ZmSUN1	SUN domain protein1	3	78.398	0	0
	Zm00001d051600	NCH1	Protein CROWDED NUCLEI 1	2	82.803	0	0
	Zm00001d043335	NCH2	Nuclear lamina component	3	28.871	0	0
	Zm00001d002723	MLKG1	OSJNBa0010H02.21-like protein	2	14.459	0	0
	Zm00001d052955	MLKS2	ARM repeat superfamily protein	2	1.2886	0	0
	Zm00001d010047	MLKT2	WPP domain-interacting tail-anchored protein 1	1	26.117	0	0
Nucleoporins	Zm00001d052282	NUA	Nuclear-pore anchor	7	87.161	0	0
	Zm00001d048023	NUP133	Nuclear pore complex protein NUP133	1	18.911	0	0
	Zm00001d046566	NUP96	Nuclear pore complex protein NUP96	1	7.253	0	0
	Zm00001d026263	NUP88	Nuclear pore complex protein NUP88	1	7.206	0	0
	Zm00001d005355	GP210	Nuclear pore complex protein GP210	1	7.206	0	0
Cytoskeletal elements	Zm00001d013410	Actin-1	ARP7 Nuclear actin	3	82.756	0	0
	Zm00001d012556	Tub6	beta tubulin6	5	40.576	0	0
	Zm00001d015348	Tub4	Tubulin beta-4 chain	1	7.253	0	0
Ribosomal proteins	Zm00001d044287	L19	Ribosomal protein L19	2	40.623	0	0
	Zm00001d017047	L6	60S ribosomal protein L6	1	30.569	0	0
	Zm00001d022463	L22-2	60S ribosomal protein L22-2	1	11.658	0	0
RAN	Zm00001d017239	RanGAP2	RAN GTPase-activating protein 2	1	14.459	0	0
	Zm00001d048223	RABA6a	Ras-related protein RABA6a	2	14.412	0	0
DNA/RNA binding	Zm00001d052820	RNA binding protein	RNA-binding (RRM/RBD/RNP motifs) family protein	2	21.665	0	0
	Zm00001d025638	bZIP5	Basic-leucine zipper domain	1	11.658	0	0
	Zm00001d042267	ARF10	Auxin response factor 10	1	11.658	0	0
	Zm00001d011297	MYBR35	Putative MYB DNA-binding domain superfamily protein	1	7.206	0	0
	Zm00001d016354	bHLH72/bHLH041	Putative transcription factor bHLH041	1	7.206	0	0
Histone	Zm00001d034479	H1a	Histone 1 (H1)	1	26.117	0	0

1213

Figure 1

Figure 2

Figure 3

Grouping:
- Viridiplantae (Green plants): all columns
- Land plants
- Seed plants
- Angiosperms (Flowering plants)
- Monocots: Zema, Sobi, Seit, Hovu, Trae, Brdi, Orsa, Zoma
- Eudicots: Arth, Potr, Vivi, Soly, Metr
- Poaceae (Grasses): Zema, Sobi, Seit, Hovu, Trae, Brdi
- Gymno: Piab
- Bryophyte: Phpa
- Algae: Chre

Gene names	Location	Group	Plaza gene family (HOMO4M)	Zema	Sobi	Seit	Hovu	Trae	Brdi	Orsa	Zoma	Arth	Potr	Vivi	Soly	Metr	Piab	Phpa	Chre
MLKG1	ONM	KASH, grass	009187	1	1	1	1	3	1	1	0	0	0	0	0	0	0	0	0
MLKG2	ONM	KASH, grass	008346	1	1	1	1	6	1	0	0	0	0	0	0	0	0	0	0
MLKS1- MLKS2	ONM	KASH, SINE	002474	2	2	2	1	6	2	2	1	2	2	1	1	1	2	7	0
MLKP1 – MLKP4	ONM	KASH, WIP	002181	4	3	3	2	7	3	3	1	3	2	1	1	2	0	0	0
MLKT1-MLKT2	ONM	WIT	000144	2	1	2	2	5	1	2	2	2	1	2	2	1	0	0	0
ZmSUN1- ZmSUN2	INM	C-term SUN	002694	2	2	2	2	6	2	2	2	2	1	1	1	1	2	0	0
ZmSUN3- ZmSUN5	INM	mid-SUN	001731	3	2	2	2	6	2	2	3	3	3	2	1	3	2	2	1
MNEAP1 – MNEAP3	INM	NEAP	002962	3	2	2	1	2	1	2	2	4	2	2	1	1	0	0	0
NCH1	NUC	NMCP/CRWN	004640	1	1	1	2	3	1	1	4	1	2	2	2	2	2	0	0
NCH2	NUC	NMCP/CRWN	003564	1	1	1	1	3	1	1	4	3	3	1	1	1	2	2	0
MKAKU41- MKAKU42	NUC	KAKU4	004991	2	1	1	1	3	1	1	2	1	2	1	1	1	0	0	0

Figure 4

A

B

Figure 5

Figure 6

Supplementary Figure legends

Figure S1. MLKP1 and MLKP2 deletion constructs. The fluorescent fusion protein (eGFP) constructs for MLKP1 and MKLP2 are depicted. (A) Schematic representation showing the location of the transmembrane and KASH domains in full length and deletion constructs. Three alanine residues added to the C-terminus of ΔKASH constructs are highlighted in yellow. (B) Pairwise protein sequence alignment (Clustal Omega) of MLKP1 and MLKP2 is shown. (C) Posterior probabilities (Phobius transmembrane topology predictor) of TM helix for C-termini of MLKP1 and MLKP2 are plotted.

Figure S2. Additional subcellular localization staining patterns of GFP-KASH constructs from *N. benthamiana* transient expression microscopy. GFP localization assays were performed as described for Figure 5. (A) Schematic overview of domain composition of the MLKP1 constructs. (B) Staining patterns in and around the nucleus (1st three columns) and a the cell periphery (last column) shows that the construct lacking the TM domain and the GFP only are distributed throughout the cytoplasm and nucleoplasm. Scale bars for panel B are 5 μm. (C) Cell periphery images show the cytoplasm near the cell cortex for full length and domain deletion constructs. Gene names are listed on the left for each row; constructs are listed on the top for each type; n.t. = not tested). The MLKP1 terminal deletions (both Δ4 and ΔKASH) result in a more soluble staining pattern compared to the full length. The other deletions retain a pattern typical of ER association. Scale bars are 10 μm.

Figure S3. FRAP images from the co-expression interaction assays.

Representative images of cells during the FRAP assays for the experiments

summarized in Figure 6. The FRAP ROIs (white boxes) are shown along with images taken at the times indicated across the top for (A) GFP-MLKG1, (B) GFP-MLKP1, and (C) GFP-MLKP2. Scale bar denotes 2μm. FRAP ROI shown by white box. All scale bars 2μm.

Figure S4. Normalized intensity FRAP curves co-plotted from all experiments performed in *N. benthamiana* leaves. The GFP fusion protein constructs (MLKG1, MKLP1, and MLKP2) are grouped by columns as labeled at top. The co-expression constructs are grouped by row as labeled at left. Individual data points (blue dots), single experiment recovery curves (black lines) and global average recovery curves (red lines) and replicate numbers (n) are shown in each panel.

Figure S5. FRAP recovery halftime values. The recovery half times (t1/2(s)) are plotted for the full-length and co-expression experiments summarized in Figure 6. The SD (blue lines) and mean (red lines) are shown for (A) GFP-MLKG1, (B) GFP-MLKP1, and (C) GFP-MLKP2. ANOVA statistical analyses with multiple comparisons are noted (ns = not significant = p≥0.05, ** = significant with P ≤0.01, **** = significant with P ≤0.0001). Replicate numbers range from n=26 to n=35, as listed in Supplemental Figure 3.

Figure S6. Co-immunoprecipitation assays of MLKP1 and MLKP2 with ZmSUN2 in *N. benthamiana*. Total protein extracts from *N. benthamiana* leaves harvested 3 days

post-infection were subjected to co-IP using dynabeads coated with anti-GFP antibody. (A) Western Blot of the co-precipitates and inputs with primary antibodies listed on the right side of the blot. The FP construct combinations used for infiltration are listed on the top of the blot. The numbers on the left are protein molecular weight standards. (B) A summary of KASH-SUN interaction evidence tabulated from this experiment (tobacco heterologous co-IP) or other experiments in this study (maize endogenous co-IP from Table 2, Heterologous FRAP form Figure 6 and Supplemental Figures 2-4).

Figure S7. Co-localization of proteins from GFP-KASH and mCherry-SUN constructs. Co-localization of GFP-MLKG1, GFP-MLKP1 and GFP-MLKP2 (green) with mCherry-ZmSUN2, mCherry-ZmSUN2ΔCC and mCherry-ZmSUN2ΔSUN (magenta) at the nuclear periphery. Size bar 5μm.

Supplemental Figure 1

A

B

C

Supplemental Figure 2

A

MLKP1 full length eGFP — MLKP1 TM KASH domain - A V P T ∗

MLKP1 ΔTM eGFP — MLKP1 ∗

B

	GFP	DAPI	Merge	Cortex
MLKP1ΔTM				
free GFP				

C

	Full length	Δ4	ΔKASH
MLKG1		(n.t.)	
MLKP1			
MLKP2			

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

Supplemental Figure 6

A **Heterologous co-IP assay of MLKP1 and MLKP2 interaction with ZmSUN2**

	Input	co-IP

B **Summary of evidence for *in planta* maize KASH-SUN2 interaction**

Gene	Endogenous co-IP (Maize earshoots)	Heterologous co-IP (Tobacco leaf)	Heterologous FRAP (Tobacco leaf)
MLKG1	**+**	(not tested)	**+**
MLKP1	**−**[a]	**+**	**−**[b]
MLKP2	**−**[a]	**+**	**+**

Notes:
[a] expression low for this gene in this tissue
[b] effect observed, but below p-value significance, most variable FRAP curves

Supplemental Figure 7