The evolutionary potential of diet-dependent effects on lifespan and fecundity in a

multi-parental population of *Drosophila melanogaster*

Enoch Ng'oma[1*], Wilton Fidelis[1], Kevin M. Middleton[2] and Elizabeth G. King[1]

[1]Division of Biological Sciences, University of Missouri, Columbia, MO 65211

[2]Department of Pathology and Anatomical Sciences, University of Missouri School of

Medicine, Columbia, MO 65212

*Corresponding author:

8   Enoch Ng'oma

9   401 Tucker Hall

10   Division of Biological Sciences

11   University of Missouri, Columbia, MO 65211

12   Tel.: 573-882-8562

13   E-mail: ngomae@missouri.edu

15 Running title: Quantitative genetics of life history in an MPP

Abstract

The nutritional conditions experienced by a population play a major role in shaping trait evolution in many taxa. Constraints exerted by nutrient limitation or nutrient imbalance can influence the maximal value that fitness components such as reproduction and lifespan attains, and organisms may shift how resources are allocated to different structures and functions in response to changes in nutrition. Whether the phenotypic changes associated with changes in nutrition represent an adaptive response is largely unknown, and whether the response of fitness components to diet even has the potential to evolve in response to selection is not generally known. In this study, we use an admixed multiparental population of *Drosophila melanogaster* reared in three different diet conditions to estimate quantitative genetic parameters for lifespan and fecundity. We develop a new image analysis technique for estimating fecundity, removing the need to hand count eggs and use this method to show there is significant genetic variation for fecundity in our population. Our results also show that lifespan has moderate to high heritabilities within diets, and genetic correlations for lifespan between diets that are less than one demonstrate a genotype by diet interaction. These findings demonstrate substantial standing genetic variation in our population that is comparable to natural populations and highlights the potential for adaptation to changing nutritional environments.

Key words: life history, heritability, genotype by environment interaction, *Drosophila melanogaster,* multiparental population

Introduction

An organism's diet has a direct influence on nearly all phenotypes by determining the amount of nutrients and energy available to build structures and perform functions. In addition to this unavoidable direct effect, organisms may also alter how resources are allocated to different traits when experiencing different diets. In an environment where resource availability varies across space and/or time, and the optimal allocation of resources changes with diet, we expect to see the evolution of such phenotypically plastic resource allocation strategies.

In many non-human populations including nematode, fruit fly (Chippindale *et al.*, 1993b), killifish (Vrtílek and Reichard, 2014), waterstrider (Kaitala, 1991) and mouse (Sprott, 1997), nutrient limitation without starvation results in lifespan extension, and is often but not always accompanied by a reduction in reproductive output (Hsin and Kenyon, 1999; Barnes *et al.*, 2006; Flatt *et al.*, 2008). These diet-dependent changes have been hypothesized to result from an increased allocation of a limited nutritional resource to somatic maintenance (van Noordwijk and de Jong, 1986; Sohal and Weindruch, 1996; Simmons and Bradley, 1997; Hughes and Reynolds, 2005), and a decreased allocation to reproduction. Previous models predict this pattern is adaptive in environments with fluctuating resource availability (Neel, 1962; Wells, 2009; Fisher *et al.*, 2010; Warren *et al.*, 2013). However, these models have not been formally tested, nor has the evolutionary potential of the focal traits been established in most systems (Ng'oma *et al.*, 2017), which is an important first step.

The capacity for a given trait to evolve in a population depends on the proportion of trait variation that is attributable to genetic variation and the intensity of selection

acting on it (Roff 2006; Brčić-Kostić, 2010). Selection can have a direct effect on trait

evolution if the trait directly affects fitness, or indirectly if the trait is correlated with a trait

under selection (Lande and Arnold, 1983; Hunt *et al.*, 2004). However, even when a

selective pressure is observed to act on the trait, the amount of evolutionary change

depends on the amount of genetic variation expressed for that trait (Fisher, 1930).

Quantitative genetic parameters, such as the heritability of a given trait, remain some of

the more informative parameters we can estimate in order to learn about the genetic

basis of a trait, to predict trait values in a population, and to predict the selection

response, even with the advances in genomics and molecular genetics in recent years.

Studies in many species have frequently demonstrated genotype by diet variation for

complex traits, such as gene expression and fitness in yeast (Gagneur *et al.,* 2013),

cuticular hydrocarbons in *D. simulans* (Ingleby *et al.*, 2013), and metabolic phenotypes

in *D. melanogaster* (Reed *et al.*, 2010, 2014; Qi *et al.*, 2012). These studies suggest

that genotype by environment interactions (GEI) are common and could be important in

the nutrition-dependent response of life history traits such as fecundity and lifespan.

There has been a significant effort to understand the genetic basis for the dietary

restriction (DR) response with both molecular genetic approaches (Mair and Dillin,

2008; Grandison *et al.*, 2009; Teleman, 2010; Nässel *et al.*, 2015) and mapping

approaches (Rikke *et al.*, 2010; Stanley *et al.*, 2017). As is the case for most complex

traits, mapping approaches to identify the genetic loci determining how lifespan changes

with diet have failed to identify the specific contributors. Studies focused on mapping in

natural populations often do not identify the regions of the genome as QTL that harbor

candidate genes identified by molecular approaches (Remolina *et al.*, 2012; Burke,

King, *et al.*, 2014; Reed *et al.*, 2014; Megan Ulmer Carnes *et al.*, 2015) and there are

several studies in which gene expression patterns do not change in the direction one

would expect from these loss-of-function studies (Min *et al.*, 2008; Giannakou *et al.*,

2008; Stanley *et al.*, 2017). In light of these studies, a possible hypothesis is that the DR

response is highly polygenic and best understood by quantifying holistic measures of

genetic variation, such as the heritability and GEI. Additionally, it is possible that the

limited genetic variation included in QTL mapping approaches limits the ability to identify

causative variants that are segregating in natural populations. The majority of QTL

studies in the past have used only two genomes to create the mapping population. More

recently, multiparent populations (MPPs) have been gaining in popularity, where

multiple (typically at least 4, and often many more) genomes are used to seed the

population, thereby including increased genetic variability (Kover *et al.*, 2009; McMullen

et al., 2009; Huang *et al.*, 2011; Aylor *et al.*, 2011; Threadgill and Churchill, 2012; King,

Macdonald, *et al.*, 2012; King, Merkes, *et al.*, 2012; Cubillos *et al.*, 2013).

In this paper, we employ a half-sibling, split environment design using an

admixed multiparent population derived from an established mapping resource, the

Drosophila Synthetic Population Resource (DSPR), to estimate quantitative genetic

parameters for lifespan and fecundity in three different diets. We are able to estimate

the evolutionary potential of how lifespan and fecundity respond to diet in order to

establish, 1) the ability of these traits to evolve in general, 2) demonstrate levels of

genetic variation that are comparable with wild derived populations, and 3) establish the

validity of the evolve and resequence approach using a synthetic population as a base

population.

Methods

Experimental setup

Generation of experimental population. We used an eight-way multi-parent

population, the DSPR, to characterize the phenotypic response of lifespan and

fecundity of mated female *D. melanogaster* to three nutritional environments and to

estimate the heritabilities of these traits. The development, and genetic properties of the

DSPR are described in detail in (King, Macdonald, *et al.*, 2012; King, Merkes, *et al.*,

2012). Briefly, the DSPR is a set of recombinant inbred lines created from 8 inbred

founder lines that were crossed for 50 generations and then inbred via full sibling mating

for 25 generations. To generate our starting population, for each of 835 recombinant

inbred lines (RILs) of the *B* sub-population of the DSPR, five young females that had

mated intra-line in vials, were randomly selected and distributed to each of six cages

(20.3 cm x 20.3 cm x 20.3 cm). Eggs were collected in milk bottles (250 mL) twice on

the next two successive days, each after a 22-hour oviposition period on maintenance

diet (Table S1). Eclosed flies were released to six cages in populations >2,000 per cage

and allowed to mate *en-masse* for five generations. A polystyrene plate, 100 mm x 15

mm (cat. No. FB0875713 Fisher Scientific, USA) containing maintenance food was

supplied to each cage three times a week (Monday, Wednesday, Friday). A separate

micro-plate (60 mm x 15 mm, cat. no. FB0875713A, Fisher Scientific, USA) with moist

cotton wool was supplied to each cage to serve as a supplemental drinking source.

Very thin slices of food with visually estimated 50-90 eggs were transferred to 30 vials

(25 mm x 95 mm, Polystyrene Reload, cat. no. 32-109RL, Genesee Scientific, USA) for

each cage for each successive generation. Egg vials from each cage for each next

generation were distributed across all cages to ensure a genetically homogenous

experimental population. We maintained all populations on a three week cycle (21 days

from oviposition to egg collection) on a cornmeal-dextrose-yeast diet for 5 generations

before the half-sibling experiment was started.

Experimental design. We employed a split-environment, half-sibling breeding

design to determine the phenotypic effects of three dietary conditions and estimate

quantitative genetic parameters. We created a set of paternal half-sibling families by

sequentially mating each of 28 males to three females (here on, sires and dams,

respectively) on a common cornmeal-dextrose-yeast (maintenance) diet (Table S1).

Each dam was housed in its own vial and each sire rotated among these three vials. In

round one, to ensure each dam had a mating opportunity early, the sire spent 2 days

with each dam. Each sire was then rotated among the three dams for two additional

rounds lasting 3.5 days each encounter. Two dams were kept waiting in media vials

when the sire was matched with the third dam. Whole families were restarted with new

flies when a sire died during crossing. In case of death of a waiting unmated dam, only

that dam was replaced, whereas if a mated dam died, a new dam family was restarted.

Starting with 61 sires and 183 dams, 28 half-sibling families and 77 maternal full-sibling

families successfully produced enough offspring to be included in the experiment (Fig.

1).

From each of the 77 dams, in most cases, we set up two replicate vials of 24

female and 6 male offspring in each of three diet treatments (described below). We

included males to provide mating opportunities to females throughout their lifetime but

chose a smaller number of males to reduce harassment of females within the vials. As

expected, even successful sire and dam pairs did not always produce enough offspring

for a full complement. Thus, some vials contained fewer than 24 females though never

below 15 females, and the dataset is not fully balanced with all sire families having all

three dams, split in all three diets, and with two replicates per diet. The realized totals of

sire half-sibling families and dam full-sibling families available for analysis in each diet

are presented in Table S2.

Figure 1: Experimental design. A set of 835 DSPR RILs were crossed for 5 generations

to create an outbred synthetic population. From this population, 28 randomly chosen

sires were each mated to 3 dams (actual N = 77 ♀). The resulting offspring from each

dam were split to each of C, DR, and HS diet treatments with two replicate vials in each

diet.

Diet treatments. Offspring of each sire were split across three experimental diets:

control (C), dietary restriction (DR), and high sugar (HS). These diets were also used in

our previous mapping study using the DSPR RILs by Stanley *et al*. (2017) and the

composition of each is detailed in Table S1. To preserve quality, diets were stored at 4

°C and used within two weeks of preparation. To permit phenotype measurement and

ensure that food did not degrade, individuals were moved to vials with fresh food three

times per week. All flies in all experiments described here were reared in a growth

chamber at 23°C, ≥ 50% relative humidity, and a 24:0 light:dark cycle.

Phenotype measurement

We measured two phenotypes: 1) lifespan - the total number of days each fly

lived from the date of eclosion, and 2) weekly fecundity - the total number of eggs a set

of females in each vial produced within a 24-hour period measured once per week. Both

phenotypes were measured from the same set of experimental vials over a full lifespan.

For lifespan, the number and sex of dead, and number of censored (i.e., escapees,

accidental deaths) events were recorded three times each week during transfer of flies

to new food until all flies in each vial had died. To estimate fecundity, females were

provided with fresh media each Monday and following a 24 hour egg laying period,

these egg vials were collected on Tuesday and stored at -20 °C until processing. We

continued this process each week until all females within a vial had died. We modified a

protocol developed in Michael Rose's laboratory by Larry Gonzaga Cabral at UC Irvine

based on personal communication and video tutorials (Rose Lab Data 2012a,b). A

vacuum apparatus was custom prepared following recommendations in the links above.

We added 2.5 mL of 50% bleach to each vial (frozen or thawed) and gently swirled for

about 2 min 50 s on an ORBITRON Rotator 1 (Boekel Scientific, USA) to separate eggs

from the media. Eggs were flushed from each vial onto a black filter disc fit in a vacuum

apparatus with a brief spray of embryo solution (12% Triton X-100, SIGMA-ALDRICH,

USA) followed by a spray of water to rinse. As a quality check, flask filtrate was re-

filtered to a new disc periodically, and processed vials inspected under a dissecting

microscope. The filter discs were prepared from black landscape fabric (DEWITT Weed

Barrier Pro, www.dewittcompany.com) to a radius of 3.65 cm. Discs bearing eggs were

photographed using a Canon EOS Rebel TS*i* (Canon Inc, Japan) camera vertically

attached to a steel clamp using the following settings: exposure time 1/25 seconds,

aperture F5.6, and ISO 100. Egg-bearing discs were mounted on a black stage placed

within a cubical photo studio 35cm x 35cm x 35cm constructed with PVC frame and

enclosed by a white fabric (Fig. S1). The entire setup was placed in a dark room lit only

by two facing white light lamps, 23W, 1200 lumens (model # LBP23PAR2/5K,

UTILITECH, China) clamped to the stage outside the studio box to allow for soft uniform

lighting across the imaging platform. Image files were uploaded to a computing system

and immediately inspected with ImageJ (Abramoff *et al.*, 2004) for quality.

Estimation of egg counts

We developed a predictive model to automate the counting of eggs from the

images we generated as described above. To train the model and assess its

performance, eggs on 345 discs were manually counted using the *Multi-point* tool in

ImageJ (Abramoff *et al.*, 2004). To ensure that all possible imaging conditions were

available for model training, manually counted discs were selected across the whole

lifespan series and across all three diet treatments. Some background reflectivity of the

discs and small debris present meant that some non-zero white area was present in

almost all images, so our training set includes an over-representation of images with

relatively few eggs, to prevent upward bias in the fecundity predictions.

Each image was manually cropped to be square at the edges of the disc, and the

background outside the disc was set to black to eliminate any reflectivity or debris

occurring outside of the disc using GNU Image Manipulation Program (GIMP v.2.8; The

GIMP team 2015; http://www.gimp.org). Subsequently, the image was binarized by

thresholding 8 bit grayscale values using the opencv (Bradski, 2000) package in python

(ver 3.6; https://github.com/EGKingLab/h2lifespan). From this thresholded image, the

area of white pixels was summed (https://github.com/EGKingLab/h2lifespan). Because

images were not all the same size due to slight variations in camera position, we

included size of the image as a covariate in the linear model predicting egg count from

white pixel area and image dimension:

$$Egg\ Count = \beta_0 + \beta_1\ \sqrt{Area} + \beta_2\ ImageSize$$

where *Area* is the sum of the white pixels and *Image Size* is the linear dimension of one

edge of the image in pixels. We use $\sqrt{Area}$ because pixel area is predicted to scale as a

square of egg count.

The optimal grayscale threshold value was determined by cross-validation using

R (v. 3.5; https://github.com/EGKingLab/h2lifespan). We tested threshold values

between 30 and 150 (for 8-bit images with 256 levels of gray), across a range of

training/testing data splits (0.5 to 0.9) and a range of proportions of data (0.5 to 1.0).

The latter was used to test the adequacy of our data. Each model was fit 1,000 times at

each combination of threshold, train/test split, and data proportion, with the root mean

squared error (RMSe) retained for each iteration

(https://github.com/EGKingLab/h2lifespan). The optimal threshold value was determined

from the mean RMSe for the 1,000 iterations at each combination (Fig. 2a, b). That

threshold was 53 for the training set, using 90% of the data each iteration for training,

and only required using 50% of the total data in any one iteration. The resulting

predictive model performed well for our training set (see Results). The trained model

was then used to predict egg counts for all 3,768 images (Fig. 2c). Images with negative

predicted egg counts were set to 0. All images were also visually inspected, and those

determined to be of poor quality, with excessive background reflectivity, were counted

manually and were not included in the training or prediction set.

Statistical analysis

All following statistical analyses were performed in R (ver. 3.5; R Core Team

2018), and all code is available online (https://github.com/EGKingLab/h2lifespan).

Survival estimates

We converted daily counts of dead and censored individuals to individual events

at a given age to perform survival analysis. Using the R package survival (ver. 2.38;

(Therneau and Grambsch, 2000; Therneau, 2015), we used the Kaplan-Meier estimator

(Kaplan and Meier, 1958) to estimate survival in each diet:

253   survfit(Surv(time, status) ~ diet)

where time is time (days), status is alive/dead, diet is diet treatment (C, DR, HS) and to

visualize survivorship curves in each diet.

To assess statistical differences between diets accounting for family

membership, we used hierarchical Bayesian inference to test for the effects of sire, dam

nested in sire, and treatment using a set of seven nested models, which tested variously

for effects of diet treatment, sire and dam, identical to the models used to test survival

estimates (Table 1). We used the brms interface (Bürkner, 2017) to the statistical

modeling language stan (Carpenter *et al.*, 2017), which samples posterior estimates of

modeled parameters using Hamiltonian Monte Carlo. Lifespan was modeled with a

Weibull distribution, which is commonly used to analyze time to failure (in this case, fly

death). We used broad priors for the Weibull shape parameter ($\Gamma(0.01, 0.01)$), Student's

$t(3, 4, 10)$ for the intercept, and Student's t(3, 0, 10) for the standard deviation, which

were quickly overwhelmed by the large amount of data (7486 observations). Models

were compared using leave-one-out (loo) cross-validation and loo model weights

(Vehtari *et al.*, 2017; Yao *et al.*, 2018), the latter which assigns relative out of sample

predictive ability to each model compared. Posterior estimates of DR and HS diet

treatments were compared to the STD diet treatment via credible intervals. We sampled

each model for 2000 iterations with 50% discarded for warm-up and 4 replicate chains,

to yield approximately 4000 total samples. We found that due to the large amount of

data, this sampling approach was adequate. Adequate sampling was determined by $\hat{R}$

values of 1 (Gelman and Rubin, 1992; Brooks and Gelman, 1998).

Fecundity estimates

We focused on two fecundity measurements from our weekly fecundity

measures. We calculated the number of eggs per female for each of our weekly

measures by dividing by the number of females alive in each vial. We then obtained an

estimate of total fecundity per female by simply summing across weeks (hereafter

referred to as total fecundity). We note that this is not strictly a measure of lifetime

fecundity as we measured fecundity only once per week, though we would expect this

estimate to be highly correlated with lifetime fecundity. Second, we considered a

snapshot of early life fecundity by choosing the time point closest to 5 days post-

eclosion (hereafter referred to as early fecundity). The actual age of the females varies

slightly as the vials were set up over several days, but our fecundity measurements

always took place on a 24 hour period beginning on Monday.

We tested for the effects of sire, dam nested in sire, and treatment using a set of

seven nested hierarchical (mixed) linear models (Gelman and Hill, 2007), which tested

variously for effects of diet treatment, sire and dam, identical to the models used to test

survival estimates (Table 1). Sampling was done on zero-centered values for early life

and total fecundity using Hamiltonian Monte Carlo using the statistical modeling

language stan (Carpenter *et al.*, 2017), via the rstanarm interface (ver. 2.13.1; (Stan

Development Team, 2016). We used mildly regularizing priors: Normal(0, 10) for

intercepts, Normal(0, 10) for diet treatment parameters, and Cauchy(0, 1) for variances.

Priors for covariance matrices were set to 1 for regularization, concentration, shape,

and scale (i.e., decov(1, 1, 1, 1)). Models were sampled for 20,000 iterations, with

10,000 discarded for burnin. Adequate sampling was assessed by $\hat{R}$, and models were

compared using leave-one-out (loo) cross-validation and loo model weights as

described above.

Heritability in single diets

To estimate variance components, heritability, and phenotypic correlations, we

used MCMCglmm (v. 2.24; Hadfield, 2010), a Bayesian inference package for R (v. 3.5;

R Core Team, 2016) to fit variations of the 'animal model' (Kruuk, 2004; Wilson *et al.*,

2010; Ingleby *et al.*, 2013). Using the animal model we first inferred, in each diet

separately, the additive genetic effect (V_A) from pedigree information where the

identities of sire and dam are included in the random effect

307   **y** ~ β + **Zu**

where **y** is the vector of phenotypes (lifespan, early fecundity, or total fecundity), β is the

mean phenotype, **Z** includes information about sire and dam identity (the "animal" term),

and **u** is the parameter vector that is estimated for the random effect, distributed as

Normal(0, $σ^2$). We used weakly informative priors: $V = 1$, $v = 0.002$, although analyses

were robust when more informative priors were tested. The models were sampled for 2

x 10^6 iterations, discarding the first 1.5 x 10^4 iterations as burnin, with a thinning interval

of 50 iterations. These sampling parameters yielded effective sample size >10,000. We

evaluated the models for convergence and autocorrelation of the samples, ensuring

autocorrelation of <0.1 for the first lag, and absence of trend in the trace plots of the

mean β (i.e., intercept term), additive variance V_A, and residual V_R (the later two

denoted in the output as 'animal' and 'units', respectively).

Genetic correlations between diets

A single phenotype that is measured in multiple environments (here diets) may

be regarded as different phenotypes in those environments (Falconer, 1952; Lynch and

Walsh, 1998)(Falconer 1952, Lynch and Walsh 1998). Based on this premise, we

computed genetic correlations for phenotype-diet combinations by fitting a Bayesian

multivariate animal model:

$$\mathbf{y} = \mathbf{X\beta} + \mathbf{Zu}$$

where **y** is a matrix of phenotypes (C, DR, HS), **β** is a vector of trait (**X**) means (with no

intercept term), **Z** includes information about sire and dam identity (the "animal" term),

and **u** is the parameter vector that is estimated for the random effect. Among the

random effects, covariance was free to vary (i.e., 'random = ~us(trait):animal') with one

restricting the variance to zero (i.e., 'random = ~idh(trait):animal'). If family genetic

effects remain the same or change in proportion between diets, then the genetic

correlation of family members between diets is 1. Genetic correlations that are credibly

smaller than 1 indicate a significant genotype x diet interaction. We therefore compared

the model above with a model in which the correlation between diets is constrained to

be 1 to test if our estimated genetic correlations were significantly different from 1. In

bivariate models, we tested several slightly stronger variations of priors (V = diag(3), v =

1.002). Model were run for 6.5×10^6 iterations, with 5×10^4 discarded as burning, with

thinning every 500 iterations. Twelve chains were run in parallel, yielding >10,000

samples total.

Results

Performance of image analysis to estimate the number of eggs

Studies measuring fecundity in *D. melanogaster* typically count the number of eggs laid by females by counting eggs on the food surface visually using a dissecting microscope, a method that requires a significant time investment. (Nouhaud *et al.*, 2018) has also developed an image-based method for counting *D. melanogaster* eggs but it relies on imaging the surface of the food and our egg densities would be very high with 24 females per vial, making it infeasible for this experiment. We employed a high throughput method for obtaining egg counts by developing a predictive model to obtain estimated egg counts from images of eggs that were separated from the food surface and filtered onto a dark fabric disc by taking advantage of the simple relationship between the number of eggs present and the amount of white area on a thresholded image of a disc (Fig. 2). We used a set of hand counted images to optimize the model and show that it performs well.

Figure 2: Performance of the model predicting egg counts from images. **a.** A representative raw image of eggs separated from fly media and filtered onto black filter fabric. **b.** The same image thresholded at the optimal threshold level identified by the model. **c.** The performance of the predictive model. The square root of image area is on the x-axis and the egg count is on the y-axis. True, hand-counted values are shown in red and the predicted values from the model are shown in blue. Pairs of values (hand-counted and predicted) for individual images are connected by a red line.

a. **b.**

Figure 2: Performance of the model predicting egg counts from images. **a.** A

representative raw image of eggs separated from fly media and filtered onto black filter

fabric. **b.** The same image thresholded at the optimal threshold level identified by the

model. **c.** The performance of the predictive model. The square root of image area is on

the x-axis and the egg count is on the y-axis. True, hand-counted values are shown in

red and the predicted values from the model are shown in blue. Pairs of values (hand-

counted and predicted) for individual images are connected by a red line.

We tested several parameters to produce an optimal model. First, by using

different proportions of our total dataset of hand counted images, we showed our hand

counted set was sufficiently large to train the predictive model and achieve a high

predictive ability. Using only 50% of the hand counted data, we achieved similar error

rates in the resulting model, demonstrating our dataset was sufficiently large to saturate

the model. We also tested a range of threshold values to identify the value that would

remove the most background noise without obscuring actual eggs. To assess the

performance of our model across different parameters, we employed cross validation to

ensure we tested the ability of the model to predict egg counts for new data (data not

used to fit the model). For each iteration, 90% of the data was used to train the model

and we assessed the ability of this model to predict the values of the remaining 10% of

the data (see Methods). Our best performing model was able to predict new data well,

with a mean absolute error of ~177 eggs and a high correlation ($r = 0.88$) between the

predicted egg count and the manual egg counts (Fig. 2c). We note that with a 24 hour

egg laying period for a vial with several females, this error rate represents a relatively

small number of eggs.

After identifying the optimal threshold value, we fit a model using our full dataset

and found similar error rates when comparing the predicted values from this model to

the hand counted images ($r = 0.88$, mean absolute error = 128 eggs; Fig. 2c). We were

then able to use this model to predict egg counts for our entire dataset of 3,768 images.

 Phenotypic response to dietary treatment

 Lifespan

We used a split family design, splitting offspring from families into three different

diets. Relative to the control diet, median survival was 24% lower on high sugar (HS: 48

days vs. C: 63 days) and 8% higher on DR diet (68 days) (Fig. 3). Lifespan trajectories

began to diverge early, from about 25 days post oviposition and remained diverged until

under 10% survivorship (Fig. 3). Individual sire family responses to diet are presented in

Fig. S2.

We compared six nested models of survival, incorporating diet treatment and

various aspects of relatedness and compared these models using leave-one-out model

weighting (Table 1). The model including sire, dam nested in sire, diet, and a sire by diet

interaction has the overwhelming model support (>95% of model weight), indicating a

significant effect of diet, the presence of significant genetic variance for survival within

the diet treatments, and significant genetic variation for the response to diet.

Specifically, the results of this favored model show that, compared to the C diet group,

individuals in the DR group were credibly less likely to die (99% credible interval = 0.01 -

0.11) and individuals in the HS treatment group were credibly more likely to die (99%

credible interval = -0.22 - -0.13).

Table 1. Comparison of models of lifespan, early fecundity, and total fecundity across

diet treatments. Models 1 and 2 do not include any aspects of pedigree, and models 3-6

use a hierarchical model. Terms in parentheses are random effects. All random effects

were "intercept" terms, allowing separate intercepts for each level of sire ID, dam ID

nested in sire ID, or sire ID nested in diet treatment. Model 1 includes only an intercept

term, allowing comparison to model 2. In all cases, model 2 was vastly superior to

model 1.

	Model	Lifespan	Early Fecundity	Total Fecundity
1	~ Intercept	3.5%	3.1%	1.1%
2	~ Diet	0%	10.3%	0.0%
3	~ Diet + (Sire ID)	0%	0.0%	2.1%
4	~ Diet + (Dam ID by Sire ID)	1.1%	86.6%	19.4%
5	~ Diet + (Sire ID) + (Dam ID by Sire ID)	0.2%	0.0%	0.0%
6	~ Diet + (Sire ID) + (Dam ID by Sire ID) + (Sire ID by Diet)	95.1%	0.0%	77.4%

Figure 3: Phenotypic response to diet treatment. The different colors represent different

diet treatments (grey = C, orange = DR, blue = HS). **a.** Fecundity per female sampled

over a 24 hour period once per week over lifespan. A smooth (loess) fit is shown for

each treatment. Individual lines represent each replicate vial. **b.** Survival probability

versus age in each of the three diet treatments. Median survival (50% survival

probability) is shown with the dotted line for each diet. Censored events are marked with

a + sign on each survivorship curve. **c.** The relationship between early life fecundity (24

hour sample period closest to 5 days post-eclosion) and median lifespan for each vial.

The average values for both within each treatment are shown by the large point outlined

in black. Error bars are the mean +/- 1 standard error. Note that the error bars all fit

within the size of the point.

427
Fecundity

Fecundity was estimated once per week in the same treatment setups from

which lifespan was recorded in order to compare the two phenotypes in the same set of

flies. As expected, in all treatments, fecundity declined with age. We used a Bayesian

model comparison approach to analyze the effects of diet and family on total fecundity

and early fecundity (Table 1). This analysis revealed a strong effect of diet for both total

fecundity and early fecundity, as models including diet were strongly preferred over a

model without diet. For both measures of fecundity, the highest values were in the C

diet and the lowest in the HS diet. There was a credible effect of family for both

fecundity measures, indicating significant genetic variation for fecundity. For early

fecundity, the model including only dam nested within sire was favored heavily. For total

fecundity, the model that included sire, dam nested in sire, and the interaction between

sire and diet was favored over the other models, indicating genetic variation for the

response to diet in addition to genetic variation for fecundity.

Estimates of Quantitative Genetic Parameters

We have a high sample size for lifespan with data for up to 48 individuals spread

across two vials for each dam family in each diet. While these same individuals

contributed to our measurements of weekly fecundity, we have a single measurement

for each vial of females at any given time point and do not have measurements per

each individual, as measuring fecundity for each individual female was not feasible for

this study. Thus, for estimating quantitative genetic parameters, our replication for

fecundity is low (2 per family per diet). The results of the above models reflect the

expected uncertainty from this low replication with wide posterior distributions. Thus,

while the models we fit above demonstrate the presence of significant genetic variation,

our estimates of heritability values have high uncertainty and wide credible intervals.

Therefore, we focus on discussing our results for lifespan below and results for

fecundity can be found in Fig. S3.

Heritability of lifespan within diets

We estimated the heritabilities of our phenotypes separately in each diet

treatment using a Bayesian approach (see Methods). Our heritability estimates for

lifespan are moderately high, ranging from 0.31 to 0.47 and do not differ substantially

across diets with all three credible intervals overlapping (Figure 4a; heritability (h^2) and

HPDI (highest posterior density interval) = C: h^2 0.47 HPDI 0.34 – 0.61; HS: h^2 0.37,

HPDI 0.25 – 0.50; DR: h^2 0.31, HPDI 0.21 – 0.43).

Correlation of lifespan across dietary treatments

To determine if there is substantial genetic variation for the response to the

environment (GEI), we fit a multivariate form of the animal model, which considers

465 lifespan within each diet as a separate phenotype (Falconer, 1952; Lynch and Walsh,

466 1998; Wilson *et al.*, 2010). When the genetic correlation between environments is less

467 than one, it indicates significant genetically based differences in the response to the

468 environment, or a significant GEI. In all pairwise diet comparisons, our estimates of the

469 genetic correlations are credibly less than one, indicating substantial genetically based

470 variation in the response to diet (genetic correlation (r_g) and HPDI = DR vs. C: r_g 0.48

471 HPDI 0.34 – 0.61; HS vs. C: r_g 0.29 HPDI 0.21 – 0.43; HS vs. DR: r_g 0.34 HPDI 0.25 –

472 0.50 (Fig. 4b). This variability is also apparent when examining the variation among sire

473 families in their response to diet for both lifespan and total fecundity (Fig. 4c & d). In

474 addition, the model comparisons we employed for survival and total fecundity both

475 favored models including a sire by diet interaction, which is consistent with the presence

476 of a significant GEI.

Figure 4: **a.** Density plots of the estimated posterior probabilities for lifespan heritability

within each diet. **b.** Density plots of the estimated posterior probabilities for the genetic

correlation between lifespan in pairs of diets. **c.** Sire by environment interaction plot for

median lifespan showing the median lifespan for each sire family in each of the three

diets. Each line represents one sire family. **d.** Sire by environment interaction plot for

total egg count per female (summed across weekly measures) showing the median

lifespan for each sire family in each of the three diets. Each line represents one sire

family.

Discussion

In this study, we used an admixed multiparent population of *D. melanogaster* to

characterize the quantitative genetics of fitness components in different nutritional

conditions. We characterized the effect of three different diets on survival and fecundity,

measured over the course of the entire lifetime of the flies. Using a quantitative genetic

breeding design, we show that these traits harbor substantial genetic variation both

within diets and in the response to diet. This result has important implications for our

understanding of the evolution of these traits in wild populations and for strategies to

uncover the genetic basis of these traits, which we discuss below.

Evolutionary potential of the coordination between nutrition and fitness components

We characterized the effects of a restricted diet in the form of a low protein diet

(DR) and of a calorie rich diet in the form of a high sugar diet (HS). Dietary restriction

has been studied extensively in *D. melanogaster* and other metazoans (e.g.,

(Chippindale *et al.*, 1993a; Good and Tatar, 2001; Vrtílek and Reichard, 2014);

reviewed in (Sohal and Weindruch, 1996; Kenyon, 2005; Kirkwood and Shanley, 2005;

Piper *et al.*, 2011; Tatar, 2011; Tatar *et al.*, 2014)). A general finding of these studies is

that reduced nutrient intake without starvation results in lifespan extension, and that this

effect is often coupled with reduced reproductive output, though there are exceptions to

this pattern (e.g., (Kaitala, 1991; Weindruch *et al.*, 1995; Kirk *et al.*, 2001; Stelzer, 2001;

Cooper *et al.*, 2004); see Ng'oma *et al.*, (2017)). Even within species that tend to show

this effect such as *D. melanogaster*, there is variability in this pattern among different

genotypes. For example, (Stanley *et al.*, 2017) found variation among inbred lines for

the amount of lifespan extension, with some lines showing little to no extension in a DR

diet, and a few showing reduced lifespan in a DR diet, a result that has also been

shown in mice (Rikke *et al.*, 2003). The effects of a high sugar diet have not been

studied as extensively, though recently some studies have shown flies reared on high

sugar diets as larvae show obesity-like phenotypes (Musselman *et al.*, 2011; Rovenko

et al., 2015), and adults reared on high sugar diets have elevated triglyceride levels,

reduced lifespan, and reduced fecundity (Skorupa *et al.*, 2008), a pattern that is

consistent with our findings.

These types of phenotypic patterns in different diet conditions have been

interpreted in different ways. The widespread observation of lifespan extension

observed in response to dietary restriction in many taxa has led to competing

hypotheses for this pattern. It is often argued that this pattern is the result of natural

selection acting in environments with fluctuating resources where it should be adaptive

to increase allocation of nutrient resources to the soma when nutrients are limiting,

however it is also possible it is the result of an unavoidable physiological constraint

(Holliday, 1989; Kirkwood and Shanley, 2005). Additionally, studies employing high

sugar and high fat diets often view the results of those studies in the context of human

populations where it is hypothesized the negative health consequences of a

Westernized diet result from a mismatch between that diet and the diet humans adapted

to in the past over the majority of their evolutionary history (Neel, 1962; Wells, 2009),

though this hypothesis has also been challenged (Speakman, 2008). In general,

assessing the potential for phenotypic effects of different diets to be adaptive patterns or

to be highly constrained physiological responses requires an understanding of the

underlying genetic variation for the response to diet in order to understand if this

response has the potential to evolve.

Here, we showed there is heritable variation for both phenotypes within diets,

and for the response the phenotypes to diet (Table 1; Fig. 4a,b). While many studies

have demonstrated GEI for different traits (Reed *et al.*, 2010; Qi *et al.*, 2012; Ingleby *et*

al., 2013), few studies have estimated GEI specifically for life history traits in response

to diet, and not all previous studies of diet have found significant GEI. For example,

(King *et al.*, 2011) estimated quantitative genetic parameters for allocation to flight

capability and to reproduction in the sand cricket, *Gryllus firmus.* These authors found

no GEI for these traits, with all genotypes allocating more to reproduction when food

levels increase, suggesting important evolutionary constraints can exist for the response

to diet and that significant GEI is not a foregone conclusion. Our results show that this

constraint does not exist for this population of *D. melanogaster* and that the response to

diet will readily evolve in response to selection. The founder lines of the DSPR have a

global distribution and were derived from wild-caught flies and thus, our population

encompasses some of the genetic variation that exists among natural populations of *D.*

melanogaster. It is plausible, therefore, that the diet dependent effects observed in this

study and others represent adaptive patterns that have evolved. However, a major

challenge that remains in interpreting these patterns is the extent to which laboratory

diets represent the nutritional conditions experienced by flies in the wild.

Prospects for multiparent population approaches

552   One of the most fundamental goals in biology is to understand the genetic basis

of complex phenotypes. This goal has proved to be quite challenging. For most traits,

some of which have been the focus of study for many years, the causative genetic

variants that have been discovered explain only a small percentage of the trait

heritability (for reviews see McCarthy *et al.*, 2008; Manolio *et al.*, 2009; Rockman, 2012;

Visscher *et al.*, 2012). For traits that are expected to be highly polygenic, such as

lifespan and fecundity, estimating quantitative genetic parameters as we have done in

this study, can be as informative about the genetic basis of a trait and its evolutionary

potential as studies aimed at uncovering specific genetic variants.

561   In recent years, multiparent populations, in which multiple inbred founder lines

are crossed for a number of generations to create a recombinant mapping population,

have increasingly been employed in studies aiming to identify the causative genetic

variants underlying complex traits (e.g. (Kover *et al.*, 2009; McMullen *et al.*, 2009;

Huang *et al.*, 2011; Aylor *et al.*, 2011; Threadgill and Churchill, 2012; King, Macdonald,

et al., 2012; King, Merkes, *et al.*, 2012; Cubillos *et al.*, 2013); reviewed in (Long *et al.*,

2014; de Koning and McIntyre, 2017). One reason for employing a mapping population

consisting of multiple parent lines is the inclusion of additional genetic variation when

compared to traditional two-line QTL mapping panels. Even with multiple parent lines

contributing, a potential criticism of these kinds of approaches when compared to

mapping approaches that sample from natural populations is that they contain limited

amounts of genetic variation. The DSPR population used in this study was started from

8 inbred parent lines. Therefore, while every individual in our admixed population has a

unique mosaic genome, at any given position, there are only 8 possible haplotypes.

Despite this simplicity, we find moderate heritabilities for lifespan in all diets (ranging

from ~0.3 - 0.5) that are comparable to estimates from natural populations. A meta-

analysis of heritability estimates from natural populations found life history traits typically

have values around 0.25 and values higher than 0.3 are not infrequent (Rose and

Charlesworth, 1981; Mousseau and Roff, 1987; Rikke *et al.*, 2010). We also found

strong evidence for a genotype by diet interaction for lifespan, indicating there is also

segregating genetic variation for the response to diet. While our evidence of a genotype

by diet interaction is not as strong for fecundity, our model comparison for total fecundity

supports a model including this interaction.

In addition to their utility for QTL mapping, multiparent populations have also

been used in evolve and resequence approaches (e.g. (Turner and Miller, 2012; Burke,

King, *et al.*, 2014; Burke, Liti, *et al.*, 2014) to identify the underlying genetic basis of

complex traits. These studies impose selection and track the resulting changes in

phenotype and genotype. This approach is potentially very powerful because 1) entire

haplotypes can be tracked over evolutionary time, 2) the simplified genetic architecture

with only a few potential haplotypes at any given location may lead to a more repeatable

response to selection increasing the likelihood of identifying causative loci, and 3) it may

allow for validation of QTL identified in mapping studies using these same populations.

A potential downside to such an approach is if the relatively simple genetic architecture

of such a population results in limited genetic variation for the trait of interest, thereby

constraining evolution. Our study shows there is substantial genetic variation for our

focal phenotypes in our admixed DSPR population, which establishes this approach as

a viable strategy for investigating both how a focal trait evolves and the underlying

genetic basis. In general, MPPs have great potential to advance our understanding of

the genetic basis of complex traits via these kinds of multi-pronged approaches.

Acknowledgments

Larry Cabral provided with valuable ideas through personal communication to establish

our fecundity protocol. Stuart Macdonald provided the DSPR RILs we used to start our

experimental colony. Elizabeth Lopresti, Michael Reed, Osvaldo Enriquez, Anna

Perinchery and Kyla Winford helped with fly husbandry, time-intensive experimental set-

ups, collection and entry of data. This work was supported by NIH grant R01 GM117135

to Elizabeth G. King, The University of Missouri, and a University of Missouri Research

Board Grant.

Conflict of interest

The authors declare no conflict of interest.

Data availability

Our raw data, including data on lifespan, fecundity, and all egg image files, have been

uploaded to Zenodo and may be retrieved at http://doi.org/10.5281/zenodo.1285237. All

scripts to reproduce all of our analyses are available at GitHub:

https://github.com/EGKingLab/h2lifespan. Supplementary information is available at

https://www.nature.com/hdy/

References

Abramoff MD, Magalhães PJ, Ram SJ (2004). Image processing with ImageJ.

Biophotonics international.

Aylor DL, Valdar W, Foulds-Mathes W, Buus RJ, Verdugo RA, Baric RS, *et al.* (2011).

Genetic analysis of complex traits in the emerging Collaborative Cross. *Genome Res*

21: 1213–1222.

Barnes AI, Boone JM, Jacobson J, Partridge L, Chapman T (2006). No extension of

lifespan by ablation of germ line in *Drosophila*. *Proc Biol Sci* **273**: 939–947.

Bradski G (2000). *The OpenCV Library*.

Brčić-Kostić K (2010). Quantitative genetics and evolution. *Periodicum biologorum* **112**:

395–402.

Brooks SP, Gelman A (1998). General methods for monitoring convergence of iterative

simulations. *Journal of Computational and Graphical Statistics* **7**: 434–455.

Burke MK, King EG, Shahrestani P, Rose MR, Long AD (2014). Genome-wide

association study of extreme longevity in *Drosophila melanogaster*. *Genome Biology*

and Evolution **6**: 1–11.

Burke MK, Liti G, Long AD (2014). Standing genetic variation drives repeatable

experimental evolution in outcrossing populations of *Saccharomyces cerevisiae*. *Mol*

Biol Evol **31**: 3228–3239.

Bürkner P-C (2017). brms: an r package for bayesian multilevel models using stan.

Journal of Statistical Software, **80**: 1–28

Cabral, L. [Rose Lab Data]. (2012, September 21). A method for gathering fecundity

data for populations of Drosophila melanogaster. [Fecundity: Animated Introduction].

Retrieved from https://www.youtube.com/watch?v=xrEBNB8bUwo&sns=em

Cabral, L., Rutledge, GA. [Rose Lab Data]. (2012, September 21). A method for

gathering fecundity data for populations of Drosophila melanogaster. [Fecundity: Live

Demonstration]. Retrieved from

https://www.youtube.com/watch?v=uqFZ4ZTC6ZY&sns=em

Carnes, MU, Campbell T, Huang W, Butler DG, Carbone MA, Duncan LH, *et al.* (2015).

The genomic basis of postponed senescence in *Drosophila melanogaster*. *PLOS ONE*

10: e0138569.

Carpenter B, Gelman A, Hoffman M, Lee D, Goodrich B, Betancourt M, *et al.* (2017).

Stan: a probabilistic programming language. *Journal of Statistical Software*, **76**: 1–32.

Chippindale A, Leroi AM, Kim S, Rose M (1993a). Phenotypic plasticity and selection in

Drosophila life-history evolution. I. Nutrition and the cost of reproduction. *J Evolution*

Biol **6**: 171–193.

Chippindale AK, Leroi AM, Kim SB, Rose MR (1993b). Phenotypic plasticity and

selection in *Drosophila* life-history evolution. I. Nutrition and the cost of reproduction.

Journal of Evolutionary Biology **6**: 171–193.

Cooper TM, Mockett RJ, Sohal BH, Sohal RS, Orr WC (2004). Effect of caloric

restriction on life span of the housefly, *Musca domestica*. *FASEB J* **18**: 1591–1593.

Cubillos FA, Parts L, Salinas F, Bergström A, Scovacricchi E, Zia A, *et al.* (2013). High-

resolution mapping of complex traits with a four-parent advanced intercross yeast

population. *Genetics* **195**: 1141–1155.

Falconer DS (1952). The problem of environment and selection. *The American*

Naturalist **86**: 293–298.

Fisher RA (1930). *The genetical theory of natural selection.* Clarendon Press: Oxford.

Fisher JAD, Frank KT, Leggett WC (2010). Breaking Bergmann's rule: truncation of

Northwest Atlantic marine fish body sizes. *Ecology* **91**: 2499–2505.

Flatt T, Min K-J, D'Alterio C, Villa-Cuesta E, Cumbers J, Lehmann R, *et al.* (2008).

Drosophila germ-line modulation of insulin signaling and lifespan. *P Natl Acad Sci USA*

105: 6368–6373.

Gagneur J, Stegle O, Zhu C, Jakob P, Tekkedil MM, Aiyar RS, *et al.* (2013). Genotype-

environment interactions reveal causal pathways that mediate genetic effects on

phenotype. *PLoS Genet* **9**: e1003803.

Gelman A, Hill J (2007). *Data analysis using regression and multilevel/hierarchical*

models. Cambridge University Press: Cambridge ; New York.

Gelman A, Rubin D (1992). A single series from the Gibbs sampler provides a false

sense of security. *Bayesian Statistics* **4**: 625–631.

Giannakou ME, Goss M, Partridge L (2008). Role of dFOXO in lifespan extension by

dietary restriction in *Drosophila melanogaster*: not required, but its activity modulates

the response. *Aging Cell* **7**: 187–198.

Good TP, Tatar M (2001). Age-specific mortality and reproduction respond to adult

dietary restriction in *Drosophila melanogaster*. *J Insect Physiol* **47**: 1467–1473.

Grandison RC, Wong R, Bass TM, Partridge L, Piper MDW (2009). Effect of a

standardised dietary restriction protocol on multiple laboratory strains of drosophila

melanogaster (H Tanimoto, Ed.). *PLOS ONE* **4**: e4067.

Hadfield JD (2010). MCMC methods for multi-response generalized linear mixed

models: the MCMCglmm R package. *Journal of Statistical Software* **33**: 1–22.

Holliday R (1989). Food, reproduction and longevity: is the extended lifespan of calorie-

restricted animals an evolutionary adaptation? *Bioessays* **10**: 125–127.

Hsin H, Kenyon C (1999). Signals from the reproductive system regulate the lifespan of

C. elegans. *Nature* **399**: 362–366.

Huang X, Paulo M-J, Boer M, Effgen S, Keizer P, Koornneef M, *et al.* (2011). Analysis

of natural allelic variation in *Arabidopsis* using a multiparent recombinant inbred line

population. *PNAS* **108**: 4488–4493.

Hughes KA, Reynolds RM (2005). Evolutionary and mechanistic theories of aging. *Annu*

Rev Entomol **50**: 421–445.

Hunt J, Bussière LF, Jennions MD, Brooks R (2004). What is genetic quality? *Trends in*

Ecology & Evolution **19**: 329–333.

Ingleby FC, Hosken DJ, Flowers K, Hawkes MF, Lane SM, Rapkin J, *et al.* (2013).

Genotype-by-environment interactions for cuticular hydrocarbon expression in

Drosophila simulans. *J Evol Biol* **26**: 94–107.

Kaitala A (1991). Phenotypic plasticity in reproductive behavior of waterstriders: trade-

offs between reproduction and longevity during food stress. *Funct Ecol* **5**: 12–18.

Kaplan EL, Meier P (1958). Nonparametric estimation from incomplete observations.

Journal of the American Statistical Association **53**: 457–481.

Kenyon CJ (2005). The plasticity of aging: insights from long-lived mutants. *Cell* **120**:

449–460.

King EG, Macdonald SJ, Long AD (2012). Properties and power of the *Drosophila*

Synthetic Population Resource for the routine dissection of complex traits. *Genetics*

191: 935–949.

King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, *et al.* (2012). Genetic

dissection of a model complex trait using the *Drosophila* Synthetic Population Resource.

Genome Res **22**: 1558–1566.

King EG, Roff DA, Fairbairn DJ (2011). The evolutionary genetics of acquisition and

allocation in the wing dimorphic cricket, *Gryllus firmus*. *Evolution* **65**: 2273–2285.

Kirk KL, Turner TL, Kirkwood TBL, Reed LK, Miller PM, Shanley DP, *et al.* (2001).

Dietary restriction and aging: comparative tests of evolutionary hypotheses. *Journal of*

Gerontology: BIOLOGICAL SCIENCES **56A**: B123-129.

Kirkwood TBL, Shanley DP (2005). Food restriction, evolution and ageing. *Mech Ageing*

Dev **126**: 1011–1016.

de Koning D-J, McIntyre LM (2017). Back to the future: multiparent populations provide

the key to unlocking the genetic basis of complex traits. *Genetics* **206**: 527–529.

Kover PX, Valdar W, Trakalo J, Scarcelli N, Ehrenreich IM, Purugganan MD, *et al.*

(2009). A multiparent advanced generation inter-cross to fine-map quantitative traits in

Arabidopsis thaliana. *PLoS Genet* **5**: e1000551.

Kruuk LEB (2004). Estimating genetic parameters in natural populations using the

'animal model'. *Philos Trans R Soc Lond B Biol Sci* **359**: 873–890.

Lande R, Arnold SJ (1983). The measurement of selection on correlated characters.

Evolution **37**: 1210–1226.

Long AD, Macdonald SJ, King EG (2014). Dissecting complex traits using the

Drosophila Synthetic Population Resource. *Trends in Genetics* **30**: 488–495.

Lynch M, Walsh B (1998). *Genetics and analysis of quantitative traits*. Sinauer

Associates, Inc.: Sunderland, MA.

Mair W, Dillin A (2008). Aging and survival: the genetics of life span extension by dietary

restriction. *Annu Rev Biochem* **77**: 727–754.

Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, *et al.* (2009).

Finding the missing heritability of complex diseases. *Nature* **461**: 747–753.

McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JPA, *et al.*

(2008). Genome-wide association studies for complex traits: consensus, uncertainty

and challenges. *Nature Reviews Genetics* **9**: 356–369.

McMullen MD, Kresovich S, Villeda HS, Bradbury P, Li H, Sun Q, *et al.* (2009). Genetic

properties of the maize nested association mapping population. *Science* **325**: 737–740.

Min K-J, Yamamoto R, Buch S, Pankratz M, Tatar M (2008). *Drosophila* lifespan control

by dietary restriction independent of insulin-like signaling. *Aging Cell* **7**: 199–206.

Mousseau TA, Roff DA (1987). Natural selection and the heritability of fitness

components. *Heredity* **59**: 181–197.

Musselman LP, Fink JL, Narzinski K, Ramachandran PV, Hathiramani SS, Cagan RL,

et al. (2011). A high-sugar diet produces obesity and insulin resistance in wild-type

Drosophila. *Disease Models & Mechanisms* **4**: 842–849.

Nässel DR, Liu Y, Luo J (2015). Insulin/IGF signaling and its regulation in *Drosophila*.

Gen Comp Endocrinol **221**: 255–266.

Neel J (1962). Diabetes mellitus: A 'thrifty' genotype rendered detrimental by 'progress'?

Am J Hum Genet **14**: 353–362.

Ng'oma E, Perinchery AM, King EG (2017). How to get the most bang for your buck: the

evolution and physiology of nutrition-dependent resource allocation strategies. *Proc Biol*

Sci **284**.

van Noordwijk AJ, de Jong G (1986). Acquisition and allocation of resources: their

influence on variation in life history tactics. *The American Naturalist* **128**: 137–142.

Nouhaud P, Mallard F, Poupardin R, Barghi N, Schlötterer C (2018). High-throughput

fecundity measurements in *Drosophila*. *Scientific Reports* **8**: 4469.

Piper MDW, Partridge L, Raubenheimer D, Simpson SJ (2011). Dietary restriction and

aging: a unifying perspective. *Cell Metab* **14**: 154–160.

Qi Q, Chu AY, Kang JH, Jensen MK, Curhan GC, Pasquale LR, *et al.* (2012). Sugar-

sweetened beverages and genetic risk of obesity. *New England Journal of Medicine*

367: 1387–1396.

R Core Team (2016). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing, Vienna, Austria.

Reed LK, Lee K, Zhang Z, Rashid L, Poe A, Hsieh B, *et al.* (2014). Systems genomics

of metabolic phenotypes in wild-type *Drosophila melanogaster*. *Genetics* **197**: 781–793.

Reed LK, Williams S, Springston M, Brown J, Freeman K, DesRoches CE, *et al.* (2010).

Genotype-by-diet interactions drive metabolic phenotype variation in *Drosophila*

melanogaster. *Genetics* **185**: 1009–1019.

Remolina SC, Chang PL, Leips J, Nuzhdin SV, Hughes KA (2012). Genomic basis of

aging and life-history evolution in *Drosophila melanogaster*. *Evolution* **66**: 3390–3403.

Rikke BA, Liao C-Y, McQueen MB, Nelson JF, Johnson TE (2010). Genetic dissection

of dietary restriction in mice supports the metabolic efficiency model of life extension.

Experimental Gerontology **45**: 691–701.

Rikke BA, Yerg JE, Battaglia ME, Nagy TR, Allison DB, Johnson TE (2003). Strain

variation in the response of body temperature to dietary restriction. *Mechanisms of*

Ageing and Development **124**: 663–678.

Rockman MV (2012). The QTN program and the alleles that matter for evolution: all

that's gold does not glitter. *Evolution* **66**: 1–17.

Roff DA (2006). *Evolutionary quantitative genetics*. In: Fox CW, Wolf JB (eds).

Evolutionary genetics: concepts and case studies. Oxford Univ. Press: New York. pp.

267–287.

Rose MR, Charlesworth B (1981). Genetics of life history in *Drosophila melanogaster*. I.

sib analysis of adult females. *Genetics* **97**: 173–186.

Rovenko BM, Perkhulyn NV, Gospodaryov DV, Sanz A, Lushchak OV, Lushchak VI

(2015). High consumption of fructose rather than glucose promotes a diet-induced

obese phenotype in *Drosophila melanogaster*. *Comp Biochem Physiol, Part A Mol*

Integr Physiol **180**: 75–85.

Simmons F, Bradley T (1997). An analysis of resource allocation in response to dietary

yeast in *Drosophila melanogaster*. *J Insect Physiol* **43**: 779–788.

Skorupa DA, Dervisefendic A, Zwiener J, Pletcher SD (2008). Dietary composition

specifies consumption, obesity, and lifespan in *Drosophila melanogaster*. *Aging Cell* **7**:

478–490.

Sohal R, Weindruch R (1996). Oxidative stress, caloric restriction, and aging. *Science*

(New York, NY).

Speakman JR (2008). Thrifty genes for obesity, an attractive but flawed idea, and an

alternative perspective: the 'drifty gene' hypothesis. *Int J Obes Relat Metab Disord* **32**:

1611–1617.

Sprott RL (1997). Diet and calorie restriction. *Experimental Gerontology* **32**: 205–214.

Stan Development Team (2016). *rstanarm: Bayesian applied regression modeling via*

Stan. R package version 2.13.1.

Stanley PD, Ng'oma E, O'Day S, King EG (2017). Genetic dissection of nutrition-

induced plasticity in insulin/insulin-like growth factor signaling and median life span in a

Drosophila multiparent population. *Genetics* **206**: 587–602.

Stelzer C (2001). Resource limitation and reproductive effort in a planktonic rotifer.

Ecology **82**: 2521–2533.

Tatar M (2011). The plate half-full: status of research on the mechanisms of dietary

restriction in *Drosophila melanogaster*. *EXG* **46**: 363–368.

Tatar M, Post S, Yu K (2014). Nutrient control of *Drosophila longevity*. *Trends Endocrin*

Met: 1–9.

Teleman AA (2010). Molecular mechanisms of metabolic regulation by insulin in

Drosophila. *Biochem J* **425**: 13–26.

Therneau TM (2015). *A package for survival analysis in S*, version 2.38.

https://CRAN.R-project.org/package=survival.

Therneau TM, Grambsch PM (2000). *Modeling survival data: Extending the cox model*.

Springer: New York.

Threadgill DW, Churchill GA (2012). Ten years of the Collaborative Cross. *G3: Genes,*

Genomes, Genetics **2**: 153–156.

Turner TL, Miller PM (2012). Investigating natural variation in *Drosophila* courtship song

by the evolve and resequence approach. *Genetics* **191**: 633–642.

Vehtari A, Gelman A, Gabry J (2017). Practical Bayesian model evaluation using leave-

one-out cross-validation and WAIC. *Stat Comput* **27**: 1413–1432.

Visscher PM, Brown MA, McCarthy MI, Yang J (2012). Five years of GWAS discovery.

The American Journal of Human Genetics **90**: 7–24.

Vrtílek M, Reichard M (2014). Highly plastic resource allocation to growth and

826 reproduction in females of an African annual fish. *Ecol Freshw Fish* **24**: 616–628.

827 Warren IA, Gotoh H, Dworkin IM, Emlen DJ, Lavine LC (2013). A general mechanism

828 for conditional expression of exaggerated sexually-selected traits. *Bioessays* **35**: 889–

829 899.

830 Weindruch R, Marriott BM, Conway J, Knapka JJ, Lane MA, Cutler RG, *et al.* (1995).

831 Measures of body size and growth in rhesus and squirrel monkeys subjected to long-

832 term dietary restriction. *Am J Primatol* **35**: 207–228.

833 Wells JCK (2009). Thrift: a guide to thrifty genes, thrifty phenotypes and thrifty norms.

834 *Int J Obes (Lond)* **33**: 1331–1338.

835 Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, *et al.* (2010).

836 An ecologist's guide to the animal model. *Journal of Animal Ecology* **79**: 13–26.

837 Yao Y, Vehtari A, Simpson D, Gelman A (2018). Using stacking to average Bayesian

838 predictive distributions. *Bayesian Anal*.