The evolutionary potential of diet-dependent effects on lifespan and fecundity in a

multi-parental population of *Drosophila melanogaster*

Enoch Ng'oma[1*], Wilton Fidelis[1], Kevin M. Middleton[2] and Elizabeth G. King[1]

[1]Division of Biological Sciences, University of Missouri, Columbia, MO 65211

[2]Department of Pathology and Anatomical Sciences, University of Missouri School of

Medicine, Columbia, MO 65212

*Corresponding author:

Enoch Ng'oma

401 Tucker Hall

Division of Biological Sciences

University of Missouri, Columbia, MO 65211

Tel.: 573-882-8562

E-mail: ngomae@missouri.edu

Running title: Quantitative genetics of life history in an MPP

Word count: 6991

Abstract

The nutritional conditions experienced by a population play a major role in shaping trait

evolution in many taxa. Constraints exerted by nutrient limitation or nutrient imbalance

can influence the maximal value that fitness components such as reproduction and

lifespan attains, and organisms may shift how resources are allocated to different

structures and functions in response to changes in nutrition. Whether the phenotypic

changes associated with changes in nutrition represent an adaptive response is largely

unknown. Further, it is unclear whether the response of fitness components to diet even

has the potential to evolve in most systems. In this study, we use an admixed

multiparental population of *Drosophila melanogaster* reared in three different diet

conditions to estimate quantitative genetic parameters for lifespan and fecundity. We

find significant genetic variation for both traits in our population and show that lifespan

has moderate to high heritabilities within diets. Genetic correlations for lifespan between

diets were significantly less than one, demonstrating a strong genotype by diet

interaction. These findings demonstrate substantial standing genetic variation in our

population that is comparable to natural populations and highlights the potential for

adaptation to changing nutritional environments.

Key words: life history, heritability, genotype by environment interaction, *Drosophila*

melanogaster, multiparental population

Introduction

An organism's diet has a direct influence on nearly all phenotypes by determining the amount of nutrients and energy available to build structures and perform functions. In addition to this unavoidable direct effect, organisms may also alter how resources are allocated to different traits when experiencing different diets. In an environment where resource availability varies across space and/or time, and the optimal allocation of resources changes with diet, we expect to see the evolution of such phenotypically plastic resource allocation strategies.

In many non-human populations including nematode (Greer and Brunet, 2009), fruit fly (Chippindale *et al.*, 1993b), killifish (Vrtílek and Reichard, 2014), waterstrider (Kaitala, 1991) and mouse (Sprott, 1997), nutrient limitation without starvation, hereafter referred to as dietary restriction (DR), results in lifespan extension, and is often but not always accompanied by a reduction in reproductive output (Hsin and Kenyon, 1999; Barnes *et al.*, 2006; Flatt *et al.*, 2008). These diet-dependent changes have been hypothesized to result from an increased allocation of a limited nutritional resource to somatic maintenance (van Noordwijk and de Jong, 1986; Sohal and Weindruch, 1996; Simmons and Bradley, 1997; Hughes and Reynolds, 2005; Kirkwood, 1977; Shanely and Kirkwood, 2000), and decreased allocation to reproduction. Previous models predict this pattern is adaptive in environments with fluctuating resource availability (Neel, 1962; Wells, 2009; Fisher *et al.*, 2010; Speakman 2011; Warren *et al.*, 2013). However, the main physiological and evolutionary drivers of the DR effect and its underlying mechanisms are still not well-understood. Several studies taking a nutritional geometry approach, which measure phenotypes across serial dilutions of diet

components, have found that lifespan and reproduction are affected primarily by the

protein-carbohydrate ratio of the diet rather than the caloric content (e.g. Skorupa *et al*,

2008; Lee *et al*, 2008; Jensen *et al*, 2015). These studies demonstrate that calorie

limitation *per se*, is not driving these phenotypic patterns, though it may be that

individual diet components act as a cue that induces an optimal allocation of resources

in wild populations. In addition, recent meta-analyses quantifying the nutritional

composition of the various types of diets that are used in DR studies have shown that

DR has greater effect on reproduction and lifespan in model species than it has on non-

model species, suggesting the effect may be due to laboratory adaptation (Speakman,

2011; Nakagawa *et al*, 2012, Moatt *et al*, 2016). However, different hypotheses for the

evolution of the DR response have rarely been formally tested (but see Zajitschek *et al,*

2016; 2018), nor has the evolutionary potential of the focal traits been established in

most systems (Ng'oma *et al.*, 2017), which is an important first step.

The capacity for a given trait to evolve in a population depends on the proportion

of trait variation that is attributable to genetic variation and the intensity of selection

acting on it (Roff 2006; Brčić-Kostić, 2010). Selection can have a direct effect on trait

evolution if the trait directly affects fitness, or indirectly if the trait is correlated with a trait

under selection (Lande and Arnold, 1983; Hunt *et al.*, 2004). However, even when a

selective pressure is observed to act on the trait, the amount of evolutionary change

depends on the amount of genetic variation expressed for that trait (Fisher, 1930).

Quantitative genetic parameters, such as the heritability of a given trait, remain some of

the more informative parameters we can estimate in order to learn about the genetic

basis of a trait, to predict trait values in a population, and to predict the selection

response, even with the advances in genomics and molecular genetics in recent years.

Recent experimental evolution studies exposing fruit flies to DR have demonstrated that

lifespan and reproduction do respond to selection under different diet conditions,

although lifespan appears to be decoupled from fecundity (Zajitschek *et al,* 2016;

2018). In addition, studies in many species have frequently demonstrated genotype by

diet variation for complex traits, such as gene expression and fitness in yeast (Gagneur

et al., 2013), cuticular hydrocarbons in *D. simulans* (Ingleby *et al.*, 2013), and metabolic

phenotypes in *D. melanogaster* (Reed *et al.*, 2010, 2014; Qi *et al.*, 2012). These studies

suggest that genotype by environment interactions (GEI) are common and could be

important in the nutrition-dependent response of life history traits such as fecundity and

lifespan.

There has been a significant effort to understand the genetic basis for the dietary

restriction (DR) response with both molecular genetic approaches (Mair and Dillin,

2008; Grandison *et al.*, 2009; Teleman, 2010; Nässel *et al.*, 2015) and mapping

approaches (Rikke *et al.*, 2010; Stanley *et al.*, 2017). As is the case for most complex

traits, mapping approaches to identify the genetic loci determining how lifespan changes

with diet have failed to identify the specific contributors. Studies focused on mapping in

natural populations often do not identify the regions of the genome as QTL that harbor

candidate genes identified by molecular approaches (Remolina *et al.*, 2012; Burke,

King, *et al.*, 2014; Reed *et al.*, 2014; Carnes *et al.*, 2015) and there are several studies

in which gene expression patterns do not change in the direction one would expect from

these loss-of-function studies (Min *et al.*, 2008; Giannakou *et al.*, 2008; Stanley *et al.*,

2017). In light of these studies, a possible hypothesis is that the DR response is highly

polygenic and best understood by quantifying holistic measures of genetic variation,

such as the heritability and GEI. Additionally, it is possible that the limited genetic

variation included in QTL mapping approaches limits the ability to identify causative

variants that are segregating in natural populations. The majority of QTL studies in the

past have used only two genomes to create the mapping population. More recently,

multiparent populations (MPPs) have been gaining in popularity, where multiple

(typically at least 4, and often many more) genomes are used to seed the population,

thereby including increased genetic variability (Kover *et al.*, 2009; McMullen *et al.*, 2009;

Huang *et al.*, 2011; Aylor *et al.*, 2011; Threadgill and Churchill, 2012; King, *et al.*, 2012;

King, *et al.*, 2012; Cubillos *et al.*, 2013).

In this paper, we employ a half-sibling, split environment design using an

admixed multiparent population derived from an established mapping resource, the

Drosophila Synthetic Population Resource (DSPR), to estimate quantitative genetic

parameters for lifespan and fecundity in three different diets. Our diets include typical

DR and control diets used in *D. melanogaster* following Bass *et al* (2007), and a high

sugar diet that contains ~7 times the amount of sugar relative to our control diet. We are

able to estimate the evolutionary potential of how lifespan and fecundity respond to

these diets in order to establish, 1) the ability of these traits to evolve in general, and

provide specific estimates of quantitative genetic parameters, 2) demonstrate levels of

genetic variation that are comparable with wild derived populations, and 3) establish the

validity of the evolve and resequence approach using a synthetic population as a base

population. We expect to find extended lifespan and reduced fecundity on the DR diet

relative to the control and reduced lifespan and fecundity in the high sugar diet relative

to the control. More importantly, however, we expect to see substantial variablity among

families in the phenotypic patterns in response to our diet treatments, with estimates of

heritabilities within diets and genetic correlations between diets showing evidence for

substantial genetic variation.

Methods

Experimental setup

Generation of experimental population. We used an eight-way multi-parent

population, the DSPR, to characterize the phenotypic response of lifespan and

fecundity of mated female *D. melanogaster* to three nutritional environments and to

estimate the heritabilities of these traits. The development, and genetic properties of the

DSPR are described in detail in King *et al.*, (2012a) and King *et al.*, (2012b). Briefly, the

DSPR is a set of recombinant inbred lines created from 8 inbred founder lines that were

crossed for 50 generations and then inbred via full sibling mating for 25 generations. To

generate our starting population, for each of 835 recombinant inbred lines (RILs) of the

B sub-population of the DSPR, five young females that had mated intra-line in vials,

were randomly selected and distributed to each of six cages (20.3 cm x 20.3 cm x 20.3

cm). Eggs were collected in milk bottles (250 mL) twice on the next two successive

days, each after a 22-hour oviposition period on maintenance diet (Table S1). Eclosed

flies were released to six cages in populations >2,000 per cage and allowed to mate *en-*

masse for five generations. A polystyrene plate, 100 mm x 15 mm (cat. No. FB0875713

Fisher Scientific, USA) containing maintenance food was supplied to each cage three

times a week (Monday, Wednesday, Friday). A separate micro-plate (60 mm x 15 mm,

cat. no. FB0875713A, Fisher Scientific, USA) with moist cotton wool was supplied to

each cage to serve as a supplemental drinking source. Very thin slices of food with

visually estimated 50-90 eggs were transferred to 30 vials (25 mm x 95 mm,

Polystyrene Reload, cat. no. 32-109RL, Genesee Scientific, USA) for each cage for

each successive generation. Egg vials from each cage for each next generation were

distributed across all cages to ensure a genetically homogenous experimental

population. We maintained all populations on a three week cycle (21 days from

oviposition to egg collection) on a cornmeal-dextrose-yeast diet for 5 generations before

the half-sibling experiment was started.

Experimental design. We employed a split-environment, half-sibling breeding

design to determine the phenotypic effects of three dietary conditions and estimate

quantitative genetic parameters. We chose to use a dietary restriction and control diet

similar to those commonly used in studies of the lifespan effects of DR in flies (Bass et

al. 2007). We also employed a high sugar diet, with the same amount of yeast as the

control diet but with ~7 times the amount of sugar. The motivation for choosing this diet

was to provide a high calorie diet that has the potential to serve as a model of some

modern human diets (i.e. a higher calorie, higher carbohydrate diet than the diet

experienced for the majority of evolutionary history). While it would have been ideal to

measure a complete range of diets with many possible combinations of yeast and

sugar, the scale of this experiment limited the number of diets that could be examined.

We created a set of paternal half-sibling families by sequentially mating each of 28

males to three females (here on, sires and dams, respectively) on a common cornmeal-

dextrose-yeast (maintenance) diet (Table S1). Each dam was housed in its own vial and

each sire rotated among these three vials. In round one, to ensure each dam had a

mating opportunity early, the sire spent 2 days with each dam. Each sire was then

rotated among the three dams for two additional rounds lasting 3.5 days each

encounter. Two dams were kept waiting in media vials when the sire was matched with

the third dam. Whole families were restarted with new flies when a sire died during

crossing. In case of death of a waiting unmated dam, only that dam was replaced,

whereas if a mated dam died, a new dam family was restarted. Starting with 61 sires

and 183 dams, 28 half-sibling families and 77 maternal full-sibling families successfully

produced enough offspring to be included in the experiment (Fig. 1).

From each of the 77 dams, in most cases, we set up two replicate vials of 24

female and 6 male offspring in each of three diet treatments (described below). We

included males to provide mating opportunities to females throughout their lifetime but

chose a smaller number of males to reduce harassment of females within the vials. As

expected, even successful sire and dam pairs did not always produce enough offspring

for a full complement. Thus, some vials contained fewer than 24 females though never

below 15 females, and the dataset is not fully balanced with all sire families having all

three dams, split in all three diets, and with two replicates per diet. The realized totals of

sire half-sibling families and dam full-sibling families available for analysis in each diet

are presented in Table S2.

Diet treatments. Offspring of each sire were split across three experimental diets:

control (C), dietary restriction (DR), and high sugar (HS). These diets were also used in

our previous mapping study using the DSPR RILs by Stanley *et al.* (2017) and the

composition of each is detailed in Table S1. We used the SAFPro Relax + YF 73050

brand of yeast typically containing 45-60g of protein and 30-38g of carbohydrate per

100g inactivated yeast (Lesaffre Yeast Corp., Milwaukee, USA). To preserve quality,

diets were stored at 4 °C and used within two weeks of preparation. To permit

phenotype measurement and ensure that food did not degrade, individuals were moved

to vials with fresh food three times per week. All flies in all experiments described here

were reared in a growth chamber at 23°C, ≥ 50% relative humidity, and a 24:0 light:dark

cycle, which are the typical maintenance conditions for the DSPR flies.

Phenotype measurement

We measured two phenotypes: 1) lifespan - the total number of days each fly

lived from the date of eclosion, and 2) weekly fecundity - the total number of eggs a set

of females in each vial produced within a 24-hour period measured once per week. Both

phenotypes were measured from the same set of experimental vials over a full lifespan.

For lifespan, the number and sex of dead, and number of censored (i.e., escapees,

accidental deaths) events were recorded three times each week during transfer of flies

to new food until all flies in each vial had died. To estimate fecundity, females were

provided with fresh media each Monday and following a 24 hour egg laying period,

these egg vials were collected on Tuesday and stored at -20 °C until processing. We

continued this process each week until all females within a vial had died. We modified a

protocol developed in Michael Rose's laboratory by Larry G. Cabral at UC Irvine based

on personal communication and video tutorials (Burke et al. 2016; Rose Lab Data

2012a,b) to filter eggs onto circular discs of black filter paper. Subsequently, discs

bearing eggs were photographed using a Canon EOS Rebel TS*i* (Canon Inc, Japan)

camera using a soft uniform lit custom studio placed in a dark room (see Ng'oma *et al*,

2018 for details). The following camera settings were used for imaging: exposure time

1/25 seconds, aperture F5.6, and ISO 100. Image files were uploaded to a computing

system and immediately inspected with ImageJ (Schindelin et al, 2012, Rueden et al

2017) for quality. A detailed description of our method can be found in Ngoma et al

(2018).

Statistical analysis

All following statistical analyses were performed in R (ver. 3.5; R Core Team

2018), and all code is available online (https://github.com/EGKingLab/h2lifespan).

Survival estimates

We converted daily counts of dead and censored individuals to individual events

at a given age to perform survival analysis. Using the R package survival (ver. 2.38;

(Therneau and Grambsch, 2000; Therneau, 2015), we used the Kaplan-Meier estimator

(Kaplan and Meier, 1958) to estimate survival in each diet:

234   survfit(Surv(time, status) ~ diet)

where time is time (days), status is alive/dead, diet is diet treatment (C, DR, HS) in

order to visualize survivorship curves in each diet.

To assess statistical differences between diets accounting for family

membership, we used hierarchical Bayesian inference to test for the effects of sire, dam

nested in sire, vial nested in dam, and treatment using a set of six nested models, which

tested variously for effects of diet treatment, sire, dam, and vial (Table 1). We used the

brms interface (Bürkner, 2017) to the statistical modeling language stan (Carpenter *et*

al., 2017), which samples posterior estimates of modeled parameters using Hamiltonian

Monte Carlo. Lifespan was modeled with a Weibull distribution, which is commonly used

to analyze time to failure (in this case, fly death). We used broad priors for the Weibull

shape parameter ($\Gamma(0.01, 0.01)$), Student's $t(3, 4, 10)$ for the intercept, and Student's

$t(3, 0, 10)$ for the standard deviation, which were quickly overwhelmed by the large

amount of data (7486 observations). Models were compared using leave-one-out (loo)

cross-validation and loo model weights (Vehtari *et al.*, 2017; Yao *et al.*, 2018), the latter

which assigns relative out of sample predictive ability to each model compared.

Posterior estimates of DR and HS diet treatments were compared to the C diet

treatment via credible intervals. We sampled each model for 2000 iterations with 50%

discarded for warm-up and 4 replicate chains, to yield approximately 4000 total

samples. We found that due to the large amount of data, this sampling approach was

adequate. Adequate sampling was determined by $\hat{R}$ values of 1 (Gelman and Rubin,

1992; Brooks and Gelman, 1998).

Fecundity estimates

We used a high through-put method to extract egg counts from images by

constructing an optimal predictive model. To accomplish this, we took advantage of the

simple relationship between number of eggs present and the amount of white area on a

thresholded image of disc and used a set of hand-counted images to optimize the

model. We were able to determine the optimal number of hand-counted images, select

an appropriate threshold value, and assess model performance across a range of

parameters. This method performs very well, with a 0.88 correlation between the model

predicted egg counts and hand counted egg counts. A detailed account of our method is

presented elsewhere (Ng'oma et al 2018).

We focused on two fecundity measurements from our weekly fecundity

measures. We calculated the number of eggs per female for each of our weekly

measures by dividing by the number of females alive in each vial. We then obtained an

estimate of total fecundity per female by simply summing across weeks (hereafter

referred to as total fecundity). We note that this is not strictly a measure of lifetime

fecundity as we measured fecundity only once per week, though we would expect this

estimate to be highly correlated with lifetime fecundity. Second, we considered a

snapshot of early life fecundity by choosing the time point closest to 5 days post-

eclosion (hereafter referred to as early fecundity). The actual age of the females varies

slightly as the vials were set up over several days, but our fecundity measurements

always took place on a 24 hour period beginning on Monday.

We tested for the effects of sire, dam nested in sire, and treatment using a set of

six nested hierarchical (mixed) linear models (Gelman and Hill, 2007), which tested

variously for effects of diet treatment, sire and dam, identical to the models used to test

survival estimates but without the vial effect given our measurements for fecundity are

averages per vial (Table 1). Sampling was done on zero-centered values for early life

and total fecundity using Hamiltonian Monte Carlo using the statistical modeling

language stan (Carpenter *et al.*, 2017), via the rstanarm interface (ver. 2.13.1; (Stan

Development Team, 2016). We used mildly regularizing priors: Normal(0, 10) for

intercepts, Normal(0, 10) for diet treatment parameters, and Cauchy(0, 1) for variances.

Priors for covariance matrices were set to 1 for regularization, concentration, shape,

and scale (i.e., decov(1, 1, 1, 1)). Models were sampled for 20,000 iterations, with

10,000 discarded for burnin. Adequate sampling was assessed by $\hat{R}$, and models were

compared using leave-one-out (loo) cross-validation and loo model weights as

described above.

     Heritability in single diets

To estimate variance components, heritability, and phenotypic correlations, we

used MCMCglmm (v. 2.24; Hadfield, 2010), a Bayesian inference package for R (v. 3.5;

R Core Team, 2016) to fit variations of the 'animal model' (Kruuk, 2004; Wilson *et al.*,

2010; Ingleby *et al.*, 2013). Using the animal model we first inferred, in each diet

separately, the additive genetic effect (V_A) from pedigree information where the

identities of sire and dam are included in the random effect

298 $$\mathbf{y} \sim \beta + \mathbf{Zu}$$

where **y** is the vector of phenotypes (lifespan, early fecundity, or total fecundity), β is the

mean phenotype, **Z** includes information about sire and dam identity (the "animal" term),

and **u** is the parameter vector that is estimated for the random effect, distributed as

Normal(0, σ^2). We did not include vial in order to estimate vial effects as with a half-

sibling design in which half-siblings do not share the same vial, these effects will not

influence the estimation of the additive genetic variance. We used weakly informative

priors: $V = 1$, $v = 0.002$ for analysis of lifespan, although analyses were robust when

more informative priors were tested. Similarly, we used weakly informative, parameter

expanded, priors for analyses of fecundity. Parameter expanded priors, which improve

prior distributions of variance components, can improve sampling in some cases

(Hadfield, 2018). The models were sampled for 2×10^6 iterations, discarding the first 1.5

$\times 10^4$ iterations as burnin, with a thinning interval of 50 iterations. These sampling

parameters yielded effective sample size >10,000. We evaluated the models for

convergence and autocorrelation of the samples, ensuring autocorrelation of <0.1 for

the first lag, and absence of trend in the trace plots of the mean β (i.e., intercept term),

additive variance V_A, and residual V_R (the later two denoted in the output as 'animal' and

'units', respectively).

Genetic correlations between diets

A single phenotype that is measured in multiple environments (here diets) may

be regarded as different phenotypes in those environments (Falconer, 1952; Lynch and

Walsh, 1998). Based on this premise, we computed genetic correlations for phenotype-

diet combinations by fitting a Bayesian multivariate animal model:

$$\mathbf{y} = \mathbf{X\beta} + \mathbf{Zu}$$

where **y** is a matrix of phenotypes (C, DR, HS), **β** is a vector of trait (**X**) means (with no

intercept term), **Z** includes information about sire and dam identity (the "animal" term),

and **u** is the parameter vector that is estimated for the random effect. Among the

random effects, covariance was free to vary (i.e., 'random = ~us(trait):animal') with one

restricting the variance to zero (i.e., 'random = ~idh(trait):animal'). If family genetic

effects remain the same or change in proportion between diets, then the genetic

correlation of family members between diets is 1. Genetic correlations that are credibly

smaller than 1 indicate a significant genotype x diet interaction. We therefore compared

the model above with a model in which the correlation between diets is constrained to

be 1 to test if our estimated genetic correlations were significantly different from 1. In

bivariate models, we tested several slightly stronger variations of priors (V = diag(3), v =

1.002). As above, we used a parameter expanded prior for fecundity. Model were run

for 6.5×10^6 iterations, with 5×10^4 discarded as burning, with thinning every 500

iterations. Twelve chains were run in parallel, yielding >10,000 samples total.

Results

Phenotypic response to dietary treatment

    Lifespan

    We used a split family design, splitting offspring from families into three different diets. Relative to the control diet, median survival was 24% lower on high sugar (HS: 48 days vs. C: 63 days) and 8% higher on DR diet (68 days) (Fig. 2b). Lifespan trajectories began to diverge early, from about 25 days post oviposition and remained diverged until under 10% survivorship (Fig. 2b, Table S3). Individual sire family responses to diet are presented in Fig. S1.

    We compared six nested models of survival, incorporating diet treatment and various aspects of relatedness and compared these models using leave-one-out model weighting (Table 1). The model including sire, dam nested in sire, vial nested in dam, diet, and a sire by diet interaction has the overwhelming model support (>95% of model weight), indicating a significant effect of diet, the presence of significant genetic variance for survival within the diet treatments, and significant genetic variation for the response to diet. Specifically, the results of this favored model show that, compared to the C diet group, individuals in the DR group were credibly less likely to die (99% credible interval of the difference between DR and C = 0.4 - 0.12) and individuals in the HS treatment group were credibly more likely to die (99% credible interval of the difference between HS and C = -0.21 - -0.13).

    Fecundity

    Fecundity was estimated once per week in the same treatment setups from which lifespan was recorded in order to compare the two phenotypes in the same set of

flies. As expected, in all treatments, fecundity declined with age (Fig. 2a). We used a

Bayesian model comparison approach to analyze the effects of diet and family on total

fecundity and early fecundity (Table 1). This analysis revealed a strong effect of diet for

both total fecundity and early fecundity, as models including diet were strongly preferred

over a model without diet. For both measures of fecundity, the highest values were in

the C diet and the lowest in the HS diet (Fig. 2a,c). There was a credible effect of family

for both fecundity measures, indicating significant genetic variation for fecundity. For

early fecundity, the model including only dam nested within sire was favored heavily.

For total fecundity, the model that included sire, dam nested in sire, and the interaction

between sire and diet was favored over the other models, indicating genetic variation for

the response to diet in addition to genetic variation for fecundity.

Estimates of quantitative genetic parameters

We have a high sample size for lifespan with data for up to 48 individuals spread

across two vials for each dam family in each diet. While these same individuals

contributed to our measurements of weekly fecundity, we have a single measurement

for each vial of females at any given time point and do not have measurements per

each individual, as measuring fecundity for each individual female was not feasible for

this study. Thus, for estimating quantitative genetic parameters, our replication for

fecundity is low (2 per family per diet). The results of the above models reflect the

expected uncertainty from this low replication with wide posterior distributions. Thus,

while the models we fit above demonstrate the presence of significant genetic variation,

our estimates of heritability values have high uncertainty and wide credible intervals.

Therefore, we focus on discussing our results for lifespan below and results for

fecundity can be found in Fig. S2.

Heritability of lifespan within diets

We estimated the heritabilities of our phenotypes separately in each diet

treatment using a Bayesian approach (see Methods). Our heritability estimates for

lifespan are moderately high, ranging from 0.31 to 0.47 (Figure 3a). Heritability (h^2) and

95% HPDI (highest posterior density interval) for each treatment were: C h^2 0.47 (0.34 –

0.61), HS h^2 0.37 (0.25 – 0.50), DR: h^2 0.31 (0.21 – 0.43). 95% HPDIs of pairwise

differences of posterior estimates of heritability did not differ credibly from 0: C-DR = -

0.02 – 0.34, C-HS = -0.09 – 0.29, and HS-DR = -0.11 – 0.23.

Correlation of lifespan across dietary treatments

To determine if there is substantial genetic variation for the response to the

environment (GEI), we fit a multivariate form of the animal model, which considers

lifespan within each diet as a separate phenotype (Falconer, 1952; Lynch and Walsh,

1998; Wilson *et al.*, 2010). When the genetic correlation between environments is less

than one, it indicates significant genetically based differences in the response to the

environment, or a significant GEI. In all pairwise diet comparisons, our estimates of the

genetic correlations are credibly less than one, indicating substantial genetically based

variation in the response to diet (Fig. 3b). Genetic correlations (r_g) and HPDIs were: DR

vs. C: r_g 0.48 (0.34 – 0.61), HS vs. C: r_g 0.29 (0.21 – 0.43), HS vs. DR: r_g 0.34 (0.25 –

0.50. This variability is also apparent when examining the variation among sire families

in their response to diet for both lifespan and total fecundity (Fig. 3c & d). In addition,

the model comparisons we employed for survival and total fecundity both favored

models including a sire by diet interaction, which is consistent with the presence of a

significant GEI. Subsequent analysis of differences of posterior estimates of genetic

correlations did not reveal credible differences: HS vs. C - DR vs. C = -0.51 – 0.12, HS

vs. C - HS vs. DR = -0.35 – 0.24, and DR vs. C - HS vs. DR = -0.19 - 0.47, indicating

similarities in genetic correlations among treatment pairs.

Discussion

In this study, we used an admixed multiparent population of *D. melanogaster* to

characterize the quantitative genetics of fitness components in different nutritional

conditions. Our study is one of only a few studies that estimate both lifespan and

lifetime fecundity across multiple diets in the same families, providing a comprehensive

picture of the interplay between the genetic basis of important life history traits and the

nutritional environment. We are able to show, not only that these traits harbor

substantial genetic variation both within diets and in the response to diet, but we also

provide concrete estimates of both the narrow-sense heritability of lifespan in multiple

diets and the genetic correlation of lifespan between diets. These results have important

implications for our understanding of the evolution of these traits in wild populations and

for strategies to uncover the genetic basis of these traits, which we discuss below.

Evolutionary potential of the coordination between nutrition and fitness components

We characterized the effects of three different diets, a dietary restriction diet

(DR), a control diet (C), and a high sugar diet (HS). Dietary restriction has been studied

extensively in *D. melanogaster* and other metazoans (e.g., Chippindale *et al.*, 1993a;

Good and Tatar, 2001; Vrtílek and Reichard, 2014; reviewed in: Sohal and Weindruch,

1996; Kenyon, 2005; Kirkwood and Shanley, 2005; Piper *et al.*, 2011; Tatar, 2011; Tatar

et al., 2014). A general finding of these studies is that reduced nutrient intake without

starvation results in lifespan extension, and that this effect is often coupled with reduced

reproductive output, though there are exceptions to this pattern (e.g., Kaitala, 1991;

Weindruch *et al.*, 1995; Kirk *et al.*, 2001; Stelzer, 2001; Cooper *et al.*, 2004; reviewed in

Ng'oma *et al.*, 2017). Even within species that tend to show this effect such as *D.*

melanogaster, there is variability in this pattern among different genotypes. For

example, Stanley *et al.* (2017) found variation among inbred lines for the amount of

lifespan extension, with some lines showing little to no extension in a DR diet, and a few

showing reduced lifespan in a DR diet, a result that has also been shown in mice (Rikke

et al., 2003). Unlike for DR, the effects of a high sugar diet have not been studied as

extensively, though recently some studies have shown flies reared on high sugar diets

as larvae show obesity-like phenotypes (Musselman *et al.*, 2011; Rovenko *et al.*, 2015),

and adults reared on high sugar diets have elevated triglyceride levels, reduced

lifespan, and reduced fecundity (Skorupa *et al.*, 2008), a pattern that is consistent with

our findings. In these studies, "high sugar" refers to a high concentration of sugar

relative to the volume of the diet, and that is how we also use the term to describe our

high sugar diet. However, nutritional geometry approaches focus on the protein to

carbohydrate ratio of the diet, and thus "high sugar" diets in this case would correspond

to a diet with a low P:C ratio. In this context, our HS diet, with a yeast-sugar ratio of

about 1:2 resulted in reduction of both lifespan and fecundity. Past studies (Skorupa *et*

al, 2008, Edgecomb *et al*, 1994) observed that flies on a diet richer in carbohydrate

tended to feed more, and that imbalance in nutrient intake resulted in shorter lifespan.

Other studies including Lee *et al* (2008) and Jensen *et al*, (2015), on the other hand,

found lifespan and lifetime fecundity maximized at much lower P:C ratios (1:16 and 1:4,

respectively). Apparent incosistence of our results (and studies cited above) with these

studies that measure specific nutrient content likely reflect diet component composition

relative to volume. We note, however, that our diets use agar to form a semi-solid media

while these studies use a liquid diet, and that our flies are maintained under 24 hour

light conditions, both of which may also lead to differences in the patterns observed.

These types of phenotypic patterns in different diet conditions have been

interpreted in different ways. The widespread observation of lifespan extension

observed in response to dietary restriction in many taxa has led to competing

hypotheses for this pattern. It is often argued that this pattern is the result of natural

selection acting in environments with fluctuating resources where it should be adaptive

to increase allocation of nutrient resources to the soma when nutrients are limiting,

however it is also possible it is the result of an unavoidable physiological constraint

(Holliday, 1989; Kirkwood and Shanley 2000; Kirkwood and Shanley, 2005).

Additionally, studies employing high sugar and high fat diets often view the results of

those studies in the context of human populations where it is hypothesized the negative

health consequences of a Westernized diet result from a mismatch between that diet

and the diet humans adapted to in the past over the majority of their evolutionary history

(Neel, 1962; Wells, 2009), though this hypothesis has also been challenged

(Speakman, 2008). Lately, a number of studies have challenged the resource allocation

framework to explain life history trade-offs (see Barnes *et al*, 2006; Flatt *et al*, 2008;

Adler *et al*, 2013). These studies (and others) suggest that the decoupling of lifespan

from fecundity that is sometimes observed, relates to hormonal signaling rather than

literal resource allocation. However, the works of Zhao and Zera (2006), Zera and Zhao

(2006), and Zera (2005) demonstrate the biochemical basis of life history trade-offs

(reviewed in Ng'oma et al 2017). In general, assessing the potential for phenotypic

effects of different diets to be adaptive patterns or to be highly constrained physiological

responses requires an understanding of the underlying genetic variation for the

response to diet in order to understand if this response has the potential to evolve.

Here, we showed there is heritable variation for both phenotypes within diets,

and for the response of phenotypes to diet (Table 1; Fig. 3a,b). Previous mapping

studies in *D. melnanogaster* have identified several QTL influencing life history traits

(reviewed in Paaby and Schmidt, 2009), including mapping QTL influencing changes in

gene expression and lifespan in different diets (Stanley et al. 2017). However, like other

complex traits, these QTL fail to explain most of the underlying genetic variance. Our

results support the hypothesis that these phenotypes are influenced by many genes and

quantify the amount of variation among individuals attributable to genetic variation in

this population. While many studies have demonstrated GEI for different traits (Reed *et*

al., 2010; Qi *et al.*, 2012; Ingleby *et al.*, 2013), few studies have estimated GEI

specifically for life history traits in response to diet, and not all previous studies of diet

have found significant GEI. For example, King *et al.* (2011) estimated quantitative

genetic parameters for allocation to flight capability and to reproduction in the sand

cricket, *Gryllus firmus.* These authors found no GEI for these traits, with all genotypes

allocating more to reproduction when food levels increase, suggesting important

evolutionary constraints can exist for the response to diet and that significant GEI is not

a foregone conclusion. Our results show that this constraint does not exist for this

population of *D. melanogaster* and that the response to diet may readily evolve in

response to selection. The founder lines of the DSPR have a global distribution and

were derived from wild-caught flies and thus, our population encompases some of the

genetic variation that exists among natural populations of *D. melanogaster.* It is

plausible, therefore, that the diet dependent effects observed in this study and others

represent adaptive patterns that have evolved. Some recent experimental evolution

studies exposing flies to constant DR, standard, and high protein diets have shown

mixed and sex-specific results in which females exposed to high protein diet evolved

increased fecundity but decreased survival on standard and DR diets (Zajitschek et al

2016), while males on DR had greater fitness compared to those on standard and high

diets (Zajitschek et al 2018). These studies establish that the responses of fecundity

and lifespan to diet do evolve, but the emergent patterns do not support the typical

hypothesis that selection on differential resource allocation is driving the pattern. It

remains a challenge to interpret how the patterns we observe in these kinds of

laboratory studies represent the nutritional conditions experienced by flies in the wild

and by what mechanisms those conditions have driven the evolution of different life

history strategies. It would be interesting to see if future studies will also find

evoltuionary potential for the response to changes in the P:C ratio.

Prospects for multiparent population approaches

One of the most fundamental goals in biology is to understand the genetic basis

of complex phenotypes. This goal has proved to be quite challenging. For most traits,

some of which have been the focus of study for many years, the causative genetic

variants that have been discovered explain only a small percentage of the trait

heritability (for reviews see McCarthy *et al.*, 2008; Manolio *et al.*, 2009; Rockman, 2012;

Visscher *et al.*, 2012). For traits that are expected to be highly polygenic, such as

lifespan and fecundity, estimating quantitative genetic parameters as we have done in

this study, can be as informative about the genetic basis of a trait and its evolutionary

potential as studies aimed at uncovering specific genetic variants.

In recent years, multiparent populations, in which multiple inbred founder lines

are crossed for a number of generations to create a recombinant mapping population,

have increasingly been employed in studies aiming to identify the causative genetic

variants underlying complex traits (e.g. (Kover *et al.*, 2009; McMullen *et al.*, 2009;

Huang *et al.*, 2011; Aylor *et al.*, 2011; Threadgill and Churchill, 2012; King *et al.*, 2012;

King, *et al.*, 2012; Cubillos *et al.*, 2013); reviewed in (Long *et al.*, 2014; de Koning and

McIntyre, 2017). One reason for employing a mapping population consisting of multiple

parent lines is the inclusion of additional genetic variation when compared to traditional

two-line QTL mapping panels. Even with multiple parent lines contributing, a potential

criticism of these kinds of approaches when compared to mapping approaches that

sample from natural populations is that they contain limited amounts of genetic

variation. The DSPR population used in this study was started from 8 inbred parent

lines. Therefore, while every individual in our admixed population has a unique mosaic

genome, at any given position, there are only 8 possible haplotypes. Despite this

simplicity, we find moderate heritabilities for lifespan in all diets (ranging from ~0.3 - 0.5)

that are comparable to estimates from natural populations. A meta-analysis of

heritability estimates from natural populations found life history traits typically have

values around 0.25 and values higher than 0.3 are not infrequent (Rose and

Charlesworth, 1981; Mousseau and Roff, 1987; Rikke *et al.*, 2010). We also found

strong evidence for a genotype by diet interaction for lifespan, indicating there is also

segregating genetic variation for the response to diet. While our evidence of a genotype

by diet interaction is not as strong for fecundity, our model comparison for total fecundity

supports a model including this interaction.

In addition to their utility for QTL mapping, multiparent populations have also

been used in evolve and resequence approaches (e.g. Turner and Miller, 2012; Burke

et al., 2014 a,b) to identify the underlying genetic basis of complex traits. These studies

impose selection and track the resulting changes in phenotype and genotype. This

approach is potentially very powerful because 1) entire haplotypes can be tracked over

evolutionary time, 2) the simplified genetic architecture with only a few potential

haplotypes at any given location may lead to a more repeatable response to selection

increasing the likelihood of identifying causative loci, and 3) it may allow for validation of

QTL identified in mapping studies using these same populations. A potential downside

to such an approach is if the relatively simple genetic architecture of such a population

results in limited genetic variation for the trait of interest, thereby constraining evolution.

Our study shows there is substantial genetic variation for our focal phenotypes in our

admixed DSPR population, which establishes this approach as a viable strategy for

investigating both how a focal trait evolves and the underlying genetic basis. It is one of

only a few studies to quantify quantitative genetic parameters for an outbred multiparent

population, providing evidence that segregating genetic variation is not limiting in these

populations. In general, MPPs have great potential to advance our understanding of the

genetic basis of complex traits via these kinds of multi-pronged approaches.

Acknowledgments

We received helpful suggestions from Larry Cabral on how to establish our fecundity

protocol. We had insightful discussions with Ian Dworkin on fitting the animal model in

the Bayesian framework. Stuart Macdonald provided the DSPR RILs used to start our

experimental colony. Elizabeth Lopresti, Michael Reed, Osvaldo Enriquez, Anna

Perinchery and Kyla Winford helped with fly husbandry, experimental set-ups, collection

and entry of data. This work was supported by NIH grant R01 GM117135 to Elizabeth

G. King, The University of Missouri, and a University of Missouri Research Board Grant.

Conflict of interest

The authors declare no conflict of interest.

Data availability

Raw data, including lifespan, fecundity, and image files, may be retrieved from Zenodo

at http://doi.org/10.5281/zenodo.1285237. Scripts to reproduce all analyses are

available at GitHub: https://github.com/EGKingLab/h2lifespan. Supplementary

information is available at https://www.nature.com/hdy/

References

Adler MI, Cassidy EJ, Fricke C, Bonduriansky R (2013). The lifespan-reproduction

trade-off under dietary restriction is sex-specific and context-dependent. *Exp Gerontol*

48: 539–548.

Aylor DL, Valdar W, Foulds-Mathes W, Buus RJ, Verdugo RA, Baric RS, *et al.* (2011).

Genetic analysis of complex traits in the emerging Collaborative Cross. *Genome Res*

21: 1213–1222.

Barnes AI, Boone JM, Jacobson J, Partridge L, Chapman T (2006). No extension of

lifespan by ablation of germ line in *Drosophila*. *Proc Biol Sci* **273**: 939–947.

Bass TM, Grandison RC, Wong R, Martinez P, Partridge L, Piper MDW (2007).

Optimization of dietary restriction protocols in Drosophila. *J Gerontol A-Biol* **62**: 1071–

1081.

Brčić-Kostić K (2010). Quantitative genetics and evolution. *Periodicum biologorum* **112**:

395–402.

Brooks SP, Gelman A (1998). General methods for monitoring convergence of iterative

simulations. *Journal of Computational and Graphical Statistics* **7**: 434–455.

Burke MK, King EG, Shahrestani P, Rose MR, Long AD (2014). Genome-wide

association study of extreme longevity in *Drosophila melanogaster*. *Genome Biology*

and Evolution **6**: 1–11.

Burke MK, Liti G, Long AD (2014). Standing genetic variation drives repeatable

experimental evolution in outcrossing populations of *Saccharomyces cerevisiae*. *Mol*

Biol Evol **31**: 3228–3239.

Burke MK, Barter TT, Cabral LG, Kezos JN, Phillips MA, Rutledge GA, *et al.* (2016).

Rapid divergence and convergence of life-history in experimentally evolved Drosophila

melanogaster. *Evolution* **70**: 2085–2098.

Bürkner P-C (2017). brms: an r package for bayesian multilevel models using stan.

Journal of Statistical Software, **80**: 1–28

Carnes, MU, Campbell T, Huang W, Butler DG, Carbone MA, Duncan LH, *et al.* (2015).

The genomic basis of postponed senescence in *Drosophila melanogaster*. *PLOS ONE*

10: e0138569.

Carpenter B, Gelman A, Hoffman M, Lee D, Goodrich B, Betancourt M, *et al.* (2017).

Stan: a probabilistic programming language. *Journal of Statistical Software*, **76**: 1–32.

Chippindale A, Leroi AM, Kim S, Rose M (1993a). Phenotypic plasticity and selection in *Drosophila* life-history evolution. I. Nutrition and the cost of reproduction. *J Evolution Biol* **6**: 171–193.

Chippindale AK, Leroi AM, Kim SB, Rose MR (1993b). Phenotypic plasticity and selection in *Drosophila* life-history evolution. I. Nutrition and the cost of reproduction. *Journal of Evolutionary Biology* **6**: 171–193.

Cooper TM, Mockett RJ, Sohal BH, Sohal RS, Orr WC (2004). Effect of caloric restriction on life span of the housefly, *Musca domestica*. *FASEB J* **18**: 1591–1593.

Cubillos FA, Parts L, Salinas F, Bergström A, Scovacricchi E, Zia A, *et al.* (2013). High-resolution mapping of complex traits with a four-parent advanced intercross yeast population. *Genetics* **195**: 1141–1155.

Edgecomb RS, Harth CE, Schneiderman AM (1994). Regulation of feeding behavior in adult Drosophila melanogaster varies with feeding regime and nutritional state. *J Exp Biol* **197**: 215–235.

Falconer DS (1952). The problem of environment and selection. *The American Naturalist* **86**: 293–298.

Fisher RA (1930). *The genetical theory of natural selection.* Clarendon Press: Oxford.

Fisher JAD, Frank KT, Leggett WC (2010). Breaking Bergmann's rule: truncation of Northwest Atlantic marine fish body sizes. *Ecology* **91**: 2499–2505.

Flatt T, Min K-J, D'Alterio C, Villa-Cuesta E, Cumbers J, Lehmann R, *et al.* (2008). *Drosophila* germ-line modulation of insulin signaling and lifespan. *P Natl Acad Sci USA* **105**: 6368–6373.

Gagneur J, Stegle O, Zhu C, Jakob P, Tekkedil MM, Aiyar RS, *et al.* (2013). Genotype-

environment interactions reveal causal pathways that mediate genetic effects on

phenotype. *PLoS Genet* **9**: e1003803.

Gelman A, Hill J (2007). *Data analysis using regression and multilevel/hierarchical*

models. Cambridge University Press: Cambridge ; New York.

Gelman A, Rubin D (1992). A single series from the Gibbs sampler provides a false

sense of security. *Bayesian Statistics* **4**: 625–631.

Giannakou ME, Goss M, Partridge L (2008). Role of dFOXO in lifespan extension by

dietary restriction in *Drosophila melanogaster*: not required, but its activity modulates

the response. *Aging Cell* **7**: 187–198.

Good TP, Tatar M (2001). Age-specific mortality and reproduction respond to adult

dietary restriction in *Drosophila melanogaster*. *J Insect Physiol* **47**: 1467–1473.

Grandison RC, Wong R, Bass TM, Partridge L, Piper MDW (2009). Effect of a

standardised dietary restriction protocol on multiple laboratory strains of drosophila

melanogaster (H Tanimoto, Ed.). *PLOS ONE* **4**: e4067.

Greer EL, Brunet A (2009). Different dietary restriction regimens extend lifespan by both

independent and overlapping genetic pathways in C. elegans. *Aging Cell* **8**: 113–127.

Hadfield JD (2010). MCMCglmm course notes. https://cran.r-

project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf.

Holliday R (1989). Food, reproduction and longevity: is the extended lifespan of calorie-

restricted animals an evolutionary adaptation? *Bioessays* **10**: 125–127.

Hsin H, Kenyon C (1999). Signals from the reproductive system regulate the lifespan of

C. elegans. *Nature* **399**: 362–366.

Huang X, Paulo M-J, Boer M, Effgen S, Keizer P, Koornneef M, *et al.* (2011). Analysis

of natural allelic variation in *Arabidopsis* using a multiparent recombinant inbred line

population. *PNAS* **108**: 4488–4493.

Hughes KA, Reynolds RM (2005). Evolutionary and mechanistic theories of aging. *Annu*

Rev Entomol **50**: 421–445.

Hunt J, Bussière LF, Jennions MD, Brooks R (2004). What is genetic quality? *Trends in*

Ecology & Evolution **19**: 329–333.

Ingleby FC, Hosken DJ, Flowers K, Hawkes MF, Lane SM, Rapkin J, *et al.* (2013).

Genotype-by-environment interactions for cuticular hydrocarbon expression in

Drosophila simulans. J Evol Biol **26**: 94–107.

Jensen K, McClure C, Priest NK, Hunt J (2015). Sex-specific effects of protein and

carbohydrate intake on reproduction but not lifespan in Drosophila melanogaster. *Aging*

Cell **14**: 605–615.

Kaitala A (1991). Phenotypic plasticity in reproductive behavior of waterstriders: trade-

offs between reproduction and longevity during food stress. *Funct Ecol* **5**: 12–18.

Kaplan EL, Meier P (1958). Nonparametric estimation from incomplete observations.

Journal of the American Statistical Association **53**: 457–481.

Kenyon CJ (2005). The plasticity of aging: insights from long-lived mutants. *Cell* **120**:

449–460.

King EG, Macdonald SJ, Long AD (2012). Properties and power of the *Drosophila*

Synthetic Population Resource for the routine dissection of complex traits. *Genetics*

191: 935–949.

King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, *et al.* (2012). Genetic

dissection of a model complex trait using the *Drosophila* Synthetic Population Resource.

Genome Res **22**: 1558–1566.

King EG, Roff DA, Fairbairn DJ (2011). The evolutionary genetics of acquisition and

allocation in the wing dimorphic cricket, *Gryllus firmus. Evolution* **65**: 2273–2285.

Kirk KL, Turner TL, Kirkwood TBL, Reed LK, Miller PM, Shanley DP, *et al.* (2001).

Dietary restriction and aging: comparative tests of evolutionary hypotheses. *Journal of*

Gerontology: BIOLOGICAL SCIENCES **56A**: B123-129.

Kirkwood TB (1977). Evolution of ageing. *Nature* **270**: 301–304.

Kirkwood TBL, Shanley DP (2005). Food restriction, evolution and ageing. *Mech Ageing*

Dev **126**: 1011–1016.

de Koning D-J, McIntyre LM (2017). Back to the future: multiparent populations provide

the key to unlocking the genetic basis of complex traits. *Genetics* **206**: 527–529.

Kover PX, Valdar W, Trakalo J, Scarcelli N, Ehrenreich IM, Purugganan MD, *et al.*

(2009). A multiparent advanced generation inter-cross to fine-map quantitative traits in

Arabidopsis thaliana. PLoS Genet **5**: e1000551.

Kruuk LEB (2004). Estimating genetic parameters in natural populations using the

'animal model'. *Philos Trans R Soc Lond B Biol Sci* **359**: 873–890.

Lande R, Arnold SJ (1983). The measurement of selection on correlated characters.

Evolution **37**: 1210–1226.

Lee KP, Simpson SJ, Clissold FJ, Brooks R, Ballard JWO, Taylor PW, *et al.* (2008).

Lifespan and reproduction in Drosophila: new insights from nutritional geometry. *Proc*

Natl Acad Sci U S A **105**: 2498–2503.

Long AD, Macdonald SJ, King EG (2014). Dissecting complex traits using the

Drosophila Synthetic Population Resource. *Trends in Genetics* **30**: 488–495.

Lynch M, Walsh B (1998). *Genetics and analysis of quantitative traits.* Sinauer

Associates, Inc.: Sunderland, MA.

Mair W, Dillin A (2008). Aging and survival: the genetics of life span extension by dietary

restriction. *Annu Rev Biochem* **77**: 727–754.

Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, *et al.* (2009).

Finding the missing heritability of complex diseases. *Nature* **461**: 747–753.

McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JPA, *et al.*

(2008). Genome-wide association studies for complex traits: consensus, uncertainty

and challenges. *Nature Reviews Genetics* **9**: 356–369.

McMullen MD, Kresovich S, Villeda HS, Bradbury P, Li H, Sun Q, *et al.* (2009). Genetic

properties of the maize nested association mapping population. *Science* **325**: 737–740.

Min K-J, Yamamoto R, Buch S, Pankratz M, Tatar M (2008). *Drosophila* lifespan control

by dietary restriction independent of insulin-like signaling. *Aging Cell* **7**: 199–206.

Moatt JP, Nakagawa S, Lagisz M, Walling CA (2016). The effect of dietary restriction on

reproduction: a meta-analytic perspective. *BMC Evolutionary Biology* **16**: 199.

Mousseau TA, Roff DA (1987). Natural selection and the heritability of fitness

components. *Heredity* **59**: 181–197.

Musselman LP, Fink JL, Narzinski K, Ramachandran PV, Hathiramani SS, Cagan RL,

et al. (2011). A high-sugar diet produces obesity and insulin resistance in wild-type

Drosophila. *Disease Models & Mechanisms* **4**: 842–849.

Nakagawa S, Lagisz M, Hector KL, Spencer HG (2012). Comparative and meta-analytic

insights into life extension via dietary restriction. *Aging Cell* **11**: 401–409.

Nässel DR, Liu Y, Luo J (2015). Insulin/IGF signaling and its regulation in *Drosophila*.

Gen Comp Endocrinol **221**: 255–266.

Neel J (1962). Diabetes mellitus: A 'thrifty' genotype rendered detrimental by 'progress'?

Am J Hum Genet **14**: 353–362.

Ng'oma E, Perinchery AM, King EG (2017). How to get the most bang for your buck: the

evolution and physiology of nutrition-dependent resource allocation strategies. *Proc Biol*

Sci **284**.

Ng'oma E, King EG, Middleton KM (2018). A model-based high throughput method for

fecundity estimation in fruit fly studies. *bioRxiv*: 382804.

Paaby AB, Schmidt PS (2009). Dissecting the genetics of longevity in *Drosophila*

melanogaster. *Fly (Austin)* **3**: 29–38.

Piper MDW, Partridge L, Raubenheimer D, Simpson SJ (2011). Dietary restriction and

aging: a unifying perspective. *Cell Metab* **14**: 154–160.

Qi Q, Chu AY, Kang JH, Jensen MK, Curhan GC, Pasquale LR, *et al.* (2012). Sugar-

sweetened beverages and genetic risk of obesity. *New England Journal of Medicine*

367: 1387–1396.

R Core Team (2016). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing, Vienna, Austria.

Reed LK, Lee K, Zhang Z, Rashid L, Poe A, Hsieh B, *et al.* (2014). Systems genomics

of metabolic phenotypes in wild-type *Drosophila melanogaster*. *Genetics* **197**: 781–793.

Reed LK, Williams S, Springston M, Brown J, Freeman K, DesRoches CE, *et al.* (2010).

Genotype-by-diet interactions drive metabolic phenotype variation in *Drosophila*

melanogaster. *Genetics* **185**: 1009–1019.

Remolina SC, Chang PL, Leips J, Nuzhdin SV, Hughes KA (2012). Genomic basis of

aging and life-history evolution in *Drosophila melanogaster. Evolution* **66**: 3390–3403.

Rikke BA, Liao C-Y, McQueen MB, Nelson JF, Johnson TE (2010). Genetic dissection

of dietary restriction in mice supports the metabolic efficiency model of life extension.

Experimental Gerontology **45**: 691–701.

Rikke BA, Yerg JE, Battaglia ME, Nagy TR, Allison DB, Johnson TE (2003). Strain

variation in the response of body temperature to dietary restriction. *Mechanisms of*

Ageing and Development **124**: 663–678.

Rockman MV (2012). The QTN program and the alleles that matter for evolution: all

that's gold does not glitter. *Evolution* **66**: 1–17.

Roff DA (2006). *Evolutionary quantitative genetics.* In: Fox CW, Wolf JB (eds).

Evolutionary genetics: concepts and case studies. Oxford Univ. Press: New York. pp.

267–287.

Rose Lab Data. (2012, September 21). A method for gathering fecundity data for

populations of Drosophila melanogaster. [Fecundity: Animated Introduction]. Viewed at

https://www.youtube.com/watch?v=xrEBNB8bUwo&sns=em

Rose Lab Data. (2012, September 21). A method for gathering fecundity data for

populations of Drosophila melanogaster. [Fecundity: Live Demonstration]. Viewed at

https://www.youtube.com/watch?v=uqFZ4ZTC6ZY&sns=em

Rose MR, Charlesworth B (1981). Genetics of life history in *Drosophila melanogaster.* I.

sib analysis of adult females. *Genetics* **97**: 173–186.

Rovenko BM, Perkhulyn NV, Gospodaryov DV, Sanz A, Lushchak OV, Lushchak VI

(2015). High consumption of fructose rather than glucose promotes a diet-induced

obese phenotype in *Drosophila melanogaster*. *Comp Biochem Physiol, Part A Mol*

Integr Physiol **180**: 75–85.

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, *et al.* (2017).

ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*

18: 529.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, *et al.*

(2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**:

676–682.

Shanley DP, Kirkwood TB (2000). Calorie restriction and aging: a life-history analysis.

Evolution **54**: 740–750.

Simmons F, Bradley T (1997). An analysis of resource allocation in response to dietary

yeast in *Drosophila melanogaster*. *J Insect Physiol* **43**: 779–788.

Skorupa DA, Dervisefendic A, Zwiener J, Pletcher SD (2008). Dietary composition

specifies consumption, obesity, and lifespan in *Drosophila melanogaster*. *Aging Cell* **7**:

478–490.

Sohal R, Weindruch R (1996). Oxidative stress, caloric restriction, and aging. *Science*

(New York, NY).

Speakman JR (2008). Thrifty genes for obesity, an attractive but flawed idea, and an

alternative perspective: the 'drifty gene' hypothesis. *Int J Obes Relat Metab Disord* **32**:

1611–1617.

Speakman JR, Mitchell SE (2011). Caloric restriction. *Mol Aspects Med* **32**: 159–221.

Sprott RL (1997). Diet and calorie restriction. *Experimental Gerontology* **32**: 205–214.

Stan Development Team (2016). *rstanarm: Bayesian applied regression modeling via*

Stan. R package version 2.13.1.

Stanley PD, Ng'oma E, O'Day S, King EG (2017). Genetic dissection of nutrition-

induced plasticity in insulin/insulin-like growth factor signaling and median life span in a

Drosophila multiparent population. *Genetics* **206**: 587–602.

Stelzer C (2001). Resource limitation and reproductive effort in a planktonic rotifer.

Ecology **82**: 2521–2533.

Tatar M (2011). The plate half-full: status of research on the mechanisms of dietary

restriction in *Drosophila melanogaster*. *EXG* **46**: 363–368.

Tatar M, Post S, Yu K (2014). Nutrient control of *Drosophila longevity. Trends Endocrin*

Met: 1–9.

Teleman AA (2010). Molecular mechanisms of metabolic regulation by insulin in

Drosophila. Biochem J **425**: 13–26.

Therneau TM (2015). *A package for survival analysis in S*, version 2.38.

https://CRAN.R-project.org/package=survival.

Therneau TM, Grambsch PM (2000). *Modeling survival data: Extending the cox model*.

Springer: New York.

Threadgill DW, Churchill GA (2012). Ten years of the Collaborative Cross. *G3: Genes,*

Genomes, Genetics **2**: 153–156.

Turner TL, Miller PM (2012). Investigating natural variation in *Drosophila* courtship song

by the evolve and resequence approach. *Genetics* **191**: 633–642.

Vehtari A, Gelman A, Gabry J (2017). Practical Bayesian model evaluation using leave-

one-out cross-validation and WAIC. *Stat Comput* **27**: 1413–1432.

Visscher PM, Brown MA, McCarthy MI, Yang J (2012). Five years of GWAS discovery.

The American Journal of Human Genetics **90**: 7–24.

van Noordwijk AJ, de Jong G (1986). Acquisition and allocation of resources: their

influence on variation in life history tactics. *The American Naturalist* **128**: 137–142.

Vrtílek M, Reichard M (2014). Highly plastic resource allocation to growth and

reproduction in females of an African annual fish. *Ecol Freshw Fish* **24**: 616–628.

Warren IA, Gotoh H, Dworkin IM, Emlen DJ, Lavine LC (2013). A general mechanism

for conditional expression of exaggerated sexually-selected traits. *Bioessays* **35**: 889–

899.

Weindruch R, Marriott BM, Conway J, Knapka JJ, Lane MA, Cutler RG, *et al.* (1995).

Measures of body size and growth in rhesus and squirrel monkeys subjected to

long-term dietary restriction. *Am J Primatol* **35**: 207–228.

Wells JCK (2009). Thrift: a guide to thrifty genes, thrifty phenotypes and thrifty norms.

Int J Obes (Lond) **33**: 1331–1338.

Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, *et al.* (2010).

An ecologist's guide to the animal model. *Journal of Animal Ecology* **79**: 13–26.

Yao Y, Vehtari A, Simpson D, Gelman A (2018). Using stacking to average Bayesian

predictive distributions. *Bayesian Anal.*

Zajitschek F, Georgolopoulos G, Vourlou A, Ericsson M, Zajitschek SRK, Friberg U, *et*

al. (2018). Evolution under dietary restriction decouples survival from fecundity in

Drosophila melanogaster females. *J Gerontol A Biol Sci Med Sci.*

Zajitschek F, Zajitschek SRK, Canton C, Georgolopoulos G, Friberg U, Maklakov AA

(2016). Evolution under dietary restriction increases male reproductive performance

without survival cost. *Proc R Soc B* **283**: 20152726.

Zera AJ (2005). Intermediary metabolism and life history trade-offs: lipid metabolism in

lines of the wing-polymorphic cricket, Gryllus firmus, selected for flight capability vs.

early age reproduction. *Integr Comp Biol* **45**: 511–524.

Zera AJ, Zhao Z (2006). Intermediary metabolism and life-history trade-offs: differential

metabolism of amino acids underlies the dispersal-reproduction trade-off in a wing-

polymorphic cricket. *Am Nat* **167**: 889–900.

 Titles and legends to figures

Figure 1: Experimental design. A set of 835 DSPR RILs were crossed for 5 generations

to create an outbred synthetic population. From this population, 28 randomly chosen

sires were each mated to 3 dams (actual N = 77 ♀). The resulting offspring from each

dam were split to each of C, DR, and HS diet treatments with two replicate vials in each

diet.

Figure 2: Phenotypic response to diet treatment. The different colors represent different

diet treatments (grey = C, orange = DR, blue = HS). **a.** Fecundity per female sampled

over a 24 hour period once per week over lifespan. A smooth (loess) fit is shown for

each treatment. Individual lines represent each replicate vial. **b.** Survival probability

versus age in each of the three diet treatments. Median survival (50% survival

probability) is shown with the dotted line for each diet. Censored events are marked with

a + sign on each survivorship curve. **c.** The relationship between early life fecundity (24

hour sample period closest to 5 days post-eclosion) and median lifespan for each vial.

The average values for both within each treatment are shown by the large point outlined

in black. Error bars are the mean +/- 1 standard error. Note that the error bars all fit

within the size of the point.

Figure 3: **a.** Density plots of the estimated posterior probabilities for lifespan heritability

within each diet. **b.** Density plots of the estimated posterior probabilities for the genetic

correlation between lifespan in pairs of diets. **c.** Sire by environment interaction plot for

median lifespan showing the median lifespan for each sire family in each of the three

diets. Each line represents one sire family. **d.** Sire by environment interaction plot for

total egg count per female (summed across weekly measures) showing the median

lifespan for each sire family in each of the three diets. Each line represents one sire

family.

Table 1. Comparison of models of lifespan, early fecundity, and total fecundity across

diet treatments. Models 1 and 2 do not include any aspects of pedigree, and models 3-6

use a hierarchical model. Terms in parentheses are random effects. All random effects

were "intercept" terms, allowing separate intercepts for each level of sire ID, dam ID

nested in sire ID, or sire ID nested in diet treatment.

	Model	Lifespan	Early Fecundity	Total Fecundity
1	~ Intercept	5%	3.1%	1.1%
2	~ Diet	0%	10.3%	0%
3	~ Diet + (Sire ID)	0%	0.0%	2.1%
4	~ Diet + (Dam ID by Sire ID) + [Vial ID by Dam ID][†]	0%	86.6%	19.4%
5	~ Diet + (Sire ID) + (Dam ID by Sire ID) + [Vial ID by Dam ID][†]	0%	0%	0%
6	~ Diet + (Sire ID) + (Dam ID by Sire ID) + (Sire ID by Diet) + [Vial ID by Dam ID][†]	95.1%	0%	77.4%

880 [†]Only the models for lifespan included this vial effect because the response variable for

881 fecundity is a vial average.