

Rhamnolipids from *Pseudomonas aeruginosa* Disperse the Biofilms of Sulfate-Reducing Bacteria

Thammajun L. Wood¹, Lei Zhu¹, James Miller³, Daniel S. Miller⁴, Bei Yin⁴, and
Thomas K. Wood^{1,2,3*}

¹Department of Chemical Engineering, ²Department of Biochemistry and Molecular Biology, and the

³Huck Institutes of the Life Sciences, Pennsylvania State University,

University Park, Pennsylvania, 16802

⁴Dow Chemical Company, Collegeville, PA, 19426

*Correspondence should be addressed to T. K. W. (tuw14@psu.edu)

Keywords: rhamnolipids, biofilm dispersal, *Pseudomonas aeruginosa*, sulfate-reducing bacteria

Running head: Rhamnolipids disperse SRB biofilms

ABSTRACT

Biofilm formation is an important problem for many industries. *Desulfovibrio vulgaris* is the representative sulfate-reducing bacterium (SRB) which causes metal corrosion in oil wells and drilling equipment, and the corrosion is related to its biofilm formation. Biofilms are extremely difficult to
5 remove since the cells are cemented in a polymer matrix. In an effort to eliminate SRB biofilms, we examined the ability of supernatants from *Pseudomonas aeruginosa* PA14 to disperse SRB biofilms. We found that the *P. aeruginosa* supernatants dispersed more than 98% of the biofilm. To determine the genetic basis of this SRB biofilm dispersal, we examined a series of *P. aeruginosa* mutants and found that mutants *rhlA*, *rhlB*, *rhlI*, and *rhlR*, defective in rhamnolipids production, had significantly reduced levels
10 of SRB biofilm dispersal. Corroborating these results, purified rhamnolipids dispersed SRB biofilms, and rhamnolipids were detected in the *P. aeruginosa* supernatants. Hence, *P. aeruginosa* supernatants disperse SRB biofilms via rhamnolipids. In addition, the supernatants of *P. aeruginosa* dispersed the SRB biofilms more readily than protease in M9 glucose minimum medium and were also effective against biofilms of *Escherichia coli* and *Bacillus subtilis*.

15

INTRODUCTION

Sulfate-reducing bacteria (SRB) are an important type of microorganisms causing iron corrosion on metal surfaces under both anaerobic and aerobic conditions¹⁻³. *Desulfovibrio vulgaris* Hildenborough is a sequenced⁴ Gram-negative SRB that has been used as an SRB model organism to study biocorrosion and bioremediation of toxic metal ions⁴ as well as biofilm formation^{5, 6} and bioimmobilization in superfund sites⁷. It is also called the “petroleum pest” because it is commonly found in oil fields and causes “souring” of petroleum and damage to topside equipment and pipelines⁸.

Biofilms are groups of bacteria that are held together in a self-produced extracellular matrix⁹ and are difficult to remove with antimicrobial agents due to their antibiotic or biocide resistance relative to planktonic cells¹⁰. *Desulfovibrio* spp. populations in biofilms have a significant role for microbial induced corrosion because of their sulfide production and electron transfer mechanism⁵, and biofilms of *D. vulgaris* have been extensively shown to cause corrosion in many types of steels and other alloys¹¹. The biofilms of *D. vulgaris* consists primarily of protein⁵, mannose⁶, fucose⁶, and *N*-acetylgalactosamine⁶. The biofilms of *Escherichia coli* consist primarily of proteinaceous curli fibres, flagella, and the polysaccharide cellulose¹², and the biofilms of *Staphylococcus aureus* are largely composed of cytoplasmic proteins¹³ and extracellular genomic DNA¹⁴.

Many Gram-negative bacteria use quorum sensing (QS) molecules or autoinducers to communicate with each other¹⁵ and to form biofilms¹⁵. The QS mechanism can control particular processes related to cell density¹⁶, and QS inhibition targeting autoinducers has been used as a method to control biofilm formation¹⁶. The opportunistic pathogen *Pseudomonas aeruginosa* has four QS systems (Las, Rhl, Pqs, and Iqs)¹⁷. Each system has its own signal and regulatory protein. For the Las system, LasI synthesizes *N*-(3-oxododecanoyl)-homoserine lactone (3oxoC12HSL), and LasR is the protein receptor¹⁷. For the Rhl system, RhlI synthesizes *N*-butyrylhomoserine lactone, and RhlR is the protein receptor. The third QS system is quinolone-based intercellular signaling; the PQS signal is 2-heptyl-3-hydroxy-4-quinolone, and it is synthesized by the products of the PQS synthesis cluster consisting of *pqsABCD*, *phnAB*, and *pqsH*. PqsR is the protein receptor¹⁷. The fourth system is called Iqs, and the QS molecule is 2-(2-

40

hydroxyphenyl)-thiazole-4-carbaldehyde¹⁷.

One aspect of biofilm formation controlled by the Rhl QS system is regulation of the synthesis of rhamnolipids, which are glycolipid biosurfactants composed of rhamnose and 3-(hydroxyalkanoyloxy)alkanoic acid (HAA)¹⁸. *P. aeruginosa* rhamnolipids affect biofilm architecture by participating in the maintenance of biofilm channels¹⁹ and reduce adhesion between cells²⁰.

There are many reasons for forming bacterial biofilms such as a defense against stress (e.g., nutrient deprivation, antibiotics, or pH changes)²¹ and a mechanism for staying stationary in a nutrient-rich area²¹. However, bacteria also must have a means to leave the biofilm (dispersal) due to environmental changes (e.g., fluctuations in oxygen levels, alterations in nutrients, or increasing of toxic products)²²; this process involves breaking the matrix to uncement the cells²³.

Here, we show that supernatants from *P. aeruginosa* PA14 (henceforth PA14) disperse the biofilms of SRB, *E. coli*, and *S. aureus*. Using QS-related mutants, we determined that the biochemical basis for this dispersal is the presence of rhamnolipids in the supernatants.

RESULTS

PA14 wild-type supernatant disperses SRB biofilm. In an effort to investigate whether there are QS compounds utilized by the representative SRB *D. vulgaris*, we tested whether its supernatants would disperse its own biofilm formed in rich medium. *D. vulgaris* supernatant did not disperse its own biofilm within 2 h.

Since there was no negative effect of SRB supernatants on its own biofilm, we investigated the effect of the supernatant of other strains (e.g., *P. aeruginosa*, *P. fluorescens*, *Bacillus subtilis*, *E. coli*). Among these strains, *P. aeruginosa* PA14 and *P. aeruginosa* PAO1 dispersed SRB biofilm the most (**Fig. 1**). The supernatants were obtained from planktonic cultures, and the SRB biofilm grown was in 96-well plates for 24 to 48 h in modified Baar's medium. Critically, the PA14 wild-type supernatant dispersed *D. vulgaris* biofilm more than 92% after 1 to 2 h of incubation. We utilized short periods of contact of the supernatants with the SRB biofilm to avoid artifacts related with growth of the bacterium.

Rhamnolipids in the supernatants disperse SRB biofilms. To determine mechanisms behind these strong dispersal results, we hypothesized that the compounds in the *P. aeruginosa* supernatant may be related to QS since PA14 is the best known strain for QS, and QS controls extracellular compounds like protease²⁴ and rhamnolipids²⁴. Hence, we tested the supernatant of mutants related to QS as well as those
70 related to virulence factors (e.g., *lasI*²⁵, *lasB*²⁶, *pelA*²⁷, *phzM*²⁸, *phzS*²⁸, *pvdF*²⁹, *pchE*³⁰, *lecA*³¹, *toxA*³², *sodM*³³, *rhlI*³⁴, *rhlR*³⁴). Similar to the supernatant of the PA14 wild-type, the supernatants of the mutants *lasI* (autoinducer), *lasB* (elastase), *pelA* (oligogalacturonide lyase), *phzM* (pyocyanin), *phzS* (pyocyanin), *pvdF* (pyoverdine), *pchE* (pyochelin), *lecA* (LecA lectin), *toxA* (exotoxinA), and *sodM* (superoxide
75 dismutase) dispersed the *D. vulgaris* biofilm like the supernatant of PA14 wild-type (**Fig. 2**), indicating the Las QS system, Pel polysaccharide, pyocyanin, pyoverdine, pyocheline, LecA lectin, exotoxinA, superoxide dismutase had no role in the biofilm dispersal.

In contrast to the other mutants, mutations in the rhamnolipids pathway, specifically in the gene encoding the autoinducer, *rhlI*, and the gene encoding the response regulator, *rhlR*, had a pronounced decrease in dispersal (**Fig. 2**). Since the *rhlI* and *rhlR* mutants were found to affect biofilm dispersal,
80 additional mutations in the rhamnolipid pathway were investigated: *rhlA*, which encodes rhamnosyltransferase 1 subunit A and *rhlB*, which encodes the catalytic subunit of the rhamnosyltransferase^{34,35}. The supernatants of both the *rhlA* and *rhlB* mutant did not disperse SRB biofilms (**Fig. 2**). Therefore, the dispersal compounds are related to rhamnolipids.

To corroborate the 96-well biofilm dispersal results with the PA14 wild-type and *rhlB* mutant,
85 confocal microscopy was used to visualize the remaining biofilm after treatment with supernatants from these two strains. As shown in **Fig. 3**, the supernatant from the wild-strain nearly completely dispersed the SRB biofilm whereas the supernatant from the *rhlB* mutant had no effect, just like the buffer negative control.

From the rhamnolipid production pathway (**Fig. 4**), RhlA synthesizes 3-(3-
90 hydroxyalkanoyloxy)alkanoic acids (HAA)³⁶, and RhlB uses HAA to make mono-rhamnolipids³⁷. Therefore, the compounds required for SRB biofilm dispersal are HAA or mono/di-rhamnolipids,

respectively. However, the results from mass spectrometry showed that there was no HAA detected in the wild-type supernatant sample. Therefore we conclude that rhamnolipids are the compounds (and not HAA) that are important for biofilm dispersal.

95 **Rhamnolipids are present in the PA14 supernatants and disperse SRB biofilms.** The genetic results clearly indicate the importance of rhamnolipids for SRB biofilm dispersal via PA14 supernatants. To corroborate these results, the supernatants of PA14, *rhlA*, *rhlB*, *rhlI*, and *rhlR* mutants were tested for the presence of rhamnolipids by mass spectrometry (**Fig. 6**). As expected, rhamnolipids were detected only in the PA14 supernatant samples and not from any of the mutant samples. There were four predominant
100 peaks of rhamnolipids shown (**Fig. 6**) which are Rha-C₁₀-C₁₀, Rha-C₁₀-C₁₂/Rha-C₁₂-C₁₀, Rha-Rha-C₁₀-C₁₀, and Rha-Rha-C₁₀-C₁₂/ Rha-Rha-C₁₂-C₁₀. By comparing with the 10 mM rhamnolipids standard from *P. aeruginosa*, we determined the concentration of rhamnolipids in the PA14 supernatant using the total peak areas is over 100 fold less than the standard (0.1 mM, **Table 1**).

To show conclusively that rhamnolipids in the supernatants are the biochemical means by which the
105 *D. vulgaris* biofilms are dispersed, purified *P. aeruginosa* rhamnolipids (10 mM) were tested. We found that the purified rhamnolipids disperse *D. vulgaris* biofilm with approximately 67% dispersal after 2 h (**Fig. 2**). The rhamnolipids standard did not disperse the biofilm as well as the PA14 supernatant though there were more rhamnolipids in the 10 mM standard sample than the amount of rhamnolipids in the supernatant (**Table 1**). This implies that the ratio of the rhamnolipids or other compounds in the
110 supernatant are important for SRB biofilm dispersal.

Addition of the supernatant of the *rhlB* mutant to the rhamnolipids standard did not increase the biofilm dispersal compared to the the rhamnolipids standard alone. This suggests the dispersal is due to the rhamnolipids alone and not some other product in the supernatants.

PA14 supernatants disperse *D. desulfuricans*, *E. coli* MG1655, and *Staphylococcus aureus* biofilms.

115 To test the ability of the PA14 supernatants to disperse other bacteria, biofilms of *D. desulfuricans*, *E. coli* MG1655, *S. aureus*, and PA14 were grown in 96-well plates, then treated with supernatants from PA14. The PA14 supernatant dispersed the biofilms of *D. desulfuricans*, *E. coli* MG1655, and *S. aureus* well

(over 90%). Although, there are some strain specific differences, these results show that the PA14 supernatants have a great deal of utility for dispersing a wide range of biofilms.

120 As expected, the supernatant from the *rhlR* mutant did not disperse biofilms well (**Fig. 5**), providing further evidence of the importance of rhamnolipids. In addition, the supernatant of PA14 did not disperse its own biofilm (data not shown).

PA14 supernatant disperses SRB biofilm grown in M9G. Since biofilm matrices can change when the growth medium is altered³⁸, we tested the ability of the PA14 supernatant to disperse SRB biofilms grown
125 in minimal glucose medium (M9G). The PA14 supernatant dispersed 90-100% of the SRB biofilm when formed in modified's Baar's medium and dispersed about 80% of the biofilm formed in M9G. In contrast, the positive control protease, which dispersed 80% of the SRB biofilm formed in modified's Baar's medium, dispersed only about 40% of the biofilm grown in M9G.

DISCUSSION

130 We have demonstrated that the supernatants of PA14 disperse *D. vulgaris* biofilms effectively and that the biochemical basis of this dispersal is the presence rhamnolipids. The PA14 supernatants dispersed the biofilm better than protease, and they are versatile because they also dispersed *D. desulfuricans*, *E. coli* MG1655, and *S. aureus* biofilm. This is significant since the matrix of *D. desulfuricans*, *E. coli*, and *S. aureus* are primarily proteins^{12,13}, carbohydrates³⁹, and some extracellular genomic DNA¹⁴,
135 respectively, which indicates that rhamnolipids are a powerful and general approach for removing biofilms.

The rhamnolipid standard did not disperse the biofilm as well as the rhamnolipids in the PA14 WT supernatant, possibly because the ratios of the rhamnolipids are critical for biofilm dispersal activity. For example, the ratio between rhamnolipids and mono rhamnolipids affects the mixture properties (e.g.,
140 foam thickness, surface electric parameters)⁴⁰ and changes the emulsification index and antimicrobial properties⁴¹. In addition, the rhamnolipid standard was not pure (90%), and the growth condition used to produce the rhamnolipids is unknown. We also surmise that purifying the rhamnolipids alters their

biofilm dispersal properties. Furthermore, previously it was reported that rhamnolipids from *P. aeruginosa* together with the QS signal 3oxoC12HSL work cooperatively to disperse *E. coli* biofilms⁴²; however, we found that there was no need for 3oxoC12HSL for supernatants to disperse SRB since the supernatants from the *lasI* mutant were just as effective as those from the wild-type strain (**Fig. 2**).

METHODS

Bacterial strains and culture conditions. *Desulfovibrio vulgaris* and *Desulfovibrio desulfuricans* were grown anaerobically in modified Baar's medium at 30°C. PA14 wild-type and its mutants were grown in lysogeny broth (LB)⁴³ or minimal medium with 0.4% glucose (M9G)⁴⁴ at 37°C. Gentamycin (Gm) 15 µg/mL was used for culturing the PA14 mutants. *Escherichia coli* TG1, *Escherichia coli* MG1655, and *Pseudomonas fluorescens* were grown in LB at 37°C. All the strains are shown in **Table 2**. The PA14 *rhlA*, *rhlB*, *rhlI*, *rhlR*, and *lasI* mutants were verified via PCR using the primers in **Table 3**.

PA14 supernatant preparation. *P. aeruginosa* PA14 wild-type and its isogenic mutants were grown in LB or LB with Gm15 µg/mL for the mutants. The overnight cultures (200 µL) were centrifuged, the cell pellets were resuspended in 50 mL of M9G, and cultured for 3 to 4 days. The supernatants were collected and filtered using a 0.22 µm filtration system. The samples were concentrated from 2x to 4x using a speed vacuum.

Biofilm dispersal assay using crystal violet. Biofilm formation was assayed in 96-well polystyrene plates using 0.1% crystal violet staining as described previously⁴⁵ with some modifications. Diluted overnight cultures (150 µL) at an initial turbidity at 600 nm of 0.05 were inoculated into 96-well-plates with modified Baar's medium, and the bacteria were cultured for 24 to 48 hours at 30°C without shaking. The planktonic cells were removed from each well, and the biofilm was washed gently using phosphate buffered saline (PBS). The supernatant or chemicals were added to the wells, and the plate was incubated at 30°C for 1 to 2 h. Then, the supernatants were removed from each well, and the remaining biofilm was washed gently using water. After the crystal violet was added to each well, the wells were rinsed and dried, and ethanol was added to dissolve the crystal violet. The total biofilm formation samples were

measured at 540 nm. At least three independent cultures were used for each strain.

Confocal Microscopy. Diluted overnight cultures (300 μ L) at an initial turbidity at 600 nm of 0.1 were
170 inoculated into an 8 chamber cell culture slide (Dot Scientific, Inc, Burton, MI) with modified Baar's
medium, and the bacteria were cultured for 48 hours at 30°C without shaking. The planktonic cells were
removed from each well, and the biofilm was washed gently using PBS. Supernatants or PBS was added
to the chambers, and the slide was incubated at 30°C for 20 minutes. Confocal microscopy images were
taken using a PlanApo60 \times /1.4 oil objective lens with Olympus Fluoview 1000 confocal laser microscope.
175 The biofilm structure images were performed using IMARIS software (Bitplane, Zurich, Switzerland),
and average biomass and thickness were obtained using COMSTAT image-processing software⁴⁶. At least
three different areas were observed, and two independent cultures were tested.

Mass spectrometric analysis of supernatants. Mass spectrometric analysis was performed on the
supernatants by using a Waters Q-TOF Premier quadrupole/time-of-flight (TOF) mass spectrometer
180 (Waters Corporation (Micromass Ltd.), Manchester, UK). Operation of the mass spectrometer was
performed using MassLynx™ software Version 4.1 (<http://www.waters.com>). Samples were introduced
into the mass spectrometer using a Waters 2695 high performance liquid chromatograph. The samples
were analyzed using flow injection analysis (FIA), in which the sample is injected into the mobile phase
flow and passes directly into the mass spectrometer, where the analytes are ionized and detected. The
185 mobile phase used was 90% acetonitrile (LC-MS grade), and 10% aqueous 10 mM ammonium acetate.
The flow rate was 0.15 mL/min. The nitrogen drying gas temperature was set to 300 °C at a flow of 7
L/min. The capillary voltage was 2.8 kV. The mass spectrometer was set to scan from 100-1000 m/z in
negative ion mode, using electrospray ionization (ESI). Rhamnolipids standard (90%) was purchased
from Sigma-Aldrich (St. Louis, MO).

190 DATA AVAILABILITY

The authors declare that data supporting the findings of this study are available within the paper.

ACKNOWLEDGMENTS

This work was supported by the Dow Chemical Company and funds derived from the Biotechnology
195 Endowed Professorship at the Pennsylvania State University.

AUTHOR CONTRIBUTIONS

Experiments were designed by TLW, LZ, DSM, BY, and TKW. The experiments were performed by
TLW except for mass spectrometry which was performed by JM. The manuscript was written by TLW
and TKW.

200

COMPETING INTERESTS

The authors declare there are no competing interests.

REFERENCES

1. Zhang, W., Culley, D.E., Nie, L. & Scholten, J.C. Comparative transcriptome analysis of *Desulfovibrio vulgaris* grown in planktonic culture and mature biofilm on a steel surface. *Appl Microbiol Biotechnol* **76**, 447-457 (2007).
2. Pankhania, I.P., Spormann, A.M., Hamilton, W.A. & Thauer, R.K. Lactate Conversion to Acetate, CO₂ and H₂ in Cell Suspensions of *Desulfovibrio Vulgaris* (Marburg): Indications for the Involvement of an Energy Driven Reaction. *Arch Microbiol* **150**, 26-31 (1988).
3. Lee, W., Lewandowski, Z., Nielsen, P.H. & Hamilton, W.A. Role of Sulfate-Reducing Bacteria in Corrosion of Mild-Steel - a Review. *Biofouling* **8**, 165-& (1995).
4. Heidelberg, J.F. et al. The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nature biotechnology* **22**, 554-559 (2004).
5. Clark, M.E., Edelmann, R.E., Duley, M.L., Wall, J.D. & Fields, M.W. Biofilm formation in *Desulfovibrio vulgaris* Hildenborough is dependent upon protein filaments. *Environ Microbiol* **9**, 2844-2854 (2007).
6. Poosarla, V.G. et al. Dispersal and inhibitory roles of mannose, 2-deoxy-d-glucose and N-acetylgalactosaminidase on the biofilm of *Desulfovibrio vulgaris*. *Environ Microbiol Rep* (2017).
7. Faybishenko, B. et al. In Situ Long-Term Reductive Bioimmobilization of Cr(VI) in Groundwater Using Hydrogen Release Compound. *Environ Sci Technol* **42**, 8478-8485 (2008).
8. Korte, H.L. et al. Independence of nitrate and nitrite inhibition of *Desulfovibrio vulgaris* Hildenborough and use of nitrite as a substrate for growth. *Environ Sci Technol* **49**, 924-931 (2015).
9. Lopez, D., Vlamakis, H. & Kolter, R. Biofilms. *Cold Spring Harb Perspect Biol* **2**, a000398 (2010).
10. Mah, T.F. & O'Toole, G.A. Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* **9**, 34-39 (2001).
11. Street, C.N. & Gibbs, A. Eradication of the corrosion-causing bacterial strains *Desulfovibrio vulgaris* and *Desulfovibrio desulfuricans* in planktonic and biofilm form using photodisinfection. *Corros Sci* **52**, 1447-1452 (2010).
12. Hoble, L., Harkins, C., MacPhee, C.E. & Stanley-Wall, N.R. Giving structure to the biofilm matrix: an overview of individual strategies and emerging common themes. *Fems Microbiol Rev* **39**, 649-669 (2015).

13. Foulston, L., Elsholz, A.K.W., DeFrancesco, A.S. & Losick, R. The Extracellular Matrix of *Staphylococcus aureus* Biofilms Comprises Cytoplasmic Proteins That Associate with the Cell Surface in Response to Decreasing pH. *Mbio* **5** (2014).
14. Rice, K.C. et al. The cidA murein hydrolase regulator contributes to DNA release and biofilm development in *Staphylococcus aureus*. *Proc Natl Acad Sci U S A* **104**, 8113-8118 (2007).
15. Dickschat, J.S. Quorum sensing and bacterial biofilms. *Nat Prod Rep* **27**, 343-369 (2010).
16. McLean, R.J.C., Pierson, L.S. & Fuqua, C. A simple screening protocol for the identification of quorum signal antagonists. *J Microbiol Meth* **58**, 351-360 (2004).
17. Lee, J. & Zhang, L.H. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein Cell* **6**, 26-41 (2015).
18. Desai, J.D. & Banat, I.M. Microbial production of surfactants and their commercial potential. *Microbiol Mol Biol Rev* **61**, 47-& (1997).
19. Davey, M.E., Caiazza, N.C. & O'Toole, G.A. Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* **185**, 1027-1036 (2003).
20. Gomes, M.Z.D. & Nitschke, M. Evaluation of rhamnolipid and surfactin to reduce the adhesion and remove biofilms of individual and mixed cultures of food pathogenic bacteria. *Food Control* **25**, 441-447 (2012).
21. Jefferson, K.K. What drives bacteria to produce a biofilm? *FEMS microbiology letters* **236**, 163-173 (2004).
22. Kostakioti, M., Hadjifrangiskou, M. & Hultgren, S.J. Bacterial Biofilms: Development, Dispersal, and Therapeutic Strategies in the Dawn of the Postantibiotic Era. *Csh Perspect Med* **3** (2013).
23. Kaplan, J.B. Biofilm dispersal: mechanisms, clinical Implications, and potential therapeutic uses. *J Dent Res* **89**, 205-218 (2010).
24. Aybey, A. & Demirkan, E. Inhibition of quorum sensing-controlled virulence factors in *Pseudomonas aeruginosa* by human serum paraoxonase. *Journal of medical microbiology* **65**, 105-113 (2016).
25. Storey, D.G., Ujack, E.E., Rabin, H.R. & Mitchell, I. *Pseudomonas aeruginosa* lasR transcription correlates with the transcription of lasA, lasB, and toxA in chronic lung infections associated with cystic fibrosis. *Infect Immun* **66**, 2521-2528 (1998).
26. Liu, P.V. The roles of various fractions of *Pseudomonas aeruginosa* in its pathogenesis. 3. Identity of the lethal toxins produced in vitro and in vivo. *J Infect Dis* **116**, 481-489 (1966).
27. Sakuragi, Y. & Kolter, R. Quorum-sensing regulation of the biofilm matrix genes (pel) of *Pseudomonas aeruginosa*. *J Bacteriol* **189**, 5383-5386 (2007).
28. Lau, G.W., Hassett, D.J., Ran, H. & Kong, F. The role of pyocyanin in *Pseudomonas aeruginosa* infection. *Trends Mol Med* **10**, 599-606 (2004).
29. Meyer, J.M., Neely, A., Stintzi, A., Georges, C. & Holder, I.A. Pyoverdinin is essential for virulence of *Pseudomonas aeruginosa*. *Infect Immun* **64**, 518-523 (1996).
30. Takase, H., Nitani, H., Hoshino, K. & Otani, T. Impact of siderophore production on *Pseudomonas aeruginosa* infections in immunosuppressed mice. *Infect Immun* **68**, 1834-1839 (2000).
31. Winzer, K. et al. The *Pseudomonas aeruginosa* lectins PA-IL and PA-III are controlled by quorum sensing and by RpoS. *J Bacteriol* **182**, 6401-6411 (2000).
32. Gambello, M.J., Kaye, S. & Iglewski, B.H. LasR of *Pseudomonas Aeruginosa* Is a Transcriptional Activator of the Alkaline Protease Gene (Apr) and an Enhancer of Exotoxin-a Expression. *Infect Immun* **61**, 1180-1184 (1993).
33. Hassett, D.J. et al. Quorum sensing in *Pseudomonas aeruginosa* controls expression of catalase and superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. *Mol Microbiol* **34**, 1082-1093 (1999).
34. Ochsner, U.A. & Reiser, J. Autoinducer-mediated regulation of rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* **92**, 6424-6428 (1995).
35. Deziel, E., Lepine, F., Milot, S. & Villemur, R. rhlA is required for the production of a novel biosurfactant promoting swarming motility in *Pseudomonas aeruginosa*: 3-(3-

- hydroxyalkanoxyloxy)alkanoic acids (HAAs), the precursors of rhamnolipids. *Microbiol-Sgm* **149**, 2005-2013 (2003).
36. Rehm, B.H., Mitsky, T.A. & Steinbuchel, A. Role of fatty acid de novo biosynthesis in polyhydroxyalkanoic acid (PHA) and rhamnolipid synthesis by pseudomonads: establishment of the transacylase (PhaG)-mediated pathway for PHA biosynthesis in *Escherichia coli*. *Appl Environ Microbiol* **67**, 3102-3109 (2001).
 37. Abdel-Mawgoud, A.M., Hausmann, R., Lepine, F., Muller, M.M. & Deziel, E. Rhamnolipids: Detection, Analysis, Biosynthesis, Genetic Regulation, and Bioengineering of Production. *Microbiol Monogr* **20**, 13-55 (2011).
 38. Flemming, H.C. & Wingender, J. The biofilm matrix. *Nature Reviews Microbiology* **8**, 623-633 (2010).
 39. Beech, I.B., Gaylarde, C.C., Smith, J.J. & Geesey, G.G. Extracellular Polysaccharides from *Desulfovibrio desulfuricans* and *Pseudomonas fluorescens* in the Presence of Mild and Stainless-Steel. *Appl Microbiol Biotechnol* **35**, 65-71 (1991).
 40. Cohen, R. & Exerowa, D. Surface forces and properties of foam films from rhamnolipid biosurfactants. *Adv Colloid Interfac* **134-35**, 24-34 (2007).
 41. Das, P., Yang, X.P. & Ma, L.Y.Z. Analysis of biosurfactants from industrially viable *Pseudomonas* strain isolated from crude oil suggests how rhamnolipids congeners affect emulsification property and antimicrobial activity. *Frontiers in microbiology* **5** (2014).
 42. Bhattacharjee, A., Nusca, T.D. & Hochbaum, A.I. Rhamnolipids Mediate an Interspecies Biofilm Dispersal Signaling Pathway. *ACS chemical biology* **11**, 3068-3076 (2016).
 43. Sambrook, J.F. & Russell, D.W. Molecular cloning: a laboratory manual, Edn. 3rd. (Cold Spring Harbor Laboratory Press, New York; 2001).
 44. Sambrook, J., Fritsch, E.F. & Maniatis, T. Molecular Cloning, A Laboratory Manual, Edn. 2. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; 1989).
 45. Fletcher, M. The effects of culture concentration and age, time, and temperature on bacterial attachment to polystyrene. *Can J Microbiol* **23**, 1-6 (1977).
 46. Heydorn, A. et al. Quantification of biofilm structures by the novel computer program COMSTAT. *Microbiol-Sgm* **146**, 2395-2407 (2000).
 47. Dusane, D.H. et al. Quorum sensing: implications on rhamnolipid biosurfactant production. *Biotechnol Genet Eng Rev* **27**, 159-184 (2010).
 48. Choi, M.H. et al. Metabolic relationship between polyhydroxyalkanoic acid and rhamnolipid synthesis in *Pseudomonas aeruginosa*: Comparative C-13 NMR analysis of the products in wild-type and mutants. *J Biotechnol* **151**, 30-42 (2011).
 49. Wittgens, A. et al. Novel insights into biosynthesis and uptake of rhamnolipids and their precursors. *Appl Microbiol Biotechnol* **101**, 2865-2878 (2017).
 50. Manner, S., Goeres, D.M., Skogman, M., Vuorela, P. & Fallarero, A. Prevention of *Staphylococcus aureus* biofilm formation by antibiotics in 96-Microtiter Well Plates and Drip Flow Reactors: critical factors influencing outcomes. *Scientific reports* **7** (2017).
 51. Deziel, E., Lepine, F., Milot, S. & Villemur, R. Mass spectrometry monitoring of rhamnolipids from a growing culture of *Pseudomonas aeruginosa* strain 57RP. *Bba-Mol Cell Biol L* **1485**, 145-152 (2000).
 52. Liberati, N.T. et al. An ordered, nonredundant library of *Pseudomonas aeruginosa* strain PA14 transposon insertion mutants. *Proc Natl Acad Sci U S A* **103**, 2833-2838 (2006).
 53. Price, K.E., Naimie, A.A., Griffin, E.F., Bay, C. & O'Toole, G.A. Tobramycin-Treated *Pseudomonas aeruginosa* PA14 Enhances *Streptococcus constellatus* 7155 Biofilm Formation in a Cystic Fibrosis Model System. *J Bacteriol* **198**, 237-247 (2015).

FIGURE LEGENDS

- Fig. 1. *D. vulgaris* biofilm dispersal by supernatants of *P. aeruginosa* PA14, *P. aeruginosa* PAO1, *B. subtilis*, *P. fluorescens*, and *E. coli* TG1.** *D. vulgaris* biofilms were grown for 2 days in modified Baar's media at 30 °C, and supernatants were concentrated to 4x and contacted with *D. vulgaris* biofilms for 2 h. * indicates no dispersal.
- Fig. 2. *D. vulgaris* biofilm dispersal by wild-type *P. aeruginosa* PA14 and its quorum sensing mutants.** *D. vulgaris* biofilms were grown for 2 days in modified Baar's media at 30 °C, and supernatants were concentrated to 4x and contacted with *D. vulgaris* biofilms for 2 h. Rhamnolipid standards were added at 10 mM. Protease 1 (Savinase) at 0.024 U was used as a positive control.
- Fig. 3. Representative images of *D. vulgaris* biofilm dispersal via *P. aeruginosa* supernatants as visualized by confocal microscopy.** *D. vulgaris* biofilms were formed on an 8 chamber cell culture slide for 48 h in modified Baar's medium, and supernatants were concentrated to 4x and contacted with *D. vulgaris* biofilms for 0.33 h. Supernatants of (A) *P. aeruginosa* PA14, (B) *P. aeruginosa* PA14 *rhlA*, and (C) PBS buffer. Scale bar is 50 μm.
- Fig. 4. Simplified rhamnolipid biosynthesis pathway in *P. aeruginosa*^{47, 48}.** RhIR binds with *N*-butanoyl-L-homoserine lactone (C4-HSL) produced by RhII to form a RhIR –C4-HSL complex. The RhIR –C4-HSL complex interacts with the *rhlA* promoter to initiate transcription of the *rhlAB* genes to produce rhamnolipids. HAA is 3-(hydroxyalkanoyloxy)alkanoic acid. *P. aeruginosa* commonly produces rhamnolipids containing fatty acids with chain lengths between C₈ and C₁₄ (n = 1-7)⁴⁹.
- Fig. 5. Supernatants of *P. aeruginosa* PA14 disperse biofilms of (A) *D. desulfuricans*, (B) *E. coli* MG1655 biofilm, and (C) *S. aureus*.** The biofilms of *D. desulfuricans* and *E. coli* MG1655 were grown for 24 h in modified Baar's medium and LB, respectively, and the biofilms of *S. aureus* were grown for 24 h in TSB⁵⁰, and *P. aeruginosa* supernatants were concentrated to 4x and contacted the biofilms for 2 h except for *S. aureus* biofilm which the 1x supernatant was used instead of 4x and the contact time was 10 min.
- Fig. 6. Detection of rhamnolipids in the *P. aeruginosa* PA14 supernatants via mass spectrometry.** *P. aeruginosa* wild-type and the *rhlA* mutant were grown planktonically for 4 days in M9G medium at 37 °C.

Table 1. Rhamnolipids in *P. aeruginosa* PA14 supernatants and in the 10 mM rhamnolipid standard. Peak areas are indicated from the mass spectrometry chromatograms.

Rhamnolipids⁵¹	Ion (m/z)	PA 14 WT	Standard
Rha-C ₁₀ -C ₁₀	503.3	31	6374
Rha-C ₁₀ -C ₁₀ /Rha-C ₁₂ -C ₁₀	531.3	3	294
Rha-Rha-C ₁₀ -C ₁₀	649.3	37	3465
Rha-C ₁₀ -C ₁₂ /Rha-C ₁₂ -C ₁₀	677.3	8	178

Table 2. Bacterial strains used in this study.

Strains	Description	Source
Strains		
<i>E. coli</i> TG1	<i>supE thi-1 Δ(lac-proAB) Δ(mcrB-hsdSM)5, (r_K⁻m_K⁻)</i> F' [<i>traD36 proAB⁺ lacI^f lacZΔM15</i>]	Jeremy Minshull
<i>E. coli</i> MG1655	wild-type	Frederick Blattner
<i>D. vulgaris</i>	wild-type	ATCC 29579
<i>D. desulfuricans</i>	wild-type	DSM 12129
<i>P. aeruginosa</i> PAO1	wild-type	Tim McDermott
<i>P. aeruginosa</i> PA14	ATCC 29579, wild-type	Frederick Ausubel
<i>P. aeruginosa</i> PA14 Δ <i>lasI</i>	PA14_45940, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>lasB</i>	PA14_16250, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>pelA</i>	PA14_24480, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>phzM</i>	PA14_09490, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>phzS</i>	PA14_09400, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>pvdF</i>	PA14_33700, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>pchE</i>	PA14_09270, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>lecA</i>	PA14_31290, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>toxA</i>	PA14_49560, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>sodM</i>	PA14_58000, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>rhlA</i>	<i>rhlA::Gm, Gm^R</i>	53
<i>P. aeruginosa</i> PA14 Δ <i>rhlB</i>	PA14_19110, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>rhlI</i>	PA14_19130, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. aeruginosa</i> PA14 Δ <i>rhlR</i>	PA14_19120, Mar2xT7 transposon insertion, Gm ^R	52
<i>P. fluorescens</i>	wild-type	Richard Frazee
<i>S. aureus</i>	ATCC 29213, wild-type	Kenneth Urish

Table 3. Primers used in this study. Forward primers are labeled with F, and rear primers are labeled with R.

Name	Sequence
rhIA F2	ATGGCCGCTGAGTTACTTGTCT
rhIA R2	CAGGCCGATGAAGGGAAATACGT
rhIB F	AAGACCCGGCCTGGCCGGGC
rhIB R	ATTGCAGTAAGCCCTGATCG
rhII F	AGAGGGCCCAGGAGTATCA
rhII R	GGCTGACGACCTCACACC
rhIR F	GTGATGCATTTTATCGATC
rhIR R	AACCTGCCAGATCTGGTAGGGC
lasI F	TGGAGGAAGTGAAGATGATCGT
lasI R	ACAGGTCCCCGTCATGAAAC




Figure 1




Figure 2




Supernatant	Figure	Total biomass ($\mu\text{m}^3/\mu\text{m}^2$)	Average thickness (μm)
A. PA14		1.33×10^{-5}	0
B. PA14 <i>rhlA</i>		1.74	3.21
C. Buffer		0.952	1.30

Figure 3






Figure 4

A. *D. desulfuricans* biofilm



B. *E. coli* biofilm



C. *S. aureus* biofilm




Figure 5




Figure 6