Analysing progress of SDG 6 in India: Past, Present, and Future

Ajishnu Roy[1], Kousik Pramanick*

Integrative Biology Research Unit (IBRU), Department of Life Sciences, Presidency

University, 86/1 College Street Kolkata – 700076

*Corresponding author

Email: kousik.dbs@presiuniv.ac.in

Abstract:

Human endeavors to meet social and economic water needs at national scale might cause

negative environmental manifestations and water stress from local to global scale. So,

appropriation of Sustainable Development Goals requires a comprehensive monitoring and

knowledge base of the water resource availability, utilization and access. Hence, scientific

research progression has a significant role to facilitate the implementation of sustainable

development goals through assessment and policy implementation from global to local scales.

India holds a key position among developing economies with a complex interconnected web

of a fast-growing population, coupled with biophysical stress, social deprivation and

economic inequality related to water and sanitation. This study addresses some of these

challenges related to monitoring and implementation of the targets of the United Nations

Sustainable Development Goal 6 in India. Acknowledging the contribution of society and

economy in sustainability paradigm, here we have chosen 28 indicators (clustered into eleven

dimensions) under two major groups, concerning biophysical and social development aspects

of water and sanitation. We have shown declining level of per capita biophysical water

resource and slow to rapidly developing social indicators related to Sustainable Development

Goal 6 in India. From past trends, we have calculated probable scenario of biophysical

consumption of India up to 2050, which shows at least 1.3 times increase. This cumulative

assessment framework contributes a tool to prioritize water resource appropriation,

management response and policy implementations to national level sustainability of water

and sanitation in India. We also advocate the necessity of restraining threats both at source

and consumption process levels in order to ensure national water security for both human and

biodiversity, keeping in mind the societal and economic development scenario.

Keywords: SDG 6; indicators; national water security; water and sanitation; India.

1. Introduction:

India is going through a time of deterioration of the environment, social deprivation and ineffective economy, combined with the rapidly growing population. This is an example of global scale initiation and progress of 'Anthropocene' (Steffen et al. 2007, 2011). During the 1980s, the concept of sustainable development was framed, which is "development that meets the needs of the present without compromising the ability of future generations to meet their own needs" (Brundtland Commission Report, 1987). In 1992 United Nations Conference on Environment and Development (UNCED, Rio de Janeiro Earth Summit), Agenda 21, calls for sustainable development indicators (SDIs) to "provide solid bases for decision-making at all levels and to contribute to a self-regulating sustainability of integrated environment and development system". Emerging from Millennium Development Goals (MDGs), United Nations has set 17 Sustainable Development Goals (SDGs) and 169 targets in 2015. SDGs incorporate three pillars of sustainable development, i.e. environmental (climate action, life below water, life on land etc.), social (zero hunger, no poverty, gender equality, peace and justice and strong institutions etc.) and economic development (reduced inequalities, decent work and economic growth etc.) (Sachs, 2012).

There is a distinct trend that was set from the 1970s, especially from 1990s, that led to the contextual understanding and necessity for this study. It can be divided into two phases.

In the first phase, the significance of managing water resource sustainably was studied and advocated. Almost all of the works, concerned only with environmental aspect of water resource management. From the 1990s, the significance of the sustainable development of water resources has been increasing steadily. Biswas (1991, 1992) emphasised an analytical framework for drivers of the water crisis, socio-environmental considerations along with institutional responses for better management. Long before the present time, Serageldin (1995) had proposed 'water resource management' as a policy tool for sustainable future.

Gleick (1996) recommended that international organizations, national and local governments,

with water providers adopt a 'basic water requirement' standard for human needs - 50 litres

per person per day and promise access to this independently of an individual's economic,

social, or political status. In another work, Gleick (1998) proposed seven 'sustainability

criteria' related to "basic water requirement (BWR)" to restructure long-term water planning.

These criteria included – basic water requirement for the need and health of both human and

ecosystems, water quality monitoring, institutional mechanism for water-related conflict

resolution, water planning and decision making etc. Postel (2000) advocated a new water

management paradigm which balances both ecological and economic functions of water by

increasing water productivity (performing more functions with less water). Vörösmarty et al.

(2000) have assessed the vulnerability of global water resource from both climate change and

population growth. Jackson et al. (2001) analysed changes in water resources and projected

future scenario for the USA. Vörösmarty et al. (2010) presented a global synthesis that jointly

considers both human and biodiversity perspective on water security.

In the second phase, both economic and societal aspects started to get their due importance in

important studies on sustainability analysis. In 2009, Rockström et al. devised a framework to

understand if anthropogenic biophysical resource consumption has already exceeded or is

going to surpass safe limits – 'planetary boundaries'. This work marked the global onset of

biophysical resource consumption monitoring in recent years. Then Steffen et al (2015)

modified and improved the framework. One of the main problems in this work is that the

indicators used in these works had not been corroborated completely with UN sustainable

development goals. Also, the socioeconomic dimension was not comprehensively discussed.

This was accomplished by Raworth (2012) through the incorporation of the social deprivation

indicators. This acted as a complementary to planetary boundary framework and together

formed 'doughnut economy' that touched all three pillars of sustainability – environment,

economy and society. Water and sanitation are strongly related to public health too. Works in

this area have also increased (Bartram et al, 2005, Moe and Rheingans, 2006, Montgomery

and Elimelech, 2007, Bartram and Cairncross, 2010, Bartram et al. 2014). There has been an

important discussion ongoing that specifically signifies water and sustainable development is

perceived from this water-perspective (Bogardi et al. 2012, Madrid et al. 2013, Bhaduri et al.

2016, Ait-Kadi, 2016).

There has been some work on the appropriation of water resources and its management in

India (Kumar et al. 2005, Shah and Koppen, 2006, Mall et al, 2006, Mujumdar, 2008). An

important lacuna is that none of these incorporate socioeconomic dimensions in sustainability

analysis of water resource. All of these studies logically imply the need for a national study

on the sustainable development of water and sanitation that includes environment, society

and economy together. That's why this study was performed.

There is no national level study present focusing on both biophysical and socioeconomic

aspects of SDG 6 i.e. sustainable development of water and sanitation. In this work, we have

tried to find answers to the following: (1) How can we downscale SDG 6 indicators to a

national scale more accurately? (2) How can we understand changes in dimensions of SDG 6

with time in order to contextualize their present values? (3) How can we use the past trends in

per capita biophysical consumptions under SDG 6 to project probable future biophysical

resource consumption at a national scale? This analysis measures the national performance of

India on SDG 6 through11dimensions, 28 indicators, segregated into 2 groups, biophysical

and social development indicators, provides important findings of the relationship between

biophysical resource use and well-being for India. Our work has been explained herein few

steps. We first present our methodology, then results of our study on India; project probable

future scenario of biophysical consumption for India; discuss limitations and scopes of this

study, the applicability of SDG 6 as a tool in policymaking and necessities for future

improvements in research.

2. Data and Method:

We collected the data related to SDG 6 of India from Aquastat, FAOSTAT and World

development indicators (world bank). As 3 main pillars of sustainable development are the

environment, society, and economy, we also clubbed all SDG 6 associated indicators into 2

broad categories – biophysical and social development indicators (Fig. 1.), following

planetary boundaries framework (Rockström et al, 2009; Steffen et al., 2015) and doughnut

economics (Raworth, 2012, 2017).

Fig. 1. Changes in biophysical and social development indicators related to sustainable

development goal 6 in India.

A and B represent the status of sustainable development of India in 2000 and 2015

respectively. Eleven indicators of biophysical stress established over biophysical ceiling

(outer chocolate ring) outwardly projected and seventeen indicators of social development

established under social foundation (inner chocolate ring) inwardly projected for India.

Biophysical indicators are – total water withdrawal (TWW), agricultural water withdrawal

(AWW), industrial water withdrawal (IWW), municipal water withdrawal (MWW), total

internal renewable water resource (TIRWR), total renewable water resource (TRWR), dam

capacity (DC), water stress (WS), produced amount of municipal wastewater (PMWW),

treated municipal wastewater (TMWW) and non-treated municipal wastewater (NTMWW).

Indicators of social development are – (1) Improved sanitation facilities, (2) Improved

sanitation facilities in rural areas, (3) Improved sanitation facilities in urban areas, (4)

Improved water source, (5) Improved water source in rural areas, (6) Improved water source

in urban areas, (7) People practicing open defecation, (8) Rural people practicing open

defecation, (9) Urban people practicing open defecation, (10) People using at least basic

drinking water services, (11) Rural people using at least basic drinking water services, (12)

Urban people using at least basic drinking water services, (13) People using at least basic

sanitation services, (14) Rural people using at least basic sanitation services, (15) Urban

people using at least basic sanitation services, (16) Rural people using safely managed

drinking water services, (17) Water use efficiency in agriculture.

Green indicates safe operating space for biophysical indicators and thresholds for indicators

of social development. Yellow indicates the zone of increasing impact for biophysical

indicators and zone of increasing deprivation for indicators of social development. Red

indicates the zone of high risk of serious impact for biophysical indicators and zone of high

level of deprivation for indicators of social development. Light blue represents indicators

without any boundary or thresholds (Unit – m^3). Violet represents water stress (Unit - %).

The area between the chocolate rings is the safe and just operating space for the sustainable

development of water and sanitation in India.

2.1. Biophysical Indicators:

We analyzed 11 indicators under 4 dimensions that indicate biophysical aspects of SDG 6

(Table 1). These 4 dimensions of biophysical indicators are - water withdrawal (4 indicators),

water availability (3 indicators), wastewater (3 indicators) and water stress, WS (1 indicator).

We calculated every biophysical indicator on per capita basis by dividing total values with

population data of India, available from FAOSTAT database (except – water stress,

expressed in percentage, %). We tried to include indicators for water pollution, water

scarcity, the condition of water-related ecosystems, water footprint etc. Either no

unanimously accepted indicator is available or there is no established database with annual

data for the significantly longer duration. So, we could not include any indicator representing

these dimensions.

No.	Dimension (related SDG target)	Indicators		Boundary	Year	Current Status	Change since initial year (+ increase, - decrease)	Data Source
1	Water withdrawal (6.1)	i.	Total water withdrawal per capita (m^3)	576.96 m^3 (2010)	1975-2010	602.3 m^3 (2010)	+3.07%	Aquastat
		ii.	Industrial water withdrawal per capita	576.96 m^3 (2010)	1975-2010	13.63 m^3 (2010)	-42.97%	Aquastat

			(m^3)				
		iii.	Agricultural water withdrawal per capita (m^3)	576.96 m^3 (2010)	1975-2010	551.62 m^3 (2010)	-0.76%	Aquastat
		iv.	Municipal water withdrawal per capita (m^3)	576.96 m^3 (2010)	1975-2010	44.9 m^3 (2010)	+60.06%	Aquastat
2	Water availability (6.1)	v.	Total internal renewable water resources per capita (m^3)	Boundary not available.	1962-2014	1103 m^3 (2014)	-64.29%	Aquastat
		vi.	Total renewable water resources per capita (m^3)	Boundary not available.	1962-2014	1458 m^3 (2014)	-64.29%	Aquastat
		vii.	Dam	Boundary	1960-	189.9	+38.12%	Aquastat

					2005	m^3 (2005)		
			capacity per capita (m^3)	not available.	2005	m^3 (2005)		
3	Wastewater (6.3.1)	viii.	Produced municipal wastewater per capita (m^3)	Boundary not available.	1979-2011	12.23% (2011)	+64.68%	Aquastat
		ix.	Treated municipal wastewater per capita (m^3)	Boundary not available.	1979-2011	3.49% (2011)	+58.86%	Aquastat
		x.	Not treated municipal wastewater per capita (m^3)	Boundary not available.	1979-2011	8.73% (2011)	+67.58%	Aquastat
4	Water Stress (6.4.2)	xi.	Water stress (%)	Boundary not available.	1979-2011	44.53% (2011)	+18.4%	Aquastat

Table 1. Biophysical indicators related to SDG 6 for India

Values for indicators that are within per capita global average freshwater use PB are shown in

green. Those have crossed the boundary, are shown in red.

2.2. Social development Indicators:

We analysed 17 indicatorsunder7 dimensions that indicate social development aspects related

to SDG 6 (Table 2). These 7 dimensions of social development indicators are - open

defecation (3 indicators), basic drinking water (3 indicators), basic sanitation services (3

indicators), safely managed drinking water services (1 indicator), improved sanitation

services (3 indicators), improved water source (3 indicators) and water use efficiency, WUE

(1 indicator). We have also set a desirable threshold for social development indicators to get a

better understanding of gaps between current status and desired status that can be reasonably

accepted as the well-developed situation of SDG 6 for India. These are – less than 10%

people for open defecation and 90% or more people for rest of 5 dimensions (excluding water

use efficiency, WUE). We have also calculated the approximate time when desired thresholds

were or will be met for social development indicators (excluding water use efficiency, WUE)

using linear interpolation at business-as-usual, BAU scenario. The details of each indicator,

along with duration, current status, change, threshold/boundary meeting time etc. are

provided in Table 1 (biophysical indicators of SDG 6) and Table 2 (social development

indicators of SDG 6).

No.	Dimensions		Indicators	Threshold	Year	Current Status	Change since initial year (+ increase, - decrease)	Data Source	Threshold meeting time (BAU)
1	Open defeca	i.	People practicing	Less than	2000 -	39.83 %	- 26.14%	World bank	2030-2035

	tion (6.2.1)		open defecation (% of population)	10% people	2015	(2015)			
		ii.	People practicing open defecation, rural (% of rural population)	Less than 10% people	2000 - 2015	55.62% (2015)	- 26.59%	World bank	2035-2045
		iii.	People practicing open defecation, urban (% of urban population)	Less than 10% people	2000 - 2015	7.41% (2015)	- 16.08%	World bank	2013
2	Basic drinking water	iv.	People using at least basic drinking	90% or more people	2000 - 2015	87.86 % (2015)	+7.15%	World bank	2020

	(6.1.1)		water services (% of population)						
		v.	People using at least basic drinking water services, rural (% of rural population)	90% or more people	2000 - 2015	84.98 % (2015)	+9.16%	World bank	2020-2025
		vi.	People using at least basic drinking water services, urban (% of urban population)	90% or more people	2000 - 2015	92.85 % (2015)	+0.44%	World bank	Before 2000

3	Basic sanitation services (6.2.1)	vii.	People using at least basic sanitation services (% of population)	90% or more people	2000 - 2015	44.15 % (2015)	+22.47 %	World bank	2045
		viii.	People using at least basic sanitation services, rural (% of rural population)	90% or more people	2000 - 2015	33.8% (2015)	+23.27 %	World bank	2050-2055
		ix.	People using at least basic sanitation services, urban (% of urban populatio	90% or more people	2000 - 2015	65.4% (2015)	+14.59 %	World bank	2040

						n)		2045
4	Safely managed drinking water services (6.1.1)	x.	People using safely managed drinking water services, rural (% of rural population)	90% or more people	2000 - 2015	49.47 % (2015)	+20.19 %	World bank	2045
5	Improved Sanitation services (6.2.1)	xi.	Improved sanitation facilities (% of population with access)	90% or more people	1990 - 2015	39.6% (2015)	+22.8%	World bank	2065-2070
		xii.	Improved sanitation facilities, rural (% of rural populatio	90% or more people	1990 - 2015	28.5% (2015)	+22.9%	World bank	2075-2080

			n with access)					**2060-2065**
		xiii.	Improved sanitation facilities, urban (% of urban populatio n with access)	90% or more people	1990 - 2015	62.6% (2015)	+13.3%	World bank	**2060-2065**
6	Impro ved water source (6.1.1)	xiv.	Improved water source (% of populatio n with access)	90% or more people	1990 - 2015	94.1% (2015)	+23.6%	World bank	2010
		xv.	Improved water source, rural (% of rural populatio n with access)	90% or more people	1990 - 2015	92.6% (2015)	+28.4%	World bank	2012

		xvi.	Improved water source, urban (% of urban population with access)	90% or more people	1990 - 2015	97.1% (2015)	+8.2%	World bank	1994
7	Water use efficiency (6.4.1)	xvii.	Water use efficiency, agriculture (water withdrawal, m^3/ GDP, current US$)	Threshold not available	1975 - 2010	2.2 m^3/ GDP, current US$ (2010)	- 76.16%	Aquastat, World bank	

Table 2. Social development indicators related to SDG 6 for India

Indicators that are going to meet threshold within UN SGD target time (2030) are shown in green. Those going meet few years after that time are shown in yellow and indicators, which are going to meet desired threshold many years after 2030 are shown in red.

2.3. Planetary boundary of freshwater use:

According to Rockström et al. (2009), the planetary boundary of freshwater use is the maximum withdrawal of 4000 km^3 y^{-1} blue water from rivers, lakes, reservoirs, and

renewable groundwater stores. We divided it with world population (from World Bank) to get

per capita global scale boundary of 576.96km^3 y^{-1} (2010).

2.4. Future scenario:

As we have calculated most of the biophysical indicators on per capita basis (except – water

stress, in %), it is possible to project probable future scenario of total consumption. We

collected future population projection (2015-2050) data (median range prediction value of

50%) of India from UN DESA (2017 Revision) and then multiplied it with 10 per capita

consumption indicators of SDG 6. We have calculated three projection series for each of 10

indicators of SDG 6, (i) with the lowest value that has happened in past year, (ii) highest

value that has happened in past year and (iii) business-as-usual, BAU scenario with latest

available data.

3. Results and Interpretation:

3.1. Biophysical Indicators:

Among the 11 indicators concerning biophysical aspects of SDG 6, 4 have decreased and 7

have increased over time (Table 1).

Both total internal renewable water resource and total renewable water resource have

significantly decreased (64.29%) which is expected, as the population is growing but the

available amount of renewable water resource is not. In future, with more population growth,

per capita available renewable water resource is likely to decrease further. This problem can

be tackled in 2 ways, (1) control of population growth in India and (2) increasing water

productivity, i.e. perform more functions with less amount of water, in another word,

becoming more water-efficient. Industrial water withdrawal has decreased significantly

(42.97%) that might prove to be good, especially for India. More water will be available for

agriculture (India has an agriculture-based economy) and domestic consumption (India is

going to be one of the highest populated countries in the world). Per capita agricultural water

withdrawal has decreased very less (0.76%) during the time (1975-2010), yet the population

has increased much more i.e. total amount of agricultural water withdrawal have increased. It

means water productivity in agriculture has not increased significantly in India. Therefore,

increasing water productivity in agriculture i.e. growing more agricultural products using less

water (direct or indirect use) should be one of the future concerns.

Total water withdrawal per capita have increased (3.07%). As population will increase in

future, at this rate of increasing water withdrawal water scarcity will be more severe unless

proper mitigation strategies are adopted. Municipal water withdrawal has also increased

significantly (60.06%) which is probably due to increasing number of urban population.

Thus, in 2050, if 50% of Indian population become urban dwelling (world urban prospects,

UN-DESA, 2014 revision), this is likely to be continued, might even at higher rate. As a

result, in future, it might not be possible to provide sufficient water for drinking and

sanitation equitably to all urban people. Bringing down indiscriminate use of water in

municipality areas is one of the probable solutions. If situation arise to more severe stage,

water tax might be considered for implementation, especially for high-income urban

localities. It is also a good time to start thinking about possible sources of freshwater in and

around urban areas for supply to increased urban population in future. Continuous monitoring

to check misuse of municipality provided water is needed. For domestic use in 2050, India

would require 111 billion m^3 water. Top 6 domestic water states (more than 8 billion m^3

water) would be – Uttar Pradesh, Bihar, Maharashtra, West Bengal, Madhya Pradesh and

Andhra Pradesh (IndiaStat 2018). (Supplementary Fig. 1). Total annual water requirement is

increasing steadily in India. In 2050, water requirement is going to be almost three times the

requirement of 1990. Water demand for power generation is going to increase more in 2025

and 2050 (IndiaStat 2018) (Supplementary Fig. 2). Dam capacity per capita have increased

(38.12%). It means, as population is increasing, more numerous and larger capacity dams are

being prepared or going to be needed to prepare in future to maintain this per capita rate. In

one hand, more hydroelectric power is necessary for growing Indian population, in another

hand, adverse effects of present and future construction of dams on aquatic life, especially

riverine biota should be kept in mind (LeRoy Poff et al. 2007, Vörösmarty et al. 2010). Not

only aquatic biodiversity is major source of livelihood of millions of people around river

basins, but also protection and conservation of aquatic biodiversity is also priority. Produced

and treated municipal wastewater has increased significantly (64.68% and 58.86%,

respectively). But not treated municipal wastewater has increased more (67.58%). It means

more amount of wastewater is remaining untreated than treated portion which brings bad

effects of wastewater in environment, soil, biodiversity etc. along with it. More wastewater

treatment plants or centres accompanied with more recycle and reuse of water are necessary

in future. Water stress has also increased (18.4%). Current water stress level in India

(44.53%) is almost 4 times more than average global level of water stress (12.82%). This

indicates water stress is steadily increasing and would become more complex problem in

future with growing number of Indian population. Changes of biophysical consumption

indicators are depicted in Fig 2.

Fig. 2. Changes in per capita biophysical indicators (1-5) and indicators of social development (6-12) related to sustainable development goal 6 in India with time.

Green indicates global per capita boundaries for biophysical indicators and thresholds for indicators of social development.

Biophysical indicators are - (1) water withdrawal (m^3), (2) water resources (m^3), (3) dam capacity (m^3), (4) municipal wastewater ((m^3) and (5) water stress (%). Indicators for social development are - (6) open defecation (%), (7) basic drinking water services (%), (8) basic sanitation services (%), (9) safely managed drinking water services (%), (10) improved sanitation facilities (%) (11) improved water source (%) and (12) water use efficiency in agriculture (m^3/GDP, current US$).

3.2. Social development Indicators:

Of the 17 indicators of social development aspects of SDG 6, 4 have decreased and 13 have increased over time (Table 2).

Open defecation rate in India has decreased in all 3 cases, overall (26.14%), rural (26.59%)

and urban (16.08%). Urban open defecation rate has already met with the desired threshold

(10% or less) in 2013. Although the success, it should be also kept in mind that with

increasing number of urban populous in coming decades, maintaining this rate might prove to

be tougher. So, necessary actions should be made parallelly for implementation in need be in

future. Remaining two will not be able to meet SDG criteria at this rate within 2030 (2030-35

and 2035-40, respectively). It means open defecation in rural areas needs to decrease more.

There are 2 possible ways for that, (1) more public and household toilets are to be constructed

in rural areas and (2) spreading awareness about adverse effects of open defecation on

environment and health in rural areas. On average, 35.09% villages in India are open-

defecation free as of 2017. Four are lowest (0%) among them (Dadra and Nagar Haveli,

Daman and Diu, Goa and Puducherry) and five have highest (100%) achievement (Haryana,

Himachal Pradesh, Kerala, Sikkim and Uttarakhand) (Supplementary Fig. 3) (IndiaStat

2018). Water use efficiency in agriculture has also decreased significantly (76.16%) which is

good, i.e. less amount of water is being withdrawn for the same amount of GDP (in current

US$, 2010).

Basic drinking water services has also increased (overall, 7.15%; rural, 9.16% and urban,

0.44%). Basic drinking water in urban areas has already met with a desirable threshold (90%

or more) before 2000. Remaining two will also be able to meet SDG criteria at this rate

within 2030 (2020 and 2020-25, respectively). Basic sanitation services have increased in

India (overall, 22.47%; rural, 23.27% and urban, 14.59%). But, none of the 3 will be able to

meet desired SDG criteria within 2030 (2045, 2050-2055 and 2040 respectively). Immediate

steps are needed to improve awareness and basic sanitation system. Safely managed drinking

water services in rural Indian population has also increased over time (20.19%). But it will

also not be able to meet desired SDG threshold within 2030 (2045). Improved sanitation

facilities have also increased over time (overall, 22.8%; rural, 22.9% and urban, 13.3%). As none of the 3 indicators will be able to meet desired SDG criteria within 2030 (2065-2070, 2075-2080 and 2060-2065 respectively), improvement of sanitation services requires serious attention in India (both rural and urban area). Improved water source has increased over time (overall, 23.6%; rural, 28.4% and urban, 8.2%). All of these 3 indicators have already meet desired SDG criteria (2010, 2012 and 1994 respectively). To sum up, in India, among 16 indicators with desirable thresholds, 7 have either already met or going to meet SDG criteria within 2030, 1 indicator will reach the target just after 2030, and remaining 8, at current rate, will be able to meet in distant future (2040-2080), requiring immediate attention. Changes in social development indicators over time are seen in Fig 2.

3.3. Planetary boundary of freshwater use at the national scale:

From this analysis, we can see that Indian per capita total water withdrawal (2010) has already crossed global average freshwater use PB (576.96 m^3, 2010) (Fig. 1). This means that with growing population it would be tougher for to achieve UN SDG 6 targets remaining under safe biophysical limits of water resource use.

3.4. Future of SDG 6:

Fig 3 shows total amount of probable biophysical consumption up to 2050 in 9 graphs. In 2050, if lowest per capita rate can be maintained, total water withdrawal would increase 18.98%, at a business-as-usual rate, it would increase 25.8% and at highest per capita rate, it would increase 26.57%. It means even with grown population level of 2050, it is possible to lower the rate of total water withdrawal 6.8-7.5% and which is a significant amount. In agricultural water withdrawal in 2050, at lowest per capita rate, it would increase 20.98%, at a business-as-usual rate, it would increase 25.8% and at highest per capita rate, it would increase 29.1%. Agricultural water withdrawal might to be 1.4 times more in 2050 than 2010. For industrial water withdrawal, in 2050, at lowest per capita rate, it would decrease 7.72%

which is a very positive scenario. But at a business-as-usual rate, industrial water withdrawal would increase 25.8% and at highest per capita rate, it would increase 57.7% in 2050. For municipal water withdrawal, at lowest per capita rate, it would decrease 46.18% in 2050 which is a positive scenario, whereas for business-as-usual (this is also highest per capita) rate, it would increase 25.8%. For produced municipal wastewater, in 2050, it is possible able to reduce it to 53.02% maintaining lowest per capita rate. Otherwise, at business-as-usual (highest per capita) rate, it would increase 24.82%. Municipal wastewater production might be 1.3 times more in 2050 than 2011. For treated municipal wastewater, it would decrease to 60.71% maintaining lowest per capita rate. But at business-as-usual (highest per capita) rate of wastewater treatment, amount of treated municipal wastewater would increase 24.82% which is also positive scenario. For non-treated municipal wastewater, it is possible to decrease it to 56.83% maintaining lowest per capita rate in 2050. On another hand, at business-as-usual (highest per capita) rate, it would increase 24.82% which paints a bleak future of India. This non-treated municipal wastewater might be 1.33 times more in 2050 than 2011. Both total renewable water resource and total internal renewable water resource, if lowest per capita (which is also business-as-usual) rate can be maintained by India, only 22% more renewable water resource will be needed in 2050. Otherwise, it would plunge to 72.15% if highest rate of renewable water resource per capita is to be maintained.

Fig. 3. Future scenario of indicators of biophysical consumption related to sustainable development goal 6 for India up to 2050.

Blue indicates changes in total values of consumption. Brown indicates projected total values at the highest rate of per capita consumption; Orange indicates projected total values at business-as-usual (BAU) rate of per capita consumption, Yellow indicates projected total values at a lowest rate of per capita consumption.

4. Discussion:

This study clearly shows that the per capita biophysical resources are decreasing over time. This trend would continue further with respect to population explosion and growing urbanisation trends in India. However, it also shows that social developments related to water and sanitation are improving, clearly at the cost of environment. To provide positive social

development and maintaining the same while the population is exploding will be tough to

handle for the government. Especially, if all of these developments come at the cost of further

degradation of the environment.

During this study, we confronted some problems along with their probable mitigation

strategies. First, availability of biophysical resource consumption data is not good enough.

For the same reason, we were unable to perform any statistical analysis. In future, analysis

and freely available data concerning biophysical consumption at state and district level

should be prepared and maintained, both for general awareness and scientific research

purposes. Second, long-term data are not available for most of the indicators which are

necessary for more accurate projection of future trends. Third, neither state-level

comprehensive data for most of the indicators are available, nor the district or any other lower

unit level. This simply creates a homogeneous outline of resource consumption and access at

national level. This is not enough to redesign policy framework or restructure governance in

accordance with scientific findings. Though it is enough for preliminary studies, would not be

enough in future, especially when adaptive policy needs to be implemented customized for

the local context. Fourth, abundance and availability of data of different biophysical

parameters are much more than different socioeconomic indicators. It shows a clear bias

towards the maintenance of biophysical data. This is not suitable for contemporary research

works and creates problem to assess sustainability for all three dimensions (i.e. environment,

society and development). Fifth, the set of indicators for sustainable development of water

and sanitation sector does not yield a very complete and comprehensive picture of SDG 6 in

India, especially in view of the socioeconomic factors that exist within the country and the

regional biophysical resource consumption that interact with them. Therefore, there is ample

scope for adapting the indicators to better suit India's specific needs in terms of reflecting

both regional and national scenario. To this end, additional indicators could be developed.

Sixth, as this study only etches the surface of multi-indicator multilevel continuous

monitoring of SDG 6, it is not rigorous enough to map synergy and trade-off between SDG 6

and other sustainable development goals (Griggs et al. 2014, Nilsson et al. 2016). Seventh,

awareness and know-how about water resource management in India is only catalysed by the

government in broad scale. Thus, government, nongovernment organizations (NGOs) and the

public, together should take more proactive steps towards promotion and utilization of water

resources to achieve sustainable development goals in water and sanitation (i.e. SDG 6) in

India. Eighth, to control misutilization and overexploitation of water resources, especially in

urban areas, regulatory instruments, such as - water utilization taxes, water recycling taxes

etc. will remain a vital component of policies, as efforts to increase water efficiency and

reduce environmental and socioeconomic impacts are usually unsupportable against a

background of low cost availability. Ninth, if consumers do not purchase devices with

improved water use efficiency and environment characteristics (e.g. agricultural transport and

household appliances etc.), when such devices are available commercially, these products

should be supplemented by more direct government intervention and investment if

advancement toward sustainable development in water resources are to be made. Tenth, since

agriculture and household are two major water consuming sector in India, utilization of

recycled and treated wastewater should be advocated. Eleventh, since agricultural water

withdrawal is highest in India among all other sectors, increasing agricultural water use

efficiency should be the highest priority through drip irrigation, hydroponics, urban terrace

and rooftop agriculture, utilization of treated household wastewater in agriculture etc.

Water resources and its utilization are intimately connected to sustainable development. For

Indian societies to attain sustainable development in water and sanitation, much effort must

be devoted not only to discover sustainable ways to consume of water resources but also to

increase the water use efficiencies (WUE) of processes utilizing water resources. Also,

correlated environmental concerns must be addressed, continuously monitored and any type or degree of declination must be solved. This proposed research framework would progress our current understanding and measurement of SDG 6. Through measurement and identification of how, when, and why changes in biophysical aspects of water and sanitation influence human socioeconomic conditions over both time and space and vice-versa, we can more fully appropriate and anticipate the impact of the bidirectional nature of human-water interactions. This understanding might also be used to monitor, update and refine the national SDG 6 framework. This would arm natural resource planning authorities with a decision support and policy management tool that more accurately reflects the trade-offs between various components of the SDG 6. As a final note, to make India water secure and sustainable in 2030, India must endeavour to increase its efforts to attain greater efficiency in utilization, consumption of water for both biophysical and socioeconomic purposes.

Acknowledgement:

This study was supported by FRPDF scheme (2013-2017) of Presidency University, Kolkata. We would like to thank Sk. Rohan Tanvir, The Institution of Engineers (India) for his kind assistance during the preparation of diagrams.

References:

Ait-Kadi, M., 2016. Water for Development and Development for Water: Realizing the

Sustainable Development Goals (SDGs) Vision. Aquat. Procedia. 6, 106-110. doi:

10.1016/j.aqpro.2016.06.013.

Bartram, J., Brocklehurst, C., Fisher, M. B., Luyendijk, R., Hossain, R., Wardlaw, T.,

Gordon, B., 2014. Global monitoring of water supply and sanitation: history, methods and

future challenges. Int. J. Environ. Res. Public Health, 11 (8), 8137-8165.

doi:10.3390/ijerph110808137.

Bartram, J., Cairncross, S., 2010. Hygiene, sanitation, and water: Forgotten foundations

of health. PLoS Med. 7 (11), 1-9. doi: 10.1371/journal.pmed.1000367.

Bartram, J., Lewis, K., Lenton, R., Wright, A., 2005. Focusing on improved water and

sanitation for health. Lancet. 365, 810-812. doi:10.1016/S0140-6736(05)17991-4.

Bhaduri, A., Bogardi, J., Siddiqi, A., Voigt, H., Vörösmarty, C., Pahl-Wostl, C., Bunn,

S.E., Shrivastava, P., Lawford, R., Foster, S., Kremer, H., Renaud, F.G., Bruns, A.,

Osuna, V.R., 2016. Achieving Sustainable Development Goals from a Water Perspective.

Front. Environ. Sci. 4, 1-13. doi: 10.3389/fenvs.2016.00064.

Biswas, A.K., 1992. Sustainable water development: A global perspective. Water Int. 17,

68-80. doi: 10.1080/02508069208686128.

Biswas, A.K., 1991. Water for Sustainable Development in the 21st Century: A Global

Perspective. Int. J. Water Resour. Dev. 7 (4), 219-224. doi:

10.1080/07900629108722516.

Bogardi, J.J., Dudgeon, D., Lawford, R., Flinkerbusch, E., Meyn, A., Pahl-Wostl, C.,

Vielhauer, K., Vörösmarty, C., 2012. Water security for a planet under pressure:

Interconnected challenges of a changing world call for sustainable solutions. Curr. Opin.

Environ. Sustain. 4, 35-43. doi: 10.1016/j.cosust.2011.12.002.

Brundtland, G.H., 1987. Our Common Future. Oxford University Press, New York.

FAO 2018. Aquastat online database. http://www.fao.org/nr/water/aquastat/main/index.stm (accessed 28.04.2018)

FAO 2018. FAOSTAT online database; http://faostat.fao.org/ (accessed 28.04.2018)

IndiaStat 2018. https://www.indiastat.com/ (accessed 22.05.2018)

Gleick, P.H., 1996. Basic water requirements for human activities: Meeting basic needs. Water Int. 21, 83-92, doi: 10.1080/02508069608686494.

Gleick, P.H., 1998. Water in crisis: paths to sustainable water use. Ecol. Appl. 8 (3), 571-579. doi: 10.1890/1051-0761(1998)008[0571:WICPTS]2.0.CO;2.

Griggs, D., Stafford-Smith, M., Gaffney, O., Nature, J.R.-, 2013. Policy: Sustainable development goals for people and planet. Nature, 495, 305-307. doi: 10.1038/495305a.

Jackson, R.B., Carpenter, S.R., Dahm, C.N., McKnight, D.M., Naiman, R.J., Postel, S.L., Running, S.W., 2001. Water in a changing world. Ecol. Appl. 11 (4), 1027-1045. doi: 10.1890/1051-0761(2001)011[1027:WIACW]2.0.CO;2.

Kumar, R., Singh, R.D., Sharma, K.D., 2005. Water resources of India. Curr. Sci. 89 (5), 794-811. doi: 10.1002/047147844X.wr243

Madrid, C., Cabello, V., Giampietro, M., 2013. Water-Use Sustainability in Socioecological Systems: A Multiscale Integrated Approach. Bioscience. 63, 14-24. doi: 10.1525/bio.2013.63.1.6.

Mall, R.K., Gupta, A., Singh, R., Singh, R.S., Rathore, L.S., 2006. Water resources and climate change: An Indian perspective. Curr. Sci. 90 (12), 1610-1626. doi: 10.1016/S0143-8166(02)00004-0.

Moe, C.L., Rheingans, R.D., 2006. Global challenges in water, sanitation and health. J. Water Health. 4 (S1), 41-57. doi: 10.2166/wh.2005.039.

Montgomery, M.A., Elimelech, M., 2007. Water And Sanitation in Developing Countries:

Including Health in the Equation. Environ. Sci. Technol. 41, 17–24. doi:

10.1021/es072435t

Mujumdar, P.P., 2008. Implications of climate change for sustainable water resources

management in India. Phys. Chem. Earth. 33, 354-358. doi: 10.1016/j.pce.2008.02.014.

Nilsson, M., Griggs, D., Visbeck, M., 2016. Policy: Map the interactions between

Sustainable Development Goals. Nature, 534, 320-322. doi: 10.1038/534320a.

Poff, N., Olden, J. D., Merritt, D. M., Pepin, D. M., 2007. Homogenization of regional

river dynamics by dams and global biodiversity implications. Proc. Natl. Acad Sci. 104

(14), 5732-5737. doi: 10.1073/pnas.0609812104.

Postel, S.L., 2000. Water and world population growth. J. Am. Water Work. Assoc. 2 (4),

131-138. doi: 10.1002/j.1551-8833.2000.tb08927.x.

Raworth, K., 2012. A safe and just space for humanity - Can we just live within the

doughnut. Oxfam Policy and Practice: Climate Change and Resilience. 8 (1), 1–26. doi:

10.1080/00420980120087081.

Raworth, K., 2017a. A Doughnut for the Anthropocene: humanity's compass in the 21st

century. Lancet Planet. Health. 1 (2), e48–e49. doi: 10.1016/S2542-5196(17)30028-1.

Raworth, K., 2017b. Doughnut economics: seven ways to think like a 21st-century

economist. Random House Business,

Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E., Lenton,

T.M., Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T.,

van der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U.,

Falkenmark, M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B.,

Liverman, D., Richardson, K., Crutzen, P., Foley, J., 2009a. A safe operating space for

humanity. Nature, 461, 472-475. doi: 10.1038/461472a.

Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F.S., Lambin, E., Lenton, T.M., Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., Foley, J., 2009b. Planetary boundaries: Exploring the safe operating space for humanity. Ecol. Soc. 14 (2), 32. doi: 10.5751/ES-03180-140232

Sachs, D. J., 2012. From millennium development goals to sustainable development goals. Lancet, 379, 2206-2211. doi: 10.1016/S0140-6736(12)60685-0

Serageldin, I., 1995. Water resources management: A new policy for a sustainable future. Water Int. 20, 15-21. doi: 10.1080/02508069508686440.

Steffen, W., Crutzen, P.J., McNeill, J.R., 2007. The Anthropocene: Are Humans Now Overwhelming the Great Forces of Nature. AMBIO A J. Hum. Environ. 36 (8), 614-621. doi: 10.1579/0044-7447(2007)36[614:TAAHNO]2.0.CO;2.

Steffen, W., Persson, Å., Deutsch, L., Zalasiewicz, J., Williams, M., Richardson, K., Crumley, C., Crutzen, P., Folke, C., Gordon, L., Molina, M., Ramanathan, V., Rockström, J., Scheffer, M., Schellnhuber, H.J., Svedin, U., 2011. The anthropocene: From global change to planetary stewardship, AMBIO A J. Hum. Environ. 40 (7), 739-761. doi: 10.1007/s13280-011-0185-x.

Steffen, W., Richardson, K., Rockstrom, J., Cornell, S.E., Fetzer, I, Bennett, E.M., Biggs, R., Carpenter, S.R., de Vries, W., de Wit, C.A., Folke, C., Gerten, D., Heinke, J., Mace, G.M., Persson, L.M., Ramanathan, V., Reyers, B., Sorlin, S., 2015. Planetary boundaries: Guiding human development on a changing planet. Science, 347 (6223), 1-10. doi: 10.1126/science.1259855.

United Nations General Assembly, 2015. Transforming our world: The 2030 agenda for

sustainable   development,

https://sustainabledevelopment.un.org/content/documents/7891Transforming%20Our%20

World. pdf. https://doi.org/10.1007/s13398-014-0173-7.2

United Nations, Department of Economic and Social Affairs, Population Division (2017).

World Population Prospects: The 2017 Revision.

United Nations, 2015. Transforming Our World: The 2030 Agenda for Sustainable

Development (A/RES/70/1).

United Nations World Population Prospects: The 2017 Revision, (Population Division,

Department of Economic and Social Affairs, United Nations, 2017).

van Koppen, B., Shah, T., 2006. Is India Ripe for Integrated Water Resources

Management? Econ Polit Wkly. 41 (31), 3413-3421.

Vorosmarty, C.J., Green, P., Salisbury, J., Lammers, R. B., 2000. Global Water

Resources: Vulnerability from Climate Change and Population Growth. Science, 289

(5477), 284-288. doi: 10.1126/science.289.5477.284.

Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,

Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global

threats to human water security and river biodiversity. Nature, 467, 555-561. doi:

10.1038/nature09440.

World Bank 2018. World Development Indicators (World Bank);

http://data.worldbank.org/. (accessed 28.04.2018)

Fig. 1. Changes in biophysical and social development indicators related to sustainable development goal 6 in India.

A and B represent the status of sustainable development of India in 2000 and 2015 respectively. Eleven indicators of biophysical stress established over biophysical ceiling (outer chocolate ring) outwardly projected and seventeen indicators of social development established under social foundation (inner chocolate ring) inwardly projected for India. Biophysical indicators are – total water withdrawal (TWW), agricultural water withdrawal (AWW), industrial water withdrawal (IWW), municipal water withdrawal (MWW), total internal renewable water resource (TIRWR), total renewable water resource (TRWR), dam capacity (DC), water stress (WS), produced amount of municipal waste water (PMWW), treated municipal waste water (TMWW) and non-treated municipal waste water (NTMWW). Indicators of social development are – (1) Improved sanitation facilities, (2) Improved sanitation facilities in rural areas, (3) Improved sanitation facilities in urban areas, (4) Improved water source, (5) Improved water source in rural areas, (6) Improved water source in urban areas, (7) People practicing open defecation, (8) Rural people practicing open defecation, (9) Urban people practicing open defecation, (10) People using at least basic drinking water services, (11) Rural people using at least basic drinking water services, (12) Urban people using at least basic drinking water services, (13) People using at least basic sanitation services, (14) Rural people using at least basic sanitation services, (15) Urban people using at least basic sanitation services, (16) Rural people using safely managed drinking water services, (17) Water use efficiency in agriculture.

Green indicates safe operating space for biophysical indicators and thresholds for indicators of social development. Yellow indicates zone of increasing impact for biophysical indicators and zone of increasing deprivation for indicators of social development. Red indicates zone of high risk of serious impact for biophysical indicators and zone of high level of deprivation for indicators of social development. Light blue represents indicators without any boundary or thresholds (Unit – m3). Violet represents water stress (Unit - %). Area between the chocolate rings is the safe and just operating space for sustainable development of water and sanitation in India.

Fig. 2. Changes in per capita biophysical indicators (1-5) and indicators of social development (6-12) related to sustainable development goal 6 in India with time.

Green indicates global per capita boundaries for biophysical indicators and thresholds for indicators of social development.

Biophysical indicators are - (1) water withdrawal (m^3), (2) water resources (m^3), (3) dam capacity (m^3), (4) municipal wastewater ((m^3) and (5) water stress (%). Indicators for social development are - (6) open defecation (%), (7) basic drinking water services (%), (8) basic sanitation services (%), (9) safely managed drinking water services (%), (10) improved sanitation facilities (%) (11) improved water source (%) and (12) water use efficiency in agriculture (m^3/GDP, current US$).

Fig. 3. Future scenario of indicators of biophysical consumption related to sustainable development goal 6 for India up to 2050.

Blue indicates changes in total values of consumption. Brown indicates projected total values at highest rate of per capita consumption; Orange indicates projected total values at business-as-usual (BAU) rate of per capita consumption, Yellow indicates projected total values at lowest rate of per capita consumption.