1   **Estimating the benefit of quarantine: eradicating invasive cane toads from islands**

3   Adam S Smart[1*], Reid Tingley[12] and Ben L Phillips[1]

4   [1]School of Biosciences, University of Melbourne, Parkville, VIC, 3010, Australia

5   [2]School of Biological Sciences, Monash University, Clayton, VIC, 3800, Australia

6   *Corresponding author: Adam Smart   email: asmart1@student.unimelb.edu.au

7   phone: +61 03 9035 7555

8   Reid Tingley   email: reid.tingley@unimelb.edu.au

9   Ben Phillips   email: phillipsb@unimelb.edu.au

10   Running title: The benefit of cane toad quarantine

11   Word Count;   summary: 342

12   main text: 3926

13   acknowledgements: 116

14   references: 1827

15   tables: 80

16   figure legends: 182

17   Number of tables and figures: **8**

18   References: **58**

20   Manuscript for consideration in *Journal of Applied Ecology*

Summary

1. Islands are increasingly used to protect endangered populations from the negative impacts of invasive species. Quarantine efforts are particularly likely to be undervalued in circumstances where a failure incurs non-economic costs. One approach to ascribe value to such efforts is by modeling the expense of restoring a system to its former state.

2. Using field-based removal experiments on two very different islands off northern Australia separated by > 400 km, we estimate cane toad densities, detection probabilities, and the resulting effort needed to eradicate toads from an island, and use these estimates to examine the financial benefit of cane toad quarantine across offshore islands prioritized for conversation management by the Australian federal government.

3. We calculate density as animals per km of freshwater shoreline, and find striking concordance of density across our two island study sites: a mean density of 353 [286, 446] individual toads per kilometer on one island, and a density of 366 [319, 343] on the second. Detection probability differed between the two islands.

4. Using a removal model and the financial costs incurred during toad removal, we estimate that eradicating cane toads would, on average, cost between $9444 (based on Horan Island; high detectability) and $18093 AUD (Indian Island; low detectability) per km of available freshwater shoreline.

5. Across islands that have been prioritized for conservation benefit within the toads'

predicted range, we provide an estimate of the value of toad quarantine on each island,

and estimate the net value of quarantine efforts to be between $27.25 – $52.20 Million

AUD. We explore a proposed mainland cane toad containment strategy – to prevent the

spread of cane toads into the Pilbara Bioregion, and estimate its potential value to be

between $33.79 – $64.74 M AUD.

6. *Synthesis and applications*. We present a modelling framework that can be used to

estimate the value of preventative management, via estimating the length and cost of an

eradication program. Our analyses suggest that there is substantial economic value in

cane toad quarantine efforts across Australian offshore islands and a proposed mainland

toad containment strategy.

Key-words: Cane Toad, density, detection probability, eradication, islands, quarantine

Introduction

It is a truth universally acknowledged that an ounce of prevention is worth a pound of cure. In invasive species management, this can be achieved by preventing human-mediated dispersal of non-indigenous species (Chen *et al*. 2018), by conducting routine surveillance programs aimed at early detection (Holden *et al*. 2015), and via translocation of endangered taxa beyond the current or predicted distributions of invaders (Woinarski *et al*. 2014; Legge *et al*. 2018; Moseby 2018) (National Species Management Plan, 2008). Despite such truisms, conservation managers rarely ascribe value to preventative management. Whilst preventative measures are increasingly being adopted to save imperiled taxa (Burns *et al*. 2012; Commonwealth of Australia 2015), without valuation, we risk falling prey to cognitive biases (e.g., immediacy bias), and so routinely commit substantially more money and effort to tactical, "cure" type approaches, than to strategic "prevention". Quarantine against invasive species is a case in point; vastly more resources are spent controlling the spread and impact of invaders than are spent on preventing their arrival and establishment (Hoffman & Broadhurst 2016).

Quarantine is particularly likely to be undervalued in circumstances in which a failure incurs non-economic costs (e.g., biodiversity loss) (Leung *et al*. 2002) or when costs or damages persist over long-time scales (Epanchin-Neill *et al*. 2015). One way to place a value on such quarantine efforts is to calculate the cost of restoring the system to its former state (Kimball *et al*. 2014; Rohr *et al*. 2016). In the case of an invasive species with primarily non-economic impacts, we can calculate the ongoing benefit of quarantine as the expense of restoring the system to this former state, i.e., a subsequent eradication program. Such a valuation is a lower bound on the benefit of quarantine for a number of reasons. First, the same quarantine effort typically protects against many potential invasive species. In addition, any impact that an invasive species has

before it is eradicated (e.g., local extinction or shifts in population structure of a native species,

altered landscape vegetation profiles formation) must be added to the cost of restoration

(Hoffmann & Broadhurst 2016, Jardine & Sanchirico 2018). Thus, the cost of eradicating a

single invader is a very conservative estimate of the true value of quarantine efforts.

Islands are important resources for conservation quarantine because they offer a natural barrier to

the spread of invasive species. Conservation biologists routinely exploit this property of islands,

not only to protect species that naturally occur on islands, but also to provide refuge for species

under threat on the mainland (Thomas 2011; Tershy *et al.* 2015; Legge *et al.* 2018). In Australia

alone, a minimum of 47 conservation translocations to islands have been carried out to date

(Department of the Environment, Water, Heritage and the Arts, 2009). In these circumstances –

where the conservation value of an island has been artificially bolstered – the subsequent arrival

of invasive species can have a larger impact than they otherwise would. Typically, island

quarantine is used by conservation managers to protect native species from invasive predators

(e.g., foxes, cats, weasels, rats). In Australia, however, islands are also used to mitigate the

impact of cane toads (*Rhinella marina*) on native predators (Moro *et al.* 2018; Ringma *et al.*

2018). Cane toads were introduced to northeastern Australia in the 1930s and, in northern

Australia, continue to spread westerly at a rate of ~50 km per year (Phillips *et al.* 2010). This

invasion has had major impacts on populations of native predators, many of which have no

resistance to the toad's toxin (Nelson *et al.* 2010; Greenlees *et al.* 2010; Llewelyn *et al.* 2014). In

response to declines of multiple predator species (e.g., dasyurids, monitors, snakes) the

Australian government implemented the Cane Toad Threat Abatement Plan (2011), which aimed

to identify, and where possible reduce, the impact of cane toads on native species

(Shanmuganathan *et al.* 2010). A lack of viable methods for broad-scale control, however, has

since led the Australian government to place an increased emphasis on containment (on the

mainland) and on quarantine (on offshore islands) to mitigate the biodiversity impacts of cane

toads.

While quarantine is currently the best available strategy, it is not a panacea: cane toads have

already established on at least 48 islands across northern Australia (McKinney *et al*. 2018 unpub

data), with potential for further natural and anthropogenic introductions. Thus, execution of the

strategy outlined in the Cane Toad Threat Abatement Plan requires ongoing quarantine,

eradication and containment efforts. Here we estimate the lower bound of the monetary value of

these ongoing efforts, by quantifying the cost of eradicating cane toads from two islands in

northern Australia. We approach this problem by estimating the density and detection probability

of toads on each island, and use these estimates to calculate the amount of time and money it

would take to remove enough toads to ensure eradication.

Materials and methods

Study Area

This study was carried out on two islands in northern Australia: Horan Island on Lake Argyle,

Western Australia (HI) and Indian Island in the Northern Territory (II). Lake Argyle is Western

Australia's largest man-made reservoir covering > 880 km^2 and is located within the East

Kimberly region. The study site is composed of exposed spinifex-covered hilltops and sparse

savanna woodland. Freshwater is available year-round, with the lake contracting from May–

November. Toads are thought to have colonized islands on the lake in the wet seasons of

2009/2010 (Somaweera & Shine 2012). Indian Island is an offshore island, 40 km west of

Darwin. It supports predominantly savanna woodland and monsoonal vine thicket, with a large

ephemeral freshwater swamp located on the northern tip of the island. Depending of the

magnitude of the wet season, standing water can be present in this swamp year-round or dry up

by late September. Toads are thought to have colonized Indian Island via rafting events around

2008. Access to Indian Island was granted by Kenbi Traditional Owners (Northern Land Council

permit 82368).

Field sampling

Cane toad surveys occurred over six nights, on each island, denoted, $t = \{0, 1, \ldots, 5\}$, during

November 2017 (HI) and October 2018 (II). Surveys commenced at sundown each evening and

lasted three hours, with ambient temperatures ranging from $24 - 35°C$. As Horan Island occurs

on a freshwater lake, the entire island was circumambulated each night by two people using

headtorches; one individual focused on the higher part of the shoreline, the other on the lower

shoreline. Indian Island is an oceanic island, with only a single freshwater swamp present in the

dry season. This swamp was navigated each night by two people using head torches. On both

islands, every toad encountered was collected and humanely killed on site in accordance with

The University of Melbourne animal ethics protocol (1714277.1) and State laws regarding

handling of non-native species. Each night, we recorded the number of individuals collected, c_t.

Surveys were conducted immediately prior to the breeding season so that only post-metamorphic

age classes were encountered.

Statistical analysis

We do not encounter every individual on a given night, and so incorporate imperfect detection.

For each island, we aim to estimate three parameters: N_0, the true number of toads on the island

at the commencement of surveys; p, the mean per-individual detection probability; and α, the

length of time (in days) required to eradicate toads from our treatment areas. The number of

individuals collected each night, c_t, can be considered a draw from a binomial distribution with:

$$c_t \sim \text{Binom}(N_t, p).$$

Where N_0, the pre-sampling population size, is a latent variable with a mean and variance equal

to λ, such that:

$$N_0 \sim \text{Poiss}(\lambda).$$

For $t > 0$:

$$N_t = N_0 - \sum_0^{t-1} c_t.$$

The length of time required to remove a population, α, from a treatment area is described via the

relationship:

$$\alpha = \left(\frac{\ln(r_{crit})}{\ln(1-p)} \right),$$

where, r_{crit}, the critical removal threshold, is equal to $\frac{1}{N_0}$ (the inverse of the pre-sampling

population size).

Models were fit with Markov chain Monte Carlo (MCMC) in JAGS v.4.6.0, run through R

v3.4.1 via the package rjags v4.6.0 (Plummer & Martyn 2013). Three model chains were run for

30,000 iterations, with the first 10,000 iterations discarded as a burn-in, which was sufficient for

the MCMC chains to converge. Convergence was checked using the Gelman-Rubin diagnostic

(Gelman & Rubin 1992); all chains produced potential scale reduction factors < 1.1, indicating

convergence of chains. The remaining samples were thinned by a factor of 2, resulting in 10,000

samples per chain for post-processing. Minimally informative prior distributions for p and λ

were specified as uniform between 0 - 1 and 0 - 10,000 respectively.

We denote a successful eradication to have occurred when only a single toad remains (i.e., no

further breeding pairs remain). As we assume that removal efforts take place on consecutive

nights until completion, we disregard breeding and immigration.

Cost analysis

We estimate the cost of eradicating toads on our study islands based on consumable, personnel,

and travel costs incurred during toad collection (see Appendix S1 in Supporting Information).

Relative to most islands across northern Australia, both Horan and Indian Islands are readily

accessible, thus our travel costs are modest. We assume that eradication is conducted by a fully-

equipped organization; thus we do not include vehicle/boat purchase or hire (i.e., set-up costs),

nor do we consider organizational in-kind associated with utilizing existing capital. Removal

efforts are carried out on subsequent nights until eradication is reached; therefore, the cost

associated with travel to and from our site is incurred only once. Travel costs include a \$85/hour

consultant rate plus the additional costs of fuel, insurance, and vehicle maintenance (an extra

\$36/hour). Thus, total travel costs are \$111/hour.

Cost Scenarios

We use our estimates of toad removal on Horan and Indian Islands (with their attendant detection

probabilities) to highlight the potential benefit of quarantine efforts on a subset of high priority

islands (Table 1). Our chosen islands are drawn from a list of 100 oceanic islands that the

Australian Commonwealth has prioritized for conservation, due to their biodiversity value and

presence of species listed under the Environment Protection and Biodiversity Conservation Act

(Department of the Environment and Energy [DEE], 1999). We refine this list to include only

islands that are ≥2 km from the Australian mainland and occur within the potential distribution

of cane toads in Australia (Kearney *et al*. 2008). For each island in our dataset, we map the

length of permanent freshwater shoreline available, using either satellite maps,

government/landholder records, or a combination of both – resulting in a net kilometer length of

shoreline for each island in our dataset. All islands were crossed-checked for the presence of

cane toads via the 'Feral Animals on Offshore Islands' database (DEE, 2016) in addition to the

presence of human settlement. In cases where islands had no permanent freshwater but did have

human settlement (or known livestock presence), a one-kilometer circumference was assumed

around dwellings and visible watering points.

In addition to the islands derived from this report, we explore the value of a potential cane toad

containment strategy outlined in a revised version of the Cane Toad Threat Abatement Plan

(Tingley *et al*. 2013).

This strategy aims to develop a 'waterless barrier' on the Australian mainland by excluding cane

toads from artificial water bodies on cattle stations between Broome and Port Hedland in

Western Australia. Using a dataset containing the presence of bore holes, cattle watering points,

dams and permanent freshwater bodies in the Pilbara bioregion (see Southwell *et al*. 2017) we

estimate the economic benefit of the proposed barrier. A one-kilometer circumference was

applied to all waterpoints, dams and pools, in addition to a per-kilometer of shoreline rate along

permanent watercourses within the region. If implemented successfully, this strategy could keep

toads out of the Pilbara (and subsequent regions) – an effective quarantine of $268,00km^2$ of the

Australian mainland (see Florance *et al*. 2011; Tingley *et al*. 2013; Southwell *et al*. 2017 for

further information).

Results

The number of cane toads removed from both Horan and Indian Island, c_t, declined over time

(Figure 1). Across the duration of our surveys, we captured and removed a total of 1550 cane

toads (1251 on HI, 299 on II). The estimated probability of detecting an individual toad on a

given night differed between our two study sites (Horan Island: mean p [95% credible interval] =

0.1 [0.07, 0.13]; Indian Island: 0.27 [0.22, 0.33]) (Figure 2). Given the site-specific detection

probability, the estimated number of toads present at the initiation of our surveys (N_0) was much

higher on Horan Island (2681 [2171, 3393]) than on Indian Island (353 [308, 408]) (Figure 3).

Horan Island – situated in a freshwater lake – has a circumference of 7.63 km, which translates

to a cane toad density of 353 [286, 446] individuals per kilometer of freshwater shoreline. The

freshwater source on Indian Island has a circumference of 1.04 km, translating to a density of

366 [319, 343] individuals per kilometer of shoreline (Figure 4). We could also express toad

density as animals per km^2 of island, in which case we calculate a density of individuals of 56 on

II and 2852 on HI.

Given the posterior estimates of p and λ, we examine the total survey effort (in days) required to

eradicate toads on both Horan and Indian Island. Inputting the distribution of N_0, leaving a single

individual is equivalent to leaving the proportion, $r_{crit} = \frac{1}{N_0}$ of the original individuals. The time

to reach this point is given by $\ln(r_{crit})/\ln(1 - p) = 75$ days [49, 110] on HI, and 19 days [10,

29] on II (Figure 5).

Cost Sensitivity

Multiplying the estimate of the number of days required to achieve eradication by our removal

costs suggests that $62 407 [$40 583, $91 104] would be required to eradicate toads from HI.

This equates to $9 748 [$6 339, $14 231] per kilometer of freshwater shoreline or $80 009 [$52

030, $116 801] per km^2 of land. In contrast, the cost to eradicate toads from II is estimated to be

$18 394 [$9 872, $28 636], equating to $17 737 [$9 523, $27 615] per kilometer of freshwater

shoreline or $2 929 [$1 572, $4 560] per km^2 of land (Figure 6).

Benefit of quarantine on Prioritized Australian Islands

Using our estimates of eradication costs per-kilometer of freshwater shoreline, we examine the

economic benefit of cane toad quarantine on all toad-free islands (by jurisdiction), as well as the

cost to restore all toad-inhabited islands to a toad-free state (Figure 7). The current economic

benefit of quarantine on all prioritized toad-free islands is estimated to be between $17.5 M (HI

estimates) and $33.3 M (based on II estimates). We estimate it would cost, on average, between

$2.8 M (HI) and $5.2 M (II) to remove toads from all prioritized islands currently occupied by

toads. Finally, we estimate the economic benefit of the 'waterless barrier' protecting the Pilbara

to be between $34.3 M (HI) and $63.6 M (II).

Discussion

As the number of invasive species requiring management increases, practitioners must identify

efficient strategies for allocating resources to various management activities. Although

conventional wisdom places emphasis on prevention measures, the practice of valuing such

actions in the face of non-economic costs can be challenging. Placing a monetary value on a

conservation benefit will most often require some value judgement as to the monetary worth of

biodiversity. Using estimates of a species detectability, population density, and subsequent

eradication costs we aim to prevent such value judgement when investigating the benefit of

quarantine measures in combatting the impact of the invasive cane toad across Australia's

prioritized offshore islands.

Despite substantial community and research effort into cane toad removal via trapping and hand

capture, there are only a handful of published detection estimates for the species (Griffiths &

McKay 2007). Our detection estimate is, of course, specific to the details of our survey.

Nonetheless, it is surprisingly low for our large-shoreline site (Horan Island). Here, the length of

shoreline meant we only passed each location once per night, and individual toads in this closed

system had, on average, a 0.1 [0.07 – 0.13] chance of being seen on any given night. This

contrasts with our small-shoreline site (Indian Island) where we were able to make multiple

passes of the same point each night. Here, individual toads had a 0.27 [0.22 – 0.33] chance of

being detected on a given survey night. Whilst individual toads are relatively easy to see when

they are active, our results suggest that this might give a misleading impression of one-pass

detectability.

We compared two density metrics: a linear density (per km) and an areal density (per km^2). Our

areal density estimate (2 852 individuals/ km^2) is similar to estimates derived from previous

studies of invasive cane toads in the Solomon Islands archipelago (1 035/km^2; Pikacha *et al*.

2015), the islands of Papua New Guinea (3 000/km^2; Zugg *et al*. 1975; Freeland *et al*. 1986), and

density estimates of an analogous invasive toad on Madagascar (3 240/km^2; Reardon *et al*.

2018). A single study conducted on the Australian mainland reported densities as high as 256

300 individuals per km² (Cohen & Alford 1993), but this estimate was predominantly of the

metamorph life stage, which occurs at very high densities prior to dispersal. Metamorphs are

strongly constrained to the edges of water bodies (Child *et al*. 2008), and typically suffer high

mortality from predation and desiccation before reaching maturity (Ward-Fear *et al*. 2010).

While an areal density would make sense in a habitat where animals are constrained by some

factor that scales with area (e.g., primary productivity), it is clear that toads in northern Australia

are often constrained by access to water in the dry season, and thus length of shoreline is more

appropriate. Length of shoreline not only defines access to water, but also the density of

infectious parasites (such as *Rhabdias pseudosphaerocephala*) that use the moist conditions and

high toad densities along shorelines as opportunities for transmission ((Kelehear *et al*. 2011;

2013). It is also likely that the survival rate of emergent metamorphs is dependent on length of

shoreline, because this will set the density of conspecifics and so moderate the rate at which

these conspecifics cannibalize each other (Pizzatto & Shine 2008).   In comparing the areal and

linear densities between our sites we find a large difference between sites in the areal metric, but

a strikingly similar density value across sites in the linear metric. Our results suggest that across

these two different systems, toads achieve a density of around 354 adults per kilometer of

shoreline.

Because toads in dry conditions require regular re-hydration (Seebacher & Alford 2002; Tingley

& Shine 2011) it is a logical step to conduct removal efforts when toads are restricted to a subset

of semi-permanent hydration points during drier sections of the year (Letnic *et al*. 2015). Given

the linear density metric is so concordant across sites, this is the best metric for calculating

eradication costs. Certainly, if we use the areal metric we find a wide gulf in the possible

eradication values in contrast to our shoreline metric (Figure 6). Costs based on the linear metric

had a greater correspondence between the two sites ($9 748 and $ 17 737 AUD per km of

shoreline). Encouragingly our cost estimates appear similar to estimates derived from a

successful eradication program associated with removing the American bullfrog from two

locations in Canada ($8 200 – $23 000 CAN per kilometer of freshwater shoreline).

To our knowledge, there is only one instance in which the cost to eradicate cane toads from an

island has been documented (Wingate 2011). Carried out on Nonsuch Island in Bermuda, this

removal occurred over six years and included countless volunteer hours and an investment of

$10 000 USD (~$14 330 AUD) to remove toads from an area of 0.6 km^2. In addition, two

successful eradications from extralimital mainland sites have been documented, occurring

beyond the southern border of the cane toads' current range in Australia (White 2010; Greenlees

et al. 2018). This handful of successful removals of the cane toad, mirrors a broad trend in the

eradication of invasive amphibian populations globally (Adams & Pearl 2007; Kraus 2009;

Beachy *et al*. 2011; Orchard 2011). As such, there is scant information available to guide policy

makers and management agencies when evaluating the feasibility of implementing amphibian

quarantine and eradication measures.

If we are to shift away from tactical, post-invasion approaches, to a preventative strategic

approach, management practitioners require an estimate of the economic value that quarantine

holds. Our analysis of the feasibility and benefit of cane toad quarantine is timely, given renewed

emphasis on Australia's offshore islands as safe-havens to buffer biodiversity against cane toad

impacts. Sixty-two Australian offshore islands designated as 'high conservation status' fall

within the cane toad's predicted distribution; 21 of those have already been colonized by toads.

Given our criteria (see Methods), we estimate the remaining value of toad quarantine across

toad-free islands in northern Australia to be up to $33 M AUD. This value is conservative for a

number of reasons. It is a reasonable expectation that as islands become home to increasing

numbers of insurance populations or endangered species, the benefit of maintaining those islands

as pest-free (measured as the cost of restoration) will increase. In addition, as toads establish

onto an increasing number of these islands, those remaining toad-free will, by their scarcity

alone, possess a greater economic and environmental value. While in reality it is unlikely that all

islands without quarantine will be invaded, the benefit of quarantine within our dataset is held

primarily in a few large islands (e.g. Melville Island, Table 1). These larger islands often have

human settlements, competing management objectives (e.g., economic growth activities, multi-

species quarantine) or more convoluted invasion pathways associated with anthropogenic

activity. In short, quarantine needs to be carefully managed on these large islands. Eradication

efforts for taxa other than toads have been successful on large islands (such as Santiago Island 5

465km^2; Cruz *et al*. 2009), but they require much greater planning, intersectional management,

and investment in post-eradication surveillance and monitoring (Moore *et al*. 2010, Rout *et al*.

2011, Carwardine *et al*. 2012). On these large islands, then, the value of prevention is very likely

underestimated.

The vanguard of the cane toad invasion is currently sweeping across Western Australia at ~50

km per annum, but recent research suggests that a waterless barrier between the Kimberley and

the Pilbara could halt the toad invasion (Florance et al. 2011; Tingley et al. 2013; Southwell et al.

2017). Applying our results to this management strategy revealed that the benefit of quarantine

over such an area ($34.3 – $63.6 million) is roughly double the value of quarantine across all

offshore islands combined ($20.3 – $38.5 million). The cost of quarantine in this case has been

rigorously estimated at around $5 million dollars over 50 years (Southwell et al. 2017), only a

fraction of what we estimate it would cost to eradicate toads from this area.

Here we demonstrate the immense benefit of toad quarantine across northern Australia. We

avoid arbitrary judgement and simply calculate the cost of eradication in the case of quarantine

failure. We acknowledge that this value is undoubtedly a lower boundary on the true benefit, but

valuing preventative management is important. It becomes more so as conservation actions

increasingly rely on offshore islands and fenced areas as cost-effective avenues to protect

biodiversity from the impacts of invasive species. Quarantine measures often protect against

multiple potential invaders but our results suggest that even when considering a single species,

the monetary value of quarantine can be substantial. Prevention, it seems, is worth more than we

might naively guess, even with aphorisms to remind us.

Author's Contributions

AS and BP contributed to all aspects of the manuscript. RT contributed to data collection and

writing of the manuscript.

Acknowledgments

We recognise and thank the Kenbi traditional Owners (Raylene and Zoe Singh) for land access

permission. We thank Chris Jolly, Sarah McGoll-Nicolsen, John 'Mango' Moreen and the Kenbi

Ranger Group for their aid in the field, and for logistical support. Corrin Everitt, John Llewelyn,

Ruchira Somaweera, and Greg Clarke provided constructive comments and advice. We also

thank Greg Smith from Lake Argyle Cruises for his input and local knowledge. All procedures

were approved by the University of Melbourne Animal Ethics Committee (1714277.1). This

research was supported by an Australian Research Council Future Fellowship to BP

(FT160100198) and an Australian Research Council DECRA to RT (DE170100601). Land

access was granted via the Northern Land Council (permit 82368).

Data Accessibility

Waterbody data are available upon request from Geographic Information Systems at the

Department of Agriculture and Food, WA, and the Spatial Data Exchange section of the

Department of water, WA. Cost data associated with our analyses can be found in the Supporting

Information.

Table 1: Islands included in analyses from the top 100 islands prioritized by the Australian Commonwealth for conservation actions

(Department of the Environment, Water, Heritage and the Arts (2009)). Estimates for the benefit of quarantine are in '000s (AUD).

Mean benefit reports the cost of removal, averaging over costs calculated with the detection probabilities of each of our island

systems.

Jurisdiction	Island Name	Toads Present	Distance to mainland (km)	Area (km^2)	Length of freshwater shoreline (km)	Mean benefit of quarantine (000s)	Lower Est.	Upper Est.
New South Wales	Lord Howe Island	No	570	11	1	18	10	28
Western Australia	Barrow Island	No	56	139	21	373	200	580
	Bernier Island	No	38	171	2	36	19	55
	East Intercourse Island	No	5.5	51	2	36	19	55
	Faure Island	No	6.1	8	2	36	19	55
Queensland	Badu Island	Yes	90	53	10	178	95	276
	Bentineck Island	Yes	25	269	5	89	48	138
	Boigu Island	Yes	7.8	6	55	977	524	1519
	Darnley Island	Yes	70	195	0	18	10	28
	Dunk Island	Yes	4	170	1	18	10	28
	Goold Island	Yes	15	101	1	18	10	28
	Hammond Island	Yes	18	104	3	53	29	83
	Horn Island	Yes	16.7	396	8	142	76	221
	Macleay Island	Yes	3	16	0.7	12	7	19
	Magnetic Island	Yes	6.3	6	2	36	19	55
	Moa Island	Yes	52	72	21	373	200	580
	Moreton Island	Yes	20	7	54	959	514	1491
	Mornington Island	Yes	29	1662	102	1812	971	2817
	North Stradbroke Island	Yes	3.8	1001	105	1865	1000	2900
	Prince of Wales Island	Yes	16	148	27	480	257	746
	Sweers Island	No	30	7	4	71	38	110

Northern Territory	Bathurst Island	No	61	235	137	2434	1305	3783
	Centre Island	Yes	7.8	64	20	355	190	552
	Croker Island	No	3	11	152	2700	1447	4197
	Groote Eylandt	No	45	42	203	3606	1933	5606
	Marchinbar Island	No	21	5	59	1048	562	1629
	Melville Island	No	24	2	1054	18724	10036	29106
	North Island	Yes	28	13	3	53	29	83
	Peron Island	No	3.4	3	3	53	29	83
	Raragala Island	No	36	52	11	195	105	304
	Vanderlin Island	Yes	7	6	68	1208	647	1878
	West Island	Yes	4	576	30	533	286	828
	Yabooma Island	No	2.7	2	3	53	29	83

375 Figure 1. Numbers of individual cane toads captured per night on Horan (gray) and Indian

376 (black) Islands.

Figure 2. Distributions of the estimated detection probabilities of cane toads on Horan and Indian

Islands.

Figure 3. Estimated distributions of cane toad population size estimates (N_0,) before removal
efforts.

387 Figure 4. Estimated density of cane toads on each island using density calculated per island, per

388 km, and per km^2. There is clear concordance across islands when we calculate a linear density

389 (per km).

Figure 5. Distributions of the estimated numbers of days required to remove cane toads from
Horan and Indian Islands.

Figure 6. Costs of eradication calculated per km of shoreline, per square kilometre, and per island. Again, we see the strong concordance in costs when we calculate them per km of shoreline.

397
398
399

Figure 7. Distribution of the benefit of cane toad quarantine across different jurisdictions within Australia. Toad present distributions denote areas where toads are known to occur, and represent the cost to remove toads. No islands in either New South Wales, Western Australia or the Pilbara Bioregion have confirmed toad presence.

402
403
404
405

Supporting Information

Supporting Table 1: Estimated costs of cane toad eradication derived from removal efforts in this study. All figures are in Australian Dollars ($AU).

Study	Cost	Mea		Low	Uppe
Horan	Consumable	6 330	±	4	9 320
	Personnel	50	±	32	74
	Travel	16	±	10	24
Indian	Consumable	1 595	±	855	2 480
	Personnel	12	±	6	19
	Travel	4 142	±	2	6 438

References

1. Adam, M. J. & Pearl, C. A. 2007. Problems and opportunities managing invasive
bullfrogs: is there any hope? In: Gherardi, F. (ed). *Biological invaders in inland waters:*
profiles, distribution, and threats, pp. 679-693. Springer, Netherlands.

2. Beachy, J. R., Neville, R. & Arnott, C. 2011. Successful control of an incipient invasive
amphibian: *Eleutherodactylus coqui* on O'ahu, Hawai'i. *Island invasives: eradication*
and management. IUCN, Gland, Switzerland pp. 140-147.

3. Burns, B., Innes, J. & Day, T. 2012. The use and potential of pest proof fencing for
ecosystem restoration and fauna conservation in New Zealand. In 'Fencing for
Conserva8tion'. (Eds M. J. Sommers & M. Hayward) pp. 65-90. (Springer: New York).

4. Carwardine, J., O'Connor, T., Legge, S., Mackay, B., Possingham, H. P. & Martin, T. G.
2012. Prioritizing threat management for biodiversity conservation. *Conservation Letters*
5: 196-204.

5. Child, T., Phillips, B. L., Brown, G. P. and Shine, R. 2008. The spatial ecology of cane
toads (*Bufo marinus*) in tropical Australia: Why do metamorph toads stay near water?
Austral Ecology **33**: 630-640.

6. Cohen, M. P & Alford, R. A. 1993. Growth, Survival and Activity Patterns of Recently
Metamorphosed *Bufo marinus*. *Wildlife research* **20**: 1-13.

7. Commonwealth of Australia (2015). Threatened species strategy. Commonwealth of
Australia, Canberra. Available at
http://www.environment.gov.au/biodiversity/threatened/publications/strategy-home.

8. Cruz, F., Carrion, V., Campell, K. J., Lavoire, C. & Donlan, J. C. 2009. Bio-Economics
of Large-scale eradication of feral goats from Santiago island, Galapagos. *Journal of*
wildlife management **73**: 191-200.

9. Cuicui, C., Epanchin0Neil, R. & Haight, R. 2018. Optimal inspection of imports to
prevent invasive pest introduction. *Risk Analysis* **38**: 603 – 619.

10. Department of the Environment and Energy (2011). Environment Protection and
Biodiversity Conservation Act 1999 (EPBC Act). Retrieved from
http://www.environment.gov.au/epbc

11. Department of Environment and Energy (2011). The biological effects, including lethal
toxic ingestion, caused by Cane Toads (*Bufo marinus*). Retrieved from
http://www.environment.gov.au/system/files/resources/2dab3eb9-8b44-45e5-b249-
651096ce31f4/files/tap-cane-toads.pdf

12. Department of the Environment, Water, Heritage and the Arts (2009). Prioritization of
high conservation status offshore islands. Retrieved from
https://www.environment.gov.au/system/files/resources/5325cdf1-b56f-43b3-8bef-
052d740d93fd/files/offshore-islands.pdf

13. Department of the Environment and Energy (2016). Feral Animals on Offshore Islands
451   Database. Retrieved from http://www.environment.gov.au/biodiversity/invasive-
452   species/feral-animals-australia/offshore-islands

453 14. Epanchin-Neill, R. & Leibhold, A. 2015. Benefits pf invasion prevention Effect of time
454   lags, spread rates, and damage persistence. *Ecological Economics* **116**: 146-153.

455 15. Freeland, W. 1986. Populations of cane toad *Bufo marinus* in relation to time since
456   colonization. Wildlife Research **13**: 321–330. doi:10.1071/ WR9860321

457 16. Florance, D., Webb, J. K, Dempster, T., Kearney, M. R, Worthing, A. and Letnic, M.
458   2011. "Excluding Access to Invasion Hubs Can Contain the Spread of an Invasive
459   Vertebrate." *Proceedings of the Royal Society B: Biological Sciences* **278**: 2900–2908.
460   doi:10.1098/rspb.2011.0032.

461 17. Gelman, A., and Rubin, D. B.1992. Inference from Iterative Simulation Using Multiple
462   Sequences. *Statistical Science* **7**: 457–511.

463 18. Greenlees, M. J., Phillips, B. L. and Shine, R. 2010. Adjusting to a Toxic Invader: Native
464   Australian Frogs Learn Not to Prey on Cane Toads. *Behavioral Ecology* **21**: 966–71.
465   doi:10.1093/beheco/arq095.

466 19. Greenlees, M. J., Harris, S., White, A. & Shine, R. 2018. The establishment and
467   eradication of an extra-limital population of invasive cane toads. *Biological Invasions* **20**:
468   2077-2089.

469 20. Griffiths, A. & McKay, J. L. 2007. Cane toads reduce the abundance and site occupancy
470   of Merten's water monitor (*Varanus mertensi*). *Wildlife Research* **34**: 609-615.

471 21. Hoffmann, B D. and Broadhurst, L. M. 2016. The Economic Cost of Managing Invasive
472   Species in Australia. *NeoBiota* **31**: 1–18. doi:10.3897/neobiota.31.6960.

473 22. Holden, M., Nyrop, J. & Ellner, S. 2016. The economic benefit of time-varying
474   surveillance effort for invasive species management. *Journal of Applied Ecology* **53**: 712
475   – 721.

476 23. Jardine, S. L. and Sanchirico, J. N. 2018. Estimating the Cost of Invasive Species
477   Control. *Journal of Environmental Economics and Management* **87**: 242–57.
478   doi:10.1016/j.jeem.2017.07.004.

479 24. Kelehear, C., Webb, J. K. & Shine, R. 2003. *Rhabdias pseudosphaerocephala* infection
480   in *Bufo marinus*: ling nematodes reduce viability of metamorph cane toads. *Parasitology*
481   **138**: 919-927.

482 25. Kelehear, C., Brown, G. P. & Shine, R. 2011. Influence of lung parasites on the growth
483   rates of free-ranging and captive adult cane toads. *Oecologia* **165**: 585-592.

484 26. Kimball, S., Lulow, M., Sorenson, Q., Balazs, K., Fang, Y., Davis, S., O'Connell, M. &
485   Huxman, T. 2014. Cost-effective ecological restoration. *Restoration Ecology* **23**.
486   10.1111/rec.12261.

27. Kraus, F. 2009. *Alien reptiles and amphibians: a scientific compendium and analysis*.
Springer Science and Business Media B. V., Dordrecht, Netherlands.

28. Legge, S., Woinarski, J., Burbidge, A., Palmer, R., Ringma, J., Radford, J., Mitchell, N.,
Bode, M., Wintle, Br., Baseler, M., Bentley, J., Copley, P., Dexter, N., Dickman, C.,
Gillespie, G., Hill, B., Latch, P., Letnic, Mi. & Tuft, K. 2018. Havens for threatened
Australian mammals: the contributions of fenced areas and offshore islands to the
protection of mammal species susceptible to introduced predators. *Wildlife Research*. **10**.
1071/WR17172.

29. Leung, B., Lodge, D. M., Finnoff, D., Shogren, J. F., Lewis, M. A. and Lamberti, G.
2002. An Ounce of Prevention or a Pound of Cure: Bioeconomic Risk Analysis of
Invasive Species. *Proceedings of the Royal Society B: Biological Sciences* **269** (1508):
2407–13. doi:10.1098/rspb.2002.2179.

30. Llewelyn, J., Schwarzkopf, L., Phillips, B. L. and Shine, R. 2014. After the Crash: How
Do Predators Adjust Following the Invasion of a Novel Toxic Prey Type?: Adjusting to a
Novel Toxic Prey Type. *Austral Ecology* **39** (2): 190–97. doi:10.1111/aec.12058.

31. Letnic, M., Webb, J. K., Jessop, T. S. & Dempster, T. 2015. Restricting access to
invasion hubs enables sustained control of an invasive vertebrate. *Journal of Applied*
Ecology **52**: 341-347.

32. Moore, J. L., Rout, T. M., Hauser, C. E., Moro, D., Jones, M., Wilcox, C. & Possingham,
H. P. 2010. Protecting islands from pest invasion: optimal allocation of biosecurity
resources between quarantine and surveillance. *Biological Conservation* **143**: 1068-1078.

33. Moro, D., Ball, D. & Bryant, S (eds) 2018. *Australian Island Arks: conservation,*
management and opportunities, CSIRO publishing, Clayton South.

34. Moseby, K., Read, J., Paton, D., Copley, P., Hill, B. & Crisp, H. 2011. Predation
determines the outcome of 10 reintroduction attempts in arid South Australia. *Biological*
Conservation **144** 2863-2872.

35. Nelson, D. W. M., Crossland, M. R. and Shine, R. 2010. Indirect Ecological Impacts of
an Invasive Toad on Predator–prey Interactions Among Native Species. *Biological*
Invasions **12** (9): 3363–9. doi:10.1007/s10530-010-9729-4.

36. Orchard, S. A. 2011. Removal of the American bullfrog *Rana (Lithobates) catesbeiana*
from a pond and a lake on Vancouver Island, British Columbia, Canada. In: Veitch, C.
R.; Clout, M. N. and Towns, D. R. (eds). 2011. *Island invasives: eradication and*
management. IUCN, Gland, Switzerland. In: Gherardi, F. (ed). *Biological invaders in*
inland waters: profiles, distribution, and threats, pp. 679-693. Springer, Netherlands.

37. Phillips, B. L., Brown, G. P. and Shine, R. 2010. Evolutionarily Accelerated Invasions:
The Rate of Dispersal Evolves Upwards During the Range Advance of Cane Toads:
Dispersal Evolution During Range Advance. *Journal of Evolutionary Biology* **23** (12):
2595–2601. doi:10.1111/j.1420-9101.2010.02118.x.

38. Pikacha, P., Lavery, T. and Leung, L. K. P. 2015. What Factors Affect the Density of
Cane Toads (Rhinella Marina) in the Solomon Islands? *Pacific Conservation Biology* **21**
(3): 200. doi:10.1071/PC14918.

39. Pizzatto, L. & Shine, R. 2008. The behavioral ecology of cannibalism in cane toads (*Bufo*
marinus). *Behavioral Ecology and Sociobiology* **63**: 123-133.

40. Plummer, Martyn. 2013. rjags: Bayesian graphical models using MCMC. R package
version 3-10. URL: http://CRAN.R-project.org/package=rjags

41. Reardon, J. T., Kraus, F., Moore, M., Rabenantenaina, L., Rabinivo, A., Nantenaina, H.,
Randrianasolo, H & Randrianasolo, R. 2018. Testing tools for eradication the invasive
toad *Duttaphynus melanosticus* in Madagascar. *Conversation Evidence* **15**: 12-19.

42. Ringma, J., Legge, S., Woinarski, J., Radford, J., Wintle, B. and Bode, M. 2018.
Australia's mammal fauna requires a strategic and enhanced network of predator-free
havens. *Nature Ecology & Evolution* **2**: 410-411.

43. Rohr, J. R., Farag, A. M., Cadotte, M. W., Clements, W. H., Smith, J. R., Ulrich, C. P., &
Woods, R. 2016. Transforming ecosystems: When, where, and how to restore
contaminated sites. *Integrated environmental assessment and management* **12**: 273-83.
541

542 44. Rout, T. M., Moore, J. L., Possingham, H. P. & McCarthy, M. 2011. Allocating
543   biosecurity resources between preventing, detecting, and eradication island invasions.
544   *Ecological Economics* **71**: 54-62.
545

546 45. Seebacher, F. & Alfrod, R. A. 2002. Shelter microhabitats determine body temperature
547   and dehydration rates of a terrestrial amphibian (*Bufo marinus*). *Journal of Herpetology*
548   **36**: 69-75.
549

46. Shanmuganathan, T., Pallister, J., Doody, S., McCallum, H., Robinson, T., Sheppard, A.,
Hardy, C., Halliday, D., Venables, D., Voysey, R., Strive, T., Hinds, L. and Hyatt, A.
2010. Biological Control of the Cane Toad in Australia: A Review: Biological Control of
Cane Toad. *Animal Conservation* **13**: 16–23. doi:10.1111/j.1469-1795.2009.00319.x.

47. Somaweera, R. & Shine, R. 2012. The (non) impact of invasive cane toads on freshwater
crocodiles at Lake Argyle in tropical Australia. *Animal Conservation* **15**: 152-163.

48. Southwell, D., Tingley, R., Bode, M., Nicholson, E. and Phillips, B. L 2017. Cost and
Feasibility of a Barrier to Halt the Spread of Invasive Cane Toads in Arid Australia:
Incorporating Expert Knowledge into Model-Based Decision-Making. *Journal of Applied*
Ecology **54** (1): 216–24. doi:10.1111/1365-2664.12744.

49. Tershy, B. R., Shen, K., Newton, K. M., Holmes, N. D. & Croll, D. A. 2015. The
importance of islands for the protection of biological and linguistic diversity. *Bioscience*
65: 592–597. doi:10.1093/biosci/biv031

50. Thomas, C. D. 2011. Translocation of Species, Climate Change, and the End of Trying to
Recreate Past Ecological Communities. *Trends in Ecology & Evolution* **26** (5): 216–21.
doi:10.1016/j.tree.2011.02.006.

51. Tingley, R. & Shine, R. 2011. Desiccation risk drives the spatial ecology in an invasive
anuran (*Rhinella marina*) in the Australian Semi-desert. *PLoS ONE* **6:** e25979.
doi.org/10.1371/journal.pone.0025979
569

52. Tingley, R., Phillips, B. L., Letnic, M., Brown, G. P., Shine, R. & Baird, S. J. E. 2013.
Identifying Optimal Barriers to Halt the Invasion of Cane Toads *Rhinella Marina* in Arid
Australia. *Journal of Applied Ecology* **50** (1): 129–37. doi:10.1111/1365-2664.12021.

53. Tingley, R., Ward-Fear, G., Schwarzkopf, L., Greenlees, M. J., Phillips, B. L., Brown, G.,
Clulow, S., Webb, J., Capon, R., Sheppard, A., Strive, T., Tizard, M. & Shine, R. 2017.
New weapons in the toad toolkit a review of methods to control and mitigate the
biodiversity impact of invasive cane toad (*Rhinella Marina*). *The Quarterly Reviews of*
Biology **92**: 123-149.

54. Ward-Fear, G., Brown, G. P. & Shine, R. 2010. Using a Native Predator (the Meat Ant,
Iridomyrmex Reburrus) to Reduce the Abundance of an Invasive Species (the Cane
Toad, *Bufo Marinus*) in Tropical Australia. *Journal of Applied Ecology* **47** (2): 273–80.
doi:10.1111/j.1365-2664.2010.01773.x.

55. White, A. 2010. Cane toad outbreak: Taren Point, 2010. Report prepared by Biosphere
Environmental Consultants Pty. Ltd. For Sutherland Shire Council, NSW.

56. Wingate, D. B. 2011. The successful elimination of Cane Toad, *Bufo marinus*, from an
island with breeding habitat off Bermuda. *Biological Invasions* **13**: 1487-1492.

57. Woinarski, J., Burbridge, A. & Harrion P. 2014. The Action Plan for Australian
Mammals 2012. CSIRO Publishing: Melbourne.

58. Zug, G., Lindgren, E. & Pippet, J. 1975. Distribution and ecology of the marine toad,
Bufo marinus, in Papua New Guinea. *Pacific Science* **29**:31–50.
590