# An efficient and robust laboratory workflow and tetrapod database for larger scale eDNA studies

Jan Axtner[1+], Alex Crampton-Platt[1], Lisa A. Hörig[1], Azlan Mohamed[1], Charles C.Y. Xu[2,3,4], Douglas W. Yu[2,5] and Andreas Wilting[1]

**Affiliations:**
[1] Leibniz Institute for Zoo and Wildlife Research (*Leibniz-IZW*), Alfred-Kowalke-Str. 17, 10315 Berlin, Germany
[2] State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, 32 Jiaochang East Road, Kunming, Yunnan 650223, China
[3] Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands
[4] Redpath Museum and Department of Biology, McGill University, Montreal, QC, Canada
[5] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ, UK

[+] corresponding author

**email addresses:**
Jan Axtner:            axtner@izw-berlin.de
Alex Crampton-Platt:  alex@naturemetrics.co.uk
Lisa A. Hörig:         lisa.hoerig@arcor.de
Azlan Mohamed:        mohamed@izw-berlin.de
Charles C.Y. Xu:      charles.cong.xu@gmail.com
Douglas W. Yu:        dougwyu@mac.com
Andreas Wilting:      wilting@izw-berlin.de

**Keywords:**
metabarcoding, iDNA, eDNA, leeches

## Abstract

### Background

The use of environmental DNA, 'eDNA,' for species detection via metabarcoding is growing rapidly. We present a co-designed lab workflow and bioinformatic pipeline to mitigate the two most important risks of eDNA: sample contamination and taxonomic mis-assignment. These risks arise from the need for PCR amplification to detect the trace amounts of DNA combined with the necessity of using short target regions due to DNA degradation.

### Findings

Our high-throughput workflow minimises these risks via a four-step strategy: (1) technical replication with two *PCR replicates* and two *extraction replicates*; (2) using multi-markers (*12S*, *16S*, *CytB*); (3) a 'twin-tagging,' two-step PCR protocol;(4) use of the probabilistic taxonomic assignment method *PROTAX*, which can account for incomplete reference databases.

As annotation errors in the reference sequences can result in taxonomic mis-assignment, we supply a protocol for curating sequence datasets. For some taxonomic groups and some markers, curation resulted in over 50% of sequences being deleted from public reference databases, due to (1) limited overlap between our target amplicon and reference sequences; (2) mislabelling of reference sequences; (3) redundancy.
Finally, we provide a bioinformatic pipeline to process amplicons and conduct *PROTAX* assignment and tested it on an 'invertebrate derived DNA' (iDNA) dataset from 1532 leeches from Sabah, Malaysia. Twin-tagging allowed us to detect and exclude sequences with non-matching tags. The smallest DNA fragment (*16S*) amplified most frequently for all samples, but was less powerful for discriminating at species rank. Using a stringent and lax acceptance criteria we found 170 (stringent) and 196 (lax) vertebrate detections of 97 (stringent) and 109 (lax) leech samples.

### Conclusions

Our metabarcoding workflow should help research groups increase the robustness of their results and therefore facilitate wider usage of e/iDNA, which is turning into a valuable source of ecological and conservation information on tetrapods.

## Introduction

Monitoring, or even detecting, elusive or cryptic species in the wild can be challenging. In recent years there has been a rise in the availability of cost-effective DNA-based methods made possible by advances in high-throughput DNA sequencing (HTS). One such method is eDNA metabarcoding, which seeks to identify the species present in a habitat from traces of 'environmental DNA' (eDNA) in substrates such as water, soil, or faeces. A variant of eDNA metabarcoding, known as 'invertebrate-derived DNA' (iDNA) metabarcoding, targets the genetic material of prey or host species extracted from copro-, sarco- or haematophagous invertebrates. Examples include tick [1] s, blow or carrion flies [2; 3; 4; 5], mosquitoes [6; 7; 8; 9] and leeches [10; 11; 12; 13]. Many of these parasites are ubiquitous, highly abundant, and easy to collect, making them an ideal source of biodiversity data, especially for terrestrial vertebrates that are otherwise difficult to detect [10; 14; 15]. In particular, the possibility for bulk collection and sequencing in order to screen large areas and minimise costs is attractive. However, most of the recent studies on iDNA studies focus on single-specimen DNA extracts and Sanger sequencing and thus are not making use of the advances of HTS and a metabarcoding framework for carrying out larger scale biodiversity surveys.

That said, e/iDNA metabarcoding also poses several challenges, due to the low quality and low amounts of target DNA available, relative to non-target DNA (including the high-quality DNA of the live-collected, invertebrate vector). In bulk iDNA samples comprised of many invertebrate specimens, this problem is further exacerbated by the variable time since each individual has fed, if at all, leading to differences in the relative amounts and degradation of target DNA per specimen. This makes e/iDNA studies similar to ancient DNA samples, which also pose the problem of low quality and low amounts of target DNA [16; 17]. The great disparity in the ratio of target to non-target DNA and the low overall amount of the former requires an enrichment step, which is achieved via the amplification of a short target sequence (amplicon) by polymerase chain reaction (PCR) to obtain enough target material for sequencing. However, this enrichment step can result in false positive species detections, either through sample cross-contamination or through volatile short PCR amplicons in the laboratory, and in false-negative results, through primer bias and low concentrations of template DNA. Although laboratory standards to prevent and control for such false results are well established in the field of ancient DNA, there are still no best-practice guidelines for e/iDNA studies, and thus few studies sufficiently account for such problems [18].

The problem is exacerbated by the use of 'universal' primers used for the PCR, which maximise the taxonomic diversity of the amplified sequences. This makes the method a powerful biodiversity assessment tool, even where little is known *a priori* about which species might be found. However, using such primers, in combination with low quality and quantity of target DNA, which often requires a high number of PCR cycles to generate enough amplicon products for sequencing, makes metabarcoding studies particularly vulnerable to false results [13; 19; 20]. The high number of PCR cycles, combined with the

high sequencing depth of HTS, also increase the likelihood that contaminants are amplified and detected, possibly to the same or greater extent as some true-positive trace DNA. As e/iDNA have been proposed as tools to detect very rare and priority conservation species such as the Saola, *Pseudoryx nghetinhensis* [10], false detection might result in misdirected conservation activities worth several hundreds of thousands of US dollars like for the ivory-billed woodpecker where most likely false evidence of the bird's existence have been overemphasized to shore up political and financial support for saving it [21]. Therefore, similar to ancient DNA studies, great care must be taken to minimise the possibility for cross-contamination in the laboratory and to maximise the correct detection of species through proper experimental and analytical design. Replication in particular is an important tool for reducing the incidence of false negatives and detection of false positives but the trade-off is increased cost, workload, and analytical complexity [19].

An important source of false positive species detections is the incorrect assignment of taxonomies to the millions of short HTS reads generated by metabarcoding. Although there has been a proliferation of tools focused on this step, most can be categorised into just three groups depending on whether the algorithm utilises sequence similarity searches, sequence composition models, or phylogenetic methods [22; 23; 24]. The one commonality among all methods is the need for a reliable reference database of correctly identified sequences, yet there are few curated databases currently appropriate for use in e/iDNA metabarcoding. Two exceptions are SILVA [25] for the nuclear markers SSU and LSU rRNA used in microbial ecology, and BOLD (Barcode of Life Database; citation) for the COI 'DNA barcode' region. For other loci, a non-curated database downloaded from the INSDC (International Nucleotide Sequence Database Collaboration, e.g. GenBank) is generally used. However, the INSDC places the burden for metadata accuracy, including taxonomy, on the sequence submitters, with no restriction on sequence quality or veracity. For instance, specimen identification is often carried out by non-specialists, which increases error rates, and common laboratory contaminant species (e.g. human DNA sequences) are sometimes submitted in lieu of the sample itself. The rate of sequence mislabelling in fungi has been assessed for GenBank where it was up to 20% [26] and it is an issue that is often neglected [27; 28]. For several curated microbial databases (Greengenes, LTP, RDP, SILVA), mislabelling rates have been estimated at between 0.2% and 2.5% [29]. Given the lack of professional curation it is likely that the true proportion of mislabelled samples in GenBank is somewhere between these numbers. Moreover, correctly identifying such errors is labour-intensive, so most metabarcoding studies simply base their taxonomic assignments on sequence-similarity searches of the whole INSDC database (e.g. with BLAST) [3; 10; 12] and thus can only detect errors if assignments are ecologically unlikely. Furthermore, reference sequences for the species that are likely to be sampled in e/iDNA studies are often underrepresented in or absent from these databases, which increases the possibility of incorrect assignment. For instance, fewer than 50% of species occurring in a tropical megadiverse rainforest are represented in Genbank (see findings below). When species-level matches are ambiguous, it might still be possible to assign a sequence to a higher

taxonomic rank by using an appropriate algorithm such as Metagenome Analyzer's
(MEGAN) Lowest Common Ancestor [30] or *PROTAX* [31].

We present here a complete laboratory workflow and complementary bioinformatics
pipeline, starting from DNA extraction to taxonomic assignment of HTS reads using a
curated reference database. The laboratory workflow allows for efficient screening of
hundreds of e/iDNA samples. The workflow includes (1) two *extraction replicates* are
separated during DNA extraction, and each is sequenced in two *PCR replicates* (Fig. 1); (2)
robustness of taxonomic assignment is improved by using up to three mitochondrial
markers; (3) a 'twin-tagged', two-step PCR protocol prevents cross-sample contamination as
no unlabelled PCR products are produced (Fig. 2) while also allowing for hundreds of PCR
products to be pooled before costly Illumina library preparation; (4) our bioinformatics
pipeline includes a standardized, automated, and replicable protocol to create a curated
database, which allows updating as new reference sequences become available, and to be
expanded to other amplicons. We provide scripts for processing raw sequence data to
quality-controlled dereplicated reads and for taxonomic assignment of these reads using
*PROTAX* [31], a probabilistic method that has been shown to be robust even when reference
databases are incomplete [23; 4] (all scripts are available from URL
https://github.com/alexcrampton-platt/screenforbio-mbc).

## Methods

### Establishment of the tetrapod reference database

*Reference database*

A custom bash script was written to generate a tetrapod reference database for up to four
mitochondrial markers – a short 93 bp fragment of *16S* rRNA (*16S*), a 389 bp fragment of
*12S* rRNA (*12S*), a 302 bp fragment of cytochrome b (*CytB*), and a 250 bp mitochondrial
cytochrome c oxidase subunit I amplicon (*COI*) that has previously been used in iDNA studies
[2]. An important time-saving step was the use of the FASTA-formatted Midori
mitochondrial database [32], which is a lightly curated subset of Genbank. Our script
updated the FASTA files with a subset of target species, removed errors and redundancy,
trimmed the sequences to include only the amplicon regions, and output FASTA files with
species names and GenBank accessions in the headers.

The script accepts four data inputs, two of which are optional. The required inputs are: (i)
the Midori sequences (December 2015 'UNIQUE', downloaded from http://www.reference-
midori.info/download.php#) for the relevant genes and (ii) an initial reference taxonomy of
tetrapods. This taxonomy is needed to find or generate a full taxonomic classification for
each sequence because the taxonomies in Midori are from Genbank and thus include
incorrect, synonymized, or incomplete taxonomies. Here we used the Integrated Taxonomic
Information System (ITIS) classification for Tetrapoda, obtained with the R package *taxize*
version 0.9.0 ([33], functions *downstream* and *classification*). The optional inputs are: (iii)
supplementary FASTA files of reference sequences that should be added to the database,

and (iv) a list of target species to be queried on GenBank to capture any sequences
published since the December 2015 Midori dataset was generated.

For this study, 72 recently published [34] and 7 unpublished partial mitochondrial mammal
genomes (Accession Numbers MH464789, MH464790, MH464791, MH464792, MH464793,
MH464794, MH464795, MH464796, MH464797, MH464798, MH464799, MH464800,
MH464801) were added as input (iii). A list of 103 mammal species known to be present in
the sampling area plus *Homo sapiens* and our positive control *Myodes glareolus* was added
as input (iv).

With the above inputs, the seven curation steps are: 1) remove sequences not identified to
species; 2) add extra sequences from optional inputs (iii) and (iv) above; 3) trim the
sequences to leave only the target amplicon; 4) remove sequences with ambiguities; 5)
compare species names from the Midori dataset to the reference taxonomy from input (ii)
and replace with a consensus taxonomy; 6) identify and remove putatively mislabelled
sequences; 7) dereplicate sequences, retaining one haplotype per species.

The script is split into four modules, allowing optional manual curation at three key steps.
The steps covered by each of the four modules are summarized in Table 2. The main
programs used are highlighted and cited in the text where relevant, but many intermediate
steps used common UNIX tools and unpublished lightweight utilities freely available from
GitHub (Table 3).

*Module 1* - The first step is to select the tetrapod sequences from the Midori database for
each of the four selected loci (input (i) above). This, and the subsequent step to discard
sequences without strict binomial species names and reduce subspecies identifications to
species-level, are made possible by the inclusion of the full NCBI taxonomic classification of
each sequence in the FASTA header by the Midori pipeline. The headers of the retained
sequences are then reformatted to include just the species name and GenBank accession
separated by underscores. If desired, additional sequences from local FASTA files are now
added to the Midori set (input (iii)). The headers of these FASTA files are required to be in
the same format. Next, optional queries are made to the NCBI GenBank and RefSeq
databases for each species in a provided list (input (iv)) for each of the four target loci, using
NCBI's Entrez Direct [35]. Matching sequences are downloaded in FASTA format, sequences
prefixed as "UNVERIFIED" are discarded, the headers are simplified as previously, and those
sequences not already in the Midori set are added. Trimming each sequence down to the
relevant target marker was carried out in a two-step process in which *usearch* (*-search_pcr*)
was used to select sequences where both primers were present, and these were in turn
used as a reference dataset for *blastn* to select partially matching sequences from the rest
of the dataset [36; 37]. Sequences with a hit length of at least 90% of the expected marker
length were retained by extracting the relevant subsequence based on the BLAST hit co-
ordinates. Sequences with ambiguous bases were discarded at this stage. In the final step in
module 1, a multiple-sequence alignment was generated with MAFFT [38; 39] for each
partially curated amplicon dataset (for the SATIVA step below). The script then breaks to

allow the user to check for any obviously problematic sequences that should be discarded
before continuing.

*Module 2* - The species labels of the edited alignments are compared with the reference
taxonomy (input (ii)). Any species not found is queried against the Catalogue of Life
database (CoL) via *taxize* in case these are known synonyms, and the correct species label
and classification is added to the reference taxonomy. The original species label is retained
as a key to facilitate sequence renaming, and a note is added to indicate its status as a
synonym. Finally, the genus name of any species not found in the CoL is searched against
the consensus taxonomy, and if found, the novel species is added by taking the higher
classification levels from one of the other species in the genus. Orphan species labels are
printed to a text file, and the script breaks to allow the user to check this list and manually
create classifications for some or all if appropriate.

*Module 3* - This module begins by checking for any manually generated classification files
(from the end of Module 2) and merging them with the reference taxonomy from Module 2.
Any remaining sequences with unverifiable classifications are removed at this step. The next
steps convert the sequences and taxonomy file to the correct formats for SATIVA [29],
which detects possibly mislabelled sequences by generating a maximum likelihood
phylogeny from the alignment in Module 1 and comparing each sequence's taxonomy
against its phylogenetic neighbors. Sequence headers in the edited MAFFT alignments are
reformatted to include only the GenBank accession, and a taxonomy key file is generated
with the correct classification listed for each accession number. In cases where the original
species label is found to be a synonym, the corrected label is used. Putatively mislabelled
sequences in each amplicon are then detected with SATIVA, and the script breaks to allow
inspection of the results. The user may choose to make appropriate edits to the taxonomy
key file or list of putative mislabels at this point.

*Module 4* - Any sequences that are still flagged as mislabelled at the start of the fourth
module are deleted from the SATIVA input alignments, and all remaining sequences are
relabelled with the correct species name and accession. A final consensus taxonomy file is
generated in the format required by *PROTAX*. Alignments are subsequently unaligned prior
to species-by-species selection of a single representative per unique haplotype. Sequences
that are the only representative of a species are automatically added to the final database.
Otherwise, all sequences for each species are extracted in turn, aligned with MAFFT, and
collapsed to unique haplotypes with *collapsetypes_4.6.pl* (zero differences allowed; [40]).
Representative sequences are then unaligned and added to the final database.

## iDNA samples

We used 242 collections of haematophagous terrestrial leeches from Deramakot Forest
Reserve in Sabah, Malaysian Borneo stored in RNA fixating saturated ammonium sulfate
solution as samples. Each sample consisted of one to 77 leech specimens (median 4). In
total, 1532 leeches were collected, exported under the permit (JKM/MBS.1000-2/3 JLD.2 (8)

issued by the Sabah Biodiversity Council), and analysed at the laboratories of the Leibniz-
IZW.

## Laboratory workflow

The laboratory workflow is designed to both minimize the risk of sample cross-
contamination and to aid identification of any instances that do occur. All laboratory steps
(extraction, pre and post PCR steps, sequencing) took place in separate laboratories and no
samples or materials were allowed to re-enter upstream laboratories at any point in the
workflow. All sample handling was carried out under specific hoods that were wiped with
bleach, sterilized, and UV irradiated for 30 minutes after each use. All labs are further UV
irradiated for four hours each night.

### DNA extraction

DNA was extracted from each sample in bulk. Leeches were cut into small pieces with a
fresh scalpel blade and incubated in lysate buffer (proteinase K and ATL buffer at a ratio of
1:10; 0.2 ml per leech) overnight at 55 °C (12 hours minimum) in an appropriately sized
vessel for the number of leeches (2 or 5 ml reaction tube). For samples with more than 35
leeches, the reaction volume was split in two and recombined after lysis.

Each lysate was split into two *extraction replicates* (A and B; maximum volume 600 µl) and
all further steps were applied to these independently. We followed the DNeasy 96 Blood &
Tissue protocol for animal tissues (Qiagen, Hilden -Germany) on 96 plates for cleanup. DNA
was eluted twice with 100 µl TE buffer. DNA concentration was measured with PicoGreen
dsDNA Assay Kit (Quant-iT, ThermoFisherScientific, Waltham -USA) in 384-well plate format
using an appropriate plate reader (200 PRO NanoQuant, Tecan Trading AG, Männedorf -
Switzerland). Finally, all samples were diluted to a maximum concentration of 10 ng/µl.

### Two-round PCR protocol

We amplified three mitochondrial markers – a short 93 bp fragment of *16S* rRNA (*16S*), a
389 bp fragment of *12S* rRNA (*12S*), and a 302 bp fragment of cytochrome b (*CytB*). For each
marker, we ran a two-round PCR protocol (Figs. 1, 2). The primers were chosen on the
expectation of successful DNA amplification over a large number of tetrapod species [41;
42], and we tested the fit of candidate primers on an alignment of available mitochondrial
sequences of 134 Southeast-Asian mammal species. Primer sequences are in Table 1.

*Primer modification.* – We modified primers of the three markers to avoid the production of
unlabelled PCR products, to allow the detection and deletion of tag-jumping events [43],
and to reduce the cost of primers and library preparation. We used two rounds of PCR. The
first round amplified the target gene and attached one of 25 different 'twin-tag' pairs (*tag*
*1*), identifying the sample within a given PCR. By 'twin-tag,' we mean that both the forward
and reverse primers were given the *same* sample-identifying sequence ('tags') added as
primer extensions (Fig. 2). The tags differed with a minimum pairwise distance of three
nucleotides ([43]; Supplemental Table 1). These primers also contained different forward

and reverse sequences (*Read 1* & *Read 2 sequence primers*) (Supplemental Table 1) to act
priming sites for the second PCR round (Fig. 2).

The second round added the Illumina adapters for sequencing and attached one of 20 twin-
tag pairs (*tag 2*) identifying the PCR, with a minimum pairwise distance of three [44]. These
primers also contained the Illumina P5 and P7 adapter sequences (Fig. 2). Thus no
unlabelled PCR products were ever produced, and the combination of *tags 1* and *2* allowed
the pooling of up to 480 (=24 X 20) samples in a single library preparation step (one *tag 1*
was reserved for controls). Twin tags allowed us later to detect and delete tag jumping
events [43] (Fig. 2).

*Cycle number considerations.* – Because we know that our target DNA is at low
concentration in the samples, we are faced with a trade-off between (1) using fewer PCR
cycles (e.g. 30) to minimise amplification bias (caused by some target DNA binding better to
the primer sequences and thus outcompeting other target sequences that bind less well
[45]) and (2) using more PCR cycles (e.g. 40) to ensure that low-concentration target DNA is
sufficiently amplified in the first place. Rather than choose between these two extremes, we
ran both low- and high-cycle protocols and sequenced both sets of amplicons.

Thus, each of the two *extraction replicates* A and B was split and amplified using different
cycle numbers (*PCR replicates* 1 and 2) for a total of four (= 2 *extraction replicates* x 2 *PCR*
*replicates* -> *A1/A2* and *B1/B2* ) replicates per sample per marker (Fig. 1). For *PCR replicates*
A1/B1, we used 30 cycles in the first PCR round to minimize the effect of amplification bias.
For *PCR replicates* A2/B2, we used 40 cycles in the first PCR round to increase the likelihood
of detecting species with very low input DNA (Fig. 1).

*PCR protocol.* – The first-round PCR reaction volume was 20 µl, including 0.1 µM primer mix,
0.2 mM dNTPs, 1.5 mM MgCl$_2$, 1x PCR buffer, 0.5 U AmpliTaq Gold™ (Invitrogen, Karlsruhe -
Germany), and 2 µl of template DNA. Initial denaturation was 5 minutes at 95°C, followed
by repeated cycles of 30 seconds at 95°C, 30 seconds at 54°C, and 45 seconds at 72°C. Final
elongation was 5 minutes at 72°C. Samples were amplified in batches of 24 plus a negative
(water) and a positive control (bank vole, *Myodes glareolus* DNA). All three markers were
amplified simultaneously for each batch of samples in a single PCR plate. Non-target by-
products were removed as required from some *12S* PCRs by purification with magnetic
Agencourt AMPure beads (Beckman Coulter, Krefeld -Germany).

In the second-round PCR, we used the same PCR protocol as above with 2 µl of the product
of the first-round PCR and 10 PCR cycles.

*Quality control and sequencing*

Amplification was visually verified after the second-round PCR by gel electrophoresis on
1.5% agarose gels. Controls were additionally checked with a TapeStation 2200 (D1000
ScreenTape assay, Agilent, Waldbronn -Germany). All samples were purified with AMPure
beads, using a bead-to-template ratio of 0.7:1 for *12S* and *CytB* products, and a ratio of 1:1
for *16S* products. DNA concentration was measured with PicoGreen dsDNA as described

above. Sequencing libraries were made by equimolar pooling of all positive amplifications;
final concentrations were between 2 and 4 nmol. Because of different amplicon lengths and
therefore different binding affinities to the flow cell, *12S* and *CytB* products were combined
in a single library, whereas positive *16S* products were always combined in a separate
library. Apart from our negative controls, we did not include samples that did not amplify, as
this would have resulted in highly diluted libraries. Up to 11 libraries were sequenced on
each run of Illumina MiSeq, following standard protocols. Libraries were sequenced with
MiSeq Reagent Kit V3 (600 cycles, 300 bp paired-end reads) and had a final concentration of
11 pM spiked with 20 to 30% of PhiX control.

## Bioinformatics workflow
### *Read processing*

Although the curation of the reference databases is our main focus, it is just one part of the
bioinformatics workflow for e/iDNA metabarcoding. A custom bash script was used to
process raw basecall files into demultiplexed, cleaned, and dereplicated reads in FASTQ
format on a run-by-run basis. All runs and amplicons were processed with the same settings
unless otherwise indicated. *bcl2fastq* (Illumina) was used to convert the basecall file from
each library to paired-end FASTQ files, demultiplexed into the separate PCRs via the *tag 2*
pairs, allowing up to 1 mismatch in each *tag 2*. Each FASTQ file was further demultiplexed
into samples via the *tag 1* pairs using *AdapterRemoval* [46], again allowing up to 1 mismatch
in each tag. These steps allowed reads to be assigned to the correct samples.

In all cases, amplicons were short enough to expect paired reads to overlap. For libraries
with more than 1000 reads pairs were merged with *usearch* (*-fastq_mergepairs*; [47; 48]),
and only successfully merged pairs were retained. For libraries with more than 500 merged
pairs the primer sequences were trimmed away with *cutadapt* [49], and only successfully
trimmed reads at least 90% of expected amplicon length were passed to a quality filtering
step with *usearch* (*-fastq_filter*). Lastly, reads were dereplicated with *usearch* (*-
derep_fulllength*), and singletons were discarded. The number of replicates that each unique
sequence represented was also added to the read header at this step (option *-sizeout*). The
number of reads processed at each step for each sample are reported in a standard tab
delimited txt-file.

### *Taxonomic assignment*

The curated reference sequences and associated taxonomy were used for *PROTAX*
taxonomic assignment of the dereplicated reads [24; 31]. *PROTAX* gives unbiased estimates
of placement probability for each read at each taxonomic rank, allowing assignments to be
made to a higher rank even when there is uncertainty at the species level. In other words,
and unlike other taxonomic assignment methods, *PROTAX* can estimate the probability that
a sequence belongs to a taxon that is not present in the reference database. This was
considered an important feature due to the known incompleteness of the reference
databases for tetrapods in the sampled location. As other studies have compared *PROTAX*

with more established methods, e.g. MEGAN [30] (see [4; 24]), it was beyond the scope of
this study to evaluate the performance of *PROTAX*.

Classification with *PROTAX* is a two-step process. Firstly, PROTAX selected a subset of the
reference database that was used as training data to parameterise a *PROTAX* model for
each marker, and secondly, the fitted models were used to assign four taxonomic ranks
(species, genus, family, order) to each of the dereplicated reads, along with a probability
estimate at each level. We also included the best similarity score of the assigned species or
genus, mined from the LAST results (see below) for each read. This was helpful for flagging
problematic assignments for downstream manual inspection, i.e. high probability
assignments based on low similarity scores (implying that there are no better matches
available) and low probability assignments based on high similarity scores (indicates
conflicting database signal from several species with highly similar sequences).

Fitting the *PROTAX* model followed Somervuo et al. [31] except that 5000 training
sequences were randomly selected for each target marker due to the large size of the
reference database. In each case, 4500 training sequences represented a mix of known
species with reference sequences (conspecific sequences retained in the database) and
known species without reference sequences (conspecific sequences omitted, simulating
species missing from the database), and 500 sequences represented previously unknown
lineages distributed evenly across the four taxonomic levels (i.e. mimicked a mix of
completely novel species, genera, families and orders). Pairwise sequence similarities of
queries and references were calculated with LAST [50] following the approach of Somervuo
et al. [31]. The models were weighted towards the Bornean mammals expected in the
sampled area by assigning a prior probability of 90% to these 103 species and a 10%
probability to all others ([31]; Supplemental Table 2). In cases of missing interspecific
variation, this helped to avoid assignments to geographically impossible taxa, especially in
case of the very short 93 bp fragment of *16S*. Maximum *a posteriori* (MAP) parameter
estimates were obtained following the approach of Somervuo et al. [24], but the models
were parameterised for each of the four taxonomic levels independently, with a total of five
parameters at each level (four regression coefficients and the probability of mislabelling).

Dereplicated reads for each sample were then classified using a custom bash script on a run-
by-run basis. For each sample, reads in FASTQ format were converted to FASTA, and
pairwise similarities were calculated against the full reference sequence database for the
applicable marker with LAST. Assignments of each read to a taxonomic node based on these
sequence similarities were made using a Perl script and the trained model for that level. The
taxonomy of each node assignment was added with a second Perl script for a final table
including the node assignment, probability, taxonomic level, and taxonomic path for each
read. Read count information was included directly in the classification output via the size
annotation added to the read headers during dereplication. All Perl scripts to convert input
files into the formats expected by *PROTAX*, *R* code for training the model following

Somervuo et al. [31], and Perl scripts for taxonomic assignment were provided by P.
Somervuo (personal communication).

*Acceptance criteria*

In total we had twelve PCR reactions per sample: two *extraction replicates A* and *B* X two
*PCR replicates* 1 and 2 per extraction replication X the three markers (Fig. 1). We applied
two different acceptance criteria to the data with different stringency regimes. One more
naive one that accepted any two positives out of the twelve *PCR replicates* (from now on
referred to as lax), and one stringent one that only accepted taxonomic assignments that
were positively detected in both *extraction replicates* (*A* & *B*, Fig. 3). Our lax approach refers
to one of the approaches of Ficetola et al. [19] where they evaluated different statistical
approaches developed to estimate occupancy in the presence of observational errors and
has been applied in other studies (e.g. [13]). The reason for conservatively omitting
assignments that appeared in only one *extraction replicate* was to rule out sample cross-
contamination during DNA extraction. In addition, we only accepted assignments with ten
or more reads per marker, if only one marker was sequenced. If a species was assigned in
more than one marker (e.g. *12S* and *16S*), we accepted the assignment even if in one
sequencing run the number of reads was below ten.

Due to the imperfect PCR amplification of markers (the small *16S* fragment amplified better
than the longer *CytB* fragment) and missing reference sequences in the database or shared
sequence motifs between species, reads sometimes were assigned to species level for one
marker but only to genus level for another marker. Thus, the final identification of species
could not be automated, and manual inspection and curation was needed. For each
assignment, three parameters were taken into consideration: number of sequencing reads,
the mean probability estimate derived from *PROTAX*, and the mean sequence similarity to
the reference sequences based on LAST.

*Shot-gun sequencing to quantify mammalian DNA content*

As the success of the metabarcoding largely depends on the mammal DNA quantity in our
leech bulk samples we quantified the mammalian DNA content in a subset of 58 of our leech
samples using shotgun sequencing. Extracted DNA was sheared with a Covaris M220
focused-ultra-sonicator to a peak target size of 100-200 bp, and re-checked for size
distribution. Double-stranded Illumina sequencing libraries were prepared according to a
ligation protocol designed by Fortes and Paijmans [51] with single 8 nt indices. All libraries
were pooled equimolarly and sequenced on the MiSeq using the v3 150-cycle kit. We
demultiplexed reads using bcl2fastq and cutadapt for trimming the adapters. We used
BLAST search to identify reads and applied Metagenome Analyzer MEGAN [30] to explore
the taxonomic content of the data based on the NCBI taxonomy. Finally we used KRONA
[52] for visualisation of the results.

## Findings & Discussion

### Database curation

The Midori UNIQUE database (December 2015 version) contains 1,019,391 sequences across the four mitochondrial loci of interest (*12S*: 66,937; *16S*: 146,164; *CytB*: 223,247; *COI*: 583,043), covering all Metazoa. Of these, 258,225 (25.3%) derive from the four tetrapod classes (Amphibia: 55,254; Aves: 51,096; Mammalia: 101,106; Reptilia: 50,769). The distribution of these sequences between classes and loci, and the losses at each curation step are shown in Figure 4. In three of the four classes, there is a clear bias towards *CytB* sequences, with over 50% of sequences derived from this locus. In both Aves and Mammalia, the *16S* and *12S* loci are severely underrepresented at less than 10% each, while for Reptilia, *COI* is the least sequenced locus in the database.

The numbers of sequences and rates of loss due to our curation steps varied among taxonomic classes and the four loci, although losses were observed between steps in almost all instances. The most significant losses followed amplicon trimming and removal of non-unique sequences. Amplicon trimming led to especially high losses in Amphibia and *16S*, indicating that data published on GenBank for this class and marker do not generally overlap with our amplicons. Meanwhile, the high level of redundancy in public databases was highlighted by the significant reduction in the number of sequences during the final step of removing redundant sequences – in all cases over 10% of sequences was discarded, with some losses exceeding 50% (Mammalia: *COI*, *CytB*, *16S*; Amphibia: *16S*).

Data loss due to apparent mislabelling ranged between 1.9% and 7.4% and was thus generally higher than similar estimates for curated microbial databases [29]. SATIVA flags potential mislabels and suggests an alternative label supported by the phylogenetic placement of the sequences, allowing the user to make an appropriate decision on a case by case basis. The pipeline pauses after this step to allow such manual inspection to take place. However, for the current database, the number of sequences flagged was large (4378 in total), and the required taxonomic expertise was lacking, so all flagged sequences from non-target species were discarded to be conservative. The majority of mislabels were identified at species level (3053), but there were also significant numbers at genus (788), family (364) and order (102) level. Two to three sequences from Bornean mammal species were unflagged in each amplicon to retain the sequences in the database. This was important as in each case these were the only reference sequences available for the species. Additionally, *Muntiacus vaginalis* sequences that were automatically synonymised to *M. muntjak* based on the available information in the Catalogue of Life were revised back to their original identifications to reflect current taxonomic knowledge.

### Database composition

The final database was skewed even more strongly towards *CytB* than was the raw database. It was the most abundant locus for each class and represented over 60% of sequences for both Mammalia and Reptilia. In all classes, *16S* made up less than 10% of the final database, with Reptilia *COI* also at less than 10%.

Figure 5 shows that most species represented in the curated database for any locus have
just one unique haplotype against which HTS reads can be compared; only a few species
have many haplotypes. The prevalence of species with 20 or more haplotypes is particularly
notable in *CytB* where the four classes have between 25 (Aves) and 265 (Mammalia) species
in this category. The coloured circles in Figure 5 also show that the species of the taxonomy
are incompletely represented across all loci, and that coverage varies significantly between
taxonomic groups. In spite of global initiatives to generate *COI* sequences [53], this marker
does not offer the best species-level coverage in any class and is a poor choice for Amphibia
and Reptilia (<15% of species included). Even the best performing marker, *CytB*, is not a
universally appropriate choice, as Amphibia is better covered by *12S*. These differences in
underlying database composition will impact the likelihood of obtaining accurate taxonomic
assignment for any one species from any single marker. Further barcoding campaigns are
clearly needed to fill gaps in the reference databases for all markers and all classes to
increase the power of future e/iDNA studies. As the costs of HTS decrease, we expect that
such gap-filling will increasingly shift towards sequencing of whole mitochondrial genomes
of specimen obtained from museum collections, trapping campaigns etc. [34], reducing the
effect of marker choice on detection likelihood. In the meantime, however, the total
number of species covered by the database can be increased by combining multiple loci
(here, up to four) and thus the impacts of database gaps on correctly detecting species can
be minimized ([54]; Fig. 6).

In the present study, the primary target for iDNA sampling was the mammal fauna of
Malaysian Borneo, and the 103 species expected in the sampling area represent an
informative case study highlighting the deficiencies in existing databases (Fig. 7). Nine
species are completely unrepresented while only slightly over half (55 species) have at least
one sequence for all of the loci. Individually, each marker covers over half of the target
species, but none achieves more than 85% coverage (*12S*: 75 species; *16S*: 68; *CytB*: 88; *COI*:
66). Equally striking is the lack of within-species diversity, as most of the incorporated
species are represented by only a single haplotype per locus. Some of the species have large
distribution ranges, so it is likely that in some cases the populations on Borneo differ
genetically from the available reference sequences, possibly limiting assignment success.
Only a few expected species have been sequenced extensively, and most are of economic
importance to humans (e.g. *Bos taurus, Bubalus bubalis, Macaca* spp, *Paradoxurus*
*hermaphroditus, Rattus* spp., *Sus scrofa*), with as many as 100 haplotypes available (*Canis*
*lupus*). Other well-represented species (≥20 haplotypes) present in the sampling area
include several Muridae (*Chiropodomys gliroides, Leopoldamys sabanus, Maxomys surifer,*
*Maxomys whiteheadi*) and the leopard cat (*Prionailurus bengalensis*).

*Laboratory workflow*

Shotgun sequencing of a subset of our samples revealed that the median mammalian DNA
content was only 0.9%, ranging from 0% to 98%. These estimates are approximate, but with
more than 75% of the samples being below 5%, this shows clearly the scarcity of target DNA

in bulk iDNA samples. The generally low DNA content and the fact that the target DNA is
often degraded make enrichment of the target barcoding loci necessary. We used PCR with
high cycle numbers to obtain enough DNA for sequencing. However, this second step
increases the risk of PCR error: artificial sequence variation, non-target amplification, and/or
raising contaminations up to a detectable level.

We addressed these problems by running two *extraction replicates*, two *PCR replicates*, and
a multi-marker approach. The need for *PCR replicates* has been acknowledged and
addressed extensively in ancient DNA studies [16] and has also been highlighted for
metabarcoding studies [19; 20; 55; 56]. Despite this, many e/iDNA studies do not carry out
multiple *PCR replicates* to detect and omit potential false sequences. In addition, *extraction*
*replicates* are seldom applied, despite the evidence that cross-sample DNA contamination
can occur during DNA extraction [57; 58; 59]. We only accepted sequences that appeared in
a minimum of two independent PCRs for the lax and for the stringent criterion, where it has
to occur in each *extraction replicate A* and *B* (Fig. 1). The latter acceptance criterion is quite
conservative and produces higher false negative rates than e.g. accepting occurrence of at
least two positives. However, it also reduces the risk of accepting a false positives compared
to it (see Supplemental Fig. 1. for a simulation of false positive and false negatives rates
within a PCR), especially with increasing risk of false positive occurrence in a PCR for e.g.
example due to higher risk of contamination etc.. Metabarcoding studies are very prone to
false negatives, and downstream analyses like occupancy models for species distributions
can account for imperfect detection and false negatives. However, methods for discounting
false positive detections are not well developed [60]. Thus we think it is more important to
avoid false positives, especially if the results will be used to make management decisions
regarding rare or endangered species. In contrast, it might be acceptable to use a relaxed
acceptance criterion for more common species, as long as the rate false-positives/true-
positives is small and does not affect species distribution estimates. Employing both of our
tested criteria researchers could flag unreliable assignments and management decisions can
still use this information, but now in a forewarned way. An alternative to our acceptance
criteria could be use the PCR replicates itself to model the detection probability within a
sample using an occupancy framework [20; 60; 61].

We used three different loci to correct for potential PCR-amplification biases. We were,
however, unable to quantify this bias in this study due to the high degradation of the target
mammalian DNA, which resulted in much higher overall amplification rates for *16S*, the
shortest of our PCR amplicons. For *16S*, 85% of the samples amplified, whereas for *CytB* and
*12S*, only 57% and 44% amplified, respectively. Also the read losses due to trimming and
quality filtering were significantly lower for the *16S* sequencing runs (1.3% and 5.3% in
average, Supplemental Table 3) compared to the sequencing runs for the longer fragments
of *12S* and *CytB* (65.3% and 44.3% in average, Supplemental Table 3). Despite the greater
taxonomic resolution of the longer *12S* and *CytB* fragments, our poorer amplification and
sequencing results for these longer fragments emphasize that e/iDNA studies should

generally focus on short PCR fragments to increase the likelihood of positive amplifications
of the degraded target DNA. In the case of mammal-focussed e/iDNA studies, developing a
shorter (100 bp) *CytB* fragment would likely be very useful.

Another major precaution was the use of twin-tagging for both PCRs (Fig. 2). This ensures
that unlabelled PCR products are never produced and allows us to multiplex a large number
of samples on a single run of Illumina MiSeq run. Just 24 sample *tags 1* and 20 plate *tags 2*
allow the differentiation of up to 480 samples with matching tags on both ends. The same
number of individual primers would have needed longer tags to maintain enough distance
between them and would have resulted in an even longer adapter-tag overhang compared
to primer length. This would have most likely resulted in lower binding efficiencies due to
steric hindrances of the primers. Furthermore, this would have resulted in increased primer
costs. Thus our approach reduced sequencing and primer purchase costs while at the same
time largely eliminating sample mis-assignment via tag jumping, because tag-jump
sequences have non-matching forward and reverse *tag 1* sequences [43]. We estimated the
rate of tag jumps producing non-matching *tag 1* sequences to be 1 to 5%, and these were
removed from the dataset (Table 4). For our sequenced PCR plates, the rate of non-
matching *tag 2* tags was 2%. These numbers are smaller than data from Zepeda-Mendoza et
al. [56] who reported on sequence losses of 19% to 23% due to unused tag combinations
when they tested their DAMe pipeline to different datasets built using standard blunt-end
ligation technique. Although their numbers might not be one-to-one comparable to our
results as they counted unique sequences, and we report on read numbers, our PCR
libraries with matching barcodes seem reduce the risk of tag jumping compared to blunt-
end ligation techniques. For the second PCR round, we used the same tag pair *tag 2* for all
24 samples of a PCR plate. In order to reduce cost we tested pooling these 24 samples prior
to the second PCR round, but we detected a very high tag jumping rate of over 40% (Table
4), which ultimately would increase cost through reduced sequencing efficiency. Twin-
tagging increases costs because of the need to purchase a larger number of primer pairs but
at the same time it increases confidence in the results.

Tagging primers in the first PCR reduces the risk of cross-contamination via aerosolised PCR
products. However, we would not be able to detect a contamination prior the second PCR
from one plate to another, as we used the same 24 tags (*tag 1*) for all plates. Nevertheless
such a contamination is very unlikely to result in any accepted false positive as it would be
improbable to contaminate both the A and B replicates, given the exchange of all reagents
and the time gap between the PCRs. Previous studies have shown that unlabelled volatile
PCR products pose a great risk of false detections [62], a risk that is greatly increased if a
high number of samples are analysed in the laboratories [13]. Also, in laboratories where
other research projects are conducted, this approach allows the detection of cross-
experiment contamination. Therefore, we see a clear advantage of our approach over
ligation techniques when it comes to producing sequencing libraries, as the Illumina tags are

 only added after the first PCR, and thus the risk of cross contamination with unlabelled PCR
612 amplicons is very low.

### Assignment results

A robust assignment of species is an important factor in metabarcoding as an incorrect
identification might result incorrect management interventions. The reliability of taxonomic
assignments is expected to vary with respect to both marker information content and
database completeness, and this is reflected in the probability estimates provided by
*PROTAX*. In a recent study, less than 10% of the mammal assignments made at species level
against a worldwide reference database were considered reliable with the short *16S*
amplicon, but this increased to 46% with full-length *16S* sequences [31]. In contrast, in the
same study over 80% of insect assignments at species level were considered reliable with a
more complete, geographically restricted database of full-length COI barcodes. A similar
pattern was observed in our data during manual curation of the assignment results – there
was more ambiguity in the results for the short *16S* amplicon than for other markers.
However, due to the limited amount of often degraded target DNA in e/iDNA samples, short
amplicons amplify much better. In our case, this had the drawback that some species lacked
any interspecific variation, and thus sequencing reads shared 99%-100% identity for several
species. For example, our only *16S* reference of *Sus barbatus* was 100% identical to *S.*
*scrofa*. But as latter species does not occur in the studied area we could assign all reads
manually to *S. barbatus*. In several cases we were able to confirm *S. barbatus* by additional
*CytB* results, highlighting the usefulness of multiple markers.

Another advantage of multiple markers is the opportunity to fill gaps in the reference
database. For example, we lacked *16S* reference sequences for *Hystrix brachyura*, and reads
were assigned by *PROTAX* only to the unknown species *Hystrix* sp.. In one sample, however,
almost 5000 *CytB* reads could be confidently assigned to *Hystrix brachyura*, and thus we
used the *Hystrix* sp. 16S sequences in the same sample to build a consensus *16S* reference
sequence for *Hystrix brachyura* for future analyses. In another example we had *CytB* reads
assigned to *Mydaus javanicus*, the Sunda stink-badger in one sample but *12S* reads assigned
to *Mydaus* sp. in another one. As we lacked a *12S Mydaus* reference and as there is only one
*Mydaus* species on Borneo we could assume that this second sample is most likely also
*Mydaus javanicus*.

We also inferred that PCR and sequencing errors resulted in reads being assigned to sister
taxa. We observed that a high number of reads of a true sequence were assigned to a
species and a lower number of noise sequences were assigned to a sister taxon. Such a
pattern was observed for ungulates, especially deer that showed little variance in *16S*. It is
hard to identify and control for such pattern automatically, and it highlights the importance
of visual inspection of the results.

For the more lax criterion (two positive *PCR replicates*) we accepted 196 species
assignments out of 109 leech samples. Under the stringent criterion (i.e. having positive
detections in both *extraction replicates A* and *B*) we accepted about 14% assignments less;

in total 170 vertebrate detections within 97 bulk samples (Table 5). For 65% of the species
frequencies did not change and almost half of the not accepted assignments were from the
most frequent species *Rusa unicolor* and *Sus barbatus*. However, with the more stringent
criterion we did not accept two species (1x *Macaca fascicularis* & 2x *Mydaus javanensis*). In
five cases the stringent criterion would not accept assignments that could be made only to
unknown species (1x *Hystrix* sp., 3x *Macaca* sp., 1x *Tragulus* sp.) (Table 5). For the all these
gernera we have two occurring species in the area. As the true occurrence of species within
our leeches was unknown we cannot evaluate how many of the additional 27 detections in
the lax criterion are false positives and how many might be false negatives for the stricter
criterion. However, by accepting only positive *AB* assignment results, we increase the
confidence of species detection, even if the total number of reads for that species was low.
When it comes to rare or threated species this outweighs the risk of reporting false positives
to our opinion. 48% of the assignments with the stringent criterion were present in all four
*A1*, *A2*, *B1* and *B2*. 35% were present in at least three replicates (e.g. A1, A2, B1).

The mean number of reads per sample used for the taxomomic assignment varied from
162,487 *16S* reads for SeqRun01 to 7,638 *CytB* reads for SeqRun05 (Supplemental Table 4).
In almost all cases, however, the number of reads of an accepted assignment was high
(median= 52,386; mean= 300,996; SD= 326,883). PCR stochasticity, primer biases, multiple
species in individual samples, and pooling of samples exert too many uncertainties that
could bias the sequencing results [63; 64]. Thus we do not believe that raw read numbers
are the most reliable indicators of tetrapod DNA quantity in iDNA samples. Replication of
detection is inherently more reliable. In contrast to our expectation that higher cycle
number might be necessary to amplify even the lowest amounts of target DNA, our data do
not support this hypothesis. Although we observed an increase in positive PCRs for A2/B2
(the 40-cycle PCR replicates), the total number of accepted assignments in *A1/B1* and *A2/B2*
samples did not differ. This indicates first that high PCR cycle numbers mainly increased the
risk of false positives and second that our multiple precautions successfully minimized the
acceptance of false detections.

## Conclusion

Metabarcoding of e/iDNA samples will certainly become a very valuable tool in assessing
biodiversity, as it allows to detect species non-invasively without the need to capture and
handle the animals [65] and because sampling effort can often be greatly reduced.
However, the technical and analytical challenges linked to sample types (low quantity and
quality DNA) and poor reference databases have so far been insufficiently recognized. In
contrast to ancient DNA studies where standardized laboratory procedures and specialized
bioinformatics pipelines have been established and are followed in most cases, there is
limited methodological consensus in e/iDNA studies, which reduces rigour. In this study, we
present a robust metabarcoding workflow for e/iDNA studies. We hope that the provided
scripts and protocols facilitate further technical and analytical developments. The use of
e/iDNA metabarcoding to study the rarest and most endangered species such as the Saola is

exciting, but geneticists bear the heavy responsibility of providing correct answers to conservationists.

## Acknowledgements

All authors thank the German Federal Ministry of Education and Research (BMBF FKZ: 01LN1301A) and the Leibniz-IZW for funding this study. We also thank the Sabah Forestry Department, especially Johnny Kissing, Peter Lagan and Datuk Sam Mannan for supporting the fieldwork and the Sabah Biodiversity Council for providing research, collection and export permits for this work. We are grateful to John Mathai, Seth Timothy Wong for conducting the field work and collecting the leeches. We are also grateful to Jörns Fickel, head of the Department Evolutionary Genetics of the Leibniz-IZW for continuous support and collaboration. Furthermore we would like to thank Sebastian Wieser for lab-support, Dorina Lenz and Anke Schmidt for their help and fruitful discussions. D.W. Yu and C.C.Y. Xu were supported by the National Natural Science Foundation of China (41661144002, 31670536, 31400470, 31500305), the Key Research Program of Frontier Sciences, CAS (QYZDY-SSW-SMC024), the Bureau of International Cooperation project (GJHZ1754), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDA20050202, XDB31000000), the Ministry of Science and Technology of China (2012FY110800), and the State Key Laboratory of Genetic Resources and Evolution at the Kunming Institute of Zoology.

## References

[1]   Gariepy TD, Lindsay R, Odgen N, Greory TR. Identifying the last supper: utility of the DNA barcode library for bloodmeal identification in ticks. Mol Ecol Res. 2012; 12: 646-52.

[2]   Lee P-S, Gan HM, Clements GR, Wilson J-J. Field calibration of blowfly-derived DNA against traditional methods for assessing mammal diversity in tropical forests. Genome 2016;59: 1008-22.

[3]   Calvignac-Spencer S, Merkel K, Kutzner N, et al.. Carrion fly-derived DNA as a tool for comprehensive and cost-effective assessment of mammalian biodiversity. Mol Ecol 2013; 22: 915-24.

[4]   Rodgers TW, Xu CCY, Giacalcone J, et al.. Carrion fly-derived DNA metabarcoding is an effective tool for mammal surveys: Evidence from a known tropical mammal community. Mol Ecol Res. 2017; 17(6):1-13

[5]   Hoffmann C, Merkel K, Sachse A, et al.. Blow flies as urban wildlife sensors. Mol Ecol Res. 2018; 18(3):502-10

[6]   Schönberger AC, Wagner S, Tuten HC, et al.. Host preferences in host-seeking and 632 blood-fed mosquitoes in Switzerland. Med Vet Entomol. 2015; 30(1): 39-52.

[7]   Townzen JS, Brower AVZ, Judd DD. Identification of mosquito bloodmeals using mitochondrial cytochrome oxidase subunit I and cytochrome b gene sequences. Med Vet Entomol. 2008; 22. 386-93.

[8]    Kocher A, Thoisy B, Catzeflies F, et al.. iDNA screening: Disease vectors as vertebrate
samplers. Mol Ecol. 2017; 26(22): 6478-86.

[9]    Taylor L, Cummings RF, Velten R, et al.. Host (Avian) Biting Preference of Southern
California Culex Mosquitoes (Diptera: Culicidae). J Med Entomol. 2012; 49(3): 687-
96.

[10]   Schnell IB, Thomsen PF, Wilkinson N, et al.. Screening mammal biodiversity using
DNA from leeches. Curr Biol. 2012, 22(8): R262—3.

[11]   Tessler M, Weiskopf SR, Berniker L, et al.. Bloodlines: mammals, leeches, and
conservation in southern Asia. Syst Biodivers. 2018; 1-9.

[12]   Weiskopf SR, McCarthy KP, Tessler M, et al.. Using terrestrial haematophagous
leeches to enhance tropical biodiversity monitoring programmes in Bangladesh. J
Appl Ecol. 2018: 1-11.

[13]   Schnell IB, Bohmann K, Schultze SE, et al.. Debugging diversity - a pan-continental
exploration of the potential of terrestrial blood-feeding leeches as a vertebrate
monitoring tool. Mol Ecol Res. 2018.

[14]   Calvignac-Spencer S, Leendertz FH, Gilbert MT, Schubert G. An invertebrate
stomach's view on vertebrate ecology: certain invertebrates could be used as
"vertebrate samplers" and deliver DNA-based information on many aspects of
vertebrate ecology. BioEssays. 2013; 35(11): 1004-13.

[15]   Schnell IB, Sollmann R, Calvignac-Spencer S, et al.. iDNA from terrestrial
haematophagous leeches as a wildlife surveying and monitoring tool – prospects,
pitfalls and avenues to be developed. Front Zool. 2015; 12:24.

[16]   Pääbo S, Poinar H, Serre D, et al.. Genetic analyses from ancient DNA. Annu Rev
Genet. 2004; 38: 645-79.

[17]   Hofreiter M, Paijmans JL, Goodchild H, et al. The future of ancient DNA: Technical
advances and conceptual shifts. BioEssays. 2015; 37(3): 284-93.

[18]   Cristescu ME, Hebert , PDN. Uses and Misuses of Environmental DNA in Biodiversity
Science and Conservation. Cristescu, Melania E und Hebert, Paul D N. 1, 2018, Annu
Rev Ecol Evol Syst 2018; 49.

[19]   Ficetola GF, Pansu J, Bonin A, et al.. Replication levels, false presences and the
estimation of the presence/absence from eDNA metabarcoding data. Mol Ecol Res.
2014; 15(3): 543-56.

[20]   Ficetola GF, Taberlet P., Coissac E. How to limit false positives in environmental DNA
and metabarcoding? Mol Ecol Res. 2016; 16(3): 604-7.

[21]   Dalton R. Still looking for that woodpecker. Nature 2010; 463: 718-9.

[22]   Bazinet AL, Cummings MP. A comparative evaluation of sequence classification
programs. BMC bioinformatics. 2012; 13(1): 92.

[23]   Richardson RT, Bengtsson-Palme J, Johnson RM. Evaluating and optimizing the
performance of software commonly used for the taxonomic classification of DNA
metabarcoding sequence data. Mol Ecol Res. 2017; 17(4): 760-9.

[24] Somervuo P, Koskela S, Pennanen J, Henrik Nilsson R, Ovaskainen O. Unbiased
probabilistic taxonomic classification for DNA barcoding. Bioinformatics. 2016;
32(19): 2920-7.

[25] Quast C, Pruesse E, Gerken J, et al.. SILVA Databases. In: Nelson KE (eds)
Encyclopedia of Metagenomics. Springer, Boston 2015; 626-35.

[26] Nilsson RH, Ryberg M. Taxonomic Reliability of DNA Sequences in Public Sequence
Databases: A Fungal Persepctive. PLoS ONE. 1, 2006; 1.

[27] Forster P. To Err is Human. Ann Hum Gen 2003; 67: 2-4.

[28] Harris JD. Can you bank on GenBank? Trends Ecol Evol. 2003; 18: 317-9.
doi:10.1016/S0169-5347(03)00150-2.

[29] Kozlov AM, Zhang J, Yilmaz P, Glöckner FO, Stamatakis A. Phylogeny-aware
identification and correction of taxonomically mislabeled sequences. Nucleic Acids
Res. 2016; 44(11): 5022-33.

[30] Huson DH, Auch AF, Qi J, Schuster SC. MEGAN analysis of metagenomic data.
Genome Res. 2007; 17(3): 377-86.

[31] Somervuo P, Yu DW, Xu CC, Ji Y, et al.. Quantifying uncertainty of taxonomic
placement in DNA barcoding and metabarcoding. Methods Ecol Evol. 2017; 8(4):
398-407.

[32] Machida RJ, Leray M, Ho SL, Knowlton N. Metazoan mitochondrial gene sequence
reference datasets for taxonomic assignment of environmental samples. Sci Data.
2017; 4: 170027.

[33] Chamberlain SA, Szöcs E. taxize: taxonomic search and retrieval in R. Version 2.
F1000Res. 2013; 2: 191.

[34] Salleh FM, Ramos-Madrigal J, Peñaloza F, et al.. An expanded mammal mitogenome
dataset from Southeast Asia. GigaScience. 2017; 6(8): 1-8

[35] Kans, Jonathan. Entrez Direct: E-utilities on the UNIX Command Line. In: Entrez
Programming Utilities Help [Internet]. Bethesda (MD): National Center for
Biotechnology Information (US). 2010.

[36] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. Journal of molecular biology. 1990; 215(3):, 403-10.

[37] Camacho C, Coulouris G, Avagyan V, et al.. BLAST+: architecture and applications.
BMC bioinformatics. 2009; 10(1): 421.

[38] Katoh K, Standley DM. (2013). MAFFT multiple sequence alignment software version
7: improvements in performance and usability. Mol Biol Evol. 2013; 30(4): 772-80.

[39] Katoh K, Misawa K, Kuma KI, Miyata T. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;
30(14): 3059-66.

[40] Chesters D. (2013) collapsetypes.pl [computer software available at
http://sourceforge.net/projects/collapsetypes/]

[41]  Kocher TD, Thomas WK, Meyer A, et al.. Dynamics of mitochondrial DNA evolution in
animals: amplification and sequencing with conserved primers. Proc. Natl. Acad. Sci.
U.S.A.. 1989; 86(16): 6196-6200.

[42]  Taylor PG. Reproducibility of ancient DNA sequences from extinct Pleistocene fauna.
Mol Biol Evol. 2996; 13(1): 283-5.

[43]  Schnell IB, Bohmann K, Gilbert MTP. Tag jumps illuminated—reducing
sequence-to-sample misidentifications in metabarcoding studies. Mol Ecol Res.
2015; 15(6): 1289-1303.

[44]  Faircloth BC, Glenn TC. Not all sequence tags are created equal: designing and
validating sequence identification tags robust to indels. PloS One. 2012; 7(8): e42543

[45]  Murray DC, Coghlan ML, Bunce M. From benchtop to desktop: important
considerations when designing amplicon sequencing workflows. PLoS One. 2015;
10(4): e0124671.

[46]  Schubert M, Lindgreen S, Orlando L. AdapterRemoval v2: rapid adapter trimming,
identification, and read merging. BMC Research Notes 2016; 9.

[47]  Edgar RC. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics. 2010; 26(19): 2460-2461.

[48]  Edgar RC, Flyvbjerg H. Error filtering, pair assembly and error correction for next-
generation sequencing reads. Bioinformatics. 2015; 31(21): 3476-82.

[49]  Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet. Jjournal. 2011; 17(1): 10-12.

[50]  Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame genomic
sequence comparison. Genome Res. 2011; 21(3): 487-493.

[51]  Fortes GG, Paijmans JLA. Analysis of Whole Mitogenomes from Ancient Samples. In:
Kroneis T (eds). Whole Genome Amplification: Methods and Protocols. Springer New
York 2015; 179-195.

[52]  Ondov BD, Bergman NH, Phillippy AM. Interactive metagenomic visualization in a
Web browser. BMC Bioinformatics 2011; 12: 385.

[53]  Ratnasingham S, Hebert PDN. BOLD: The Barcode of Life Data System
(www.barcodinglife.org). Mol Ecol Notes. 2007; 3: 355-64.

[54]  Evans NT, Li Y, Renshaw MA, et al. Fish community assessment with eDNA
metabarcoding: effects of sampling design and bioinformatic filtering. Can J Fish
Aquat Sci. 2017; 74(9):, 1362-74.

[55]  Bonin A, Taberlet P, Zinger L, Coissac E. Environmental DNA: For Biodiversity
Research and Monitoring. 1st ed. Oxford University Press; 2018.

[56]  Zepeda-Mendoza ML, Bohmann K, Baez A, Gilbert M., DAMe: a toolkit for the initial
processing of datasets with PCR replicates of double-tagged amplicons for DNA
metabarcoding analyses. BMC Res Notes 2016; 9.

[57]  Racimo F, Renaud G, Slatkin M. Joint estimation of contamination, error and
demography for nuclear DNA from ancient humans. PLoS Genet. 2016; 12(4):
e1005972.

[58]   Orlando L, Gilbert MTP, Willerslev E. Reconstructing ancient genomes and
epigenomes. Nat Rev Genet 2015; 16(7): 395

[59]   Laurin-Lemay S, Brinkmann H, Philippe H. Origin of land plants revisited in the light
of sequence contamination and missing data. Current Biology. 2012; 22(15): R593-4.

[60]   Lahoz-Monfort JJ, Guillera-Arroita G, Tingley R. Statistical approaches to account for
false-positive errors in environmental DNA samples. Mol Ecol Res. 2015; 16: 673-85.

[61]   Dorazio RM, Erickson RA. ednaoccupancy: An r package for multiscale occupancy
modelling of environmental DNA data. Mol Ecol Res. 2018; 18: 368-80.

[62]   Kwok S, Higuchi R. Avoiding false positives with PCR. Nature. 1989; 339: 237-8.

[63]   Bush A, Sollmann R, Wilting A, et al.. Connecting Earth observation to high-
throughput biodiversity data. Nat Ecol Evol 2017; 1(7): 0176

[64]   Piñol J, Mir G, Gomez-Polo P, Agustí N. Universal and blocking primer mismatches
limit the use of high-throughput DNA sequencing for the quantitative metabarcoding
of arthropods., Mol Ecol Res.2014; 15: 819-30.

[65]   Nichols RV, Vollmers C, Newsom L, et al.. Minimizing polymerase biases in
metabarcoding. Mol Ecol Res. 2018; 18: 927-39.

**Table 1:** Sequence motifs that compose the 25 different target primers for the first and the
second PCR. First PCR primers consist of target specific primer followed by an overhang out
of sample specific *tag 1* and *read 1* and *read 2* sequencing primer, respectively. The second
PCR primers consist of the *read 1* or the *read 2* sequencing primer followed by an plate
specific *tag 2* and the P5 and P7 adapters, respectively (see also Fig. 2).

| Name | Sequence | Reference |
|---|---|---|
| tag A | TGCAT | Faircloth & and Glenn 2012 |
| tag B | TCAGC | Faircloth & and Glenn 2012 |
| tag C | AAGCG | Faircloth & and Glenn 2012 |
| tag D | ACAAG | Faircloth & and Glenn 2012 |
| tag E | AGTGG | Faircloth & and Glenn 2012 |
| tag F | TTGAC | Faircloth & and Glenn 2012 |
| tag G | CCTAT | Faircloth & and Glenn 2012 |
| tag H | GGATG | Faircloth & and Glenn 2012 |
| tag I | CTAGG | Faircloth & and Glenn 2012 |
| tag K | CACCT | Faircloth & and Glenn 2012 |
| tag L | GTCAA | Faircloth & and Glenn 2012 |
| tag M | GAAGT | Faircloth & and Glenn 2012 |
| tag N | CGGTT | Faircloth & and Glenn 2012 |
| tag O | ACCGA | Faircloth & and Glenn 2012 |
| tag P | ACGTC | Faircloth & and Glenn 2012 |
| tag Q | AGACT | Faircloth & and Glenn 2012 |
| tag R | AGGAA | Faircloth & and Glenn 2012 |
| tag S | ATTCC | Faircloth & and Glenn 2012 |
| tag T | CAATC | Faircloth & and Glenn 2012 |
| tag V | CATGA | Faircloth & and Glenn 2012 |
| tag W | CCACA | Faircloth & and Glenn 2012 |
| tag X | GCTTA | Faircloth & and Glenn 2012 |
| tag Y | GGTAC | Faircloth & and Glenn 2012 |
| tag Z | AACAC | Faircloth & and Glenn 2012 |
| Tag Control | ATCTG | Faircloth & and Glenn 2012 |
| *CytB*-fw | AAAAAGCTTCCATCCAACATCTCAGCATGATGAAA | Kocher et al. 1989 |
| *CytB*-rv | AAACTGCAGCCCCTCAGAATGATATTTGTCCTCA | Kocher et al. 1989 |
| *16S*-fw | CGGTTGGGGTGACCTCGGA | Taylor 1996 |
| *16S*-rv | GCTGTTATCCCTAGGGTAACT | Taylor 1996 |
| *12S*-fw | AAAAAGCTTCAAACTGGGATTAGATACCCCACTAT | Kocher et al. 1989 |
| *12S*-rv | TGACTGCAGAGGGTGACGGGCGGTGTGT | Kocher et al. 1989 |
| Read 1 sequence primer | ACACTCTTTCCCTACACGACGCTCTTCCGATCT | Illumina Document # 1000000002694 v03 |
| Read 2 sequence primer | GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT | Illumina Document # 1000000002694 v03 |
| P5 adapter | AATGATACGGCGACCACCGAGATCTACAC | Illumina Document # 1000000002694 v03 |
| P7 adapter | CAAGCAGAAGACGGCATACGAGAT | Illumina Document # 1000000002694 v03 |

873  **Table 2:** Main steps undertaken by each module of the database curation script.

| MODULE | STEPS |
| --- | --- |
| Module 1 | Extract subset of raw Midori database for query taxon and loci. |
| | Remove sequences with non-binomial species names, reduce subspecies to species labels |
| | Add local sequences (optional) |
| | Check for relevant new sequences for list of query species on NCBI (GenBank and RefSeq) (optional) |
| | Select amplicon region and remove primers |
| | Remove sequences with ambiguous bases |
| | Align |
| | End of module: Optional check of alignments |
| Module 2 | Compare sequence species labels with taxonomy |
| | Non-matching labels queried against Catalogue of Life to check for known synonyms |
| | Remaining mismatches kept if genus already exists in taxonomy, otherwise flagged for removal |
| | End of module: Optional check of flagged species labels |
| Module 3 | Discard flagged sequences |
| | Update taxonomy key file for sequences found to be incorrectly labelled in Module 2 |
| | Run SATIVA |
| | End of module: Optional check of putatively mislabelled sequences |
| Module 4 | Discard flagged sequences |
| | Finalise consensus taxonomy and relabel sequences with correct species label and accession number |
| | Select one representative sequence per haplotype per species |

874 **Table 3:** GNU core utilities and other lightweight tools used for manipulation of text and
875 sequence files

| TOOL | FUNCTION | SOURCE |
| --- | --- | --- |
| awk, cut, grep, join, sed, sort, tr | Processing text files | GNU core utilities |
| seqbuddy | Processing FASTA/Q files | https://github.com/biologyguy/BuddySuite |
| seqkit | Processing FASTA/Q files | https://github.com/shenwei356/seqkit |
| seqtk | Processing FASTA/Q files | https://github.com/lh3/seqtk |
| tabtk | Processing tab-delimited text files | https://github.com/lh3/tabtk |

877 **Table 4:** Number of reads per sequencing run and the numbers of reads with matching, non-matching or unidentifiable tags for seven of the
878 eight sequencing runs*.

| | total reads | matching tag 2 reads | non-matching tag 2 reads | %[1] | matching tag 1 reads | non-matching tag 1 reads | %[2] | erroneous tag 1 reads | %[2] |
|---|---|---|---|---|---|---|---|---|---|
| **SeqRun01** | 18,438,517 | 18,102,702 | 282,419 | 1.5 | 17,514,515 | 451,028 | 2.5 | 137,159 | 0.8 |
| **SeqRun02** | 25,385,558 | 24,596,380 | 626,245 | 2.5 | 23,426,084 | 612,045 | 2.5 | 558,251 | 2.3 |
| **SeqRun03** | 14,875,796 | 14,393,884 | 343,528 | 2.3 | 13,766,187 | 426,181 | 3.0 | 201,516 | 1.4 |
| **SeqRun04** | 2,027,794 | 1,935,149 | 56,077 | 2.8 | 1,806,655 | 88,307 | 4.6 | 40,187 | 2.1 |
| **SeqRun05** | 18,221,504 | 17,500,366 | 421,588 | 2.3 | 16,793,851 | 482,365 | 2.8 | 161,458 | 0.9 |
| **SeqRun06** | 20,718,202 | 19,874,913 | 429,048 | 2.1 | 19,317,305 | 371,048 | 1.9 | 81,422 | 0.4 |
| **SeqRun07** | 24,604,610 | 23,746,938 | 663,730 | 2.7 | 22,446,187 | 497,366 | 2.1 | 803,385 | 3.4 |
| **Total** | 124,271,981 | 120,150,332 | 2,822,635 | 2.3 | 115,070,784 | 2,928,340 | 2,5 | 1,983,378 | 1,7 |
| **IndexRun** | 10,276,093 | 10,116,808 | NA | NA | 5,841,190 | 4,186,688 | 41.4 | 88,930 | 0.9 |

[1] refers to total reads
[2] refers to matching tag 2

879 *Sequencig run SeqRun08 run contained libraries of another project, thus we were unable to provide a number of raw reads.

880 **Table 5:** Number of accepted species assignments with two different acceptance criteria the
881 more stringent criterion accepting only assignments occurring in both *extraction replicates*
882 (A & B), and the more lax criterion accepting assignment two or more positives in any of the
883 twelve *PCR replicates*.

| | stringent | lax | change |
|---|---|---|---|
| *Aonyx cinereus* | 1 | 1 | 0 |
| *Arctictis binturong* | 1 | 1 | 0 |
| *Bos Javanicus* | 9 | 11 | +2 |
| *Echinosorex gymnura* | 5 | 6 | +1 |
| *Felis catus* | 2 | 2 | 0 |
| *Helarctos malayanus* | 5 | 6 | +1 |
| *Hemigalus derbyanus* | 3 | 3 | 0 |
| *Hystrix brachyura* | 8 | 8 | 0 |
| *Hystrix crassipinis* | 1 | 1 | 0 |
| *Hystrix sp.* | 1 | 2 | +1 |
| *Kalophrynus pleurostigma* | 1 | 1 | 0 |
| *Macaca fascicularis* | | 1 | +1 |
| *Macaca nemestrina* | 1 | 2 | +1 |
| *Macaca sp.* | | 3 | +3 |
| *Manis javanicus* | 2 | 2 | 0 |
| *Muntiacus atherodes* | 6 | 6 | 0 |
| *Muntiacus muntjak* | 2 | 2 | 0 |
| *Muntiacus sp.* | 10 | 10 | 0 |
| *Mydaus javanensis* | | 1 | +1 |
| *Mydaus sp.* | | 1 | +1 |
| *Pongo pygmaeus* | 5 | 5 | 0 |
| *Rusa unicolor* | 61 | 69 | +8 |
| *Sus barbatus* | 17 | 22 | +5 |
| *Tragulus javanicus* | 3 | 3 | 0 |
| *Tragulus napu* | 10 | 11 | +1 |
| *Tragulus sp.* | | 1 | +1 |
| *Trichys fasciculata* | 4 | 4 | 0 |
| *Viverra tangalunga* | 12 | 12 | 0 |
| **total accepted assignments** | 170 | 197 | +27 |

**Figure 1:** laboratory scheme; during DNA extraction the sample is split into two extraction replicates A & B. Our Protocol consists of two rounds of PCR that were the sample tags, the necessary sequencing primer and sequencing adapters are added to the the amplicons. For each extraction replicate we ran a low cycle PCR and a high cycle PCR for each marker that we have twelve independent PCR replicates per sample. All PCR products were sequenced and the obtained reads were taxonomically identified with PROTAX.

**Figure 2:** Scheme to build double 'twin-tagged' PCR libraries. The first round of PCR uses target-specific primers (*12S*, *16S*, or *CytB*, dark grey) that have both been extended with the same (i.e. 'twin') sample-identifying *tag* sequences *tag 1* (yellow) and then with the different *read 1* (dark blue) and *read 2* (light blue) sequence primers. The second round of PCR uses the priming sites of the *read 1* and *read 2* sequencing primers to add twin plate-identifying *tag* sequences *tag 2* (orange) and the P5 (dark red) and P7 (light red) Illumina adapters.

**Figure 3:** For the stringent acceptance criterion we only accepted taxonomic assignments that were positively detected in both *extraction replicates* A and B (green colour). The numbers (1 & 2) refer to the two PCR replicates for each extraction replicate.

**Figure 4:** Data availability and percentage loss at each major step in the database curation procedure for each target amplicon and class of Tetrapoda. The number of sequences decreases between steps except "Extra sequences added" where additional target sequences are included for Mammalia and there is no change for the other three classes.

**Figure 5:** Haplotype number by species (frequency distribution) and the total number of species with at least one haplotype, shown relative to the total number of species in the taxonomy for that category (bubbles), shown for each marker and class of Tetrapoda. The proportion of species covered by the database varies between categories but in all cases a majority of recovered species are represented by a single unique haplotype.

**Figure 6:** The percentage of the full taxonomy covered by the final database at each taxonomic level for each class of Tetrapoda. Includes the percentage of taxa represented by each marker and all markers combined. In all cases taking all four markers together increases the proportion of species, genera and families covered by the database, but it remains incomplete when compared with the full taxonomy.

**Figure 7:** The number of unique haplotypes per marker for each of the 103 mammal species expected in the study area. Bubble size is proportional to the number of haplotypes and varies between 0 and 100. Only 55 species have at least one sequence per marker and nine species are completely unrepresented in the current database.

**Supplemental Figure 1:** The rates of accepted false negatives (upper graph) and false positives (lower graph) for both our used acceptance criteria for varying PCR detection probabilities. The red line always denotes the stringent acceptance criterion that a positive is only accepted if it is present in at least one A and one B replicate. The lax criterion (blue)

931 accepted at any two positives out of the twelve replicates. The stringent criterion poses a
932 higher risk of accepting a false negative but it reduces clearly the risk of false positives,
933 especially with increasing detection probability due to higher risk of contamination.

## 1.-round PCR

<<amplification | target | tag 1 | read 2

DNA

DNA

read 1 | tag 1 | target | amplification>>

## 1.-round product:

## 2.-round PCR

<<amplification | read 2 | tag 2 | P7

P5 | tag 2 | read 1 | amplification>>

## 2.-round product: