Selective inhibition of CA IX over CA XII in breast cancer cells using benzene sulfonamides: Disconnect between CA activity and growth inhibition

Mam Y. Mboge[1], Zhijuan Chen[1], Alyssa Wolff[1], John V. Mathias[1], Chingkuang Tu[1], Kevin D. Brown[1], Murat Bozdag[2], Fabrizio Carta[2], Claudiu T. Supuran[2], Robert McKenna[1], and Susan C. Frost[1]*

From the [1]Department of Biochemistry and Molecular Biology, University of Florida, 1200 Newell Drive, Gainesville, FL 32610, USA; [2]University of Florence, NEUROFARBA Department, Sezione di Farmaceutica e Nutraceutica, Via Ugo Schiff 6, 50019 Sesto Fiorentino (Florence), Italy.

Running title: Selective inhibition of CA IX over CA XII activity

*Corresponding author

*sfrost@ufl.edu (SCF)

Abstract

Carbonic anhydrases (CAs) have been linked to tumor progression, particularly membrane-bound CA isoform IX (CA IX). The role of CA IX in the context of breast cancer is to regulate the pH of the tumor microenvironment. In contrast to CA IX, expression of CA XII, specifically in breast cancer, is associated with better outcome despite performing the same catalytic function. In this study, we have structurally modeled the orientation of bound ureido-substituted benzene sulfonamides (USBs) within the active site of CA XII, in comparison to CA IX and cytosolic off-target CA II, to understand isoform specific inhibition. This has identified specific residues within the CA active site, which differ between isoforms that are important for inhibitor binding and isoform specificity. The ability of these sulfonamides to block CA IX activity in breast cancer cells is less effective than their ability to block activity of the recombinant protein (by one to two orders of magnitude depending on the inhibitor). The same is true for CA XII activity but now they are two to three orders of magnitude less effective. Thus, there is significantly greater specificity for CA IX activity over CA XII. While the inhibitors block cell growth, without inducing cell death, this again occurs at two orders of magnitude above the K_i values for inhibition of CA IX and CA XII activity in their respective cell types. Surprisingly, the USBs inhibited cell growth even in cells where CA IX and CA XII expression was ablated. Despite the potential for these sulfonamides as chemotherapeutic agents, these data suggest that we reconsider the role of CA activity on growth potentiation.

Introduction

Breast cancer remains the second most diagnosed and one of the leading causes of cancer-related deaths among women in the United States. Gene expression patterns of dissected human breast tumors have provided distinct "molecular portraits" allowing classification into subtypes based solely on differences in these patterns [1]. These subtypes fall into three major groups: those that express estrogen (ER)/progesterone receptors (PR) (the luminal subtype), those that express HER2 (ERBB2+), and those that express neither (the basal-like, including the "triple negative" phenotype, TNBC). About 74% of breast cancer patients have ER-positive tumors [2]. For these patients, adjuvant therapy with tamoxifen (a selective estrogen receptor modulator) has contributed to the 30% decrease in breast cancer mortality in the past two decades, making it the most successful targeted cancer therapy to date [3]. Yet, about half of the tamoxifen-treated women will develop endocrine-resistant disease [4]. HER2 positive breast cancers are observed in about 15% of patients (some of which are also ER/PR positive), who in recent years has been successfully treated using a monoclonal antibody against the HER2 receptor. Triple negative breast cancer (TNBC) also accounts for about 15% of breast cancer patients, but is responsible for a disproportionate number of breast cancer deaths [5]. The higher mortality rate in women with TNBC is attributed to lack of targeted therapies coupled with the inability to treat metastatic disease [5-7]. That tumors develop resistance to radiation and/or chemotherapies complicates treatment.

Changes in the tumor microenvironment (TM), induced partly by hypoxia, underlie aggressive and resistant cancer phenotypes [8]. It is known that the hypoxic

TM contributes to tumorigenic characteristics such as tumor cell proliferation,

angiogenesis, cell motility, and invasiveness that translate to an overall poor prognosis

in cancer patients [8, 9]. Hypoxia induces a global change in gene expression through

stabilization of hypoxia inducible factor one alpha (HIF1α) [10]. Stabilization of this

transcription factor leads to the upregulation of genes important for cell survival [10].

Hypoxia-induced upregulation of genes in glycolysis cause elevated glucose uptake,

which is a hallmark for actively dividing cells and correlates closely with transformation

[11-13]. Increased glycolysis results in high rates of extracellular acidification (pH_e)

through the export of H^+ in an attempt to maintain a more basic/neutral intracellular pH

(pH_i) [14-16]. This increase in H^+ production and export to the extracellular space

induces apoptosis in normal cells surrounding tumors [17]. In contrast, by mechanisms

not understood, cancer cells establish as new "set-point" which allows them to tolerate

low pH_e [8, 17-19]. The stress of the TM also induces resistance to therapeutic drugs,

preventing uptake as pH gradients change across the membrane [12]. This, in turn,

induces the counter transport of drugs reducing concentrations inside cells and thus

effectiveness. This is true not only for TNBC patients treated with cytotoxic drugs but

also estrogen receptor (ER)- and HER2-positive patients treated with targeted therapies

[3].

Important modulators of the hypoxic TM include membrane-bound carbonic

anhydrases (CAs), which are zinc metalloenzymes that catalyze the reversible hydration

of CO_2 [16, 20-26]. In breast cancer, carbonic anhydrase IX (CA IX) expression is

upregulated in hypoxic tissue and is considered a prognostic marker for the TNBC

phenotype [27-30]. In normal cells, CA IX expression is limited primarily to gut

epithelium, but is upregulated in many hypoxic tumors including breast cancer [28, 31-

43]. CA IX is not only a marker for hypoxia but may also serve as a therapeutic target

for the treatment of aggressive breast cancer. CA XII expression, on the other hand, is

not regulated by hypoxia and is an indicator of better patient outcome in several types of

cancers, including breast cancer [44]. Its expression is under the control of the ER and

associated with ER-positive breast carcinomas [44-47]. CA XII is also expressed more

abundantly across tissue types, including normal breast tissue [41, 48].

The role of these carbonic anhydrases in cancer progression are under intense

investigation, particularly that of CA IX. CA IX was originally identified as a subunit of

the MaTu protein and a short time later characterized as a novel member of the

carbonic anhydrase family [49, 50]. While it is normally localize to gut epithelium, it is

strongly upregulated in tumors, generally in more aggressive phenotypes [51, 52]. CA

IX is a transmembrane protein, organized as a dimer, with the catalytic domain facing

the extracellular space. This enzyme catalyzes the reversible hydration of CO_2 forming

a bicarbonate ion and a proton, in the presence of water. The pericellular space, with

which CA IX associates, is rich in metabolic acids such as lactate, protons, and CO_2. It

has been postulated that CA IX cooperates with bicarbonate transporters by providing

bicarbonate for intracellular alkalinization at the same time that it acidifies the tumor

microenvironment [53]. Data supporting this hypothesis have been provided through

studies in cell lines and cancer cell-derived spheroids [34, 54, 55]. However, CA IX, like

other members of the family, is a reversible enzyme. We, and others, have

hypothesized that CA IX may stabilize extracellular pH at a value that favors cancer

cells growth and proliferation [27, 56]. New data from studies with CA IX expressing

xenografts support this notion that CA IX acts as an extracellular pH-stat [57]. This adds

to the therapeutic rationale for pharmacological interference with CA IX activity.

Sulfonamides are the most studied class of CA inhibitors (CAIs) with several

compounds in clinical use for the treatment of diseases including glaucoma, altitude

sickness, and epileptic seizures [25, 58-62]. Although these sulfonamides are potent

CAIs, most lack isoform selectivity and do not specifically target the tumor-associated

isoforms CA IX and CA XII [60]. This lack of specificity is attributable to the highly

conserved residues within the active site of CAs [63]. These similarities have challenged

the development of isoform-specific inhibitors, causing competition between isoforms,

which reduce efficacy. This is especially true in the case of CA II. This isoform has the

widest tissue distribution and is abundantly expressed in red blood cells (representing

5% of all proteins) [64, 65]. As most drugs are delivered systemically and are

membrane permeable, it is likely that CA II will sequester CA inhibitors reducing

circulating concentrations limiting exposure within tumors. That said, none of the clinical

CAIs have been used for the treatment of cancer. Ureido-substituted benzene

sulfonamides (USBs) are potent and selective inhibitors of recombinant CA IX and CA

XII activity [66-68]. X-ray crystallographic studies recently provided insight into the

mechanism of USB-mediated selective inhibition of CA IX over CA II [66]. Those studies

showed that the USBs interact more favorably with residues in the active site of CA IX

versus CA II [66]. These inhibitors also showed both anti-proliferative and anti-

metastatic properties in preclinical models [29, 68]. One of the USBs (SLC-0111)

recently completed phase I clinical trials and is scheduled to begin phase II trials for the

treatment of solid tumors that over express CA IX (clinical trials.gov, NCT02215850).

The goal of this study was to distinguish the mechanistic effects of USB-mediated

inhibition of CA IX and CA XII in the context of breast cancer cells. To accomplish this,

three USBs (including SLC-0111) were studied, using *in-silico* modeling to understand

binding mechanisms and their efficacy was tested in cell lines representing TNBC and

ER-positive breast cancers. With this approach, differences in residues within the active

sites of CA IX and CA XII indicated that these drive isoform-specific interactions

affecting affinity of the USBs. The USB compounds were also shown to inhibit CA

activity with surprising specificity for CA IX in TNBC cells, relative to CA XII in ER-

positive cells given similar K_i values for purified protein. This same specificity appears to

drive USB-mediated growth inhibition. Yet the K_i values for inhibiting catalytic activity in

CA IX and CA XII in cells is two and three orders, respectively, less efficacious than

would be expected based on activity studies using purified protein and binding kinetics

based on structures. In addition, growth inhibition requires even higher concentrations

of inhibitors than that needed for inhibition of activity in cells. Further, cells that lack

expression of CA IX or CA XII are still sensitive to USB-mediated inhibition of cell

growth. Despite the excellent inhibition of the USBs against purified CA IX and CA XII,

and the specificity for CA IX over CA XII in cells, we must consider the possibility that

CA activity may not be the target for sulfonamide action with respect to cell growth

inhibition.

Materials and methods

Ureido-substituted benzene sulfonamide (USB) inhibitors were prepared in dimethyl sulfoxide (DMSO) at stock concentrations of 100 mM and further diluted to appropriate concentrations in cell culture medium before use. Control experiments contain DMSO only. CA II antibody was obtained from Novus Biological (Littleton, CO), CA IX (M75) antibody was a gift from Egbert Oosterwijk, CA XII and GAPDH antibodies were purchased from R&D systems (Minneapolis, MN) and Cell Signaling Technology (Beverly, MA), respectively.

In Silico Modeling

X-ray crystallography structures of USB compounds in complex with purified CA II and a CA IX-mimic (analogous site directed mutagenesis of residues in the active site of CA II to resemble CA IX) were previously obtained and published elsewhere [66], but shown for comparison to the modeled CA XII complex in the present study (Figure 1). Each of the USB compounds were superimposed into CA XII structures (PDB ID: 1JCZ) based on the crystal structures of the compounds using Coot [69, 70] and adjusted based on steric hindrance constraints. These structures were also energy minimized using the Chimera Minimize structure feature (data not shown) to ensure that these modeled compounds were at their energy minima.

Cell Culture

MCF 10A, UFH-001, and T47D cells were maintained as previously described [58]. For hypoxia treatment, cells were exposed to 1% O_2, 5% CO_2 and balanced N_2 for 16 h

using a Billups Rothenberg Metabolic Chamber. UFH-001 cells are a newly

characterized line that exhibits the triple negative (TNBC) phenotype [71, 72]. These

cells do not express ER, the progesterone receptor (PR), or HER2. However, they do

express EGFR, also typical of the TNBC phenotype. In addition, these cells express CA

IX but not CA XII. The T47D cells are a luminal ER-positive line that expresses CA XII

but not CA IX. The MCF 10A line is used as a control and expresses CA IX only under

hypoxic conditions. All cell lines were authenticated prior to conducting experiments.

CA IX deletion by CRISPR/Cas9

Ablation of CA IX in UFH-001 cells has been described previously [72].

pLentiCRISPR v2 was a gift from Dr. Feng Zhang (Addgene, Cambridge, MA) and was

used to knockout CA IX in UFH-001 cells. Briefly, guide RNA sequences within the first

coding exon were identified downstream of the CA IX translational start site using an

online design tool (crispr.cos.uni-heidelberg.de) and three non-overlapping gRNAs were

chosen within this region of the *CA9* gene. Complementary double-stranded

oligonucleotides were cloned into BsmBI digested pLentiCRISPR v2, recombinant

clones identified by restriction analysis and confirmed by automated Sanger

sequencing. Lentivirus was formed from these CA IX sgRNA plasmids and empty

lentiCRISPR v2 plasmid as previously outlined [72]. UFH-001 cells were transduced

and selected with puromycin (2 µg/mL) for 3 weeks. Stably transduced cells were

harvested, and CA IX depletion was confirmed by western blotting.

Knockdown of CA XII by shRNA lentiviral particles

Knockdown of CA XII expression in T47D cells has also been previously described [72]. Briefly, cells were transfected with short hairpin RNA (shRNA) lentiviral particles obtained from Thermo Fisher Scientific (Waltham, MA) against CA XII. Cells were seeded at $5x10^4$ cells/well in 24-well plates and grown for 24 h. Then, cells were infected with lentivirus in serum-free medium for 6 h. The cells were further incubated with normal growth medium for 24 h. GFP expression was monitored to confirm the efficiency of transduction. Stable cells were established by puromycin (2 µg/mL) selection (Sigma Aldrich, St. Louis, MO). CA XII knockdown was confirmed by western blotting.

Western Blot Analysis

Western blot analysis was performed, as previously described [28]. Briefly, cells were collected after drug treatments for 48 h and washed 3x with ice-cold phosphate buffered saline (PBS) and lysed in RIPA buffer (containing 1% orthovanadate, 25 mM NaF) supplemented with protease inhibitor for 15 min on ice. Lysates were collected and clarified by centrifugation at 55,000 rpm for 60 min at 4°C. Clarified supernatants were collected and stored at -20°C. Protein concentration was determined using the Markwell modification of the Lowry procedure. Equal amounts of cell lysate (30 µg) were subjected to SDS-polyacrylamide electrophoresis, prepared as described by Laemmli et al. [73]. Samples were electrotransferred from SDS-PAGE gels to nitrocellulose membranes in transfer buffer at 200 mA for 2 h at 4°C. After incubation with selective primary and secondary antibodies, proteins were visualized using ECL reagent

253 according to the manufacturer's directions (GE Healthcare, Wauwatosa, WI). Images

254 were scanned and cropped using Adobe Photoshop version 11.

Membrane Inlet for Mass Spectrometry

CA catalysis was measured by the exchange of ^{18}O from species of $^{13}CO_2$ into water

determined by membrane inlet mass spectrometry (MIMS) [74, 75]. The application of

this method in demonstrating exofacial CA activity in breast cancer cells has been

described previously [27] and more recently in Chen et al. [72]. To decrease

inaccuracies arising from ^{12}C-containing CO_2 in cell preparations, ^{13}C- and ^{18}O-enriched

CO_2/HCO_3 were used and the rate of depletion of ^{18}O in ^{13}C-containing CO_2 measured.

Specifically, the atom fraction of ^{18}O in extracellular CO_2 was determined using peak

heights from the mass spectrometer representing CO_2 [½ (47) + (49)] /[(45) + (47) +

(49)] where the numbers in parentheses represent the peak heights of the

corresponding masses of specific species of isotopic CO_2. The MIMS assay actually

measures CA activity in the dehydration direction, because only isotopic variations in

CO_2 are measured by the mass spec (although activity in both directions is required for

the decrease in the isotopic distribution of ^{18}O in CO_2). A membrane barrier (to the mass

spec) prevents bicarbonate concentration from being measured by the mass spec.

When added to the reaction vessels, the cells are exposed to an equilibrated solution of

$^{13}C^{18}O_2:HC^{18}O_3^-$ at a ratio of about 5%: 95%, respectively (25 mM, total). The total

amount of CO_2 does not change over the course of the experiment, just the isotopic

enrichment. Mass spectra were obtained on an Extrel EXM-200 mass spectrometer

using electron ionization (70 eV) at an emission current of a 1 mA. First order rate

constants (k) were determined for CA activity. These were calculated from the change

in atom fraction enrichment (^{18}O in CO_2) using: $A = A_0\, e^{-kt}$, where (A) is the fraction of

enrichment at a specific end point, and (A_0) is the starting enrichment at a specific point.

Cells were collected from culture plates using cell release buffer (Gibco, Waltham,

MA), washed with bicarbonate-free DMEM buffered containing 25 mM Hepes (pH 7.4),

and counted. Cells (5.0×10^5 cells/mL, unless otherwise stated) were added to a

reaction vessel containing 2 mL of Hepes-buffered, bicarbonate-free DMEM at 16°C in

which ^{18}O-enriched $^{13}CO^2$ /H$^{13}CO_3$ at 25 mM total $^{13}CO_2$ species was dissolved.

Catalysis by CA, in the presence or absence of CA inhibitors, was measured after

addition of cells (exposed to normoxic or hypoxic conditions) by the exchange of ^{18}O

species from species of $^{13}CO_2$ into water determined by measuring CO_2 mass by MIMS

as described above. This exchange is a specific measure of CA activity during the time

frame of the assay. A general example of this assay is provided in Supporting

Information (S1 Fig) showing the effect of two classical sulfonamide based inhibitors on

CA activity in UFH-001 and T47D cells, which is further discussed in the results.

Ki values were calculated using the following equation:

(V - unc) = (Vo - unc) / (1 + I / Ki),

where (V) is the rate at a specific inhibitor concentration, (unc) is the uncatalyzed rate,

(Vo) is rate in the absence of inhibitor, and (I) is inhibitor concentration. For tight binding

inhibitors, IC_{50} = [enzyme]/2 + Ki (app). However, when the K_i >> greater than the

enzyme concentration, $IC_{50} \simeq K_i$. In our experiments, enzyme concentration in cells is

in the low nM range, based on titrations with purified CA II. The K_i values that we

calculate are much higher than [enzyme] for either UFH-001 or T47 D cells, despite the

difference in enzyme concentration between the two cell lines.

Cell Growth Assay

Cell growth in the presence or absence of compounds was analyzed using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT assays). Briefly, cells were seeded in 96-well plates and treated with different concentrations of compounds after 24 h. The cells were then incubated under normoxic or hypoxic conditions under the different treatment conditions for 48 h and/or 96 h. The medium was replaced every two days. MTT reagent (10% v/v of 5 mg/mL) was added to the cells after each time point and the cells were incubated for an additional 4 h. The medium was removed without disturbing the cells and DMSO was added to each well. The DMSO dissolved cells were incubated while shaking for 15 min in the dark. Absorbance was measured at 570 nm using an Epoch microplate reader (Biotek, Winooski, VT). IC_{50} values (inhibitor concentration where the response is reduced by half) were calculated using Prism 7.0c for Macs. Non-linear regression analysis was performed on % dose response with log plots interpolating the value at 50% from the least squares fit of the growth curves.

Lactate Dehydrogenase Release Assay

Cytotoxicity was estimated using the release of lactate dehydrogenase (LDH) activity (Sigma-Aldrich, St. Louis, MO). Cells were plated in 96-well plates. USB compounds were added the next day. After 48 h, medium was collected from each well to measure the amount of released LDH activity. The amount of LDH released from each sample was measured at room temperature at 450 nm using a BioTek Epoch microplate reader. LDH activity is reported as nmol/min/mL of medium.

Caspase-3 Activity Assay

Caspase-3 activity assay was performed using an Apo-alert colorimetric caspase assay kit (BioVision, Milpitas, CA). Cells were plated onto 10 cm dishes and treated with compounds 24 h later. 48 h after various treatments, cells were collected, and lysates prepared. Protein concentration was determined, and equal volumes of lysates were used for caspase-3 activity assay, measured at 405 nm in a microtiter plate. The reader detected chromophore p-nitroaniline (pNA) after its cleavage by caspase-3 from the labeled caspase-3 specific substrate, DEVD-pNA. These data are presented as pmol of pNA per µg of cell lysate per hour of incubation. Staurosporine (Sigma-Aldrich, St. Louis, MO) was used as a positive control.

Analysis of Cell Cycle Progression

Cells were seeded in 24-well plates. After 24 h, USB compounds were added to the respective wells and incubated for 48 h. Cells were released with trypsin, harvested, fixed with 500 µL of 50% ethanol in tubes and collected by centrifugation at 13,000 rpm for 5 min. Cell pellets were re-suspended in 500 µL. Propidium Iodine (10 µg/mL) containing 300 µg/mL RNase (Invitrogen, Waltham, MA). Then, cells were incubated at room temperature in the dark for 15 min. Cell cycle distribution data in Figure 8C-8E (gated for live, diploid cells only) was analyzed using the FCS Express Version 5 from De Novo Software (Glendale, CA) from data obtained using FACS caliber (Becton Dickson, CA). The representative images in the text (Fig 8A and 8B) were generated by ModFit LT.

Cell Migration and Invasion Assays

Serum starved cells were plated at a density of 50,000 cells in 300 µL per insert in 24-well cell migration and invasion plates (Cell BioLabs, San Diego, CA). The cells were allowed to migrate (24 h) or invade (48 h) from the insert containing serum free medium, with or without USB compounds, towards the well that had medium containing 10% FBS. Fixing and staining cells terminated the assay. Images were then collected and analyzed. All cell lines were maintained at $37^{\circ}C$ in humidified air with 5% CO_2 for the duration of the experiment.

Spheroid Formation Assay

UFH-001, UFH-001 CA IX KO, T47D, and T47D CA XII KO cells (10,000 cells in a total volume of 100 µL) were plated into 96-well, low attachment microplates (Corning, NY). Cells were incubated up to 96 h in 5% CO_2 at 37°C. Spheroids were imaged at specific intervals using the EVOS microscope system (Thermo Fisher Scientific, Waltham, MA).

Statistical Analysis

Data are presented as means ± SEM. Differences between the treatment groups were analyzed using Student's t-test in GraphPad Prism, version 7.0a: p-values < 0.05 was considered statistically significant.

Results

Isoform selective binding and interaction of USB compounds

to CA IX and CA XII

Cytosolic CA II exists as a monomer (Fig 1A), while both membrane-bound CA IX

and CA XII have been shown to form dimers. The catalytic domains of these dimers are

shown in Fig 1C and 1E. The structural formulae of the USB inhibitors used in this study

and their respective K_i values for recombinant CA II, IX and XII were determined

elsewhere and provided in Table 1 [66-68]. These previously published data show that

two of the three USB compounds (U-CH$_3$ and U-F, also called SLC-0111) selectively

bind to and inhibit the activity of the tumor-associated isoforms CA IX and XII relative to

the off-target CA II [66-68]. Isoform selectivity towards recombinant CA IX and CA XII

was evident in the differences in K_i values for these isoforms relative to CA II, obtained

using Applied Photophysics stopped-flow kinetics of purified recombinant proteins.

Isoform selectivity is important *in vivo*, where inhibition of CA II (one of the predominant

isoforms in red blood cells) might cause unintended consequences and sequester

inhibitor reducing the concentration within the tumor.

To investigate the underlying mechanism of binding, we previously determined the

crystallographic structures of all three compounds bound to CA II (Fig 1B) and the CA

IX-mimic (Fig 1D) [66]. As expected, each compound was shown to directly interact

with the zinc ion, via the sulfonamide moiety [66]. The extended tail moieties of the USB

compounds also interacted with residues, between 10-15 Å away from the zinc, favoring

the hydrophobic region of the active site (Fig 1B and 1D). This region is located close to

the entrance of the active site and has been termed the "selective pocket" [76, 77]. This

is because it has a clustering of several amino acid residues that are different among

CA isoforms (Table 2). For example, residue 131, located within the selective pocket, is

different among the isoforms. In CA II, this residue is a phenylalanine (CA II residue

numbering will be used throughout this manuscript). Its bulky side chain causes steric

hindrance and therefore induces conformation changes in the tail groups of the bound

compounds (Fig 1B, Table 1). In CA IX, residue 131 is a valine, which allows for more

favorable binding and therefore greater isoform selectivity towards CA IX compared to

CA II (Fig 1D, Table 1). We believe that the differences in active site residues, located

within the selective pocket, are responsible for the observed differences in K_i values for

bound compounds (Table 1). By comparison, the classical CA inhibitor, acetazolamide,

can only interact with the active site zinc ion and with residues within 10 Å of the zinc,

because of its shorter tail moiety [78]. Residues in this region of the active site are

among the most conserved between isoforms [63].

To compare the CA II and CA IX structural data with that of CA XII, we have now

performed *in silico* modeling of the three USB compounds (U-CH$_3$, U-F and U-NO$_2$) into

the active site of CA XII (Fig 1F). Residue 131 in CA XII is an alanine (Table 2), which

like valine (in CA IX) has a small hydrophobic side chain that results in favorable binding

and selectivity towards this isoform (Table 1). In addition, CA XII has a serine (a polar

uncharged side chain) at position 132, which is also located close to the entrance of the

active site. As such, this residue also causes steric hindrance that induces

conformational changes in all three sulfonamides, enhancing binding and inhibition

(Table 1). Additionally, serine 132 and also serine 135 (both charged residues and also

unique to CA XII) create a narrow hydrophobic region within the CA XII active site (Fig

1F). This causes an energetically favorable change, especially, in the conformation of

compound U-F (relative to CA II and CA IX) within the active site of CA XII. These

residues also prevent steric clashes with residue 131 and facilitate compound

interactions in the narrow hydrophobic region within the selective pocket. These

interactions and the resulting conformational change in U-F will be specific to CA XII,

because neither CA II nor CA IX has serine residues at positions 132 or 135 (Fig 1 and

Table 2).

Fig 1. USBs bound in the active site of CAs. Panel A. Ribbon diagram of monomeric
CA isoform II (gray). Panel C. Ribbon diagram of the catalytic domains of dimeric CA
isoform IX (cyan). Panel E. Ribbon diagram of the catalytic domains of dimeric CA
isoform XII (wheat). Surface representation of compounds U-CH$_3$ (pink), U-F (green)
and U-NO$_2$ (yellow) in complex with monomers of CA II (Panel B), the CA IX-mimic
(Panel D), and modeled into the active site of CA XII (Panel F). Catalytic zinc (magenta
sphere), hydrophilic (blue) and hydrophobic (orange) residues are as shown. Red
double-headed arrows indicate isoform specificity relative to residue 131 (labeled in
white). These arrows also show flexibility in tail conformations seen in CA II and CA IX
but not in CA XII. Previously published K$_i$ values of each compound bound to purified
CA II, CA IX and CA XII are noted next to the inhibitor name [67, 68]. Figures were
designed using PyMol.
436

Effect of USB inhibitors on CA IX and CA XII activity in breast

cancer cells

One of the goals of this study was to determine if these high affinity inhibitors (for CA

IX and CA XII) were equally effective in the context of the cellular environment. To

determine the effect of USB compounds on CA activity in breast cancer cells, cell lines

that only express CA IX or CA XII at the cell surface were selected. Based on previous

work, two lines were selected based on their strong expression of CA IX or CA XII. The

T47D line is a well-studied ER-positive, luminal line. UFH-001 cells represent a new

line, with a triple negative phenotype, arising from MCF 10A cells. MCF 10A cells were

derived from a patient with fibrocystic disease, which spontaneously immortalized in cell

culture [79, 80]. These are considered by many as the control line for breast cancer

cells, although these cells map with other lines that have the basal/triple negative

phenotype [81]. The UFH-001 cells are an especially aggressive that, unlike MCF 10A

cells, form tumors in nude mice. Characterization of the UFH-001 cells is described in

two recent publications [71, 72]. When compared to control MCF 10A cells, UFH-001

cells show greater expression of CA IX protein (Fig 2A) and mRNA (S2A Fig) under

normoxic conditions. However, under hypoxic conditions both MCF 10A and UFH-001

cells upregulate CA IX expression (Fig 2A). It can also be noted that there is a

difference in the migration patterns of CA IX between the two cell lines. In UFH-001

cells, CA IX migrates as a doublet with molecular weights of 54 and 58 kDa, both of

which are glycosylated [82]. However, in MCF 10A cells, CA IX migrates as a singlet

with a molecular weight slightly lower than 58 kDa. The observed differences in

molecular weight may result from differences or deletions in primary sequence. CA XII

expression is limited to the T47D cells (Fig 2B and S2B Fig) and is independent of

hypoxia (Fig 2B). Expression of cytosolic CA II protein is only observed in UFH-001

cells and is also not sensitive to hypoxia (Fig 2A). The upregulation of CA IX in hypoxic

UFH-001 cells underlies the increase in cell surface CA activity in hypoxic versus

normoxic cells (Fig 2C) [27]. That there is no change in CA activity in T47D cells in

response to hypoxia relative to normoxia also correlates with the lack of change in CA

XII expression (Fig 2D).

Fig 2. CA expression and activity in breast cell lines. Panel A. Protein expression in
a normal immortalized basal type breast cell line (MCF 10A) and a triple negative breast
cancer (TNBC) cell line (UFH-001), using western blot analysis (under normoxic (N) or
hypoxic conditions (H) for 16 h, respectively). Panel B. Immunoblots for CAs II, IX and
XII protein expression in MCF 10A and the luminal ER positive breast cancer cell line
(T47D). Panels C and D. UFH-001 and T47D cells were grown for 3 days at which point
they were exposed to normoxia or 16 h of hypoxia and the cells assayed for CA IX and

XII activity, respectively, using the MIMS assay. First order rate constants for CA activity
in UFH-001 cells (Panel A) and T47D cells (Panel B), as described in the methods, are
shown. These data represent three independent experiments and are reported as the
mean ± SEM.
478

The inhibitory capacity of USB compounds U-CH$_3$, U-F, and U-NO$_2$ on CA IX and

CA XII activity in UFH-001 and T47D cells, respectfully, were studied under normoxic

and hypoxic conditions, but only the representative normoxic plots are shown (Fig 3).

Because UFH-001 cells express both CA II and CA IX, the progress curves that follow

CA activity are biphasic on a semi log plot (Fig 3A-3C). The first phase (phase I, 20-40

sec after addition of cells) reflects CA II activity. The second phase (phase II, 100-400

sec) reflects CA IX activity. Phase I represents the rapid diffusion of $^{13}C^{18}O_2$ into cells

where CA II catalytic cycles deplete ^{18}O from $H^{13}C^{18}O_3$ followed by efflux of $^{13}CO_2$

species from the cell. The concentration of CO$_2$ and the isotopic forms of CO$_2$ are

measured by the mass spec. There is little change in the slope of this phase between

normoxic and hypoxic cells, although the length of time in this phase is shortened (Fig

3A-3C). Phase II represents the hydration-dehydration activity of CO_2/HCO_3^- outside of

the cell representing CA IX activity. The first order rate constant of phase II is greater for

hypoxic cells compared with normoxic cells (Fig 2C). This indicates a higher rate of CO$_2$

hydration/dehydration on the extracellular face of hypoxic UFH-001 cells, which we

associate with enhanced expression of exofacial CA IX (Fig 2A and 2C).

The results in Fig 3 also show that USB compounds have limited effects on phase I

in the UFH-001 progress curves under either normoxic (Fig 3A-3C) or hypoxic

conditions (data not shown). This means that during the time course of these

experiments, the inhibitors did not permeate the membrane to inhibit CA II except at

high concentrations of U-F (SLC-0111) which was observed as an increase in the

atomic fraction of ^{18}O in $^{13}CO_2$ (starting at ~150 seconds, Fig 3B). All three CA inhibitors

had significant effects on phase II of the progress curve indicative of a decrease in CA

IX activity (Fig 3A-3C). In other words, the slopes of the phase II progress curve (used

to calculate first order rate constants in Fig 2) decline in the presence of the inhibitors.

Based on K_i values in normoxic and hypoxic UFH-001 cells, the most potent inhibitor

was U-F (SLC-0111) and the least inhibitory was U-NO_2 (Fig 3G). Hypoxia did not

significantly affect the efficacy of any of the USB compounds. These plots are

representative of multiple experiments but show a limited number of inhibitor

concentrations. The type of experiments that are used to actually calculate K_i values

are shown in S3 Fig where an extensive range of concentrations (for U-NO_2 in these

cases) was utilized. In addition, we have included the effect of two classical sulfonamide

inhibitors on CA activity in UFH-001 and T47D cells, that differ in their cell permeability

(S1 Fig). Acetazolamide (ACZ) is impermeant during the time course of the experiment

and is expected to block only exofacial CA activity, like CA IX and CA XII.

Ethoxzolamide (EZA) is permeant, and quickly inhibits both exofacial and intracellular

CAs. Activity in UFH-001 cells (S1A Fig), is, once again, observed as a biphasic

progress curve. The presence of ACZ blocks only exofacial CA activity which confirms

that the second phase of the progress curve (from about 200 to 600 sec) represents CA

IX activity. The activity in the presence of EZA represents the spontaneous (non-

catalytic) interconversion of CO_2 and HCO_3^-, which is essentially an extension of the

activity before addition of cells. First order rate constants are shown for each of the

progress curves (S1A Fig). Finally, we present data on CA activity in cells in which CA

IX is ablated (Fig 4A). Here, we show that CA activity is reduced in CA IX KO cells

relative to the EV (hypoxic) controls. However, inhibition by the impermeant

sulfonamide, N-3500, is better at blocking CA activity than is CA IX ablation. It is

possible that there are other exofacial CA's present on the surface of UFH-001 cells.

We have only tested for the expression of CA XIV, which is not detected under

conditions in which we observe CA IX. While microarray data (S2A Fig) show CA XIV

mRNA expression in UFH-001 cells, earlier northern blot analysis revealed no CA XIV

mRNA [23]. Furthermore, no difference in CA XIV mRNA expression was observed

between MCF 10A and UFH-001 cells, under normoxic conditions (S2A Fig). As far as

we are aware, no one has shown positive expression of either CA XIV or CA IV (the

other membrane bound CA) at the protein level in breast cancer cells.

T47D cells, which express membrane-bound CA XII (Fig 2B), exhibit single-phase

progress curves (Fig 3D-3F), because they lack CA II. This activity specifically reflects

exofacial CA XII activity. This activity was not sensitive to hypoxia (Fig 2D), which is

consistent with lack of CA XII induction by hypoxia (Fig 2B). Like CA IX, USB

compounds inhibit CA XII activity (Fig 3D-3F). Again, the slope of the progress curve

decreases in the presence of inhibitor. However, there is a striking difference in efficacy

that is reflected in a significant increase in the K_i values obtained in T47D cells versus

UFH-001 cells (Fig 3G). Again, an extensive range of concentrations are used to

calculate K_i and shown for one of the inhibitors (U-NO$_2$) in S3 Fig. Unlike UFH-001 cells,

there were only slight differences in inhibitory effects of the different USB's in T47D

cells, which again was not affected by normoxic or hypoxic conditions (Fig 3G). S1B

Figure shows the progress curves in the presence of ACZ and EZA. Catalysis in the

absence of inhibitors is linear. Inhibition by ACZ generates a progress curve that

resembles that of USB compounds. The activity in the presence of EZA once again

represents the spontaneous (non-catalytic) interconversion of CO_2 and HCO_3^-. These

experiments were performed at 16°C, as were all of the MIMS experiments in this

manuscript, so can be directly compared. CA XII knockdown significantly reduced CA

activity in T47D cells relative to EV controls (Fig 4B). Inhibition of CA activity with N-

3500 is only slightly more effective than the knockdown of CA XII expression.

Fig 3. Effects of USBs on CA IX and CA XII activity in breast cancer cell lines.
Panels A-C. UFH-001 cells were grown for 4 days under normoxic conditions. CA IX
activity (0.5×10^6 cells/mL, unless otherwise indicated) was assayed using MIMS in the
presence or absence of U-CH$_3$ (Panel A), U-F (Panel B) or U-NO$_2$ (Panel C). Panels D-
F. T47D cells prepared similarly to UFH-001 cells, but cultured for 6 days, were also
assayed for CA XII activity (0.5×10^6 cells/mL) in the absence or presence of the same
USB based inhibitors: U-CH$_3$ (Panel D), U-F (Panel E) or U-NO$_2$ (Panel F). Atom
fractions of ^{18}O in CO$_2$ were collected continuously. For ease of illustration, data points
at 25-s intervals are shown. Phase I in UFH-001 cells indicate CA II activity and Phase
II indicate CA IX activity. In T47D cells, the progress curves are linear and
representative of CA XII activity. Arrows indicate time point at which cells were added.
Panel G. K$_i$ values of each compound in UFH-001 and T47D cells, under normoxic or
hypoxic (16 h) conditions, are shown. Data are representative of the average of
triplicate experiments ± SEM.
566
Fig 4. Effect of CA IX and CA XII knockdown on CA activity. CA IX knockout in
UFH-001 cells was accomplished using Crispr/Cas9 technology. CA XII knockdown in
T47D cells was performed using shRNAi lentiviral strategies. Cells were grown similarly
to those described in Fig 3. Panel A. CA activity was measured in 1 x 10^6 UFH-001 cells
(EV controls, CA IX KO, or cells treated with the pegylated sulfonamide, N-3500).
Panel B. CA activity was measured in 5 x 10^5 T47D cells (EV controls, CA XII KO, or
cells treated with N-3500. Data represent duplicate experiments.
574

Impact of USB-based inhibitors on breast cancer cell growth,

cytotoxicity, and apoptosis

MTT, lactate dehydrogenase (LDH), and caspase activity assays were used to test

the efficacy of CA IX and CA XII inhibition on breast cancer cell growth, viability and

apoptosis, respectively. These results show that all three compounds decreased cell

growth (MTT assay) in the different cell lines under normoxic conditions (Fig 5, Table 3).

No significant trend in compound effectiveness was established between the cells

treated for 48 h compared to 96 h (Fig 5 and Table 3). The most effective compound at

inhibiting growth was U-NO$_2$ in UFH-001 cells (IC$_{50}$ ~25 μM). This compound also

inhibited the growth of MCF 10A cells in addition to that of the T47D cells, but requiring

nearly three times the concentration compared to UFH-001 cells (Table 3). The MCF

10A cells were the most resistant to compounds U-CH$_3$ and U-F, with IC$_{50}$ values

observed in the range of 500 µM. The concentration required to block growth of T47D

cells was also in this range at 48 h but improved somewhat at 96 h of exposure (Table

3). We repeated the experiments under hypoxic conditions (Fig 6) with similar results.

In addition, we tested the effects of the USB's on cells in which CA IX or CA XII was

knocked out in UFH-001 (UFH-001 KO) or T47D (T47D KO) cells, respectively, which

have been previously characterized [72]. Surprisingly, the result was the same, i.e., the

USB's still blocked cell growth at high concentrations even in the absence of CA IX or

CA XII expression.

Fig 5. **Effects of USBs on breast cancer cell growth.** Breast cancer cell lines grown
under normal culture conditions for 24 h were exposed to compounds for 48 h [U-CH$_3$
(Panel A), U-F (Panel B), U-NO$_2$ (Panel D)] or 96 h [U-CH$_3$ (Panel D), U-F (Panel E), or
U-NO$_2$ (Panel F)]. MTT assay was performed at 48 h and 96 h, respectively. Data
shown are an average of at least three independent experiments and are represented
as the mean ± SEM.
601
Fig 6. Effects of USBs on cell growth in the presence or absence of hypoxia or in
CA knockout cells. One day after plating, breast cancer cell lines were exposed to
normoxic or hypoxic conditions for 48 h in the presence of U-CH$_3$, U-F, or U-NO$_2$ at the
given concentrations in UFH-001 empty vector control (EV) cells (Panel A), UFH-001
CA IX KO (UFH-001 KO) cells (Panel B), T47D EV cells (Panel C) and T47D CA XII KO
(T47D KO) cells (Panel D). MTT assay was performed and shown as % cell growth.
Data shown represent at least three independent experiments and are shown as the
mean ± SEM.
610

Both UFH-001 and T47D cells are able to form spheroids (S4 Fig). UFH-001 cells

form spheroids as early as 24 h after plating (even when the plating density is low). The

spheroids appear dense and round. The T47D cells also form spheroids, which are

somewhat larger than UFH-001 spheroids. Neither the loss of CA IX nor CA XII affects

spheroid formation over the 96 h observation period. We were unable to measure

activity in these spheroids because of the limited number of cells.

To specifically test for cytotoxicity, LDH release over 48 h in the presence or

absence of the USB compounds was measured. The inhibitors caused only limited

cytotoxicity when exposed to cells (Fig 7), relative to a positive control (S5 Fig). UFH-

001 cells were the most sensitive, but only at high concentrations of compound U-NO$_2$

(Fig 7). The MCF 10A cells were also somewhat sensitive to U-NO$_2$ but again at high

concentrations (Fig 7A). T47D cells showed no loss of membrane permeability in

response to inhibitor treatment (Fig 7C).

Fig 7. **Effects of USBs on breast cancer cell viability.** The cytotoxic effects of USBs
were evaluated using the LDH release assay. MCF 10A (Panel A), UFH-001 (Panel B)
and T47D (Panel C) cells were grown in 96-well plates and exposed to U-CH$_3$, U-F or
U-NO$_2$ for 48 h, under normoxic conditions. LDH release was assayed after treatment,
results were evaluated, and data analyzed using Prism. Data shown are representative
of three independent experiments and as the mean ± SEM, *p < 0.05.
630

These observations led us to question whether the growth arrest and (limited)

cytotoxicity observed with the inhibitors was due to the activation of apoptosis. To test

this, caspase activity assays were employed. These data revealed that apoptosis was

not activated, as caspase activity did not change in either UFH-001 or T47D cells in the

presence of USB compounds (S6A and S6B Fig, respectively).

Because of the USB-induced growth arrest, albeit at high concentrations, the effects

on cell cycle transition were evaluated after exposure to inhibitors for 48 h, under

normoxic conditions (Fig 8). Of the diploid population, ~ 40% of the UFH-001 cells were

in the G1 phase of the cell cycle (Dp G1) and a much lower percentage were in G2

phase (Dp G2). The number of T47D cells in G1 (~ 60%) was significantly higher than

the number of cells in the G2 phase of the cell cycle (~ 20%). A significantly higher

percentage of cells were observed in the G1 phase in T47D compared to UFH-001

cells, while cells in G2 phase were much higher in UFH-001 than T47D cells (Fig 8C). In

the presence of USB compounds, there was no change in the number of UFH-001 cells

in S phase when compared to the control cells (Fig 8D). However, a significant

decrease in the number of T47D cells in S phase was noted with exposure to U-CH$_3$

and U-F, but not U-NO$_2$ (Fig 8E). Higher concentrations of U-CH$_3$ and U-F where used

for experiments with T47D cells, because both the K$_i$ values obtained from the MIMS

experiments (Fig 3G) and the IC$_{50}$ values obtained from the cell growth assays (Table 3)

were much higher for these compounds relative to compound U-NO$_2$, and by

comparison to the more sensitive UFH-001 cells.

Fig 8. **Effects of USBs on cell cycle transition in breast cancer cells.** Cell cycle
analysis was performed in UFH-001 (Panel A) and T47D (Panel B) cells treated with
varying concentrations of USB compounds for 48 h, under normoxic conditions. Post
treatment, cells were stained with Propidium Iodine containing RNase A, data was
obtained using the FACS caliber instrument and results analyzed with FCS Express and
ModFit LT softwares. The percentage of breast cancer cells in G-phase (Panel C), all
phases of the cell cycle in UFH-001 (Panel D) and T47D cells (Panel E), were quantified
using Prism. Data are represented as mean of at least three independent experiments ±
SEM and NC = negative control.

Effect of USB inhibitors on CA expression in breast cancer

cells

We next examined if inhibition of CA IX and CA XII activity induced an upregulation

of the targets or expression of other CA isoforms as a compensatory mechanism. To

achieve this, UFH-001 and T47D cells were treated with increasing concentrations of

USB compounds for 48 h, under normoxic conditions. Immunoblotting with cell lysates

was performed using antibodies against CA II, CA IX and CA XII. No upregulation of

other CA isoforms was observed in response to USB treatment (S7 Fig). However,

there were changes in the endogenously expressed CAs. For instance with U-CH$_3$

treatment, both UFH-001 and T47D cells showed increases in the expression of CA IX

and CA XII, respectively, compared to controls (S7A and S7B Fig, respectively).

Treatment with compound U-F, at lower concentrations (1 μM and 10 μM) increased CA

IX and CA XII expression, in UFH-001 and T47D cells, respectively when compared to

the control. However, at higher concentrations (100 μM) a decrease in CA IX and CA XII

expression was observed (S7A Fig) and in some cases even to a lesser extent than the

control cells (S7B Fig). Similar to U-F, compound U-NO$_2$ also increased, somewhat, the

expression of CA IX and CA XII at lower drug concentrations but ultimately reduced

expression, more so with CA IX than CA XII, at concentrations of 100 μM of the

compound, when compared to the untreated controls (S7A and S7B Fig).

Effect of USB inhibitors on cell mobility

While cells grown on polystyrene plates provide useful models, studies of cells

grown on more compliant surfaces allow them to recapitulate, in part, their behavior *in*

vivo. We have taken advantage of Boyden Chambers to monitor migration and invasion

of UFH-001 and T47D cells in the presence or absence of USB compounds. The results

show that only the UFH-001 cells have the ability to migrate and invade which was

tested at 24 and 48 h, respectively (Fig 9A and 9C). All three USB compounds blocked

UFH-001 cell migration at high concentrations: U-CH$_3$ and U-NO$_2$ were the most

effective. Significant inhibition of UFH-001 cell invasion was observed with only U-F

treatment and at the highest concentration, well above their K_i values for inhibition of

activity. Not unexpectedly, we did not observe T47D cell migration and invasion in either

the absence or presence of the USBs (Fig 9B and 9D) and thus we are unable to make

a definitive statement regarding any potential effect of the sulfonamide inhibitors on

cells that do not possess these features.

Fig 9. Effects of USBs on breast cancer cell migration and invasion. Serum starved
UFH-001 and T47D cells were allowed to either migrate for 24 h or invade for 48 h
towards a chemoattractant in the presence or absence of USB compounds. Bright-field
images of migrating and invading cells are shown in Panel A (UFH-001 cells) and Panel
B (T47D cells). Data are quantified for UFH-001 and T47D cells Panel C (migration) and
Panel D (invasion). NC= negative control. Data shown are an average of duplicate
experiments ± SEM. *p < 0.05, and ***p < 0.001.

Discussion

In the current study, we investigated the binding of a class of USB-based compounds that were specifically designed to inhibit the cancer-associated isoforms, CA IX and CA XII, over the off-target CA II. We have shown, as others have previously, that residue 131 is important for isoform selectivity of these compounds towards CA IX and CA XII [66-68, 77, 83]. In addition, our model shows that serine residues 132 and 135 located within the active site of CA XII are required for USB selectivity (compared to the other CAs), towards this isoform. This then raises the question: Can we relate these structural data to the K_i values published for recombinant CAs [68]? Data in Table 1 show that compounds U-CH$_3$ and U-F exhibit K_i values for CA II that are one to two orders of magnitude greater than for CA IX or CA XII. Our structural data illustrate that these differences are caused by residue 131, which is a phenylalanine in CA II, forcing a lower energy conformation of U-CH$_3$ and U-F in the active site of CA II [66]. This differs from the more favorable conformation adopted by inhibitors in CA IX [66] and predicted in CA XII (Fig 1). U-NO$_2$ conformation is similar among the three CA isoforms, hence the similar K_i values. Therefore, we must conclude that residue 131 does not cause steric hindrance in the active site of CA II for U-NO$_2$ but clearly does create steric hindrance for the other two inhibitors. This may be related to physiochemical properties or flexibility of the tail moiety.

In the physiological setting of intact cells, inhibition of CA IX activity in UFH-001 cells by USBs was significantly more efficient than inhibition of CA XII in T47D cells (Fig 3). This was unexpected based on the relatively similar K_i values obtained using recombinant proteins (Table 1) [68]. This difference in efficacy was about an order of magnitude between the cell line that expresses CA IX versus the cell line that expresses

CA XII. Because of the difference in CA IX and CA XII as prognosticators for patient

outcome, this result suggests the potential for selective therapeutic targeting for CA IX,

in the clinical setting, despite the similar efficacy of the drugs with the purified proteins.

However, there was also a difference in K_i values between purified protein and those in

intact cells. For CA IX, this difference varied across inhibitors ranging from 3-fold (U-F)

to 500-fold higher (U-NO$_2$) (i.e., inhibition of CA IX in cells versus recombinant protein).

For CA XII, this difference was nearly three orders of magnitude for all USBs. It is

possible that part of this difference lies in the use of different techniques for analyzing

activity. Recombinant protein activity was measured using stop flow kinetics [55] while

enzymatic activity in cells was measured using MIMS, a technique originally developed

to define the catalytic mechanisms of purified CAs [75, 82], including that for CA IX [84].

That differences in the K_i values are based on technique seems unlikely because the

rate constants calculated for recombinant CA IX are similar across these techniques

[84-86]. Further, rate constants determined in isolated membranes containing either CA

IX or CA XII [27, 72] show strikingly similar values, suggesting that the membrane

environment does not influence catalytic activity. Yet, we must also consider that the

cellular environment (intact cells) adds layers of complexity relative to even isolated

membranes. The cells are in suspension during the MIMS assay, having been released

from plates using a non-enzymatic procedure. Thus, many of the cells are still in

"colonies" which means that the extracellular matrix is still intact. This layer, comprised

of glycoproteins, proteoglycans, and fibrous proteins like collagen, may provide a barrier

for incoming molecules (like USBs). Yet, the permeant sulfonamide, ethoxzolamide,

inhibits activity even inside the cell and with no time delay. In addition, the USBs are

impermeant over the course of the assay, and thus inhibitor concentration is preserved

outside of the cells.

The effect of the USB inhibitors on cell growth followed a trend similar to that

observed for activity, but is more problematic. For both the UFH-001 and T47D cells,

the inhibitors were less effective by about two orders of magnitude, based on IC_{50}

values. We initially conducted these experiments under normoxic conditions, even

though aggressive forms of cancer exist in hypoxic environments. Our rationale for

using normoxia was based on the lack of difference in the K_i values of the USB

compounds for CA IX and CA XII between normoxic and hypoxic cell culture conditions.

However, Elena et al. showed that U-F (SLC-0111) activated apoptotic and necrotic

programs only in acidified medium [87], a condition associated with hypoxia. But again,

these effects were seen at only high concentrations of inhibitor, orders of magnitude

above the K_i values that we have measured for inhibition of CA activity. Regardless, we

repeated the growth studies with cells exposed to hypoxia, but the data were identical.

Several studies have shown that these USB compounds are effective at blocking tumor

growth [68, 88, 89], so this is clearly a useful strategy for treating breast cancer patients.

So, we must ask why there is a disconnect between the loss of CA activity, particularly

that of CA IX, and the inhibition of cell growth/cytotoxicity. The activity assay is

conducted for about 6 min, while the growth, migration/invasion, and cytotoxicity

experiments are conducted for up to 96 h. Under these later conditions it is possible that

the inhibitors undergo chemical inactivation by oxidation, reduction, or cleavage

reducing their ability to block the CAs. It is difficult to measure these parameters in the

context of cell culture. It is also possible that the drugs are transported into cells over

the long term, reducing the local concentration around exofacial catalytic sites of CA IX

or CA XII. However, the studies in which we tested USB compounds, in cells where CA

IX and CA XII were ablated, showed the same inhibitory effect of sulfonamides on cell

growth. This suggests that, at very high concentrations of the USB's, they are acting

independently of their inhibition of exofacial CA activity. Indeed, others have recently

observed this as well [90, 91]. In these latter studies, investigators identified a potential

new sulfonamide target, RMB39 (RNA binding motif protein 39) that is also called

CAPERα. They demonstrated that indisulam (an aryl sulfonamide drug in phase II

clinical trials for the treatment of advanced stage solid tumors) promotes the recruitment

of RBM39 to the CUL4-DCAF15 E3 ubiquitin ligase leading to RBM39 ubiquitination and

proteasomal degradation [90, 91]. Mutations in RBM39 that prevent its recruitment

increase its stability, conferring resistance to indisulam's inhibition of cell growth and

cytotoxicity [90]. Another series of studies used a naphthalene sulfonamide to bind to

the pY705 in STAT3 (signal transducer and activator of transcription 3). This binding

inhibits STAT3 phosphorylation and dimerization, which is required for its interaction

with DNA [92, 93]. *In vivo*, this sulfonamide blocks tumor growth in a breast cancer

xenograft. With these "off target" effects of sulfonamides, we must consider that

exofacial CA activity, alone, does not regulate cell growth.

In conclusion, we have shown that specific residues within the catalytic sites of CA

IX and CA XII determine the binding affinities of the USB inhibitors. These interactions,

particularly at residue 131, provide the rationale for selective inhibition of recombinant

CA IX and CA XII over the off-target CA II. Based on K_i values, determined in the

context of cells, we conclude that USB inhibition of activity shows strong specificity for

CA IX relative to CA XII. This was surprising based on the similar K_i values between

recombinant CA IX and CA XII. Because there are now data to further prove that CA IX

acts to stabilize pH in the tumor setting, blocking its activity may alter the

microenvironment, specifically in those tumors that express CA IX over CA XII. While

we observe the same trend in cell growth inhibition, (i.e., UFH-001 cells are more

sensitive to USB inhibition than T47D cells), the inhibitor IC_{50} values increase by

another two orders of magnitude. This leads us to question the role of CA activity in cell

growth. Indeed, the presence of CA IX or CA XII is not required for USB-mediated cell

growth inhibition. Obviously, the inhibitors do block cell growth in culture and tumor

growth in xenograft and metastatic models, so their value as chemotherapeutic agents

may still be considerable. That said, the mechanism of action of sulfonamides need

further investigation.

831
832
833
834

835 # Acknowledgements

References

1.   Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. Proc Natl Acad Sci U S A. 2001;98(19):10869-74. doi: 10.1073/pnas.191367098. PubMed PMID: 11553815; PubMed Central PMCID: PMCPMC58566.

2.   Harvey JM, Clark GM, Osborne CK, Allred DC. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. J Clin Oncol. 1999;17(5):1474-81. doi: 10.1200/JCO.1999.17.5.1474. PubMed PMID: 10334533.

3.   Abe O, Abe R, Enomoto K, Kikuchi K, Koyama H, Masuda H, et al. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet. 2005;365(9472):1687-717. PubMed PMID: WOS:000229082300022.

4.   Musgrove EA, Sutherland RL. Biological determinants of endocrine resistance in breast cancer. Nat Rev Cancer. 2009;9(9):631-43. doi: 10.1038/nrc2713. PubMed PMID: 19701242.

5.   Schneider BP, Winer EP, Foulkes WD, Garber J, Perou CM, Richardson A, et al. Triple-negative breast cancer: risk factors to potential targets. Clin Cancer Res. 2008;14(24):8010-8. doi: 10.1158/1078-0432.CCR-08-1208. PubMed PMID: 19088017.

6.   Dawood S, Hu R, Homes MD, Collins LC, Schnitt SJ, Connolly J, et al. Defining breast cancer prognosis based on molecular phenotypes: results from a large cohort study. Breast Cancer Res Treat. 2011;126(1):185-92. doi: 10.1007/s10549-010-1113-7. PubMed PMID: 20711652; PubMed Central PMCID: PMCPMC3026074.

7.   Stead LA, Lash TL, Sobieraj JE, Chi DD, Westrup JL, Charlot M, et al. Triple-negative breast cancers are increased in black women regardless of age or body mass index. Breast Cancer Res. 2009;11(2):R18. doi: 10.1186/bcr2242. PubMed PMID: 19320967; PubMed Central PMCID: PMCPMC2688946.

8.   Fang JS, Gillies RD, Gatenby RA. Adaptation to hypoxia and acidosis in carcinogenesis and tumor progression. Semin Cancer Biol. 2008;18(5):330-7. doi: 10.1016/j.semcancer.2008.03.011. PubMed PMID: 18455429; PubMed Central PMCID: PMCPMC2953714.

9.   Verduzco D, Lloyd M, Xu L, Ibrahim-Hashim A, Balagurunathan Y, Gatenby RA, et al. Intermittent hypoxia selects for genotypes and phenotypes that increase survival, invasion, and therapy resistance. PLoS One. 2015;10(3):e0120958. doi: 10.1371/journal.pone.0120958. PubMed PMID: 25811878; PubMed Central PMCID: PMC4374837.

10.   Sadri N, Zhang PJ. Hypoxia-inducible factors: mediators of cancer progression; prognostic and therapeutic targets in soft tissue sarcomas. Cancers (Basel). 2013;5(2):320-33. doi: 10.3390/cancers5020320. PubMed PMID: 24216979; PubMed Central PMCID: PMC3730324.

11.   Semenza GL. Hypoxia-inducible factors: mediators of cancer progression and
targets for cancer therapy. Trends Pharmacol Sci. 2012;33(4):207-14. doi:
10.1016/j.tips.2012.01.005. PubMed PMID: 22398146; PubMed Central PMCID:
PMC3437546.
12.   Brown JM. Tumor hypoxia in cancer therapy. Methods Enzymol. 2007;435:297-
321. doi: 10.1016/S0076-6879(07)35015-5. PubMed PMID: 17998060.
13.   Chiche J, Brahimi-Horn MC, Pouyssegur J. Tumour hypoxia induces a metabolic
shift causing acidosis: a common feature in cancer. J Cell Mol Med. 2010;14(4):771-94.
doi: 10.1111/j.1582-4934.2009.00994.x. PubMed PMID: 20015196; PubMed Central
PMCID: PMC3823111.
14.   Gatenby RA, Gillies RJ. Why do cancers have high aerobic glycolysis? Nat Rev
Cancer. 2004;4(11):891-9. doi: 10.1038/nrc1478. PubMed PMID: 15516961.
15.   Neri D, Supuran CT. Interfering with pH regulation in tumours as a therapeutic
strategy. Nat Rev Drug Discov. 2011;10(10):767-77. doi: 10.1038/nrd3554. PubMed
PMID: WOS:000295396100025.
16.   Swietach P, Vaughan-Jones RD, Harris AL. Regulation of tumor pH and the role
of carbonic anhydrase 9. Cancer Metastasis Rev. 2007;26(2):299-310. doi:
10.1007/s10555-007-9064-0. PubMed PMID: 17415526.
17.   Gottlieb RA, Giesing HA, Zhu JY, Engler RL, Babior BM. Cell acidification in
apoptosis - granulocyte-colony-stimulating factor delays programmed cell-death in
neutrophils by up-regulating the vacuolar H+-Atpase. P Natl Acad Sci USA.
1995;92(13):5965-8. doi: DOI 10.1073/pnas.92.13.5965. PubMed PMID:
WOS:A1995RF05000043.
18.   Gatenby RA, Gawlinski ET, Gmitro AF, Kaylor B, Gillies RJ. Acid-mediated tumor
invasion: a multidisciplinary study. Cancer Res. 2006;66(10):5216-23. doi:
10.1158/0008-5472.CAN-05-4193. PubMed PMID: 16707446.
19.   Stubbs M, McSheehy PM, Griffiths JR, Bashford CL. Causes and consequences
of tumour acidity and implications for treatment. Mol Med Today. 2000;6(1):15-9.
PubMed PMID: 10637570.
20.   Chegwidden WR, Dodgson SJ, Spencer IM. The roles of carbonic anhydrase in
metabolism, cell growth and cancer in animals. Exs. 2000;90(90):343-63. Epub
2001/03/28. PubMed PMID: 11268523.
21.   Frost SC. Physiological functions of the alpha class of carbonic anhydrases.
Subcell Biochem. 2014;75:9-30. doi: 10.1007/978-94-007-7359-2_2. PubMed PMID:
24146372.
22.   Gilmour KM. Perspectives on carbonic anhydrase. Comp Biochem Physiol A Mol
Integr Physiol. 2010;157(3):193-7. doi: 10.1016/j.cbpa.2010.06.161. PubMed PMID:
20541618.
23.   McKenna R, Frost SC. Overview of the carbonic anhydrase family. Subcell
Biochem. 2014;75:3-5. doi: 10.1007/978-94-007-7359-2_1. PubMed PMID: 24146371.
24.   Pastorekova S, Pastorek J. Cancer-related carbonic anhydrase isozymes and
their inhibition. In: Supuran CT, Scozzafava A, Conway J, editors. Carbonic Anhydrase:
Its Inhibitors and Activators. Bocal Raton: CRC Press; 2004. p. 255-81.
25.   Supuran CT. Carbonic anhydrases--an overview. Current pharmaceutical design.
2008;14(7):603-14. Epub 2008/03/14. PubMed PMID: 18336305.
26.   Supuran CT. Structure and function of carbonic anhydrases. Biochem J.
2016;473(14):2023-32. doi: 10.1042/BCJ20160115. PubMed PMID: 27407171.

27.   Li Y, Tu C, Wang H, Silverman DN, Frost SC. Catalysis and pH control by
membrane-associated carbonic anhydrase IX in MDA-MB-231 breast cancer cells. J
Biol Chem. 2011;286(18):15789-96. doi: 10.1074/jbc.M110.188524. PubMed PMID:
21454639; PubMed Central PMCID: PMC3091188.
28.   Li Y, Wang H, Oosterwijk E, Tu C, Shiverick KT, Silverman DN, et al. Expression
and activity of carbonic anhydrase IX is associated with metabolic dysfunction in MDA-
MB-231 breast cancer cells. Cancer Invest. 2009;27(6):613-23. doi:
10.1080/07357900802653464. PubMed PMID: 19367501; PubMed Central PMCID:
PMCPMC2873695.
29.   Lock FE, McDonald PC, Lou Y, Serrano I, Chafe SC, Ostlund C, et al. Targeting
carbonic anhydrase IX depletes breast cancer stem cells within the hypoxic niche.
Oncogene. 2013;32(44):5210-9. doi: 10.1038/onc.2012.550. PubMed PMID: 23208505.
30.   Lou Y, McDonald PC, Oloumi A, Chia S, Ostlund C, Ahmadi A, et al. Targeting
tumor hypoxia: suppression of breast tumor growth and metastasis by novel carbonic
anhydrase IX inhibitors. Cancer Res. 2011;71(9):3364-76. doi: 10.1158/0008-
5472.CAN-10-4261. PubMed PMID: 21415165.
31.   Atkins M, Regan M, McDermott D, Mier J, Stanbridge E, Youmans A, et al.
Carbonic anhydrase IX expression predicts outcome of interleukin 2 therapy for renal
cancer. Clinical Cancer Res. 2005;11(10):3714-21. doi: Doi 10.1158/1078-0432.Ccr-04-
2019. PubMed PMID: WOS:000229086600016.
32.   Bui MH, Seligson D, Han KR, Pantuck AJ, Dorey FJ, Huang YD, et al. Carbonic
anhydrase IX is an independent predictor of survival in advanced renal clear cell
carcinoma: Implications for prognosis and therapy. J Urology. 2003;169(4):172-.
PubMed PMID: WOS:000181721400663.
33.   Chen LQ, Howison CM, Spier C, Stopeck AT, Malm SW, Pagel MD, et al.
Assessment of carbonic anhydrase IX expression and extracellular pH in B-cell
lymphoma cell line models. Leuk Lymphoma. 2015;56(5):1432-9. doi:
10.3109/10428194.2014.933218. PubMed PMID: WOS:000356532000053.
34.   Chiche J, Ilc K, Laferriere J, Trottier E, Dayan F, Mazure NM, et al. Hypoxia-
inducible carbonic anhydrase IX and XII promote tumor cell growth by counteracting
acidosis through the regulation of the intracellular pH. Cancer Res. 2009;69(1):358-68.
doi: 10.1158/0008-5472.Can-08-2470. PubMed PMID: WOS:000262273100044.
35.   Dorai T, Sawczuk IS, Pastorek J, Wiernik PH, Dutcher JP. The role of carbonic
anhydrase IX overexpression in kidney cancer. Eur J Cancer. 2005;41(18):2935-47. doi:
10.1016/j.ejca.2005.09.011. PubMed PMID: 16310354.
36.   Genega EM, Ghebremichael M, Najarian R, Fu Y, Wang Y, Argani P, et al.
Carbonic anhydrase IX expression in renal neoplasms: correlation with tumor type and
grade. Am J Clin Pathol. 2010;134(6):873-9. doi: 10.1309/AJCPPPR57HNJMSLZ.
PubMed PMID: 21088149; PubMed Central PMCID: PMCPMC3778911.
37.   Generali D, Fox SB, Berruti A, Brizzi MP, Campo L, Bonardi S, et al. Role of
carbonic anhydrase IX expression in prediction of the efficacy and outcome of primary
epirubicin/tamoxifen therapy for breast cancer. Endocr Relat Cancer. 2006;13(3):921-
30. doi: 10.1677/erc.1.01216. PubMed PMID: 16954440.
38.   Kivela AJ, Parkkila S, Saarnio J, Karttunen TJ, Kivela J, Parkkila AK, et al.
Expression of transmembrane carbonic anhydrase isoenzymes IX and XII in normal
human pancreas and pancreatic tumours. Histochem Cell Biol. 2000;114(3):197-204.
PubMed PMID: 11083462.

39.   Kock L, Mahner S, Choschzick M, Eulenburg C, Milde-Langosch K, Schwarz J, et
al. Serum carbonic anhydrase IX and its prognostic relevance in vulvar cancer. Int J
Gynecol Cancer. 2011;21(1):141-8. doi: 10.1097/IGC.0b013e318204c34f. PubMed
PMID: 21330838.
40.   Leibovich BC, Sheinin Y, Lohse CM, Thompson RH, Cheville JC, Zavada J, et al.
Carbonic anhydrase IX is not an independent predictor of outcome for patients with
clear cell renal cell carcinoma. J Clin Oncol. 2007;25(30):4757-64. doi:
10.1200/JCO.2007.12.1087. PubMed PMID: 17947723.
41.   Liao SY, Lerman MI, Stanbridge EJ. Expression of transmembrane carbonic
anhydrases, CAIX and CAXII, in human development. BMC Dev Biol. 2009;9:22. doi:
10.1186/1471-213X-9-22. PubMed PMID: 19291313; PubMed Central PMCID:
PMC2666674.
42.   Woelber L, Kress K, Kersten JF, Choschzick M, Kilic E, Herwig U, et al. Carbonic
anhydrase IX in tumor tissue and sera of patients with primary cervical cancer. BMC
Cancer. 2011;11:12. doi: 10.1186/1471-2407-11-12. PubMed PMID: 21223596;
PubMed Central PMCID: PMCPMC3027191.
43.   Wykoff CC, Beasley NJ, Watson PH, Turner KJ, Pastorek J, Sibtain A, et al.
Hypoxia-inducible expression of tumor-associated carbonic anhydrases. Cancer Res.
2000;60(24):7075-83. PubMed PMID: 11156414.
44.   Watson PH, Chia SK, Wykoff CC, Han C, Leek RD, Sly WS, et al. Carbonic
anhydrase XII is a marker of good prognosis in invasive breast carcinoma. Br J Cancer.
2003;88(7):1065-70. doi: 10.1038/sj.bjc.6600796. PubMed PMID: 12671706; PubMed
Central PMCID: PMC2376369.
45.   Barnett DH, Sheng S, Charn TH, Waheed A, Sly WS, Lin CY, et al. Estrogen
receptor regulation of carbonic anhydrase XII through a distal enhancer in breast
cancer. Cancer Res. 2008;68(9):3505-15. doi: 10.1158/0008-5472.CAN-07-6151.
PubMed PMID: 18451179.
46.   Creighton CJ, Cordero KE, Larios JM, Miller RS, Johnson MD, Chinnaiyan AM,
et al. Genes regulated by estrogen in breast tumor cells in vitro are similarly regulated in
vivo in tumor xenografts and human breast tumors. Genome Biol. 2006;7(4):R28. doi:
10.1186/gb-2006-7-4-r28. PubMed PMID: 16606439; PubMed Central PMCID:
PMC1557996.
47.   Deroo BJ, Korach KS. Estrogen receptors and human disease. J Clin Invest.
2006;116(3):561-70. doi: 10.1172/JCI27987. PubMed PMID: 16511588; PubMed
Central PMCID: PMC2373424.
48.   Kummola L, Hamalainen JM, Kivela J, Kivela AJ, Saarnio J, Karttunen T, et al.
Expression of a novel carbonic anhydrase, CA XIII, in normal and neoplastic colorectal
mucosa. BMC Cancer. 2005;5:41. doi: 10.1186/1471-2407-5-41. PubMed PMID:
15836783; PubMed Central PMCID: PMC1097719.
49.   Pastorek J, Pastorekova S, Callebaut I, Mornon JP, Zelnik V, Opavsky R, et al.
Cloning and characterization of MN, a human tumor-associated protein with a domain
homologous to carbonic anhydrase and a putative helix-loop-helix DNA binding
segment. Oncogene. 1994;9(10):2877-88. PubMed PMID: 8084592.
50.   Pastorekova S, Zavadova Z, Kostal M, Babusikova O, Zavada J. A novel quasi-
viral agent, MaTu, is a two-component system. Virology. 1992;187(2):620-6. PubMed
PMID: 1312272.

51.   Pastorekova S, Parkkila S, Parkkila AK, Opavsky R, Zelnik V, Saarnio J, et al.
Carbonic anhydrase IX, MN/CA IX: analysis of stomach complementary DNA sequence
and expression in human and rat alimentary tracts. Gastroenterology. 1997;112(2):398-
408. PubMed PMID: 9024293.
52.   Saarnio J, Parkkila S, Parkkila AK, Haukipuro K, Pastorekova S, Pastorek J, et
al. Immunohistochemical study of colorectal tumors for expression of a novel
transmembrane carbonic anhydrase, MN/CA IX, with potential value as a marker of cell
proliferation. Am J Pathol. 1998;153(1):279-85. doi: 10.1016/S0002-9440(10)65569-1.
PubMed PMID: 9665489; PubMed Central PMCID: PMC1852958.
53.   Potter C, Harris AL. Hypoxia inducible carbonic anhydrase IX, marker of tumour
hypoxia, survival pathway and therapy target. Cell Cycle. 2004;3(2):164-7. PubMed
PMID: 14712082.
54.   Swietach P, Wigfield S, Cobden P, Supuran CT, Harris AL, Vaughan-Jones RD.
Tumor-associated carbonic anhydrase 9 spatially coordinates intracellular pH in three-
dimensional multicellular growths. J Biol Chem. 2008;283(29):20473-83. doi:
10.1074/jbc.M801330200. PubMed PMID: 18482982.
55.   Swietach P, Wigfield S, Supuran CT, Harris AL, Vaughan-Jones RD. Cancer-
associated, hypoxia-inducible carbonic anhydrase IX facilitates CO_2 diffusion. BJU Int.
2008;101 Suppl 4:22-4. doi: 10.1111/j.1464-410X.2008.07644.x. PubMed PMID:
18430118.
56.   Stubbs M, McSheehy PM, Griffiths JR. Causes and consequences of acidic pH in
tumors: a magnetic resonance study. Adv Enzyme Regul. 1999;39:13-30. PubMed
PMID: 10470364.
57.   Lee SH, McIntyre D, Honess D, Hulikova A, Pacheco-Torres J, Cerdan S, et al.
Carbonic anhydrase IX is a pH-stat that sets an acidic tumour extracellular pH in vivo.
Br J Cancer. 2018. doi: 10.1038/s41416-018-0216-5. PubMed PMID: 30206370.
58.   Supuran C, Dedhar S, Carta F, Winum JY, Mcdonald PC. Carbonic anhydrase
inhibitors with antimetastatic activity. Google Patents; 2012.
59.   Supuran CT. Structure-based drug discovery of carbonic anhydrase inhibitors. J
Enzyme Inhib Med Chem. 2012;27(6):759-72. doi: 10.3109/14756366.2012.672983.
PubMed PMID: 22468747.
60.   Supuran CT. How many carbonic anhydrase inhibition mechanisms exist? J
Enzyme Inhib Med Chem. 2016;31(3):345-60. doi: 10.3109/14756366.2015.1122001.
PubMed PMID: 26619898.
61.   Supuran CT. Advances in structure-based drug discovery of carbonic anhydrase
inhibitors. Expert Opin Drug Discov. 2017;12(1):61-88. doi:
10.1080/17460441.2017.1253677. PubMed PMID: 27783541.
62.   Supuran CT, Winum JY. Designing carbonic anhydrase inhibitors for the
treatment of breast cancer. Expert Opin Drug Discov. 2015;10(6):591-7. doi:
10.1517/17460441.2015.1038235. PubMed PMID: 25891195.
63.   Pinard MA, Mahon B, McKenna R. Probing the surface of human carbonic
anhydrase for clues towards the design of isoform specific inhibitors. Biomed Res Int.
2015;2015:453543. doi: 10.1155/2015/453543. PubMed PMID: 25811028; PubMed
Central PMCID: PMC4355338.
1086
1087

64.   Carter MJ, Parsons DS. The purification and properties of carbonic anhydrases
from guinea-pig erythrocytes and mucosae of the gastrointestinal tract. Biochem J.
1970;120(4):797-808. PubMed PMID: 4992954; PubMed Central PMCID:
PMCPMC1179673.
65.   Lindskog S. Purification and properties of bovine erythrocyte carbonic anhydrase.
Biochimica Et Biophysica Acta. 1960;39(2):218-26. doi: Doi 10.1016/0006-
3002(60)90156-6. PubMed PMID: WOS:A1960WF20900004.
66.   Mboge MY, Mahon BP, Lamas N, Socorro L, Carta F, Supuran CT, et al.
Structure activity study of carbonic anhydrase IX: Selective inhibition with ureido-
substituted benzenesulfonamides. Eur J Med Chem. 2017;132:184-91. doi:
10.1016/j.ejmech.2017.03.026. PubMed PMID: 28363153.
67.   Pacchiano F, Aggarwal M, Avvaru BS, Robbins AH, Scozzafava A, McKenna R,
et al. Selective hydrophobic pocket binding observed within the carbonic anhydrase II
active site accommodate different 4-substituted-ureido-benzenesulfonamides and
correlate to inhibitor potency. Chem Commun (Camb). 2010;46(44):8371-3. doi:
10.1039/c0cc02707c. PubMed PMID: 20922253.
68.   Pacchiano F, Carta F, McDonald PC, Lou Y, Vullo D, Scozzafava A, et al.
Ureido-substituted benzenesulfonamides potently inhibit carbonic anhydrase IX and
show antimetastatic activity in a model of breast cancer metastasis. J Med Chem.
2011;54(6):1896-902. doi: 10.1021/jm101541x. PubMed PMID: 21361354.
69.   Debreczeni JE, Emsley P. Handling ligands with Coot. Acta Crystallogr D Biol
Crystallogr. 2012;68(Pt 4):425-30. doi: 10.1107/S0907444912000200. PubMed PMID:
22505262; PubMed Central PMCID: PMC3322601.
70.   Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta
Crystallogr D Biol Crystallogr. 2004;60(Pt 12 Pt 1):2126-32. doi:
10.1107/S0907444904019158. PubMed PMID: 15572765.
71.   Chen Z, Ai L, Mboge MY, McKenna R, Frost CJ, Heldermon CD, et al. UFH-001
cells: A novel triple negative, CAIX-positive, human breast cancer model system.
Cancer Biol Ther. 2018:1-32. doi: 10.1080/15384047.2018.1449612. PubMed PMID:
29561695.
72.   Chen Z, Ai L, Mboge MY, Tu C, McKenna R, Brown DB, et al. Differential
expression and function of CAIX and CAXII in breast cancer: A comparison between
tumorgraft models and cells. PLoS One. 2018.
73.   Laemmli UK. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. Nature. 1970;227(5259):680-5. PubMed PMID: 5432063.
74.   Silverman DN. Carbonic anhydrase: oxygen-18 exchange catalyzed by an
enzyme with rate-contributing proton-transfer steps. Methods Enzymol. 1982;87:732-52.
PubMed PMID: 6294458.
75.   Silverman DN, Lindskog S. The catalytic mechanism of carbonic anhydrase -
implication of a rate limiting protolysis of water. Acc Chem Res. 1988;21:30-6.
76.   Aggarwal M, Kondeti B, McKenna R. Insights towards sulfonamide drug
specificity in alpha-carbonic anhydrases. Bioorg Med Chem. 2013;21(6):1526-33. doi:
10.1016/j.bmc.2012.08.019. PubMed PMID: 22985956; PubMed Central PMCID:
PMC3593968.
1132
1133

77.   Pinard MA, Boone CD, Rife BD, Supuran CT, McKenna R. Structural study of
interaction between brinzolamide and dorzolamide inhibition of human carbonic
anhydrases. Bioorg Med Chem. 2013;21(22):7210-5. doi: 10.1016/j.bmc.2013.08.033.
PubMed PMID: 24090602.
78.   Fisher SZ, Aggarwal M, Kovalevsky AY, Silverman DN, McKenna R. Neutron
diffraction of acetazolamide-bound human carbonic anhydrase II reveals atomic details
of drug binding. J Am Chem Soc. 2012;134(36):14726-9. doi: 10.1021/ja3068098.
PubMed PMID: 22928733; PubMed Central PMCID: PMCPMC3524527.
79.   Soule HD, Maloney TM, Wolman SR, Peterson WD, Jr., Brenz R, McGrath CM,
et al. Isolation and characterization of a spontaneously immortalized human breast
epithelial cell line, MCF-10. Cancer Res. 1990;50(18):6075-86. PubMed PMID:
1975513.
80.   Tait L, Soule HD, Russo J. Ultrastructural and immunocytochemical
characterization of an immortalized human breast epithelial cell line, MCF-10. Cancer
Res. 1990;50(18):6087-94. PubMed PMID: 1697506.
81.   Chavez KJ, Garimella SV, Lipkowitz S. Triple negative breast cancer cell lines:
one tool in the search for better treatment of triple negative breast cancer. Breast Dis.
2010;32(1-2):35-48. doi: 10.3233/BD-2010-0307. PubMed PMID: 21778573; PubMed
Central PMCID: PMCPMC3532890.
82.   Li Y, Wang H, Tu C, Shiverick KT, Silverman DN, Frost SC. Role of hypoxia and
EGF on expression, activity, localization and phosphorylation of carbonic anhydrase IX
in MDA-MB-231 breast cancer cells. Biochim Biophys Acta. 2011;1813(1):159-67. doi:
10.1016/j.bbamcr.2010.09.018. PubMed PMID: 20920536; PubMed Central PMCID:
PMCPMC3014402.
83.   Mahon BP, Hendon AM, Driscoll JM, Rankin GM, Poulsen SA, Supuran CT, et al.
Saccharin: a lead compound for structure-based drug design of carbonic anhydrase IX
inhibitors. Bioorg Med Chem. 2015;23(4):849-54. doi: 10.1016/j.bmc.2014.12.030.
PubMed PMID: 25614109; PubMed Central PMCID: PMC4352949.
84.   Wingo T, Tu C, Laipis PJ, Silverman DN. The catalytic properties of human
carbonic anhydrase IX. Biochem Biophys Res Commun. 2001;288(3):666-9. doi:
10.1006/bbrc.2001.5824. PubMed PMID: 11676494.
85.   Alterio V, Vitale RM, Monti SM, Pedone C, Scozzafava A, Cecchi A, et al.
Carbonic anhydrase inhibitors: X-ray and molecular modeling study for the interaction of
a fluorescent antitumor sulfonamide with isozyme II and IX. J Am Chem Soc.
2006;128(25):8329-35. doi: 10.1021/ja061574s. PubMed PMID: 16787097.
86.   Vullo D, Franchi M, Gallori E, Pastorek J, Scozzafava A, Pastorekova S, et al.
Carbonic anhydrase inhibitors: inhibition of the tumor-associated isozyme IX with
aromatic and heterocyclic sulfonamides. Bioorg Med Chem Lett. 2003;13(6):1005-9.
PubMed PMID: 12643899.
87.   Andreucci E, Peppicelli S, Carta F, Brisotto G, Biscontin E, Ruzzolini J, et al.
Carbonic anhydrase IX inhibition affects viability of cancer cells adapted to extracellular
acidosis. J Mol Med (Berl). 2017;95(12):1341-53. doi: 10.1007/s00109-017-1590-9.
PubMed PMID: 28929255.
88.   Lou Y, McDonald PC, Oloumi A, Chia S, Ostlund C, Ahmadi A, et al. Targeting
tumor hypoxia: Suppression of breast tumor growth and metastasis by novel carbonic
anhydrase IX inhibitors. Cancer Res. 2011;71:3364-76.

89.   Touisni N, Maresca A, McDonald PC, Lou Y, Scozzafava A, Dedhar S, et al.
Glycosyl coumarin carbonic anhydrase IX and XII inhibitors strongly attenuate the
growth of primary breast tumors. J Med Chem. 2011;54(24):8271-7. Epub 2011/11/15.
doi: 10.1021/jm200983e. PubMed PMID: 22077347.
90.   Han T, Goralski M, Gaskill N, Capota E, Kim J, Ting TC, et al. Anticancer
sulfonamides target splicing by inducing RBM39 degradation via recruitment to
DCAF15. Science. 2017;356(6336). doi: 10.1126/science.aal3755. PubMed PMID:
28302793.
91.   Uehara T, Minoshima Y, Sagane K, Sugi NH, Mitsuhashi KO, Yamamoto N, et al.
Selective degradation of splicing factor CAPERalpha by anticancer sulfonamides. Nat
Chem Biol. 2017;13(6):675-80. doi: 10.1038/nchembio.2363. PubMed PMID: 28437394.
92.   Yu W, Li C, Zhang W, Xia Y, Li S, Lin JY, et al. Discovery of an orally selective
inhibitor of signal transducer and activator of transcription 3 using advanced multiple
ligand simultaneous docking. J Med Chem. 2017;60(7):2718-31. doi:
10.1021/acs.jmedchem.6b01489. PubMed PMID: 28245116.
93.   Yu W, Xiao H, Lin J, Li C. Discovery of novel STAT3 small molecule inhibitors via
in silico site-directed fragment-based drug design. J Med Chem. 2013;56(11):4402-12.
doi: 10.1021/jm400080c. PubMed PMID: 23651330.

1227   **Table 1.** Inhibition constant (K_i) values for recombinant human CA II, IX, and XII with ureido-
1228   substituted benzene sulfonamide (USB) compounds **U-CH₃, U-F,** and **U-NO₂** obtained from
1229   Applied Photophysics stopped-flow kinetics experiments.
1230

USB Compound	K_i (nM) ± 5% error		
	CA II	CA IX	CA XII
U-CH₃	1765	7	6
U-F	960	45	4
U-NO₂	15	1	6

1231   *K_i values were previously published by Pacchiano et al. [67, 68].*

1232

1233   **Table 2.** Resides within the active site of human CA II, CA IX and CA XII that differ
1234   among all three isoforms and make up the selective pocket (*CA II numbering*).
1235

Residues in Selective Region	CA II	CA IX	CA XII
67	Asn	Gln	Lys
91	Ile	Leu	Thr
131	Phe	Val	Ala
132	Gly	Ala	Ser
135	Val	Leu	Ser

1236   *Information adapted from Pinard et al. [63]*

1237

1238

1239

1240

1241

1242 **Table 3.** IC_{50} values for compounds **U-CH₃, U-F,** and **U-NO₂** in MCF 10A, UFH-001 and
1243 T47D cells obtained from MTT assay experiments.
1244

Cell Lines	MCF 10A Cells		UFH-001 Cells		T47D Cells	
Treatment	48 h	96 h	48 h	96 h	48 h	96 h
U-CH₃ (μM)	491 ± 10	508 ± 10	80 ± 2	74 ± 1	330 ± 10	101 ± 10
U-F (μM)	550 ± 10	550 ± 10	62 ± 1	51± 1	537± 10	252 ± 20
U-NO₂ (μM)	64 ± 1	69 ± 1	26 ± 1	20 ± 1	61 ± 1	78 ± 1

Supporting information

S1 Fig. Effect of classical sulfonamide inhibitors on CA activity in UFH-001 and T47D cells. CA activity was measured in UFH-001 cells (Panel A) and T46D cells (Panel B) using the MIMS assay in the absence or presence of acetazolamide (ACZ) or ethoxzolamide (EZA). Data are representative of two independent experiments. First order rate constants were calculated according to the formula described in the Methods. Is noted that the scale on the y-axis is different between these two representative plots. This difference represents the different isotopic enrichments of CO_2, but the concentration of CO_2 is identical between the two experiments.

S2 Fig. CA mRNA expression in breast cells lines. Panel A: mRNA expression (from microarray data) in a normal immortalized basal type breast cell line (MCF10A) compared to a triple negative breast cancer cell lline (UFH-001) and Panel B: MCF10A versus T47D cells were analyzed using data mining techniques. Accession numbers, GSE107209 (for comparison between MCF10A and UFH-001 cell lines) and NCI-60 data sets for T47D cells, were used for this comparison.

S3 Fig. Effect of U-NO$_2$ on CA activity. CA activity was measured in normoxic and hypoxic cells UFH-001 cells (Panel A) or normoxic and hypoxic T47D cells (Panel B) in the presence of U-NO$_2$ to determine K_i values across an extensive range of inhibitor concentrations.

S4 Fig. Effect of CA knockdown on spheroid growth. Western blots of lysates from UFH-001 cells (EV controls and KO cells) exposed to normoxic or hypoxic conditions (Panel A) were compared to lysates from T47D cells (EV controls and KO cells) exposed to normoxic and hypoxic conditions (Panel B). Panel C shows spheroid development of UFH-OO1 cells (EV controls and KO cells) while Panel D shows spheroid development of T47D cells (EV controls and KO cells) over 96 h in culture. GAPDH and actin were used as loading controls.

S5 Fig. Total LDH activity released by breast cell lines. Cells were grown in 96 well plates for 24 h at which point they were treated with a drug which is cytotoxic (β-caryophyllene) as a positive control or left untreated (NC) under normoxic conditions. LDH assays were performed after 48 h of treatment, results were evaluated at 450 nm (absorbance), and data was analyzed using Prism. Total LDH activity (nmol/min) ws assessed in Panel A) MCF10A cells; Panel B UFH-001 cells; and Panel C t47D cells. Data represent the mean ± SEM of 3 independent experiments.

S6 Fig. Effect of USBs on activation of apoptosis. Activation of apoptotic pathways was evaluated using the caspase activity assay in Panel A) UFH-001 and Panel B) T47D cells after 48 h of treatment with either absence (negative control, NC) or presence of USB-based compounds, under normoxic conditions. These data were compared to the presence of staurosporine (positive control, PC). Data shown for the USB-treated cells are the averages of at least three independent experiments. For the PC-treated cells, these data represent the average of two independent experiments.

S7 Fig. Effects of USB compounds on CA expression in breast cancer cells.

Immunoblotting is shown for CA IX, CA XII, and CA II from cells grown for 2-3 days and then treated with compounds U-CH3, U-F, or U-NO2 for 48 h, under normoxic conditions. GAPDH was used as a loading control. Data are representative of 3 independent experiments. Panel A, UFH-001 cells. Panel B, T47D cells.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6

Fig.7

Fig.8

Fig.9