Structural evolutionary analysis predicts functional sites in the artemisinin resistance malaria protein K13

Romain Coppée[1], Daniel C. Jeffares[2], Audrey Sabbagh[1,*], Jérôme Clain[1,3,*]

[1]UMR 216 MERIT, Institut de Recherche pour le Développement, Université Paris Descartes, Sorbonne Paris Cité, Paris, France

[2]Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

[3]Centre National de Référence du Paludisme, Hôpital Bichat-Claude Bernard, Assistance Publique des Hôpitaux de Paris, France

*__Corresponding authors:__ audrey.sabbagh@parisdescartes.fr, jerome.clain@parisdescartes.fr

Abstract

K13 is an essential Plasmodium falciparum protein that plays a key role in malaria resistance to artemisinins. Although K13 resembles BTB- and Kelch/propeller-containing proteins involved in ubiquitin ligase complexes, its functional sites remain uncharacterized. Using evolutionary and structural information, we searched for the most conserved K13 sites across Apicomplexa species evolution to identify sub-regions of K13 that are likely functional. An amino acid electropositive "patch" in the K13 propeller domain has a dense concentration of extraordinarily conserved positions located at a shallow pocket, suggesting a role as binding surface. When applied to experimentally-characterized BTB-Kelch proteins, our strategy successfully identifies the validated substrate-binding residues within their own propeller shallow pocket. Another patch of slowly evolving sites is identified in the K13 BTB domain which partially overlaps the surface that binds to Cullin proteins in BTB-Cullin complexes. We provide candidate binding sites in K13 propeller and BTB domains for functional follow-up studies.

Introduction

Current efforts to control malaria are threatened by the spread in Southeast Asia (SEA) of *Plasmodium falciparum* parasites that are resistant to artemisinin derivatives (ARTs)[1]. Treatment failures are now reported in some geographic areas of SEA for the current front-line ART-based combination therapies[1–3]. In Africa, ART resistance (ART-R) is not yet established[4,5], although some *P. falciparum* parasites from Uganda and Equatorial Guinea exhibiting high survival rate have been described [6,7].

In parasites from SEA, ART-R is primarily conferred by single non-synonymous mutations in the *P. falciparum k13* (*pfk13*) gene[8,9]. Multiple *pfk13* ART-R mutations have emerged concomitantly in the early 2000's until a specific, multidrug-resistant lineage carrying the C580Y mutation became the most common, especially in the East Thailand-Cambodia-Lao PDR-Vietnam region[4,8,10–12]. The ART-R phenotype is defined as parasites exhibiting *in vivo* a delayed clearance time following an ART-based treatment[1] and *in vitro* an increased survival rate following a brief exposure to a high dose of ART[13].

The *pfk13* gene encodes a 726 amino acid protein (PfK13) which is essential at least during the intraerythrocytic parasite blood stage[14,15]. Both the gene ontology annotation of *pfk13* and the study of ART-resistant parasites carrying a *pfk13* mutation suggest that this protein has some regulatory functions at the protein level[16,17]. At the cell level, *pfk13* mutant parasites decelerate their development during the early intraerythrocytic stage[13,18]. At the molecular level, they exhibit an increased expression of unfolded protein response pathways[19], lower levels of ubiquitinated proteins[17,18], and phosphorylation of the parasite eukaryotic initiation factor-2α (eIF2α) which correlates with ART-induced latency[20]. There are some indications of the interactors with PfK13 that partially clarify its function. For example, PfK13 was immunoprecipited with the phosphatidylinositol 3-kinase (PI3K)[17]. Ubiquitination of PI3K is also decreased in *pfk13* C580Y mutants, resulting in wide phosphatidylinositol 3-phosphate-related cellular alterations[17,21] which roles in ART-R are still to be characterized.

PfK13 is related to the BTB-Kelch structural subgroup of Kelch-repeat proteins[8,22,23]. It possesses a BTB domain (Broad-complex, tramtrack and bric-à-brac; also known as BTB/POZ; amino acids 350-

437) and a C-terminal propeller domain (also known as Kelch domain; amino acids 443-726)

composed of six repeated Kelch motifs (PDB code 4YY8, unpublished). However, PfK13 exhibits

specific features when compared to typical BTB-Kelch proteins, such as a poorly conserved

Apicomplexa-specific N-terminal region, a coiled-coil-containing (CCC) domain (amino acids 212-

341) located upstream of BTB, and an absence of BACK domain (for BTB And C-terminal Kelch)

often found between BTB and Kelch domains[23–25]. Both BTB and propeller are protein domains

known to carry binding functions. Nearly all *pfk13* mutations associated with ART-R – including the

C580Y mutation – cluster in the propeller domain[8,11], suggesting a major functional role of this

domain.

Many proteins harboring a BTB domain are found in multi-subunit Cullin-RING E3 ligase

complexes in which a substrate protein will be ubiquitinated and then degraded by the proteasome[24,26–

28]. In those complexes, BTB mediates varying oligomerization architectures and also contributes to

Cullin recruitment. The propeller domain often serves as the substrate protein receptor in BTB-Kelch

and other proteins[23,24]. It usually comprises four to six repeated β-stranded Kelch motifs – also named

blades – arranged in a circular symmetry (the six blades in PfK13 are named I to VI). The loops

protruding at the bottom face of propeller form a shallow pocket involved in the binding of the

substrate protein subsequently ubiquitinated and targeted for degradation[24]. For example, the propeller

shallow pocket of BTB-Kelch KEAP1 and KLHL3 directly binds to the transcription factor Nrf2 and

the kinase WNK, respectively, and controls their ubiquitination[29,30]. PfK13 may exhibit similar

functions, however, its binding regions and functionally important sites remain poorly characterized.

Here, we hypothesize that functionally important sites of K13 have evolved under stronger

purifying selection and could be identified by the analysis of *k13* molecular evolution across 43

Apicomplexa species. To examine this, we inferred substitution rates for each amino acid site of the

K13 sequence, taking into account the *k13* phylogeny and the spatial correlation of site-specific

substitution rates in the protein tertiary structure when possible. We identified a major functional patch

of slowly evolving sites located at the bottom face of K13 propeller that form part of the shallow

pocket. To show the relevance of our approach, we applied it to the propeller domain of four well-

characterized BTB-Kelch proteins (KEAP1, KLHL2, KLHL3 and KLHL12) and successfully

identified the functionally and structurally validated substrate-binding residues located in the pocket of

these propeller domains. Another patch of slowly evolving sites was also identified in the BTB domain

of K13 which partially overlaps with the surface that binds to Cullin proteins in known BTB-Cullin

complexes[27]. Altogether, these findings support a crucial role of K13 in binding partner molecules and

predict specific binding sites.

Results

K13 sequence sampling, multiple alignment and phylogenetic reconstruction. Forty-three

complete amino acid sequences from distinct *Apicomplexa* species were unambiguously identified as

orthologous to PfK13 in sequence databases, encompassing 21 *Plasmodium* and 22 other *Apicomplexa*

K13 sequences (*Cryptosporidia*, n = 7; *Piroplasmida*, n = 7; and *Coccidia*, n = 8). The length of K13

protein sequences ranged from 506 (*Eimeria brunetti*) to 820 (*Hammondia hammondi*) amino acids

(Supplementary Table 1). By visual inspection, the three annotated domains of K13 (CCC, BTB and

propeller) were conserved (see the K13 sequence alignment in Supplementary Fig. 1), whereas the N-

terminal region preceding the CCC domain appeared much more variable among sequences, even

being absent in some of them. Since K13 sequences aligned poorly in that region, the first 234 amino

acid positions of the K13 multiple alignment were removed, along with other positions showing a high

divergence level and/or a gap enrichment among sequences (32 amino acid positions; Supplementary

Fig. 1). The final K13 multiple sequence alignment contained 514 amino acid positions which covered

the whole CCC, BTB and propeller domains. The average pairwise sequence identity in that cleaned

K13 sequence alignment was 64.6%, ranging from 48.6% for the *Babesia bigemina-Cryptosporidium*

ubiquitum comparison to 99.2% for the *P. chabaudi* spp. pair.

The maximum-likelihood phylogenetic tree built from the curated alignment of the corresponding

Apicomplexa k13 cDNA sequences revealed four monophyletic groups: *Cryptosporidia, Plasmodium,*

Piroplasmida and *Coccidia*, all supported by high bootstrap values (≥ 98%; Supplementary Fig. 2).

The group of *Hematozoa k13* sequences appeared as paraphyletic (bootstrap value = 100%), with

Piroplasmida unexpectedly clustering with *Coccidia* (Supplementary Fig. 2). The phylogenetic

relationships of *Plasmodium k13* sequences were largely consistent with the acknowledged phylogeny

of *Plasmodium* species, except for bird-infecting *Plasmodium k13* sequences (*P. gallinaceum* and *P.*

relictum), which appeared related to human-infecting *P. ovale* spp. sequences, although this grouping

was poorly supported (bootstrap value = 47%; Supplementary Fig. 2).

The *k13* sequence has evolved under strong purifying selection. To evaluate the selective pressure

acting on *k13*, we used codon substitution models to estimate the rate of non-synonymous to

synonymous substitutions, $\omega = d_N/d_S$, across codon sites of the *k13* sequence (site models) and

branches of the *k13* phylogeny (branch models). A series of nested likelihood ratio tests (LRTs) using

different sets of site and branch models were carried out using the codeml tool from the PAML

package[31]. When applied to the *k13* codon alignment, LRTs of codon and branch models indicated

varying substitution rates ω both among codon sites of the *k13* sequence (M0-M3 comparison, $p = 3.3$

$\times 10^{-225}$) and among branches of the *k13* phylogeny (M0-FR, $p = 1.9 \times 10^{-53}$; Table 1 and

Supplementary Table 2). This suggests that *k13* has evolved under a variable selective regime both

across codon sites and lineages. No evidence of positive selection was found in any of the tree

branches (Supplementary Fig. 3). Similarly, site models incorporating positive selection (M2a and

M8) provided no better fit to the data than those including only purifying selection and neutral

evolution (M1a and M7: $2\Delta\ln L = 0$ in both cases; Table 1), thus supporting an absence of detectable

adaptive selection events at any *k13* codon site over the long time scale of *Apicomplexa* evolution.

Altogether, the data indicate that much of the K13 protein, except the N-terminal region, has been

strongly conserved over evolutionary time.

When considering the one-ratio PAML values of 3,256 protein-coding genes previously estimated

by Jeffares and colleagues using six *Plasmodium* species[32], *k13* ranked among the 5% most conserved

protein-coding genes of the *Plasmodium* proteome (rank: 140/3,256; Supplementary Fig. 4a). Since a

significant correlation between protein length and one-ratio PAML value was evidenced in the whole

dataset (Spearman's rank correlation: $p = 9.2 \times 10^{-82}$, r = 0.33; Supplementary Fig. 4b), we repeated

the analysis by considering only those protein sequences whose length was included in an interval of ±

100 amino acids centered on the PfK13 protein length (Spearman's rank correlation: $p = 0.83$, r =

0.01). Again, *k13* ranked among the most conserved protein-coding genes of the *Plasmodium*

proteome (sized-rank: 6/393), whereas four other five- or six-bladed Kelch protein-coding sequences

showed much less intense levels of conservation than K13 (Fig. 1).

Variable levels of amino acid conservation between the annotated domains of K13. We next

compared the conservation level between the annotated domains of K13 (CCC, BTB and propeller)

using ω estimates obtained under the best fitted PAML model that indicates a variable selective

regime among sites (model M3; Supplementary Table 2). First, we noted that the three domains have

evolved under strong purifying selection with most sites being highly conserved during evolution (Fig.

2). BTB was however found to evolve under more intense purifying selection than either CCC ($p = 1.6$

$\times 10^{-4}$, Mann-Whitney U test) or propeller ($p = 1.0 \times 10^{-3}$, Mann-Whitney U test), but no difference in

ω estimates was detected between CCC and propeller ($p = 0.75$, Mann-Whitney U test; Fig. 2 and

Supplementary Table 3). To confirm these results, we inferred the site-specific substitution rate at the

protein level using the FuncPatch server which takes into account the spatial correlation of site-

specific substitution rates in the protein tertiary structure[33] (hereafter called λ substitution rate). λ could

not be inferred for the CCC domain (because of the lack of a resolved 3D structure), but the analysis

confirmed that BTB was more conserved than propeller over evolutionary time ($p = 5.4 \times 10^{-5}$, Mann

Whitney U test; Supplementary Table 4).

The BTB domain of K13 resembles that of KCTD proteins and exhibits a predicted functional

patch. We then performed a more extensive study of the BTB and propeller domains of K13 because

of their likely role in mediating K13 functions and the availability of their tertiary structures. To detect

patches of slowly evolving amino acid sites in the BTB-propeller structure, we focused on the site-

specific, spatially correlated, substitution rate λ at the amino acid level. This rate has been shown to

provide a more reliable estimation of the conservation level at amino acid sites compared to standard

substitution estimates, especially in the case of highly conserved proteins[33,34].

Although K13 is related to the BTB-Kelch structural subgroup of proteins[8], the BTB domain of

K13 exhibits atypical features compared to BTB-Kelch proteins. First, the K13 BTB fold appeared

shortened, lacking the A6 helix and the N- and C-terminal extensions[27], similar to Elongin C (Fig. 3a).

Second, the primary sequence of K13 BTB grouped with those of the KCTD protein family rather than

of other BTB-containing protein families (Fig. 3b). Finally, K13 BTB exhibited a higher similarity in

tertiary structure with the BTB domain of KCTD17 compared to those of Elongin C and KEAP1: the

root-mean-square deviations (RMSDs) of atomic positions for BTB domains of K13-KCTD17, K13-

Elongin C and K13-KEAP1 were 1.13 Ångström (Å), 2.33 Å and 2.17 Å, respectively.

To identify putative functional sites in K13 BTB, we examined whether a spatial correlation of the

site-specific substitution rates λ is present in the K13 BTB-propeller tertiary structure. Despite a low

standard deviation of substitution rates across amino acid sites, a significant spatial correlation was

found with a log Bayes factor drastically > 8 and a 5 Å characteristic length (Table 2). The 10% most

conserved sites predicted by FuncPatch formed one clearly bounded patch located at the surface of

BTB (Fig. 3c). The BTB patch contained sites located at both B2-B3 and B4-A4 loops and at the A4

helix (positions/residues 355-358/NVGG, 397-399/DRD and 402-403/LF using the PfK13 sequence

numbering; Fig. 3c). To test whether a similarly located, conserved patch was also found in the BTB

domain of KCTD proteins, to which K13 BTB is the most similar, we inferred site-specific

substitution rates λ from 124 and 139 orthologous sequences of SHKBP1 and KCTD17, respectively.

For both proteins, the 10% most conserved BTB positions formed a patch that partially overlapped

with the one of K13, with positions 357 (B2-B3 loop), 397 and 398 (B4-A4 loop) being shared

between the three patches (PfK13 sequence numbering; Fig. 3d and Table 2). These positions are

usually involved in BTB-BTB interactions in KCTDs[28] or in BTB-Cullin interactions in some other

BTB-containing protein families[27], as in the X-ray structure of the Elongin C-Cullin2 complex

(Supplementary Fig. 5). By functional annotation transfer, this indicates that these K13 BTB sites may

be implied in some protein-protein interactions.

The propeller domain of K13 exhibits a conserved shallow pocket. In Kelch-containing proteins,

the propeller domain usually serves as the receptor for a substrate further ubiquitinated[23,24]. Before

examining its conservation level, we first re-evaluated the architecture of the K13 propeller domain

using its resolved 3D structure (PDB code 4YY8, chain A, unpublished). The PfK13 propeller

structure is composed of six repeats of the Kelch motif (or blade)[8,22]. As expected, each blade is a β-

sheet secondary structure involving four twisted antiparallel β-strands (numbered A to D). The

innermost A strands line the central channel of the propeller fold whereas the outermost D strands are

part of the propeller outside surface. The top face of the domain, containing the central channel

opening, is formed by residues from several strands and AB and CD loops. The bottom face of

propeller is composed of residues from the DA and BC loops and contains a shallow pocket, similar to

other propeller structures[24]. Since there is no conventional definition for the shallow pocket

delineation in the propeller fold, we defined it as the amino acids forming the surface plan of the

pocket and protruding out of the plan (n = 19 positions; Fig. 4a and Supplementary Fig. 6).

To characterize the pattern of conservation within the domain, we superimposed the site-specific

substitution rates λ inferred by FuncPatch onto an amino acid sequence alignment of the six blades

(Fig. 4b), custom-produced from a structural alignment of the blades using PyMOL (Fig. 4a and

Supplementary Fig. 7). We found that: *i)* the conservation level significantly differed between the six

blades of propeller ($p = 3.0 \times 10^{-3}$, Kruskal-Wallis *H* test), with blade VI exhibiting the lowest

conservation (Fig. 4b, Supplementary Fig. 8 and Supplementary Table 5); *ii)* loops were more

conserved than strands ($p = 5.0 \times 10^{-3}$, Mann-Whitney *U* test; Fig. 4b and Supplementary Table 5); *iii)*

the solvent-exposed A and D strands were less conserved than the buried B and C strands ($p = 1.3 \times$

10^{-6}, Kruskal-Wallis *H* test; Fig. 4b and Supplementary Table 5); and *iv)* the conservation level was

the strongest at the blade positions 7-10 (DA loops) and 23-25 (BC loops; Fig. 4b), which altogether

formed the surface and underlying layer of the shallow pocket in the PfK13 propeller tertiary structure

(Fig. 4a). Of note, similar results were obtained with the ω estimates inferred under the best fitted

PAML model (Supplementary Table 5).

With one exception (position PfK13 514), the 10% most conserved sites in K13 propeller (n = 29

out of 284) were all located at the bottom side of the propeller fold (Fig. 5a). Among them, the sites

exposed at the surface (n = 9) formed a statistically significant patch located within the shallow pocket

(Table 2 and Fig. 5a). The nineteen positions forming the shallow pocket of K13 propeller were

significantly enriched in the 10% most conserved propeller amino acid sites ($p = 1.6 \times 10^{-5}$, chi-

squared test; Fig. 5a, Tables 3 and 4). A similar trend was observed using the ω PAML values

(Supplementary Fig. 9).

Using the PfK13 propeller structure as reference, we also identified two remarkable features of this

conserved patch. First, it overlapped with a region of the shallow pocket harboring an electropositive

surface potential energy, in contrast to the overall electronegative one of the propeller bottom surface

(Fig. 5b). Second, it contained several arginine and serine residues strictly conserved in *Apicomplexa*

(PfK13 R529, S576, S577, R597 and S623; Fig. 5a and Table 3), which are known to mediate protein-

protein interactions in the pocket of other propeller domains[35,36]. Altogether, our evolutionary analysis

of K13 propeller revealed that the shallow pocket is extremely conserved and may bind a substrate

molecule.

The conserved propeller patch predicted by FuncPatch is related to propeller binding activities

in well-characterized BTB-Kelch proteins. To evaluate the reliability of FuncPatch to infer

conserved functional sites in the context of propeller domains, we studied four other BTB-Kelch

proteins found in mammals and which are functionally and structurally well-characterized (KEAP1,

KLHL2, KLHL3 and KLHL12; Supplementary Fig. 10). All these proteins are known to bind a

substrate protein through validated binding residues located in their propeller shallow pocket

(substrates: Nrf2 for KEAP1[37], WNK for KLHL2 and KLHL3[35], and Dishevelled for KLHL12[38]).

Using large sets of orthologous amino acid sequences (ranging from 129 sequences for KLHL12 to

162 sequences for KLHL2), a statistically significant spatial correlation of site-specific substitution

rates λ was detected for each propeller fold (Table 2). In each case, the 10% most conserved propeller

positions clustered in the shallow pocket (highest $p = 1.4 \times 10^{-7}$ for KEAP1, chi-squared test; Fig. 5c

and Table 4). The shallow pocket of KEAP1, KLHL2 and KLHL3 propeller structures also showed a

markedly electropositive surface potential energy while the one of KLHL12 was much more variable

(Fig. 5c).

Therefore, the conserved functional sites predicted by FuncPatch in propeller domains from several

BTB-Kelch proteins are consistent with the findings of previous experimental studies, demonstrating

the reliability of our approach.

Lack of association of ART-R mutations with some long-term evolutionary or structural facets

of K13 propeller. Numerous non-synonymous mutations in PfK13 propeller segregating at variable

frequencies have been reported to confer ART-R to *P. falciparum* parasites from SEA[4,8–11]. Here, we

hypothesized that they may be associated with specific patterns of evolutionary- and/or structure-based

parameters despite their large distribution across the propeller fold (Supplementary Fig 11). K13

propeller positions were classified as associated (n = 27) or not (n = 257) with an ART-R mutation, on

the basis of the last World Health Organization (WHO) status report on ART-R[5]. No difference in the

inter-species site-specific substitution rates λ was observed between the two groups ($p = 0.96$, Mann-

Whitney U test; Supplementary Fig. 12). Importantly, we noted that no ART-R mutation has been

reported at the positions located at the surface of the shallow pocket, although this trend was not

statistically confirmed ($p = 0.23$, Fisher's exact test; 0/19 ART-R mutations for the shallow pocket

positions, and 27/265 ART-R mutations for the remaining K13 propeller positions). Two structure-

based parameters associated with site evolutionary rate were also estimated: the relative solvent

accessibility or RSA, and the side-chain weighted contact number or WCN_{sc}[39]. Again, no association

was found between these structural parameters and the propeller positions associated with ART-R

mutations ($p = 0.44$ and $p = 0.46$, respectively, Mann-Whitney U test; Supplementary Fig. 12). In

conclusion, none of the evolutionary- and structure-based parameters tested here were associated with

ART-R K13 propeller positions.

Discussion

Combining evolutionary and tertiary structure information provides a powerful and efficient way to

gain insight into the functionality of protein sites[39]. The favored strategy usually screens sites that have

evolved more rapidly than expected under a neutral model and interprets them as a signature of

adaptive evolution corresponding to a gain of new function(s)[40,41]. Here, we focused on the most

slowly evolving, 3D-bounded sites to identify highly conserved sub-regions of K13 across species

evolution that are likely to play a functional role. Because of the extreme conservation of the *k13*

gene[10] and its essential function[14,15,42,43], we made inter- rather than intra-species comparisons. In the

context of a sustained and intense purifying selection operating in all annotated domains of K13, our

analysis of sequence evolution coupled to BTB-propeller tertiary structure identified two patches of

particularly slowly evolving sites.

The most striking conserved patch is found in the K13 propeller domain. It includes specific

positions from the inter-blade DA and intra-blade BC loops, within a shallow pocket on the bottom

face of the propeller structure. Several lines of evidence suggest that this patch contains functional

sites. First, the shallow pocket is highly enriched in conserved positions, whereas solvent-exposed

sites in proteins are usually less conserved than buried ones[39]. Second, the shallow pocket of well-

characterized BTB-Kelch proteins directly mediates propeller's binding activities (KEAP1, KLHL2,

KLHL3, KLHL12)[24,35,36] and are also predicted as conserved patches in our analyses. Finally, the

conserved patches at the shallow pocket of BTB-Kelch proteins and K13 share interesting properties:

they display a markedly electropositive surface potential energy and they are enriched in arginine and

serine residues (PfK13 R529, R597, S576, S577 and S623, all strictly conserved in *Apicomplexa*;

Table 3). In KEAP1, KLHL2 and KLHL3, the corresponding residues bind to acidic peptides derived

from their substrates, Nrf2 and WNK, respectively[35,36] (Fig. 5c). According to the electropositive

region surrounding the K13 propeller conserved patch, the K13 substrate molecule(s) may harbor an

acidic binding motif. Altogether, these results indicate that the shallow pocket of K13 propeller

exhibits several properties of a binding surface, and we speculate that it may be critical for the

recognition of substrate molecule(s).

In *P. falciparum*, PI3K is a likely candidate as it is immunoprecipitated with full-length PfK13, and

its ubiquitination and proteasomal degradation are altered by the *pfk13* propeller C580Y mutation[17].

Another candidate may be the PK4 kinase which phosphorylates eIF2α, a key mediator of translation-

mediated latency involved in ART-R[20].

However, whether the *pfk13*-mediated ART-R mechanism in *P. falciparum* is related to the

predicted binding activity of the propeller shallow pocket remains elusive. Based on the last WHO

status report on ART-R[5], none of the 27 validated or candidate ART-R mutations is located at the

surface of the propeller shallow pocket. Several candidates are found in its underlayer (Supplementary

Fig. 11) and some others have a preferential localization at positions proximal to the A and B

strands[10]. Polymorphisms which have an uncharacterized phenotype were reported at 9 out of the 19

positions forming the shallow pocket, but found in only one or two parasite samples from population

surveys (Table 3). Therefore, we speculate that amino acid changes at the positions contributing

directly to the pocket surface are too damaging for the K13 native function to provide a long-term

competitive advantage. In this context, rather than directly altering the binding residues, ART-R

mutations may either induce long-range conformational changes propagating to the pocket surface or

decrease the propeller domain stability, as reported for pathogenic mutations in KEAP1 W544C and

KLHL3 S410L, respectively[44,45]. Furthermore, putative propeller conformational changes associated

with ART-R mutations might be induced by specific cellular conditions, for example increased

oxidative stress[46].

BTB appeared here as the most conserved domain of K13 during *Apicomplexa* evolution, and

therefore likely carries critical activities. It most resembles the BTB of the KCTD protein family in

primary sequence, tertiary structure and short domain size. The shortened BTB of KCTDs could still

mediate protein oligomerization[28,47], consistent with the dimer observed in the solved PfK13 BTB-

propeller crystal structures (PDB codes 4YY8 and 4ZGC, unpublished). K13 BTB harbors a predicted,

functional patch – located at the B2-B3 and B4-A4 loops and at the A4 helix – that overlaps with the

one of KCTD proteins, suggesting that they share similar functional sites. In KCTDs, these sites make

BTB-BTB contacts in tetrameric or pentameric assemblies when BTB is solved as an isolated

domain[28]. However, the PfK13 BTB-propeller structure forms a dimer and none of the highly

conserved positions make BTB-BTB contacts (PDB codes 4YY8 and 4ZGC, unpublished). Finally,

amino acids of the B4-A4 loop (corresponding to positions 397-399 in PfK13) are exposed at the

BTB-Cullin binding interface in several solved complexes[27]. These discrepancies in the role of the

predicted BTB patch could be due to the fact that additional domains (such as propeller) or partner

proteins might constraint the folding of the BTB domain into oligomers or complexes. Altogether, data

from the literature however support that the BTB predicted patch of K13 mediates protein-protein

interactions, possibly with a Cullin protein.

Interestingly, K13 also contains a highly conserved CCC domain, located before BTB, which

therefore likely carries critical activities. Consistent with this hypothesis, three *pfk13* mutations

conferring a moderate ART-R are located in CCC (PfK13 R239Q, E252Q and D281V)[3,11,48]. Coiled-

coils are ubiquitous protein-protein interaction domains composed of two or more α-helices coiling

together[49]. A CCC domain was reported in a few Kelch-containing proteins (including some KCTDs)

involved in cell morphogenesis but these CCC have a different domain organization than the one of

K13[22]. The CCC of K13 may participate in K13 oligomerization and/or serve as a binding interface

with other proteins[50].

In conclusion, through evolutionary and structural analyses, we identified the shallow pocket of the

K13 propeller domain as a candidate surface for binding a substrate molecule. We also detected in the

BTB domain of K13 a conserved patch of sites that are involved in protein-protein interactions in

known BTB-Cullin and BTB-BTB complexes. Efforts should now focus on the identification of

molecule(s) that bind(s) to the pocket of K13 propeller, which may help clarify the link between K13

function and ART-R.

Materials and Methods

Collection of *k13* orthologous sequences from genomic databases. The amino acid sequence of

PfK13 (PlasmoDB code PF3D7_1343700) was queried against the specialized eukaryotic pathogen

database (EuPathDB release 33)[51] and the NCBI non-redundant protein database using blastp and

tblastn searches[52] (BLOSUM62 matrix). A protein was considered as a likely orthologous sequence if

the sequence identity was ≥ 30% and the e-value below the 10^{-3} cutoff. Forty three K13 sequences –

and corresponding *k13* cDNA sequences – were retrieved from distinct *Apicomplexa* species including

21 *Plasmodium* species. A detailed bioinformatics analysis was performed on each protein sequence to

confirm the presence of the three annotated domains of K13 (CCC, BTB and propeller) using

InterPro[53].

k13 **sequence alignment.** Considering the greater divergence of coding nucleotide sequences as

compared to protein sequences due to the genetic code redundancy, a K13 protein sequence alignment

was first generated using Mafft version 7[54] (E-INS-I strategy with BLOSUM62 scoring matrix, gap

opening penalty 2.0 and offset 0.1). The output alignment was visually inspected and manually edited

with BioEdit v7.2.5[55]. The positions containing gaps in at least 30% of all sequences were removed, as

suggested by PAML' authors[31]. Then, the *k13* nucleotide sequence alignment was generated with

PAL2NAL[56] using the cleaned K13 amino acid alignment as template.

Phylogenetic analysis of *k13*.   The phylogenetic relationships of *k13* nucleotide sequences were

inferred using the maximum-likelihood method implemented in PhyML v3.0[57], after determining the

best-fitting nucleotide substitution model using the Smart Model Selection (SMS) package[58]. A

general time-reversible model with optimized equilibrium frequencies, gamma distributed among-site

rate variation and estimated proportion of invariable sites (GTR + G + I) was used, as selected by the

Akaike Information Criterion. The nearest neighbor interchange approach was chosen for tree

improving, and branch supports were estimated using the approximate likelihood ratio aLRT SH-like

method[59]. The *k13* phylogeny was rooted using *Cryptosporidia* species as outgroup.

Molecular evolutionary analysis of *k13*. To investigate the evolutionary regime that has shaped the

k13 protein-coding DNA sequence during species evolution, we analyzed the non-synonymous (d_N) to

synonymous (d_S) substitution rate ratio ω (= d_N/d_S), estimated by maximum-likelihood using the

codeml tool from PAML v.4.8[31,61]. ω provides a sensitive measure of selective pressure at the amino

acid level by comparing substitution rates with statistical distribution and considering the phylogenetic

tree topology. Typically, $\omega < 1$ indicates purifying selection, while $\omega = 1$ and $\omega > 1$ indicate neutral

evolution and positive selection, respectively.

The heterogeneity of ω among lineages of the *k13* phylogenetic tree (branch models) was tested by

comparing the free-ratio (FR) model, which assumes as many ω parameters as the number of branches

in the tree, to the one-ratio (M0) model which supposes only one ω value for all branches[62,63].

The variation of ω among codon sites was then evaluated using codon models M1a, M2a, M3, M7 and

M8[62,63]. M1a allows codon sites to fall into two site classes, either with $\omega < 1$ (purifying selection) or

$\omega = 1$ (neutral evolution), whereas model M2a extends model M1a with a further site class as $\omega > 1$

(positive selection). Model M3 includes a discrete distribution of independent ω with k classes of sites

($k = [3, 4, 5]$ in this study), with ω values and corresponding proportions estimated from the dataset.

Model M7 assumed a β-distribution of ten ω ratios limited to the interval [0, 1] with two shape

parameters p and q, whereas model M8 adds an additional site class with ω possibly > 1 as M2a does.

The heterogeneity of ω across codon sites was tested by comparison of models M0-M3, while

comparison of paired models M1a-M2a and M7-M8 allowed to detect positive selection[62].

Model comparisons were made using likelihood ratio tests (LRTs)[64]. For each of the LRTs, twice

the log-likelihood difference between alternative and null models ($2\Delta\ln L$) was compared to critical

values from a chi-squared distribution with degrees of freedom equal to the difference in the number

of estimated parameters between both models[40]. Candidate sites for positive selection were pinpointed

using the Bayes empirical Bayes (BEB) inference which calculates the posterior probability that each

codon site falls into a site class affected by positive selection (in models M2a and M8), as described

by Yang and colleagues[65]. For model M3, in which no BEB approach is implemented yet, the naïve

empirical Bayes (NEB) approach was used to identify those sites evolving under positive selection.

Three codon substitution models were used and compared for all models: F1x4 and F3x4, which

assume equal nucleotide frequencies and individual codon frequencies at all codon positions,

respectively, and the parameter-rich model F61, which estimates codon frequencies separately for each

codon[61,66]. Since the three codon substitution models yielded similar results (Supplementary Fig. 13),

we only presented those obtained with the most widely used F3x4 codon model. The analyses were

run multiple times with different ω starting values to check the consistency of the results.

For PAML model M3 with k site classes of ω ratios, the posterior mean of ω value at each codon

site was calculated as the average of the ω ratios across the k ω site classes weighted by their posterior

probabilities[31].

In addition to *k13*, four other Kelch protein-coding sequences were considered to compare their ω

with those estimated for the whole *Plasmodium* proteome. We used the ω values previously estimated

by Jeffares and colleagues[32] with PAML under the one-ratio model for each of the 3,256 orthologous

protein-coding genes among six *Plasmodium* species: *P. falciparum*, *P. berghei*, *P. chabaudi*, *P. vivax*,

P. yoelii and *P. knowlesi*. A full description of the procedure is presented in the original paper[32].

Inferring site-specific substitution rates considering their spatial correlation in the K13 BTB-

propeller tertiary structure. Most methods – including PAML – assume that site-specific

substitution rates are independently distributed across sites[67]. However, it is widely acknowledged that

amino acids located close to each other in protein tertiary structures are more likely to carry out

similar functions, suggesting a site interdependence in amino acid sequence evolution attributed to

tertiary structure[67,68]. Consequently, the substitution rate at the protein level (named λ in this study)

was inferred using the FuncPatch server[33]. FuncPatch requires an amino acid sequence alignment, a

phylogenetic tree and a protein tertiary structure to estimate the conservation level during species

evolution and the characteristic scale (in Å) of spatially correlated site-specific substitution rates λ. We

used the X-ray structure at 1.81 Å resolution of PfK13 BTB-propeller as the reference structure which

does not contain the conserved CCC domain (PDB code 4YY8, unpublished). Beforehand, a

Ramachandran analysis was performed to validate the quality of the structure using MolProbity[69]:

96.9% and 3.1% of the amino acids were in favored and allowed regions, respectively, and there were

no outliers. FuncPatch only accepts monomeric proteins as input whereas BTB-propeller of PfK13

dimerizes in crystal structure. To take into account the dimeric organization of PfK13, its tertiary

structure was edited using customized python scripts (Python v2.7.13) in order to merge the two

monomers (chains A and B) and the K13 sequence was duplicated in the K13 protein sequence

alignment. The analysis was also done using either one of the other monomeric BTB-propeller tertiary

structure and also using a disulfide-bonded version of PfK13 BTB-propeller (PDB code 4ZGC,

unpublished). All these control analyses yielded similar results (data not shown). The spatial

correlation of the site-specific substitution rates λ in the K13 tertiary structure was tested using a

Bayesian model comparison, where a null model (model 0), in which no spatial correlation of site-

specific substitution rates λ is present, is compared to the alternative model (model 1). As suggested by

FuncPatch' authors, the spatial correlation was considered as significant if the estimated log Bayes

factor (model 1 versus model 0) was larger than 8 in the dataset (conservative cutoff)[33].

Delineation of K13 propeller blades and secondary structures. The propeller domain of PfK13 is

composed of six blades having slightly different amino acid lengths. To get an accurate blade

alignment at the primary amino acid sequence level, we first sought to align the six blade structures.

The PDB propeller structure was obtained from the PfK13 BTB-propeller structure (PDB code 4YY8,

chain A, unpublished) and was then divided into six parts, each one containing the atomic coordinates

of one blade. The six blade structures were then aligned by minimizing the RMSD of atomic positions

using the *align* function in PyMOL Molecular Graphics System[70] so as to identify the amino acids

from the six blades that are located at exactly the same blade position. This structure alignment was

then used to align the six blades at the primary amino acid sequence level. The delineation of the

strands and loops was obtained directly from the PDB file (PDB code 4YY8, chain A, unpublished).

Definition of ART-R mutations. We used the last status report on ART-R provided by the WHO to

classify the positions of the PfK13 propeller domain as associated or not with an ART-R mutation[5].

Evolutionary analysis of the BTB and propeller domains in other BTB- and Kelch-containing

proteins. To characterize the BTB domain of K13, we arbitrarily retrieved some members belonging

to the main BTB-containing protein families (BTB-ZF, BTB-Kelch, RhoBTB, BTB-NPH3, MATH-

BTB, KCTD, KCNA and SKP1 and Elongin C proteins; full list provided in Supplementary Table 6).

A multiple protein alignment was generated using Mafft version 7[54] (default parameters) and was then

manually edited with BioEdit v7.2.5[55] to retain only the region referring to the BTB core fold. The

phylogenetic relationships were inferred with the PhyML[57] procedure using the best-fitting protein

substitution model as determined by the SMS package[58].

For further comparisons with the K13 BTB and propeller domains, site-specific substitution rates λ

were inferred with FuncPatch for the BTB and propeller domains of several mammalian KCTD and

BTB-Kelch proteins, respectively. In the present study, the proteins were selected on the basis of their

sequence homology with K13, the availability of a solved 3D structure, and their known implication in

a Cullin-RING E3 ligase complex as suspected for K13. In addition, only well-characterized ligand-

binding function and the presence of a six-bladed propeller structure similar to the one of K13 were

considered for BTB-Kelch proteins. After a careful review of the literature, we selected two KCTD

proteins: SHKBP1 (UniProt code Q8TBC3) which regulates the epidermal growth factor receptor

(EGFR) signaling pathway[71]; and KCTD17 (Q8N5Z5) which mediates the ubiquitination and

proteasomal degradation of the ciliogenesis down-regulation TCHP protein[72]. Considering BTB-Kelch

proteins, we focused on: KEAP1 (Q14145) which interacts with its client protein Nrf2 for the

induction of cytoprotective responses to oxidative stress[37]; KLHL2 (O95198) and KLHL3 (Q9UH77)

which both participate in the ubiquitination and degradation of WNK substrates regulating blood

pressure[35]; and KLHL12 (Q53G59) which negatively regulates the WNT-beta-catenin pathway

through the degradation of Dishevelled proteins[38]. First, each *Homo sapiens* KCTD and BTB-Kelch

sequence was successively submitted as query sequence for a blastp search[52] (BLOSUM62 scoring

matrix, max target sequences fixed at 1,000) against the NCBI non-redundant protein database to

retrieve orthologous sequences from a large amount of species. The output lists were then filtered

according to specific criteria so as to keep only sequences having an unambiguous description (*i.e.* a

description that includes the name of the queried KCTD or BTB-Kelch protein), and that aligned with

$\geq 80\%$ sequence coverage and had $\geq 60\%$ sequence identity with the query sequence. The multiple

protein alignment of each set of orthologous sequences was then generated using Mafft version 7[54] (E-

INS-I strategy with BLOSUM62 scoring matrix, gap opening penalty 2.0 and offset 0.1). A second

filtering step was performed to remove incomplete or miss-annotated sequences, *i.e.* the sequences that

did not contain all the annotated domains (using the domain annotation automatically generated by the

Uniprot Knowledgebase) and/or that included a gapped position located in one of the annotated

domains. The final multiple protein alignments included: *i*) 124 sequences $\times$ 103 aligned positions for

SHKBP1; *ii*) 139 sequences $\times$ 102 aligned positions for KCTD17; *iii*) 135 sequences $\times$ 285 aligned

positions for KEAP1; *iv*) 162 sequences $\times$ 286 aligned positions for KLHL2; *v*) 158 sequences $\times$ 286

aligned positions for KLHL3; and *vi*) 129 sequences $\times$ 289 aligned positions for KLHL12. The full list

of orthologous sequences used for each mammalian KCTD and BTB-Kelch protein is provided in

Supplementary Table 7. Then, the phylogenetic relationships were inferred using PhyML[57] after

determining the best-fitting protein substitution model with the SMS package[58]. The 3D structures of

KCTD BTB and BTB-Kelch propeller domains were retrieved from the PDBsum database under the

following accession numbers: 4CRH for SHKBP1 (resolution: 1.72 Å)[28], 5A6R for KCTD17

(resolution: 2.85 Å)[28], 2FLU for KEAP1 (resolution: 1.50 Å, in complex with a Nrf2 peptide)[73], 4CHB

for KLHL2 (resolution: 1.56 Å, in complex with a WNK4 peptide)[35], 4CH9 for KLHL3 (resolution:

1.84 Å, in complex with a WNK4 peptide)[35], and 2VPJ for KLHL12 (resolution: 1.85 Å)[24].

Beforehand, the quality of each structure was validated using MolProbity[69]: none of the structures had

amino acids identified as outliers, and approximately 98% of the amino acids of each structure were in

favored regions.

Evaluation of structural properties. The electrostatic potential energy of each propeller structure

was calculated using the Adaptive Poisson-Boltzmann Solver (APBS) method[74]. Beforehand, the

required pqr input files were prepared using PDB2PQR v.2.1.1[75]. The missing charges were added

using the *Add Charge* function implemented in USCF Chimera[76]. A grid-based method was used to

solve the linearized Poisson-Boltzmann equation at 298 K, with solute (protein) and solvent dielectric

constant values fixed at 2 and 78.5, respectively. The contact surface selection was mapped using a

radius of 1.4 Å in a scale of -8 *kT/e* to +8 *kT/e*.

The relative solvent accessibility (RSA) was estimated as the accessible surface area of amino acids

using DSSP[77], then normalized with the maximal accessible surface area of each amino acid[78]. The

side-chain weighted contact number (WCN_{sc}) of each amino acid was calculated using a customized

python script provided by Sydykova and colleagues[79]. All structural properties were assessed using the

aforementioned PDB files.

Structure visualization. All molecular drawings were generated using the UCSF Chimera software[76].

Statistical analyses. Substitution rates among partitions were compared using non-parametric Mann-

Whitney *U* or Kruskal-Wallis *H* tests. When focusing on the propeller shallow pocket, contingency

tables were produced and statistically tested with chi-squared test. We used $p < 0.05$ as the cutoff for

significance in all statistical tests.

 Tables

 Table 1 – Lack of detectable positive selection in the *k13* gene using codon substitution models

Models	Purpose of the test	$2\Delta\ln L$	df	*p value*
M0-FR	Heterogeneity of ω among lineages of the phylogenetic tree	457.0	82	1.9×10^{-53}
M0-M3	Heterogeneity of ω across codon sites	1,046.3	4	3.3×10^{-225}
M1a-M2a	Detection of sites evolving under positive selection	0	2	1
M7-M8	Detection of sites evolving under positive selection	0	2	1

 Note – LRT, likelihood ratio test; ω, non-synonymous (d_N) to synonymous (d_S) substitution rate ratio
 (d_N/d_S); $2\Delta\ln L$, log-likelihood ratio of the two tested models; df, degrees of freedom. $2\Delta\ln L$ was
 compared to chi-squared table to determine the significance of the LRTs. The M1a-M2a comparison is
 reported to be more stringent than the M7-M8 comparison[31].

 Table 2 – Spatial correlation of amino acid substitution rates in the BTB and propeller domains of
 K13 and other proteins estimated by FuncPatch analysis

Protein domain	l	σ	log Bayes factor
K13$_{\text{BTB-propeller}}$	5	0.9	30.84
K13$_{\text{propeller}}$	3	0.9	20.67
SHKBP1$_{\text{BTB}}$	11	1.5	13.27
KCTD17$_{\text{BTB}}$	7	1.3	9.69
KEAP1$_{\text{propeller}}$	5	1.7	44.06
KLHL2$_{\text{propeller}}$	7	1.5	93.54
KLHL3$_{\text{propeller}}$	7	1.3	79.47
KLHL12$_{\text{propeller}}$	7	1.5	25.18

 Note – l, strength of the spatial correlation of site-specific substitution rates λ over the protein tertiary
 structures, in Å; σ, signal standard deviation measuring the variability of λ substitution rates across
 amino acid sites; log Bayes factor, statistical significance of the spatial correlation of site-specific
 substitution rates λ. A log Bayes factor ≥ 8 identifies a significant spatial correlation of site-specific
 substitutions rates λ in protein tertiary structure (conservative cutoff as suggested by FuncPatch'
 authors[33]).

551 **Table 3 –** Conservation level of the amino acids located at the surface of the K13 propeller shallow
552 pocket
553

Amino acid position	Reference amino acid (*P. fal*)	PfK13 mutant [a]	Sequence variation across species		λ-based propeller rank n, (%) [b]
			Plasmodium (n = 21)	Non-*Plasmodium* (n = 22)	
451	F	—	F (21)	H (16); C (4); T (2); S(1); N (1)	132 (46.48)
456	Y	—	Y (21)	H (12); I (7); P (2); Y (1)	125 (44.01)
482	Y	S[4] (Africa)	Y (21)	Y (17); H (5)	37 (13.03)
498	N	I[80] (Africa)	N (21)	Q (13); Y (7); H (2)	44 (15.49)
529	**R**	**K[80] (Africa)**	**R (21)**	**R (22)**	**1 (0.35)**
530	**N**	**K/Y[80] (Africa)**	**N (21)**	**N (22)**	**9 (3.17)**
546	**Y**	**—**	**Y (21)**	**F (22)**	**13 (4.58)**
551	I	—	I (21)	I (19); V (3)	86 (30.28)
576	**S**	**L[81] (Africa)**	**S (21)**	**S (22)**	**3 (1.06)**
577	**S**	**P[4] (Africa)**	**S (21)**	**S (22)**	**10 (3.52)**
593	**T**	**S[10] (Africa)**	**T (15); A (6)**	**T (22)**	**19 (6.69)**
597	**R**	**—**	**R (21)**	**R (22)**	**12 (4.23)**
623	**S**	**C[8] (Asia)**	**S (21)**	**S (22)**	**2 (0.70)**
624	**S**	**—**	**S (21)**	**A (15); G (7)**	**23 (8.10)**
640	I	V[4] (Africa)	I (21)	I (19); M (2); V(1)	42 (14.79)
671	M	—	M (21)	M (17); L (3); I (1); F (1)	73 (25.70)
672	N	—	N (21)	D (22)	47 (16.55)
688	E	—	E (21)	Q (22)	45 (15.85)
717	F	—	F (21)	Y (19); S (3)	33 (11.62)

554
Note – [a] None of the PfK13 mutant positions located at the surface of the K13 propeller shallow
pocket was reported to confer the ART-R phenotype. All PfK13 mutants presented here have been
described in one or two parasite samples from population surveys. [b] The rank attributed to the shallow
pocket positions was based on the site-specific substitution rates λ (Supplementary Data 1) estimated
for the 284 studied propeller positions using FuncPatch[33]. The lower the rank, the more conserved was
the position. Positions in bold are those belonging to the 10% most conserved positions of propeller.
561
562

564 **Table 4 –** Distribution of the conserved propeller shallow pocket amino acids for K13 and some
565 members of the BTB-Kelch protein family
566

Protein	Conservation level [a]	Shallow pocket positions [b]	Others positions	*p value*
PfK13$_{propeller}$	Highly conserved[10%]	9	20	1.60×10^{-5}
	Others	10	245	
KEAP1$_{propeller}$	Highly conserved[10%]	12	17	1.44×10^{-7}
	Others	12	244	
KLHL2$_{propeller}$	Highly conserved[10%]	13	16	1.26×10^{-10}
	Others	6	251	
KLHL3$_{propeller}$	Highly conserved[10%]	12	17	1.36×10^{-9}
	Others	6	251	
KLHL12$_{propeller}$	Highly conserved[10%]	14	15	2.94×10^{-12}
	Others	5	255	

567
Note – [a] The propeller positions were categorized into two groups: the *highly conserved* group
contained the 10% most conserved positions according to the λ substitution rates, the *other* group
included the remaining propeller positions. [b] The shallow pocket of K13 propeller was defined as the
positions forming the surface plan of the pocket and protruding out of the plan. A view and delineation
of the PfK13 shallow pocket is provided in Supplementary Fig. 6. The same structural delineation of
the propeller shallow pocket was applied to the four other BTB-Kelch propeller domains. Contingency
tables were subjected to chi-squared test.

 Figure legends

Fig 1 – Conservation level of *k13* and four other Kelch-containing protein-coding sequences compared
to the all *Plasmodium* protein-coding genes. The conservation level (d_N/d_S) was estimated for 3,256
orthologous genes among six *Plasmodium* species (*P. falciparum*, *P. berghei*, *P. chabaudi*, *P. vivax*,
P. yoelii and *P. knowlesi)* under the one-ratio PAML model by Jeffares and colleagues[32]. Histograms
show the distribution of one-ratio PAML values for protein-coding genes whose length is comprised in
an interval of ± 100 amino acids centered on each Kelch-containing protein length (sized-rank).
PlasmoDB accession numbers of the five Kelch-containing proteins investigated are provided above
each histogram. Red bars indicate the position of each of these Kelch-containing proteins in their
respective d_N/d_S distribution, with vertical dashed lines showing the five percent cutoff of the most and
less conserved protein-coding genes. The table, shown below the histograms, provides the precise rank
and sized-rank of each Kelch-containing protein-coding sequence. This latter corrects for a correlation
between protein length and one-ratio PAML value.

Fig 2 – Conservation level between the three annotated domains of K13. The scatter plot (*left*) shows
the conservation level of *k13* codon sites using the *pfk13* sequence as reference for the site numbering
(starting at codon 213). White circles correspond to inter-domain positions. All codon sites were
reported to evolve under strong purifying selection, with ω drastically < 1. The box-whisker plot
(*right*) shows that BTB evolves under more intense purifying selection compared to either CCC or
propeller, using non-parametric Mann-Whitney U test ($p < 0.05$). Box boundaries represent the first
and third quartiles and the length of whiskers correspond to 1.5 times the interquartile range. We used
the ω estimates obtained under the best fitted PAML model that indicates a variable selective regime
among sites.

Fig. 3 – Conservation and structure homology of the K13 BTB fold. **a.** Linear schematic
representation of the BTB fold of some BTB-containing protein families. Yellow arrows and cyan
cylinders represent strands and helices, respectively. Stars in magenta correspond to the 10% most
conserved K13 BTB amino acid sites (based on the λ substitution rate ranking, FuncPatch analysis,
Supplementary Data 1). Structural elements are labelled. **b.** Unrooted phylogenetic tree of the BTB
protein superfamily using a few reference sequences per BTB-containing protein family. Each BTB-
containing protein family forms a monophyletic group, identified with a colored background. K13
BTB is represented in red and clusters with the KCTD protein family of BTB-containing proteins. **c.**
Patch of slowly evolving amino acid sites in a three-dimensional view of PfK13 BTB. The amino acid
sites are labelled using the PfK13 sequence as reference. **d.** Superposition of the K13 BTB fold with
two members of the KCTD protein family. The most conserved amino acid sites for each protein
structure based on FuncPatch analysis are shown (magenta, orange and red for K13 BTB (white),
SHKBP1 (green) and KCTD17 (cornflower blue), respectively). The patch of slowly evolving amino
acid sites shared similarities between these three BTB-containing proteins, with three common
positions: 357 on the B2-B3 loop and 397-398 on the B4-A4 loop.

Fig 4 – Structural organization of amino acid conservation across K13 propeller. **a.** Structure-based
amino acid alignment of the six blade repeats of PfK13 propeller. The *x axis* shows the position in the
structure-based amino acid alignment of the six blades; the left *y axis* shows the blade number,
followed by the number of the first and last blade position. The strands of PfK13 propeller are colored
in grey; the AB and CD loops forming the upper face of the propeller are colored in cyan; the BC and
DA loops architecting the bottom face of the propeller are colored in yellow and green, respectively. A
consensus sequence of the blades was defined: strict consensus amino acids are shown in bold capital

letters, and highly conserved amino acids are shown in standard lowercase. A mapping of the different strands and loops onto the three-dimensional structure of PfK13 propeller is shown as surface above the structure-based amino acid alignment: side view (*left*), bottom view (*middle*) and upper view (*right*). The outline of the shallow pocket surface was delineated with a black line on the bottom view structure (*middle*) and the amino acid sites forming at the shallow pocket surface were surrounded and written in bold in the structure-based amino acid alignment. **b.** Mapping of the λ substitution rates onto the structure-based amino acid alignment of the six blade repeats of PfK13 propeller. *Upper graph*: heat map showing the λ substitution rate for each amino acid site. Black boxes correspond to the gaps in the structure-based amino acid alignment. The median λ substitution rate value was first calculated (white) to produce a scale ranging from the lowest (magenta) to the highest (green) site-specific substitution rate λ. *Lower graph*: Plot of the mean and 95% confidence interval of the λ substitution rates along the structure-based amino acid alignment of the six blades. The positions including one or more gaps were discarded.

Fig. 5 – Conservation level and electrostatic potential across the propeller structures of K13 and other BTB-Kelch proteins. **a.** Location of the 10% most conserved amino acid sites (magenta) on the three-dimensional structure of PfK13 propeller. The conservation level of positions was defined using the site-specific substitution rates λ (Supplementary Data 1). The propeller structure is shown from the side view as cartoon (*left* structure) and from the bottom view as surface (*right* structure). The amino acid sites forming the surface of the shallow pocket and belonging to the 10% most conserved sites are labelled. **b.** Electrostatic surface potential of the PfK13 propeller structure, estimated with the APBS method. Electrostatic potential values are in units of *kT/e* at 298 K, on a scale of -8 *kT/e* (red) to +8 *kT/e* (blue). White color indicates a neutral potential. The missing charges were added using the *Add Charge* function implemented in USCF Chimera. **c.** Location of the 10% most conserved amino acid sites and electrostatic potential for KEAP1, KLHL2, KLHL3 and KLHL12 propeller structures. The color code and structure orientation are the same as for K13 in Fig. 4a, b. For KEAP1, KLHL2 and KLHL3, the key amino acids interacting with their respective protein substrates are labelled. The PDB IDs and protein substrates are provided above each propeller structure.

References

1. Dondorp, A. M. *et al.* Artemisinin resistance in *Plasmodium falciparum* malaria. *N. Engl. J. Med.* **361,** 455–467 (2009).

2. Duru, V. *et al. Plasmodium falciparum* dihydroartemisinin-piperaquine failures in Cambodia are associated with mutant K13 parasites presenting high survival rates in novel piperaquine in vitro assays: retrospective and prospective investigations. *BMC Med.* **13,** 305 (2015).

3. Phyo, A. P. *et al.* Declining efficacy of artemisinin combination therapy against *P. falciparum* malaria on the Thai-Myanmar border (2003-2013): the role of parasite genetic factors. *Clin. Infect. Dis.* **63,** 784–791 (2016).

4. Ménard, D. *et al.* A worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. *N. Engl. J. Med.* **374,** 2453–2464 (2016).

5. WHO. Status report on artemisinin and ACT resistance, april 2017. http://www.who.int/malaria/publications/atoz/artemisinin-resistance-april2017/en/ (2017).

6. Lu, F. *et al.* Emergence of indigenous artemisinin-resistant *Plasmodium falciparum* in Africa. *N. Engl. J. Med.* **376,** 991–993 (2017).

7. Ikeda, M. *et al.* Artemisinin-resistant *Plasmodium falciparum* with high survival rates, Uganda, 2014-2016. *Emerging Infect. Dis.* **24,** 718–726 (2018).

8. Ariey, F. *et al.* A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature* **505,** 50–55 (2014).

9. Straimer, J. *et al.* K13-propeller mutations confer artemisinin resistance in *Plasmodium falciparum* clinical isolates. *Science* **347,** 428–431 (2015).

10. MalariaGEN Plasmodium falciparum Community Project. Genomic epidemiology of artemisinin resistant malaria. *Elife* **5,** pii: e08714 (2016).

11. Anderson, T. J. C. *et al.* Population parameters underlying an ongoing soft sweep in Southeast Asian malaria parasites. *Mol. Biol. Evol.* **34,** 131–144 (2017).

12. Cerqueira, G. C. *et al.* Longitudinal genomic surveillance of *Plasmodium falciparum* malaria

parasites reveals complex genomic architecture of emerging artemisinin resistance. *Genome Biol.*

18, 78 (2017).

13. Witkowski, B. *et al.* Novel phenotypic assays for the detection of artemisinin-resistant

Plasmodium falciparum malaria in Cambodia: *in-vitro* and *ex-vivo* drug-response studies. *Lancet*

Infect. Dis. **13,** 1043–1049 (2013).

14. Birnbaum, J. *et al.* A genetic system to study *Plasmodium falciparum* protein function. *Nat.*

Methods **14,** 450–456 (2017).

15. Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite *Plasmodium*

falciparum by saturation mutagenesis. *Science* **360,** pii: eaap7847 (2018).

16. The Gene Ontology Consortium. Expansion of the gene ontology knowledgebase and resources.

Nucleic Acids Res. **45,** D331–D338 (2017).

17. Mbengue, A. *et al.* A molecular mechanism of artemisinin resistance in *Plasmodium falciparum*

malaria. *Nature* **520,** 683–687 (2015).

18. Dogovski, C. *et al.* Targeting the cell stress response of *Plasmodium falciparum* to overcome

artemisinin Resistance. *PLoS Biol.* **13,** e1002132 (2015).

19. Mok, S. *et al.* Population transcriptomics of human malaria parasites reveals the mechanism of

artemisinin resistance. *Science* **347,** 431–435 (2015).

20. Zhang, M. *et al.* Inhibiting the *Plasmodium* eIF2α kinase PK4 prevents artemisinin-induced

latency. *Cell Host & Microbe* **22,** 766–776.e4 (2017).

21. Bhattacharjee, S. *et al.* Remodeling of the malaria parasite and host human red cell by vesicle

amplification that induces artemisinin resistance. *Blood* **131,** 1234–1247 (2018).

22. Adams, J., Kelso, R. & Cooley, L. The kelch repeat superfamily of proteins: propellers of cell

function. *Trends Cell Biol.* **10,** 17–24 (2000).

23. Dhanoa, B. S., Cogliati, T., Satish, A. G., Bruford, E. A. & Friedman, J. S. Update on the Kelch-

like (KLHL) gene family. *Hum. Genomics* **7,** 13 (2013).

24. Canning, P. *et al.* Structural basis for Cul3 protein assembly with the BTB-Kelch family of E3

ubiquitin ligases. *J. Biol. Chem.* **288,** 7803–7814 (2013).

25. Tilley, L., Straimer, J., Gnädig, N. F., Ralph, S. A. & Fidock, D. A. Artemisinin action and

resistance in *Plasmodium falciparum. Trends Parasitol.* **32,** 682–696 (2016).

26. Geyer, R., Wee, S., Anderson, S., Yates, J. & Wolf, D. A. BTB/POZ domain proteins are putative

substrate adaptors for Cullin 3 ubiquitin ligases. *Mol. Cell* **12,** 783–790 (2003).

27. Stogios, P. J., Downs, G. S., Jauhal, J. J., Nandra, S. K. & Privé, G. G. Sequence and structural

analysis of BTB domain proteins. *Genome Biol.* **6,** R82 (2005).

28. Pinkas, D. M. *et al.* Structural complexity in the KCTD family of Cullin3-dependent E3 ubiquitin

ligases. *Biochem. J.* **474,** 3747–3761 (2017).

29. Furukawa, M. & Xiong, Y. BTB protein Keap1 targets antioxidant transcription factor Nrf2 for

ubiquitination by the Cullin 3-Roc1 ligase. *Mol. Cell. Biol.* **25,** 162–171 (2005).

30. Shibata, S., Zhang, J., Puthumana, J., Stone, K. L. & Lifton, R. P. Kelch-like 3 and Cullin 3

regulate electrolyte homeostasis via ubiquitination and degradation of WNK4. *Proc. Natl. Acad.*

Sci. U.S.A. **110,** 7838–7843 (2013).

31. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24,** 1586–1591

(2007).

32. Jeffares, D. C., Tomiczek, B., Sojo, V. & dos Reis, M. A beginners guide to estimating the non-

synonymous to synonymous rate ratio of all protein-coding genes in a genome. *Methods Mol.*

Biol. **1201,** 65–90 (2015).

33. Huang, Y.-F. & Golding, G. B. FuncPatch: a web server for the fast Bayesian inference of

conserved functional patches in protein 3D structures. *Bioinformatics* **31,** 523–531 (2015).

34. Huang, Y.-F. & Golding, G. B. Phylogenetic Gaussian process model for the inference of

functionally important regions in protein tertiary structures. *PLoS Comput. Biol.* **10,** e1003429

(2014).

35. Schumacher, F. R., Sorrell, F. J., Alessi, D. R., Bullock, A. N. & Kurz, T. Structural and

biochemical characterization of the KLHL3–WNK kinase interaction important in blood pressure

regulation. *Biochem. J.* **460,** 237–246 (2014).

36. Canning, P., Sorrell, F. J. & Bullock, A. N. Structural basis of Keap1 interactions with Nrf2. *Free*

Radic. Biol. Med. **88,** 101–107 (2015).

37. Cullinan, S. B. *et al.* Nrf2 Is a Direct PERK substrate and effector of PERK-dependent cell

survival. *Mol. Cell. Biol.* **23,** 7198–7209 (2003).

38. Angers, S. *et al.* The KLHL12–Cullin-3 ubiquitin ligase negatively regulates the Wnt–β-catenin

pathway by targeting Dishevelled for degradation. *Nat. Cell Biol.* **8,** 348–357 (2006).

39. Echave, J., Spielman, S. J. & Wilke, C. O. Causes of evolutionary rate variation among protein

sites. *Nat. Rev. Genet.* **17,** 109–121 (2016).

40. Yang, Z., Nielsen, R., Goldman, N. & Pedersen, A. M. Codon-substitution models for

heterogeneous selection pressure at amino acid sites. *Genetics* **155,** 431–449 (2000).

41. Sawyer, S. L., Wu, L. I., Emerman, M. & Malik, H. S. Positive selection of primate TRIM5alpha

identifies a critical species-specific retroviral restriction domain. *Proc. Natl. Acad. Sci. U.S.A.*

102, 2832–2837 (2005).

42. Sidik, S. M. *et al.* A genome-wide CRISPR screen in *Toxoplasma* identifies essential

apicomplexan genes. *Cell* **166,** 1423–1435.e12 (2016).

43. Bushell, E. *et al.* Functional profiling of a *plasmodium* genome reveals an abundance of essential

genes. *Cell* **170,** 260–272.e8 (2017).

44. Hast, B. E. *et al.* Cancer-derived mutations in KEAP1 impair NRF2 degradation but not

ubiquitination. *Cancer Res.* **74,** 808–817 (2014).

45. Mori, Y. *et al.* Decrease of WNK4 ubiquitination by disease-causing mutations of KLHL3

through different molecular mechanisms. *Biochem. Biophys. Res. Commun.* **439,** 30–34 (2013).

46. Haldar, K., Bhattacharjee, S. & Safeukui, I. Drug resistance in *Plasmodium*. *Nat. Rev. Microbiol.*

16, 156–170 (2018).

47. Correale, S. *et al.* Molecular organization of the cullin E3 ligase adaptor KCTD11. *Biochimie* **93,**

715–724 (2011).

48. Boullé, M. *et al.* Artemisinin-resistant *Plasmodium falciparum* K13 mutant alleles, Thailand–

Myanmar border. *Emerg. Infect. Dis.* **22,** 1503–1505 (2016).

49. Hartmann, M. D. Functional and structural roles of coiled coils. *Subcell. Biochem.* **82,** 63–93

(2017).

50. de Moura, T. R. *et al.* Prp19/Pso4 is an autoinhibited ubiquitin ligase activated by stepwise

assembly of three splicing factors. *Mol. Cell* **69,** 979–992.e6 (2018).

51. Aurrecoechea, C. *et al.* EuPathDB: the eukaryotic pathogen genomics database resource. *Nucleic*

Acids Res. **45,** D581–D591 (2017).

52. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search

tool. *J. Mol. Biol.* **215,** 403–410 (1990).

53. Hunter, S. *et al.* InterPro: the integrative protein signature database. *Nucleic Acids Res.* **37,** D211–

D215 (2009).

54. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30,** 772–780 (2013).

55. Hall, T. BioEdit: a user-friendly biological sequence alignment editor and analysis program for

Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41,** 95–98 (1999).

56. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34,** W609–W612 (2006).

57. Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies:

assessing the performance of PhyML 3.0. *Syst. Biol.* **59,** 307–321 (2010).

58. Lefort, V., Longueville, J.-E. & Gascuel, O. SMS: smart model selection in PhyML. *Mol. Biol.*

Evol. **34,** 2422–2424 (2017).

59. Anisimova, M., Bielawski, J. P. & Yang, Z. Accuracy and power of the likelihood ratio test in

detecting adaptive molecular evolution. *Mol. Biol. Evol.* **18,** 1585–1592 (2001).

60. Parfrey, L. W., Lahr, D. J. G., Knoll, A. H. & Katz, L. A. Estimating the timing of early

eukaryotic diversification with multigene molecular clocks. *Proc. Natl. Acad. Sci. U.S.A.* **108,**

13624–13629 (2011).

61. Goldman, N. & Yang, Z. A codon-based model of nucleotide substitution for protein-coding DNA

sequences. *Mol. Biol. Evol.* **11,** 725–736 (1994).

62. Nielsen, R. & Yang, Z. Likelihood models for detecting positively selected amino acid sites and

applications to the HIV-1 envelope gene. *Genetics* **148,** 929–936 (1998).

63. Yang, Z. Likelihood ratio tests for detecting positive selection and application to primate

lysozyme evolution. *Mol. Biol. Evol.* **15,** 568–573 (1998).

64. Vuong, Q. H. Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica*

57, 307–333 (1989).

65. Yang, Z., Wong, W. S. W. & Nielsen, R. Bayes empirical bayes inference of amino acid sites

under positive selection. *Mol. Biol. Evol.* **22,** 1107–1118 (2005).

66. Muse, S. V. & Gaut, B. S. A likelihood approach for comparing synonymous and nonsynonymous

nucleotide substitution rates, with application to the chloroplast genome. *Mol. Biol. Evol.* **11,** 715–

724 (1994).

67. Rodrigue, N., Lartillot, N., Bryant, D. & Philippe, H. Site interdependence attributed to tertiary

structure in amino acid sequence evolution. *Gene* **347,** 207–217 (2005).

68. Robinson, D. M., Jones, D. T., Kishino, H., Goldman, N. & Thorne, J. L. Protein evolution with

dependence among codons due to tertiary structure. *Mol. Biol. Evol.* **20,** 1692–1704 (2003).

69. Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography.

Acta Crystallogr. D Biol. Crystallogr. **66,** 12–21 (2010).

70. Delano, W. The PyMOL molecular graphics system. http://www.pymol.org/ (2002).

71. Feng, L., Wang, J.-T., Jin, H., Qian, K. & Geng, J.-G. SH3KBP1-binding protein 1 prevents

epidermal growth factor receptor degradation by the interruption of c-Cbl-CIN85 complex. *Cell*

Biochem. Funct. **29,** 589–596 (2011).

72. Kasahara, K. *et al.* Ubiquitin-proteasome system controls ciliogenesis at the initial step of

axoneme extension. *Nat. Commun.* **5,** 5081 (2014).

73. Lo, S.-C., Li, X., Henzl, M. T., Beamer, L. J. & Hannink, M. Structure of the Keap1:Nrf2

interface provides mechanistic insight into Nrf2 signaling. *EMBO J.* **25,** 3605–3617 (2006).

74. Baker, N. A., Sept, D., Joseph, S., Holst, M. J. & McCammon, J. A. Electrostatics of

nanosystems: application to microtubules and the ribosome. *Proc. Natl. Acad. Sci. U.S.A.* **98,**

10037–10041 (2001).

75. Dolinsky, T. J., Nielsen, J. E., McCammon, J. A. & Baker, N. A. PDB2PQR: an automated

pipeline for the setup of Poisson-Boltzmann electrostatics calculations. *Nucleic Acids Res.* **32,**

W665–W667 (2004).

76. Pettersen, E. F. *et al.* UCSF Chimera—A visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25,** 1605–1612 (2004).

77. Kabsch, W. & Sander, C. Dictionary of protein secondary structure: pattern recognition of

hydrogen-bonded and geometrical features. *Biopolymers* **22,** 2577–2637 (1983).

78. Tien, M. Z., Meyer, A. G., Sydykova, D. K., Spielman, S. J. & Wilke, C. O. Maximum allowed

solvent accessibilites of residues in proteins. *PLoS One* **8,** e80635 (2013).

79. Sydykova, D. K., Jack, B. R., Spielman, S. J. & Wilke, C. O. Measuring evolutionary rates of

proteins in a structural context. *F1000Res.* **6,** 1845 (2017).

80. de Laurent, Z. R. *et al.* Polymorphisms in the K13 gene in *Plasmodium falciparum* from different

malaria transmission areas of Kenya. *Am. J. Trop. Med. Hyg.* **98,** 1360–1366 (2018).

81. Kamau, E. *et al.* K13-propeller polymorphisms in *Plasmodium falciparum* parasites from Sub-

Saharan Africa. *J. Infect. Dis.* **211,** 1352–1355 (2015).

Acknowledgements

We thank YF. Huang and GB. Golding for help with FuncPatch analysis. We thank O. Mercereau-Puijalon, M. Miteva, and R. Duval for helpful discussions. We thank A. Sissoko and F. Palstra for proofreading the manuscript. R. Coppée is supported by PhD grant funding from the École Doctorale MTCI.

Author contributions

R.C., A.S. and J.C. designed the study. D.J. provided data. R.C. performed most of the analyses. R.C., A.S., and J.C. contributed to interpretations. All authors have participated in paper writing and approved it prior to submission.

Competing interests

The authors declare no competing financial interests.

Accession number	Description	one-ratio PAML value	Protein length (aa)	Rank (%)	Sized-rank (%)
PF3D7_1343700	kelch protein K13	0.015	726	140/3,256 (4.3)	6/393 (1.5)
PF3D7_0724800	kelch domain-containing protein, putative	0.083	407	1,532/3,256 (47.1)	388/742 (52.3)
PF3D7_1022600	kelch protein K10	0.068	877	1,171/3,256 (36.0)	83/300 (27.3)
PF3D7_1125700	kelch domain-containing protein, putative	0.107	515	1,949/3,256 (59.9)	407/647 (62.9)
PF3D7_0724800	kelch domain-containing protein, putative	0.083	407	1,532/3,256 (47.1)	388/742 (52.3)