Computer Simulations of the Humoral Immune System Reveal How Imprinting Can Affect Responses to Influenza HA Stalk with Implications for the Design of Universal Vaccines

Christopher S. Anderson[1], Mark Y. Sangster[1], Hongmei Yang[2], Sidhartha Chaudhury[3], David J. Topham[1*]

[1]New York Influenza Center of Excellence at David Smith Center for Immunology and Vaccine Biology, Department of Microbiology and Immunology, University of Rochester School of Medicine and Dentistry, Rochester, NY. [2]Department of Biostatistics and Computational Biology, University of Rochester Medical Center, Rochester, NY. [3]BHSAI/U.S. Army MRMC, Frederick, MD.

*Corresponding author. E-mail: David_Topham@URMC.rochester.edu

Abstract

Background: Successful vaccination against the H1N1 Influenza A virus has required the continuous development of new vaccines that are antigenically similar to currently circulating strains. Vaccine strategies that can increase the cross-reactivity of the antibody response, especially to conserved regions, are essential to creating long-lasting immunity to H1N1 viruses. How pre-existing immunity affects vaccine-induced antibody cross-reactivity is still not well understood.

Methods: An immunological shape space of antigenic sites of hemagglutinin (HA) was constructed using viral sequence data. A Gillespie Algorithm-based model of the humoral immune system was used to simulate B cell responses to A/California/07/2009 (CA09) HA antigen after prior immunization with an antigenically similar or dissimilar strain. The effect of pre-existing memory B cells and antibody on the resulting antibody responses was interrogated.

Results: We found increased levels of highly-cross-reactive antibodies after immunization with antigenically dissimilar strains. This increase was dependent on pre-existing memory B cells. Furthermore, pre-existing antibody also interfered with the cross-reactive antibody response, but this effect occurred irrespective of the priming antigen.

Conclusion: These findings suggest that vaccination by divergent strains will boost highly-cross-reactive antibodies by selectively targeting memory B cells specific to conserved antigenic sites and by reducing the negative interference caused by pre-existing antibody.

Introduction

Vaccination strategies are needed that induced highly-cross-reactive antibodies capable of binding a broad range of antigenically distinct influenza virus strains[1]. The seasonal H1N1 influenza virus vaccine requires continuous updating to compensate for antigenic drift in order to be effective[2]. Due to antigenic shift, the season H1N1 vaccine was not effective against the 2009 "Swine flu" pandemic[3]. In order to overcome these challenges novel vaccination strategies are needed[4].

The 2009 pandemic virus vaccine was capable of significantly boosting cross-

reactivity, but only in younger age groups[5]. Serum antibody from these individuals

bound a broad range of seasonal viruses and highly divergent avian influenza

strains[5,6]. Additionally, there was an induction of antibodies towards the conserved

stalk region of HA[6,7]. Additionally studies have suggested a role of Memory B cells in

these highly cross-reactive antibody responses[8,9].

The age-specific differences seen after vaccination (or infection) with CA09 are

thought to result from differences in the strains individuals have been previously

exposed. Both pre-existing antibody and memory b cell specificity have been implicated

in interfering with the cross-reactivity of secondary immune responses[9-14]. How pre-

existing antibody and memory B cells contributed to the differences in cross-reactivity

seen in different age groups is not well understood.

To gain insights into the role of memory B cells and pre-existing antibodies on

cross-reactivity, we simulated secondary immune responses to CA09 HA antigen after

priming with antigenically similar or distinct HA antigens. We investigated the effect pre-

exposure to antigenically distinct antigens had on the cross-reactivity of a secondary

antibody response.

Materials and Methods

HA Protein Sequences

Influenza HA protein sequences used in the model were obtained from Genbank: A/California/07/2009 (CA09) [NC_026433], A/Brisbane/59/2007 (BR07) [KP458398], A/South Carolina/01/1918 (SC18) [AF117241], A/Beijing/262/1995 (BE95) [AAP34323], A/Brazil/11/1978 (BR78) [A4GBX7], A/Chile/1/1983 (CH83) [A4GCH5], A/New Caledonia/20/99 (NC99) [AY289929], A/Singapore/6/1986 (SI86) [ABO38395], A/Solomon Islands/3/2006 (SI06) [ABU99109], A/USSR/90/1977 (US77) [P03453], A/New Jersey/11/1976 (NJ76) [ACU80014].

Model Description

Simulations were performed using a Gillespie algorithm and a set of rate equations. The rate equations represent biological processes of B cells that occur during exposure to viral antigen. Together, these biological processes represent a simplified humoral immune system that can responds to antigen, produce antibodies, and remove antigen from the system. The model is identical to those described by Chaudhury et al. [15] accept for two modifications: (1) the number of antigenic sites representing each antigen was increased from 2 to 6 (2) long-lived plasma cells were added to the model [9] (Fig 1; Sup. Methods).

Fig 1. Schematic Representation of Humoral Immune System Model.

81   Schematic is adapted from Chaudhury et al. [15]

Simulating Immune Responses to the 2009 Pandemic Vaccine

Two scenarios were modeled with 50 simulations carried out for each scenario.

In scenario one, the model was "primed" (antigen added to the system) with the SC18

HA antigen, then the immune response was allowed to resolve for 365 days, during

which antibody level to returned close to baseline and subsequently immunized with

CA09 HA antigen (Fig 2A). Scenario two was identical to scenario one except the model

was primed with the BR07 HA antigen. B cell and antibody counts, genotype, and

antigen specificities were tracked during the simulation allowing quantification of

antigenic-site-specific B cells and antibodies during the simulation.

Perturbed Models

Two perturbed models were created in order to assess the contribution of pre-

existing antibody and memory B cells on the increase in stalk-specific antibodies. For

the first model ("No Clearance"), the antibody clearance rate equation (which removes

antigen form the system) was removed such that only basal decay of the antigen

occurred. For the second model ("No Memory"), the memory B cell activation rate

equation (which initiates germinal centers from memory B cells) was removed from the

simulation such that only naïve B cells contributed to the germinal center reactions.

Simulations were performed identical to the original model.

2009 H1N1 Vaccine Clinical Trial Human Serum

Samples from a previous clinical trial were used to test specific predictions of the simulations. The study was conducted under a protocol approved by the University of Rochester Research Subjects Review Board. Informed written consent was obtained from each participant. ClinicalTrials.gov identifier NCT01055184. Healthy adults and children were enrolled as previously described and results of this clinical trial have been published previously[6]. Subjects received a single intramuscular (i.m.) injection of inactivated influenza A/California/07/2009 (H1N1) monovalent subunit vaccine (Novartis). Each 0.5-ml dose contained 15µg of HA antigen. Administration of the vaccine (study day 0) took place from January 2010 to March 2010. Serum was collected before and 28 days after vaccination.

Enzyme-linked Immunosorbent Assay

In order to test predictions of the model, we measured human serum antibody levels by Enzyme-linked immunosorbent assays (ELISA). Serum antibody levels were measured against recombinant HA proteins derived from historical antigens. ELISAs were performed using recombinant HA proteins coated on MaxiSorb 96-well plates (ThermoSci; 439454) overnight at 4°C. Plates were blocked with 3% bovine serum albumin (BSA) in phosphate buffered saline (PBS) for 1hr at room temperature. Serum was diluted 1:1000 in PBS/0.5% BSA/0.05% Tween-20. Plates were washed and incubated with alkaline phosphatase (AP)-conjugated secondary antibody for 2 hrs

at room temperature. Plates were washed and developed using AP substrate

(ThermoSci 34064). Recombinant HA proteins were obtained from Influenza Reagent

Resource (Cat#: FR-67, FR-692, FR-65, FR-180, FR-699) and BEI Resources (Cat#

NR-19240, NR-48873). Stalk specific antibody responses were measured using

chimeric HA proteins that contained an "exotic" HA head but retained the conserved

stalk region (cH9.1 and cH6.1). We report the relative change in antibody levels

(d28/d0).

Statistics

Two sample, two-tailed, t-test using the *t.test* function was performed using the

base packages in R. A p-value of 0.05 or less was considered statistically significant.

For group comparisons of the ELISA, a multivariate linear model (*lm* function in R) was

performed to determine statistically significant differences between groups.

Results

Antigenic Distance Determination

We first determined the antigenic distance (AD) between HA antigens from 11

influenza virus strains. AD was determined using the H1N1 HA sequence-based

antigenic distance approach previously described[16]. Given that each HA in the model

contains 6 antigenic sites, and each antigenic site in the model contains 20 characters,

the maximum epitopic distance (antigenic-site-specific antigenic distance; ED) is 20 and

the maximum AD for each antigen is 120. 0verall, the real-life HA antigens of SC18 and

CA09 had the greatest estimate of antigenic similarity with an AD of 21 (Table 1). BR07

and CA09 HA antigens had the greatest dissimilarity with an AD of 53. For SC18 and

CA09 HA antigens, four of the five head epitopes had an ED of less than or equal to

seven (Sa = 2, Sb = 3, Ca1 = 5, Ca2 =8, Cb = 3), with the Sa antigenic site having the

least distance. Alternatively, BR07 and CA09 HA antigens had only one cross-reactive

antigenic site (Sa = 8, Sb = 15, Ca1 = 7, Ca2 = 10, Cb = 13), with an ED of seven.

Thus, in the model SC18 HA antigen was antigenically more similar to CA09 while the

BR07 HA antigen was largely antigenically distinct from CA09.

Table 1 Antigenic Distances

	SC18	PR34	NJ76	US77	BR78	SI86	CH83	BE95	NC99	SI06	BR07	CA09
SC18	0											
PR34	43	0										
NJ76	16	43	0									
US77	50	46	50	0								
BR78	50	46	50	0	0							
SI86	45	48	53	18	18	0						
CH83	50	48	53	4	4	16	0					
BE95	53	48	53	22	22	21	23	0				

NC99	48	45	55	31	31	18	29	10	0			
SI06	47	50	54	35	35	23	33	18	10	0		
BR07	49	50	56	34	34	22	32	17	7	8	0	
CA09	21	48	24	55	55	50	55	58	53	51	53	0

Antigenic-Site-Specific Antibody Responses

Antigenic-site-specific antibodies and memory B cells reactive to the priming Ag (SC18 or BR07) were measured throughout the simulation. We found that counts of antibody and memory B cells specific for the priming antigen were similar across head antigenic sites and between groups (Fig 2A, S1 Fig A-D) and these similarities remained up until boosting (S1 Fig E-F). Stalk-specific antibody and memory B cell counts were significantly less than head antigenic sites making up about 10% of the total response and were similar for both groups (S1 Fig E and F). Therefore, antibody and memory B cell counts and specificities to their priming antigen were similar for both BR07-primed and SC18-primed groups.

Fig 2. Immune Responses After Prime and Boost.

(A) Antigenic-site-specific antibody titers to the priming antigen for the SC18 primed, CA09 boosted group (left) and BR07 primed, CA09 boosted group (right). Curves represent average titers for 50 simulations and colored area represents the standard deviation. Arrows represent times simulation was primed and boosted. (B) Average

antibody titers and (C) memory B cells cross-reactive to CA09 pre-boost (Day 365). Statistic represents result of two sample t-test. (D) Affinity (antigenic distances 1-7) of memory B cells to CA09 HA antigen. Statistic represents result of two sample t-test. (E) Percent of stalk antigenic site-specific antibody for each priming group.

Given that immune responses to the priming antigens were similar, we sought to determine if the cross-reactivity to CA09 differed between groups prior to boosting. We found a statistically significantly (p-values < 0.001) higher antibody and memory B cell levels cross-reactive to CA09 in the SC18 primed group compared to the BR07 primed group with a greater than 2-fold difference (Fig 2B-C). Although the amount of memory B cells with immunoglobulin receptors with low Hamming distance (high affinity) to CA09 HA antigen did not differ between groups, modest and high Hamming distance (low affinity) memory B cells were significantly increased in the SC18 group (Fig 2D). Taken together, priming-antigen-specific antibody and memory B cells were similar between groups, but were their cross-reactivity to CA09 was different.

After boosting with CA09 total antibody levels reactive to CA09 in the SC18 group were slightly higher, although this difference did not reach significance (S2 Fig A). For the SC18 primed group, antibodies to the Sa-antigenic-site of CA09 dominated with Sb and Cb antigenic-site specific antibodies also boosted (Fig 2A). Stalk antigenic site antibody were also boosted but to a lesser extent compared to head epitopes and comprised about 15% of the total antibody response (Fig 2E).

Unlike the SC18-primed group, stalk-specific antibody responses dominated for the BR07 primed group (Fig 2A) comprising 35% of the total antibody response (Fig 2E). A moderate increase in other antigenic-site-specific antibodies was also observed

(Fig 2A). Antigenic site-specific differences between groups generally corresponded to

differences in epitopic distances between the priming and CA09 antigens with those

with closer epitopic distances showing a greater boost. Antibodies to the stalk antigenic

site, which has the same epitopic distance in both groups, also demonstrated

differences between groups. Despite being subdominant during priming, the large

increase in stalk specific antibodies in the BR07-primed group demonstrates that the

shorter ED at this site more than compensated for the decreased immunogenicity.

Pre-Exposure Affects Cross-Reactivity of Secondary Responses

We found that after boosting with CA09 both groups had strong antibody

responses to the antigens to which they had been previously exposed, but differed

largely in responses to other strains (Fig 3A). This occurred despite the fact that

antigenic distances to the 11 strains were not significantly different between BR07 and

SC18 (two-sample t-test, p-value = 0.362). Generally, the SC18-primed group was

cross-reactive to strains antigenically similar to CA09, while the BR07-primed group

antibody response demonstrated greater cross-reactive to other strains. The BR07-

primed group had a statistically significant increase in the number of stalk antibodies

which were cross-reactive to all HA antigens (Fig 3B). Cross-reactive antibody levels in

the SC18 primed group correlated well with the antigenic distance from CA09, while the

BR07 primed group antibody cross-reactivity showed no linear correlation with antigenic

distance (pval = 0.0001, pval = 0.4983; respectively). Therefore, although antigenic

distance was a good predictor of cross-reactivity during the primary response of the

simulation, for secondary immune responses antigenic distance alone was not sufficient

to predict cross-reactive immune responses in individuals with different antigen

exposure histories.

Fig 3. Cross-reactivity After Boosting with CA09.

(A) Antibody levels to HA antigens from representative strains that circulated from 1918-

2009 for the SC18-primed group or BR07-primed group. Antibody levels were taken at

30 days post-boost (day 395) and log transformed. Values are averages of 50

simulations. (B) Of all antibodies present at 30 days post-boost with CA09, pie-chart

represents the number of those antibodies that are cross-reactive to other HA antigens

(1-11 HA antigens). Number in parenthesis represents percentage rounded to the

nearest whole number. Asterisk represents statistically significant difference (p-value <

0.05) between SC18 and BR07 groups as determined by two sample t-test.

Contribution of Pre-Existing Memory B cells and Antibody on Cross-Reactivity

For the SC18-primed group, removal of memory B cell activation significantly

increased antibody levels to the historical antigens (two sample t-test, p-value = 8.5 x

10^{-26}) compared to the original unperturbed ("Normal") model (Fig 4A). Interestingly, this

increase occurred despite a decrease in stalk-specific antibody (Fig 4C). Removal of

antibody clearance for the SC18-primed group also significantly increased antibody

cross-reactivity (two sample t-test, p-value = 2.4 x 10^{-15}), but this was to a lesser extent.

For the BR07-primed group, removal of antibody clearance also significantly increased

the cross-reactive response (two sample t-test, p-value = 4 x 10^{-29}), but unlike the

SC18-primed group, removal of memory B cells from the germinal centers statistically

decreased (two sample t-test, p-value = 3 x 10^{-76}) the cross-reactive response (Fig 4B).

Stalk-specific antibody was also significantly increased in the "No Clearance" model, but

significantly decreased in the "No Memory" model. Taken together, pre-existing

antibody decreased cross-reactive antibody responses in both groups, while the effect

of memory B cells had the opposite effect on cross-reactivity across groups. The

increased cross-reactivity found in the BR07 primed group after boosting with CA09

resulted from activation of pre-existing memory B cells cross-reactive CA09 and not

primarily from naive B cell responses.

Fig. 4 Antibody Levels After Perturbation of the Model

(A) Antibody levels measured at day 30 post-boost with CA09 for the SC18-primed

group or (B) BR07-primed group for perturbed and normal models. Values were log

transformed. (C) Stalk-specific antibody levels 30 days post boost with CA09 for each

model. Average of 50 simulations, error bars represent standard deviation. Statistics

were determined by two-sample t-test.

Cross-Reactivity of Age-Stratified Serum

Finally, we sought to support our finding that cross-reactivity would be affected

by prior exposure. Using serum from a previously described clinical trial, we measured

antibody binding to a range of historical strains. Unsupervised clustering showed

stratification by age group although this grouping was not exact (Fig 5A). Cross-

reactivity antibody levels were generally increased in the 18-32-year-old age group, but

not to all HA antigens (Fig 5B). Importantly, the HA antigen, CH6.1, showed a

statistically significant increase in antibody levels for the 18-32-year-old age group

compared to the 60+ age group (multivariate regression model, p-value = 0.04358).

Alternatively, NC99 antibody binding was modestly increased in the 60+ age group,

although the difference did not pass our significance threshold (multivariate regression

model, p-value = 0.09916). Taken together, these finding support that cross-reactivity of

secondary immune responses are affected by exposure history.

Fig 5 Serum Antibody Levels for Age-Stratified Cohort

(A) Hierarchical clustering of antibody binding measured against recombinant HA

proteins using ELISA. (B) Heatmap of ELISA antibody binding data. Data represents the

relative change in binding. Data was log transformed and standardized, values

represent column z-scores.

Discussion

Protection against antigenically drifting or shifting H1N1 viruses by vaccination

requires continuous reformulation of the vaccine. Increasing the cross-reactivity induced

by vaccination is a hallmark of "universal" vaccine efforts[4]. Here we computationally

simulated immunization with SC18 HA antigen or BR07 HA antigen and subsequent

immunization with CA09 HA antigen. We also evaluated the effect of pre-existing

antibody and memory B cells on the levels and cross-reactivity of antibodies after

immunization with CA09 HA antigen.

Elucidating the combined effect of differences in HA antigenic site conservation,

pre-existing immunity, epitope dominance, and B cell and antibody specificities is a

daunting task for the experimentalist. However, computational models allow explicit

manipulation and observability of biological processes that are not possible with typical

animal and human models. Perelson et al. hypothesized that B cell receptor repertoires

(paratopes) exist in an immunological shape space and antigen binding differences

between them are represented as distance in shape space[17]. Smith et al.

subsequently derived the parameters of such an immunological shape space for

influenza viruses[18]. Moreover, Smith et al. developed a computational model of the

humoral immune system and demonstrated that such a model can be used to

understand secondary immune responses to influenza antigen[9]. Recently, Chaudhury

et al. developed a stochastic simulation model using the parameters developed by

Smith et al. and expanded the model to include multiple antigenic sites of different

conservation[15]. Here we adapted the Chaudhury et al. model to better simulate

immune responses to the H1N1 influenza HA antigen.

In the current study, we aimed to understand how prior exposure to influenza

virus antigens affects the antibody specificity during secondary exposures. This work

was an extension of the work originally performed by Smith et al.[19] and the theory of

Shape Space originally developed by Perelson et al.[17]. Consist with Smith et al.

findings, we found that the antigenic relationship between the first and secondary

exposure antigens largely affect the specificity of the antibody response. Moreover,

during secondary immune responses in the model, antigen was removed from the

system more quickly in the group previously exposed to an antigenically similar strain

during the primary exposure, consistent with the notion of antibody mediated negative

interference[9]. Moreover, the increase in stalk-specific antibodies in the BR07 group

during boosting is consistent with weakening of negative interference from head-specific

antibodies. Additionally, the increased antibody response to the CA09 strain in the

SC18 exposed group after boosting supports the notion of positive interference, in

which antibody responses from pre-existing memory B cells are increased. Taken

together, our findings are consistent with the Antigenic Distance Hypothesis described

by Smith et al.[9] .

The expansion of the Shape Space-based model to include multiple antigenic

sites by Chaudhury et al. was a major advancement in use of the model to understand

B cell specificity across complex antigens[20]. By incorporating multiple antigenic sites,

the model creates competition for antigen between B cells complementary to different

antigenic sites on the same antigen. Although Chaudhury et al. modeled a multivalent

vaccine, our findings are consistent with their finding that antibody responses to a

normally sub-dominant stalk antigenic site will dominate when the antigenic distance

between head antigens are large, even when immunogens are given successively.

Additionally, the large increase in stalk-specific antibodies in the BR07 group is

consistent with reports on universal vaccine development that apply a similar strategy to

boost stalk specific antibodies[4,21].

One of the most significant findings of the 2009 pandemic was the ability of 2009

pandemic vaccine to induce antibodies able to bind antigenically distinct viruses[22,23].

The results of our simulations corroborate these findings by demonstrating that BR07-

primed individuals had increased antibody reactivity to 1918-like HA antigens (CA09,

SC18, NJ76) as well as seasonal H1N1 viruses after exposure to CA09, while SC18-primed individuals produced antibodies primarily reactive to 1918-like HA antigens. These findings are consistent with the reports suggesting that original virus exposures, not age, affected the vaccine response to the 2009 vaccine[5,24]. Furthermore, although only slightly different, SC18 antibody titers were higher than CA09 titers after boosting with CA09 in the SC18 group but CA09 titers were higher than BR07 in the BR07 primed group. The SC18-primed group demonstrated antibody responses consistent with original antigenic sin (OAS), where individuals display antibody levels highest against viruses that circulated when they were young (and first exposed to influenza) despite subsequent exposure to drifted influenzas viruses[25]. The BR07-primed group did not show the same OAS phenotype as CA09 antibodies were highest after boosting, but antibody levels to BR07 were highest in the BR07 exposed group.

Pre-boost antibody levels of the SC18 primed group were almost 3-fold greater for CA09 than those primed for BR07, similar to what has been reported[26]. Additionally, the fold-change increases in the antibody response to the stalk is consistent with published reports[6]. We found that the antigenic site (Sa), which had the least antigenic difference among SC18 and CA09 HA head antigenic sites, dominated the antibody response after boosting with CA09 in the SC18-primed group. The Sa antigenic site dominance is the SC18 group is consistent with experimental data showing that antibody responses from the 60+ year old individuals had antibody responses focused on the Sa site of CA09[27]. Furthermore fold change titers (pre-boost/post-boost) were decreased in the SC18 primed group suggesting it is important

to take into account priming history of the elderly when trying to assess

immunosenescence or predict responses in different age groups[9,11,28,29].

Here we showed that cross-reactivity can be boosted with sequential

immunization by antigenically distinct antigens. Differences in cross-reactivity after

sequential vaccination have been previous demonstrated in the context of pandemic

virus vaccines where more highly cross-reactive antibodies were observed in subjects

primed with an A/Hong Kong/97 H5 vaccine and later boosted with an A/Vietnam/04

vaccine, who then subsequently mounted antibody responses recognizing both vaccine

strains, as well as a third H5 strain (A/Indonesia/05) not included in either vaccination

[30].

It is worth noting that we found a decrease in stalk specific antibodies when the

number of head antigenic sites was increased. This finding may help answer how the

ratio of head to stalk epitopes of HA affects the subdominance of the stalk antigenic

site. If in reality the head contains more antigenic sites than the stalk, the model predicts

that stalk-antigenic site response will be decreased. Our analysis suggests that stalk

specific antibody truly decreases with the addition of head antigenic sites, and it was not

that stalk-specific antibodies remain constant and only the relative amount compared to

the head is changed. It also suggests that the immunologic subdominance of the stalk

does not necessarily mean it is inherently less immunogenic, having implications for

targeting this domain in universal vaccination efforts.

Lastly, the work described here demonstrates the limitations with the current

vaccine selection process that relies only on antigenic and phylogenetic distances

between strains. Here, the shorter antigenic distance between SC18 and CA09

compared to BR07 and CA09 led to two different outcomes. For instance, the SC18

primed group had low titers to US77 after boost with CA09, while the BR07 primed

group had greater titers. Therefore, although the antigenic distance between CA09 and

US77 is fixed, previously exposed individuals produce antibody responses inconsistent

with these antigenic distance estimates. This suggests that serum samples are not

'impartial observers' of antigenic similarity and they are highly biased by their own

immune histories. This is an inherent challenge with the current influenza vaccine

approach and highlights the need to consider prior exposure histories when trying to

predict antibody specificities after vaccination.

In conclusion, our findings are consistent with other studies that point to negative

and positive interference as a mechanism affecting this enhancement of cross-reactivity

after sequential immunization[9-11,31]. Therefore, immunization regimens that can

relieve negative interference while increasing positive interference (especially to

conserved regions on an antigen) may act to broaden cross-reactive immunity to the

Influenza virus[15,32].

Acknowledgments

We thank the Center for Integrated Research Computing and the Health

Sciences Center for Computational Innovation for computational assistance and

resources. Thank you Carrie A. Anderson and Elaine Smolock for help with the

manuscript. Thank you Alan Perelson, Martin Zand, Juilee, Thakar, John Treanor, Jim

Miller, and Paige Lawrence for supervision of this work. Thank you Derek Smith for

helpful comments early on in this project. Thank you Anthony DiPiazza, Matt Brewer, and Patrick McCall for discussions on the project. Chimera proteins were a kind gift from Dr. Florian Krammer. Funding for this work was supported by the New York Influenza Center of Excellence NIH/NIAID/DMID, HHSN272201400005C and the University of Rochester Pulmonary training grant T32-HL066988.

REFERENCES

1. Chiu C, Wrammert J, Li G-M, McCausland M, Wilson PC, Ahmed R. Cross-reactive humoral responses to influenza and their implications for a universal vaccine. Ann N Y Acad Sci. 2013;1283: 13–21. doi:10.1111/nyas.12012

2. Russell CA, Jones TC, Barr IG, Cox NJ, Garten RJ, Gregory V, et al. Influenza vaccine strain selection and recent studies on the global migration of seasonal influenza viruses. Vaccine. 2008;26 Suppl 4: D31–4.

3. Garten RJ, Davis CT, Russell CA, Shu B, Lindstrom S, Balish A, et al. Antigenic and Genetic Characteristics of Swine-Origin 2009 A(H1N1) Influenza Viruses Circulating in Humans. Science. American Association for the Advancement of Science; 2009;325: 197–201. doi:10.1126/science.1176225

4. Krammer F, García-Sastre A, Palese P. Is It Possible to Develop a "Universal" Influenza Virus Vaccine? Toward a Universal Influenza Virus Vaccine: Potential Target Antigens and Critical Aspects for Vaccine Development. Cold Spring Harbor Perspectives in Biology. 2017. doi:10.1101/cshperspect.a028845

5. Andrews SF, Huang Y, Kaur K, Popova LI, Ho IY, Pauli NT, et al. Immune history profoundly affects broadly protective B cell responses to influenza. Science Translational Medicine. American Association for the Advancement of Science; 2015;7: –316ra192. doi:10.1126/scitranslmed.aad0522

6. Sangster MY, Baer J, Santiago FW, Fitzgerald T, Ilyushina NA, Sundararajan A, et al. B Cell Response and Hemagglutinin Stalk-Reactive Antibody Production in Different Age Cohorts following 2009 H1N1 Influenza Virus Vaccination. Clinical and Vaccine Immunology. 2013;20: 867–876. doi:10.1128/CVI.00735-12

7. Nachbagauer R, Choi A, Izikson R, Cox MM, Palese P, Krammer F. Age Dependence and Isotype Specificity of Influenza Virus Hemagglutinin Stalk-Reactive Antibodies in Humans. MBio. American Society for Microbiology; 2016;7: e01996–15. doi:10.1128/mBio.01996-15

8.   Tesini BL, Kanagaiah P, Wang J, Hahn M, Halliley JL, Chaves FA, et al. Broad
hemagglutinin-specific memory B cell expansion by seasonal influenza virus
infection reflects early-life imprinting and adaptation to the infecting virus. J Virol.
2019. doi:10.1128/JVI.00169-19

9.   Smith DJ, Forrest S, Ackley DH, Perelson AS. Variable efficacy of repeated
annual influenza vaccination. Proceedings of the National Academy of Sciences.
1999;96: 14001–14006.

10.   Ndifon W, Wingreen NS, Levin SA. Differential neutralization efficiency of
hemagglutinin epitopes, antibody interference, and the design of influenza
vaccines. Proceedings of the National Academy of Sciences. National Acad
Sciences; 2009;106: 8701–8706. doi:10.1073/pnas.0903427106

11.   Fonville JM, Wilks SH, James SL, Fox A, Ventresca M, Aban M, et al. Antibody
landscapes after influenza virus infection or vaccination. Science. 2014;346: 996–
1000. doi:10.1126/science.1256427

12.   Hoskins TW, Davies JR, Smith AJ, Miller CL, Allchin A. Assessment of inactivated
influenza-A vaccine after three outbreaks of influenza A at Christ's Hospital.
Lancet. 1979;1: 33–35. doi:10.1016/S0140-6736(79)90468-9

13.   Keitel WA, Cate TR, Couch RB, Huggins LL, Hess KR. Efficacy of repeated
annual immunization with inactivated influenza virus vaccines over a five year
period. Vaccine. 1997;15: 1114–1122.

14.   Zhang Y, Meyer-Hermann M, George LA, Figge MT, Khan M, Goodall M, et al.
Germinal center B cells govern their own fate via antibody feedback. J Exp Med.
2013;210: 457–464. doi:10.1084/jem.20120150

15.   Chaudhury S, Reifman J, Wallqvist A. Simulation of B cell affinity maturation
explains enhanced antibody cross-reactivity induced by the polyvalent malaria
vaccine AMA1. J Immunol. American Association of Immunologists; 2014;193:
2073–2086. doi:10.4049/jimmunol.1401054

16.   Anderson CS, McCall PR, Stern HA, Yang H, Topham DJ. Antigenic cartography
of H1N1 influenza viruses using sequence-based antigenic distance calculation.
BMC Bioinformatics. BioMed Central; 2018;19: 51. doi:10.1186/s12859-018-2042-
4

17.   Perelson AS, Oster GF. Theoretical studies of clonal selection: Minimal antibody
repertoire size and reliability of self-non-self discrimination. Journal of Theoretical
Biology. 1979;81: 645–670. doi:10.1016/0022-5193(79)90275-3

18.   Smith DJ, Forrest S, Hightower RR, Perelson AS. Deriving shape space
parameters from immunological data. Journal of Theoretical Biology. 1997;189:
141–150. doi:10.1006/jtbi.1997.0495

19.   Smith DJ. Derek J. Smith Thesis Dissertation. 1997. pp. 1–124.

20.   Chaudhury S, Reifman J, Wallqvist A. Simulation of B Cell Affinity Maturation
468   Explains Enhanced Antibody Cross-Reactivity Induced by the Polyvalent Malaria
469   Vaccine AMA1. The Journal of Immunology. 2014;193: 2073–2086.
470   doi:10.4049/jimmunol.1401054

471   21.   Pica N, Palese P. Toward a Universal Influenza Virus Vaccine: Prospects and
472   Challenges. Annu Rev Med. 2013;64: 189–202. doi:10.1146/annurev-med-
473   120611-145115

474   22.   Li G-M, Chiu C, Wrammert J, McCausland M, Andrews SF, Zheng N-Y, et al.
475   Pandemic H1N1 influenza vaccine induces a recall response in humans that
476   favors broadly cross-reactive memory B cells. Proc Natl Acad Sci USA. National
477   Acad Sciences; 2012;109: 9047–9052. doi:10.1073/pnas.1118979109

478   23.   Wrammert J, Koutsonanos D, Li GM, Edupuganti S, Sui J, Morrissey M, et al.
479   Broadly cross-reactive antibodies dominate the human B cell response against
480   2009 pandemic H1N1 influenza virus infection. Journal of Experimental Medicine.
481   2011;208: 181–193. doi:10.1084/jem.20101352

482   24.   Li Y, Myers JL, Bostick DL, Sullivan CB, Madara J, Linderman SL, et al. Immune
483   history shapes specificity of pandemic H1N1 influenza antibody responses. J Exp
484   Med. 2013;210: 1493–1500. doi:10.1084/jem.20130212

485   25.   Francis T. On the doctrine of original antigenic sin. 1960. doi:10.2307/985534

486   26.   Hancock K, Veguilla V, Lu X, Zhong W, Butler EN, Sun H, et al. Cross-reactive
487   antibody responses to the 2009 pandemic H1N1 influenza virus. N Engl J Med.
488   2009;361: 1945–1952. doi:10.1056/NEJMoa0906453

489   27.   Krause JC, Tumpey TM, Huffman CJ, McGraw PA, Pearce MB, Tsibane T, et al.
490   Naturally occurring human monoclonal antibodies neutralize both 1918 and 2009
491   pandemic influenza A (H1N1) viruses. J Virol. 2010;84: 3127–3130.
492   doi:10.1128/JVI.02184-09

493   28.   Thomas Francis J. On the Doctrine of Original Antigenic Sin. Proceedings of the
494   American Philosophical Society. 2012;: 1–8.

495   29.   Höpping AM, McElhaney J, Fonville JM, Powers DC, Beyer WEP, Smith DJ. The
496   confounded effects of age and exposure history in response to influenza
497   vaccination. Vaccine. 2016;34: 540–546. doi:10.1016/j.vaccine.2015.11.058

498   30.   Goji NA, Nolan C, Hill H, Wolff M, Noah DL, Williams TB, et al. Immune
499   responses of healthy subjects to a single dose of intramuscular inactivated
500   influenza A/Vietnam/1203/2004 (H5N1) vaccine after priming with an antigenic
501   variant. Journal of Infectious Diseases. 2008;198: 635–641. doi:10.1086/590916

31.   Ndifon W. A simple mechanistic explanation for original antigenic sin and its
alleviation by adjuvants. J R Soc Interface. 2015;12. doi:10.1098/rsif.2015.0627

32.   Wu JT, Wein LM, Perelson AS. Optimization of Influenza Vaccine Selection.
Operations Research. INFORMS; 2005;53: 456–476.
doi:10.2307/25146881?ref=search-
gateway:1675630d61600269146c9e3a07b04691

Supplimental Figure 1: Immune Responses To Priming Antigen
(A) Representative simultion of B cell, antibody, and antigen kinetics after priming for SC18 primed group
(B) or BR07 primed group. (C)Average antibody titers to priming antigen measured at day 30 post immuniza-
tion for each group. Two groups of 50 simulations were immunized with either (SC18 or BR07). (D)
Epitope-specific antibody titers to priming antigen for each epitope in the model, day 30 . (E) Average counts
of memory B cells specific to the priming antigen for each group measured at day 30. (F) Epitope-specific
memory B cell counts specific to priming antigen measured at day 30. (G) Epitope-specific antibody titers to
priming antigen and (H) memory B cells measured pre-boost (Day 365) .

Supplimental Figure 2
(A) Total antibody to CA09 measured day 30 after boosting with CA09. (B) Antigen clearance kinetics post-boosting with CA09 antigen for all subjects in each priming group.