Low-dose cadmium potentiates lung inflammatory response to 2009 pandemic H1N1

influenza virus in mice

Joshua D. Chandler[*], Xin Hu[*], Eunju Ko[†], Soojin Park[†], Jolyn Fernandes[*], Young-Tae Lee[†],

Michael L. Orr[*], Li Hao[*], M. Ryan Smith[*], David C. Neujahr[*], Karan Uppal[*], Sang-Moo Kang[†],

Dean P. Jones[*‡] and Young-Mi Go[*‡]

[*] Division of Pulmonary, Allergy and Critical Care Medicine, Emory University School of

Medicine, Atlanta, GA, 30322, [†] Center for Inflammation, Immunity and Infection, Georgia

State University, Atlanta, GA, 30303.

Running Head: Cadmium potentiates H1N1 lung inflammation in mice

[‡] Corresponding author: Whitehead Biomedical Research Building, 615 Michael St, Room 225

Atlanta, GA, 30322. Phone: 404-727-5984. Fax: 404-727-2974. E-mail: dpjones@emory.edu

Acknowledgements

Dr. Young-Mi Go and Dr. Dean P. Jones share equal senior authorship in this collaborative

research. We thank the Emory Integrated Genomics Core, Dana B. Barr PhD and Priya E.

D'Souza MPH for technical assistance. This study was supported by NIEHS Grant R01

ES023485 (DPJ and YMG), R21 ES025632 (DPJ and YMG), NIH S10 OD018006 (DPJ),

NIH/NIAID grants R01 AI105170 (SMK), R01 AI093772 (SMK), and R21 AI119366 (SMK),

and NHLBI F32 1F32HL132493 (JDC) and Cystic Fibrosis Foundation CHANDL16F0 (JDC).

Author Contributions

• Designed research – YMG, DPJ, SMK

• Performed research – JDC, JF, XH, YTL, EJK, SJP, DCN, MRS, MLO, LH

• Contributed new reagents or analytic tools – KU

• Analyzed data – JDC, YTL, SJP, EJK, JF, XH, MRS

• Wrote the paper – JDC, YMG, DPJ

Declaration of Competing Financial Interests

The authors declare they have no actual or potential competing financial interests.

Abbreviations

Cd, Cadmium; Ctl, control; Cys, cysteine; CySS, cystine; FACS, fluorescence-activated cell

sorting; FDR, False discovery rate; GSEA, Gene set enrichment analysis; H1N1, pandemic 2009

influenza A (H1N1) virus; HRM, High resolution metabolomics; ICP-MS, Inductively coupled

plasma mass spectrometer; interferon gamma, IFNγ; interleukin-1beta, IL-1β; LC, liquid

chromatograph; MS, mass spectrometer; *m/z*, mass to charge; RT, retention time; sPLS, sparse

partial least squares; Trx, thioredoxin.

Abstract

BACKGROUND: Cadmium (Cd) is a toxic, pro-inflammatory metal ubiquitous in the diet that

accumulates in body organs due to inefficient elimination. Many individuals exposed to dietary

Cd are also infected by seasonal influenza virus. The H1N1 strain causes mild to severe

pneumonia which can be fatal.

OBJECTIVES: To determine the influence of low-dose Cd exposure on inflammatory

responses to H1N1 influenza A virus.

METHODS: We exposed mice to low-dose (1 mg $CdCl_2$/l) Cd or vehicle (water) for 16 weeks

prior to infection with a sub-lethal dose of H1N1. Lung inflammation was assessed by

histopathology and flow cytometry. We used a combination of transcriptomics, metabolomics

and bioinformatics to determine the molecular associations of inflammatory cells important in

Cd-exacerbated responses.

RESULTS: Cd-treated mice had increased lung tissue inflammatory cells, including neutrophils,

monocytes, T lymphocytes and dendritic cells, following H1N1 infection. Lung genetic

responses to infection (increasing TNF-α, interferon and complement, and decreasing

myogenesis) were also exacerbated. Global correlations with immune cell counts, leading edge

gene transcripts and metabolites revealed that Cd increased correlation of myeloid immune cells

with pro-inflammatory genes, particularly interferon-γ, and metabolites in amino acid,

nucleobase, glycerophospholipid and vitamin B3 pathways.

DISCUSSION: Cd burden in mice increased inflammation in response to sub-lethal H1N1

challenge, which was coordinated by genetic and metabolic responses, and could provide new

targets for intervention against lethal inflammatory pathology of clinical H1N1 infection.

Keywords: Environmental safety, exposome, heavy metals, influenza A virus, public health.

Introduction

Recent studies of the environmental toxic metal cadmium (Cd) show that levels found in

human lungs potentiate pro-inflammatory signaling by the redox-sensitive transcription factor

NF-κB (Chandler et al. 2016b; Olszowski et al. 2012). Cd is a commercially important metal

used in NiCd batteries, plating, pigments and plastics (Kataranovski et al. 2009; Waalkes 2003;

Zhang et al. 2014). Earlier occupational toxicities from Cd have been largely eliminated by

occupational hygiene, and tobacco use remains as a major avoidable source of exposure. In non-

smokers, diet is the most important source of human Cd burden, on average representing about

half of the exposure of individuals with active tobacco use (Riederer et al. 2013; Satarug and

Moore 2004). Although Cd is poorly absorbed, there is no effective mechanism for elimination

so that the biological half-life in humans is over 10 years (Satarug and Moore 2004; Suwazono et

al. 2009). Our mechanistic studies show that low concentrations of Cd cause oxidation of

thiol/disulfide redox systems (Go et al. 2011; Go et al. 2013a; Go et al. 2014), disrupt the actin-

cytoskeleton regulation (Go et al. 2013a), cause translocation of thioredoxin-1 (Trx1) into the

nucleus (Go et al. 2013a), increase NF-κB activity and increase pro-inflammatory cytokines (Go

et al. 2011; Go et al. 2013b). Cd causes a dose-dependent oxidation of plasma cysteine/cystine

(Cys/CySS) redox state (E_h CySS, Y.-M. Go, unpublished), the major low molecular weight

thiol/disulfide system in plasma previously associated with pro-inflammatory signaling (Go and

Jones 2005, 2011). Additionally, studies with a transgenic NLS-thioredoxin-1 (Trx1) mouse

showed that increased nuclear Trx1 substantially potentiated adverse outcomes from H1N1

influenza infection (Go et al. 2011) and also increased oxidation of thiol/disulfide redox systems

(Go et al. 2011).

H1N1 and other influenza A strains are highly infectious (Liu et al. 2015; Xing et al.

2015) and cause variable severity of disease (Baigent and McCauley 2003a, b; Safronetz et al.

2011a, b; Taubenberger and Kash 2010a, b). Host genetics impact severity of response (Horby et

al. 2012, 2013), but inter-individual differences in sensitivity have mainly been linked to

demographic factors, such as age and general health (Gagnon et al. 2013; Wilson et al. 2014). In

most cases, the pandemic 2009 influenza A (H1N1) virus caused an uncomplicated respiratory

tract illness with symptoms similar to those caused by seasonal influenza viruses. However,

severe responses also occurred in a fraction of patients and were linked to increased levels of

pro-inflammatory cytokines, termed a "cytokine storm" (Gautam 2012; Liu et al. 2016), as well

as to excess activity of the complement system (Monsalvo et al. 2011). Cytokine storm included

high levels of IL-6 and was strongly associated with decreased lung function and death

(Bermejo-Martin et al. 2009; To et al. 2010).

The current study was designed to test whether Cd burden, at a level found in lungs of

non-smokers, enhanced severity of HIN1 influenza virus infection in a mouse model. Mice were

given water ad libitum with or without $CdCl_2$ (1 mg/l) for 16 weeks, and 10 days before sacrifice

each mouse was also intranasally exposed to either sub-lethal H1N1 or mock (sterile saline)

treatment. Lung histopathology and immune cell counts were used to evaluate inflammatory

responses and pathology. Gene expression analysis and high-resolution metabolomics were used

to identify Cd-dependent associations of inflammatory cells using the open-source software,

xMWAS (Uppal et al. 2018). Targeted analysis of IFNγ protein correlations with immune cells

was used to confirm *xMWAS* results.

Materials and Methods

Animals

All experimental protocols for animal studies were approved by Emory University and Georgia State University Internal Care and Use Committees. Experimental methods were carried out in accordance with the relevant guidelines and regulations. Male C57BL6 mice (n = 11-13 per group) aged 8 weeks (Jackson Labs, Bar Harbor, ME, USA) were maintained in clean facilities and given sterile-filtered drinking water with 1 mg/l $CdCl_2$ (Sigma-Aldrich, St. Louis, MO, USA) or vehicle for 16 weeks (including the 10-day duration of H1N1 infection). Food content of Cd (62 ± 1 ng/g food) was negligible compared to the Cd derived from water (Chandler et al. 2016b). Ten days prior to study completion, mice were inoculated with H1N1 (A/California/04/2009; dose of 0.6 x 10^3 pfu) under isoflurane anesthesia. Day 10 was chosen as the time point of maximal sub-lethal morbidity but also as a clinically relevant time point for severe and persistent cases. People infected with influenza are generally asymptomatic for 1-4 days, then exhibit symptoms for 3-7 days (https://www.cdc.gov/flu/professionals/acip/clinical.htm). Virus titer from mouse lung homogenate is expressed as the $\log_{10}$ of the 50% egg infective dose (EID_{50}) per ml homogenate.

For the main study, mice were weighed every 7 days until infection, then daily for the remaining 10 days. Mice were observed daily for other signs of influenza morbidity, e.g., a ruffled hair coat or labored breathing, and no apparent differences were observed between H1N1 and Cd+H1N1. Airway reactivity as measured by airway enhanced pause (PenH) was used to evaluate possible airway effects (Chandler et al. 2016a; Hwang et al. 2016). The PenH procedure does not require additional compound exposures (e.g., methacholine) which could impact the lung metabolome.

Cd measurement by Inductively-Coupled Plasma Mass Spectrometry (ICP-MS)

Lung tissue [114]Cd was assayed as previously described (Chandler et al. 2016b). ICP-MS

procedures conformed to accuracy (100 ± 10%) and precision standards (relative standard

deviation < 12%). Cd values were normalized to tissue mass. Human lung tissue samples were

collected in the Emory Transplant Center (Emory IRB protocol 000006248) and maintained

frozen at -80 ˚C until analysis.

Flow cytometry and staining

Immune cells in lung tissue were quantified by fluorescence-activated cell sorting (FACS). Lung

tissues were homogenized in DMEM and applied to Percoll gradients before washing with PBS.

Total viable cells were counted using a hemocytometer with 0.4% trypan blue for dye exclusion.

A subset of cells for FACS analysis was treated with anti-surface marker antibodies including

CD11b (M1/70), CD45 (30-F11), B220 (RA3-6B2), Siglec F (E50-2440) (BD Pharmingen,

Franklin Lakes, NJ, USA); CD11c (N418), F4/80 (BM8) (eBioscience, Thermo Fisher Scientific,

Waltham, MA, USA); and CD103 (2E7), Ly6c (HK1.4), MHC-II (M5/114.15.2), CD3 (17A2),

CD4 (GK1.5), and CD8 (53-5.8) (BioLegend, San Diego, CA, USA). Data were acquired using

an LSR-II/Fortessa (Becton-Dickinson, Franklin Lakes, NJ, USA) and analyzed using FlowJo v.

10.1 (Tree Star, Inc., Ashland, OR, USA). $CD11b^+$ DCs were gated as those with low CD103

expression, and vice-versa; we did not analyze a mixed population ($CD11b^+$/$CD103^+$). Absolute

counts were expressed from the percentage from FACS analysis multiplied by viable cell counts

obtained with a hemocytometer.

Cytokine and chemokine assays

IFNγ was measured as previously described (Ko et al. 2016; Lee et al. 2015; Lee et al. 2016).

Briefly, right lung lobe extracts were prepared by grinding and filter straining prior to IFNγ

measurement according to manufacturer instructions (eBioscience, Thermo Fisher Scientific,

Waltham, MA, USA).

Analysis of glutathione and cysteine

Glutathione (GSH), oxidized glutathione (GSSG), cysteine (Cys) and oxidized form of Cys

(cystine, CySS) in lung tissue and plasma were analyzed by HPLC with fluorescence detection

(Go and Jones 2005; Jones et al. 2002; Jones and Liang 2009). Protein assay (bicinchonicic acid)

was used to standardize the results to intracellular concentrations. Redox states of GSH/GSSG

(E_h[GSSG]) and Cys/CySS (E_h[CySS]) were calculated using the Nernst equation (Go and Jones

2005; Jones et al. 2002; Jones and Liang 2009).

Histopathology

The lung collected at 10 days after challenge was fixed with 10% normal buffered formalin

(NBF) in PBS. After fixation NBF tissues were transferred to 70% alcohol and processed to

paraffin using a Shandon Excelsior AS (Thermo Fisher Scientific). A standard program of

dehydration was carried out in ethanol 70%, 80%, 90%, 95%, 100%, 100% 1 hour each,

followed by xylenes twice for 30 min each. Finally, the tissue was soaked in paraffin for 30 min

and repeated once, then held in the second paraffin bath for 1 hour. Left lung tissue sections (4

mice per group) were stained with hematoxylin and eosin (H&E) to assess histopathological

changes as described (Ko et al. 2016). Images were acquired by Axiovert 100 (Zeiss,

Oberkochen, Germany) at 100X magnification, and inflammation was blindly scored as

previously described (Ko et al. 2016).

Transcriptomics

20-25 mg of lung tissue was stored in RNAlater (Qiagen, Hilden, Germany) at -80 ˚C prior to

lysis and RNA isolation using the miRNEasy kit (Qiagen, Hilden, Germany). 250 ng of RNA

was prepared and hybridized to MoGene ST 2.0 arrays (Affymetrix, Thermo Fisher Scientific,

Waltham, MA, USA). CEL files were converted to a Robust-Multi Array (RMA) and loaded in

the gene set enrichment analysis (GSEA) Java applet (v. 2.2; Broad Institute) to test gene set

enrichment. Annotations were made using MSigDB file 'MoGene_2_0_st.chip', accessed from

the GSEA applet. Redundant annotations were collapsed using maximum intensities.

High resolution metabolomics (HRM)

Lung tissue metabolites were extracted from 20-30 mg tissue in 15 µl/mg of 2:1

acetonitrile:water containing internal standards per mg tissue (Chandler et al. 2016a). Tissues

were sonicated, vortexed and incubated on ice for 30 minutes before centrifugation at 16,000 g

and 4 ˚C. Five µl of supernatant was injected on a Dionex Ultimate 3000 LC system coupled to

QExactive High Field mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA); each

analysis was performed with three technical replicates. Chromatographic separation was

achieved with Accucore HILIC chromatography (100 x 2.1 mm, 2.6 µm, 80Å; Thermo Fisher

Scientific, Waltham, MA, USA). Untargeted data were extracted with XCMS (Smith et al.

2006) and *xMSanalyzer* (Uppal et al. 2013), recovering 2,956 features (high resolution *m/z* peak

paired with retention time). Feature intensities were log$_2$-transformed prior to integrative

network analysis using *xMWAS* (Uppal et al. 2018). The pathway enrichment analysis by

mummichog software (Li et al. 2013) was used to identify metabolic pathways enriched in mass

spectral features obtained from *xMWAS* analyses (Uppal et al. 2018); this approach was

previously validated for relevant metabolites by coelution with pure standards and MS^2 spectra

(Frediani et al. 2014; Uppal et al. 2017).

Data integration and network analysis

Immune cells (Cd+H1N1 significantly different from H1N1 alone), gene transcripts (from

leading edges of gene sets significantly different between Cd+H1N1 and H1N1 alone) and lung

m/z features (all) were selected for *xMWAS* analysis to identify Cd-dependent associations during

H1N1 infection. *xMWAS* (Uppal et al. 2018) uses the sparse partial least squares (sPLS)

regression method and the *network( )* function implemented in R package mixOmics (Gonzalez

et al. 2012; Le Cao et al. 2009) for generating the association matrices between datasets, with

Student's t-test used to evaluate the statistical significance of pairwise association scores.

Statistics

Data are presented as the mean and standard error of the mean. One-way ANOVA with Holm-

Sidak's post-test was used to test statistical significance at a significance level of $p < 0.05$. The

GSEA applet (v. 2.2; Broad Institute) produced all statistics for gene set enrichment

(Subramanian et al. 2005). Genes were ranked according the "Signal2Noise" statistic with a

significance threshold of false discovery rate (FDR)-adjusted $q < 0.05$. In *xMWAS*, $p < 0.05$ was

the significance threshold for correlations.

Data Sharing

In addition to Supplementary Figures, data corresponding to the Figures and Supplementary

Figures have been provided as numbered Supplementary Files. Datasets corresponding to

metabolomics and transcriptomics used for *xMWAS* analysis have also been included (see List of

Supplementary Files).

Results

Selection of sub-lethal H1N1 dose

Selection of H1N1 virus titer for mouse infection was critical to achieve maximal sub-lethal
morbidity as evaluated by body mass decrease and increased lung inflammation at Day 10 post-
infection. We compared the effects of 5 different virus doses on severity of lung infection,
measuring body weight change over 10 days and observing lung inflammatory histopathology at
Day 10 (**Supplementary Figure 1A**). The results showed that the virus doses of 0.6-1.2 x 10^3
pfu caused sustained body weight loss up to Day 10 post-infection. Lower doses exhibited body
weight recovery after Day 8 or 9. Doses of 0.6-1.2 x 10^3 pfu also caused extensive inflammatory
peribronchiolar histopathology (**Supplementary Figure 1B**). We reasoned that the lowest dose
that caused sustained body weight loss (0.6 x 10^3 pfu) would enhance the ability to detect
second-hit effects of Cd and selected this dose for the main study involving Cd pre-treatment
before infection. Consistent with previous low-dose Cd studies (Thijssen et al. 2007), Cd pre-
treatment had no effect on body mass or weight gain compared to vehicle treatment in the weeks
prior to infection (**Supplementary Figure 1C**).

Impact of Cd treatment on pathology of H1N1 infection

Four groups of mice were maintained for the treatment period, half receiving vehicle (sterile
drinking water) and half receiving low-dose Cd. At 10 days prior to termination, mice were
intranasally exposed either to mock treatment with sterile saline (Ctl, Cd groups) or 0.6 x 10^3 pfu
of H1N1 (H1N1, Cd+H1N1). We examined lung histopathology to determine the extent of
inflammation in airway (**Figure 1B**), blood vessels (**Figure 1C**) and interstitial spaces (**Figure
1D**). As expected, H1N1 infection caused inflammation to increase compared to control in all

compartments (H1N1: airways, $p = 0.0016$; blood vessels, $p = 0.0095$; interstitial spaces, $p =$

0.0127; **Figure 1B-D**). Cd treatment alone did not significantly increase histopathological

scoring, and combination of Cd treatment with H1N1 did not significantly increase

histopathology score compared to H1N1 with vehicle treatment. However, the combination of

Cd+H1N1 had consistently lower p-values compared to control than H1N1 alone, for each

histological compartment (Cd+H1N1: airways, $p = 2.6 \times 10^{-4}$; blood vessels, $p = 0.0011$;

interstitial spaces, $p = 9.8 \times 10^{-4}$; **Figure 1B-D**). Mice treated with Cd exhibited increased lung

Cd burden compared to vehicle treatment (**Supplementary Figure 1D**). To compare our model

to human lung Cd, we used the same analytical methods to measure Cd in available lung samples

from both smokers (S) and non-smokers (NS) (**Supplementary Figure 1D**). Cd content (pmol

Cd/g wet tissue, mean ± SE) in Cd-treated mouse lung tissue [Cd (n = 7), 241 ± 32; Cd+H1N1 (n

= 7), 220 ± 29] overlapped the range of non-smokers (n = 5, 157 ± 29) and lower portion of the

range for smokers (n = 14, 568 ± 137). Lung tissue Cd content in vehicle-treated mice was lower

than in Cd-treated mice [Ctl (n = 7), 137 ± 70; H1N1 (n = 8), 24 ± 7]. Treatment with Cd had no

effect on infection-mediated body weight loss (Day 10 percentage of Day 0 body weight at

inoculation: H1N1, 78 ± 2%; Cd+H1N1, 79 ± 2%, **Supplementary Figure 1E**), infection-

mediated airway enhanced paused (PenH; mean Day 10 results as percentage of Day 0 value:

H1N1, 700 ± 85%; Cd+H1N1, 669 ± 56%, **Supplementary Figure 1F**) or lung virus titer

(H1N1, $10.0 ± 0.3 \log_{10}[EID_{50}/ml]$; Cd+H1N1, $9.6 ± 0.4 \log_{10}[EID_{50}/ml]$, **Supplementary**

Figure 1G).

Cd increased H1N1-mediated lung tissue inflammatory cells

Specific cellular contributions to lung inflammation were quantified using flow cytometry

(**Figure 2**). Inflammatory immune cells were increased in lung tissue of Cd+H1N1 compared to

H1N1, including neutrophils ($p = 0.0002$), monocytes ($p = 0.0011$), CD11b$^+$ dendritic cells

(DCs; $p = 0.0001$), CD103$^+$ DCs ($p = 3\text{x}10^{-6}$), plasmacytoid DCs (pDCs) ($p = 7\text{x}10^{-6}$), CD4$^+$ T

lymphocytes ($p = 0.0001$) and CD8$^+$ T lymphocytes ($p = 0.0019$) (**Figure 2A-G**). In contrast, we

did not observe significant differences in the numbers of lung tissue alveolar macrophages or

eosinophils when comparing Cd+H1N1 and H1N1 (**Supplementary Figure 2**).

Impact of Cd on glutathione and cysteine in lung tissue and plasma

We measured the steady-state redox potentials of glutathione ($E_h[GSSG]$) and cysteine

($E_h[CySS]$) in plasma and lung tissue to establish whether low dose Cd treatment impacted these

thiol-disulfide antioxidant systems in the context of sub-lethal influenza infection. H1N1

treatment did cause significant oxidation of $E_h[CySS]$ in lung tissue (**Supplementary Figure**

3B), while no effects were observed for lung glutathione or for plasma (**Supplementary Figure**

3A,C-D). No potentiation of the effect noted in lung tissue $E_h[CySS]$ by low dose Cd was

observed.

Cd enhanced pro-inflammatory gene set expression in H1N1-infected mice

To determine whether Cd impacted gene expression responses to H1N1 infection, we used

Affymetrix MoGene ST 2.0 arrays followed by Gene Set Enrichment Analysis (GSEA)

(Subramanian et al. 2005) to study effects on transcript abundances among the treatment groups,

comparing overrepresentation of gene sets between experimental groups (n = 5). We selected

GSEA gene sets with $p < 0.05$ and FDR $q < 0.05$ for gene set enrichment. Seven gene sets

passed these criteria when comparing Cd+H1N1 and H1N1 (**Table 1**), including six that were

overrepresented in Cd+H1N1 compared to H1N1 alone (1. allograft rejection, 2. complement

signaling, 3. inflammatory response, 4. IFNα response, 5. IFNγresponse, 6. TNF-α signaling via

NF-κB; each comprised predominantly of pro-inflammatory genes). The seventh gene set

(myogenesis; comprised of cytoskeletal subunit genes) was overrepresented in H1N1 compared

to Cd+H1N1. Each of the six gene sets overrepresented in Cd+H1N1 compared to H1N1 was

also overrepresented in either of the H1N1-infected groups compared to mock-treated groups. In

contrast, the myogenesis gene set was overrepresented in Ctl, Cd and H1N1 groups compared to

Cd+H1N1.

The leading edge genes from each of the gene sets in **Table 1** were selected for global

correlation analysis by *xMWAS* (Uppal et al. 2018). After accounting for redundancies across

gene sets, this yielded 293 leading edge genes (**Supplementary File S3**). Two-hundred and

twenty of these genes belonged to the pro-inflammatory gene sets in Table 1 that were increased

in Cd+H1N1 compared to H1N1. Pro-inflammatory leading edge genes were predominantly

comprised of cytokines, chemokines, receptors of cell activation and signaling proteins. The

highest ranked leading edge gene from these sets (contributing most to their enrichment) was

'interleukin 6 (interferon, beta 2)' (*Il6*). The remaining 73 leading edge genes belonged to the

myogenesis gene set with cytoskeletal subunit and regulatory genes including 'actin, alpha 1'

(*Acta1*)*,* 'creatine kinase, muscle' (*Ckm*) and 'collagen 1A1' (*Col1a1*). The lowest ranked gene

from this gene set (contributing most to the "negative enrichment" of the gene set in Cd+H1N1

compared to H1N1 alone; **Table 1**) was 'myosin, light chain 7, regulatory' (*Myl7*).

Global correlation analysis of Cd effect on lung response to H1N1 influenza infection

To reveal organization of the network structure and pinpoint molecules critical to Cd-altered

lung function, global correlations were made for immune cell counts, leading edge gene

transcripts and metabolites using *xMWAS* (Uppal et al. 2018). *xMWAS* is a data-driven

integration and network analysis tool which uses community detection to improve understanding

of complex interactions. As input, we used the 293 leading edge genes from Table 1 gene sets

(**Supplementary File S3**) paired with inflammatory cells as measured in **Figure 2A-G**

(**Suppelementary File S4**) and lung tissue metabolites measured by HRM (total of 2,956

metabolic features; **Supplementary File S5**) to produce experimental group-specific association

networks, with the explicit goal to compare the respective associations of the H1N1 and

Cd+H1N1 networks (**Figure 3**; n = 5). Student's t-test was used to determine the association

score (absolute value) threshold of 0.88 (determined by *xMWAS* software based on $p < 0.05$

criteria and n = 5 samples per group). The resulting network of H1N1 was comprised of 426

metabolic features, 160 gene transcripts and 7 immune cell types, while the network of

Cd+H1N1 was comprised of 478 metabolic features, 158 gene transcripts and 6 immune cell

types (**Figure 3A**; full network membership given in **Supplementary File S6**).

To quantify the relative importance of "nodes" (cells, transcripts or metabolites) in each

network, *xMWAS* calculates an eigenvector centrality score that can be compared across the

groups (**Figure 3B**). Respective delta centrality scores showed that monocytes (+0.419), CD11b$^+$

DCs (+0.380) and neutrophils (+0.301) had increased centrality in the Cd+H1N1 network

compared to H1N1 (**Figure 3B**). Notably, these are all myeloid cell populations, so this result

indicates an increase in myeloid cell importance in the Cd+H1N1 model. This result can also be

qualitatively detected in the model visualizations, in which neutrophils, a major myeloid cell

type, shift from peripheral (H1N1) to central (Cd+H1N1) network locations between the two

experimental conditions, whereas CD4$^+$ and CD8$^+$ T lymphocytes, canonical lymphoid cells, do

the opposite (**Figure 3A**). Additionally, CD103$^+$, a lymphoid dendritic cell sub-population, was

not found in the Cd+H1N1 network (indicating no strong correlations were made with this cell

type in the network (**Figure 3A**). Thus, Cd treatment altered biological interactions of immune

cells with gene transcript and metabolites in lung tissue following sub-lethal H1N1 infection,

causing myeloid cell associations to be strengthened.

Previous research showed that influenza-induced IFNγ restricts protective innate

lymphoid cell group II function in mouse lung following challenge with H1N1 (Califano et al.

2018). We therefore performed targeted tests to determine whether correlations with IFNγ

protein, measured in lung tissue homogenate by ELISA, differed for myeloid and lymphoid cells

in Cd+H1N1 compared to H1N1. Results showed that each myeloid cell count correlated with

IFNγ only in Cd+H1N1 mice [monocytes (Cd+H1N1, $r = 0.97$, $p = 3\times10^{-7}$; H1N1, $r = 0.21$, $p = $

0.4862; **Figure 3C**), CD11b$^+$ DCs (Cd+H1N1, $r = 0.92$, $p = 2\times10^{-5}$; H1N1, $r = -0.05$, $p = 0.8657$;

Figure 3D) and neutrophils (Cd+H1N1, $r = 0.89$, $p = 9\times10^{-5}$; H1N1, $r = 0.22$, $p = 0.4773$;

Figure 3E)]. Additionally, pDC exhibited a modest correlation with IFNγ in Cd+H1N1 not

observed in H1N1 (Cd+H1N1, $r = 0.69$, $p = 0.0122$; H1N1, $r = -0.12$, $p = 0.6869$), while none of

the lymphoid cell populations tested (CD8$^+$, CD4$^+$, CD103$^+$ DCs) exhibited significant

correlations with IFNγ in either H1N1 or Cd+H1N1 groups (**Supplementary Figure 4**). The

results confirm the global correlation network analysis showing that Cd burden increases

centrality of myeloid cells with pro-inflammatory pathways following H1N1 infection.

Furthermore, the strength of the IFNγ correlations in Cd+H1N1 across cell types followed the

same rank order as delta centrality (**Figure 3B**). These results implicate IFNγ signaling as highly

relevant to increased inflammation and myeloid cell dominance caused by Cd treatment.

Gene sets and metabolic pathways associated with Cd effect on immune cell responses to

H1N1

We further examined the molecular composition of each of the correlation networks to determine

which gene transcripts and metabolic pathways were central in each group. The overall network

composition of Cd+H1N1 and H1N1 overlapped by >50% in both leading edge genes and

metabolic features so we focused on the most divergent relationships between Cd+H1N1 and

H1N1 by analyzing genes and metabolites with absolute delta eigenvector

centrality, $|\text{Centrality}_{Cd+H1N1} - \text{Centrality}_{H1N1}|, \geq 0.1$. (If a gene or metabolic feature was not

present in one of the correlation networks, its eigenvector centrality was considered 0.) Results

for the transcriptome data showed 83 leading edge genes of ≥ 0.1 absolute delta centrality were

present in the Cd+H1N1 and H1N1 networks (**Supplementary File S8**), with the IFNγ response

gene set (27.4% of total gene counts; **Figure 4A**) being most represented. IFNα response

(17.8%) followed, and myogenesis was least represented (5.2%). The IFNγ transcript, *Ifng*, had

differential centrality of +0.435 for Cd+H1N1 network compared to H1N1. The transcript with

most positive change for gene centrality (1.0) was immediate early response 2, *Ier2*, encoding a

DNA-binding protein, and the transcript with most negative change for gene centrality (-0.88)

was Dnaj (Hsp40) homolog, subfamily B, member 4, *DnajB4*, a chaperone protein targeted to the

plasma membrane. Thus, the transcript data are consistent with Cd effects occurring through

IFNγ-dependent signaling mechanisms and also involving more complex interactions of nuclear

as well as plasma membrane systems.

Prior research has shown that low-dose Cd has widespread effects on metabolism,

especially related to oxidative stress, mitochondrial β-oxidation of fatty acids and amino acid

metabolism (Go et al. 2015). We analyzed 165 metabolic features with delta eigenvector

centrality ≥ 0.1 (**Supplementary File S9**), using the metabolic pathway analysis software,

mummichog 1.0.10 (Li et al. 2013). The top metabolic pathway was methionine and cysteine

metabolism (**Figure 4B**). The largest positive centrality change for a single metabolite was for

S-adenosylhomocysteine (+0.415), an intermediate in methylation processes linked to

methionine. Other pathways included vitamin B3 (niacin/NAD), urea cycle and

glycerophospholipid metabolism, all of which are connected to mitochondrial function and

known to be perturbed by Cd. Other metabolic pathways included amino acid, pyrimidine and

purine pathways that have previously been associated with inflammation and/or cell proliferation

and turnover (**Figure 4B**). Consistent with this, the metabolite with the most negative centrality

change was adenine (-0.953). The results are consistent with Cd potentiation of inflammation to

H1N1 infection through widespread effects of low-dose Cd exposure on oxidative stress,

mitochondrial function and regulation of immune cell populations through IFNγ.

Discussion

A previous study showed that blood concentrations of toxic metals including Cd were significantly higher in H1N1 cases with pneumonia versus pneumonia not caused by H1N1, and two H1N1 cases having highest Cd concentrations did not recover from illness (Moya et al. 2013). Although limited by the small number of cases, the study suggested that Cd could be a risk factor for inflammatory response and enhanced severity of H1N1 virus infection-induced respiratory diseases. In the present study using a mouse model of H1N1 infection, we show that lung Cd burden, at a value present in human lung, exacerbated inflammation and lung injury from H1N1 infection in mice. Comparing Cd+H1N1 to H1N1-infected mice, Cd increased the Day 10 post-infection lung burden of myeloid (monocytes, neutrophils), lymphoid (CD4$^+$ and CD8$^+$ T cells) and multi-origin (DCs) bone marrow lineages of immune cells. Integrated network analysis of omics data further showed that Cd burden shifted the immune response from one centered on both lineages to one with myeloid cells (including CD11b$^+$ DCs) as a dominant central hub. Thus, high Cd burden appears to contribute to worse outcomes following H1N1 infection by increasing the immune response with suppression of lymphoid and enhancement of myeloid cell participation in response to infection.

The global correlation network analysis of immune cells with gene transcript and metabolite abundances revealed the centrality of myeloid cells (monocytes, CD11b$^+$ DCs [considered a myeloid DC subtype (Chistiakov et al. 2015)] and neutrophils) in the Cd+H1N1 group. Transcriptional responses suggest that Cd primes myeloid cells for tight coordination with IFNγ signaling. The finding that a unique lysosomal thiol reductase, IFNγ-inducible lysosomal thiol reductase, controls conditions for degradation of proteins to small peptides in lysosomes (Hastings and Cresswell 2011; Phan et al. 2002), raises the possibility that Cd specifically impacts IFNγ-dependent functions through interaction with, or oxidation of, the

thioredoxin-like active site involved in disulfide reduction. Low dose Cd causes oxidation and

disruption of the actin-cytoskeleton (Go et al. 2013a), suggesting that targeted effects on a

critical signaling system could occur even without effects on major GSH or thioredoxin systems.

Effects of low-dose Cd on the actin-cytoskeleton system triggers translocation of Trx1 to cell

nuclei (Go et al. 2013a) and increase NF-κB activation (Go et al. 2011; Go et al. 2013b) and

monocyte adhesion (Go and Jones 2005). Thus, these processes also could contribute to higher

immune cell counts in Cd+H1N1 compared to H1N1. The GSEA results showing effects of Cd

on pro-inflammatory gene sets (**Table 1**), as well as increased cytokine levels and immune cell

counts, support such a Cd-dependent mechanism.

Alternative mechanisms for Cd effects on inflammatory pathology of H1N1 could

involve cytoskeletal functions, myogenesis and elastogenesis in the inflamed and injured lungs.

It is noteworthy that the combination of Cd pre-treatment with H1N1 infection was the

experimental condition to cause a significant negative enrichment, or decrease in overall

transcript levels, of the myogenesis gene set. Cytoskeletal regulation by tight junctions and

adherens junctions is a key element of airway permeability to infiltrating immune cells (Rezaee

and Georas 2014). We previously found that low dose Cd disruption of actin cytoskeleton

regulation (Go et al. 2013a; Hu et al. 2017) triggered nuclear translocation of thioredoxin-1 and

caused cell death (Go et al. 2013b). A study by Desouza *et al*. also showed that actin-binding

proteins contribute to cell death, and actin dynamics plays a key role in regulating apoptosis

signaling (Desouza et al. 2012). Studies of low dose Cd effects on actin cytoskeleton and cell

death in fibroblasts, HeLa and HT-29 cells [(Go et al. 2013a; Hu et al. 2017); Y.-M. Go, D.P.

Jones, unpublished] suggest that this could be a general cell response and contribute to barrier

dysfunction. Thus, Cd could also contribute to severity of response through mechanisms

involving impaired myogenesis, elastogenesis, increased cell death and failure to clear cell debris

in the inflamed and injured lungs.

In contrast to the coherent picture provided by Cd effects on proinflammatory signaling,

immune cell accumulation in the lungs and amplified effects on myeloid compared to lymphoid

cells, metabolic effects associated with Cd effects were diverse and complex; the results

suggested more of a generalized adaptive response than a causal role in exacerbation of

inflammation. The top pathway from pathway enrichment analysis was for sulfur-containing

amino acids which serve to support GSH synthesis and antioxidant activities. This would appear

to be a protective mechanism, but maladaptive effects could also occur at specific subcellular

sites. Pathway enrichment analysis also showed that pathways of several amino acids,

nucleosides and phospholipids were overrepresented in the networks with differential

eigenvector centrality ≥ 0.1 (**Figure 4B**). Overall, these results suggest that Cd burden causes a

complex array of metabolic responses.

Inhaled Cd causes acute pneumonitis and lung edema (Nemery 1990), and intraperitoneal

injection of Cd causes lung inflammation in rats (Kataranovski et al. 2009). However, chronic

ingestion of Cd at levels mimicking those derived from the human diet has not previously been

directly linked to inflammatory lung disease. The present research with 16 weeks of Cd in

drinking water showed little evidence for inflammation without the second stimulus of H1N1

infection. Thus, Cd accumulation with advanced age (> 65 y), tobacco use, metabolic syndrome

and/or obesity, may create a sub-clinical state poised to over-react to influenza virus infection-

induced inflammation, which may in turn lead to worse morbidity or mortality (Choi et al. 2014;

Fiore et al. 2011; Lee and Kim 2016; Olsson et al. 2002; Ramsey and Kumar 2013). For future

research, the dose of virus used in our study permits recovery to study later complications such

as fibrosis caused by low-dose Cd (Hu et al. 2017).

526   Finally, in light of common consumption of foods with relatively high Cd content

527 (potatoes, cereals), the results indicate that studies in humans are needed to evaluate whether Cd

528 burden is associated with severity of response to influenza infection. Cd burden can be readily

529 evaluated by determination of Cd/creatinine level in urine (Haddam et al. 2011; Larsson and

530 Wolk 2016). While there are no approved strategies to prevent Cd accumulation or decrease its

531 burden, mechanisms discussed above could be targeted to prevent Cd-dependent potentiation of

532 influenza severity. For instance, the mechanism for nuclear Trx1 translocation could be targeted

533 to block excessive activation of transcriptional responses. Alternatively, Txnip, a vitamin D_3-

534 binding protein that controls nuclear Trx1 activity (Wang et al. 2002; Yamawaki et al. 2005),

535 could provide a target for therapeutic manipulation. Trp14 catalyzes reduction of cystine

536 (oxidized form of cysteine) (Pader et al. 2014) and could provide a target to protect against

537 oxidation of proteins and downstream signaling and recruitment of immune cells. Cd also

538 interacts with trace elements including selenium, zinc and iron (Gocmen et al. 2000; Riederer et

539 al. 2013), providing additional strategies for supplement use (Choi et al. 2014; Moya et al. 2013).

540   In summary, this study showed that mice given Cd at a level achieved in humans from

541 dietary intake had worse lung inflammation and inflammatory pathology after H1N1 influenza

542 infection, in association with enhanced myeloid-dominant pro-inflammatory gene and metabolic

543 responses. Studies are needed to evaluate whether Cd burden contributes to influenza severity in

544 humans. Combined analyses of phenotypic measures with transcriptome and metabolome

545 analyses enable global surveillance of complex systems responses to identify candidate

546 mechanisms and potential therapeutic targets.

List of Supplementary Figures and Data Files

Supplementary Figure 1. Virus dose response, response to Cd pre-treatment, and body mass,

PenH and virus titer endpoints of main study.

Supplementary Figure 2. Flow cytometry-quantified eosinophils and alveolar macrophages.

Supplementary Figure 3. Steady-state redox potentials of glutathione and cysteine in lung

tissue and plasma.

Supplementary Figure 4. Correlations of additional cell types with IFNγ.

Supplementary File S1. Corresponds to Figure 1 histopathology data.

Supplementary File S2. Corresponds to Figure 2 flow cytometry data.

Supplementary File S3. Leading edge genes selected for *xMWAS* analysis.

Supplementary File S4. Flow cytometry results selected for *xMWAS* analysis.

Supplementary File S5. Metabolomics dataset selected for *xMWAS* analysis.

Supplementary File S6. Corresponds to Figure 3 *xMWAS* analysis data and IFNγ correlations.

Supplementary File S7. Corresponds to Figure 4 enrichment tests data.

Supplementary File S8. Leading edge genes used for proportion testing following *xMWAS*.

Supplementary File S9. Metabolite features used for *mummichog* testing following *xMWAS*.

Supplementary File S10. Full Robust Multi-Array dataset from lung tissue Affymetrix array.

Supplementary File S11. Corresponds to Figure S1 data.

Supplementary File S12. Corresponds to Figure S2 data.

Supplementary File S13. Corresponds to Figure S3 data.

Supplementary File S14. Corresponds to Figure S4 data.

References

Baigent SJ, McCauley JW. 2003a. Influenza type a in humans, mammals and birds:
Determinants of virus virulence, host-range and interspecies transmission. Bioessays
25:657-671.

Baigent SJ, McCauley JW. 2003b. Influenza type a in humans, mammals and birds:
Determinants of virus virulence, host-range and interspecies transmission. Bioessays
25:657-671.

Bermejo-Martin JF, Ortiz de Lejarazu R, Pumarola T, Rello J, Almansa R, Ramirez P, et al.
2009. Th1 and th17 hypercytokinemia as early host response signature in severe
pandemic influenza. Crit Care 13:R201.

Califano D, Furuya Y, Roberts S, Avram D, McKenzie ANJ, Metzger DW. 2018. Ifn-gamma
increases susceptibility to influenza a infection through suppression of group ii innate
lymphoid cells. Mucosal Immunol 11:209-219.

Chandler JD, Hu X, Ko EJ, Park S, Lee YT, Orr M, et al. 2016a. Metabolic pathways of lung
inflammation revealed by high-resolution metabolomics (hrm) of h1n1 influenza virus
infection in mice. Am J Physiol Regul Integr Comp Physiol 311:R906-R916.

Chandler JD, Wongtrakool C, Banton SA, Li S, Orr ML, Barr DB, et al. 2016b. Low-dose oral
cadmium increases airway reactivity and lung neuronal gene expression in mice. Physiol
Rep 4.

Chistiakov DA, Sobenin IA, Orekhov AN, Bobryshev YV. 2015. Myeloid dendritic cells:
Development, functions, and role in atherosclerotic inflammation. Immunobiology
220:833-844.

Choi SM, Jeong YJ, Park JS, Kang HJ, Lee YJ, Park SS, et al. 2014. The impact of lifestyle
behaviors on the acquisition of pandemic (h1n1) influenza infection: A case-control
study. Yonsei Med J 55:422-427.

Desouza M, Gunning PW, Stehn JR. 2012. The actin cytoskeleton as a sensor and mediator of
apoptosis. Bioarchitecture 2:75-87.

Fiore AE, Fry A, Shay D, Gubareva L, Bresee JS, Uyeki TM. 2011. Antiviral agents for the
treatment and chemoprophylaxis of influenza --- recommendations of the advisory
committee on immunization practices (acip). MMWR Recomm Rep 60:1-24.

Frediani JK, Jones DP, Tukvadze N, Uppal K, Sanikidze E, Kipiani M, et al. 2014. Plasma
metabolomics in human pulmonary tuberculosis disease: A pilot study. PLoS One
9:e108854.

Gagnon A, Miller MS, Hallman SA, Bourbeau R, Herring DA, Earn DJ, et al. 2013. Age-specific
mortality during the 1918 influenza pandemic: Unravelling the mystery of high young
adult mortality. PLoS One 8:e69586.

Gautam M. 2012. Endothelial cells as regulators of cytokine storms during influenza infection.
Thorax 67:617-617.

Go YM, Jones DP. 2005. Intracellular proatherogenic events and cell adhesion modulated by
extracellular thiol/disulfide redox state. Circulation 111:2973-2980.

Go YM, Jones DP. 2011. Cysteine/cystine redox signaling in cardiovascular disease. Free Radic
Biol Med 50:495-509.

Go YM, Kang SM, Roede JR, Orr M, Jones DP. 2011. Increased inflammatory signaling and
lethality of influenza h1n1 by nuclear thioredoxin-1. PLoS One 6:e18918.

Go YM, Orr M, Jones DP. 2013a. Actin cytoskeleton redox proteome oxidation by cadmium.
Am J Physiol Lung Cell Mol Physiol 305:L831-843.

Go YM, Orr M, Jones DP. 2013b. Increased nuclear thioredoxin-1 potentiates cadmium-induced
cytotoxicity. Toxicological sciences 131:84-94.

Go YM, Roede JR, Orr M, Liang Y, Jones DP. 2014. Integrated redox proteomics and
metabolomics of mitochondria to identify mechanisms of cd toxicity. Toxicological
sciences 139:59-73.

Go YM, Sutliff RL, Chandler JD, Khalidur R, Kang BY, Anania FA, et al. 2015. Low-dose
cadmium causes metabolic and genetic dysregulation associated with fatty liver disease
in mice. Toxicological sciences 147:524-534.

Gocmen C, Kumcu EK, Secilmis A, Ucar P, Dikmen A, Baysal F. 2000. Restorative effects of
zinc and selenium on nitrergic relaxations impaired by cadmium in the mouse corpus
cavernosum. Toxicol Lett 111:229-234.

Gonzalez I, Cao KA, Davis MJ, Dejean S. 2012. Visualising associations between paired 'omics'
data sets. BioData Min 5:19.

Haddam N, Samira S, Dumont X, Taleb A, Lison D, Haufroid V, et al. 2011. Confounders in the
assessment of the renal effects associated with low-level urinary cadmium: An analysis in
industrial workers. Environ Health 10:37.

Hastings KT, Cresswell P. 2011. Disulfide reduction in the endocytic pathway: Immunological
637   functions of gamma-interferon-inducible lysosomal thiol reductase. Antioxid Redox
638   Signal 15:657-668.

639   Horby P, Nguyen NY, Dunstan SJ, Baillie JK. 2012. The role of host genetics in susceptibility to
640   influenza: A systematic review. PLoS One 7:e33180.

641   Horby P, Nguyen NY, Dunstan SJ, Baillie JK. 2013. An updated systematic review of the role of
642   host genetics in susceptibility to influenza. Influenza Other Respir Viruses 7 Suppl 2:37-
643   41.

644   Hu X, Fernandes J, Jones DP, Go YM. 2017. Cadmium stimulates myofibroblast differentiation
645   and mouse lung fibrosis. Toxicology 383:50-56.

646   Hwang HS, Lee YT, Kim KH, Park S, Kwon YM, Lee Y, et al. 2016. Combined virus-like
647   particle and fusion protein-encoding DNA vaccination of cotton rats induces protection
648   against respiratory syncytial virus without causing vaccine-enhanced disease. Virology
649   494:215-224.

650   Jones DP, Mody VC, Jr., Carlson JL, Lynn MJ, Sternberg P, Jr. 2002. Redox analysis of human
651   plasma allows separation of pro-oxidant events of aging from decline in antioxidant
652   defenses. Free Radic Biol Med 33:1290-1300.

653   Jones DP, Liang Y. 2009. Measuring the poise of thiol/disulfide couples in vivo. Free Radic Biol
654   Med 47:1329-1338.

655   Kataranovski M, Mirkov I, Belij S, Nikolic M, Zolotarevski L, Ciric D, et al. 2009. Lungs:
656   Remote inflammatory target of systemic cadmium administration in rats. Environ Toxicol
657   Pharmacol 28:225-231.

658   Ko EJ, Lee YT, Kim KH, Jung YJ, Lee Y, Denning TL, et al. 2016. Effects of mf59 adjuvant on
659   induction of isotype-switched igg antibodies and protection after immunization with t-
660   dependent influenza virus vaccine in the absence of cd4+ t cells. J Virol 90:6976-6988.

661   Larsson SC, Wolk A. 2016. Urinary cadmium and mortality from all causes, cancer and
662   cardiovascular disease in the general population: Systematic review and meta-analysis of
663   cohort studies. Int J Epidemiol 45:782-791.

664   Le Cao KA, Gonzalez I, Dejean S. 2009. Integromics: An r package to unravel relationships
665   between two omics datasets. Bioinformatics 25:2855-2856.

Lee BK, Kim Y. 2016. Association of blood cadmium level with metabolic syndrome after
adjustment for confounding by serum ferritin and other factors: 2008-2012 korean
national health and nutrition examination survey. Biol Trace Elem Res 171:6-16.

Lee YN, Kim MC, Lee YT, Kim YJ, Lee J, Kim C, et al. 2015. Co-immunization with tandem
repeat heterologous m2 extracellular proteins overcomes strain-specific protection of split
vaccine against influenza a virus. Antiviral Res 122:82-90.

Lee YT, Ko EJ, Lee Y, Lee YN, Bian Z, Liu Y, et al. 2016. Cd47 plays a role as a negative
regulator in inducing protective immune responses to vaccination against influenza virus.
J Virol 90:6746-6758.

Li S, Park Y, Duraisingham S, Strobel FH, Khan N, Soltow QA, et al. 2013. Predicting network
activity from high throughput metabolomics. PLoS Comput Biol 9:e1003123.

Liu M, Zhao X, Hua S, Du X, Peng Y, Li X, et al. 2015. Antigenic patterns and evolution of the
human influenza a (h1n1) virus. Sci Rep 5:14171.

Liu Q, Zhou YH, Yang ZQ. 2016. The cytokine storm of severe influenza and development of
immunomodulatory therapy. Cell Mol Immunol 13:3-10.

Monsalvo AC, Batalle JP, Lopez MF, Krause JC, Klemenc J, Hernandez JZ, et al. 2011. Severe
pandemic 2009 h1n1 influenza disease due to pathogenic immune complexes. Nat Med
17:195-199.

Moya M, Bautista EG, Velazquez-Gonzalez A, Vazquez-Gutierrez F, Tzintzun G, Garcia-
Arreola ME, et al. 2013. Potentially-toxic and essential elements profile of ah1n1 patients
in mexico city. Sci Rep 3:1284.

Nemery B. 1990. Metal toxicity and the respiratory tract. Eur Respir J 3:202-219.

Olsson IM, Bensryd I, Lundh T, Ottosson H, Skerfving S, Oskarsson A. 2002. Cadmium in
blood and urine--impact of sex, age, dietary intake, iron status, and former smoking--
association of renal effects. Environ Health Perspect 110:1185-1190.

Olszowski T, Baranowska-Bosiacka I, Gutowska I, Chlubek D. 2012. Pro-inflammatory
properties of cadmium. Acta Biochim Pol 59:475-482.

Pader I, Sengupta R, Cebula M, Xu J, Lundberg JO, Holmgren A, et al. 2014. Thioredoxin-
related protein of 14 kda is an efficient l-cystine reductase and s-denitrosylase. Proc Natl
Acad Sci U S A 111:6964-6969.

Phan UT, Lackman RL, Cresswell P. 2002. Role of the c-terminal propeptide in the activity and

maturation of gamma -interferon-inducible lysosomal thiol reductase (gilt). Proc Natl

Acad Sci U S A 99:12298-12303.

Ramsey CD, Kumar A. 2013. Influenza and endemic viral pneumonia. Crit Care Clin 29:1069-

1086.

Rezaee F, Georas SN. 2014. Breaking barriers. New insights into airway epithelial barrier

function in health and disease. Am J Respir Cell Mol Biol 50:857-869.

Riederer AM, Belova A, George BJ, Anastas PT. 2013. Urinary cadmium in the 1999-2008 u.S.

National health and nutrition examination survey (nhanes). Environ Sci Technol 47:1137-

1147.

Safronetz D, Rockx B, Feldmann F, Belisle SE, Palermo RE, Brining D, et al. 2011a. Pandemic

swine-origin h1n1 influenza a virus isolates show heterogeneous virulence in macaques. J

Virol 85:1214-1223.

Safronetz D, Rockx B, Feldmann F, Belisle SE, Palermo RE, Brining D, et al. 2011b. Pandemic

swine-origin h1n1 influenza a virus isolates show heterogeneous virulence in macaques. J

Virol 85:1214-1223.

Satarug S, Moore MR. 2004. Adverse health effects of chronic exposure to low-level cadmium

in foodstuffs and cigarette smoke. Environ Health Perspect 112:1099-1103.

Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. 2006. Xcms: Processing mass

spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and

identification. Anal Chem 78:779-787.

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. 2005. Gene

set enrichment analysis: A knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci U S A 102:15545-15550.

Suwazono Y, Kido T, Nakagawa H, Nishijo M, Honda R, Kobayashi E, et al. 2009. Biological

half-life of cadmium in the urine of inhabitants after cessation of cadmium exposure.

Biomarkers 14:77-81.

Taubenberger JK, Kash JC. 2010a. Influenza virus evolution, host adaptation, and pandemic

formation. Cell Host Microbe 7:440-451.

Taubenberger JK, Kash JC. 2010b. Influenza virus evolution, host adaptation, and pandemic

formation. Cell Host Microbe 7:440-451.

Thijssen S, Maringwa J, Faes C, Lambrichts I, Van Kerkhove E. 2007. Chronic exposure of mice
to environmentally relevant, low doses of cadmium leads to early renal damage, not
predicted by blood or urine cadmium levels. Toxicology 229:145-156.
To KK, Hung IF, Li IW, Lee KL, Koo CK, Yan WW, et al. 2010. Delayed clearance of viral
load and marked cytokine activation in severe cases of pandemic h1n1 2009 influenza
virus infection. Clin Infect Dis 50:850-859.
Uppal K, Soltow QA, Strobel FH, Pittard WS, Gernert KM, Yu T, et al. 2013. Xmsanalyzer:
Automated pipeline for improved feature detection and downstream analysis of large-
scale, non-targeted metabolomics data. BMC Bioinformatics 14:15.
Uppal K, Walker DI, Jones DP. 2017. Xmsannotator: An r package for network-based annotation
of high-resolution metabolomics data. Anal Chem 89:1063-1067.
Uppal K, Ma C, Go YM, Jones DP, Wren J. 2018. Xmwas: A data-driven integration and
differential network analysis tool. Bioinformatics 34:701-702.
Waalkes MP. 2003. Cadmium carcinogenesis. Mutat Res 533:107-120.
Wang Y, De Keulenaer GW, Lee RT. 2002. Vitamin d(3)-up-regulated protein-1 is a stress-
responsive gene that regulates cardiomyocyte viability through interaction with
thioredoxin. J Biol Chem 277:26496-26500.
Wilson N, Oliver J, Rice G, Summers JA, Baker MG, Waller M, et al. 2014. Age-specific
mortality during the 1918-19 influenza pandemic and possible relationship to the 1889-92
influenza pandemic. J Infect Dis 210:993-995.
Xing ZH, Sun X, Xu L, Wu Q, Li L, Wu XJ, et al. 2015. Thin-section computed tomography
detects long-term pulmonary sequelae 3 years after novel influenza a virus-associated
pneumonia. Chin Med J (Engl) 128:902-908.
Yamawaki H, Pan S, Lee RT, Berk BC. 2005. Fluid shear stress inhibits vascular inflammation
by decreasing thioredoxin-interacting protein in endothelial cells. J Clin Invest 115:733-
738.
Zhang W, Zhi J, Cui Y, Zhang F, Habyarimana A, Cambier C, et al. 2014. Potentiated interaction
between ineffective doses of budesonide and formoterol to control the inhaled cadmium-
induced up-regulation of metalloproteinases and acute pulmonary inflammation in rats.
PLoS One 9:e109136.
757
758

 Figure Legends

Figure 1. Impact of low-dose Cd on H1N1 infection lung histopathology. Mice were given ad

libitum drinking water without or with $CdCl_2$ (1 mg/l) for 16 weeks; ten days prior to

completion, mice in both groups were given one intranasal dose of either sterile saline (Ctl, Cd)

or H1N1 influenza virus (H1N1, Cd+H1N1). Lung tissue was collected and formalin-fixed upon

sacrifice at Day 10 after infection. **(A)** Representative images of hematoxylin and eosin-stained

sections of lung tissue are shown for each group (scale bar: 100 μm). **(B)** Airway, **(C)** blood

vessel and **(D)** interstitial spaces were assessed for inflammatory foci in blinded fashion. n = 4-5

mice per group. *$p < 0.05$ compared to Ctl.

Figure 2. Cd increased lung immune cell response to H1N1 infection. Cells were isolated from

lung tissue homogenates, labeled with appropriate antibodies for **(A)** neutrophils (PMNs), **(B)**

monocytes, **(C)** $CD11b^+$ dendritic cells (DCs), **(D)** $CD103^+$ DCs, **(E)** plasmacytoid DCs (pDCs),

(F) $CD4^+$ T cells, **(G)** $CD8^+$ T cells and quantified by flow cytometry. Bar plots show mean ±

SEM of cell counts (n = 11-13, except for $CD4^+$ and $CD8^+$ T cells, n = 7-8). *$p < 0.05$ compared

to Ctl, #$p < 0.05$ compared to H1N1 by one-way ANOVA with Holm-Sidak's post-test.

Figure 3. Cd promotes mouse lung myeloid cell responses to H1N1 infection. Inflammatory

lung tissue cell counts (7 cell types, **Fig. 2**), leading edge gene intensities (293 leading edge

genes selected by GSEA, **Supplementary File 3**;) and lung tissue metabolites measured by

untargeted LC-MS (total of 2956 metabolic features, **Supplementary File 5**) were correlated

using sparse PLS regression implemented by *xMWAS* (n = 5/group). **(A)** Networks were

constructed of variables with inter-dataset absolute association score >0.88 at $p < 0.05$ (n = 5) for

H1N1 or Cd+H1N1 groups. Symbols depict metabolic features (orange rectangle), gene

transcripts (green circle) or cell counts (blue triangle), and lines indicate relationship between

symbols (red, positive correlation; blue, negative correlation). The blue triangles have been

enlarged for easier viewing. Abbreviations for cells are as follows: Monocytes, 'Mono';

neutrophils, 'PMN'; $CD11b^+$ dendritic cells, '$CD11b^+$ DC'; $CD103^+$ dendritic cells, '$CD103^+$

DC'; plasmacytoid dendritic cells, 'pDC'; CD4 T lymphocytes, '$CD4^+$'; and CD8 T

lymphocytes, '$CD8^+$'. Note, $CD103^+$ DC is not found in the Cd+H1N1 network. **(B)** The

absolute value of difference in eigenvector centrality (Delta Centrality) of each immune cell type

was contrasted between the networks of Cd+H1N1 and H1N1 networks.   Pearson's correlation

(r) was analyzed for the abundance of lung IFNγ protein and lung monocytes **(C)**, $CD11b^+$ DCs

(D) and neutrophils **(E)**. Open circles, H1N1 alone; closed circles, Cd+H1N1. The resulting

statistics of correlation (r) and significance (p) were as follows: **C**, monocytes (<u>Cd+H1N1</u>, r =

0.97, p = 3×10^{-7}; H1N1, r = 0.21, p = 0.49); **D**, $CD11b^+$ DCs (Cd+H1N1, r = 0.92, p = 2×10^{-5};

H1N1, r = -0.05, p = 0.86); **E**, neutrophils (Cd+H1N1, r = 0.89, p = 9×10^{-5}; H1N1, r = 0.22, p =

0.48). n = 12-13.

Figure 4. Cd potentiation of immune response to H1N1 infection through widespread effects of

low Cd exposure on inflammation, mitochondrial function and regulation of immune cell

populations through IFNγ. **(A)** Percentage of gene sets represented in the 83 leading edge genes

(**Supplementary Table 2**) with delta eigenvector centrality (absolute value), |$Centrality_{Cd+H1N1}$ −

$Centrality_{H1N1}$|, ≥ 0.1 between Cd+H1N1 and H1N1 networks. **(B)** Metabolic pathway

enrichment analysis was performed on 165 metabolic features with delta eigenvector centrality $\geq$

0.1 between networks of Cd+H1N1 and H1N1 using *mummichog*. The negative $\log_{10}$ of the

pathway p-value is plotted as a bar graph for each significant pathway ($p < 0.05$, at least 3

metabolite matches in pathway).

831 **Table 1.** Cd increases pro-inflammatory gene enrichment in response to H1N1 challenge.

Gene Sets		Enrichment		
Name	Number of Genes	*H1N1 vs. Ctl*	*Cd+H1N1 vs. Ctl*	*Cd+H1N1 vs. H1N1*
Allograft rejection	189	3.02	3.21	2.31
Complement	181	2.36	2.56	1.73
Inflammatory response	199	2.61	2.75	1.84
Interferon-α response	89	2.79	2.88	2.04
Interferon-γ response	195	3.09	3.36	2.49
TNF-α signaling via NF-κB	196	2.60	2.82	2.06
Myogenesis	195	NA	-1.66	-2.14

Gene set normalized enrichment scores that are statistically significant comparing Cd+H1N1 and H1N1

($p < 0.05$ and FDR $q < 0.05$) are shown. Normalized enrichment scores compared to controls (also $p <$

0.05 and FDR $q < 0.05$) are shown for H1N1 alone or Cd+H1N1. Negative enrichment values indicate

lower gene abundances compared to the reference group (Ctl or H1N1, indicated in the column header).

NA, gene set enrichment did not pass significance testing threshold.

Figure 1

853   **Figure 2**

   Figure 3

879   **Figure 4**