1 **Title:**
2
3 Identifying, understanding, and correcting technical biases on the sex chromosomes in
4 next-generation sequencing data
5
6 **Authors and Affiliations:**
7

8 Timothy H. Webster[1], Madeline Couse[2], Bruno M. Grande[3], Eric Karlins[4], Tanya Phung[5],
9 Phillip A. Richmond[6,7], Whitney Whitford[8], Melissa A. Wilson Sayres[1,9]
10
[1]School of Life Sciences, Arizona State University
[2]Child and Family Research Institute, University of British Columbia
[3]Department of Molecular Biology and Biochemistry, Simon Fraser University
[4]Division of Cancer Epidemiology and Genetics, National Cancer Institute, National
Institutes of Health
[5]Interdepartmental Program in Bioinformatics, UCLA
[6]Centre for Molecular Medicine and Therapeutics, University of British Columbia
[7]BC Children's Hospital
[8]School of Biological Sciences, The University of Auckland
[9]Center for Evolution and Medicine, Arizona State University
21
22 **Corresponding Authors:**
23
24 Timothy H. Webster
25 School of Life Sciences
26 Arizona State University
27 Tempe, AZ, USA 85281
28 Timothy.h.webster@asu.edu
29
30 Melissa A. Wilson Sayres
31 School of Life Sciences
32 Arizona State University
33 Tempe, AZ, USA 85281
34 melissa.wilsonsayres@asu.edu
35
36

Abstract
Mammalian X and Y chromosomes share a common evolutionary origin and retain
regions of high sequence similarity. This sequence homology can cause the mismapping
of short sequencing reads derived from the sex chromosomes and affect variant calling
and other downstream analyses. Understanding, and correcting, this problem is critical
for medical genomics and population genomic inference. Here, we characterize how
sequence homology can affect analyses on the sex chromosomes and present XYalign, a
new tool that: 1) aids in the inference of sex chromosome complement from next-
generation sequencing data, 2) corrects erroneous read mapping on the sex chromosomes,
and 3) tabulates and visualizes important metrics for quality control such as mapping
quality, sequencing depth, and allele balance. We show how these metrics can be used to
identify XX and XY individuals across diverse sequencing experiments, including low
and high coverage whole genome sequencing, and exome sequencing. We also show that
the default steps taken by XYalign correct many mismapped reads on the sex
chromosomes, resulting in more accurate variant calling. Finally, we discuss how the
flexibility of XYalign's framework can be leveraged for other use cases including the
identification of aneuploidy on the autosomes. XYalign is available open source under
the GNU General Public License (version 3).
55

56   **Keywords**
57   X chromosome; Y chromosome; ploidy; aneuploidy; genomics; variant calling; mapping

Introduction

Accurate genotyping and variant calling are priorities in medical genetics, including molecular diagnostics, and population genomics (Taylor *et al.*, 2015; Ashley, 2016). Despite the availability of numerous powerful tools developed to infer genotypes from sequencing data, sequence homology among genomic regions still presents a major challenge to genome assembly, short read mapping, and variant calling. Specifically, similar sequence content can confound the mapping of short next-generation sequencing reads to a reference genome and lead to technical artifacts in downstream analyses and applications. Heteromorphic sex chromosomes, in particular, present a case of sequence homology likely to affect all individuals in a given species.

Sex chromosomes in therians—the clade containing eutherian mammals and marsupials—share a common evolutionary origin as a pair of homologous autosomes (Glas *et al.*, 1999). Approximately 180 to 210 million years ago, they began differentiating from each other through a series of recombination suppression events and subsequent gene loss on the Y chromosome (Rens *et al.*, 2007; Lahn and Page, 1999; Livernois *et al.*, 2012; Wilson Sayres and Makova, 2013). This pattern, however, is not unique to mammalian evolution or even XX/XY systems, and occurs often across taxa with genetic sex determination (Bergero and Charlesworth, 2009; Wilson and Makova, 2009). This shared origin and complex history characteristic of sex chromosomes lead to unique challenges for genome assembly and analysis, including large blocks of homologous sequence between the sex chromosomes—called gametologous sequence—that we hypothesize can lead to the mismapping of reads between the sex chromosomes. Further, the sex chromosomes of many species contain pseudoautosomal regions (PARs; of which humans have two: PAR1 and PAR2)—regions that have not differentiated between the sex chromosomes and are identical in sequence between the two sex chromosomes (Simmler *et al.*, 1985; Ross *et al.*, 2005). A reference genome that includes the entire sequence content from both sex chromosomes will thus duplicate the PARs and substantially reduce mapping quality in these regions because most reads will identically map to two regions in the reference assembly. This stands in contrast to autosomal sequence, for which each diploid autosome is represented just once in the reference genome.

The technical challenges presented by the biological realities of the sex chromosomes can lead to erroneous genotype calls, so the sex chromosomes are routinely excluded from genome-wide analyses (e.g., Wise *et al.*, 2013). This is unfortunate because the sex chromosomes contribute to phenotype and disease etiology (e.g., Chang *et al.*, 2014) and are useful in population genetic inference of demography and patterns of natural selection (Webster and Wilson Sayres, 2016; Wilson Sayres, 2018; Vicoso and Charlesworth, 2006; Ellegren, 2009; Meisel and Connallon, 2013).

A number of tools, methods, and frameworks have been developed to aid in the identification of sex-linked sequence (e.g., Muyle *et al.*, 2016), inference of an individual's sex chromosome complement (e.g., Madel *et al.*, 2016), and handling of some of the technical challenges sex chromosomes present in genome-wide association studies (e.g., Gao *et al.*, 2015). However, to our knowledge, there is no tool that simultaneously aids in the identification of sex chromosome complement and corrects for associated technical biases for the purposes of short read mapping and variant calling.

Out of the urgent need to understand the effects of sex chromosome homology on
next-generation sequencing analyses, in this paper we first test whether sequence
homology between sex chromosomes can confound aspects of read mapping and lead to
downstream errors in sequence analysis. We then present XYalign, a tool developed to
perform three major tasks: (1) aid in the characterization of an individual's sex
chromosome complement; (2) identify and correct for technical artifacts arising from sex
chromosome sequence homology; and (3) tabulate and visualize important metrics for
quality control such as mapping quality, sequencing depth, and allele balance. We show
how XYalign can be used to identify XX and XY individuals across sequencing depths
and capture techniques. We also show that the default steps taken by XYalign correct
many mismapped reads on the sex chromosomes, resulting in more accurate variant
calling. Finally, because XYalign is designed to be both scalable and customizable, we
discuss how it can be used in a variety of situations including genetic sex identification in
both XX/XY and ZZ/ZW systems, identification of sex-linked sequences and
pseudoautosomal regions in new draft genomes, correction of technical biases in genomic
and transcriptomic data, detection of aneuploidy, and investigation of mapping success
across arbitrary chromosomes.
120
Methods
Implementation
XYalign is implemented in Python and uses a number of third-party Python
packages including Matplotlib (Hunter, 2007), NumPy (Oliphant, 2006), Pandas
(McKinney, 2010), PyBedTools (Quinlan and Hall, 2010; Dale *et al.*, 2011), PySam
(https://github.com/pysam-developers/pysam), and SciPy (Jones *et al.*, 2001). It further
wraps the following external tools: repair.sh and shuffle.sh from BBTools
(https://sourceforge.net/projects/bbmap/), BWA (Li, 2013), Platypus (Rimmer *et al.*,
2014), Sambamba (Tarasov *et al.*, 2015), and SAMtools (Li *et al.*, 2009).
XYalign is composed of six modules that can be called individually or serve as
steps in a full pipeline: PREPARE_REFERENCE, CHROM_STATS, ANALYZE_BAM,
CHARACTERIZE_SEX_CHROMS, STRIP_READS, and REMAPPING. Below, we
discuss each module as a step in the full XYalign pipeline using human samples (XX/XY
sex determination) as an example. Note, however, that XYalign will work with other sex
chromosome systems (e.g., ZZ/ZW) and on arbitrary chromosomes (e.g., detecting
autosomal aneuploidy). We describe examples of XYalign commands in the section titled
"Use Cases."
The PREPARE_REFERENCE module generates two versions of the same
reference genome: one for the homogametic sex (e.g., XX) and one for the heterogametic
sex (e.g, XY). In the simplest case, it will completely hard-mask the Y chromosome with
Ns in the XX version of the reference. However, it will also accept one or more BED
files containing regions to hard mask in both reference versions. If pseudoautosomal
regions (PARs) are present on both sex chromosome sequences in the reference, we
strongly suggest masking the PARs on the Y chromosome, allowing reads from these
regions to map exclusively to the X chromosome in XY individuals. In XYalign, we use
hard masks, rather than omitting the Y chromosome in the XX reference version because
these hard masks allow files from both references to share the same sequence dictionaries

and indices, thus permitting seamless integration of files from both references into
downstream analyses (e.g., joint variant calling).
The CHROM_STATS module provides a relatively quick comparison of mapping
quality and sequencing depth across one or more chromosomes and over multiple BAM
files. While this provides a less detailed perspective than ANALYZE_BAM or
CHARACTERIZE_SEX_CHROMS (detailed below), we envision it to be especially
useful in at least two different cases. First, in well-characterized systems (e.g., human),
comparing chromosome-wide values of mean mapping quality and depth represent a
quick and often sufficient way to identify the sex chromosome complement (e.g., XX or
XY) of individuals across a population. Second, in uncharacterized systems, the
CHROM_STATS output provides information that can help with the identification of
sex-linked scaffolds. It is important to note, however, that results for both cases will vary
based on ploidy and with differences in the degree of sequence homology between the
sex chromosomes.
The ANALYZE_BAM module runs a series of analyses designed to aid in the
identification of sex-linked sequence and characterize the sex chromosome content of an
individual. In doing so, it provides more detailed metrics than CHROM_STATS. For
ANALYZE_BAM, XYalign runs Platypus (Rimmer *et al.*, 2014) across multiple threads,
if permitted, to call variants. It then parses the output VCF file containing the variants,
applies filters for site quality, genotype quality, and read depth, and plots the read balance
at variant sites. Here, we define read balance at a given site as the number of reads
containing the alternate allele (i.e., nonreference allele) divided by the total number of
reads mapped to the position. XYalign produces plots and tables for read balance per site,
as well as mean read balance and variant count per genomic bin or window across a
chromosome. We anticipate these data will not only be useful for masking regions
containing incorrect genotypes, but will also aid in the identification of PARs as well.
XYalign next traverses the BAM file, calculating mean mapping quality and an
approximation of mean depth in windows across the genome. During traversal, secondary
and supplementary read mappings are ignored, and depth is calculated as the total length
of all reads mapping to a genomic window divided by the total length of the window. We
have found that this heuristic approximation is very similar to calculations of exact depth,
particularly as window sizes increase, and much faster to compute across entire
chromosomes. XYalign will output a table containing genomic coordinates, mean depth,
and mean mapping quality for each window. It will then filter windows based on user-
defined thresholds of mean depth and mapping quality and output two BED files
containing windows that passed and failed these thresholds, respectively, which can be
used for additional masking in downstream applications. Finally, XYalign will output
plots of mapping quality and depth in each window across each chromosome.
After running ANALYZE_BAM, the windows meeting thresholds can be used by
the CHARATERIZE_SEX_CHROMS module to systematically compare mean depth in
pairs of chromosomes using three different approaches. The first is a bootstrap analysis
that provides 95% confidence intervals of mean window depth for each of the
chromosomes in a given pair to test for overlap. The second is a permutation analysis to
test for differences in depth between the two chromosomes. The third is a two-sample
Kolmogorov-Smirnov test (Massey Jr., 1951). Though all three tests are implemented in
XYalign, we only present results from the bootstrap analyses in this manuscript. Further,

while we present analyses pairing sex chromosomes with an autosome (here we use
chromosome 19), the chromosome pairs are arbitrary and can feature any scaffolds or
chromosomes in a reference genome, depending on a user's needs.
Finally, the REMAPPING module will infer the presence or absence of a Y
chromosome based on the results of CHARACTERIZE_SEX_CHROMS. If a Y
chromosome is not detected, XYalign's STRIP_READS module will iteratively remove
reads from the sex chromosomes by read group ID using SAMtools (Li *et al.*, 2009),
writing FASTQ files for each. XYalign will use repair.sh from BBTools to sort and re-
pair paired-end reads or shuffle.sh from BBTools to sort single-end reads for each read
group. The REMAPPING module then, by read group, maps reads with BWA-MEM (Li,
2013) and sorts alignments with SAMtools (Li *et al.*, 2009). If more than one read group
is present, the resulting BAM files are merged using SAMtools (Li *et al.*, 2009). Finally,
XYalign uses SAMBAMBA (Tarasov *et al.*, 2015) to isolate all scaffolds not associated
with sex chromosomes from the original BAM file and SAMtools (Li *et al.*, 2009) and
merge this file with the bam file containing the new sex chromosome mappings.
When run as a full pipeline on a sample, XYalign will first call
PREPARE_REFERENCE to generate XX and XY reference genomes with appropriate
masks. Next, it will call ANALYZE_BAM and CHARACTERIZE_SEX_CHROMS to
preliminarily analyze the unprocessed input BAM file. Then, based on the results of
CHARACTERIZE_SEX_CHROMS, XYalign will call STRIP_READS to extract reads
from the sex chromosomes and REMAPPING to remap to the appropriate reference
genome output from PREPARE_REFERENCE. Finally, XYalign will re-run the
ANALYZE_BAM module to analyze the remapped BAM file and provide metrics to
allow a before-and-after comparison.
While we anticipate this full pipeline will be useful in certain situations, it is
neither the only nor the best-suited option for most users. Rather, we expect that most
users will call modules individually. We give examples of other implementations below
and provide recommendations for incorporating XYalign into bioinformatics pipelines in
the discussion.
223
Operation
XYalign is available via PyPI (https://pypi.python.org/pypi), Bioconda (Grüning
et al., 2017), and Github (https://github.com/WilsonSayresLab/XYalign), with
documentation hosted at Read the Docs (http://xyalign.readthedocs.io/en/latest/). A full
environment containing all dependencies can be most easily installed and managed using
Anaconda (https://www.continuum.io/) and Bioconda (Grüning *et al.*, 2017). It has been
tested on a variety of UNIX operating systems (including Linux and MacOS), but is not
currently supported for the Windows operating system.
XYalign is typically invoked from the command line, but it can be imported into
Python scripts for more customized use cases. The next section lists a number of example
commands that illustrate how to call the full pipeline as well as individual modules.
235
Use Cases
To highlight some of XYalign's features and flexibility, we used two datasets
from publically available sources (Supplemental Table S1): (1) exome, low-coverage
whole-genome, and high-coverage whole-genome data from one male (HG00512) and

one female (HG00513) from the 1000 Genomes Project (Dataset 1; (The 1000 Genomes
Project Consortium, 2015); and (2) 24 high-coverage whole-genomes from the 1000
Genomes Project (Dataset 2; (Sudmant *et al.*, 2015). For Dataset 1, we mapped reads to
the hg19 version of the human reference genome (International Human Genome
Sequencing Consortium, 2001) using BWA MEM (Li, 2013), marked duplicates with
SAMBLASTER (Faust and Hall, 2014), and used SAMtools (Li *et al.*, 2009) to sort,
index, and merge BAM files. The publicly available bam files for Dataset 2 were
previously mapped using a different version of hg19 (from the Broad Institute's GATK
Resource Bundle; https://software.broadinstitute.org/gatk/download/bundle), which we
used for analyses involving this dataset.
With these datasets, we examined three potential uses of XYalign. First, to
explore the effects of simple corrections for technical biases arising from sequence
homology on the sex chromosomes, we ran the full XYalign pipleline on all six BAM
files from the Dataset 1. In all cases below, the exact commands are included in the
Supplementary Material and in a Snakemake (Köster and Rahmann, 2012) pipeline
available with the XYalign software distribution (Webster *et al.*, 2018), and templates are
shown here for ease in reading. Because we were using the same output directory for
these analyses, we avoided conflicts by initially preparing separate XX and XY
references using the following command:
259
260 *xyalign --PREPARE_REFERENCE --ref <hg19 reference genome> --xx_ref_out*
261 *hg19.XXonly.fasta --xy_ref_out hg19.XY.fasta --output_dir <output_directory> --*
262 *x_chromosome chrX --y_chromosome chrY --bwa_index True,*
263
264 where *<hg19 reference genome>* was the path to the FASTA file containing the hg19
265 reference, *<input bam file>* was a sorted BAM file, and *<output directory>* was the
266 directory for XYalign to write output. We then ran the full pipeline on all six files using
267 the following command template:
268
xyalign --ref <hg19 reference genome> --bam <input bam file> --output_dir <output
directory> --sample_id <sample ID> --cpus 4 --reference_mask
hg19_PAR_Ymask_startEnd.bed --window_size 5000 --chromosomes chr19 chrX chrY --
x_chromosome chrX --y_chromosome chrY --xmx 4g --fastq_compression 4 --
min_depth_filter 0.2 --max_depth_filter 2 --xx_ref_in hg19.XXonly.fasta --xy_ref_in
ref_out hg19.XY.fasta,
275
where *<sample ID>* was the identification code for a given sample,
hg19_PAR_Ymask_startEnd.bed was a BED file containing the genomic coordinates of
the PARs in the hg19 assembly, and hg19.XXonly.fasta and hg19.XY.fasta were the two
fasta formatted reference genomes prepared in the previous step.
Next, we examined how XYalign's metrics can be used to identify the sex
chromosome complement of individuals with both datasets. Here, we used the
CHARACTERIZE_SEX_CHROMS module of XYalign. This was automatically done
for Dataset 1 when running the full pipeline (see above). For Dataset 2, we used the
following command template for BAM files:
285

286 *xyalign --CHARACTERIZE_SEX_CHROMS --ref <1000 genomes reference genome> --*
287 *bam <input bam file> --output_dir <output directory> --sample_id <sample ID> --cpus*
288 *4 --window_size 5000 --chromosomes 19 X Y --x_chromosome X --y_chromosome Y*
289
290 Finally, we explored the utility of the CHROM_STATS module for identifying
291 sex chromosome complement and potentially sex-linked scaffolds with both datasets
292 using the following command template for BAM files:
293
294 *xyalign --CHROM_STATS --chromosomes chr1 chr8 chr19 chrX chrY chrM --bam*
295 *<input_bam_1> <input_bam_2> <input_bam_3> --ref null --sample_id*
296 *<name_of_analysis> --output_dir <output_dir>*
297
298 We additionally ran CHROM_STATS using the above command with the
299 addition of the "--use_counts" flag to calculate metrics using only the number of reads
300 mapping to each chromosome.
301
302 We visualized all CHROM_STATS results using XYalign's plot_count_stats utility,
303 using the template:
304
305 *plot_count_stats --input <chrom_stats output file> --output_prefix <output prefix>--*
306 *meta <metadata text file> --exclude_suffix <suffix> --first_chr chrX --second_chr chrY -*
307 *-const_chr chr19 --var1_marker color --var1_marker_vals darklateblue thistle --*
308 *var2_marker shape --var2_marker_vals o s v --marker_size 1700 --legend_marker_scale*
309 *0.4*
310
311 where *<chrom_stats_output_file>* was either the count, mapq, or depth output of
312 CHROM_STATS, *<metadata text file>* was the appropriate metadata text file, and
313 *<suffix>* was the string to remove from filenames.
314
Sex chromosome coordinates
To better understand the genomic context of technical artifacts, we explored
variation in mapping quality and depth in association with genomic features on the X and
Y chromosomes. On the Y chromosome, we used coordinates from Poznik et al. (2013)
based on Skaletsky et al. (2003) (provided by D. Poznik, personal communication). On
the X chromosome, we obtained coordinates for campliconic regions from Cotter et al.
(2016) and all other regions from the UCSC Table Browser (Karolchik *et al.*, 2004). We
define the X-transposed region (XTR) on the X chromosome as beginning at the start of
DXS1217 and ending at the end of DXS3 (Mumm *et al.*, 1997).
To count variants falling in major genomic regions, we intersected a BED file
containing coordinates with VCF files using BEDTools (Quinlan and Hall, 2010). We
first filtered VCF files using BCFtools (Li *et al.*, 2009) with the following command
template:
328
329 *bcftools filter --include 'INFO/MQ>=30 && %QUAL>=30' <input_vcf>*
330

331   We then found variants unique to each file through iterations of the "subtract" command
332   in BEDtools (Quinlan and Hall, 2010):

333
334   *bedtools subtract -header -a <first_vcf> -b <second_vcf>*

335
336   Finally, in each region, we counted variants present in a given filtered VCF file using the
337   BEDtools (Quinlan and Hall, 2010) "intersect" command:

338
339   *bedtools intersect -c -a <BED file> -b <vcf_file>*

340
341   where *<BED_file>* is the BED file containing genomic coordinates (Supplemental Table
342   S2).

343
344   *Specific commands*
345   We provide Snakemake (Köster and Rahmann, 2012) workflows for all assembly
346   and analysis steps on Github (https://github.com/WilsonSayresLab/XYalign), Zenodo
347   (Webster *et al.*, 2018), and in the Supplemental Material.

348
Results and Discussion
We observed a number of artifacts stemming from several methodological
challenges presented by the human sex chromosomes. First, PAR1 and PAR2 on both sex
chromosomes are clearly identifiable in genomic scatter plots of mapping quality and
depth in all datasets (Figures 1-3). While these results are not surprising given the
sequence homology in these regions (Ross *et al.*, 2005), they highlight the fact that these
measures can help identify other similarly problematic areas. For example, there is a
region of reduced mapping quality on the X chromosome beginning near 88.4 Mb and
ending near 92.3 Mb (Figure 2). This corresponds to the X-transposed region (XTR),
which arose by a duplication from the X to the Y chromosome in the human lineage since
its divergence with the chimpanzee-bonobo lineage (Page *et al.*, 1984; Ross *et al.*, 2005).
This region retains more than 98% sequence similarity between the X and Y chromosome
(Ross *et al.*, 2005), likely leading to the reduction in mapping quality. Interestingly, we
observe a similar decrease in mapping quality on the Y chromosome beginning near 2.9
Mb and ending near 6.6 Mb, corresponding with known coordinates of the XTR on the Y
chromosome (Figure 3). In fact, the combination of mapping quality and depth
recapitulates well-known major genomic features of both sex chromosomes (e.g.,
ampliconic regions, PARs, and XTRs) described in previous studies (Figures 1-3; Poznik
et al., 2013; Mueller *et al.*, 2013). This suggests that, in at least some cases, the output of
XYalign can be used to quickly explore broad patterns of genomic architecture as well as
mask regions likely to introduce technical difficulties in genomic analyses.

370
371

Figure 1. Sequencing depth on chromosome X before and after XYalign. Mean sequencing depth in 5 kb windows across the X chromosome before (A) and after (B) XYalign processing. Changes in depth (D) are presented as the sign of the difference times the absolute value of the $\log_{10}$ difference, where the difference is depth after XYalign minus depth before XYalign. The chromosome map (C) presents the location of X chromosome genomic features depicted in the legend. X chromosome coordinates are identical in all plots.

Figure 2. Mapping quality on chromosome X before and after XYalign. Mean mapping quality (MAPQ) in 5 kb windows across the X chromosome before (A) and after (B) XYalign processing. Changes in MAPQ (D) are presented as the difference is MAPQ after XYalign minus MAPQ before XYalign. The chromosome map (C) presents the location of X chromosome genomic features depicted in the legend. X chromosome coordinates are identical in all plots.

389
390 **Figure 3. Y chromosome sequencing depth and quality**. Mean sequencing depth (A)
391 and mapping quality (MAPQ; C) in 5 kb windows across the Y chromosome. The
392 chromosome map (B) presents the location of Y chromosome genomic features depicted
393 in the legend. Y chromosome coordinates are identical in all plots.
394

By hard-masking the Y chromosome in the XX reference genome, and the
pseudoautosomal regions (PAR1 and PAR2) in the reference genome for the XY
reference genome, we observed clear improvements in read mapping (Figures 1-2). On
the X chromosome, all metrics exhibited striking improvements in PAR1, PAR2, and
XTR (Figures 1-2). Further, the Y chromosome of the XX individual no longer exhibited
any variant calls or mapped reads, though many passed filters before processing (variants
before: 4266; variants after: 0; mapped reads before: 5,729,007; reads mapped after: 0).
While this is expected given the hard masking of the Y chromosome, it is worth
emphasizing that this matches the biological state of the individual.
We found that these improvements in mapping on the X chromosome after
masking the Y chromosome substantially impacted downstream variant calling (Table 1).
Unsurprisingly, the effect was most pronounced in the PARs, in which thousands of
variants were callable after masking the identical sequences present on the Y
chromosome in the reference assembly. The XTR also had a large number of variants
become callable after Y masking—an average of 85.4 variants per megabase of sequence
(Table 1). However, effects were not limited to these regions of well-documented
homology: both the X-added region (XAR) and X-conserved region (XCR) contained
hundreds of affected variants, suggesting effects of more extensive homology across the
sex chromosomes.
414
415 **Table 1. The effect of sex chromosome homology on variant calling on the X**
416 **chromosome.**

Region[a]	Length[b]	False positives (per Mb)[c]	False negatives (per Mb)[d]
PAR1	2,589,520	0 (0)	7563 (2920.6)
PAR2	329,516	0 (0)	633 (1921)
XTR	4,287,237	40 (9.3)	366 (85.4)
XAR	55,982,492	299 (5.3)	400 (7.2)
XCR	89,011,795	610 (6.9)	523 (5.9)
Total	*152,250,560*	*949 (6.2)*	*9485 (62.3)*

417
[a]PAR1: pseudoautosomal region 1; PAR2: pseudoautosomal region 2; XTR: X-
transposed region; XAR: X-added region; XCR: X-conserved region.
[b]Total sequence length of region in base pairs.
[c]Total number of false positive variants after filtering, defined as being present before but
not after Y chromosome masking. Variants per Mb of sequence are presented in
parentheses.
[d]Total number of false negative variants after filtering, defined as being present after but
not before Y chromosome masking. Variants per Mb of sequence are presented in
parentheses.
427
428

Figure 4. Read balance in XY and XX samples. Histograms of read balance for an XY
sample (Left Column; A, C, and E) and XX sample (Right Column; B and D) across
chromosome 19 (Top; A and B), chromosome X (Middle; C and D), and chromosome Y
(Bottom; E). Read balance at a given site is defined as the number of reads containing a

434 non-reference allele divided by the total number of reads mapped to a site. Read balances
435 between 0.05 and 1.0 are presented.
436
Inferring Genetic Sex
In our analyses, the most striking measure for assessing and individual's sex
chromosome complement was the distribution of read balances across a chromosome
(Figure 4). Specifically, when we plotted the distribution of the fraction of reads
containing a nonreference allele at a given variant site, we observed that diploid
chromosomes (e.g., autosomes, and chromosome X in XX individuals) exhibited peaks
both around 0.5 and 1.0, consistent with the presence of heterozygous sites and sites
homozygous for a nonreference allele, respectively (Figure 4). In the case of the X
chromosome in XY individuals, we observed a single peak near 1.0, consistent with an
expected haploid state (i.e., no heterozygous sites; Figure 4). We observed one exception
to this pattern: the Y chromosome exhibited a peak around 0.2 in addition to the one near
1.0 (Figure 4). All variants included in analyses met thresholds for depth, site quality, and
genotype quality, so quality does not appear to be a driving factor of this pattern. This
pattern also remained after genomic windows of low mapping quality and irregular depth
were removed. We are currently unable to explain these results and more work is thus
required to understand the factors responsible for this pattern.
Across datasets, we observed variation in relative depth of the X and Y
chromosomes in XX and XY individuals, particularly among different sequencing
strategies: exome, low-coverage whole-genome, and high-coverage whole-genome
sequencing (Figure 5A). However, within datasets, XX and XY individuals were clearly
differentiated (Figure 5; Supplemental Figure S1). This pattern suggests that a general
threshold for assigning different genetic sexes across a range of organisms and
sequencing experiments might be difficult to implement. That being said, within species,
some combination of depth, mapping quality, and read balance is likely to be informative.
For example, in humans, relative mapping quality appears to be informative in some
sequencing strategies, particularly exome sequencing (Figure 5B). This should be
explored in each experiment, however, as we did not observe this differentiation in the
uncorrected 1000 Genomes high-coverage samples (Supplemental Figure S2).
Generating these results for all individuals in a study is easy to do with XYalign:
one can iteratively run the CHARACTERIZE_SEX_CHROMS module on preliminarily
mapped BAM files. Then, the results from all individuals can be analyzed together (see
the Supplementary Material for an example of such analysis). At least with human
samples, for which X and Y chromosomes are very differentiated, this process can be
sped up significantly with the CHROM_STATS module. In our data, read counts on the
X and Y chromosomes quickly and clearly clustered male and female samples within
sequencing strategies (i.e., exome, low-coverage whole-genome, and high-coverage
whole-genome; Supplemental Figures S3-S4). However, the success of this procedure
likely depends on the degree of differentiation between sex chromosomes; other
organisms might require the statistics output as part of the
CHARACTERIZE_SEX_CHROMS module.
477
478
479

Figure 5. Relative sequencing depth and mapping quality on the X and Y
chromosomes across different sequencing strategies. Values of relative (A) sequencing
depth and (B) mapping quality come from exome (circles), low-coverage whole-genome
sequencing (squares), and high-coverage whole-genome sequencing (triangles) for a
single male (green) and female (blue) individual. Mean depth and MAPQ on
chromosome 19 was used to normalize the sex chromosomes.

Recommendations for researchers

Based on these results, we can make the following recommendations for
researchers. For organisms with multiple sex chromosomes assembled (e.g., both X and
Y or both Z and W) and included in reference assemblies (e.g., human, chimpanzee,
rhesus macaque, gorilla, mouse, rat, chicken, *Drosophila*), *if the genetic sex of every*
individual is known: 1) prepare separate assemblies for the different sexes using
XYalign's PREPARE_REFERENCE module, 2) map and process reads according to
user's typical pipeline (mapping individuals by sex to their corresponding reference), 3)
confirm genetic sex using XYalign's CHROM_STATS module, and 4) remap any
incorrectly assigned individuals, and 5) move to downstream analyses. *If genetic sexes of*
individuals are unknown, 1) prepare separate assemblies for the different sexes using
XYalign's PREPARE_REFERENCE module, 2) map and process a suitable number of
reads (e.g., whole dataset for exome or a single lane of WGS) according to user's typical
pipeline using the reference genome of the heterogametic sex (i.e., XY or ZW), 3) infer
the sex chromosome complement using either/both CHROM_STATS (for well-
characterized and highly divergent sex chromosomes) or
CHARACTERIZE_SEX_CHROMS, 4) map and process all reads using the prepared
reference genome corresponding to the inferred sex of each individual, and 5) run
downstream analyses.

For individuals of the homogametic sex (i.e., XX or ZZ), the above
recommendations will likely completely remove artifacts stemming from sex
chromosome homology, assuming only a single unmasked sex chromosome is left after

XYalign processing. However, homology is unavoidable for individuals of the
heterogametic sex (i.e., XY or ZW) because both sex chromosomes are required in the
reference assembly for mapping. In this case, a more local masking or filtering approach
is likely the most promising option. For studies investigating specific variants, for which
false negatives are preferable to false positives, we suggest strict variant filtering that
includes high thresholds for mapping quality (e.g., thresholds of 55 or higher are required
to eliminate the effects of homology in the X-transposed region). However, for studies
investigating invariant sites as well (e.g., measures of genetic diversity require
information from all monomorphic and polymorphic sites), we recommend filtering
entire regions based on, at the very least, mapping and depth metrics. These masks are
output by the BAM_ANALYSIS module in XYalign, and for this use, we recommend
using small windows (e.g, 1 kb to 5 kb) and exploring a variety of depths. Finally, in all
cases, if pseudoautosomal regions are present in the reference genome, they should be
masked in the heterogametic sex's assembly output by the PREPARE_REFERENCE
module.
526
Additional uses for XYalign
While the development of XYalign was motivated by challenges surrounding
erroneous read mapping and variant calling due to sex chromosome homology in human
sequencing experiments, the software can be utilized in a number of additional scenarios.
First, it can be applied to any species with heteromorphic sex chromosomes to identify
relative quality and depth. The results output by CHROM_STATS, ANALYZE_BAM,
and CHARACTERIZE_SEX_CHROMS can be used to identify sex-linked scaffolds,
characterize sex chromosome complements, and determine the most appropriate
remapping strategy. Second, XYalign can be used to detect relative sequencing depth,
mapping quality, and read balance on any chromosome, not just the sex chromosomes. In
addition to exploring mapping success and biases, we anticipate that this will aid in
detection of aneuploidy in the autosomes. However, we note that many programs exist to
calculate depth of coverage (e.g., Quinlan and Hall, 2010; Pedersen and Quinlan, 2018;
McKenna *et al.*, 2010) and identify structural variation within statistical frameworks
(e.g., Chen *et al.*, 2016; Layer *et al.*, 2014; Abyzov *et al.*, 2011; Roller *et al.*, 2016), so
XYalign might not be the most appropriate option for detecting local phenomena such as
copy number variants. Finally, XYalign may also be extended to other types of data,
including RNA sequencing data, where the same fundamental challenge (gametologous
sequence between the X and Y) can affect mapping and variant calling. In particular, we
expect biases to manifest in differential expression and biased-allelic expression, and
suggest that the PREPARE_REFERENCE module be considered for all RNA sequencing
experiments in systems with sex chromosomes.
549
Conclusion
We showed that the complex evolutionary history of the sex chromosomes creates
mapping biases in next-generation sequencing data that have downstream effects on
variant calling and other analyses. These technical artifacts are likely present in most
genomic datasets of species with chromosomal sex determination. However, many of
these biases can be corrected through the strategic use of masks during read mapping and
the filtering of variants. We developed XYalign, a tool that aids in the characterization of

an individual's sex chromosome complement and implements this masking strategy to correct these technical biases. We illustrated how XYalign can be used to identify the presence or absence of a Y chromosome, characterize mapping biases across the genome, and correct for these mapping biases. XYalign provides a framework to generate more robust short read mapping and improve variant calling on the sex chromosomes.

Software Availability
XYalign is available on Github (https://github.com/WilsonSayresLab/XYalign) under a GNU General Public License (version 3). We have also deposited a static version of the source code used for analyses in this paper at Zenodo (Webster *et al.*, 2018).

Author Contributions
All authors participated in the conception and initial design of the software. THW developed and implemented the software. BG, EK, TP, WW, and THW tested the software. THW carried out data analysis. THW and MAWS wrote the manuscript. All authors were involved in the revision of the manuscript and have agreed to the final content.

Competing Interests

No competing interests were disclosed.

Grant Information

This study was supported by startup funds from the School of Life Sciences and the Biodesign Institute at Arizona State University to MAWS. Furthermore, this study was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM124827 to MAWS. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements

We thank the organizers of Hackseq 2016 for facilitating this project and supporting this collaboration; members of the Wilson Sayres lab for helpful comments; and ASU Research Computing for computational resources.

References
Abyzov, A. *et al.* (2011) CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Res.*, **21**, 974–984.
Ashley, E. A. (2016) Towards precision medicine. *Nat. Rev. Genet.*, **17**, 507–522.
Bergero, R. and Charlesworth, D. (2009) The evolution of restricted recombination in sex chromosomes. *Trends Ecol. Evol.*, **24**, 94–102.
Chang, D. *et al.* (2014) Accounting for eXentricities: analysis of the X chromosome in GWAS reveals X-linked genes implicated in autoimmune diseases. *PloS One*, **9**,

e113684.

Chen, X. *et al.* (2016) Manta: rapid detection of structural variants and indels for
germline and cancer sequencing applications. *Bioinformatics*, **32**, 1220–1222.

Cotter, D. J. *et al.* (2016) Genetic Diversity on the human X chromosome does not
support a strict pseudoautosomal boundary. *Genetics*, **203**, 485–492.

Dale, R. K. *et al.* (2011) Pybedtools: a flexible Python library for manipulating genomic
datasets and annotations. *Bioinformatics*, **27**, 3423–3424.

Ellegren, H. (2009) The different levels of genetic diversity in sex chromosomes and
autosomes. *Trends Genet.*, **25**, 278–284.

Faust, G. G. and Hall, I. M. (2014) SAMBLASTER: fast duplicate marking and structural
variant read extraction. *Bioinformatics*, **30**, 2503–2505.

Gao, F. *et al.* (2015) XWAS: a software toolset for genetic data analysis and association
studies of the X chromosome. *J. Hered.*, **106**, 666–671.

Glas, R. *et al.* (1999) Cross-species chromosome painting between human and marsupial
directly demonstrates the ancient region of the mammalian X. *Mamm. Genome*
Off. J. Int. Mamm. Genome Soc., **10**, 1115–1116.

Grüning, B. *et al.* (2017) Bioconda: a sustainable and comprehensive software
distribution for the life sciences. *bioRxiv*.

Hunter, J. D. (2007) Matplotlib: a 2D graphics environment. *Comput. Sci. Eng.*, **9**, 90–95.

International Human Genome Sequencing Consortium (2001) Initial sequencing and
analysis of the human genome. *Nature*, **409**, 860–921.

Jones, E. *et al.* (2001) SciPy: open source scientific tools for Python.

Karolchik, D. *et al.* (2004) The UCSC Table Browser data retrieval tool. *Nucleic Acids*
Res., **32**, D493–D496.

Köster, J. and Rahmann, S. (2012) Snakemake--a scalable bioinformatics workflow
engine. *Bioinformatics*, **28**, 2520–2522.

Lahn, B. T. and Page, D. C. (1999) Four evolutionary strata on the human X
chromosome. *Science*, **286**, 964–967.

Layer, R. M. *et al.* (2014) LUMPY: a probabilistic framework for structural variant
discovery. *Genome Biol.*, **15**, R84.

Li, H. (2013) Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *arXiv*, **1303.3997**.

Li, H. *et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics*,
25, 2078–2079.

Livernois, A. M. *et al.* (2012) The origin and evolution of vertebrate sex chromosomes
and dosage compensation. *Heredity*, **108**, 50–58.

Madel, M.-B. *et al.* (2016) TriXY-Homogeneous genetic sexing of highly degraded
forensic samples including hair shafts. *Forensic Sci. Int. Genet.*, **25**, 166–174.

Massey Jr., F. J. (1951) The Kolmogorov-Smirnov test for goodness of fit. *J. Am. Stat.*
Assoc., **46**, 68–78.

McKenna, A. *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.*, **20**, 1297–1303.

McKinney, W. (2010) Data structures for statistical computing in Python., *Proceedings*
of the 9th Python in Science Conference, 51–56.

Meisel, R. P. and Connallon, T. (2013) The faster-X effect: integrating theory and data.
Trends Genet., **29**, 537–544.

Mueller, J. L. *et al.* (2013) Independent specialization of the human and mouse X
chromosomes for the male germ line. *Nat. Genet.*, **45**, 1083.
Mumm, S. *et al.* (1997) Evolutionary Features of the 4-Mb Xq21.3 XY Homology
Region Revealed by a Map at 60-kb Resolution. *Genome Res.*, **7**, 307–314.
Muyle, A. *et al.* (2016) SEX-DETector: a probabilistic approach to study sex
chromosomes in non-model organisms. *Genome Biol. Evol.*, **8**, 2530–2543.
Oliphant, T. E. (2006) A Guide to NumPy. Trelgol Publishing, USA.
Page, D. C. *et al.* (1984) Occurrence of a transposition from the X-chromosome long arm
to the Y-chromosome short arm during human evolution. *Nature*, **311**, 119–123.
Pedersen, B. S. and Quinlan, A. R. (2018) Mosdepth: quick coverage calculation for
genomes and exomes. *Bioinformatics*, **34**, 867-868.
Poznik, G. D. *et al.* (2013) Sequencing Y chromosomes resolves discrepancy in time to
common ancestor of males versus females. *Science*, **341**, 562–565.
Quinlan, A. R. and Hall, I. M. (2010) BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics*, **26**, 841–842.
Rens, W. *et al.* (2007) The multiple sex chromosomes of platypus and echidna are not
completely identical and several share homology with the avian Z. *Genome Biol.*,
8, R243.
Rimmer, A. *et al.* (2014) Integrating mapping-, assembly- and haplotype-based
approaches for calling variants in clinical sequencing applications. *Nat. Genet.*,
46, 912.
Roller, E. *et al.* (2016) Canvas: versatile and scalable detection of copy number variants.
Bioinformatics, **32**, 2375–2377.
Ross, M. T. *et al.* (2005) The DNA sequence of the human X chromosome. *Nature*, **434**,
325–337.
Simmler, M. C. *et al.* (1985) Pseudoautosomal DNA sequences in the pairing region of
the human sex chromosomes. *Nature*, **317**, 692–697.
Skaletsky, H. *et al.* (2003) The male-specific region of the human Y chromosome is a
mosaic of discrete sequence classes. *Nature*, **423**, 825–837.
Sudmant, P. H. *et al.* (2015) An integrated map of structural variation in 2,504 human
genomes. *Nature*, **526**, 75.
Tarasov, A. *et al.* (2015) Sambamba: fast processing of NGS alignment formats.
Bioinformatics, **31**, 2032–2034.
Taylor, J. C. *et al.* (2015) Factors influencing success of clinical genome sequencing
across a broad spectrum of disorders. *Nat. Genet.*, **47**, 717–726.
The 1000 Genomes Project Consortium (2015) A global reference for human genetic
variation. *Nature*, **526**, 68–74.
Vicoso, B. and Charlesworth, B. (2006) Evolution on the X chromosome: unusual
patterns and processes. *Nat. Rev. Genet.*, **7**, 645–653.
Webster, T. H. *et al.* (2018) XYalign: Version 1.1.3 Zenodo.
Webster, T. H. and Wilson Sayres, M. A. (2016) Genomic signatures of sex-biased
demography: progress and prospects. *Curr. Opin. Genet. Dev.*, **41**, 62–71.
Wilson, M. A. and Makova, K. D. (2009) Genomic analyses of sex chromosome
evolution. *Annu. Rev. Genomics Hum. Genet.*, **10**, 333–354.
Wilson Sayres, M. A. (2018) Genetic Diversity on the Sex Chromosomes. *Genome Biol.*
Evol., **10**, 1064–1078.

Wilson Sayres, M. A. and Makova, K. D. (2013) Gene Survival and Death on the Human
Y Chromosome. *Mol. Biol. Evol.*, **30**, 781–787.
Wise, A. L. *et al.* (2013) eXclusion: toward integrating the X chromosome in genome-
wide association analyses. *Am. J. Hum. Genet.*, **92**, 643–647.
697