# Dissecting the neural focus of attention reveals distinct processes for spatial attention and object-based storage in visual working memory.

Nicole Hakim[1,2,*], Kirsten C. S. Adam[1,2], Eren Gunseli[3],
Edward Awh[1,2,4] & Edward K. Vogel[1,2,4]

1. Department of Psychology, University of Chicago, Chicago, Il.
2. Institute for Mind and Biology, University of Chicago, Chicago, Il.
3. Department of Psychology, Columbia University, New York, NY
4. Grossman Institute for Neuroscience, Quantitative Biology, and Human Behavior, University of Chicago, Chicago, Il.
* Corresponding author

**Running Head**: DISSECTING THE FOCUS OF ATTENTION

**Figures:** 5

**Tables:** 0

**Key words:** visual working memory, attention, electroencephalogram, contralateral delay activity

**Contributions:** All authors conceived and designed the experiments. N.H. and K.A. collected the data, performed analyses, and drafted the manuscript; N.H., K.A., E.A., and E.V. revised the manuscript.

**Funding:** Research was supported by NIMH grant 5R01 MH087214-08 and Office of Naval Research grant N00014-12-1-0972.

**Data Availability:** Datasets for all experiments will become available online on Open Science Framework upon acceptance of the manuscript or reviewer request: https://osf.io/ws3j9/?view_only=40b7f0018df44d5a862b51601d4599ed.

**Conflicts of Interest:** none

**Correspondence to:**
Nicole Hakim
University of Chicago
940 E 57th St, Chicago, IL 60637
(773)-795-4784
nhakim@uchicago.edu

**Abstract**
Working memory (WM) maintains relevant information in an accessible state, and is
composed of an active focus of attention and passive offline storage. Here, we dissect the
focus of attention by showing that distinct neural signals index the online storage of objects
and sustained spatial attention. We recorded EEG activity during two tasks that employed
identical stimulus displays while the relative demands for object storage and spatial
attention varied. Across four experiments, we found dissociable delay-period signatures for
an attention task (which only required spatial attention) and WM task (which invoked both
spatial attention and object storage). Although both tasks required active maintenance of
spatial information, only the WM task elicited robust contralateral delay activity that was
sensitive to the number of items in the array. Thus, we argue that the focus of attention is
maintained via the combined operation of distinct processes for covert spatial orienting
and object-based storage.

**Introduction**

93
Working memory (WM) facilitates the temporary maintenance of small amounts of
information so that it can be manipulated or acted upon. Contemporary theories of WM
have coalesced on variations of embedded process models (Cowan, 1999; Oberauer, 2002)
in which performance in WM tasks depends upon memory mechanisms that represent
information in two distinct states: an online, active state ("focus of attention"); and an
offline, passive state ("silent WM"). The focus of attention generally refers to information
that is currently "in mind", whereas silent WM is information that was recently within the
focus but can still be rapidly accessed. These two states of WM representation have been
proposed to be implemented at the neural level through persistent neural firing for items
within the focus (Curtis & D'Esposito, 2003) and via rapid synaptic plasticity that allows
recently attended items to quickly be reinstated (Jonides, Lacey, & Nee, 2015; Lewis-
Peacock, Drysdale, Oberauer, & Postle, 2012; Rose et al., 2016; Stokes, 2015; Wolff, Jochim,
Akyurek, & Stokes, 2017).
Here, we seek to characterize the neural mechanisms supporting the focus of
attention. Broad neuroscientific support for focus of attention-related activity has been
observed in sustained neural firing in monkey electrophysiological studies (Buschman,
Siegel, Roy, & Miller, 2011; Funahashi, Chafee, & Goldman-Rakic, 1993), uni- and multi-
variate measurements of BOLD in human fMRI studies (Todd & Marois, 2004; Xu & Chun,
2006), and sustained electrical and magnetic fluctuations in human EEG and MEG studies
(van Dijk, van der Werf, Mazaheri, Medendorp, & Jensen, 2010; Vogel & Machizawa, 2004).
Within EEG and MEG studies, two candidate measures are consistent with the focus of
attention construct. The first is alpha power (8-12hz), which shows sustained modulations
during the retention period and has been shown to contain precise spatial information
about the remembered/attended stimulus (Foster, Bsales, Jaffe, & Awh, 2017; Foster,
Sutterer, Serences, & Awh, 2016). Another candidate is the Contralateral Delay Activity
(CDA), which is a sustained negativity over the hemisphere contralateral to the positions of
to-be-remembered items. CDA amplitude is modulated by the number of items held in WM,
reaches an asymptote once WM capacity is exhausted, and predicts individual differences
in WM capacity (Unsworth, Fukuda, Awh, & Vogel, 2014; Vogel & Machizawa, 2004; Vogel,
Mccollough, & Machizawa, 2005). A prevailing view of the CDA is that it tracks the number
of task-relevant objects that are stored in WM (Balaban & Luria, 2017; Luria, Balaban, Awh,
& Vogel, 2016).
While the literatures on the CDA and alpha power have largely developed
independently, recent proposals claim that they reflect isomorphic measures of the focus of
attention. Specifically, van Dijk, et al. (2010) argued that the CDA is essentially an averaging
artifact of trial-level alpha modulation, and, therefore, reflects attention to the spatial
positions of the memoranda, rather than representations of items in WM. A similar
proposal was made by Berggren & Eimer (2016), who found that when two arrays were
presented sequentially in different hemifields, CDA amplitude tracked the positions of the
most recently seen items (but also see: Feldmann-Wüstefeld & Vogel, 2018). Such spatial
attention accounts make two broad, but untested assertions regarding neural measures of
the focus of attention. First, that sustained EEG activity reflecting the focus of attention
exclusively represents the current regions of attended space, rather than the online
maintenance of the items that occupy those regions of space (Berggren & Eimer, 2016).
Second, that such neural measures amount to a monolithic "focus of attention," rather than

a collection of distinct but overlapping mechanisms that together comprise the focus of
attention.
Here, we provide evidence that the focus of attention in WM is not a monolithic
construct, but rather, involves at least two neurally separable processes: (1) attention to
regions in space (2) representations of objects that occupy the attended regions (i.e., object
files). Alpha activity, but not the CDA, tracked attention to relevant spatial positions.
Conversely, when participants stored object representations, lateralized alpha activity that
tracked the attended positions was accompanied by robust, load-sensitive CDA activity.
These results suggest the neural focus of attention can be dissected into at least two
complementary, but distinct facets of activity: a map of prioritized space and
representation of object information in active memory.

150                  **General Methods**

**Experimental Design**
Our broad strategy was to compare delay period activity across two tasks that
employed physically identical displays but distinct cognitive requirements. We designed
distinct "Attention" and "WM" tasks to disentangle the neural correlates of hypothesized
sub-components of the focus of attention. Both tasks are known to recruit sustained spatial
attention (i.e. representation of a spatial priority map), but only the WM task required
online storage of items (i.e. representation of the objects which occupied the attended
locations). For all experiments, participants completed both a WM task and an attention
task, and the sequence of physical stimuli was identical for both tasks; the attention and
WM tasks differed only in the instructions given to participants and in the response
mapping to keys. In Experiment 1, the WM task required that participants remember the
color of the items in the sample array, whereas the attention condition required
participants to direct spatial attention towards the locations of the items in the sample
array (item color was irrelevant). Although highly similar, one key difference between the
tasks in Experiment 1 was that participants were required to remember non-spatial
features only in the WM task. To test whether the requirement to remember non-spatial
features was responsible for our findings in Experiment 1, we eliminated this difference to
make the tasks even more similar in Experiment 2; the WM task required that participants
*store the spatial positions of items* in the sample array, and the attention task required that
participants *covertly attend spatial positions* in anticipation of rare targets during the delay.

**Participants**
Experimental procedures were approved by the University of Chicago Institutional
Review Board. All participants gave informed consent and were compensated for their
participation with cash payment ($15 per hour); participants reported normal color vision
and normal or corrected-to-normal visual acuity. Participants were recruited from the
University of Chicago and surrounding community. For each sub-experiment (e.g., Exp. 1a),
we set a minimum sample size of 20 subjects (after attrition and artifact rejection). This
minimum sample size was chosen to ensure that we would be able to robustly detect set-
size dependent delay activity. Prior work employing sample sizes of 10 to 20 subjects per
experiment can robustly detect set-size dependent CDA activity (Vogel & Machizawa, 2004;
Vogel et al., 2005), and differences in CDA amplitude between novel experimental
conditions (Balaban & Luria, 2017). We chose a minimum sample size toward the upper
end of this conventional  range.
A total of 63 and 54 participants were run in Experiments 1 and 2, respectively. Due

to a technical error, EEG activity was not recorded for 3 participants in Experiment 1. In
addition, data from some participants was excluded because of excessive EEG artifacts
(<120 trials remaining in any of the four experimental conditions) or poor behavioral
performance. This left 48 subjects in Experiment 1 (28 in Experiment 1a, 20 in Experiment
1b), and 49 subjects in Experiment 2 (20 in Experiment 2a, 29 in Experiment 2b).

**EEG Acquisition**

Participants were seated inside an electrically shielded chamber, with their heads
resting on a padded chin-rest 74 cm from the monitor. We recorded EEG activity from 30
active Ag/AgCl electrodes (Brain Products actiCHamp, Munich, Germany) mounted in an
elastic cap positioned according to the International 10-20 system [Fp1, Fp2, F7, F8, F3, F4,
Fz, FC5, FC6, FC1, FC2, C3, C4, Cz, CP5, CP6, CP1, CP2, P7, P8, P3, P4, Pz, PO7, PO8, PO3, PO4,
O1, O2, Oz]. Two additional electrodes were affixed with stickers to the left and right
mastoids, and a ground electrode was placed in the elastic cap at position Fpz. Data were
referenced online to the right mastoid and re-referenced offline to the algebraic average of
the left and right mastoids. Incoming data were filtered [low cut-off = .01 Hz, high cut-off =
80 Hz, slope from low- to high-cutoff = 12 dB/octave] and recorded with a 500 Hz sampling
rate. Impedance values were kept below 10 kΩ.

Eye movements and blinks were monitored using electrooculogram (EOG) activity
and eye-tracking. We collected EOG data with 5 passive Ag/AgCl electrodes (2 vertical EOG
electrodes placed above and below the right eye, 2 horizontal EOG electrodes placed ~1 cm
from the outer canthi, and 1 ground electrode placed on the left cheek). We collected eye-
tracking data using a desk-mounted EyeLink 1000 Plus eye-tracking camera (SR Research
Ltd., Ontario, Canada) sampling at 1,000 Hz. Usable eye-tracking data were acquired for 25
out of 28 participants in Experiment 1a, 19 out of 20 participants in Experiment 1b, 17 out
of 20 participants in Experiment 2a, and 29 out of 29 participants in Experiment 2b.

**Artifact rejection**

Eye movements, blinks, blocking, drift, and muscle artifacts were first detected by
applying automatic detection criteria. After automatic detection, trials were manually
inspected to confirm that detection thresholds were working as expected. Subjects were
excluded if they had fewer than 120 total trials remaining in any of the 4 conditions. In
Experiment 1a, we rejected an average 25% of trials across all four conditions. This left us
with an average of 282 trials in WM set size 2 condition, 275 trials in the WM set size 4
condition, 302 trials in the Attention set size 2 condition, and 302 trials in the Attention set
size 4 condition.  In Experiment 1b, we rejected an average of 32% of trials across all four
conditions. This left us with an average of 291 trials in the WM set size 2 condition, 285
trials in the WM set size 4 condition, 320 trials in the attention set size 2 condition, and 320
trials in the attention set size 4 condition. In Experiment 2a, we rejected an average of 22%
of trials across all four conditions. This left us with an average of 302 trials in the WM set
size 2 condition, 301 trials in the WM set size 4 condition, 322 trials in the attention set size
2 condition, and 323 trials in the attention set size 4 condition. In Experiment 2b, we
rejected an average of 27% of trials across all four conditions. This left us with an average
of 283 trials in the WM set size 2 condition, 283 trials in the WM set size 4 condition, 298
trials in the attention set size 2 condition, and 295 trials in the attention set size 4
condition.

**Eye movements.** We used a sliding window step-function to check for eye
movements in the HEOG and the eye-tracking gaze coordinates. For HEOG rejection, we

used a split-half sliding window approach (window size = 100 ms, step size = 10 ms,
threshold = 20 μV). We only used the HEOG rejection if the eye tracking data were bad for
that trial epoch. We slid a 100 ms time window in steps of 10 ms from the beginning to the
end of the trial. If the change in voltage from the first half to the second half of the window
was greater than 20 μV, it was marked as an eye movement and rejected. For eye-tracking
rejection, we applied a sliding window analysis to the x-gaze coordinates and y-gaze
coordinates (window size = 100 ms, step size = 10 ms, threshold = 0.5$^o$ of visual angle).
**Blinks.** We used a sliding window step function to check for blinks in the VEOG
(window size = 80 ms, step size = 10 ms, threshold = 30 μV). We checked the eye-tracking
data for trial segments with missing data-points (no position data is recorded when the eye
is closed).
**Drift, muscle artifacts, and blocking.** We checked for drift (e.g. skin potentials) by
comparing the absolute change in voltage from the first quarter of the trial to the last
quarter of the trial. If the change in voltage exceeded 100 μV, the trial was rejected for drift.
In addition to slow drift, we checked for sudden step-like changes in voltage with a sliding
window (window size = 100 ms, step size = 10 ms, threshold = 100 μV). We excluded trials
for muscle artifacts if any electrode had peak-to-peak amplitude greater than 200 μV
within a 15 ms time window. We excluded trials for blocking if any electrode had at least
30 time-points in any given 200-ms time window that were within 1μV of each other.
**Analysis of Horizontal Gaze Position**
We rejected all trials that had eye movements greater than 0.5$^o$ of visual angle.
Nevertheless, participants could still move their eyes within the 0.5$^o$ of visual angle
threshold (e.g. microsaccades). To compare eye movements in the two tasks, we compared
the horizontal gaze position recorded by the eye tracker. We were most concerned with
horizontal eye movements, as these could contaminate our lateralized EEG measures. We
drift-corrected gaze position data by subtracting the mean gaze position measured 200 ms
before the pre-cue to achieve optimal sensitivity to changes in eye position (Cornelissen,
Peters, & Palmer, 2002). We then took the mean change in gaze position (in degrees of
visual angle) for left and right trials during same time-window that we used in the CDA
analysis, 400 to 1450 ms after stimulus onset. Eye gaze values from left trials were sign-
reversed so that left and right trials could be combined together. As such, positive values
indicate eye movements toward the remembered side, and negative values indicate eye
movements away from the remembered side. Importantly, not all participants had eye
tracking with adequate quality to be included in this analysis. Therefore, only 25
participants from Experiment 1a, 19 participants from Experiment 1b, 17 participants from
Experiment 2a, and 27 participants from Experiment 2b were included in the analysis.
**Analysis of Pupil Dilation**
As an additional metric of task difficulty, we compared task-evoked pupil dilation
between the WM and attention tasks. Many studies have demonstrated that task-evoked
pupil dilation correlates with cognitive load; the pupil dilates more when there are higher
attentional and working memory demands (Beatty, 1982; Steinhauer & Hakerem, 1992).
Since we were most interested in assessing the relative difficulty of the two tasks, we
collapsed the data across set size within each task. For our analysis, pupil dilation data
were baselined from 400 to 0 ms before the onset of the colored squares. Differences in
pupil dilation between the WM and attention tasks (collapsed across set sizes) were

calculated by comparing pupil size during the same time-window as is used in the CDA
analysis (400 to 1450 ms after stimulus onset). Just as in the analysis of horizontal gaze
position, not all participants had eye tracking that was good enough to be included in this
analysis. The same participants were included in both the horizontal gaze position and
pupil dilation analyses.
**Analysis of contralateral delay activity**
EEG activity was baselined from 400 ms to 0 ms before the onset of the stimulus
array. Trials containing targets for the attention task were excluded. Event-related
potentials were calculated by averaging baselined activity at each electrode across all
accurate trials within each condition (Set-Size 2 WM, Set-Size 4 WM, Set-Size 2 Attention,
and Set-Size 4 Attention). We calculated amplitude of contralateral and ipsilateral activity
for five posterior and parietal pairs of electrodes chosen a priori based on prior literature:
O1/O2, PO3/PO4, PO7/PO8, P3/P4, and P7/P8. Statistical analyses were performed on
data that was not filtered beyond the .01 – 80 Hz online data acquisition filter; we low-pass
filtered data (30 Hz) for illustrative purposes in paper figures.
**Analysis of lateralized alpha power**
EEG signal processing was performed in MATLAB 2015a (The MathWorks, Natick,
MA). We band-pass filtered trial epochs in the alpha band (8-12 Hz) using a bandpass filter
from the FieldTrip toolbox (Oostenveld, Fries, Maris, & Schoffelen, 2011;
'ft_preproc_bandpass.m') and then extracted instantaneous power by applying a Hilbert
transform ('hilbert.m') to the filtered data. Trials containing targets for the attention task
were excluded. We calculated alpha power for the same five posterior and parietal pairs of
electrodes as CDA activity: O1/O2, PO3/PO4, PO7/PO8, P3/P4, and P7/P8.
<div align="center">**Experiment 1**</div>
<div align="center">**Materials & Methods**</div>
**Stimuli & Procedures**
Stimuli in all experiments were presented on a 24-inch LCD computer screen (BenQ
XL2430T; 120 Hz refresh rate) on a Dell Optiplex 9020 computer. Participants were seated
with a chin-rest 74 cm from the screen. Stimuli were presented on a gray background, and
participants fixated a small black dot with a diameter of approximately 0.2 degrees of
visual angle. We ran two very similar versions of Experiment 1 (hereafter referred to as
Experiments 1a and 1b). The stimuli and procedures for Experiments 1a and 1b were
almost identical, with the exception of the differences noted below. Each trial began with a
blank inter-trial interval (750 ms), followed by a diamond cue (300 ms) indicating the
relevant side of the screen (right or left). This diamond cue (0.65$^{o}$ maximum width, 0.65$^{o}$
maximum height) was centered 0.65$^{o}$ above the fixation dot and was half green (RGB = 74,
183, 72) and half pink (RGB = 183, 73, 177). Half of the participants were instructed to
attend the green side and the other half were instructed to attend the pink side. After the
cue, 2 or 4 colored squares (Exp. 1: 1.1$^{o}$ by 1.1$^{o}$; Exp. 2: 1$^{o}$ by 1$^{o}$) briefly appeared in each
hemifield (150 ms) then disappeared for a 1,300 ms delay period. Squares could appear
within a subset of the display subtending 3.1$^{o}$ to the left or right of fixation and 3.5$^{o}$ degrees
above and below fixation. Colors for the squares were selected randomly from a set of 9
possible colors (Red = 255 0 0; Green = 0 255 0; Blue = 0 0 255; Yellow = 255 255 0;
Magenta = 255 0 255; Cyan = 0 255 255; Orange = 255 128 0; White = 255 255 255; Black =
1 1 1). Colors were chosen without replacement within each hemifield, and colors could be
repeated across, but not within, hemifields. On 10% of trials, two small, black lines (0.02$^{o}$

wide, 0.4º long) appeared (66.7 ms), one at the location of a colored square in the cued
hemifield, and one at the location of a colored square in the un-cued hemifield. The lines
could appear at any point during the delay period from 100 to 1200 ms after the offset of
the stimuli. Each line could be tilted 31.3 degrees to the left or 31.3 degrees to the right. At
test, a probe display appeared until response, consisting of 1 colored square in each
hemifield.
Participants in both Experiment 1a and 1b completed a WM task and an attention
task (Figure 1). Within each experiment, the sequence of physical stimuli was identical for
both tasks. Differences in procedures for Experiment 1a and 1b are described below. The
attention and WM tasks differed only in the instructions given to participants and in the
keys used to respond. Task order (attention first or WM first) and relevant cue color (pink
or green) were counterbalanced across participants. Participants completed 20 blocks of
80 trials each (1,600 trials total, 400 per condition).
335

336
**Figure 1**. Working memory and attention tasks for all experiments. At the start of each trial, a cue appeared
on the screen for 300 ms, which cued participants to attend one side of the screen. Then, an array of 2 or 4
colored squares (Exp. 1a & 1b) or circles (Exp. 2a & 2b) briefly appeared (150 ms). On 10% of trials, during
the blank retention interval (1300 ms), two small lines appeared for 66 ms between 100 and 1200 ms after
memory array offset. In Experiment 1a, one line appeared in each hemifield. In all other experiments, both
lines appeared in the same hemifield, one in an attended location and one in an unattended location. After the
retention interval, a response screen appeared with one square (Exp. 1a & 1b) or circle (Exp. 2a & 2b) in each
hemifield. In the WM task, to respond, participants reported whether the square (Exp. 1a & 1b) or circle (Exp.
2a & 2b) that reappeared in the attended hemifield was the same color (Exp. 1a & 1b) or in the same location
(Exp. 2a & 2b). Participants pressed "z" if it was the same color (Exp. 1a & 1b) or location (Exp. 2a & 2b) and
"/?" if it was different. In the Attention task, if a line was not present during the delay period, participants
pressed "spacebar." If a line was present during the delay, participants had to report the orientation of the
line that appeared in one of the cued locations. If the line was tilted left, participants pressed "z" and if it was
tilted right, participants pressed "/?." The response screen remained visible until a response was made.
351

**Working Memory Task.** Participants were instructed to remember the colors of
the presented squares in the cued hemifield and to ignore the lines that might flash during
the middle of the delay period. At test, participants were asked to identify whether the
color presented at the relevant probed location was the same as the color held in mind
(same trial) or different (change trial). The colors changed on 50% of trials. Participants
pressed the "z" key to indicate the response "same" and pressed the "/" key to indicate

"different".
**Attention Task.** Participants were instructed to maintain their attention at the
locations of the presented squares in the cued hemifield in order to identify the orientation
of a small line that appeared at one of the attended locations on 10% of trials. Participants
were instructed to press the "z" key if the line appeared and was tilted left, and the "/" key
if the line appeared and was tilted right. On 90% of trials, no line was presented, and
participants were instructed to press the "space" key to indicate that there was no target
present. The physical stimulus displays were identical to the memory task; thus, one
colored square appeared in each hemifield at the end of the attention trials. Participants
were told that the appearance of the test display indicated that it was time to respond, and
that the location and the color of the squares were irrelevant to the task.
Stimuli and procedures for Experiment 1b differed only for the target-present trials
(10% of trials). Specifically, during target-present trials, we presented both a relevant and
an irrelevant line within the cued hemifield. One line always appeared at the same location
as one of the colored squares; the second line appeared at a foil location where no colored
square had been presented (a minimum distance of 0.75 items' width from any of the
colored squares' locations). Thus, the participants were required to maintain their
attention at precise locations within the relevant hemifield so that they knew which line to
report. We reasoned that the inclusion of an irrelevant item in the cued hemifield in
Experiment 1b would encourage subjects to orient attention more precisely. However,
subsequent analyses revealed no main effect or interactions associated with the changes in
procedure between Experiment 1a and 1b. Therefore, data were collapsed across these two
versions of the task.
381                                   **Results**
**Rationale for Collapsing Experiment 1a and 1b**
In a preliminary analysis, we examined whether the small variations in the task
design between Experiments 1a and 1b had an effect on the observed results. For this
purpose, we ran repeated-measures ANOVAs for each analysis (e.g. Behavior, CDA, etc.)
with the within-subjects factors Task (WM, Attention) and Set Size (2, 4 items) and the
between-subjects factor Experiment (1a, 1b). With one exception (described next) there
was no main effect of Experiment and no interaction of Experiment with any other factor,
$p>=.49$. For the CDA analysis, there was a significant interaction of Laterality, Task, and
Experiment, $F(46)=4.28$, $p=.04$, $\eta_p^2 =.09$. To explore whether task lateralization varied
across experiments, we ran a repeated-measures ANOVA with the factors Laterality
(contra, ipsi) and Task (WM, Attention) for Experiment 1a and 1b separately. For both
experiments, there was a significant interaction of Laterality and Task, $p<.001$. Specifically,
there was a larger difference in laterality between the two tasks for Experiment 1a (M=-.50,
SD=.33) than for Experiment 1b (M=-.31, SD=.31). The triple interaction between
Laterality, Task and Experiment was not important for our central question which focused
on the cognitive processes that yield lateralized activity. Thus, the data were collapsed
across Experiments 1a and 1b.
**Behavior**
*Working memory*
WM performance (Figure 2) was converted to a capacity score, K, calculated as K = N
x (H-FA). N is the set-size; H is the hit rate; and FA is the false alarm rate (Cowan, 2001). We
only analyzed "target absent" trials, as we excluded "target present" trials (10% of total

trials) from the CDA and alpha analyses. For the WM trials, we compared performance
between set size 2 and set size 4 using a two-tailed, paired-samples t-test. There was a
significant difference in K score between set size 2 and 4, $t(47)= -3.14$, $p=.003$. Although the
effect was smaller than usual, participants remembered significantly more items on set size
4 trials (M=1.73, SD=.77) than set size 2 trials (M=1.52, SD=.40).
*Attention*
410           Accuracy (Figure 2) was essentially at ceiling for detecting whether a line was
411    present (Set Size 2: M=.99, SD=.01; Set Size 4: M=.97, SD=.02). On the rare trials in which
412    lines were presented, participants correctly reported the orientation of the target line more
413    frequently on set size 2 (M=.82, SD=.13) trials than on Set Size 4 (M=.78, SD=.15) trials,
414    $t(47)=2.56$, $p=.01$. Thus, monitoring four locations was more difficult than monitoring two
415    locations.

416
417    **Figure 2**. Behavioral performance. (A-C) Average K score in the WM task for each experiment. The
418    distribution of K scores for all participants is represented by the violin plot. Dots and light gray lines
419    represent one participant's performance. (D-F) Average accuracy in the Attention tasks for each experiment.
420

**Contralateral Delay Activity**

421
422           Recall that the WM condition required participants to direct attention to one side
423    and store the colors of the objects in WM, while the attention task required only the
424    deployment of spatial attention without storage of the objects in the sample array. Thus,
425    the central goal of Experiment 1 was to test whether CDA activity entails the storage of
426    individuated items, or whether the deployment of spatial attention is sufficient to generate
427    a CDA. As shown in Figure 3, CDA activity was observed only in the WM condition.
428           The CDA manifests as an increased negativity contralateral to the items stored in
429    working memory. Thus, to characterize the apparent differences in CDA amplitude across
430    task, we ran a 2x2x2 repeated-measures ANOVA with the factors Laterality (contra, ipsi),
431    Task (WM, attention), and Set Size (2, 4 items) on data averaged from 400 to 1450 ms after
432    stimulus onset. In this analysis, a significant effect of Laterality (i.e., greater negativity

contralateral than ipsilateral to the sample array) provides evidence for reliable CDA activity. This analysis revealed a significant main effect of Laterality, $F(1,47)=36.26$, $p<.001$, $\eta_p^2=.44$, and Set Size, $F(1,47)=12.85$, $p=.001$, $\eta_p^2=.21$. There was also a significant 2-way interaction of Laterality and Task, $p<.001$, and a significant three-way interaction of Laterality, Task, and Set Size, $F(1,47)=4.96$, $p=.03$, $\eta_p^2=.10$. No other effects were significant, $p>=.60$.

To characterize the significant interactions, we ran separate repeated-measures ANOVAs for the WM and Attention tasks, with the within-subjects factors Laterality (contra, ipsi) and Set Size (2, 4 items). For the WM task, there was a significant main effect of Set Size, $F(1,47)=6.56$, $p=.01$, $\eta_p^2=.12$, and of Laterality, $F(1,47)=66.59$, $p<.001$, $\eta_p^2=.59$. The main effect of Set Size does not provide information about the CDA component, because the effect was found in data collapsed across Laterality. The main effect of Laterality, however, demonstrates that there was reliable CDA activity in the WM condition. For the Attention task, the repeated measures ANOVA revealed a significant main effect of Set Size, $F(1,47)=10.07$, $p=.003$, $\eta_p^2=.18$. Again, this effect is not diagnostic regarding CDA activity, because the data were collapsed across laterality.  Critically, there was no main effect of Laterality in the Attention task, $F(1,47)=.95$, $p=.33$, $\eta_p^2=.02$, suggesting that deploying covert attention to the locations of the sample items was not sufficient to drive CDA activity.

Finally, the three-way interaction between Laterality, Set Size, and Task reflects the finding that set size effects in the attention task went modestly in the opposite direction of a typical CDA, with greater negativity in the set size 2 condition (M=-.07, SD=.29)  than in the set size 4 condition (M=.001, SD=.29). This was not the case for the WM condition. One unusual finding was that we did not observe a significant interaction between Set Size and Laterality on the CDA amplitude in the WM task, despite many past demonstrations that CDA activity is sensitive to changes in mnemonic load (Luria et al., 2016). CDA activity was numerically higher with the larger set size, but this was not statistically reliable. We think this may be due to unusually low WM performance for this group of subjects. While participants stored significantly more items in the set size 4 condition compared to the set size 2 condition, this difference was quite small (about 0.1 items); this modest difference in the number of items stored in the two conditions probably explains why a reliable change in CDA amplitude was not observed. Nevertheless, Experiment 2 will replicate the core empirical pattern while revealing a clear effect of WM load on CDA amplitude. Furthermore, aggregate analysis of CDA activity across Experiments 1 and 2 will also show a robust set size effect in line with past findings in the literature.

468
469
470  **Figure 3**. Contralateral delay activity. (A) CDA amplitude (μV) over time for Experiment 1. Timepoint zero
471  marks the onset of the memory array, and timepoint 1450 marks the onset of the response array. (B) Average
472  CDA amplitude (μV) for Experiment 1 during the time window of interest, 400 to 1450 ms. The distribution of
473  CDA amplitudes for all participants is represented by the violin plot. Dots and light gray lines represent one
474  participant's CDA amplitude. (C, D) CDA for Experiment 2. (E, F) CDA for All Experiments combined.
475
To summarize the CDA analysis for Experiment 1, the pattern of CDA activity was
strikingly different for the WM and Attention tasks. When participants were instructed to
encode and maintain object representations, we observed a robust CDA. However, when
participants were instructed to deploy covert attention to the locations of the same squares
to perform a demanding target discrimination task, we saw no evidence of CDA activity.
**Lateralized Alpha Power**
A broad array of studies have shown that alpha power is reduced in electrodes
contralateral to attended locations (e.g. Foster, Sutterer, Serences, Vogel, & Awh, 2017;
Kelly, Lalor, Reilly, & Foxe, 2006; Sauseng et al., 2005; Thut et al., 2006). In addition, the
scalp topography of alpha power has been shown to track the locations of stimuli held in
WM (Foster et al., 2016), even when location is not a relevant feature (Foster, Bsales, et al.,
2017). Thus, a key question for the current study is whether both the WM and attention
tasks elicited this signature of covert spatial orienting.

As Figure 4 shows, we observed the typical suppression of alpha power
contralateral to the relevant hemifield in both tasks, though it was larger in the WM task.
We confirmed these impressions with a repeated-measures ANOVA on the average alpha
power in the same window as the CDA with the factors Laterality (contra, ipsi), Task
(attention, WM) and Set Size (2, 4 items). The analysis revealed a main effect of Laterality,
$F(1,47)=22.99$, $p<.001$, $\eta_p^2 =.33$, and a significant interaction of Laterality and Task,
$F(1,47)=15.28$, $p<.001$, $\eta_p^2 =.25$. All other effects were not significant, $p>=.23$. Mean alpha
lateralization was stronger in the WM (M=-.46, SD=.45) than in the Attention (M=-.04,
SD=.29) condition. Nevertheless, follow-up two-way paired-samples t-tests revealed that
alpha power was significantly lateralized in all conditions, WM set size 2: $t(47)=-5.25$,
$p<.001$; WM set size 4: $t(47)=-4.91$, $p<.001$; Attention set size 2: $t(47)=-3.97$, $p<.001$;
Attention set size 4: $t(47)=-3.29$, $p=.002$.  Thus, in both the WM and Attention conditions,
we saw clear evidence of reduced alpha power in posterior contralateral electrodes,
consistent with the hypothesis that both the WM and Attention tasks recruited covert
spatial attention to the locations of the sample items. However, we also observed an effect
of task, with more robust lateralization of alpha power in the WM task. This pattern of
activity is somewhat similar to the effect of task on CDA amplitude. However, the aggregate
analysis below will show that whereas CDA activity tracks set size, lateralized alpha power
did not.  Therefore, even though there is a main effect of task on lateralized alpha power,
this signal is dissociable from the CDA. We further speculate about whether lateralized
alpha power may be modulated by the requirement to form object representations in the
General Discussion.

**Figure 4**. Lateralized Alpha Power. (A) Lateralized alpha power (μV²) over time for Experiment 1. Timepoint zero marks the onset of the memory array, and timepoint 1450 marks the onset of the response array. (B) Average lateralized alpha power (μV²) for Experiment 1 during the time window of interest, 400 to 1450 ms. The distribution of lateralized alpha power for all participants is represented by the violin plot. Dots and light gray lines represent one participant's alpha power. (C, D) Lateralized alpha power for Experiment 2. (E, F) Lateralized alpha power for all experiments combined.

## Assessing Cognitive Effort with Pupil Dilation

The CDA analysis for Experiment 1 revealed robust CDA activity in the WM task, but no such evidence for a lateralized negativity in the Attention task. We considered whether this difference could have been a reflection of differential cognitive effort in the WM and Attention tasks. We note that subjects were well below ceiling in discriminating the orientation of the targets in the Attention task, suggesting that it was challenging. Nevertheless, it would be valuable to compare effort between the WM and Attention tasks using a common metric. Pupil dilation provides such an opportunity. When bottom-up stimulus energy is controlled, pupil dilation has been shown to be a sensitive measure of relative cognitive effort (Kahneman & Beatty, 1966). Thus, because the WM and Attention tasks employed identical stimulus displays (when rare target trials were removed from both tasks), we were able to use pupil dilation to assess relative effort in the two tasks. To compare pupil dilation (Figure 5) between the WM and Attention tasks collapsed across set size, we ran a two-way paired-samples t-test on the averaged pupil dilation from the same

time window as the CDA. We did this separately for each task. There was no significant
difference in average pupil dilation between the WM (M=.65, SD=2.00) and Attention
(M=.86, SD=1.78) tasks, $t(43)$=-.83, $p$=.41. Thus, pupil dilation provided no indication that
stronger lateralized activity in the WM task was due to increased effort. Indeed, an
aggregate analysis below will provide evidence that this correlate of cognitive effort was
actually higher in the Attention task.

**Figure 5**. Task evoked pupil response. (A) Pupil dilation (arbitrary units) over time for Experiment 1.
Timepoint zero marks the onset of the memory array, and timepoint 1450 marks the onset of the response
array. (B) Average pupil dilation (arbitrary units) for Experiment 1 during the time window of interest, 400 to
1450 ms. The distribution of pupil dilation for all participants is represented by the violin plot. Dots and light
gray lines represent one participant's pupil dilation. Pupil dilation for Experiment 2 (C, D) and all
experiments combined (E, F).

**Horizontal Gaze Position**

Because we were comparing lateralized EEG activity across tasks, it is critical to rule
out eye movements as a potential source of differences between the tasks. We used high-
resolution eye tracking to test for such differences. To compare horizontal gaze position
between the tasks (during the same time window used to measure the CDA) we ran a 2x2
repeated measures ANOVA on horizontal eye position with the factors Task (WM,
Attention) and Set Size (2, 4 items). This analysis revealed that there was no main effect of
Task or Set Size, and no significant interaction between these two factors, $p$>=.12 for all
effects. Thus, participants moved their eyes the same amount in all conditions. In all
conditions, participants moved their eyes less than 0.017 degrees of visual angle, which is

smaller than the size of the fixation dot.
**Experiment 1 Summary**
Both the WM and Attention tasks produced reliable evidence of reduced alpha
power in electrodes contralateral to the relevant hemifield, in line with a large literature
showing that the topography of alpha power tracks the relevant spatial positions in WM
and attention paradigms. Critically, the findings with alpha contrasted sharply with those
observed with CDA activity, where lateralized activity was observed only when
participants were storing object representations in WM. These findings highlight the
possibility that CDA activity and alpha power play distinct functional roles in WM storage.
567                                                   **Experiment 2**
568                                              **Materials  & Methods**
Using identical stimulus displays, Experiment 1 showed that CDA activity was
strongly dependent on the voluntary storage of object information and not just the
requirement to deploy covert spatial attention. When subjects stored the objects in the
sample display, the lateralized CDA signal was easily observed. By contrast, the precise
deployment of covert attention to the same physical displays yielded no detectable CDA
activity. Although this empirical pattern is consistent with the hypothesis that CDA activity
is tied to item storage *per se*, the WM and Attention tasks differed in more than one way.
Specifically, the WM task required participants to remember both the *color* and the location
of the sample stimuli, whereas only the location of those stimuli was relevant in the
Attention task. Thus, the goal of Experiment 2 was to more precisely manipulate item
storage requirements by holding constant the relevant feature in the WM and Attention
tasks. To this end, the WM condition required participants to store the locations of the
items in the sample display and to detect whether any item's position had changed at the
end of the delay period. In the attention task, participants were instructed to maintain
covert attention at the positions of the sample items to facilitate the detection and
discrimination of rare line targets. Critically, while both tasks required participants to
attend the location of the sample items, only the WM task encouraged subjects to store
those object representations in memory. In the Attention task, by contrast, subjects
attended the position of those items in anticipation of a separate target stimulus. Thus, if
CDA activity is contingent on the online storage of individuated object representations,
then only the WM task should elicit robust CDA activity.
**Stimuli & Procedures**
We ran two versions of Experiment 2 that were almost identical except that sample
stimuli were either black circles (Experiment 2a) or colored squares (Experiment 2b). In
addition, there were some minor differences in stimulus sizes. These differences are
described in detail below. Because preliminary analyses revealed no impact of these
differences, subsequent analyses collapsed across the two versions of the procedure.
**Experiment 2a**. Stimuli were similar to Experiment 1b with the following
exceptions. Participants were presented with 2 or 4 black circles ($0.611^o$ diameter; RGB= 1
1 1) in each hemifield with a minimum of $1.53^o$ degrees (2.5 objects)  between each item.
These circles could appear within a subset of the display subtending 2.44 degrees to the left
or right of fixation and 3.06 degrees above and below fixation.  On target-present trials, the
two small lines that were presented briefly during the retention interval were $0.04^o$ wide
and $0.76^o$ long.
**Experiment 2b**. Stimuli were similar to Experiment 2a with the following

exceptions. Participants were presented with 2 or 4 colored circles (0.84$^o$ diameter) in each
hemifield with a minimum of 2.10$^o$ (2.5 objects) between each item.  Colors for the circles
were randomly selected without replacement within each hemifield from a set of 10
possible colors (Red = 255 0 0; Green = 0 255 0; Blue = 0 0 255; Yellow = 255 255 0;
Magenta = 255 0 255; Cyan = 0 255 255; Orange = 255 128 0; Brown = 102 51 0; White =
255 255 255; Black = 1 1 1). On target-present trials, the two small lines that were
presented briefly during the retention interval were 0.04$^o$ wide and 0.99$^o$ long.
Just as in Experiment 1, participants in both Experiment 2a and 2b completed both
an attention task and a WM task. The timing of trial events was identical to Experiment 1b.
The key difference between Experiment 1 and 2 is that participants completed a spatial
change detection task rather than a color change detection task.
**Working Memory Task.** Procedures for the WM task were very similar to the
procedure from Experiment 1, except that participants were asked to identify whether the
location of the presented circle in the attended hemifield was in the same or different
location as the any of the original circles.
**Attention Task.** Procedures and instructions for the attention task were identical to
Experiment 1b. Only the visual stimuli differed, so as to match the visual stimuli presented
in the WM task.

<div align="center">

**Results**

</div>

### Rationale for Collapsing Experiment 2a and 2b

In a preliminary analysis, we examined whether the small variations in task design
between Experiments 2a and 2b had an effect on the observed results. For this purpose, we
ran repeated-measures ANOVAs for each analysis (e.g. Behavior, CDA, etc.) with the within-
subjects factors Task (WM, Attention) and Set Size (2, 4 items) and the between-subjects
factor Experiment (2a, 2b). There was no main effect of Experiment for any of the analyses,
$p>=.10$, and no interaction of Experiment with any other factor ($p>.10$). Therefore,
subsequent analyses collapsed the data across 2a and 2b.

### Behavior

As we did for Experiment 1, we separately analyzed performance for the WM and
attention tasks. For the WM task, there was a significant difference in K score between Set
Size 2 and 4, $t(48)=-4.1$, $p<.001$. Participants remembered significantly more items on set
size 4 (M=1.43, SD=.69) than set size 2 trials (M=1.23, SD=.49).
In the Attention task, participants had a high rate of detecting whether a line was
present (Set Size 2: M=.97, SD=.02; Set Size 4: M=.96, SD=.02). To compare performance
between set size 2 and 4 when participants had to discriminate the orientation of the target
line, we used a two-tailed, paired-samples t-test. Participants correctly reported the
orientation of the target line more frequently on Set Size 2 (M=.75, SD=.10) than on Set Size
4 (M=.69, SD=.11) trials, $t(48)=4.00$, $p<.001$. Thus, monitoring four locations was more
difficult than monitoring two locations.

### Contralateral Delay Activity

As shown in Figure 3, CDA activity was different across the WM and Attention tasks,
with more robust lateralized activity and effects of set size in the WM task. This impression
was confirmed with a repeated-measure ANOVA with the factors Task (WM, Attention),
Laterality (contra, ipsi) and Set Size (2, 4 items). The repeated measures ANOVA revealed
significant main effects of Laterality, $F(1,48)=39.38$, $p<.001$, $\eta p^2 =.45$, and Set Size,
$F(1,48)=21.03$, $p<.001$, $\eta_p^2 =.31$. Critically, there was a reliable interaction between

Laterality and Task $F(1,48)=23.68$, $p<.001$, $\eta_p^2 =.33$, showing that CDA activity differed
between the WM and Attention tasks. There were also significant interactions between
Task and Set Size, $F(1,48)=5.72$, $p=.02$, $\eta_p^2 =.11$, and between Laterality and Set Size
$F(1,48)=16.88$, $p<.001$, $\eta_p^2 =.26$; because these interactions refer to data collapsed across
Laterality or Task, they are not informative regarding the central question of which task
requirements generate CDA activity. Finally, given our hypothesis that CDA activity may be
specifically tied to the formation of object representations, we also predicted a triple
interaction between Laterality, Task and Set Size, because load-dependent CDA activity
should be more pronounced in the WM task. This interaction was trending but did not
reach conventional thresholds for significance, $F(1,48)=3.72$, $p=.06$, $\eta_p^2 =.07$. We re-visit
this question below with a more sensitive aggregate analysis of Experiments 1 and 2
together.
662            To further delineate the key interaction between Laterality and Task, we ran a
follow-up paired-samples t-tests, collapsed across set size. Examining the difference
between contralateral and ipsilateral activity in each task, we observed reliable effects of
Laterality for both the WM and Attention tasks, $p<=.03$ for all conditions. Critically,
however, the effect of Laterality was substantially larger in the WM task, (M=-.49, SD=.50)
than in the Attention task (M=-.17, SD=.35), $t(48)=-4.88$, $p<.001$. Thus, Experiment 2
replicated the broad empirical pattern in Experiment 1; CDA activity was far stronger when
participants were instructed to store object representations in WM than when they were
instructed to attend those locations in anticipation of upcoming targets.
671            The marked difference in CDA activity between the WM and Attention tasks is
particularly striking in light of their strong similarity. Indeed, while the labels for the WM
and Attention tasks highlight their respective storage and selection requirements, both
tasks actually require the sustained maintenance of spatial information. In the WM task,
participants held the positions of the sample items in mind to facilitate change detection,
while in the Attention task, participants held a sustained focus of spatial attention at
specific positions. Nevertheless, CDA activity was more than twice as high in the WM task,
motivating the conclusion that CDA activity is tied to object representations held in WM *per*
*se*, and not simply the deployment of covert attention – even when an attention task
requires the maintenance of spatial information. If this is correct, however, it raises the
question of why we saw reliable CDA activity in the Attention task. Our working hypothesis
is that this modest effect of Laterality could reflect the occasional storage of object
representations during the Attention task, but the current study does not provide a clear
way to test this possibility. Nevertheless, while this question can't be fully answered,
Experiment 2 did replicate the striking divergence in CDA activity between the WM and
Attention tasks, in line with the hypothesis that this neural signal may be specifically tied to
item storage in visual WM.
**Lateralized Alpha Power**
689            Just as in Experiment 1, we observed the typical suppression of alpha power
contralateral to the relevant hemifield in both tasks. Once again, this effect was larger in the
WM task. We confirmed these impressions with a repeated-measures ANOVA on average
alpha power with the factors Laterality (contra, ipsi), Task (attention, WM) and Set Size (2,
4 items). The analysis revealed a significant main effect of Laterality, $F(1,48)=24.17$,
$p<.001$, $\eta_p^2 =.34$, and a significant interaction between Laterality and Task, $F(1,48)=13.38$,
$p<.001$, $\eta_p^2 =.22$. Follow-up two-way paired samples t-tests on the difference between

contralateral and ipsilateral activity for each condition revealed significant alpha power
suppression in all conditions, $p<=.001$ for both set sizes in both the WM and Attention
tasks. No other effects were significant, $p>=.10$.
To characterize the interaction of Laterality and Task, we collapsed Laterality into a
difference wave (contra minus ipsi). We then compared this difference wave for the
Attention and WM conditions, averaged across Set Size, with a two-way paired samples t-
test. This analysis revealed a significant difference in alpha power lateralization between
the WM and Attention tasks, $t(48)=-3.66$, $p=.001$. Alpha power was more suppressed in
contralateral relative to ipsilateral electrodes during the WM (M=-9.16, SD=13.38) than
during the Attention (M=-4.01, SD=6.73) task.
To summarize, in both tasks, we found clear evidence of sustained covert orienting
towards the relevant hemifield. This makes the important point that the lack of a robust
CDA in the Attention task is not due to a failure to maintain attention towards the relevant
hemifield. In addition, replicating the finding from Experiment 1, we observed reliably
stronger lateralization of alpha power when participants were instructed to store the
sample items in WM. Thus, both neural signals suggest a distinction between the
maintenance of items in working memory, and the maintenance of spatial attention at the
position of those items.
**Horizontal Gaze Position**
To determine whether horizontal gaze position varied across tasks, we ran a 2x2
repeated-measures ANOVA with the factors Task (WM, Attention) and Set Size (2, 4 items).
This analysis revealed a significant main effect of Task, $F(1,42)=4.78$, $p=.03$, $\eta_p^2=.10$.
Participants moved their eyes more during the WM (M=.03, SD=.03) than the Attention
(M=.02, SD=.02) task. We would like to point out that the difference in eye movements
between these two tasks is only about .01 degrees, which is smaller than the diameter of
the fixation point. Nevertheless, to determine whether these differences in horizontal gaze
position drive the differences in the CDA that we observe, we compared horizontal gaze
position during the time window when the CDA initially emerges, 400 to 925 ms after
stimulus onset. The 2x2 repeated measures ANOVA with factors Task (WM, Attention) and
Set Size (2, 4 items) for the average horizontal gaze position during this time window
revealed that there was no significant main effect of Task or Set Size and no interaction of
these two factors, $p>=.06$ for all effects. This indicates that during the time window when
the CDA is ramping up, participants moved their eyes the same amount in all conditions.
Therefore, any differences in the CDA that we observe are not driven by differences in
horizontal eye movements. As a follow-up analysis we also analyzed horizontal eye
movement averaged over the end of the trial, 926 to 1450 ms, with another 2x2 ANOVA
with factors Task (WM, Attention) and Set Size (2, 4 items). This analysis revealed that
differences in horizontal eye movements between the WM and Attention tasks emerged
toward the end of the trial, significant main effect of Task, $F(1,43)=4.40$, $p=.04$, $\eta_p^2=.09$.  All
other effects and interactions were not significant, $p>=.06$.
**Pupil Dilation**
As with Experiment 1, to compare the task-evoked pupil response between the WM
and Attention tasks collapsed across set size, we ran a two-way paired-samples t-test. This
analysis revealed greater task-evoked pupil dilation in the Attention task (M=.60, SD=.29)
than in the WM task (M=-.07, SD=2.36), $t(43)=-3.65$, $p=.001$, suggesting that the Attention
task elicited greater cognitive effort. Therefore, the robust CDA in the WM condition is

unlikely to reflect greater cognitive effort in the WM than the Attention task.

**Aggregate analysis of Experiments 1 and 2**

In the analysis reported below, we aggregated data across Experiments 1 and 2 to
provide the most power for understanding the distinctions between the WM and Attention
tasks. This analysis included 97 participants, and clearly reinforces the robustness of the
key empirical patterns. In this aggregate analysis, we focus on CDA, alpha power, and pupil
size because aggregate analyses of behavior and eye position were not central to our
primary arguments and merely echoed all the main effects and interactions that were
observed in the individual experiments.

**Preliminary analysis of the effect of Experiment**

In a preliminary analysis, we examined whether the small variations in task design
between Experiments 1 and 2 had an effect on the observed results. For this purpose, we
ran repeated-measures ANOVAs for each analysis (i.e., CDA, alpha power, and pupil size)
with the within-subjects factors Task (WM, Attention) and Set Size (2, 4 items) and the
between-subjects factor Experiment (1, 2). For all analyses, there was no main effect of
Experiment, $p>=.16$. Therefore, it was justified to collapse data across the two experiments.
For the horizontal gaze position and the lateralized alpha analyses, none of the
factors significantly interacted with Experiment, $p>=.19$. However, for the pupil dilation
analysis, there was a significant interaction of Task and Experiment, $F(1,86)=10.76$, $p=.002$,
$\eta p^2 =.11$. This significant interaction is explained by greater pupil dilation in the Attention
task than in the WM task in Experiment 2, but not in Experiment 1.
For the CDA analysis, there was a significant 3-way interaction of Laterality, Set Size,
and Experiment, $F(1,95)=6.73$, $p=.01$, $\eta_p^2 =.07$. To further delineate this three-way
interaction, we ran follow-up ANOVAs with the factors Laterality (contra, ipsi) and Set Size
(2, 4 items) for Experiment 1 and Experiment 2 separately. These follow-up analyses
revealed that there was a significant interaction of Laterality and Set Size for Experiment 2,
$F(1,48)=16.88$, $p<.001$, $\eta_p^2 =.26$, but not for Experiment 1, $F(1,47)=.08$, $p=.78$, $\eta_p^2 =.002$.

**Contralateral Delay Activity**

Using all data from Experiments 1 and 2 together, we ran a repeated-measures
ANOVA with the factors Task (WM, Attention) and Set Size (2, 4 items). This analysis
revealed significant main effects of Laterality, $F(1,96)=74.41$, $p<.001$, $\eta_p^2 =.44$, and Set Size,
$F(1,96)=33.27$, $p<.001$, $\eta_p^2 =.26$. Because these main effects were collapsed across Task,
they are not informative for our central question of how storage-related neural signals
differ across tasks. Thus, the first important finding was a significant 2-way interaction
between Laterality and Task, $F(1,96)=81.27$, $p<.001$, $\eta_p^2 =.46$ that reflected a greater
laterality effect in the WM than in the Attention task. To confirm this impression, we ran a
follow-up 2-way paired-samples t-test that compared contralateral to ipsilateral activity
separately for the WM and Attention and each set size (e.g. WM ss2, WM ss4, ATT ss2, ATT
ss4). This analysis revealed that the CDA was significantly more lateralized in the WM (Set
Size 2: M =-.38, SD=.44; Set Size 4: M=-.54, SD=.54) than in the Attention (Set Size 2: M=-.09,
SD=.34; Set Size 4: M=-.10, SD=.37) task for both set sizes (Set Size 2: $t(98)=-6.71$, $p<.001$; Set
Size 4: $t(98)=-8.57$, $p<.001$). We note, however, that there was reliable lateralized activity for
both tasks, $p<=.007$.
The most important opportunity afforded by this aggregate analysis of CDA activity
was that we now had sufficient power to examine whether increased mnemonic load had a
differential effect on CDA activity in the WM and Attention task. Indeed, there was a reliable

triple interaction between Task, Laterality and Set Size $F(1,96)=8.75$, $p=.004$, $\eta_p^2 =.08$. As
Figure 3 shows, CDA activity was set size dependent in the WM task, but not in the
Attention task. To verify this impression, we ran separate follow-up repeated-measures
ANOVAs for each task (WM and Attention) with the factors Laterality (contra, ipsi) and Set
Size (2, 4 items). This analysis revealed that there was a significant interaction of Laterality
and Set Size for the WM task, $F(1,96)=14.39$, $p<.001$, $\eta_p^2 =.13$, but not the Attention task,
$F(1,96)=.07$, $p=.79$, $\eta_p^2 =.001$. Thus, while data from the WM task showed that CDA
amplitude was larger for set size 4 (M=-.55, SD=.54) than set size 2 (M=-.39, SD=.44), data
from the Attention task showed no evidence of a difference in CDA amplitude between Set
Size 2 (M=-.10, SD=.34) and Set Size 4 (M=-.11, SD=.37).
To summarize, the aggregate analysis confirmed that CDA activity was substantially
stronger in the WM than in the Attention task. Moreover, this analysis had sufficient power
to show that CDA activity tracked the increase in mnemonic load from two to four items,
while the CDA signal in the Attention task – in addition to being over four times smaller
than in the WM task – showed no effect of mnemonic load at all, a defining feature of the
CDA. This core result motivates our conclusion that CDA activity is directly linked with the
online maintenance of object representations in WM, and not the deployment of attention
to the positions of the sample items.
**Lateralized Alpha Power**
The aggregate analysis of Experiments 1 and 2 confirmed the results of the
individual experiments. As Figure 4 shows, we observed the typical suppression of alpha
power contralateral to the relevant hemifield in both tasks, though it was larger in the WM
task. Just as with the individual experiments, we confirmed these impressions with a
repeated-measures ANOVA on the average alpha power with the factors Laterality (contra,
ipsi), Task (attention, WM) and Set Size (2, 4 items). This analysis revealed a significant
main effect of Laterality, $F(1,96)=45.57$, $p<.001$, $\eta_p^2 =.32$, and a significant interaction
between Laterality and Set Size, $F(1,96)=9.75$, $p=.002$, $\eta_p^2 =.09$. Paired t-tests confirmed
that this interaction reflects a stronger lateralization of alpha power in the set size 2
condition (M=-12.24, SD=17.17) than in the set size 4 condition (M=-10.04, SD=16.06)
($t(96)=-3.123$, $p=.002$). Thus, the strength of lateralized alpha activity varied with the
number of stored or attended positions in both the WM and Attention tasks. Critically,
however, the effect of set size on lateralized alpha power was in the opposite direction
from the effect we observed with CDA activity. CDA activity was stronger for set size 4 than
for set size 2 whereas alpha lateralization was stronger for set size 2 than for set size 4.
These findings support the hypothesis that CDA and alpha activity reflect distinct aspects of
online storage in visual WM.
The aggregate analysis also revealed a significant interaction between Laterality and
Task, $F(1,96)=27.22$, $p<.001$, $\eta_p^2 =.22$, that reflected the greater lateralization of alpha
power in the WM than in the Attention task. This impression was confirmed with a two-
way paired samples t-test that revealed a significant difference in alpha power
lateralization between the WM (M=-7.79, SD=11.61)  and Attention (M=-3.35, SD=5.69)
tasks, $t(96)=-5.22$, $p<.001$. Critically, both tasks showed clear evidence of lateralized alpha
power in both set sizes ($p<.001$ for all conditions), confirming that covert attention was
deployed to the position of sample items in a sustained fashion in both tasks. The greater
lateralization of alpha power in the WM than Attention task is a robust empirical pattern
that is present in both experiments and in the aggregate analysis. Though we did not expect

this pattern *a priori*, this reliable difference in alpha lateralization between the two tasks
may reflect a direct influence of online object representations on the deployment of spatial
attention.
**Pupil Dilation**
We argue that the WM task encouraged online storage of object representations
while the Attention task did not. Thus, the restriction of load-dependent CDA activity to the
WM task could reflect a direct link between the CDA and item storage in WM. A clear
alternative hypothesis, however, is that the WM task may differ from the Attention task in
terms of the intensity or effort applied to the task rather than the specific cognitive
operations that were invoked. While accuracy was similar (and off of ceiling) in the two
tasks, this does not provide strong evidence for equivalent effort. Fortunately, pupil
dilation measurements have been shown to provide a sensitive index of cognitive effort
and arousal when bottom-up stimulus factors are controlled. Thus, we ran a two-way
paired-samples t-test to examine whether pupil size differed during the time window in
which CDA activity was measured. This analysis revealed a greater level of pupil dilation in
the Attention task (M= .63, SD=1.89) compared to the WM task (M= .29, SD=2.20), $t(87)$=-
3.13, $p$=.002, suggesting that the Attention task recruited greater levels of cognitive effort.
Thus, our finding that CDA activity was far larger in the WM task cannot be explained by
increased effort in the WM task. Indeed, pupil analysis of the aggregated data suggests that
the WM task was the easier of the two. These findings argue for a difference in the nature of
the cognitive operations evoked by the WM and Attention tasks, rather than in the degree
to which similar operations were carried out.
**Aggregate Analysis Summary**
With 97 subjects, this aggregate analysis reinforced the key conclusions of
Experiments 1 and 2, and provided strong statistical power for documenting how neural
activity differed between the WM and Attention tasks. CDA activity was more than four
times larger in the WM task than in the Attention task. Moreover, CDA activity in the WM
task clearly tracked changes in mnemonic load whereas CDA activity in the Attention task
showed no evidence of load sensitivity. Thus, given that these tasks employed identical
stimulus displays, we conclude that CDA activity may be directly tied to the unique *object*
*representation* requirements in the WM task and not covert attentional orienting to the
sample array positions.
The WM and Attention tasks differ in terms of object storage, but past work suggests
that both WM and attention tasks may call upon a common spatial attention process that
elicits orderly changes in the scalp topography of alpha power. In addition to past studies
showing the broad involvement of alpha activity across a wide range of attention and
memory paradigms (Canolty & Knight, 2010; Fries, 2005; Klimesch, 2012), more recent
work has also established that the topography of alpha activity on the scalp can be used to
precisely track the locus of covert attention (Foster, Sutterer, et al., 2017; Rihs, Michel, &
Thut, 2007) and locations stored in WM (Foster, Bsales, et al., 2017; Foster et al., 2016). In
line with this work, there was clear evidence from both the WM and Attention tasks that
alpha power in posterior electrodes was reduced contralateral to the sample array.
Importantly, the aggregate analysis also had enough power to reveal a reliable effect of set
size on the strength of alpha lateralization, such that greater lateralization was observed in
the set size 2 condition compared to the set size 4 condition. This effect may not generalize,
however, as some previous research has not found an effect of set size on lateralization

(Fukuda, Mance, & Vogel, 2015), while others have found greater lateralization for larger
set sizes (Sauseng et al., 2009). Nevertheless, the fact that CDA activity showed the opposite
pattern, with higher CDA activity for the larger set size, highlights the possibility that these
two neural signals (measured from within the same set of electrodes) index distinct aspects
of maintenance within the focus of attention.
Finally, we examined whether these differences in neural activity were a
consequence of differential effort or arousal in the two tasks. Because the stimulus displays
were identical, we were able to use task-evoked pupil dilation to obtain a sensitive metric
of cognitive effort and arousal. The aggregate analysis revealed that the Attention task
elicited reliably larger pupil size than the WM task, suggesting that the Attention task
elicited greater effort. In line with this conclusion, we also note that while behavioral data
from the Attention task showed that monitoring four locations was more difficult than
monitoring two locations, CDA activity in the Attention task was unaffected by set size.
Together, these findings argue strongly against the hypothesis that stronger delay period
signals in the WM task were a consequence of greater cognitive effort.
895                                            **Discussion**
The focus of attention refers to the small set of mental representations that can be
held in an *online* or readily accessible state. Motivated by its central role in intelligent
behaviors, there has been a longstanding effort to elucidate the neural signals that track the
contents of this internally attended information. This body of work has tended to treat the
focus as a monolithic entity, but here we extend the growing evidence that the focus of
attention may be implemented via multiple component processes playing distinct
functional roles: one that represents currently prioritized space (alpha); and another that
reflects item storage within the focus of attention (CDA).  This proposal converges with
other findings that suggest a dissociation between spatial attention and WM storage (Tas et
al., 2016; Sheremata et al 2018)
**CDA activity and lateralized alpha power: Distinct components of the focus of**
**attention**
van Dijk et al. (2010) proposed that asymmetric modulations of alpha power at the
trial-level can generate a CDA-like negative slow wave in an event-related average.
However, there is growing evidence that these two measures can be clearly dissociated. For
example, Fukuda et al. (2016) used a lateralized change detection task where they cued
participants to one side of the screen, but had a longer than normal (1,000 ms) SOA
between the cue and the memory array. During this blank cue period, participants knew
which hemifield would contain memory items, but no items had yet appeared. During this
time, there was robust alpha power lateralization but no CDA. However, after the memory
array appeared, the CDA and alpha power lateralization appeared in concert during the
memory maintenance period (1,000 ms). These results suggest lateralized alpha power,
and thus attention, can be shifted to empty space, but that the CDA necessitates object
storage (see also Fukuda, Mance & Vogel, 2015).
**CDA activity as an index of item-based storage in working memory**
What was the critical difference between the WM and Attention tasks? Despite the
fact that they employed identical stimulus displays, the amplitude of the CDA was more
than four times larger in the WM than in the Attention task, and only the WM task elicited
load-dependent CDA activity. Both tasks elicited covert orienting to the positions of the
items in the sample array, as shown by sustained lateralized alpha power modulations.

Moreover, despite distinct monikers, *both* tasks required the sustained maintenance of
spatial information across a blank delay. This storage requirement is obvious for the WM
change detection task. But even for the Attention task, subjects must have maintained the
cued positions so that they could be distinguished from lures. Indeed, in Experiment 2,
change detection in the WM task required precisely the same spatial discriminations as did
target identification in the Attention task. Thus, we propose that the critical difference
between the WM and Attention tasks was that the WM task encouraged the continued
*representation of the items* in the sample array, while in the Attention task participants
directed spatial attention to those positions without maintaining the items themselves.
**CDA activity as a neural index of object file maintenance**
Our interpretation of the CDA as an index of continued representations of object
files critically hinges on a distinction between the maintenance of items in working
memory and the maintenance of spatial information without an accompanying item
representation. While some may view this as provocative, recent work has shown
dissociable patterns of activity in parietal lobe between WM and spatial attention demands
(Sheremata, Somers, & Shomstein, 2018). Additionally, we note that there is a longstanding
precedent for a distinction between the representation of an object and the representation
of the features or identifying labels associated with that object. Kahneman, Treisman, &
Gibbs (1992) elucidated this idea with the *object file* construct which proposes two
separable stages of processing. The first involves the parsing of the scene into a set of
individuated items that are indexed based on their spatial and temporal coordinates.
Subsequently, the specific feature values (e.g., color and orientation) are processed and
incorporated into the associated object file. Thus, object files anchor the episodic
representation in a specific time and place, and are distinct from the specific feature values
that are bound together by virtue of an object file. In the present context, an intriguing
possibility is that CDA activity indexes the maintenance of object files in WM. This proposal
is consistent with recent work showing that the CDA is sensitive to objecthood cues
(Balaban & Luria, 2016) and tracks the number of encoded objects, not the number of
features within objects (Luria & Vogel, 2011). Thus, even though the Attention task
required the sustained maintenance of location information, CDA activity was minimal or
absent (and insensitive to mnemonic load) because the task did not encourage the
maintenance of the object files that were created during the encoding of the sample array.
**Open question on the impact of "object files" on the allocation of spatial attention**
In this series of experiments, lateralized alpha power was a useful tool to illustrate
that participants sustained their attention to the cued side even when the CDA was
completely absent (Exp 1). However, we also observed a main effect of our task
manipulation on lateralized alpha power. When task demands required participants to
encode object representations, alpha power was significantly more lateralized than when
they only had to sustain their attention to empty space. Though we did not predict this
pattern *a priori*, it was reliable in both experiments. This suggests that, like the CDA,
lateralized alpha power respects the dissociation between forming object representations
and maintaining a spatial priority map. One possible interpretation of this effect is that
object representations serve as "anchors" for the allocation of spatial attention, thus
amplifying the effects of attention and leading to increased alpha power lateralization.
While future work is needed to investigate the complex interrelationship between
lateralized alpha power and task demands, the present work clearly suggests that

lateralized alpha power does not directly generate, and is dissociable from, the CDA.

**Conclusions**

A growing body of evidence has shown that CDA activity and alpha power are tightly
linked with the maintenance of information in the focus of attention. Here, we present new
evidence that these two neural signals represent distinct facets of this online system. A
topographic distribution of alpha power indexes the current locus of spatial attention, a
process that is integral to both visual selection and the voluntary storage of items in WM.
By contrast, CDA activity tracks the active maintenance of object files, the item-based
representations that allow observers to integrate the ensemble of features and labels that
are associated with visual objects. The dissociable activity of the CDA and alpha power
suggests that the focus of attention is composed of at least two distinct but complementary
neural processes, a conclusion with strong implications for both cognitive and neural
models of this online storage system.

**References**

Balaban, H., & Luria, R. (2016). Integration of Distinct Objects in Visual Working Memory
Depends on Strong Objecthood Cues Even for Different-Dimension Conjunctions.
*Cerebral Cortex*, *26*(5), 2093–2104.
Balaban, H., & Luria, R. (2017). Neural and Behavioral Evidence for an Online Resetting
Process in Visual Working Memory. *The Journal of Neuroscience*, *37*(5), 1225–1239.
Beatty, J. (1982). Task-Evoked Pupillary Responses, Processing Load, and the Structure of
Processing Resources, *91*(2), 276–292.
Berggren, N., & Eimer, M. (2016). Does Contralateral Delay Activity Reflect Working
Memory Storage or the Current Focus of Spatial Attention within Visual Working
Memory? *Journal of Cognitive Neuroscience*, *28*(12).
Buschman, T. J., Siegel, M., Roy, J. E., & Miller, E. K. (2011). Neural substrates of cognitive
capacity limitations. *Proceedings of the National Academy of Sciences*, *108*(27), 1–4.
Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling.
*Trends in Cognitive Sciences*, *14*(11), 506–515.
Cornelissen, F. W., Peters, E. M., & Palmer, J. (2002). The Eyelink Toolbox: Eye tracking with
MATLAB and the Psychophysics Toolbox. *Behavior Research Methods, Instruments, and*
*Computers*, *34*(4), 613–617.
Cowan, N. (1999). *An embedded-process model of working memory*.
Cowan, N. (2001). The magical number 4 in short term memory. A reconsideration of
storage capacity. *Behavioral and Brain Sciences*, *24*(4), 87–186.
Curtis, C. E., & D'Esposito, M. (2003). Persistent activity in the prefrontal cortex during
working memory. *Trends Cogn. Sci.*, *7*, 415–423.
Feldmann-Wüstefeld, T., & Vogel, E. K. (2018). Neural Evidence for the Contribution of
Active Suppression During Working Memory Filtering. *Cerebral Cortex*.
Foster, J. J., Bsales, E. M., Jaffe, R. J., & Awh, E. (2017). Alpha-Band Activity Reveals
Spontaneous Representations of Spatial Position in Visual Working Memory. *Current*
*Biology*, *27*(20), 3216–3223.e6.
Foster, J. J., Sutterer, D. W., Serences, J. T., & Awh, E. (2016). The topography of alpha-band
activity tracks the content of spatial working memory. *Journal of Neurophysiology*,
*115*(1), 168–77.
Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K., & Awh, E. (2017). Alpha-Band
Oscillations Enable Spatially and Temporally Resolved Tracking of Covert Spatial
Attention. *Psychological Science*, *28*(7), 929–941.
Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through
neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.
Fukuda, K., Mance, I., & Vogel, E. K. (2015). Alpha Power Modulation and Event-Related
Slow Wave Provide Dissociable Correlates of Visual Working Memory. *Journal of*
*Neuroscience*, *35*(41), 14009–14016.
Funahashi, S., Chafee, M. V., & Goldman-Rakic, P. S. (1993). Prefrontal neuronal activity in
rhesus monkeys performing a delayed anti-saccade task. *Nature*, *365*(6448), 753–756.
Jonides, J., Lacey, S. C., & Nee, D. E. (2015). Processes of Working Memory in Mind and
Brain. *Society*, *14*(1), 12–15.
Kahneman, D., & Beatty, J. (1966). Pupil Diameter and Load on Memory. *Science*.
Kahneman, D., Treisman, A., & Gibbs, B. J. (1992). The reviewing of object files: Object-

specific integration of information. *Cognitive Psychology*, *24*(2), 175–219.

Kelly, S. P., Lalor, E., Reilly, R., & Foxe, J. J. (2006). Increases in Alpha Oscillatory Power Reflect an Active Retinotopic Mechanism for Distracter Suppression During Sustained Visuospatial Attention. *Journal of Neurophysiology*, *95*(6), 3844–3851.

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences*, *16*(12), 606–617.

Lewis-Peacock, J. A., Drysdale, A. T., Oberauer, K., & Postle, B. R. (2012). Neural Evidence for a Distinction between Short-term Memory and the Focus of Attention. *Journal of Cognitive Neuroscience*, *24*(1), 61–79.

Luria, R., Balaban, H., Awh, E., & Vogel, E. K. (2016). The contralateral delay activity as a neural measure of visual working memory. *Neuroscience and Biobehavioral Reviews*, *34*(5), 352–359.

Luria, R., & Vogel, E. K. (2011). Shape and color conjunction stimuli are represented as bound objects in visual working memory. *Neuropsychologia*, *49*(6), 1632–1639.

Oberauer, K. (2002). Access to Information in Working Memory: Exploring the Focus of Attention. *Journal of Experimental Psychology: Learning Memory and Cognition*, *28*(3), 411–421.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*(1–9).

Rihs, T. A., Michel, C. M., & Thut, G. (2007). Mechanisms of selective inhibition in visual spatial attention are indexed by alpha-band EEG synchronization. *European Journal of Neuroscience*, *25*(2), 603–610.

Rose, N. S., LaRocque, J. J., Riggall, A. C., Gosseries, O., Starrett, M. J., Mayering, E. E., & Postle, B. R. (2016). Reactivation of latent working memories with transcranial magnetic stimulation. *Science*, *354*(6316), 1136–1139.

Sauseng, P., Klimesch, W., Heise, K. F., Gruber, W. R., Holz, E., Karim, A. A., ... Hummel, F. C. (2009). Brain Oscillatory Substrates of Visual Short-Term Memory Capacity. *Current Biology*, *19*(21), 1846–1852.

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., ... Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with human EEG alpha activity. *European Journal of Neuroscience*, *22*(11), 2917–2926.

Sheremata, S. L., Somers, D. C., & Shomstein, S. (2018). Visual short-term memory activity in parietal lobe reflects cognitive processes beyond attentional selection. *The Journal of Neuroscience*, *38*(6), 1716–17.

Steinhauer, S., & Hakerem, G. (1992). The Pupillary Response in Cognitive Psychophysiology and Schizophrenia. *Annals of the New York Academy of Sciences*, *658*(1), 182–204.

Stokes, M. G. (2015). "Activity-silent" working memory in prefrontal cortex: A dynamic coding framework. *Trends in Cognitive Sciences*, *19*(7), 394–405.

Tas, A. C., Luck, S. J., & Hollingworth, A. (2016). The relationship between visual attention and visual working memory encoding: A dissociation between covert and overt orienting. *Journal of Experimental Psychology: Human Perception and Performance*, *42*(8), 1121–1138.

Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). Alpha Band Electroencephalographic Activity over Occipital Cortex Indexes Visuospatial Attention

Bias and Predicts Visual Target Detection. *Journal of Neuroscience*, *26*(37), 9494–9502.

Todd, J. J., & Marois, R. (2004). Capacity limit of visual short-term memory in human posterior parietal cortex. *Nature*, *428*, 751–754.

Unsworth, N., Fukuda, K., Awh, E., & Vogel, E. K. (2014). Working memory and fluid intelligence: Capacity, attention control, and secondary memory retrieval. *Cognitive Psychology*, *71*, 1–26.

van Dijk, H., van der Werf, J., Mazaheri, A., Medendorp, W. P., & Jensen, O. (2010). Modulations in oscillatory activity with amplitude asymmetry can produce cognitively relevant event-related responses. *Proceedings of the National Academy of Sciences*, *107*(2), 900–905.

Vogel, E. K., & Machizawa, M. G. (2004). Neural activity predicts individual differences in visual working memory capacity. *Nature*, *428*, 748–751.

Vogel, E. K., Mccollough, A. W., & Machizawa, M. G. (2005). Neural measures reveal individual differences in controlling access to working memory. *Nature*, *438*(7067), 500–503.

Wolff, M. J., Jochim, J., Akyurek, E. G., & Stokes, M. G. (2017). Dynamic hidden states underlying working memory guided behaviour. *Nature Neurosci.*, 1–35.

Xu, Y., & Chun, M. M. (2006). Dissociable neural mechanisms supporting visual short-term memory for objects. *Nature*, *440*, 91–95. Retrieved from