# Chromosomal fusion and life history-associated genomic variation contribute to within-river local adaptation of Atlantic salmon.

Kyle Wellband[1,2], Claire Mérot[2], Tommi Linnansaari[1,3], J. A. K. Elliott[4], R. Allen Curry[1,3], Louis Bernatchez[2]

[1] Canadian Rivers Institute, Department of Biology, University of New Brunswick, Fredericton, NB, E3B 5A3

[2] Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec, QC, G1V 0A6

[3] Faculty of Forestry and Environmental Management, University of New Brunswick, Fredericton, NB, E3B 5A3

[4] Cooke Aquaculture Inc., Oak Bay, NB, E3L 3S7

Correspondence: Kyle Wellband, kwellban@unb.ca

Running title: Fusion and outlier SNPs structure salmon pops

Keywords: structural variation, six6, Atlantic salmon, chromosomal fusion, local adaptation

## Abstract

Genomic data provide the means to investigate the genetic basis of local adaptation and provide information about the forces structuring populations. Atlantic salmon are typically characterized by hierarchical population structure at multiple spatial scales; however, this is not the case for a large river in North America. To resolve genetic relationships between, and the forces structuring, Atlantic salmon populations in this system we applied a 50K SNP array to genotype 728 juvenile salmon from throughout the watershed. Consistent with previous work, we report extremely weak overall population structuring (Global $F_{ST}$ = 0.004) and failed to support for hierarchical structure between the river's two main branches. In contrast, outlier variation between populations was concentrated in a 250 Kb region of chromosome 9, including the gene *six6*, that has previously been linked to age-at-maturity and run-timing for this species. A polygenic framework for characterizing adaptive divergence identified 198 genes that were associated with an elevation and climatic gradient. We further identified a fusion between chromosomes 8 and 29 that was polymorphic among individuals. Fusion genomic characteristics included high LD, reduced heterozygosity in the fused homokaryotes, and strong divergence between the fused and the unfused rearrangement. The frequency of the fusion was associated with summer precipitation which supports the hypothesis that this rearrangement may contribute local adaptation despite weak neutral differentiation. Overall our results indicate adaptive processes, independent of major river branching, are more important than neutral process for structuring these populations.

**Introduction**

Conservation and management of species benefit from understanding of fine-scale demographic and evolutionary processes, which aid in delimiting appropriate population-scale units for management and conservation of genetic diversity (Fraser & Bernatchez, 2001; Funk, McKay, Hohenlohe, & Allendorf, 2012; Palsbøll, Bérubé, & Allendorf, 2007; R. S. Waples & Gaggiotti, 2006). Population genomic methods provide increased power for delineating genetic structure over traditional approaches and can identify putatively adaptive variation among populations that may reflect local adaptation (Funk et al., 2012). Genomic variation associated with environmental factors can indicate relevant biotic or abiotic forces possibly driving local adaptation (e.g. Benestan et al., 2016; Bourret, Dionne, Kent, Lien, & Bernatchez, 2013; Hecht, Matala, Hess, & Narum, 2015). Furthermore, the genomic basis of important life history variation relevant to maintaining within-population diversity can be identified (Barson et al., 2015; Cauwelier, Gilbey, Sampayo, Stradmeyer, & Middlemas, 2017). Knowledge of genomic variation linked to various locally-adaptive phenotypes will facilitate improved population monitoring and more effective conservation and management of species (Aykanat, Lindqvist, Pritchard, & Primmer, 2016).

Beyond applied questions, population genomics has provided important insights into the evolution of biological diversity, helping to disentangle the relative roles of demographic processes, genomic features, and selection in shaping adaptation (Rogers, Xu, & Schlüter, 2017; Tiffin & Ross-Ibarra, 2014). In particular, population genomics have provided insight into the basis of phenotypic parallelism within- and among-species (Bernatchez, 2016; Elmer & Meyer, 2011). The architecture of these traits has been shown to reflect both conserved and divergent sets of genes and generated questions regarding the conditions and constraints driving the evolution of parallelism vs. novelty at a genomic level (Yeaman, Gerstein, Hodgins, & Whitlock, 2018).

Population genomics has also helped reveal the role of chromosomal rearrangements in eco-evolutionary processes (Wellenreuther & Bernatchez, 2018). Large rearrangements (e.g. chromosomal inversions, fusions, fissions, or translocations) disturb homologous chromosomal pairing during meiosis that results in tight linkage among genes within rearranged regions detectable by genomic methods (Wellenreuther & Bernatchez, 2018). Chromosomal rearrangements function like "supergenes" and have been implicated in important adaptive phenotypes of several species (e.g., salinity in cod: Barth et al., 2017; climatic gradients in *Drosophila*: Kapun, Fabian, Goudet, & Flatt, 2016; crypsis in stick insects: Lindtke et al., 2017; migratory behaviour in cod: Sinclair-Waters et al., 2017). Furthermore, they have been shown to facilitate environmental adaptation in the face of high levels of gene flow (Barth et al., 2017).

Atlantic salmon, *Salmo salar*, is an economically and culturally important anadromous fish on both sides of the Atlantic Ocean. The species exhibits a wide variety of life history strategies that are shared throughout large portions of their range and believed to reflect local adaptations (Garcia de Leaniz et al., 2007). They exhibit hierarchical population structure in both Europe and North America (Bourret, Kent, et al., 2013; Dionne, Caron, Dodson, & Bernatchez, 2008; Perrier, Guyomard, Bagliniere, & Evanno, 2011; Sinclair-Waters et al., 2017; Verspoor et al., 2005). Regional structure is driven by a combination of geographic proximity as well as environmental gradients (e.g. temperature) suggesting that adaptive processes are important for structuring regional gene pools (Bourret, Dionne, et al., 2013; Dionne, Miller, Dodson, Caron, & Bernatchez, 2007; Jeffery et al., 2017). At finer scales, rivers are differentiated from one another and population structure can be present among tributaries within certain river systems (Aykanat et al., 2015; Cauwelier et al., 2017; Dionne, Caron, Dodson, & Bernatchez, 2009; Garant, Dodson, &

Bernatchez, 2000; Primmer et al., 2006; Pritchard et al., 2018; Vähä, Erkinaro, Niemelä, & Primmer, 2007). Within-river structuring also reflects local adaptation because even weak genetic structure within river systems corresponds to meaningful biological differences in growth and survival among populations (Aykanat et al., 2015; Mobley et al., 2018; Pritchard et al., 2018).

The Miramichi River, New Brunswick, Canada, presents a conundrum for the expected patterns of population structuring inherent to Atlantic salmon in a large and complex river system. It is a large watershed with a complex river network that has no human barriers for fish movement (Cunjak & Newbury, 2005). Despite harbouring a large population of Atlantic salmon with some evidence of demographic independence between the two main branches of the river at least (DFO, 2017), several studies have failed to characterize genetic population structure in this system (Dionne et al., 2009; Dodson & Colombani, 1997; Moore et al., 2014; but see Moller, 2005). These studies were based on limited numbers of neutral markers and modest sampling designs, leaving uncertainties about the source of negative results (panmixia) and without the possibility of inferring adaptive variation. Here we revisit these questions and employ both an extensive sampling design and high-resolution genotyping to characterize within-river population structure and identify putative adaptive differentiation for Atlantic salmon in the Miramichi River. This work contributes to our knowledge of the patterns and processes of local adaptation in large river systems.

## Methods

*Sampling and SNP array genotyping*

Juvenile Atlantic salmon (age 1+ parr) were collected from sites representing 15 major tributaries throughout the Miramichi River watershed from August 29 to October 13, 2016 (Figure 1, Supplemental Table 1). Samples were humanely euthanized and transported on ice to the Miramichi Salmon Association Conservation Rearing Facility at South Esk, New Brunswick. Fin clips were removed and preserved in 95% ethanol. The carcases were supplied to DFO Gulf region for additional analyses.

DNA was extracted from fin clips by the Center for Aquaculture Technologies Canada (Souris, PEI) using a standard DNA extraction procedure and samples were genotyped using a custom 50K SNP Axiom genotyping array designed for North American Atlantic Salmon at the Centre for Integrative Genetics (CIGENE, Norwegian University of Life Sciences, Norway). Raw data from the arrays was processed following the Affymetrix Best Practices Workflow using thresholds of >0.8 DQC and >0.97 QC call rate for high-quality samples. A number of samples (163/766) failed to meet these thresholds and thus we attempted to recover data from these sample by re-implementing the Affymetrix Best Practices Workflow with thresholds of >0.8 DQC and >0.90 QC call rate. Samples not exceeding these reduced thresholds (23 samples) were excluded from further analysis. We then compared the minor allele frequency (MAF) of each SNP in all retained individuals under the reduced quality thresholds with the frequency estimated for the high-quality samples and high-quality thresholds. Any SNPs demonstrating significant shifts in allele frequency between the two quality thresholds were imputed as missing genotypes for the recovered samples. SNPs were ordered following their genomic position in the Atlantic salmon reference genome (ICSASG v2; NCBI Genbank accession: GCA_000233375.4; Lien et al., 2016). Due to the commercial nature of the SNP array, SNP positions in the dataset have been shifted randomly from their exact location on the reference genome to protect intellectual property while maintaining genomic order and approximate genomic location. We removed both samples and

SNPs with greater than 10% missing data as well as SNPs with a minor allele frequency less than 0.01 from the data set using PLINK v1.9 (Chang et al., 2015).

For population genetic analyses that assume a set of neutral and unlinked markers, we created a reduced dataset. We first used BayeScan v2.1 (Foll & Gaggiotti, 2008) with relaxed settings (prior odds = 10) to characterize $F_{ST}$ outlier markers among the tributary samples. These markers were first removed from the dataset and we then removed SNPs with high physical linkage using a sliding-window approach in PLINK where SNPs with a variance inflation factor greater than two (VIF > 2) were removed from 50 SNP windows shifted by 5 SNPs after each iteration. This neutral and LD pruned dataset is hereafter referred to as the neutral dataset and, unless otherwise stated, was used as the primary dataset for population genetic analyses.

*Detection and characterization of structural variation*

We used principal components analysis (PCA) implemented in PLINK to visually explore patterns of allelic variation for the whole SNP dataset. The first two PCs of a PCA on the entire dataset revealed a striking pattern of three groups of samples that could not be explained by biological characteristics or geographic grouping of the samples (see results). We investigated this pattern further, first using pcadapt v3.0.4 (Luu, Bazin, & Blum, 2017) to characterize divergence between these three groups. Visual inspection of PC axes 3+ indicated that only axes 1 and 2 were involved in generating the three groups and thus we set the number of PCs retained for pcadapt to K = 2. This analysis revealed two strongly divergent genomic regions localized to the beginnings of chromosomes 8 and 29 ($F_{ST} > 0.6$). North American populations of Atlantic salmon are thought to be polymorphic for a fusion between chromosomes 8 and 29 (Brenna-Hansen et al., 2012). Thus, we further investigated to see if variation in chromosome structure could explain the observed pattern in our data. We first calculated pairwise linkage ($r^2$) among all SNPs located on chromosomes 8 and 29 using PLINK. Based on the pattern observed, we reversed the order of chromosome 8 and merged it with chromosome 29. We then used the package inveRsion v1.22 (Cáceres, Sindi, Raphael, Cáceres, & González, 2012) in R v3.3.3 (R Core Team 2017) to estimate the break points delineating a chromosomal rearrangement. Using the identified break points, we then assigned a rearrangement genotype (karyotype) to each individual using the R package invClust v1.0 (Cáceres & González, 2015). We then used karyotype as a grouping factor and investigated patterns of genetic diversity across chromosomes 8 and 29. Minor allele frequency and heterozygosity for each karyotype, as well as pairwise $F_{ST}$ among karyotypes were calculated using PLINK.

Based on individual karyotype assignments, we calculated the frequency of the fusion (called allele "A") and the relative proportion of the three karyotypes (AA: homokaryote fused, AB: heterokaryote, BB: homokaryote unfused) for each tributary. Heterogeneity in fusion frequencies between tributaries was tested using an analysis of deviance on a generalized linear model (GLM) with binomial logistic transformation. Within each tributary, HWE was tested using a chi-square test. We also tested for population structure among tributaries based on frequencies of fusion karyotypes as detailed below for neutral markers.

*Tributary genetic diversity and population structure*

Observed and expected heterozygosities as well as inbreeding coefficients were calculated for each of these populations using PLINK. The effective number of breeders (Nb) for the sampled cohort was estimated using the bias-corrected linkage disequilibrium method of Waples and Do (2010), as implemented in the software NeEstimator v2.1 (Do et al., 2014). Genomic datasets with

thousands of markers are known to introduce a downward bias for Nb estimates due to linkage disequilibrium caused by the inclusion of comparisons of physically linked markers (Waples, Larson, & Waples, 2016). Despite our creation of a neutral dataset with low physical linkage, we used genomic position information of the SNPs to only compare linkage among SNPs located on different chromosomes and thus absolutely not in physical linkage. Due to the evidence for a fusion between chromosomes 8 and 29, we further excluded these chromosomes from the dataset used to calculate Nb. We then corrected Nb to account for overlapping generations in the parents of the sampled cohort following Waples et al. (2014) and using parameters derived by Ferchaud et al. (2016) for Atlantic salmon.

We calculated Weir and Cockerham's (1984) unbiased estimate of $F_{ST}$ among all pairs of tributary populations and the statistical significance of $F_{ST}$ estimates was assessed using 10,000 permutations of the data in Arlequin v3.5.2.2 (Excoffier & Lischer, 2010). To explicitly test for hierarchical population structure, we used an AMOVA where tributary populations were grouped according to their location in the two main branches of the river (Northwest and Southwest). The site NWM was excluded from this analysis because it is the donor stream for residual stocking product (i.e. it receives hatchery-derived salmon from mixed and untraceable stocks). AMOVA calculations were conducted in Arlequin v3.5.2.2 and 10,000 permutations were used to assess the statistical significance of population structure.

To test for a pattern of isolation by distance among tributary populations, we used a Mantel test as well as the redundancy analysis (RDA) approach recommended by Meirmans (2015). We used the riverdist v0.15.0 (Tyers 2017) package in R v3.3.3 (R Core Team 2017) to calculate the shortest linear river distances between tributary sampling sites. We tested the association of pairwise linear river distance with linearized pairwise genetic distance ($F_{ST}/1-F_{ST}$) using a Mantel test with 1,000 permutations as implemented in the R package vegan v2.4-5 (Oksanen et al. 2017). We also decomposed the genetic variance explained by geographic distance using RDA as implemented in vegan v2.4-5. We first calculated distance-based Moran's eigenvector maps (MEMs) from the linear river distances among sites using the R package adespatial v0.1-0 (Dray et al. 2017) to represent the sampling sites in Cartesian coordinates that accounted for dispersal restrictions induced by the constraints of the dendritic structure of the river. The MEMs with a positive value of Moran's I were then used as predictor variables in an RDA where the tributary minor allele frequencies were used as the dependent variables. We used a forward step procedure to only select MEMs that explained significant (1,000 permutations, $p < 0.05$) variance in allele frequencies. Significance of the final model, constrained axes, and the marginal significance of selected MEMs were assessed using 1,000 permutations. The proportion of among-population genetic variation explained by river distance is thus the proportion of total RDA variance that is explained by constrained axes in the final model.

*Clustering analyses*

We used several naïve clustering routines to also test for population structure. We first used Admixture v1.3 (Alexander, Novembre, & Lange, 2009) to estimate the number of genetic clusters in the neutral dataset. Admixture was with 5-fold cross-validation for a range of putative populations (K = 1 to K = 10). The number of genetic groups (K) was chosen based on the lowest cross-validation error. We also used Discriminant Analysis of Principal Components (DAPC; Jombart, Devillard, & Balloux, 2010) to provide a non-model-based approach for estimating genetic clusters. The number of genetic clusters was determined using k-means clustering of principal components transformed allele frequency data, as implemented in the 'find.clusters'

function of the adegenet v2.1.0 package (Jombart & Ahmed, 2011) in R v3.3.3 (R Core Team 2017). The number of clusters was determined based on the profile of the Bayesian information criterion (BIC) across a range of putative populations ($K = 1$ to $K = 10$). Finally, we used principal components analyses, as implemented in PLINK, to visually investigate population groupings in multivariate allelic space.

*Population assignment*

We performed population assignment analyses using the frequency-based likelihood criterion of Paetkau et al. (1995). Likelihoods of assigning to a population were calculated based on the home likelihood ($L_{Home}$) using a leave-one-out procedure where missing alleles in a population were input at a frequency of 0.005 to prevent undefined likelihood values. To discriminate between miss-assignments and recent immigrants we used 1000 Monte Carlo permutations (Paetkau, Slade, Burden, & Estoup, 2004) to generate null distributions of $L_{Home}$ and excluded individuals from their sampled population if their $L_{Home}$ fell in the lower 5% of the null distribution for that population. Loci for conducting the assignment were chosen based on their $F_{ST}$ rank among all loci. We tested a range of SNP loci used for assignment (500, 1000, 2000, 3000, 4000, 5000) to assess the potential for increased power with increasing numbers of markers. To avoid high-grading bias (Anderson, 2010), we randomly split populations in half where one half represented a training set that was used to calculate global $F_{ST}$ among tributaries and rank loci and the other half represented a holdout set that was used to assess the accuracy of genotype assignment. We repeated the training-set-based marker selection, genotype assignment and holdout-set assignment accuracy assessment 10 times to evaluate the repeatability of our genotype assignment results.

*Outlier analyses*

Fractal landscapes, like the dendritic ecological networks of rivers, restrict dispersal of organisms relative to that predicted by traditional 'island' or 'stepping-stone' models of population connectivity. This restricted dispersal results in inflated variance in $F_{ST}$ at neutral markers compared to that expected under traditional population genetic models and leads to increased false positive outliers if these demographic relationships are not properly take into account (Fourcade, Chaput-Bardy, Secondi, Fleurant, & Lemaire, 2013). To reduce the detection of false positive outliers we implemented two outlier tests that take into account the genetic relatedness of populations: the modified FLK test of Bonhomme et al. (2010) and BayPass v2.1 (Gautier, 2015). These test control for demography by taking into account the genetic relatedness of populations and thus should exhibit reduced false-positives relative to tests assuming an island model (e.g. BayeScan). For FLK, we first estimated Reynold's distance among populations based on the neutral SNP dataset (21599 SNPs) and then calculated the $T_{F-LK}$ statistic for each SNP as implemented in R following Bonhomme et al. (2010). R code for the FLK analysis was obtained from: https://qgsp.jouy.inra.fr (date accessed: December 15, 2017). Statistical significance of outliers was assessed by comparing the test statistic to the Chi-squared distribution and correcting the p-values using FDR of 0.05. BayPass does not provide p-values and thus we used the ranking of loci based on the XtX statistic to corroborate results from other tests. Due the lack of hierarchical structure we also conducted outliers tests using Bayescan v2.1 (Foll & Gaggiotti, 2008) with program defaults but with the prior odds set to 10000 to reduce the incidence of false positives (Lotterhos & Whitlock, 2014) and OutFLANK was used also (Whitlock & Lotterhos, 2015). Loci putatively experiencing divergent selection were identified as those with a false discovery rate

(FDR) of less than 0.05. We identified a consensus set of outliers as the intersection of significant outliers across all outlier approaches. To identify genes that are potential targets of selection we used BEDTools v2.26.0 (Quinlan & Hall, 2010) to search the Atlantic salmon genome for genes that occurred within a 10 Kb window on either side of each outlier.

*Multivariate selection and environmental association*

To investigate the role of environmental variation in driving differentiation and the potential for multivariate selection, we examined associations of climate (i.e. temperature and precipitation) and physiography (i.e. river distance and elevation above sea level) with tributary population allele frequencies using Redundancy Analysis (RDA). We also tested for correlations of the climate and physiography variables with the frequency of the fusion and outlier SNPs allele frequencies using logistic regression.

We used GPS locations of the sampling sites to extract temperature and precipitation normals (1970 – 2000) from the WorldClim 2.0 Bioclimatic database (Fick & Hijmans, 2017), as well as elevation (NRCAN 2017) using QGIS v2.14 software (QGIS Development Team 2016). We calculated distance to the river mouth using the riverdist v0.15.0 package in R. Because tributaries are represented by more than one sampling site, we summarized the environmental data for each tributary by taking an average of the sampling site level environmental data weighted by the number of individuals at a site for each tributary. Many of the climate and physiography variables were highly correlated across tributaries ($|r| > 0.8$; Figure S1), so we first performed a variable reduction step by using two PCAs to summarize variation for the combined temperature/elevation/distance variables and the precipitation variables separately. For each PCA, we retained PC axes based on the broken-stick distribution (Frontier, 1976). Site scores of the retained PC axes represent summaries of environmental variation and were subsequently used as predictors of genomic variation.

Multivariate polygenic selection was tested using RDA with the three environmental summary variables (see results) as predictors and tributary allele frequencies for the whole SNP dataset as dependent variables. We performed forward step selection of the environmental variables and retained those that explained significant (1000 permutations, $p < 0.05$) variance in allele frequencies. Variance inflation factors were below 10 for all variables indicating no multicollinearity between the predictors. Significance of the final model was assessed using 1000 permutations. To identify SNPs influenced by environmental variation we scaled and centered the raw scores for SNPs on each constrained RDA axis. We then used a robust Mahalanobis distance metric similar to that applied in the outlier software pcadapt (Luu et al., 2017) to identify SNPs significantly loaded on each constrained axis. To control for population structure while testing for environmental associations, we also performed a partial RDA as just described but using the significant MEMs from the isolation by distance analysis as conditioning variables.

To test the individual association between the chromosome 8 and 29 fusion or outlier SNP frequencies and environment, we used a GLM with a logistic link function for binomial data, the response variable being the number of individuals carrying/not carrying the fusion arrangement or the alternative allele and the explanatory variable being an environmental variable. Given that outlier SNPs on the same chromosome were in strong linkage disequilibrium and that the distribution of frequencies was highly correlated (Figure S2), we performed the analysis on the mean MAF across outlier SNPs from the same chromosome.

*Functional enrichment*

To test for functional enrichment of genes associated with reduced recombination region of the fusion as well as the significant environmental variables from the RDA we conducted Gene Ontology enrichment tests. We first extracted gene ontology (GO) information for all Atlantic salmon genes from SalmoBase (Samy et al., 2017; www.salmobase.org, accessed: November 25, 2017). We then determined reference sets of genes for each analysis as all of the Atlantic salmon genes for the fusion and all genes within 10 Kb of a SNP on our array for the environmental outliers (the same window used to associate genes with significant SNPs). We tested for functional enrichment of GO biological process terms using the "weight01" algorithm to account for the structure of the GO term relationships as implemented in the topGO v2.26 package in R (Alexa, Rahnenführer, & Lengauer, 2006).

## Results

*Sampling and SNP genotyping*

A total of 766 age 1+ Atlantic salmon were collected throughout the Miramichi River watershed. The size and weight distributions of the collections were unimodal and indicate the collections represented a single cohort (mean ± sd; fork length = 85.1 ± 6.7 mm; weight = 7.4 ± 1.7 g; Figure S3). Following all quality filtering, we retained data for 728 individuals at 52466 SNPs. The overall genotyping rate was high with less than 2% missing data in the full dataset. Following position-based pruning for linkage disequilibrium, 21599 SNPs were retained in the neutral dataset that was used for most population genetic inferences.

*Detection and characterization of structural variation*

Exploratory analyses of the entire dataset using PCA revealed three prominent clusters that did not obviously reflect geographic (i.e. tributaries) or biological characteristics (i.e. sex, length, weight) of the samples (Figure 2A). Divergence among these three groups was driven entirely by two regions located on chromosomes 8 and 29 (Figure 2B). Mean divergence between these three groups within this region ($F_{ST} = 0.27$) was nearly three orders of magnitude greater than mean divergence between these groups across the rest of the genome ($F_{ST} = 0.0003$). Pairwise LD among SNPs on chromosomes 8 and 29 revealed two prominent linkage blocks, each spanning several Mb, within both chromosomes and strong linkage (mean pairwise $r^2$ within the region = 0.17 vs. mean pairwise $r^2$ outside the region = 0.007) between 7 Mb of chromosome 8 and 3 Mb of chromosome 29 (Figure 2D). This pattern reflects the fusion between chromosomes 8 and 29 that has been previously described in North American Atlantic salmon using linkage mapping and cytogenetic techniques (Brenna-Hansen et al., 2012). Our patterns of linkage suggest that the p-arm of chromosome 8, containing primarily rDNA repeated units (Lien et al., 2016; Pendás, Morán, & Garcia-Vázquez, 1993), fused with the centromere of the acrocentric chromosome 29 (Figure 2D).

This fusion has not been fixed in North American populations and thus individuals may carry two 8-29 fused chromosomes (AA), two unfused sets of chromosomes 8 and 29 (BB) or one fused 8-29 and one unfused set of chromosomes 8 and 29 (AB). Genetic diversity within each of the three karyotype groups revealed that fused homokaryotes have dramatically reduced allelic diversity and heterozygosity in the highly linked region while unfused homokaryotes do not exhibit this pattern (Mean heterozygosity within vs. outside linked region: AA = 0.08 vs. 0.27, t = 22.3, p << 0.001; BB = 0.31 vs. 0.29, t = 1.8, p = 0.11; Figure 2C). These results are consistent with observations that recombination is reduced near the fusion point in fused chromosomes but not in

unfused chromosomes (Bidau, Giménez, Palmer, & Searle, 2001; Dumas & Britton-Davidian, 2002).

The fusion karyotypes were in Hardy-Weinberg equilibrium in all tributary populations and there was no sex bias in the distribution of fusion ($\chi^2 = 0.35$, df = 2, p = 0.84). The fusion was polymorphic in all tributary samples and its frequency was significantly heterogeneous between tributaries (deviance = 39, df = 14, p < 0.001), ranging from 38% to 66%. However, there was no overall difference in fusion frequency between the two main branches of the river (NW vs. SW, $\chi^2 = 1.95$, df = 2, p = 0.38). Further population structure and environmental associations with the fusion karyotype are presented below with the results for SNPs.

*Tributary genetic diversity and population structure*

Genetic diversity was consistent amongst the tributary samples and levels of inbreeding were low (Table 1). The effective number of breeders ranged over two orders of magnitude from approximately 80 in Taxis River to over 4000 in Clearwater Brook (Table 1). Genetic divergence among tributaries was low, but statistically significant (Global $F_{ST} = 0.004$, pairwise $F_{ST}$ range = 0.001 – 0.014; Figure 3). Three tributary populations from upstream reaches exhibited consistent stronger differentiation from all other tributaries, i.e., in the Southwest Miramichi: Rocky Brook (mean $F_{ST} = 0.01$), and Taxis River (mean $F_{ST} = 0.007$) and in the Northwest Miramichi: Upper Northwest Miramichi River (mean $F_{ST} = 0.006$; Figure 3). Additionally, Northwest Millstream which empties into the tidal reaches of the Northwest Miramichi and regularly receives hatchery residuals of untraceable origin exhibited similar levels of differentiation ($F_{ST} = 0.007$). Omitting these populations reduced the magnitude of overall structure by half ($F_{ST} = 0.002$). Hierarchical structure was not supported by the AMOVA as no genetic variance was explained by the two main branches of the Miramichi River ($F_{ST} = 0.0001$, p = 0.35; Table 2); however, the AMOVA did support population structure among tributaries within the main branches of the river consistent with the pairwise $F_{ST}$ results ($F_{ST} = 0.004$, p < 0.001; Table 2). The Mantel test for isolation by river distance was non-significant (p = 0.17, Mantel's r = 0.09); however, the spatial RDA identified the first MEM as explaining significant among-population variation in allele frequencies (p = 0.03; Figure 4). The spatial RDA resulted in a single constrained axis that explained 1.1% of the among-population variation in allele frequencies. This weak pattern of spatial structure was driven by differentiation of the tributaries in the upstream reaches of the Southwest from the downstream Southwest tributaries and Northwest tributaries (Figure 4).

The population structure among tributaries based on the frequency of the chromosomal fusion was five times stronger than that exhibited by SNPs (global $F_{ST} = 0.019$ vs. 0.004) and was in the upper 5% of the distribution of global $F_{ST}$ estimates for all SNPs (Figure S4). The estimates of pairwise tributary divergence were not correlated between the fusion and SNP markers (Mantel's r = 0.19; p = 0.16). AMOVA based on the fusion karyotype variation also failed to support the hierarchical structuring between the main branches (NW vs. SW; $F_{ST} = -0.0004$, p = 0.66; Table 2) of the Miramichi River but did identify structuring among tributaries within the main branches ($F_{ST} = 0.019$, p < 0.001; Table 2). The fusion karyotype also did not exhibit a pattern of isolation by distance (Mantel's r = -0.09, p = 0.83).

*Clustering analyses*

Principal components analysis of the putatively neutral dataset revealed groupings of samples in multivariate allelic space that reflected the geographic location of sampling (Figure 4). The first three PC axes explained 7.3%, 6.7%, and 6.2% of the total variation, respectively, and

were sufficient to visually separate the Taxis River, Rocky Brook, Upper Northwest Miramichi River, and, to a lesser extent, Northwest Millstream from each other and from the rest of the samples (Figure 3). The PCA results were consistent with the population structure analyses, where most individuals were weakly differentiated but these four tributary populations exhibited elevated divergence relative to the other tributaries. Inspection of additional axes as well as the removal of these populations and subsequent PCA on reduced datasets failed to reveal additional groupings of individuals. Neither Admixture nor DAPC supported genetic clustering of samples likely due to the weak overall structure we observed in this system (Figures S5 – S7).

*Population assignment*

The assignment of individuals in the holdout datasets was highly variable across tributary populations with only five tributary populations having successful assignment of greater than 80% (Figure S8). Of these, three were previously identified as the weakly differentiated tributaries. The low assignment success for most tributaries reflects the lack of differentiation in the system. Accounting for individuals that could be excluded as potential migrants (<5% probability of assignment based on 1000 permutations) had a positive effect on the assignment accuracy but did not change the overall pattern of results. Mis-assigned individuals were generally mis-assigned as coming from the Little Southwest Miramichi River (LSW; Figure S9). This tributary had the largest sample size among the group of weakly differentiated tributaries and likely reflects a more accurate allele frequency estimation.

*Outlier analyses*

We detected 21, 23, and 15 outliers using Bayescan, OutFlank, and FLK, respectively (Figure S10). In total, 13 of these SNPs were identified as statistically significant by all methods and all were among the top 17 most divergent SNPs according to the XtX statistic of BayPass. Two of the common outliers were found on chromosome 3, one on chromosome 6, and the remaining 10 outliers were clustered in a region of chromosome 9 that spanned approximately 250 kb. We searched the *Salmo salar* genome annotation for genes that fell within 10 kb on either side of each outlier SNP. Twelve different genes were in proximity to outlier SNPs and could represent putative targets of selection (Table 3). Of particular note, the region on chromosome 9 includes the gene *six6*, which has previously been linked to age- and possibly size-at-maturity in Atlantic salmon in Europe (Barson et al., 2015; Johnston et al., 2014) as well as run-timing (Cauwelier et al., 2017; Pritchard et al., 2018).

*Multivariate selection and environmental association*

The PCA-based reduction of environmental variables resulted in three summary variables (Figure S11) that generally described: 1) temp./elevation, representing the first PC of geographic and temperature data, which explained 73% of variation and was positively associated with temperature (throughout the year) and negatively associated with elevation/distance to river mouth, 2) winter precipitation, PC1 of precipitation data, which explained 52% of variation and was positively associated with precipitation during the coldest/driest months, and 3) summer precipitation, PC2 of precipitation data, which explained 40% of variation and was positively associated with annual precipitation and precipitation during the warmest/wettest months.

When assessing their associations with genomic data, the forward step procedure of the RDA selected only the temp./elevation variable as explaining significant allelic variation and generated one significant discriminant axis (p = 0.001) explaining 3.1% of the total allelic

variation. There were 226 SNPs that were significantly associated with variation in temp./elevation. When controlling for underlying population structure using the significant MEM from the spatial RDA, the signal of polygenic selection associated with temp./elevation remained significant ($p = 0.014$), explaining 2.6% of the allelic variance. After controlling for population structure, variation in temp./elevation correlated with 198 SNPs, including eight outliers (out of 13) previously detected using single locus tests (Table 3; Figure 5).

Consistent with the RDA results, allele frequency variation for outlier SNPs on chromosomes 3 and 6 were significantly associated with variation in temp./elevation (chromosome 3: GLM $z = -5.9$, $p < 0.001$, chromosome 6: $z = -5.7$, $p < 0.001$, Figure 6A, S13A), which explained 36% (chromosome 3) and 30% (chromosome 6) of allele frequency variation between tributaries in these chromosomal regions. A marginal but significant effect of winter precipitation (GLM $z = -2.6$, $p = 0.009$) explains an additional 3-5% of variance for chromosome 6 (Figure 6). On chromosome 9, variation in allele frequency was also significantly associated with temp./elevation (GLM, $z = -3.7$, $p < 0.001$) and winter precipitation (GLM, $z = 4.3$, $p < 0.001$) and those combined predictors explained 11% of the variation between tributaries on this chromosome.

Variation in fusion frequency between tributaries was best predicted by variation in summer precipitation, which explained 21% of variance (Figure 6), and by variation in temp./elevation, which explained an additional 5% of variance. The fusion was significantly more frequent in tributaries with higher summer precipitation (GLM, $z = 3.0$, $p = 0.002$; Figure 6B) and tributaries at higher elevation characterized by lower temperatures (GLM, $z = 2.0$, $p = 0.02$; Figure S12B). Variation in fusion frequencies primarily corresponded to a replacement of one homokaryote by the other homokaryote (Figure S12B). These gradients occurred in parallel for both main branches of the river (Southwest: GLM, $z = 3.6$, $p = 0.002$; Northwest: GLM, $z = 1.7$, $p = 0.08$; Figure 6B). None of the environmental associations with genetic markers (outlier SNPs or chromosome 8-29 fusion) reflected differences between the NW and SW branches of the river.

*Functional enrichment*

We identified 213 genes located within the linked region associated with the chromosomal fusion. These genes represented functional enrichment for 26 GO biological processes ($p < 0.01$, Table 4). Of particular interest are several processes related to neural development and behaviour, phototransduction, and cell adhesion and cell-cell communication processes involved in contractile tissues. For the 198 genes associated with temp./elevation, 13 GO biological processes were over-represented ($p < 0.01$, Table 4). Processes related to development, memory, and sexual maturation dominated the enriched functions.

**Discussion**

Overall, population genomic analysis of 728 juvenile Atlantic salmon collected throughout the Miramichi River system generated three primary conclusions. First, population structure of salmon in the Miramichi River is surprisingly weak and not hierarchical as would be expected for organisms inhabiting dendritic systems like rivers. Second, we detected outlier variation among tributaries that was correlated with a gradient of elevation and temperature. Significantly, the detected outliers were clustered in a region on chromosome 9 that has been previously linked with both age-at-maturity and run-timing in European populations of Atlantic salmon. These traits vary among tributaries in the Miramichi River and suggest that locally adaptive processes are more important than neutral processes for structuring populations. Third, we identified population

structure for an important chromosomal rearrangement. This variation correlated with summer precipitation suggesting that variation in the presence of the fusion has functional importance for local adaptation of tributary populations. Furthermore, our dense SNP dataset provided one of the first insight into the genomic patterns related to a fusion rearrangement.

*Neutral population structure (or lack thereof?)*

Our results revealed remarkably weak neutral population structure for the largest Atlantic salmon river in North America, the Miramichi River (Cunjak & Newbury, 2005). There was no evidence of hierarchical structuring of populations between the two main branches of the river and spatial structure only explained approximately 1% of the total allelic variation. Our results are consistent with two previous population genetic studies in the Miramichi River system based on a few microsatellites that also found weak structure and no support for hierarchical structuring of populations (Dionne et al., 2009; Moore et al., 2014). Large rivers containing populations of Atlantic salmon on both sides of the Atlantic Ocean frequently exhibit genetically distinct populations with levels of differentiation up to an order of magnitude greater than we report (e.g. Aykanat et al., 2015; Dionne et al., 2009; Primmer et al., 2006; Vähä et al., 2007). Strong philopatry and natal homing behaviours are defining traits of salmonids and are expected to result in hierarchical population structuring (Fraser, Weir, Bernatchez, Hansen, & Taylor, 2011). Thus, the pattern of population structuring we have reported is at odds with expectations based on the species' biology.

Population divergence ($F_{ST}$) is a function of the product of the effective population size and the number of migrants per generation (Allendorf & Phelps, 1981). Within the Miramichi River, two of the more divergent tributaries we identified had the lowest Nb estimates suggesting that higher genetic drift may explain the elevated neutral divergence in these cases. Yet, for the other tributaries, our adjusted Nb estimates were not larger than those reported in other Atlantic salmon rivers that exhibit more pronounced genetic divergence (Ferchaud et al., 2016; Palstra, O'Connell, & Ruzzante, 2007, 2009), suggesting that particularly large effective population size is an unlikely explanation for the overall low levels of divergence we observed.

Alternatively, high rates of straying by spawning adults could explain the overall low divergence in the system. Straying for Atlantic salmon populations is typically at a rate of 6 – 10 % (Hendry, Castric, Kinnison, & Quinn, 2004; Keefer & Caudill, 2014; Pess, 2009); however, straying rates in the Miramichi River may be as high as 21% (Chaput, Moore, Hayward, Sheasgreen, & Dubee, 2001). Additionally, the Miramichi River is stocked annually and has been for more than 100 years (Labadie, 2016). Salmonids reared in hatcheries can have straying rates higher than 20% (Keefer & Caudill, 2014). What would cause higher rates of straying in the Miramichi River compared to many other salmon rivers is unclear, but past stocking activities may play a role.

Stocking to enhance Atlantic salmon populations in the Miramichi River dates back to the 1870s. Current enhancement activities collect adults from specific tributaries and return offspring back to those tributaries (Labadie, 2016). These stocking efforts are believed to account for only a small fraction of contemporary adult returns (Chaput et al., 2001; Wallace & Curry, 2017). At times, as many as 4000 adult salmon broodstock are known to have been collected in the estuary, bred in the hatchery, and their offspring released throughout the watershed (M. Hambrook, Miramichi Salmon Association, pers. comm.). Depending on the frequency and duration of this stocking, these activities could give the appearance of highly elevated gene flow and may have erased historical signals of population structure. Stocking can disrupt the genetic structure of

natural salmonid populations by reducing genetic divergence and increasing admixture among previously structured groups (Hansen, Fraser, Meier, & Mensberg, 2009; Pearse, Martinez, & Garza, 2011; Perrier, Baglinière, & Evanno, 2013; Perrier et al., 2011; Valiquette, Perrier, Thibault, & Bernatchez, 2014). Additionally, Atlantic salmon suffered a major anthropogenic insult in the 1950-60s, namely widespread pollution from DDT that resulted in mass mortalities (Elson, 1967; MacDonald, 1971), for which long-lasting effects on population structure are unclear. A combination of tagging studies, demographic modelling, and genotyping of archival samples to assess temporal trends in population structure would all be useful approaches to help resolve the underlying causes of weak population structuring in the Miramichi River.

For the three tributaries that show elevated divergence in this system, two of them exhibited small effective population sizes (Taxis River and Rocky Brook). In some years, Taxis River may be inaccessible to migrating adults due to low water levels (M. Hambrook, Miramichi Salmon Association, pers. comm.) suggesting that access to this system may drive the elevated genetic drift we observed. For the other two systems (Rocky Brook and Upper Northwest), there is no obvious reason why their connectivity to the rest of the system may be reduced. Both of these systems exhibited high alternate allele frequencies for the outlier region on chromosome 9 suggesting that selection may play a role in limiting the connectivity between these systems and the rest of the tributaries.

*Adaptive variation*

In contrast to the weak neutral population structure of Atlantic salmon in the Miramichi River, we identified genetic distinctiveness along an environmental gradient of elevation / temperature. Most of the outliers we detected were localized to a 250 Kb region of chromosome 9 that has previously been implicated in important life history variation for this species. In European Atlantic salmon, a particular a gene in this region, *six6*, has been associated with age-at-maturity in Atlantic salmon in Europe (Barson et al., 2015; Johnston et al., 2014), differences in run-timing within rivers, and local adaptation between tributaries (Cauwelier et al., 2017; Pritchard et al., 2018). Our study is the first to report direct evidence of within-river variation at this locus in a North American population of Atlantic salmon. Previously, Kusche et al. (2017) failed to find an association of *six6*-linked SNPs with age-at-maturity in four Atlantic salmon rivers from Québec. Their failure to repeat this association may have been a result of *six6* influencing a phenotype (e.g. size-at-maturity or run-timing) that is only correlated with age-at-maturity in certain rivers, as has been suggested by Barson et al. (2015).

In our study, variation in *six6* allele frequencies varied along an elevation gradient suggesting that run-timing, or a related trait, in the Miramichi River may too be influenced by variation at this locus. Run-timing in the Miramichi River has historically been bimodal with peaks that represent early (June – July) and late (Sept. – Nov.) returning fish respectively (Chaput, Douglas, & Hayward, 2016; Saunders, 1967). Adults spawning in the upper reaches of rivers return earlier and ascend higher than late returning salmon (Chaput et al., 2016). In particular, the fish from Rocky Brook are known to return early and they also exhibited the highest alternate allele frequencies for *six6* associated SNPs of any tributary. Fish in this river are morphologically distinct and exhibit earlier migration timing from those occupying lower reaches of the Miramichi River (Riddell & Leggett, 1981). They exhibit more fusiform bodies and longer paired fins consistent with life in higher velocity streams (Riddell & Leggett, 1981). These differences are heritable and persist when grown in common environments with salmon from lower elevation sites in the Miramichi River (Riddell, Leggett, & Saunders, 1981). There is evidence of similar trait variation

between salmon from the upper and lower sections of the NW Miramichi River (Elson, 1973). In Europe, Pritchard et al. (Pritchard et al., 2018) found an association of this locus with a synthetic variable explaining "flow volume" for Atlantic salmon in the Teno/Tana River in Norway. Our results for a North American system thus suggest that flow velocity may be a convergent mechanism across continents that is driving adaptive divergence among upstream and downstream populations.

In addition to single locus tests, we employed a polygenic framework to investigate correlated allelic changes among tributaries to environmental predictors. The same temperature / elevation variable was associated with correlated changes in allele frequency for 198 SNPs including 5 of those previously identified on chromosome 9. Gene function enrichment revealed roles in morphological development, memory, and sexual maturation for these genes that are consistent with the previously discussed differences in life history between tributaries. In total, nine of our variants detected by the RDA were located in proximity (<10 Kb) to variants associated with flow volume by Pritchard et al. (Pritchard et al., 2018). This represented 14% of the flow-volume-associated SNPs detected by Pritchard et al. (Pritchard et al., 2018) suggesting there is a level of genomic parallelism across continents. Of note, we too identified the neuronal growth regulator 1 gene (negr-1) in our temperature / elevation associated genes that has previously been suggested as a candidate locus for further investigation of its role in salmonid life history variation (Pritchard et al., 2018).

*Chromosomal rearrangement polymorphism: a possible role in local adaptation?*

Beyond single-locus variation, we identified a major polymorphic chromosomal rearrangement (fusion between chromosome 8 and 29). Interestingly, among-population variation in the frequency of the fusion did not mirror patterns of neutral variation, and rather co-varied with environmental variation in summer precipitation and elevation, suggesting a possible role in local adaptation. Chromosomal rearrangements, in particular inversions, frequently display such polymorphism in natural populations, with variations of frequencies along environmental gradients, which is generally interpreted as correlative evidence for a role of the inversion in local adaptation (e.g. Kapun et al., 2016). In fact, inversions underlie important phenotypic and life history variation in a wide variety of organisms (Wellenreuther & Bernatchez, 2018). In fish, inversions have been linked with a suite of traits differentiating anadromous and resident ecotypes of rainbow trout (Pearse, Miller, Abadia-Cardoso, & Garza, 2014), migratory phenotypes in Atlantic cod on both sides of the Atlantic Ocean (Berg et al., 2016; Sinclair-Waters et al., 2017), and adaptation to salinity gradients in Atlantic cod (Barth et al., 2017; Berg et al., 2015). Remarkably, the chromosomal rearrangement in cod allows fjord cod to maintain local adaptation to low salinity conditions despite high levels of immigration from offshore populations (Barth et al., 2017). By limiting recombination between the standard and inverted rearrangements and by trapping together a set of alleles, inversions indeed provide an important mechanism that facilitates local adaptation in the face of high levels of gene flow. Such mechanisms are also expected for fusions (Guerrero & Kirkpatrick, 2014) and may be particularly relevant for Atlantic salmon populations in the Miramichi River, which experience contrasting environments but weak neutral differentiation.

In contrast with the accumulation of studies on inversions, the role of fusions in adaptation is poorly documented. Karyotypic variation for fusions are widely reported within and between species (Dobigny, Britton-Davidian, & Robinson, 2017) and for some taxa, such as the house mouse, can represent extreme levels of within-species karyotypic variation (Piálek, Hauffe, &

Searle, 2005). Yet, very few studies report the distribution of fusion polymorphism in natural populations with possible links to fitness, phenotype, or the environment (Dobigny et al., 2017). Theoretical work suggests that chromosomal fusions may evolve in response to local selective forces (Guerrero & Kirkpatrick, 2014). To our knowledge, the best empirical support for this prediction is the grasshopper *Dichroplus pratensis*, for which fusion has been linked to phenotype and where fusion frequency varies with abiotic conditions, suggesting a link with adaptation to environmental extremes (Bidau, Miño, Castillo, & Martí, 2012). In such a context, the patterns of polymorphism and the environmental association we have observed for the chromosome 8 and 29 fusion in Miramichi River populations of Atlantic salmon represent a new opportunity to better understand whether, like inversions, fusions play a key role in adaptation in the face of gene flow.

Further investigation would be needed to properly test the link between the 8-29 fusion and phenotype/fitness in Atlantic salmon. Given the correlation with summer precipitation and elevation, we propose that the fusion may be associated with the ability to handle high flow velocity during the critical first few months of post-hatching juvenile salmon. Several studies in salmonids have also identified precipitation as an important variable underlying genomic variation among populations (Bourret, Dionne, et al., 2013; Hecht et al., 2015; Matala, Ackerman, Campbell, & Narum, 2014; Olsen et al., 2011). Alternatively, this variation in precipitation could be related to triggering and the timing of spawning migrations (Leggett, 1977). Indeed, enriched gene functions detected in the fused region were related to neural development, behaviour, and phototransduction that may be involved the timing of this life history variation.

The dense SNP dataset employed in our study also provides one of the first analyses of the genomic features associated with a chromosomal fusion and provides a complementary picture to previous knowledge based on cytology. For instance, patterns of allelic variation indicated that one group of homokaryotes, namely the fused one, exhibit much reduced diversity compared with the other homokaryotes. This result is consistent with meiotic observations that reported a reduced frequency of recombination events in fused chromosomes and a shift in the distribution of recombination toward the distal ends of chromosomes in fused homokaryotes and to a lesser extent in heterokaryotes (Bidau et al., 2001; Dumas & Britton-Davidian, 2002).

The patterns of linkage we report between these chromosomes also allow delimiting the regions affected by the reduction of recombination associated with the rearrangement. Previous work by Brenna-Hansen et al. (2012), which described this fusion, reported difficulties in determining the architecture of the fused chromosome from genetic mapping data. On the basis of fluorescence in-situ hybridization analysis, they had suggested that the fusion represented a tandem joining of the q-arm of chromosome 8 to the centromere of the acrocentric chromosome 29. Our linkage data suggests instead that it is the p-arm of chromosome 8 that has fused with the centromere of chromosome 29. The p-arm of chromosome 8 contains a dense cluster of rDNA repeats (Brenna-Hansen et al., 2012; Lien et al., 2016) and repeated elements of rDNA have been associated with chromosomal fusions and fissions in a wide variety of species (e.g. Barros et al., 2017; Cazaux, Catalan, Veyrunes, Douzery, & Britton-Davidian, 2011; Huang, Ma, Yang, Fei, & Li, 2008; Stahl et al., 1983) suggesting they play an important role in chromosomal rearrangements and thus may explain the occurrence of the fusion we have observed. Ultimately long-read sequencing data will be necessary to verify the architecture of this chromosomal rearrangement.

*Management implications*

Our data have soundly rejected the presence of hierarchical structuring of Atlantic salmon populations in the Miramichi River that had been previously hypothesized. Instead our results

support the relative isolation of upper watersheds from those in the lower river and a primary role for adaptive processes in structuring populations in this system. These results have important relevance for the management of Atlantic salmon. Run-timing rather than geography appears to be more important for structuring populations. Given the recent trend of run-timing homogenization (Douglas, Chaput, Hayward, & Sheasgreen, 2015), more attention must be paid to the underlying mechanisms generating this pattern and their role in population declines in this system. Furthermore, ongoing and proposed supplementation activities to enhance populations must continue to respect the potential local adaptation of tributary populations.

*Conclusions*

We have generated the largest and highest resolution dataset for Atlantic salmon in the Miramichi River to date. We reported unexpectedly weak neutral population structure and a complete absence of support for hierarchical structuring in this large river system. Instead we found important outlier variation that correlated with the environment and is consistent with both known life-history variation in this and in multiple other systems throughout the species' range. Our results contribute to a growing body of literature implicating parallel selection at the *six6* gene for explaining within-river local adaptation of Atlantic salmon on both sides of the Atlantic Ocean. We further characterized variation for a chromosomal fusion that revealed associations with environmental parameters orthogonal to that of other genomic variation, suggesting that this fusion rearrangement may contribute to local adaptation in the face of gene flow. Our results represent significant progress toward the resolution of a long-standing debate over the presence of genetic structure for Atlantic salmon in the Miramichi River. They reveal that adaptive processes appear to be more important for structuring populations than neutral processes. Whether this reflects intrinsic properties of the populations (i.e. large effective population size) or a legacy of anthropogenic influence is an open and important question that is in urgent need of resolution for the improved conservation and management of this species.

**Acknowledgements**

We thank the contributions of sampling teams from the Collaboration for Atlantic Salmon Tomorrow and Fisheries and Oceans Canada who collected the samples used in this study. We also acknowledge staff at CIGENE in Norway for conducting the SNP genotyping.

**Literature Cited**

Alexa, A., Rahnenführer, J., & Lengauer, T. (2006). Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. *Bioinformatics*, *22*(13), 1600–1607. doi:10.1093/bioinformatics/btl140

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655–1664. doi:10.1101/gr.094052.109

Allendorf, F. W., & Phelps, S. R. (1981). Use of Allelic Frequencies to Describe Population Structure. *Canadian Journal of Fisheries and Aquatic Sciences*, *38*(12), 1507–1514. doi:10.1139/f81-203

Anderson, E. C. (2010). Assessing the power of informative subsets of loci for population assignment: Standard methods are upwardly biased. *Molecular Ecology Resources*, *10*(4), 701–710. doi:10.1111/j.1755-0998.2010.02846.x

Aykanat, T., Johnston, S. E., Orell, P., Niemelä, E., Erkinaro, J., & Primmer, C. R. (2015). Low but significant genetic differentiation underlies biologically meaningful phenotypic divergence in a large Atlantic salmon population. *Molecular Ecology*, *24*(20), 5158–5174. doi:10.1111/mec.13383

Aykanat, T., Lindqvist, M., Pritchard, V. L., & Primmer, C. R. (2016). From population genomics to conservation and management: a workflow for targeted analysis of markers identified using genome-wide approaches in Atlantic salmon Salmo salar. *Journal of Fish Biology*, *89*(6), 2658–2679. doi:10.1111/jfb.13149

Barros, A. V., Wolski, M. A. V., Nogaroto, V., Almeida, M. C., Moreira-Filho, O., & Vicari, M. R. (2017). Fragile sites, dysfunctional telomere and chromosome fusions: What is 5S rDNA role? *Gene*, *608*, 20–27. doi:10.1016/j.gene.2017.01.013

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., … Primmer, C. R. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, *528*(7582), 405–408. doi:10.1038/nature16062

Barth, J. M. I., Berg, P. R., Jonsson, P. R., Bonanomi, S., Corell, H., Hemmer-Hansen, J., … André, C. (2017). Genome architecture enables local adaptation of Atlantic cod despite high connectivity. *Molecular Ecology*, *26*(17), 4452–4466. doi:10.1111/mec.14207

Benestan, L., Quinn, B. K., Maaroufi, H., Laporte, M., Clark, F. K., Greenwood, S. J., … Bernatchez, L. (2016). Seascape genomics provides evidence for thermal adaptation and current-mediated population structure in American lobster ( Homarus americanus ). *Molecular Ecology*, *25*(20), 5073–5092. doi:10.1111/mec.13811

Berg, P. R., Jentoft, S., Star, B., Ring, K. H., Knutsen, H., Lien, S., … André, C. (2015). Adaptation to Low Salinity Promotes Genomic Divergence in Atlantic Cod ( Gadus morhua L.). *Genome Biology and Evolution*, *7*(6), 1644–1663. doi:10.1093/gbe/evv093

Berg, P. R., Star, B., Pampoulie, C., Sodeland, M., Barth, J. M. I., Knutsen, H., … Jentoft, S. (2016). Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. *Scientific Reports*, *6*(December 2015), 1–12. doi:10.1038/srep23246

Bernatchez, L. (2016). On the maintenance of genetic variation and adaptation to environmental change: considerations from population genomics in fishes. *Journal of Fish Biology*, *89*(6), 2519–2556. doi:10.1111/jfb.13145

Bidau, C. J., Giménez, M. D., Palmer, C. L., & Searle, J. B. (2001). The effects of Robertsonian fusions on chiasma frequency and distribution in the house mouse (Mus musculus domesticus) from a hybrid zone in northern Scotland. *Heredity*, *87*(3), 305–313. doi:10.1046/j.1365-2540.2001.00877.x

Bidau, C. J., Miño, C. I., Castillo, E. R., & Martí, D. A. (2012). Effects of Abiotic Factors on the Geographic Distribution of Body Size Variation and Chromosomal Polymorphisms in Two Neotropical Grasshopper Species ( Dichroplus : Melanoplinae: Acrididae). *Psyche: A Journal of Entomology*, *2012*, 1–11. doi:10.1155/2012/863947

Bonhomme, M., Chevalet, C., Servin, B., Boitard, S., Abdallah, J., Blott, S., & SanCristobal, M. (2010). Detecting selection in population trees: The Lewontin and Krakauer test extended. *Genetics*, *186*(1), 241–262. doi:10.1534/genetics.110.117275

Bourret, V., Dionne, M., Kent, M. P., Lien, S., & Bernatchez, L. (2013). Landscape genomics in Atlantic salmon ( salmo salar ): Searching for gene-environment interactions driving local adaptation. *Evolution*, *67*(12), 3469–3487. doi:10.1111/evo.12139

Bourret, V., Kent, M. P., Primmer, C. R., Vasemägi, A., Karlsson, S., Hindar, K., … Lien, S.

(2013). SNP-array reveals genome-wide patterns of geographical and potential adaptive divergence across the natural range of Atlantic salmon (Salmo salar). *Molecular Ecology*, *22*(3), 532–551. doi:10.1111/mec.12003

Brenna-Hansen, S., Li, J., Kent, M. P., Boulding, E. G., Dominik, S., Davidson, W. S., & Lien, S. (2012). Chromosomal differences between European and North American Atlantic salmon discovered by linkage mapping and supported by fluorescence in situ hybridization analysis. *BMC Genomics*, *13*(1). doi:10.1186/1471-2164-13-432

Cáceres, A., & González, J. R. (2015). Following the footprints of polymorphic inversions on SNP data: from detection to association tests. *Nucleic Acids Research*, *43*(8), e53–e53. doi:10.1093/nar/gkv073

Cáceres, A., Sindi, S. S., Raphael, B. J., Cáceres, M., & González, J. R. (2012). Identification of polymorphic inversions from genotypes. *BMC Bioinformatics*, *13*, 28. doi:10.1186/1471-2105-13-28

Cauwelier, E., Gilbey, J., Sampayo, J., Stradmeyer, L., & Middlemas, S. J. (2017). Identification of a single genomic region associated with seasonal river return timing in adult Scottish Atlantic salmon (Salmo salar L.) identified using a genome-wide association study. *Canadian Journal of Fisheries and Aquatic Sciences*, cjfas-2017-0293. doi:10.1139/cjfas-2017-0293

Cazaux, B., Catalan, J., Veyrunes, F., Douzery, E. J., & Britton-Davidian, J. (2011). Are ribosomal DNA clusters rearrangement hotspots? A case study in the genus Mus (Rodentia, Muridae). *BMC Evolutionary Biology*, *11*(1), 124. doi:10.1186/1471-2148-11-124

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), 1–16. doi:10.1186/s13742-015-0047-8

Chaput, G., Douglas, S. G., & Hayward, J. (2016). Biological Characteristics and Population Dynamics of Atlantic Salmon ( Salmo salar ) from the Miramichi River , New Brunswick , Canada. *Canadian Science Advisory Secretariat Research Document 2016/029*.

Chaput, G., Moore, D., Hayward, J., Sheasgreen, J., & Dubee, B. (2001). Stock status of Atlantic Salmon (Salmo salar) in the Miramichi River, 2000. *Canadian Science Advisory Secretariat Research Document 2001/008*, 1–88.

Cunjak, R. A., & Newbury, R. W. (2005). Atlantic Coast Rivers of Canada. In A. C. Benke & C. E. Cushing (Eds.), *Rivers of North America* (pp. 939–980). San Diego, CA: Elsevier Inc. (Academic Press).

DFO. (2017). Update of indicators to 2017 of adult Altantic salmon for the Miramichi River (NB), Salmon Fishing Area 16, DFO Gulf Region. *Canadian Science Advisory Secretariat Science Response 2017/043*, 1–7.

Dionne, M., Caron, F., Dodson, J. J., & Bernatchez, L. (2008). Landscape genetics and hierarchical genetic structure in Atlantic salmon: The interaction of gene flow and local adaptation. *Molecular Ecology*, *17*(10), 2382–2396. doi:10.1111/j.1365-294X.2008.03771.x

Dionne, M., Caron, F., Dodson, J. J., & Bernatchez, L. (2009). Comparative survey of within-river genetic structure in Atlantic salmon; relevance for management and conservation. *Conservation Genetics*, *10*(4), 869–879. doi:10.1007/s10592-008-9647-5

Dionne, M., Miller, K. M., Dodson, J. J., Caron, F., & Bernatchez, L. (2007). Clinal variation in MHC diversity with temperature: Evidence for the role of host-pathogen interaction on local adaptation in Atlantic salmon. *Evolution*, *61*(9), 2154–2164. doi:10.1111/j.1558-5646.2007.00178.x

Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014).

NeEstimator v2: Re-implementation of software for the estimation of contemporary effective population size (Ne) from genetic data. *Molecular Ecology Resources*, *14*(1), 209–214. doi:10.1111/1755-0998.12157

Dobigny, G., Britton-Davidian, J., & Robinson, T. J. (2017). Chromosomal polymorphism in mammals: an evolutionary perspective. *Biological Reviews*, *92*(1), 1–21. doi:10.1111/brv.12213

Dodson, J., & Colombani, F. (1997). The genetic identity of the Clearwater Brook population of Atlantic salmon ( Salmo salar ); a temporal and spatial study of Atlantic salmon population genetic structure in the Miramichi, St . John and Margaree Rivers. *Atlantic Salmon Federation Report*, 1–24.

Douglas, S. G., Chaput, G., Hayward, J., & Sheasgreen, J. (2015). Assessment of Atlantic Salmon (Salmo salar) in Salmon Fishing Area 16 of the southern Gulf of St. Lawrence to 2013. *Canadian Science Advisory Secretariat Research Document 2015/049*. Retrieved from http://www.dfo-mpo.gc.ca/csas-sccs/

Dray, S., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., Jombart, T., Larocque, G., Legendre, P., Madi, N., & Wagner, H. (2018). adespatial: Multivariate Multiscale Spatial Analysis. R package version 0.1-1. https://CRAN.R-project.org/package=adespatial

Dumas, D., & Britton-Davidian, J. (2002). Chromosomal rearrangements and evolution of recombination: Comparison of chiasma distribution patterns in standard and Robertsonian populations of the house mouse. *Genetics*, *162*(3), 1355–1366.

Elmer, K. R., & Meyer, A. (2011). Adaptation in the age of ecological genomics: insights from parallelism and convergence. *Trends in Ecology & Evolution*, *26*(6), 298–306. doi:10.1016/j.tree.2011.02.008

Elson, P. F. (1967). Effects on Wild Young Salmon of Spraying DDT over New Brunswick Forests. *Journal of the Fisheries Research Board of Canada*, *24*(4), 731–767. doi:10.1139/f67-066

Elson, P. F. (1973). Genetic polymorphism in Northwest Miramishi salmon, in relation to season of river ascent and age at maturation and its implications for management of the stocks. *International Commission for Northwest Atlantic Fisheries Research Document*, *73*(76).

Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, *10*(3), 564–567. doi:10.1111/j.1755-0998.2010.02847.x

Ferchaud, A. L., Perrier, C., April, J., Hernandez, C., Dionne, M., & Bernatchez, L. (2016). Making sense of the relationships between Ne, Nb and Nc towards defining conservation thresholds in Atlantic salmon (Salmo salar). *Heredity*, *117*(4), 268–278. doi:10.1038/hdy.2016.62

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. doi:10.1002/joc.5086

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. *Genetics*, *180*(2), 977–993. doi:10.1534/genetics.108.092221

Fourcade, Y., Chaput-Bardy, A., Secondi, J., Fleurant, C., & Lemaire, C. (2013). Is local selection so widespread in river organisms? Fractal geometry of river networks leads to high bias in outlier detection. *Molecular Ecology*, *22*(8), 2065–2073. doi:10.1111/mec.12158

Fraser, D. J., & Bernatchez, L. (2001). Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Molecular Ecology*, *10*(12), 2741–52. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/11903888

Fraser, D. J., Weir, L. K., Bernatchez, L., Hansen, M. M., & Taylor, E. B. (2011). Extent and scale of local adaptation in salmonid fishes: review and meta-analysis. *Heredity*, *106*(3), 404–420. doi:10.1038/hdy.2010.167

Frontier, S. (1976). Étude de la décroissance des valeurs propres dans une analyse en composantes principales: Comparaison avec le moddle du bâton brisé. *Journal of Experimental Marine Biology and Ecology*, *25*(1), 67–75. doi:10.1016/0022-0981(76)90076-9

Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012). Harnessing genomics for delineating conservation units. *Trends in Ecology and Evolution*, *27*(9), 489–496. doi:10.1016/j.tree.2012.05.012

Garant, D., Dodson, J. J., & Bernatchez, L. (2000). Ecological determinants and temporal stability of the within-river population structure in Atlantic salmon (Salmo salar L.). *Molecular Ecology*, *9*(5), 615–628. doi:10.1046/j.1365-294X.2000.00909.x

Garcia de Leaniz, C., Fleming, I. A., Einum, S., Verspoor, E., Jordan, W. C., Consuegra, S., … Quinn, T. P. (2007). A critical review of adaptive genetic variation in Atlantic salmon: implications for conservation. *Biological Reviews*, *82*(2), 173–211. doi:10.1111/j.1469-185X.2006.00004.x

Gautier, M. (2015). Genome-Wide Scan for Adaptive Divergence and Association with Population-Specific Covariates. *Genetics*, *201*(4), 1555–1579. doi:10.1534/genetics.115.181453

Guerrero, R. F., & Kirkpatrick, M. (2014). Local adaptation and the evolution of chromosome fusions. *Evolution*, *68*(10), 2747–2756. doi:10.1111/evo.12481

Hansen, M. M., Fraser, D. J., Meier, K., & Mensberg, K.-L. D. (2009). Sixty years of anthropogenic pressure: a spatio-temporal genetic analysis of brown trout populations subject to stocking and population declines. *Molecular Ecology*, *18*(12), 2549–2562. doi:10.1111/j.1365-294X.2009.04198.x

Hecht, B. C., Matala, A. P., Hess, J. E., & Narum, S. R. (2015). Environmental adaptation in Chinook salmon ( Oncorhynchus tshawytscha ) throughout their North American range. *Molecular Ecology*, *24*(22), 5573–5595. doi:10.1111/mec.13409

Hendry, A. P., Castric, V., Kinnison, M. T., & Quinn, T. P. (2004). The Evolution of Philopatry and Dispersal: Homing Versus Straying in Salmonids. In A. P. Hendry & S. C. Stearns (Eds.), *Evolution Illuminated: Salmon and thier relatives* (pp. 52–91). New York, NY: Oxford University Press.

Huang, J., Ma, L., Yang, F., Fei, S. Z., & Li, L. (2008). 45S rDNA regions are chromosome fragile sites expressed as gaps in vitro on metaphase chromosomes of root-tip meristematic cells in Lolium spp. *PLoS ONE*, *3*(5), 1–7. doi:10.1371/journal.pone.0002167

Jeffery, N. W., Stanley, R. R. E., Wringe, B. F., Guijarro-Sabaniel, J., Bourret, V., Bernatchez, L., … Bradbury, I. R. (2017). Range-wide parallel climate-associated genomic clines in Atlantic salmon. *Royal Society Open Science*, *4*(11), 171394. doi:10.1098/rsos.171394

Johnston, S. E., Orell, P., Pritchard, V. L., Kent, M. P., Lien, S., Niemelä, E., … Primmer, C. R. (2014). Genome-wide SNP analysis reveals a genetic basis for sea-age variation in a wild population of Atlantic salmon (Salmo salar). *Molecular Ecology*, *23*(14), 3452–3468. doi:10.1111/mec.12832

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide SNP data. *Bioinformatics*, *27*(21), 3070–3071. doi:10.1093/bioinformatics/btr521

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. *BMC Genetics*, *11*(1), 94. doi:10.1186/1471-2156-11-94

Kapun, M., Fabian, D. K., Goudet, J., & Flatt, T. (2016). Genomic Evidence for Adaptive Inversion Clines in Drosophila melanogaster. *Molecular Biology and Evolution*, *33*(5), 1317–1336. doi:10.1093/molbev/msw016

Keefer, M. L., & Caudill, C. C. (2014). Homing and straying by anadromous salmonids: A review of mechanisms and rates. *Reviews in Fish Biology and Fisheries*, *24*(1), 333–368. doi:10.1007/s11160-013-9334-6

Kusche, H., Côté, G., Hernandez, C., Normandeau, E., Boivin-Delisle, D., & Bernatchez, L. (2017). Characterization of natural variation in North American Atlantic Salmon populations (Salmonidae: Salmo salar ) at a locus with a major effect on sea age. *Ecology and Evolution*, *7*(15), 5797–5807. doi:10.1002/ece3.3132

Labadie, H. (2016). *Miramichi Salmon and Trout Restoration - Stocking 2016*. Miramichi River Salmon Association Technical Report. https://miramichisalmon.ca/programs/stocking/

Leggett, W. C. (1977). The ecology of fish migrations. *Annual Review of Ecology and Systematics*, (8), 285–308.

Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., … Davidson, W. S. (2016). The Atlantic salmon genome provides insights into rediploidization. *Nature*, *533*(7602), 200–205. doi:10.1038/nature17164

Lindtke, D., Lucek, K., Soria-Carrasco, V., Villoutreix, R., Farkas, T. E., Riesch, R., … Nosil, P. (2017). Long-term balancing selection on chromosomal variants associated with crypsis in a stick insect. *Molecular Ecology*, *26*(22), 6189–6205. doi:10.1111/mec.14280

Lotterhos, K. E., & Whitlock, M. C. (2014). Evaluation of demographic history and neutral parameterization on the performance of FST outlier tests. *Molecular Ecology*, *23*(9), 2178–2192. doi:10.1111/mec.12725

Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt : an R package to perform genome scans for selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1), 67–77. doi:10.1111/1755-0998.12592

MacDonald, J. R. (1971). Delayed mortality of Atlantic salmon (Salmo salar) parr after forest spraying with DDT. *Canadian Fisheries and Forestry Resource Development Branch Progress Report No. 2*, 1–26.

Matala, A. P., Ackerman, M. W., Campbell, M. R., & Narum, S. R. (2014). Relative contributions of neutral and non-neutral genetic differentiation to inform conservation of steelhead trout across highly variable landscapes. *Evolutionary Applications*, *7*(6), 682–701. doi:10.1111/eva.12174

Meirmans, P. G. (2015). Seven common mistakes in population genetics and how to avoid them. *Molecular Ecology*, *24*(13), 3223–3231. doi:10.1111/mec.13243

Mobley, K. B., Granroth-wilding, H., Ellmen, M., Vähä, J., Johnston, S. E., Orell, P., … Primmer, C. R. (2018). Home ground advantage : selection against dispersers promotes cryptic local adaptation in wild salmon. *BioRxiv*. doi:10.1101/311258

Moller, D. (2005). Genetic studies on serum transferrins in Atlantic salmon. *Journal of Fish Biology*, *67*(s1), 55–67. doi:10.1111/j.0022-1112.2005.00839.x

Moore, J. S., Bourret, V., Dionne, M., Bradbury, I., O'Reilly, P., Kent, M., … Bernatchez, L. (2014). Conservation genomics of anadromous Atlantic salmon across its North American range: Outlier loci identify the same patterns of population structure as neutral loci. *Molecular Ecology*, *23*(23), 5680–5697. doi:10.1111/mec.12972

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs E., & Wagner H. (2017). vegan: Community Ecology Package. R package version 2.4-5. https://CRAN.R-project.org/package=vegan

Olsen, J. B., Crane, P. A., Flannery, B. G., Dunmall, K., Templin, W. D., & Wenburg, J. K. (2011). Comparative landscape genetic analysis of three Pacific salmon species from subarctic North America. *Conservation Genetics*, *12*(1), 223–241. doi:10.1007/s10592-010-0135-3

Paetkau, D., Calvert, W., Stirling, I., & Strobeck, C. (1995). Microsatellite analysis of population structure in Canadian polar bears. *Molecular Ecology*, *4*(3), 347–354. doi:10.1111/j.1365-294X.1995.tb00227.x

Paetkau, D., Slade, R., Burden, M., & Estoup, A. (2004). Genetic assignment methods for the direct, real-time estimation of migration rate: a simulation-based exploration of accuracy and power. *Molecular Ecology*, *13*, 55–65. doi:10.1046/j.1365-294X.2003.02008.x

Palsbøll, P. J., Bérubé, M., & Allendorf, F. W. (2007). Identification of management units using population genetic data. *Trends in Ecology and Evolution*, *22*(1), 11–16. doi:10.1016/j.tree.2006.09.003

Palstra, F. P., O'Connell, M. F., & Ruzzante, D. E. (2007). Population structure and gene flow reversals in Atlantic salmon (Salmo salar) over contemporary and long-term temporal scales: Effects of population size and life history. *Molecular Ecology*, *16*(21), 4504–4522. doi:10.1111/j.1365-294X.2007.03541.x

Palstra, F. P., O'Connell, M. F., & Ruzzante, D. E. (2009). Age structure, changing demography and effective population size in Atlantic salmon (Salmo salar). *Genetics*, *182*(4), 1233–1249. doi:10.1534/genetics.109.101972

Pearse, D. E., Martinez, E., & Garza, J. C. (2011). Disruption of historical patterns of isolation by distance in coastal steelhead. *Conservation Genetics*, *12*(3), 691–700. doi:10.1007/s10592-010-0175-8

Pearse, D. E., Miller, M. R., Abadia-Cardoso, A., & Garza, J. C. (2014). Rapid parallel evolution of standing variation in a single, complex, genomic region is associated with life history in steelhead/rainbow trout. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1783), 20140012–20140012. doi:10.1098/rspb.2014.0012

Pendás, A. M., Morán, P., & Garcia-Vázquez, E. (1993). Ribosomal RNA genes are interspersed throughout a heterochromatic chromosome arm in Atlantic salmon. *Cytogenetic and Genome Research*, *63*(2), 128–130. doi:10.1159/10.1159/000133517

Perrier, C., Baglinière, J. L., & Evanno, G. (2013). Understanding admixture patterns in supplemented populations: A case study combining molecular analyses and temporally explicit simulations in Atlantic salmon. *Evolutionary Applications*, *6*(2), 218–230. doi:10.1111/j.1752-4571.2012.00280.x

Perrier, C., Guyomard, R., Bagliniere, J.-L., & Evanno, G. (2011). Determinants of hierarchical genetic structure in Atlantic salmon populations: environmental factors vs. anthropogenic influences. *Molecular Ecology*, *20*(20), 4231–4245. doi:10.1111/j.1365-294X.2011.05266.x

Pess, G. R. (2009). *Patterns and Processes of Salmon Colonization*. University of Washington, Seattle WA.

Piálek, J., Hauffe, H. C., & Searle, J. B. (2005). Chromosomal variation in the house mouse. *Biological Journal of the Linnean Society*, *84*(3), 535–563. doi:10.1111/j.1095-8312.2005.00454.x

Primmer, C. R., Veselov, A. J., Zubchenko, A., Poututkin, A., Bakhmet, I., & Koskinen, M. T.

(2006). Isolation by distance within a river system: Genetic population structuring of Atlantic salmon, Salmo salar, in tributaries of the Varzuga River in northwest Russia. *Molecular Ecology*, *15*(3), 653–666. doi:10.1111/j.1365-294X.2005.02844.x

Pritchard, V. L., Mäkinen, H., Vähä, J.-P., Erkinaro, J., Orell, P., & Primmer, C. R. (2018). Genomic signatures of fine-scale local selection in atlantic salmon suggest involvement of sexual maturation, energy homeostasis, and immune defence-related genes. *Molecular Ecology*. doi:10.1111/mec.14705

QGIS Development Team (2016). QGIS Geographic Information System. Open Source Geospatial Foundation Project. http://qgis.osgeo.org

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, *26*(6), 841–842. doi:10.1093/bioinformatics/btq033

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org

Riddell, B. E., & Leggett, W. C. (1981). Evidence of an Adaptive Basis for Geographic Variation in Body Morphology and Time of Downstream Migration of Juvenile Atlantic Salmon ( *Salmo salar* ). *Canadian Journal of Fisheries and Aquatic Sciences*, *38*(3), 308–320. doi:10.1139/f81-042

Riddell, B. E., Leggett, W. C., & Saunders, R. L. (1981). Evidence of adaptive polygenic variation between two populations of Atlantic salmon (Salmo salar) native to tributaries of the SW Miramichi River, N. B. *Canandian Journal of Fisheries and Aquatic Science*, *38*(5605 1), 321–333.

Rogers, S. M., Xu, S., & Schlüter, P. M. (2017). Introduction: integrative molecular ecology is rapidly advancing the study of adaptation and speciation. *Molecular Ecology*, *26*(1), 1–6. doi:10.1111/mec.13947

Samy, J. K. A., Mulugeta, T. D., Nome, T., Sandve, S. R., Grammes, F., Kent, M. P., … Våge, D. I. (2017). SalmoBase: An integrated molecular data resource for Salmonid species. *BMC Genomics*, *18*(1), 1–5. doi:10.1186/s12864-017-3877-1

Saunders, R. L. (1967). Seasonal Pattern of Return of Atlantic Salmon in the Northwest Miramichi River, New Brunswick. *Journal of the Fisheries Research Board of Canada*, *24*(1), 21–32. doi:10.1139/f67-003

Sinclair-Waters, M., Bradbury, I. R., Morris, C. J., Lien, S., Kent, M. P., & Bentzen, P. (2017). Ancient chromosomal rearrangement associated with local adaptation of a postglacially colonized population of Atlantic Cod in the northwest Atlantic. *Molecular Ecology*, (November), 1–13. doi:10.1111/mec.14442

Stahl, A., Luciani, J. M., Hartung, M., Devictor, M., Bergé-Lefranc, J. L., & Guichaoua, M. (1983). Structural basis for Robertsonian translocations in man: association of ribosomal genes in the nucleolar fibrillar center in meiotic spermatocytes and oocytes. *Proceedings of the National Academy of Sciences of the United States of America*, *80*(19), 5946–5950. doi:10.1073/pnas.80.19.5946

Tiffin, P., & Ross-Ibarra, J. (2014). Advances and limits of using population genetics to understand local adaptation. *Trends in Ecology & Evolution*, *29*(12), 673–680. doi:10.1016/j.tree.2014.10.004

Tyers, M. (2017). riverdist: River Network Distance Computation and Applications. R package version 0.15.0. https://CRAN.R-project.org/package=riverdist

Vähä, J.-P., Erkinaro, J., Niemelä, E., & Primmer, C. R. (2007). Life-history and habitat features influence the within-river genetic structure of Atlantic salmon. *Molecular Ecology*, *16*(13),

2638–2654. doi:10.1111/j.1365-294X.2007.03329.x

Valiquette, E., Perrier, C., Thibault, I., & Bernatchez, L. (2014). Loss of genetic integrity in wild lake trout populations following stocking: insights from an exhaustive study of 72 lakes from Québec, Canada. *Evolutionary Applications*, *7*(6), 625–644. doi:10.1111/eva.12160

Verspoor, E., Beardmore, J. A., Consuegra, S., Garcia de Leaniz, C., Hindar, K., Jordan, W. C., … Cross, T. F. (2005). Population structure in the Atlantic salmon: insights from 40 years of research into genetic protein variation. *Journal of Fish Biology*, *67*, 3–54. doi:10.1111/j.1095-8649.2005.00838.x

Wallace, B. G., & Curry, R. A. (2017). Assessing the outcomes of stocking hatchery-reared juveniles in the presence of wild Atlantic salmon. *Environmental Biology of Fishes*, *100*(7), 877–887. doi:10.1007/s10641-017-0613-2

Waples, R. K., Larson, W. A., & Waples, R. S. (2016). Estimating contemporary effective population size in non-model species using linkage disequilibrium across thousands of loci. *Heredity*, *117*(4), 233–240. doi:10.1038/hdy.2016.60

Waples, R. S., Antao, T., & Luikart, G. (2014). Effects of Overlapping Generations on Linkage Disequilibrium Estimates of Effective Population Size. *Genetics*, *197*(2), 769–780. doi:10.1534/genetics.114.164822

Waples, R. S., & Do, C. (2010). Linkage disequilibrium estimates of contemporary Ne using highly variable genetic markers: A largely untapped resource for applied conservation and evolution. *Evolutionary Applications*, *3*(3), 244–262. doi:10.1111/j.1752-4571.2009.00104.x

Waples, R. S., & Gaggiotti, O. (2006). What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, *15*(6), 1419–1439. doi:10.1111/j.1365-294X.2006.02890.x

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the analysis of population structure. *Evolution*, *38*(6), 1358–1370. doi:10.2307/2408641

Wellenreuther, M., & Bernatchez, L. (2018). Eco-evolutionary genomics of chromosomal inversions. *Trends in Ecology and Evolution*.

Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible for Local Adaptation: Inference of a Null Model through Trimming the Distribution of F ST. *The American Naturalist*, *186*(S1), S24–S36. doi:10.1086/682949

Yeaman, S., Gerstein, A. C., Hodgins, K. A., & Whitlock, M. C. (2018). Quantifying how constraints limit the diversity of viable routes to adaptation. *BioRxiv*, 279661. doi:10.1101/279661

**Data Accessibility**

The SNP genotype data in PLINK format and the environmental data for sites will be available on Dryad following article acceptance. Scripts used to analyze the data are available as a GitHub repository: https://github.com/kylewellband/ssa_cast2016.

**Author Contributions:**

TL, AC, and LB conceived the study, TL organized sample collection, JE provided the SNP genotyping platform, KW and CM analyzed the data, KW wrote the manuscript, and all authors contributed to the final version.

Table 1: Genetic diversity summary statistics for Atlantic salmon populations from 15 tributaries of the Miramichi River. Populations ordered geographically from North to South and grouped according to the two main branches of the river.

| Tributary | Code | N | $H_O$ | $H_E$ | $F_{IS}$ | $N_b^{Cor}$ |
|---|---|---|---|---|---|---|
| *Northwest branch* | | | | | | |
| Upper Northwest | UNW | 77 | 0.287 | 0.283 | -0.02 | 723 |
| Sevogle | SEV | 48 | 0.289 | 0.285 | -0.03 | 635 |
| Lower Northwest | LNW | 62 | 0.290 | 0.286 | -0.02 | 630 |
| Little Southwest | LSW | 133 | 0.294 | 0.286 | -0.02 | 494 |
| Northwest Millstream | NWM | 47 | 0.301 | 0.280 | -0.08 | 145 |
| *Southwest branch* | | | | | | |
| Renous | REN | 48 | 0.291 | 0.284 | -0.03 | 440 |
| Dungarvon | DUN | 49 | 0.290 | 0.285 | -0.03 | 1783 |
| Bartholomew | BAR | 25 | 0.293 | 0.284 | -0.07 | 556 |
| Cains | CAN | 57 | 0.293 | 0.287 | -0.03 | 678 |
| Southwest | MSW | 71 | 0.295 | 0.287 | -0.03 | 531 |
| Taxis | TAX | 44 | 0.299 | 0.281 | -0.07 | 77 |
| Rocky Brook | RBR | 46 | 0.290 | 0.282 | -0.05 | 108 |
| Clearwater | CWB | 63 | 0.293 | 0.285 | -0.03 | 4300 |
| Burnthill Brook | BHB | 26 | 0.294 | 0.281 | -0.07 | 81 |
| Upper Southwest | USW | 51 | 0.290 | 0.284 | -0.03 | 168 |

N = number of individuals, $H_O$ = observed heterozygosity, $H_E$ = expected heterozygosity, $F_{IS}$ = inbreeding coefficient, $N_b^{Cor}$ = corrected effective number of breeders.

Table 2: Analysis of molecular variance (AMOVA) of tributary populations in the Miramichi River based on neutral SNPs and chromosome 8 and 29 fusion karyotype.

| Source of variation | % Variation | F-statistic | P-value |
|---|---|---|---|
| *Neutral SNPs* | | | |
| Among main river branches | 0.01 | 0.0001 | 0.35 |
| Among tributaries within river branches | 0.42 | 0.0042 | <0.001 |
| Among individuals within tributaries | 99.57 | - | - |
| *Chromosome 8 & 29 fusion* | | | |
| Among main river branches | -0.40 | -0.004 | 0.66 |
| Among tributaries within river branches | 1.90 | 0.019 | <0.001 |
| Among individuals within tributaries | 98.50 | - | - |

Table 3: Genes that occur within 10 Kb of common outlier SNPs detected by Bayescan, OutFlank, FLK, and BayPass.

| Chr | Pos (Mb) | Ssalar Accession # | Gene ID | Name | $N_{SNP}$ |
|---|---|---|---|---|---|
| 3 | 86.287 | CIGSSA_017654 | gtf2f1 | General transcription factor IIF subunit 1 | 1(1) |
| 3 | 86.288 | CIGSSA_019541 | fcl | GDP-L-fucose synthetase | 1(1) |
| 3 | 86.305 | CIGSSA_022019 | gsonmt00000635001 | Uncharacterized protein | 1(1) |
| 6 | 21.845 | CIGSSA_032420 | elfn2 | Uncharacterized protein | 1 |
| 9 | 24.641 | CIGSSA_042815 | tdrd9 | Putative ATP-dependent RNA helicase TDRD9 | 2 |
| 9 | 24.656 | CIGSSA_044891 | rd3 | Protein RD3 | 1 |
| 9 | 24.681 | CIGSSA_044892 | rtn1b | Reticulon | 1 |
| 9 | 24.793 | CIGSSA_042817 | pcnxl4 | Pecanex-like protein 4 | 2(2) |
| 9 | 24.794 | CIGSSA_044893 | dhrs7 | Dehydrogenase/reductase SDR family member 7 | 2(2) |
| 9 | 24.820 | CIGSSA_042818 | ppm1aa | Protein phosphatase type 2C alpha 2 | 5(4) |
| 9 | 24.853 | CIGSSA_047207 | gsonmt00044825001 | Uncharacterized protein | 1(1) |
| 9 | 24.903 | CIGSSA_042819 | six6 | Homeobox protein SIX6 | 2(2) |

Chr = Chromosome number, Pos = approximate position on chromosome in Mb, $N_{SNP}$ = number of outlier SNPs within 10 Kb of the gene (number of SNPs also detected in the polygenic selection framework in parentheses).

Table 4: Gene ontology enrichment for genes located within the region of reduced recombination associated with the fusion of chromosomes 8 and 29 and the 198 genes identified as potentially under polygenic selection by their association with an elevation and temperature gradient. $N_{anno}$ = number of genes in reference set annotated with the specific GO function, $N_{sig}$ = number of genes in fusion region with specific GO annotation, and $N_{exp}$ = number of genes in fusion region with specific GO annotation expected if randomly distributed.

| GO ID | Description | $N_{anno}$ | $N_{sig}$ | $N_{exp}$ | p-value |
|---|---|---|---|---|---|
| *Chromosome 8 and 29 fusion* | | | | | |
| GO:2001045 | negative regulation of integrin-mediated signaling pathway | 15 | 3 | 0.05 | <0.001 |
| GO:0071681 | cellular response to indole-3-methanol | 17 | 3 | 0.05 | <0.001 |
| GO:0021942 | radial glia guided migration of Purkinje cell | 19 | 3 | 0.06 | <0.001 |
| GO:2000146 | negative regulation of cell motility | 850 | 4 | 2.74 | <0.001 |
| GO:0007217 | tachykinin receptor signaling pathway | 28 | 3 | 0.09 | <0.001 |
| GO:0006653 | lecithin metabolic process | 6 | 2 | 0.02 | <0.001 |
| GO:0016264 | gap junction assembly | 43 | 3 | 0.14 | <0.001 |
| GO:0045880 | positive regulation of smoothened signaling pathway | 107 | 4 | 0.34 | <0.001 |
| GO:0030828 | positive regulation of cGMP biosynthetic process | 50 | 3 | 0.16 | 0.001 |
| GO:0007406 | negative regulation of neuroblast proliferation | 51 | 3 | 0.16 | 0.001 |
| GO:0009103 | lipopolysaccharide biosynthetic process | 17 | 2 | 0.05 | 0.001 |
| GO:0007603 | phototransduction, visible light | 377 | 5 | 1.21 | 0.001 |
| GO:0007568 | aging | 1457 | 11 | 4.69 | 0.002 |
| GO:0060134 | prepulse inhibition | 84 | 3 | 0.27 | 0.003 |
| GO:0070474 | positive regulation of uterine smooth muscle contraction | 24 | 2 | 0.08 | 0.003 |
| GO:0043297 | apical junction assembly | 284 | 5 | 0.91 | 0.003 |
| GO:0051823 | regulation of synapse structural plasticity | 87 | 3 | 0.28 | 0.003 |
| GO:2001241 | positive regulation of extrinsic apoptotis in absence of ligand | 87 | 3 | 0.28 | 0.003 |
| GO:0010460 | positive regulation of heart rate | 139 | 4 | 0.45 | 0.004 |
| GO:0071504 | cellular response to heparin | 33 | 2 | 0.11 | 0.005 |
| GO:0035290 | trunk segmentation | 34 | 2 | 0.11 | 0.005 |
| GO:0043278 | response to morphine | 117 | 3 | 0.38 | 0.006 |
| GO:0006776 | vitamin A metabolic process | 38 | 2 | 0.12 | 0.007 |
| GO:0090136 | epithelial cell-cell adhesion | 122 | 3 | 0.39 | 0.007 |
| GO:0007210 | serotonin receptor signaling pathway | 40 | 2 | 0.13 | 0.007 |
| GO:0072047 | proximal/distal nephron patterning | 37 | 2 | 0.12 | 0.009 |
| *Polygenic selection* | | | | | |
| GO:0030705 | cytoskeleton-dependent intracellular transport | 270 | 5 | 1.78 | <0.001 |
| GO:0048870 | cell motility | 2920 | 27 | 19.23 | <0.001 |
| GO:0045137 | development of primary sexual characteristics | 790 | 4 | 5.2 | 0.002 |
| GO:0016559 | peroxisome fission | 10 | 2 | 0.07 | 0.002 |
| GO:0043585 | nose morphogenesis | 14 | 2 | 0.09 | 0.004 |
| GO:0003342 | proepicardium development | 15 | 2 | 0.1 | 0.004 |

| GO:0045163 | clustering of voltage-gated potassium channels | 17 | 2 | 0.11 | 0.005 |
|---|---|---|---|---|---|
| GO:0001675 | acrosome assembly | 17 | 2 | 0.11 | 0.005 |
| GO:2001224 | positive regulation of neuron migration | 57 | 3 | 0.38 | 0.006 |
| GO:0046839 | phospholipid dephosphorylation | 57 | 3 | 0.38 | 0.006 |
| GO:0007166 | cell surface receptor signaling pathway | 4416 | 42 | 29.09 | 0.007 |
| GO:0007616 | long-term memory | 120 | 4 | 0.79 | 0.008 |
| GO:0007175 | inhibition of epidermal growth factor receptor activity | 21 | 2 | 0.14 | 0.008 |

Figure 1: Map of sampling locations for Atlantic salmon and tributary delineation in the Miramichi River (see Table 1 for description of tributary codes).

Figure 2: Characterization of a fusion between Atlantic salmon chromosomes 8 and 29. (A) Principal components analysis revealed three prominent groups corresponding to homokaryote fused (purple / top right), homokaryote unfused (green / bottom left), and heterokaryote (orange / middle) individuals. (B) Genetic differentiation ($F_{ST}$) between the three karyotypes is localized on chromosomes 8 and 29. (C) Genetic diversity (minor allele frequency = MAF, observed heterozygosity = Obs. Het.) is reduced for the fused homokaryotes (AA, purple) and genetic divergence ($F_{ST}$) between homokaryotes (AA vs. BB, green) is elevated in a region of high linkage among chromosomes (grey shading). (D) Linkage disequilibrium ($R^2$) summarized in 1 Mb windows within and between chromosomes 8 and 29. Patterns of linkage suggests the p-arm of chromosome 8 containing rDNA repeated elements fused to the centromere of the acrocentric chromosome 29.

Figure 3: Genetic structure of Atlantic salmon in the Miramichi River based on neutral LD-pruned SNPs. (A) Pairwise $F_{ST}$ among 15 tributary populations. Populations arranged in geographical order from upstream in the northwest branch (NW) to upstream in the southwest branch (SW). See Table 1 for description of site codes. Grey boxes indicate $F_{ST}$ comparisons that are not statistically different from zero based on 10000 permutations of the data. (B) Principal components analysis of allelic variation. Color indicates tributary of sampling, see Table 1 for description of site codes.

Figure 4: Weak spatial structuring of neutral genetic diversity. The first Moran's eigenvector map (MEM1) of pairwise river distance among tributaries explains 1.1% (RDA, 1000 permutations, p = 0.03) of the allelic variation among tributaries. Site loadings on MEM1 (color scale) generally separate tributaries in the upper reaches of the Southwest Miramichi River (dark grey shading) from tributaries in the lower Southwest and Northwest Miramichi River (medium grey shading).

Figure 5: Evidence for polygenic selection among Miramichi River Atlantic salmon populations based on partial redundancy analysis. (A) RDA triplot of the association between tributaries (Northwest = black, Southwest = grey), SNPs, and environmental predictors. The Temp./Elev. summary variable explained 2.6% of the among-population variation in allele frequencies (1000 permutations, p = 0.011). (B) zoom of RDA triplot highlighting 198 outliers (blue points) significantly associated with the Temp./Elev. summary variable (C) Genomic distribution of the strength of SNP association with the Temp./Elev. summary variable (outliers highlighted with blue points).

Figure 6: (A) Statistical associations between each environmental predictor and the frequency of fusion rearrangement or the allelic frequency of outlier SNPs. The outlier SNPs on each chromosome were highly linked and thus only one SNP per chromosome is shown (see Figure S11 for specific correlations with each SNP). Strength and direction of the statistical association (GLM) are indicated by the shape of the ellipse and its colour (red: positive, blue: negative). Stars denote significance at 0.05 level. (B) Summer precipitation predicts fusion frequency (solid black line, GLM z = 3.0, p = 0.002, $R^2$ = 0.21). This pattern of association is present in both the Southwest branch (grey dashed line, GLM z = 3.6, p = 0.002) and Northwest branch (black dashed line, GLM z = 1.7, p = 0.08) of the river.