E-cadherin represses anchorage-independent growth in sarcomas through both signaling and mechanical mechanisms

<u>Running title:</u> E-cadherin represses anchorage-independent growth

Mohit Kumar Jolly[1#], Kathryn E. Ware[2#], Shengnan Xu[2], Shivee Gilja[2], Samantha Shetler[2], Yanjun Yang[1, 3], Xueyang Wang[4], R. Garland Austin[2], Daniella Runyambo[2], Alexander J. Hish[2], Suzanne Bartholf DeWitt[5], Jason T. George[1,6,7], R. Timothy Kreulen[8], Mary-Keara Boss[9], Alexander L. Lazarides[8], David L. Kerr[8], Drew G. Gerber[2], Dharshan Sivaraj[2], Andrew J. Armstrong[10,11], Mark W. Dewhirst[12], William C. Eward[8], Herbert Levine[1,6], and Jason A. Somarelli[2]

[1]Center for Theoretical Biological Physics, Rice University, Houston, Texas, USA; [2]Department of Medicine, Duke University Medical Center, Durham, North Carolina, USA; [3]Department of Applied Physics, Rice University, Houston, Texas, USA; [4]School of Medicine, Johns Hopkins University, Baltimore, Maryland, USA; [5]Department of Pathology, Duke University Medical Center, Durham, North Carolina, USA; [6] Department of Bioengineering, Rice University, Houston, Texas, USA; [7]Medical Scientist Training Program, Baylor College of Medicine, Houston, TX, USA; [8]Department of Orthopaedics, Duke University Medical Center, Durham, North Carolina, USA; [9]Flint Animal Cancer Center, Colorado State University, Fort Collins, Colorado, USA; [10]Solid Tumor Program and [11]Duke Prostate Center, Duke University Medical Center, Durham, North Carolina, USA; [12]Department of Radiation Oncology, Duke University Medical Center, Durham, North Carolina, USA

#These authors contributed equally to this work.

*Address correspondence to:
Jason A Somarelli
Director of Research, Duke Comparative Oncology Group
Department of Medicine, Duke Cancer Institute, Box 103861
905 Lasalle St., GSRBI Room 3044, Durham, North Carolina 27710
Phone: +1 - 919-613-8319
Email: jason.somarelli@duke.edu

Conflict of Interest Statement
The authors declare no potential conflicts of interest.

Key words: anoikis resistance, phenotypic plasticity, E-cadherin, TBX2, CREB

Abstract

E-cadherin, an epithelial-specific cell-cell adhesion molecule, plays multiple roles in maintaining adherens junctions, regulating migration and invasion, and mediating intracellular signaling. Downregulation of E-cadherin is a hallmark of epithelial-mesenchymal transition (EMT) and correlates with poor prognosis in multiple carcinomas. Conversely, upregulation of E-cadherin is prognostic for improved survival in sarcomas. Yet, despite the prognostic benefit of E-cadherin expression in sarcoma, the mechanistic significance of E-cadherin in sarcomas remains poorly understood. Here, by combining mathematical models with wet-bench experiments, we identify the core regulatory networks mediated by E-cadherin in sarcomas, and decipher their functional consequences. Unlike in carcinomas, E-cadherin overexpression in sarcomas does not induce a mesenchymal-epithelial transition (MET). However, E-cadherin acts to reduce both anchorage-independent growth and spheroid formation of sarcoma cells. Ectopic E-cadherin expression acts to downregulate phosphorylated CREB (p-CREB) and the transcription factor, TBX2, to inhibit anchorage-independent growth. RNAi-mediated knockdown of TBX2 phenocopies the effect of E-cadherin on p-CREB levels and restores sensitivity to anchorage-independent growth in sarcoma cells. Beyond its signaling role, E-cadherin expression in sarcoma cells can also strengthen cell-cell adhesion and restricts spheroid growth through mechanical action. Together, our results demonstrate that E-cadherin inhibits sarcoma aggressiveness by preventing anchorage-independent growth.

61
62

Introduction

Sarcomas – deadly cancers that arise from tissues of a mesenchymal lineage – are highly aggressive, with five year survival rates of just 66% (1). Despite their mesenchymal origin, some sarcomas undergo phenotypic plasticity in which they gain "epithelial-like" traits (2–4). While this transition to a more epithelial-like state is now being recognized as a feature of multiple subtypes of soft tissue sarcoma and osteosarcoma (2–4), there are also a number of sarcoma subtypes that are classically known to exhibit epithelioid features pathologically, including synovial sarcoma (5), epithelioid sarcoma (6), and adamantinoma (7). One might expect the acquisition of epithelial-like traits to be of little relevance in mesenchymal tumors, yet that is not the case. Phenotypic plasticity is clinically important in sarcoma patients: Sarcoma patients whose tumors express epithelial-like biomarkers have improved outcomes relative to patients with more "mesenchymal-like" tumors (2–4,8).

Phenotypic plasticity observed in sarcomas is reminiscent of the phenomenon of epithelial plasticity in carcinomas. Epithelial plasticity refers to reversible transitions between epithelial and mesenchymal phenotypes. In carcinomas, the phenotypic transition to a more mesenchymal-like state via an epithelial-mesenchymal transition (EMT) promotes migratory and invasive gene expression programs that facilitate cancer cell invasion and metastatic seeding (9). Subsequent to metastatic dissemination, a reversion to an epithelial-like state via mesenchymal - epithelial transition (MET) re-awakens proliferative signals within the metastatic niche to enable metastatic colonization (9).

In carcinomas, the gene expression programs that control EMT/MET are regulated at multiple stages, including through epigenetics (10), transcription (11), microRNAs (12), alternative splicing (13,14), and post-translational protein stability (15). These regulatory mechanisms control genes involved in cell polarity, cytoskeletal architecture, cell-substrate adhesion, and cell-cell adhesion. One of these genes, E-cadherin, is an epithelial-specific cell-cell adhesion molecule that has multiple functions in maintenance of adherens junctions (16), cytoskeletal organization (17), migration (18,19), and intracellular signaling (20). Downregulation of E-cadherin is a marker of poor prognosis in multiple cancers of an epithelial origin (21,22). In addition, loss-of-function germline mutations in E-cadherin predispose individuals to familial gastric cancer (23), early onset colorectal cancer (24), and hereditary lobular breast cancer (25).

Consistent with its known tumor suppressor role in carcinomas, E-cadherin upregulation is also prognostic for improved survival in sarcomas (8). However, despite the prognostic importance of E-cadherin in sarcomas, little is known about the molecular mechanisms that underlie improved outcomes of E-cadherin upregulation in mesenchymally-derived cancers. Here, we use a combined theoretical-experimental approach to decipher the gene regulatory networks driven by E-cadherin in sarcomas. Though not a generalized phenomenon, in some carcinomas E-cadherin is sufficient to induce a more epithelial-like phenotype (26); however, our study demonstrates E-cadherin expression is not sufficient to alter epithelial plasticity biomarkers, migration, or invasion. E-cadherin expression did, however, significantly inhibit

both anchorage-independent growth and spheroid growth in sarcoma cells. Non-cancer cells that
become detached from the normal tissue architecture undergo a cell death program known as
anoikis. Resistance to anoikis is a hallmark of cancer progression and of an aggressive
phenotype. E-cadherin-mediated repression of anchorage-independent growth was accompanied
by downregulation of phospho-CREB and the transcription factor, TBX2. TBX2 knockdown led
to reductions in total and phospho-CREB and phenocopied E-cadherin-mediated inhibition of
anchorage-independent growth. In addition to its signaling role, E-cadherin repressed spheroid
size through increased cell-cell adhesion. Together, these results indicate that E-cadherin acts
through both signaling and mechanical roles to inhibit sarcoma aggressiveness by suppressing
both anchorage-independent growth and spheroid growth.
121
Materials and Methods
<u>Analysis of the prognostic impact of E-cadherin in clinical data sets</u>
The prognostic significance of E-cadherin in osteosarcoma was analyzed using the R2:
Genomics Analysis and Visualization Platform (https://hgserver1.amc.nl/cgi-bin/r2/main.cgi).
Kaplan Meier plots of metastasis-free survival and overall survival were constructed using the
Kaplan Meier option with the 'mixed osteosarcoma' dataset (27). For soft tissue sarcomas,
Kaplan Meier plots were constructed in JMP13.0 as previously described (Somarelli et al. 2016)
using E-cadherin RNA-Seq and protein levels from The Cancer Genome Atlas (TCGA).
Correlations between E-cadherin levels and EMT score were performed using RNA-Seq data
from TCGA and analyzed in JMP13.0.
132
To calculate EMT scores, raw TCGA sarcoma gene expression data for 259 samples was
accessed from the R2: Genomics Analysis and Visualization Platform (http://r2.amc.nl). EMT
quantification was applied by using a small set of EMT-predictors in addition to a normalization
signature of EMT-uncorrelated transcripts as described previously (28). Each sample was
assigned an EMT score, $0 < \mu < 2$, with E scores $\mu_E < 0.5$, M scores $\mu_M > 1.5$ and hybrid E/M
scores $0.5 \leq \mu_{E/M} \leq 1.5$ intermediary to both E and M scores. Samples were partitioned into E-
cadherin^{low} and E-cadherin^{high} categories based on median E-cadherin expression. The
probability distribution for each group was estimated by spline interpolation of the empirical
EMT score histogram.
142
<u>Cell culture conditions, ectopic E-cadherin expression, and flow cytometry</u>
143B, U2OS, Abrams, and RD cells were cultured in high glucose Dulbecco's Modified Eagle's
Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1%
penicillin/streptomycin (pen-strep) at 37°C and 5% CO_2 in a humidified incubator. 143B, U2OS,
and RD cells were obtained from the Duke University Cell Culture Facility, which performs
routine mycoplasma testing and verifies cell identity by analysis of short tandem repeats. Abrams
cells were a gift of Dr. Douglas Thamm (Colorado State University). To generate sarcoma cells
stably expressing E-cadherin, 143B, U2OS, and RD cells were stably transfected with an empty

vector (pcDNA3) or hE-cadherin-pcDNA3 (a gift from Barry Gumbiner; Addgene plasmid #
45769; (29)). E-cadherin positive cells were sorted by flow cytometry using a PerCP710 anti-E-
cadherin antibody (eBioscience cat. 46-3249). To do this, confluent cells in a T75 flask were
washed with phosphate buffered saline (PBS), scraped, pelleted by centrifugation at 250 x g for
10 minutes, resuspended in 5% bovine serum albumin in PBS, incubated for 30 minutes,
centrifuged as above, resuspended in anti-E-cadherin antibody dilution at 5 μL in 100 μL of
DMEM with 5% FBS per 1 x 10^6 cells, incubated for 30 minutes, centrifuged again, washed with
5 ml of PBS, spun, resuspended in 0.5 ml of DMEM supplemented with 5% FBS, and filtered
through a 30 μM filter (Partek) into a flow cytometry tube. E-cadherin+ cells were sorted into
fully supplemented DMEM at the Duke University Flow Cytometry Shared Resource.

<u>Knockdown by siRNAs</u>
A total of 20 nM of Allstars non-silencing (Qiagen) or target gene siRNA (Qiagen) was diluted
in Opti-MEM and mixed with Lipofectamine RNAiMax diluted in Opti-MEM. The RNAiMax-
siRNA mixtures were incubated for 20 minutes at room temperature, followed by plating of the
mixtures in 24-well plates. A total of 25,000 cells/well diluted in fully supplemented DMEM
were seeded atop the RNAiMax-siRNA mixtures and incubated overnight at 37°C and 5% CO_2
in a humidified incubator. Media was replaced the next day. After 72 hours of incubation, cells
were collected in the appropriate buffer or fixative for downstream applications.
169
<u>Gene expression analysis by qRT-PCR</u>
RNA was extracted from cultured cells using the Zymo Quick RNA Miniprep Kit following the
manufacturer's protocol. Reverse transcription (RT) reactions were performed as described
(Somarelli et al. 2016) using the ABI High Capacity cDNA Reverse Transcription Kit
(ThermoFisher). RT reactions were incubated following the manufacturer's protocol in a
SimpliAmp thermocycler (Life Technologies). RT reactions were diluted 1:5 in nuclease-free
H2O, and RT-qPCR was performed as described (Somarelli et al. 2016) in a Vii7 real time-PCR
detection system (Applied Biosystems).
178
<u>Western blotting, immunofluorescence staining, phospho-kinase arrays, and ELISAs</u>
Western blotting and immunofluorescence staining were performed as previously described
(Somarelli et al. 2016). A complete list of primary antibodies and their dilutions is provided in
the Supplementary Text. To prepare lysates for phospho-kinase arrays, E-cadherin⁻ and E-
cadherin⁺ cells were sorted by flow cytometry prior to each assay, seeded into 10 cm dishes, and
allowed to grow for 48-72 hours until they reached 80% confluence. Cells were then washed
twice with ice-cold PBS and scraped gently in 2 ml of PBS using cell scrapers. Cells were
counted using the CountessII system (Life Technologies), and cells were solubilized at 1 x 10^7
cells/ml in Lysis Buffer 6 provided as part of the Proteome Profiler Human Phospho-Kinase
Array Kit (R&D Systems). Total protein concentrations in 143B+empty vector and 143B+E-
cadherin lysates were estimated using a combination of both Bradford assays and western
blotting (to analyze consistent loading of intact protein with β-actin as a loading control). Lysates

were used in the Proteome Profiler Human Phospho-Kinase Array (R&D cat. ARY003b)
according to the manufacturer's instructions. Membranes were exposed to x-ray films and
visualized on a light box. The same procedure was used to generate lysates for ELISAs, and p-
CREB levels were quantified using DuoSet IC Human/Mouse/Rat Phospho-CREB (S133)
ELISA kits (R&D Systems).
196
<u>Migration, invasion, proliferation, soft agar, spheroid growth assays and cell-cell adhesion assays</u>
For migration and invasion assays, a total of 500 μL of DMEM (with 10% FBS and 1%
pen/strep) was added to the top and bottom of Boyden chamber wells for two hours to hydrate
the transwell membranes. Next, 50,000 cells per well diluted in 500 μL of serum-free DMEM
was added to the top of the chambers, with 500 μL of media containing 10% FBS in the bottom
chambers. Cells were incubated for 24 hours, after which the tops of the chambers were scrubbed
with sterile cotton swabs, and the bottoms were fixed in 4% paraformaldehyde (PFA) for 15
minutes, permeabilized in 0.2% Triton X-100 in PBS for 30 minutes, blocked in 5% BSA in PBS
for 30 minutes, and stained with Hoechst dye (1:2,000 dilution) for one hour. Images were
captured at 40X total magnification via epifluorescence microscopy, and cells were counted
using ImageJ. Invasion was calculated as the proportion of cells migrating through Matrigel-
coated chambers divided by the number of cells migrating through uncoated chambers.
Scratch wound migration assays were performed in 96-well plates for 143B cells and in
24-well plates for RD cells. 143B cells were seeded at 20,000 cells per well and incubated
overnight, after which scratches were made in the wells using a WoundMaker (Essen
Biosciences). For RD cells, 200,000 cells per well were seeded and incubated overnight.
Scratches were made using a 20-200 μl pipette tip. For both cell types, wells were washed with
PBS, and replaced with fully supplemented DMEM. Images were captured every two hours
using the IncuCyte Zoom Live Cell Imaging System. Scratch wound migration of 143B cells was
analyzed using the migration analysis module in the IncuCyte Zoom software, and migration of
RD cells was analyzed using ImageJ.
For monolayer proliferation assays, 25,000 cells per well were plated in 500 μL of
DMEM in 24-well plates, and percent confluence was analyzed using the IncuCyte Zoom
software, with imaging at two hour intervals at 100X total magnification.
Soft agar anchorage-independent growth assays were performed as previously described
(30). Briefly, 25,000 cells were embedded in a 0.3% agarose layer atop a bottom layer of 0.5%
agarose and cultured for approximately three weeks in fully supplemented DMEM in 6-well
plates. Media was replenished weekly, after which colonies were stained with Nitrotetrazoleum
Blue overnight, plates were imaged, and colonies counted using ImageJ.
For spheroid growth assays, cells were scraped in fully supplemented DMEM, and 1,000
cells/well were seeded in ultra-low attachment plates (Costar, cat. #7007). Spheroids were
quantified by manual counting after approximately 2 weeks in culture.
For cell-cell adhesion assays, 12-well plates were incubated overnight with
0.5%BSA/Fraction V in CMFS buffer at 4°C. CMFS buffer contains 4 g NaCl, 0.2g KCl, 0.03g

Na_2HPO_4, and 0.5 g glucose diluted in 500 ml PBS (-/-). Plates were washed with CMFS buffer

and 200,000 cells were immediately seeded in 1mL of media. Plates were incubated on an orbital

shaker at 100 rpm at 37°C for 3 hours, followed by the addition of 1.5 mL of 4% PFA and

incubated at room temperature for 30 minutes. Cells aggregates were immediately quantified in

size categories using ImageJ.

<u>Statistical analyses</u>

All data were analyzed for statistically significant differences using the Student's t-test (for two

comparisons) or analysis of variance with Tukey's post-hoc correction (for multiple

comparisons) in JMP Pro 13. Any p-value < 0.05 was considered statistically significant.

<u>Quantitative modeling for simulating the mechanical effects of E-cadherin</u>

We use a two-dimensional subcellular elements model (31–33) to simulate the effects of

different levels of E-cadherin, where each cell is represented by two subcellular elements, the

front and rear element (inserted cartoon in Fig. 7B (32)). Each subcellular element is self-

propelled with a self-propulsion force m, which balances the intracellular contraction f_{contr}

between the front and rear element. We use a long range attractive and short range repulsive

intercellular force $f_{rep/adh}$ to model cell-cell adhesion (E-cadherin level) and volume exclusion,

respectively. There is a friction between each subcellular element and the substrate with a

constant coefficient ξ. The velocity and position for each subcellular element is updated by: v =

1/ξ (m + f_{contr} + $f_{rep/adh}$), and x = vdt. In our simulation, cells can divide with a probability based

on its size (32). Once the cell length exceeds a certain threshold, it divides at a given probability.

Upon division, two new subcellular elements are inserted, forming two new cells. This process

simulates the cell cycle: the expansion of cell culture is driven by contact inhibition of

locomotion, and cells are elongated during the expansion.

<u>Quantitative mathematical modeling of E-cadherin regulatory network in mediating anchorage</u>

<u>independence</u>

A mathematical model representing the interactions among E-cadherin, TBX2, NRAGE, and

their effects on anchorage independence was constructed. The set of coupled ordinary

differential equations (ODEs) representing the dynamics of these molecular species was

simulated in MATLAB (Mathworks Inc.). The details of model construction and parameters used

is given in supplemental methods.

Results

<u>Aberrant E-cadherin expression in sarcomas is prognostic for improved clinical outcomes</u>

Our previous work found that phenotypic plasticity in sarcomas leads to a more epithelial-like

phenotype (2). This transition to a more epithelial gene expression program has prognostic

relevance: Compared to patients whose sarcomas are more strongly mesenchymal-like, patients

with epithelial-like sarcomas have improved overall survival (2). One of the hallmarks of this
mesenchymal-epithelial transition (MET) is upregulation of the epithelial cell-cell adhesion
molecule, E-cadherin. To further understand the prognostic relevance of E-cadherin in sarcomas,
we stratified osteosarcoma patients into E-cadherin^{high} and E-cadherin^{low} groups using publicly-
available RNA-Seq data (27). E-cadherin^{high} patients had improved metastasis-free survival
(**Figure 1A**) and overall survival compared to E-cadherin^{low} patients (**Figure 1B**). Likewise, soft
tissue sarcoma (STS) patients from The Cancer Genome Atlas with higher E-cadherin mRNA
trended toward better overall survival than patients with lower E-cadherin (**Figure 1C**) while
patients with higher E-cadherin protein had significantly improved overall survival compared to
patients with lower E-cadherin protein levels (**Figure 1D**). These analyses suggest that E-
cadherin has prognostic utility in sarcomas.

<u>Ectopic E-cadherin expression in sarcomas does not promote MET</u>
While it is clear that E-cadherin has prognostic relevance in sarcomas, the underlying molecular
mechanism(s) and cellular phenotypes by which E-cadherin contributes to a less aggressive
tumor remain unknown. Given the important role of E-cadherin as a mediator of the epithelial
phenotype, we hypothesized that E-cadherin may promote an MET-like phenotype in sarcomas,
including suppression of migration and invasion pathways. Consistent with this hypothesis, data
from The Cancer Genome Atlas indicate E-cadherin expression in sarcomas is inversely
correlated with an EMT signature; sarcomas with a more mesenchymal-like gene expression
program have lower E-cadherin while those with a more epithelial-like gene expression program
have higher E-cadherin (**Figure 2A**). To test the hypothesis that E-cadherin inhibits sarcoma
aggression by promoting an MET-like phenotypic switch, we ectopically expressed E-cadherin
in RD human rhabdomyosarcoma and 143B human osteosarcoma cells and assayed for changes
in cell growth. We found E-cadherin expression had no effect on the growth of cells in
monolayer (**Supplementary Figure 1B, 1D**). We next assayed for changes in EMT biomarkers
(Snail, Slug, Twist, Zeb1, Vimentin) by qRT-PCR. With the exception of a statistically
significant two-fold reduction in Slug in 143B cells, E-cadherin had no significant effect on
EMT markers (**Figure 2B, Supplementary Figure 2A**). Consistent with these results, E-
cadherin expression did not alter scratch wound migration (**Figure 2C**) or invasion in 143B cells
(**Figure 2D**) or in RD cells (**Supplementary Figure 2B-C**). We next asked if E-cadherin
status changed the overall EMT score using our previously-derived scoring metric for epithelial,
mesenchymal and hybrid epithelial/mesenchymal states (George, Jolly et al. 2017). Using
empirical probably density functions, we calculated EMT scores for E-cadherin^{high} vs. E-
cadherin^{low} expressing osteosarcomas. The scoring metric was strongly mesenchymal for all
sarcomas analyzed, regardless of E-cadherin status, suggesting that E-cadherin was not capable
of lineage reprogramming toward an epithelial-like state in sarcomas (**Figure 2E**). Visualization
of the EMT score distribution shows only a slight increase in the number of samples predicted as
epithelial or hybrid E/M in cases of E-cadherin^{high} samples relative to the E-cadherin^{low} samples
(**Figure 2E**). Together, using both computational and experimental approaches, our results

suggest that E-cadherin overexpression need not be sufficient to induce MET in sarcomas, and that E-cadherin acts independently of migration/invasion programs to inhibit sarcoma aggressiveness.

E-cadherin suppresses anchorage-independent growth in sarcomas

To explore alternative mechanisms by which E-cadherin may impact the aggressiveness of sarcoma tumors, we analyzed the effect of E-cadherin on anchorage-independent growth. Given the role of E-cadherin as a cell adhesion molecule, we posited that E-cadherin may alter the anchorage-independent growth phenotype of sarcoma cells. To test this hypothesis, cells expressing an empty vector (EV) or E-cadherin were cultured in soft agar to simulate anchorage-independent growth. Interestingly, E-cadherin expression significantly inhibited the number of colonies formed in soft agar growth assays in 143B cells (**Figure 3A**). This effect was independent of differences in monolayer growth, as E-cadherin expressing cells showed no difference in 2D growth (**Supplementary Figure 1B, D**). We observed similar results in a second sarcoma cell line, RD cells. Ectopic expression of E-cadherin did not induce MET by gene expression, migration, or invasion (**Supplementary Figure 2A-C**). However, consistent with 143B cells, E-cadherin overexpression inhibited anchorage independent growth (**Supplementary Figure 2D, E**). We next measured spheroid formation to assess the impact of E-cadherin on anchorage-independent growth and morphology. For these experiments, we noted that, despite consistent downregulation of soft agar growth in both RD and 143B cells, E-cadherin was not expressed at the cell membrane of RD cells (**Supplementary Figure 1A**), while E-cadherin was localized at the membrane of 143B cells (**Supplementary Figure 1C**). Because membrane E-cadherin is the most typically-characterized aspect of E-cadherin function, we focused on 143B cells for which E-cadherin was correctly localized to the cell membrane. We also included U2OS cells as a second model for which E-cadherin was localized to the cell membrane (**Supplementary Figure 1E**). Consistent with our result in soft agar, E-cadherin expressing spheroids formed as tight aggregates and significantly reduced the size of spheroid formation in both 143B and U2OS cells (**Figure 3B, C**). These results indicate that sarcomas with E-cadherin expression have suppressed anchorage-independent growth as indicated by their deficient ability to grow suspended in soft agar.

E-cadherin-mediated sensitivity to anchorage-independent growth acts, in part, through CREB

To determine the signaling pathway(s) through which E-cadherin inhibits anchorage-independent growth in sarcomas, we performed phospho-kinase arrays on 143B osteosarcoma cells with and without E-cadherin expression. The phospho-kinase arrays contain antibodies against 43 different kinases (R&D Systems). From these arrays, we identified phospho (p)-CREB as consistently downregulated in E-cadherin expressing 143B cells as compared to 143B cells expressing an empty vector lacking E-cadherin (**Figure 4A**). The inhibition of phospho- and total CREB in E-cadherin expressing 143B cells was validated by enzyme linked immunosorbent assays (ELISAs) and western blotting (**Figure 4B, C**). Consistent with this, we also observed

lower levels of phospho- and total CREB in Abrams canine osteosarcoma cells expressing E-cadherin as compared to cells expressing an empty vector lacking E-cadherin (**Figure 4D**).

We next asked if E-cadherin-mediated CREB inhibition was responsible for the reductions in anchorage-independent growth capacity of E-cadherin expressing cells. To do this, we knocked down CREB using two independent siRNAs and verified knockdown at the mRNA and protein levels in 143B cells, with siRNA_5 providing a stronger knockdown (**Figure 4E, F**). We noted a consistent reduction in soft agar growth with both CREB siRNAs compared to a non-silencing siRNA (**Figure 4G**); however, there was only a significant inhibition of soft agar growth in cells treated with the most effective siRNA_5 (**Figure 4G**). Importantly, CREB knockdown had no effect on E-cadherin expression (Supplementary Figure 3A). We conclude from these data that E-cadherin inhibition of anchorage-independent growth is mediated through CREB, and downregulation of CREB partially phenocopies E-cadherin expression, although other factors are likely at play in driving anchorage-independent growth in sarcomas.

During our analysis of MET biomarkers and phenotypes in E-cadherin expressing cells, we noted that groups of E-cadherin+ 143B cells lost cell surface N-cadherin expression (**Supplementary Figure 4A**). Interestingly, the reduction in cell surface N-cadherin was not due to changes in N-cadherin expression at either the mRNA or protein levels (**Supplementary Figure 4B, C**). Cadherin switching has been observed in carcinomas in which an E-cadherin to N-cadherin switch is prognostic for poorer clinical outcomes (34–36). In addition, N-cadherin has been shown to promote anchorage-independent growth in carcinomas (37). Thus, we tested if a reduction of cell surface N-cadherin was responsible for the downregulation of anchorage-independent growth in E-cadherin+ cells. Knockdown of N-cadherin with two independent siRNAs (**Supplementary Figure 4D**) had no effect on 143B colony formation in soft agar (**Supplementary Figure 4E**). Likewise, N-cadherin knockdown did not alter levels of p-CREB protein, as determined by p-CREB ELISAs and western blotting (**Supplementary Figure 4F, G**). These results suggest that E-cadherin inhibits anchorage-independent growth through a pathway that is independent of N-cadherin.

An E-cadherin/TBX2/CREB axis modulates anchorage-independent growth of sarcomas
To elucidate further the signaling axis that modulates E-cadherin/CREB-mediated suppression of anchorage-independent growth, we investigated additional signaling axes that may be involved with E-cadherin. One of these axes, TBX2, is a transcription factor that can directly repress E-cadherin (38), can be modulated by CREB (39), and is overexpressed in rhabdomyosarcoma (40,41). To determine if E-cadherin regulates TBX2 in sarcomas, we measured levels of TBX2 in E-cadherin expressing 143B cells. TBX2 mRNA was significantly downregulated in E-cadherin+ 143B cells (**Figure 5A**). Similarly, U2OS cells ectopically expressing E-cadherin (**Supplementary Figure 1E**) had significantly lower levels of TBX2 mRNA (**Figure 5B**). To further delineate the directionality of the E-cadherin/CREB/TBX2 signaling network, we performed siRNA-mediated CREB knockdown and measured TBX2 mRNA levels. Interestingly, CREB knockdown had no effect on TBX2 expression (**Figure 5C,**

390 **Supplementary Figure 3B**), suggesting that CREB is downstream or independent of TBX2 in
391 this gene regulatory network. Consistent with these data, TBX2 knockdown (**Figure 5D**) led to
392 significant reductions in CREB mRNA (**Figure 5E**) and protein (**Figure 5F**). Taken together,
393 these results suggest that E-cadherin inhibits TBX2, which leads to CREB downregulation and
394 suppression of anchorage-independent growth.
395
<u>Mathematical modeling of E-cadherin/TBX2 axis regulating anchorage-independent growth</u>
By integrating our quantitative RT-PCR result showing E-cadherin-mediated suppression of
TBX2 mRNA with previous reports suggesting E-cadherin to be a transcriptional target of TBX2
(38), and regulation of anchorage-independent growth by E-cadherin and/or TBX2 (40,42,43),
we constructed a mechanism-based mathematical model to decode the dynamics of the E-
cadherin/TBX2 axis in mediating anchorage-independent growth (**Figure 6A**). This model
includes the following interactions: (i) mutual repression between E-cadherin and TBX2, (ii)
inhibition of NRAGE (neurotrophin receptor-interacting melanoma antigen) by E-cadherin
(Kumar et al., 2011), and (iii) inhibition of anchorage-independent growth by the TBX2
complex, which is promoted in the presence of NRAGE (42). Such interconnected regulatory
loops can often obviate an intuitive understanding of the emergent outcomes of these interactions
(44). Thus, mechanism-based mathematical models can be a powerful tool for both helping
explain previous experimental observations, and for making novel predictions to guide future
experimental design (45).
Our model predicted that TBX2 knockdown would restrict anchorage-independent
growth in a similar way as that by E-cadherin overexpression (**Figure 6B**). To test this
prediction, we used two independent siRNA to TBX2 to knockdown expression and measured
anchorage independent growth. Consistent with our prediction, TBX2 knockdown using two
independent siRNAs significantly abrogated colony growth in 143B cells (**Figure 6C**). Further,
the model predicted that knockdown of TBX2 under ectopic expression of E-cadherin
accentuates sensitivity to anchorage independence (**Figure 6B**). However, TBX2 knockdown
had no effect on spheroid formation, when E-cadherin was overexpressed in U2OS cells (**Figure
6D**).
The inconsistency in model predictions vs. experimental observations drew our attention
to multiple underlying assumptions of the mathematical model – (a) TBX2-mediated inhibition
of E-cadherin affects both mRNA and protein levels of E-cadherin, (b) endogenous production of
E-cadherin is enough to inhibit TBX2. The first assumption is based on observations in breast
epithelial cells MCF10A (38); similarly, the second assumption also is likely to be more typical
of carcinomas than of sarcomas. Given that our model was constructed using data exclusively
taken from carcinomas, we reasoned that the discrepancy between the model and the
experimental validation may be due to differences in regulatory networks between carcinomas
and sarcomas. Therefore, to determine if similar regulatory networks between TBX2 and E-
cadherin exist in sarcomas, we knocked down TBX2 with two independent siRNAs in U2OS
cells and measured E-cadherin mRNA and protein levels. TBX2 knockdown led to a modest,

albeit significant, two- to four-fold upregulation of E-cadherin mRNA levels **(Figure 6E)**. More
importantly, this modest change in expression in a cell line with relatively low levels of E-
cadherin mRNA expression at baseline was not enough of an increase in mRNA to show a
change at the protein level **(Figure 6F)**. The relative lack of E-cadherin re-activation upon TBX2
knockdown led us to hypothesize that sarcoma cells had alterations in the promoter of E-cadherin
that prevented robust increases in expression upon TBX2 downregulation. Consistent with this
hypothesis, analysis of sarcoma tumors from The Cancer Genome Atlas (TCGA) revealed that
the E-cadherin promoter was significantly more methylated in sarcomas as compared to that in
carcinomas **(Figure 6G)**. Moreover, the degree of E-cadherin promoter methylation was lowest
in synovial sarcoma (**Figure 6H**), which corresponded with a relative increase in E-cadherin
expression (**Figure 6I**). Interestingly, synovial sarcoma often displays epithelioid histological
features, including E-cadherin upregulation. This suggests that one potential difference between
the E-cadherin/TBX2 axis in sarcomas vs. carcinomas is the ability of TBX2 to access the E-
cadherin promoter, which may impact the mathematical models of networks based from largely
studies in carcinoma models. Incorporating this new experimental data, we revised our
mathematical model to include a weaker inhibition of E-cadherin by TBX2. The revised model
predicted that, consistent with our experimental results, the effect of TBX2 knockdown on
anchorage-independent growth in the context of E-cadherin overexpression was reduced
(**Supplementary Figure 5**).
449
<u>A mechanical model relates E-cadherin mediated cell-cell adhesion to spheroid formation</u>
Our experimental results suggest E-cadherin inhibits soft agar colony formation mediated by
suppression of CREB and TBX2 in sarcoma cells. Interestingly, E-cadherin expression alone is
sufficient to suppress the size of spheroid formation while TBX2 has no effect on spheroid
growth **(Figure 6D)**. The differences in response between these phenotypic assays suggest that
E-cadherin may act through different molecular mechanisms to suppress anchorage-independent
growth and spheroid growth. Given the well-described role for E-cadherin as an adhesion
molecule, we hypothesized that E-cadherin can regulate spheroid formation through its
mechanical role as a regulator of cell-cell adhesion. Indeed, E-cadherin mediated cell-cell
adhesion has been proposed to be crucial in mediating formation of cohesive multicellular units
such as tumor emboli and invasive collective migration in breast cancer (19,46–48). Consistent
with this, U2OS cells overexpressing E-cadherin formed tight multicellular clusters at 24 hours
of spheroid formation as compared to control cells **(Figure 7A)**.
To quantitatively capture the relationship between cell-cell adhesion and spheroid size,
we developed a two-dimensional mechanical model of sphere formation. This model represents
each cell by two subcellular elements connected to each other with a contractile force, and
incorporates various interconnected biomechanical aspects of cellular aggregation, such as
intracellular contraction, intercellular adhesion, cell proliferation, self-propulsion forces of the
cells, and friction between the cells and the substrate (31–33). We initialized our simulations
with a seed colony of 233 cells arranged as a two-dimensional circular sheet, and simulated the

colony growth for varying levels of intercellular adhesion, while keeping all other parameters the
same. The intercellular interaction is repulsive at short distances, attractive at longer distances
and becomes zero further away, which models the volume exclusion and cell-cell adhesion. For
different levels of adhesion, the attractive part of the intercellular force follows different curves
(Figure 7B). This mechanical model predicts a decrease in final spheroid size and the total
number of cells with an increase in E-cadherin levels **(Figure 7C)**, and a concomitant increase in
cell density **(Supplementary Figure 6)**. Thus, the model predicts that cell-cell adhesion can
regulate colony size through mechanical factors.
To test if E-cadherin modulates cell-cell adhesion in sarcomas, we quantified the
numbers of single cells and cellular aggregates (two or more cells clumped together) using cell-
cell adhesion assays for both 143B and U2OS cells upon ectopic expression of an empty vector
or E-cadherin. Consistent with our model, we observed that E-cadherin overexpression drives a
significant enrichment of multicellular aggregates in both 143B and U2OS cells **(Figure 7D-E),**
suggesting that E-cadherin levels can play a fundamental role in enabling cell-cell adhesion in
sarcomas. Together, our results integrating gene regulatory network models, mechanical models,
and experimental validations indicate that E-cadherin plays a dual role in suppressing sarcoma
aggression that is independent of EMT through both signaling-related inhibition of anchorage-
independent growth and mechanics-based inhibition of spheroid formation.
488
489 **Discussion**
490
Our integrated computational-experimental study illustrates that E-cadherin inhibits sarcoma
aggressiveness in at least the following two ways: first, by suppressing anchorage-independent
growth through a CREB/TBX2 signaling axis; and second, by increasing cell-cell adhesion and
restricting spheroid growth. Anchorage independent growth is a hallmark of cancer and a
contributor to metastatic growth (50,51). Although normal cells require attachment to the ECM
to receive cell survival signals, cancer cells can overcome the death signals that are driven by
loss of cell-ECM attachment (49). This can lead to anchorage-independent survival of otherwise
adherent cells, including at sites where a unique ECM composition prevents cellular attachment
(49). Interestingly, anchorage-independent growth has often been studied in the context of EMT
in multiple epithelial cancers (52–54); however, ours is among the minority of studies connecting
E-cadherin with anchorage independence in sarcomas. Indeed, our results pinpoint an E-
cadherin/p-CREB/TBX2 axis that mediates anchorage-independent growth in sarcoma cells. This
signaling axis is independent of MET, as neither ectopic E-cadherin expression nor TBX2
knockdown had any effect on MET markers in 143B or RD cells. This observation reinforces
previous results that overexpression of transcription factor GRHL2 – an activator of E-cadherin –
was sufficient to induce MET in breast cancer cells having undergone EMT, but not in sarcoma
(2,55). Highly methylated promoters of both E-cadherin (**Figure 6G**) and GRHL2 (2) in
sarcomas relative to carcinomas, as observed from analysis of TCGA data, can potentially
explain this difference. In this scenario, epigenetic silencing of epithelial promoters would

prevent lineage reprogramming and MET. However, it is possible that some sarcoma subtypes
are more likely to undergo MET; for instance, E-cadherin expression (**Supplementary Figure**
1A), and an epithelial-like phenotype in general (56), appears to be more frequently upregulated
in certain soft tissue sarcoma subtypes, such as synovial sarcomas, malignant peripheral nerve
sheath tumors, and leiomyosarcomas (**Figure 6I**).

While E-cadherin inhibits anchorage-independent growth through a signaling axis, the
role of E-cadherin in suppressing spheroid growth is mediated, at least in part, through its
mechanical activity of increasing cell-cell adhesion and cellular aggregation (**Figure 7**). This
observation is consistent with previous reports in which knock down of E-cadherin led to less
cell-cell adhesion and lower multicellular aggregation in colorectal cancer cells (57). However,
unlike our study, this study in colorectal cancer also found that E-cadherin loss led to increased
migration and invasion. Yet, despite the fact that E-cadherin did not change migration or
invasion in sarcomas, E-cadherin may still inhibit invasiveness of some tumors not by
maintaining an epithelial state, but by increasing cell-cell adhesion. In this way, E-cadherin
would indirectly reduce invasion by preventing cells from detaching from the bulk tumor without
changing the EMT factors involved in migration and/or invasion.
The signaling and mechanical modes of action we observed in suppressing anchorage-
independent growth and spheroid formation, respectively, need not be independent of each other,
especially given the interconnections among mechanosensitive and biochemical signaling
pathways. For instance, Notch signaling can be mechanosensitive (58), give rise to multiple
multicellular patterns of cells in varying phenotypic states (59,60), and be involved in anoikis
(61). Similarly, E-cadherin can sense and provide mechanical cues for epithelial homeostasis and
collective cell migration (62–64). Future studies will be aimed at elucidating the crosstalk
between inter-cellular mechano-sensitive and intra-cellular p-CREB/TBX2 modules to yield new
insights into the mechanisms underlying non-cell autonomous control of anoikis, as recently
observed (65). Moreover, it would be intriguing to compare the results for soft agar colony
growth and spheroid formation assays for RD cells where E-cadherin exerts an effect on
anchorage-independent growth in the absence of membrane localization.
Overall, our results provide a mechanistic basis for the observation that E-cadherin is a
prognostic biomarker for improved clinical outcomes in sarcomas. Although E-cadherin
upregulation in the clinical setting is likely to be a harbinger of a broader change in EMT/MET
programs, our data argue that E-cadherin upregulation also has important effects on sarcoma
aggressiveness that are independent of MET induction. Together, these results indicate that
targeting the EMT/MET axis in sarcomas may have multiple benefits, including downregulation
of stemness, migration, invasion, and anchorage-independent growth pathways. One way this
might be achieved is through epigenetic reprogramming. Indeed, multiple reports in carcinomas
suggest that treatment with epigenetic modifiers, including histone deacetylases inhibitors, can
upregulate E-cadherin (66–69). Consistent with this hypothesis, when anchorage-dependent and
anchorage-independent subpopulations of osteosarcoma cells are compared, the anchorage-
independent subpopulations are significantly more susceptible to epigenetic modifying therapies

(70). A similar strategy could be used to drive sarcomas toward a more epithelial-like phenotype
and lower their aggressiveness by targeting many aggressive phenotypes simultaneously. Such
combinatorial therapies may limit the adaptive phenotypic plasticity of cancer cells – a crucial
determinant of therapeutic resistance (71).
554

Acknowledgments
The results published here are in part based upon data generated by the TCGA Research
Network: http://cancergenome.nih.gov. The authors acknowledge the Duke University Cell
Culture Facility and the Duke Cancer Institute Flow Cytometry Shared Resource. HL was
supported by NSF Center for Theoretical Biological Physics (NSF PHY-1427654) and NSF
DMS-1361411, and as a CPRIT (Cancer Prevention and Research Institute of Texas) Scholar in
Cancer Research of the State of Texas at Rice University. MKJ received a training fellowship
from the Gulf Coast Consortia on the Computational Cancer Biology Training Program (CPRIT
grant no. RP170593). KEW was supported by the NIH F32 CA192630. JAS wishes to
acknowledge support from Meg and Bill Lindenberger, the Paul and Shirley Friedland Fund, the
Triangle Center for Evolutionary Medicine, and funds raised in memory of Muriel E.
Rudershausen (riding4research.org).
567

References
1.   Siegel R, Miller K, Jemal A. Cancer statistics , 2017 . CA Cancer J Clin. 2017;67:7–30.

2.   Somarelli JA, Shetler S, Jolly MK, Wang X, Bartholf Dewitt S, Hish AJ, et al.
Mesenchymal-epithelial transition in sarcomas is controlled by the combinatorial
expression of microRNA 200s and GRHL2. Mol Cell Biol. 2016;36:2503–13.

3.   Yin K, Liao Q, He H, Zhong D. Prognostic value of Twist and E-cadherin in patients with
osteosarcoma. Med Oncol. 2012;29:3449–55.

4.   Tian W, Wang G, Yang J, Pan Y, Ma Y. Prognostic role of E-cadherin and Vimentin
expression in various subtypes of soft tissue leiomyosarcomas. Med Oncol. 2013;30:401.

5.   Thway K, Fisher C. Synovial sarcoma: defining features and diagnostic evolution. Ann
Diagn Pathol. 2014;18:369–80.

6.   Chang J, Erler J. Hypoxia-mediated metastasis. Adv Exp Med Biol2. 2014;772:55–81.

7.   Sarita-Reyes CD, Greco MA, Steiner GC. Mesenchymal-epithelial differentiation of
adamantinoma of long bones: an immunohistochemical and ultrastructural study.
Ultrastruct Pathol. 2012;36:23–30.

8.   Wang N, He Y-L, Pang L-J, Zou H, Liu C-X, Zhao J, et al. Down-regulated E-cadherin
expression is associated with poor five-year overall survival in bone and soft tissue
sarcoma: Results of a meta-analysis. PLoS One. 2015;10:e0121448.

9.   Nieto MA, Huang RY, Jackson RA, Thiery JP. EMT: 2016. Cell. 2016;166:21–45.

10.   Tam WL, Weinberg RA. The epigenetics of epithelial-mesenchymal plasticity in cancer.
588   Nat Med. 2013;19:1438–49.

589   11.   De Craene B, Berx G. Regulatory networks defining EMT during cancer initiation and
590   progression. Nat Rev Cancer. 2013;13:97–110.

591   12.   Lu M, Jolly MK, Levine H, Onuchic JN, Ben-Jacob E. MicroRNA-based regulation of
592   epithelial–hybrid–mesenchymal fate determination. Proc Natl Acad Sci. 2013;110:18144–
593   9.

594   13.   Brown RL, Reinke LM, Damerow MS, Perez D, Chodosh L a., Yang J, et al. CD44 splice
595   isoform switching in human and mouse epithelium is essential for epithelial-mesenchymal
596   transition and breast cancer progression. J Clin Invest. 2011;121:1064–74.

597   14.   Jia D, Jolly MK, Tripathi SC, Hollander P Den, Huang B, Lu M, et al. Distinguishing
598   Mechanisms Underlying EMT Tristability. Cancer Converg. 2017;1:2.

599   15.   De Herreros AG, Peiró S, Nassour M, Savagner P. Snail family regulation and epithelial
600   mesenchymal transitions in breast cancer progression. J Mammary Gland Biol Neoplasia.
601   2010;15:135–47.

602   16.   Lewis JE, Jensen PJ, Johnson KR, Wheelock MJ. E-cadherin mediates adherens junction
603   organization through protein kinase C. J Cell Sci. 1995;107:3615–21.

604   17.   Chen A, Beetham H, Black MA, Priya R, Telford BJ, Guest J, et al. E-cadherin loss alters
605   cytoskeletal organization and adhesion in non-malignant breast cells but is insufficient to
606   induce an epithelial-mesenchymal transition. BMC Cancer. 2014;14:552.

607   18.   Chen W, Obrink B. Cell-cell contacts mediated by E-cadherin (uvomorulin) restrict
608   invasive behavior of L-cells. J Cell Biol. 1991;114:319–27.

609   19.   Jolly MK, Boareto M, Debeb BG, Aceto N, Farach-Carson MC, Woodward WA, et al.
610   Inflammatory Breast Cancer: a model for investigating cluster-based dissemination. NPJ
611   Breast Cancer. 2017;3:21.

612   20.   Orsulic S, Huber O, Aberle H, Arnold S, Kemler R. E-cadherin binding prevents beta-
613   catenin nuclear localization and beta-catenin/LEF-1-mediated transactivation. J Cell Sci.
614   1999;112:1237–45.

615   21.   Li Z, Yin S, Zhang L, Liu W, Chen B. Prognostic value of reduced E-cadherin expression
616   in breast cancer: a meta-analysis. Oncotarget [Internet]. 2017;8:16445–55. Available
617   from: http://www.oncotarget.com/abstract/14860

618   22.   Hong S-M, Li A, Olino K, Wolfgang CL, Herman JM, Schulick RD, et al. Loss of E-
619   cadherin expression and outcome among patients with resectable pancreatic
620   adenocarcinomas. Mod Pathol [Internet]. 2011;24:1237–47. Available from:
621   http://www.nature.com/doifinder/10.1038/modpathol.2011.74

622   23.   Guilford P, Hopkins J, Harraway J, McLeod M, McLeod N, Harawira P, et al. E-cadherin
623   germline mutations in familial gastric cancer. Nature. 1998;392:402–5.

24. Richards FM, McKee SA, Rajpar MH, Cole TRP, Evans DGR, Jankowski JA, et al.
Germline E-cadherin gene (CDH1) mutations predispose to familial gastric cancer and
colorectal cancer. Hum Mol Genet. 1999;8:607–10.

25. Corso G, Intra M, Trentin C, Veronesi P, Galimberti V. CDH1 germline mutations and
hereditary lobular breast cancer. Fam Cancer. Springer Netherlands; 2016;15:215–9.

26. Chao YL, Shepard CR, Wells A. Breast carcinoma cells re-express E-cadherin during
mesenchymal to epithelial reverting transition. Mol Cancer. 2010;9:179.

27. Kuijjer ML, Rydbeck H, Kresse SH, Buddingh EP, Lid AB, Roelofs H, et al.
Identification of osteosarcoma driver genes by integrative analysis of copy number and
gene expression data. Genes Chromosom Cancer. 2012;51:696–706.

28. George JT, Jolly MK, Xu S, Somarelli JA, Levine H. Survival Outcomes in Cancer
Patients Predicted by a Partial EMT Gene Expression Scoring Metric. Cancer Res.
2017;77:6415–28.

29. Gottardi CJ, Wong E, Gumbiner BM. E-cadherin suppresses cellular transformation by
inhibiting β-catenin signaling in an adhesion-independent manner. J Cell Biol.
2001;153:1049–59.

30. Borowicz S, Van Scoyk M, Avasarala S, Karuppusamy Rathinam MK, Tauler J,
Bikkavilli RK, et al. The Soft Agar Colony Formation Assay. J Vis Exp [Internet].
2014;e51998. Available from: http://www.jove.com/video/51998/the-soft-agar-colony-
formation-assay

31. Basan M, Elgeti J, Hannezo E, Rappel W-JW-J, Levine H. Alignment of cellular motility
forces with tissue flow as a mechanism for efficient wound healing. Proc Natl Acad Sci U
S A. 2013;110:2452–9.

32. Zimmermann J, Camley BA, Rappel W-J, Levine H. Contact inhibition of locomotion
determines cell – cell and cell – substrate forces in tissues. Proc Natl Acad Sci.
2016;113:2660–5.

33. Yang Y, Levine H. Role of the supracellular actomyosin cable during epithelial wound
healing. Soft Matter. 2018;14:4866–73.

34. Gravdal K, Halvorsen OJ, Haukaas SA, Akslen LA. A switch from E-cadherin to N-
cadherin expression indicates epithelial to mesenchymal transition and is of strong and
independent importance for the progress of prostate cancer. Clin Cancer Res.
2007;13:7003–11.

35. Hazan RB, Qiao R, Keren R, Badano I, Suyama K. Cadherin switch in tumor progression.
Ann N Y Acad Sci. 2004;1014:155–63.

36. Jager T, Becker M, Eisenhardt A, Tilki D, Totsch M, Schmid KW, et al. The prognostic
value of cadherin switch in bladder cancer. Oncol Rep. 2010;23:1125–32.

37. Ko H, Kim S, Jin C-H, Lee E, Ham S, Yook JI, et al. Protein Kinase Casein Kinase 2-

Mediated Upregulation of N-Cadherin Confers Anoikis Resistance on Esophageal
Carcinoma Cells. Mol Cancer Res. 2012;10:1032–8.

38.   Wang B, Lindley LE, Fernandez-Vega V, Rieger ME, Sims AH, Briegel KJ. The T box
transcription factor TBX2 promotes epithelial-mesenchymal transition and invasion of
normal and malignant breast epithelial cells. PLoS One. 2012;7:e41355.

39.   Liu F, Cao J, Lv J, Dong L, Pier E, Xu GX, et al. TBX2 expression is regulated by PAX3
in the melanocyte lineage. Pigment Cell Melanoma Res. 2013;26:67–77.

40.   Zhu B, Zhang M, Byru S, Tackett AJ, Davie JK. TBX2 blocks myogenesis and promotes
proliferation in rhabdomyosarcoma cells. Int J Cancer. 2014;135:785–97.

41.   Zhu B, Zhang M, Williams EM, Keller C, Mansoor A, Davie JK. TBX2 represses PTEN
in rhabdomyosarcoma and skeletal muscle. Oncogene. 2016;35:4212–24.

42.   Kumar S, Park SH, Cieply B, Schupp J, Killiam E, Zhang F, et al. A Pathway for the
Control of Anoikis Sensitivity by E-Cadherin and Epithelial-to-Mesenchymal Transition.
Mol Cell Biol. 2011;31:4036–51.

43.   Lingbeek ME, Jacobs JJL, Van Lohuizen M. The T-box repressors TBX2 and TBX3
specifically regulate the tumor suppressor gene p14ARF via a variant T-site in the
initiator. J Biol Chem. 2002;277:26120–7.

44.   Jolly MK, Tripathi SC, Somarelli JA, Hanash SM, Levine H. Epithelial/mesenchymal
plasticity: how have quantitative mathematical models helped improve our understanding?
Mol Oncol. 2017;11:739–54.

45.   Mobius W, Laan L. Physical and Mathematical Modeling in Experimental Papers. Cell.
2015;163:1577–83.

46.   Elisha Y, Kalchenko V, Kuznetsov Y, Geiger B. Dual role of E-cadherin in the regulation
of invasive collective migration of mammary carcinoma cells. Sci Rep. 2018;8:4986.

47.   Shamir ER, Pappalardo E, Jorgens DM, Coutinho K, Tsai WT, Aziz K, et al. Twist1-
induced dissemination preserves epithelial identity and requires E-cadherin. J Cell Biol.
2014;204:839–56.

48.   Shamir ER, Ewald AJ. Adhesion in mammary development: novel roles for E-cadherin in
individual and collective cell migration. Curr Top Dev Biol. 2015;112:353–82.

49.   Paoli P, Giannoni E, Chiarugi P. Anoikis molecular pathways and its role in cancer
progression. Biochim Biophys Acta - Mol Cell Res. 2013;1833:3481–98.

50.   Simpson CD, Anyjwe K, Schimmer AD. Anoikis resistance and tumor metastasis. Cancer
Lett. 2008;272:177–85.

51.   Taddei ML, Giannoni E, Fiaschi T, Chiarugi P. Anoikis: An emerging hallmark in health
and diseases. J Pathol. 2012;226:380–93.

52.   Huang RY-J, Wong MK, Tan TZ, Kuay KT, Ng a HC, Chung VY, et al. An EMT

697   spectrum defines an anoikis-resistant and spheroidogenic intermediate mesenchymal state
698   that is sensitive to e-cadherin restoration by a src-kinase inhibitor, saracatinib (AZD0530).
699   Cell Death Dis. 2013;4:e915.

700   53.   Frisch SM, Schaller M, Cieply B. Mechanisms that link the oncogenic epithelial–
701   mesenchymal transition to suppression of anoikis. J Cell Sci [Internet]. 2013;126:21–9.
702   Available from: http://jcs.biologists.org/lookup/doi/10.1242/jcs.120907

703   54.   Xu S, Yang Y, Dong L, Qiu W, Yang L, Wang X, et al. Construction and characteristics
704   of an E-cadherin-related three-dimensional suspension growth model of ovarian cancer.
705   Sci Rep. 2014;4:5646.

706   55.   Cieply B, Riley IV P, Pifer PM, Widmeyer J, Addison JB, Ivanov A V., et al. Suppression
707   of the epithelial-mesenchymal transition by grainyhead-like-2. Cancer Res.
708   2012;72:2440–53.

709   56.   Yang J, Eddy J a, Pan Y, Hategan A, Tabus I, Wang Y, et al. Integrated proteomics and
710   genomics analysis reveals a novel mesenchymal to epithelial reverting transition in
711   leiomyosarcoma through regulation of slug. Mol Cell Proteomics. 2010;9:2405–13.

712   57.   Stadler M, Scherzer M, Walter S, Holzner S, Pudelko K, Riedl A, et al. Exclusion from
713   spheroid formation identifies loss of essential cell-cell adhesion molecules in colon cancer
714   cells. Sci Rep. 2018;8:1151.

715   58.   Riahi R, Sun J, Wang S, Long M, Zhang DD, Wong PK. Notch1-Dll4 signalling and
716   mechanical force regulate leader cell formation during collective cell migration. Nat
717   Commun. Nature Publishing Group; 2015;6:6656.

718   59.   Sprinzak D, Lakhanpal A, Lebon L, Santat LA, Fontes ME, Anderson GA, et al. Cis-
719   interactions between Notch and Delta generate mutually exclusive signalling states.
720   Nature. Nature Publishing Group; 2010;465:86–90.

721   60.   Boareto M, Jolly MK, Goldman A, Pietilä M, Mani SA, Sengupta S, et al. Notch-Jagged
722   signalling can give rise to clusters of cells exhibiting a hybrid epithelial/mesenchymal
723   phenotype. J R Soc Interface. 2016;13:20151106.

724   61.   Stylianou S, Clarke RB, Brennan K. Aberrant activation of Notch signaling in human
725   breast cancer. Cancer Res. 2006;66:1517–25.

726   62.   De Beco S, Perney JB, Coscoy S, Amblard F. Mechanosensitive adaptation of E-cadherin
727   turnover across adherens junctions. PLoS One. 2015;10:1–14.

728   63.   Le Duc Q, Shi Q, Blonk I, Sonnenberg A, Wang N, Leckband D, et al. Vinculin
729   potentiates E-cadherin mechanosensing and is recruited to actin-anchored sites within
730   adherens junctions in a myosin II-dependent manner. J Cell Biol. 2010;189:1107–15.

731   64.   Cai D, Chen SC, Prasad M, He L, Wang X, Choesmel-Cadamuro V, et al. Mechanical
732   feedback through E-cadherin promotes direction sensing during collective cell migration.
733   Cell. 2014;157:1146–59.

65. Kim SY, Hong SH, Basse PH, Wu C, Bartlett DL, Kwon YT, et al. Cancer Stem Cells
Protect Non-Stem Cells From Anoikis: Bystander Effects. J Cell Biochem.
2016;2301:2289–301.

66. Mateen S, Raina K, Agarwal C, Chan D, Agarwal R. Silibinin synergizes with histone
deacetylase and DNA methyltransferase inhibitors in upregulating E-cadherin expression
together with inhibition of migration and invasion of human non-small cell lung cancer
cells. J Pharmacol Exp Ther. 2013;345:206–14.

67. Kim NH, Kim SN, Kim YK. Involvement of HDAC1 in E-cadherin expression in prostate
cancer cells; its implication for cell motility and invasion. Biochem Biophys Res
Commun. 2011;404:915–21.

68. Choi SY, Kee HJ, Kurz T, Hansen FK, Ryu Y, Kim GR, et al. Class I HDACs specifically
regulate E-cadherin expression in human renal epithelial cells. J Cell Mol Med.
2016;20:2289–98.

69. Law ME, Corsino PE, Jahn SC, Davis BJ, Chen S, Patel B, et al. Glucocorticoids and
histone deacetylase inhibitors cooperate to block the invasiveness of basal-like breast
cancer cells through novel mechanisms. Oncogene. 2013;32:1316–29.

70. Foley JM, Scholten DJ, Monks NR, Cherba D, Monsma DJ, Davidson P, et al. Anoikis-
resistant subpopulations of human osteosarcoma display significant chemoresistance and
are sensitive to targeted epigenetic therapies predicted by expression profiling. J Transl
Med. 2015;13:110.

71. Jolly MK, Kulkarni P, Weninger K, Orban J, Levine H. Phenotypic plasticity, bet-
hedging, and androgen independence in prostate cancer: Role of non-genetic
heterogeneity. Front Oncol. 2018;8:50.

Figure Legends

Figure 1. E-cadherin upregulation is prognostic for improved outcomes in sarcoma. A-B.
Osteosarcomas with elevated E-cadherin have better metastasis-free survival (**A**) and overall
survival (**B**) as compared to tumors with low/no E-cadherin expression. **C-D.** Soft tissue
sarcomas (STS) from The Cancer Genome Atlas with higher E-cadherin mRNA (**C**) and protein
expression (**D**) have improved overall survival as compared to tumors with low or no E-cadherin.

Figure 2. Ectopic E-cadherin expression in sarcoma cells does not alter EMT. A. E-cadherin
expression significantly correlates with the EMT score of sarcoma tumors in The Cancer
Genome Atlas. **B.** Ectopic expression of E-cadherin in 143B human osteosarcoma cells has no
influence on mesenchymal markers (Snail, Slug, Twist, Zeb1, Vimentin). **C.** E-cadherin
expression has no effect on migration of 143B cells. Images (left) were collected every two hours

and quantified (right) using the IncuCyte Zoom system. **D.** E-cadherin expression does not
change invasion in 143B cells. **E.** Using mRNA expression of CDH1 (E-cadherin), empirical
probably density functions of EMT scores for CDH1-high (red) and CDH1-low (blue) TCGA
sarcoma sample were constructed by interpolation of the EMT score histogram (E<0.5,
0.5≤E/M≤1.5, M>1.5). EMT score distribution showed no significant difference when separated
for E-cadherin^{high} vs. E-cadherin^{low} expression from analysis of publicly-available sarcoma data
sets.
778
779 **Figure 3. E-cadherin inhibits anchorage-independent growth of sarcomas. A.** Anchorage-
780 independent growth of 143B cells expressing E-cadherin was significantly inhibited. **B-C.** E-
781 cadherin expression leads to reduces spheroid size in **B.** 143B and **C.** U2OS cells.
782
Figure 4. Ectopic expression of E-cadherin in sarcoma cells inhibits phospho-CREB levels.
A. A phospho-kinase array revealed that E-cadherin led to downregulation of phospho-CREB. **B-**
C. E-cadherin-mediated phospho-CREB inhibition was verified by **B.** ELISAs and **C.** western
blotting. **D.** Abrams canine osteosarcoma cells exhibited reduced phospho-CREB in E-cadherin
over-expressing cells. **E.** QRT-PCR confirmed knockdown of CREB with two independent
siRNAs. **F.** Western blotting to confirm knockdown of CREB in 143B cells. **G.** CREB
knockdown led to a modest downregulation of 143B colony growth in soft agar.
790
791 **Figure 5. TBX2 knockdown phenocopies E-cadherin-mediated CREB inhibition. A-**
792 **B.** TBX2 mRNA is downregulated in E-cadherin-expressing **A.** 143B cells and **B.** U2OS cells.
793 **C.** CREB knockdown with two independent siRNAs had no effect on TBX2 mRNA. **D-F.**
794 Conversely, TBX2 knockdown, verified in **D.** led to a significant reduction in CREB mRNA (**E**)
795 and protein level (**F**).
796
Figure 6. E-cad/TBX2 interplay mediates anchorage-independent growth. A. Intracellular
regulatory circuit modulating anchorage-independent growth. **B.** Mathematical model predicts
similar effects of Ecad-OE and TBX2-KD on anchorage-independent growth. **C.** TBX2
knockdown using two independent siRNAs inhibited anchorage-independent growth of 143B
cells. **D.** TBX2 downregulation did not affect sphere formation in the presence of ectopic E-
cadherin expression**. E.** TBX2 knockdown upregulates E-cadherin mRNA. **F.** Western blot
analysis of U2OS cells upon TBX2 knockdown indicates E-cadherin protein is not upregulated
by loss of TBX2. Prostate cancer (PC) cell line, LNCaP, was used as a positive control for E-
cadherin expression. **G.** The E-cadherin promoter is differentially methylated in sarcomas as
compared to carcinomas. **H.** Synovial sarcomas, which often display epithelioid
histopathological features, have the lowest levels of E-cadherin promoter methylation of all
sarcoma subtypes. **I.** Among soft tissue sarcoma histological subtypes E-cadherin mRNA
expression is highest in synovial sarcomas. Letters indicate statistically significant associations
between groups. Undifferentiated pleomorphic sarcoma (UPS), liposarcoma (LPS),

leiomyosarcoma (LMS), synovial sarcoma (SS), malignant peripheral nerve sheath tumors
(MPNSTs), and Myxofibrosarcoma (MFS).

Figure 7. E-cadherin inhibits spheroid formation by increased cell-cell adhesion. A. E-
cadherin causes tighter clustering of cells during spheroid formation. **B.** A mechanical model
illustrates the relationship between particle distance to adhesion. **C.** A mechanical model predicts
E-cadherin drives down spheroid size through an increase in cell-cell adhesion. **D-E.** As
predicted in the model, ectopic E-cadherin expression increases cell-cell adhesion in 143B (**D**)
and U2OS (**E**) cells.

Figures

Figure 1

Figure 2

864
865
866   Figure 3

 Figure 4

878   Figure 5
879

896   Figure 6

916   Figure 7