How a well-adapting immune system remembers

Andreas Mayer,[1,2] Vijay Balasubramanian,[3] Thierry Mora,[4,5] and Aleksandra M. Walczak[2,5]

[1]*Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA*
[2]*Laboratoire de physique théorique, CNRS, Sorbonne Université and École Normale Supérieure (PSL), 75005 Paris, France*
[3]*David Rittenhouse Laboratories, University of Pennsylvania, Philadelphia, PA 19104;*
and Initiative for the Theoretical Sciences, The Graduate Center,
The City University of New York, NY 10016, USA
[4]*Laboratoire de physique statistique, CNRS, Sorbonne Université,*
Université Paris-Diderot, and École normale supérieure (PSL), 75005 Paris, France
[5]*These authors contributed equally. Please send correspondence to* tmora@lps.ens.fr, awalczak@lpt.ens.fr
(Dated: June 14, 2018)

An adaptive agent predicting the future state of an environment must weigh trust in new observations against prior experiences. In this light, we propose a view of the adaptive immune system as a dynamic Bayesian machinery that updates its memory repertoire by balancing evidence from new pathogen encounters against past experience of infection to predict and prepare for future threats. This framework links the observed initial rapid increase of the memory pool early in life followed by a mid-life plateau to the ease of learning salient features of sparse environments. We also derive a modulated memory pool update rule in agreement with current vaccine response experiments. Our results suggest that pathogenic environments are sparse and that memory repertoires significantly decrease infection costs even with moderate sampling. The predicted optimal update scheme maps onto commonly considered competitive dynamics for antigen receptors.

I. INTRODUCTION

All living systems sense the environment, learn from the past, and adapt predictively to prepare for the future. Their task is challenging because environments change constantly, and it is impossible to sample them completely. Thus a key question is how much weight should be given to new observations versus accumulated past experience. Because evidence from the world is generally uncertain, it is convenient to cast this problem in the language of probabilistic inference where past experience is encapsulated in a prior probability distribution which is updated according to sampled evidence. This framework has been successfully used to understand aspects of cellular [1–4] and neural [5–8] sensing. Here, we propose that the dynamics of the adaptive immune repertoires of vertebrates can be similarly understood as a system for Bayesian sequential forecasting [9] of the pathogen environment.

The adaptive immune system relies on a diverse repertoire of B and T cell receptors to protect the host organism from a wide range of pathogens. These receptors are expressed on clones of receptor-carrying cells present in varying copy numbers. A defining feature of the adaptive immune system is its ability to change its clone composition throughout the lifetime of an individual, in particular via the formation of memory repertoires of B and T cells following pathogen encounters [10–15]. In detail, after a proliferation event that follows successful recognition of a foreign antigen, some elements of the newly expanded clone are flagged as memory cells [16]. These cells make up the memory repertoire compartment that is governed by its own homeostasis, separate from the inexperienced naive cells from which they came. Upon reinfection by a similar antigen, the memory repertoire guarantees a fast immune response. With time, our memory becomes specific to our history of infections, and adapted to the environments we live in. However, the commitment of part of the repertoire to maintaining memory must be balanced against the need to also provide broad protection from as yet unseen threats. What is more, memory will lose its usefulness over time as pathogens evolve to evade recognition.

How much benefit can immunological memory provide to an organism? How much memory should be kept to minimize harm from infections? How much should each pathogen encounter affect the distribution of receptor clones? To answer these questions we extend a framework for predicting optimal repertoires given pathogen statistics [17] by explicitly considering the inference of pathogen frequencies as a Bayesian forecasting problem [9]. We derive the optimal repertoire dynamics in a temporally varying environment and analyze its features, linking the amount of memory production to the variability of environment. We also show how biologically realistic population dynamics can approximate the optimal inference process, and analyze conditions under which memory provides a benefit. Comparing predictions of our theory to experiment, we argue for a view in which the adaptive immune system can be interpreted as a dynamical framework for learning a highly sparse distribution of antigens.

Learning the antigenic environment to form memory is equivalent to estimating the frequencies of the pathogens that surround us, and using this information to construct an adapted conjugate response. This frequency estimation problem is encountered generally throughout science, e.g., in situations where we must estimate the number of species (macroscopic or microbial) in different ecological niches. Our results suggest a dynamical fre-

quency estimation procedure that will be useful in these other settings.

II. THEORY OF OPTIMAL IMMUNE PREDICTION

At any given time, the space of possible pathogens is huge and unknown. We consider an antigenic space of K different pathogens with time-varying frequencies $\boldsymbol{Q}(t) = (Q_1(t), ..., Q_K(t))$. These frequencies are unknown to the organism, and evolve stochastically. Their dynamics is formally described by a continuous evolution encoding the rates of jumping from one state of the pathogenic frequencies to another (Fig. 1C). For concreteness we consider a classical neutral population dynamics with genetic drift due to demographic noise, and immigration from a reservoir (Eq. 9 in Methods).

The pathogenic environment is enormous and the immune system can only sample it sparsely, as pathogens enter into contact with it at some rate λ. The probability of encountering each pathogen is determined by $\boldsymbol{Q}(t)$. We reason that the immune system should efficiently use the information available through these encounters, along with prior knowledge of how pathogens evolve encoded in the system dynamics, to build an internal representation of the environment (Fig. 1D). Based on this representation of the world, the immune system should organize its defenses to minimize harm from future infections (Fig. 1E). The actual dynamics of the immune system manifests several notable features: (1) Immune memory increases by orders of magnitude after a first encounter with a pathogen, leading to substantial decreases in serious infections in early life (Fig. 1A), (2) The magnitude of memory increase is much lower after subsequent encounters, (3) The memory pool undergoes attrition over time, and (4) There is a mid-life plateau in the fraction of the immune repertoire occupied by memory (Fig. 1B).

The internal representation of the environment can be regarded as a system of beliefs, or guesses, about pathogen frequencies. Formally, these beliefs can be represented in the form of a posterior probability distribution function $B(\boldsymbol{Q}, t)$ over pathogen frequencies, which the host implicitly computes using all the information it has garnered over time. Optimally, these beliefs are computed by the rules of Bayesian statistics, by combining the memory of past encounters with knowledge of the stochastic rules under which the pathogenic environment evolves (Methods). Upon encountering a pathogen a at time t, the Bayesian belief distribution jumps from a pre-encounter value $B(\boldsymbol{Q}, t^-)$ to a post-encounter belief $B(\boldsymbol{Q}, t^+)$ according to:

$$B(\boldsymbol{Q}, t^+) = \frac{Q_a\, B(\boldsymbol{Q}, t^-)}{\int \mathrm{d}\boldsymbol{Q}'\, Q_a'\, B(\boldsymbol{Q}', t^-)}. \tag{1}$$

Between encounters, beliefs simply follow an evolution operator $\mathcal{A}$ which encodes the dynamics of pathogen frequency changes:

$$\frac{\mathrm{d}B(\boldsymbol{Q}, t)}{\mathrm{d}t} = \mathcal{A}B(\boldsymbol{Q}, t). \tag{2}$$

This prediction step, which is performed in the absence of any new information, relies on the immune system implicitly "knowing" the probability laws governing the stochastic evolution of the environment—but not, of course, the actual path that it takes. Optimally, the belief distribution should be initialized at birth to reflect the steady state of the dynamics, $B(\boldsymbol{Q}, 0) = \rho_s(Q)$, where $\mathcal{A}\rho_s = 0$.

How could the immune system leverage a representation of beliefs about pathogen frequencies to provide effective immunity? Each lymphocyte of the adaptive immune system expresses on its surface a single receptor r out of L possible receptors. This receptor endows the lymphocyte with the ability to specifically recognize pathogens (labeled a) with probability $f_{a,r}$. The immune repertoire is defined by frequencies of these receptors across the lymphocyte population, denoted by $\boldsymbol{P} = (P_1, ...P_L)$. These frequencies sum up to one, which implies a resource allocation trade-off between the different receptor types – having more of one in the repertoire implies having less of others. How much harm an infection is expected to inflict depends on the *coverage*, i.e. the fraction $\tilde{P}_a(t) = \sum_a f_{a,r} P_r$ of the repertoire specific to antigen a. Given the pathogen frequencies $\boldsymbol{Q}(t)$ and repertoire distribution $\boldsymbol{P}(t)$, the mean harm cause by the next infection is given by $\sum_a Q_a \cdot c(\tilde{P}_a)$, where c is decreasing function of the fraction of the repertoire specific to the infection [17]. The host organism does not know $\boldsymbol{Q}$ with certainty, but has an internal belief $B(\boldsymbol{Q}, t)$ about the frequencies learned through sampling during previous infections. An optimal immune system can then distribute its resources to minimize the expected harm of future of infections:

$$\boldsymbol{P}^\star(t) = \underset{\boldsymbol{P}}{\mathrm{argmin}} \sum_a \hat{Q}_a(t) \cdot c(\tilde{P}_a) \equiv G(\hat{\boldsymbol{Q}}(t)), \tag{3}$$

where $\hat{\boldsymbol{Q}}(t) \equiv \langle \boldsymbol{Q} \rangle_{B(\boldsymbol{Q},t)}$ are the expected frequencies of pathogens. Although the function G may be complicated, it generally implies that receptors that are specific to frequent infections (high $\hat{Q}_a$) should be well represented in the optimal repertoire (high $\tilde{P}_a^\star$) ([17] and Methods). In this framework we have assumed that infections, their clearing by the immune system, and the subsequent update of the repertoire, are all fast compared to changes in the environment, which occur over a time τ, and to the mean time between pathogenic encounters (λ^{-1}).

We are proposing an interpretive framework for understanding adaptive immunity as a scheme for sequential inference. This view provides two key insights. First, it suggests how new experience should be balanced against previous memory and against unknown threats in order for adaptive immunity to work well. Second, it suggests

FIG. 1: Sketch of a model of immune repertoire dynamics as a sequential inference process about a time-varying pathogen distribution. (A) Infectious disease hospitalization rates and (B) median percentage of T cells showing memory markers as a function of age. (C) The organism lives in a pathogenic environment with frequencies of different pathogen strains that change over time. (D) Past pathogen encounters provide an avenue for the immune system to learn the pathogen distribution. Using sequential Bayesian inference provides an optimal way to update the beliefs about the frequencies of different pathogens over time. (E) Bases on its beliefs about the prevalence of pathogens the optimal immune dynamics allocates lymphocytes across different pathogens to minimizes the expected harm from infections. Broadly, the more frequent a pathogen is the more the organism should be covered. This resource allocation maps the changes in beliefs to the changes in the repertoire composition. Data on infectious disease hospitalization rate from [18], T cell data from [19] and [20].

a particular dynamics of implicit belief updates that can globally reorganize the immune repertoire to minimize harm from the pathogenic environment. In Results we will show a plausible strategy that the immune system could follow to approximate optimal Bayesian inference, and we will compare the resulting dynamics with specific features of the adaptive immune system.

III. RESULTS

A. A lymphocyte dynamics for approximating optimal sequential inference

For specificity, we assume neutral stochastic evolution of pathogens driven by genetic drift happening on a characteristic timescale τ and immigration from an external pool with immigration parameters $\boldsymbol{\theta} = (\theta_1, \ldots, \theta_K)$ (Eq. 9 in Methods). In this case, we show that optimal Bayesian belief update dynamics can be approximated by maintaining a memory of an effective count of previous encounters $\boldsymbol{n} = (n_1, \ldots, n_K)$, initialized to the immigration rates $\boldsymbol{n}(0) = \boldsymbol{\theta}$, and subject to the update rules (see

SI Text A 2):

$$n_a(t^+) = n_a(t^-) + 1 \quad \text{upon encountering } a \qquad (4)$$

$$\tau \frac{\mathrm{d}\boldsymbol{n}}{\mathrm{d}t} = -\frac{1}{2}\left(|\boldsymbol{n}| - 1\right)\left(\boldsymbol{n} - \boldsymbol{\theta}\right), \qquad (5)$$

where $|\boldsymbol{n}| = \sum_{a'} n_{a'}$. The expected frequency of each pathogen (used in Eq. 3) is estimated from these counts as:

$$\hat{Q}_a(t) \approx \frac{n_a}{|\boldsymbol{n}|}, \qquad (6)$$

We checked the accuracy of this approximation explicitly by comparing it to an exact solution computed by spectrally expanding the generator of the stochastic dynamics (see SI Text A 3 and Fig. S1). Intuitively, in the first equation (4), encountering a pathogen a increases its count by one. In the second equation (5), old evidence is discarded through relaxation dynamics. Encounters are gradually forgotten, which is beneficial in a changing environment where old information becomes obsolete. In the extreme case where the relaxation goes undisturbed by further pathogenic encounters, Eq. 5 converges to the normalized immigration rates $\boldsymbol{\theta}$, consistent with their interpretation as the relative frequencies of pathogenic epitopes learned over evolutionary timescales.

An optimal immune system should should map the counts above into a receptor repertoire $\boldsymbol{P}^\star$ as in Eq. 3.

The repertoire then follows a memoryless dynamics derived from Eqs. 4-5 (see SI Text B). In this way, the repertoire itself is a representation of the host's belief about the structure of the pathogenic environment inferred from the history of encounters. To understand this, consider a cost function $c(\tilde{P}_a) = -\log \tilde{P}_a$ and uniquely specific receptors, $f_{a,r} = \delta_{a,r}$. In this case the mapping Eq. 3 is the identity, i.e. $\boldsymbol{P}^\star(t) = \hat{\boldsymbol{Q}}(t)$ ([17] and Methods). Then the optimal repertoire dynamics can be achieved simply by having clone sizes of different receptors follow Eq. 4,5 up to some scaling. More generally the optimal repertoire is some non-linear mapping of the encounter counts, but only requires information that can be represented in population sizes of different clones, which are quantities regulated by the actual biological dynamics.

The implied dynamics of the optimal repertoire qualitatively mirrors adaptive immune system function where past experiences are encoded by memory cells guaranteeing higher protection after an infection. In our framework, memory is created by the change in $\boldsymbol{P}^\star$ that follows from applying the resource allocation rule in Eq. 3 to the incremented count following an encounter with a pathogen. In this way, the frequency of pathogens is learned by the immune system through encounters, leading to increase in the part of the repertoire that is specific to common infections (Fig. 1E), in parallel with actual immune function.

Our theory suggests that there is an optimal timescale for forgetting about old infections which is set by the duration over which the environment varies. Given the long timescales over which many relevant pathogens change, this view predicts that memory should generally be long-lived (with the timescale of decay being of the order of years or decades). Indeed, despite the relatively short life span of memory cells [21], constant balanced turnover keeps elevated levels of protection relatively stable for decades after an infection, even in the absence of persistent antigens [22–24]. Yet, the environment is not constant and dynamic changes of the repertoire are necessary to follow its changes. Over longer timescales, and in the absence of re-infection, the repertoire in our theory converges back towards its broadly distributed initial state $\boldsymbol{P}^\star(0)$ (Eq. 5; Fig. 1E, red lines). This relaxation in the optimal dynamics allows continued learning of the changing environment. Biologically, immune memory diminishes over long times despite its active maintenance. Our theory proposes that such attrition of memory might be adaptive and that there exists an optimal timescale of attrition, which is determined by the time over which pathogen frequencies are correlated.

It is natural to interpret the primordial repertoire, $\boldsymbol{P}^\star(0)$, as the naive repertoire. In our Bayesian theory, it is a function of the long-term expected prevalence of different antigenic epitopes in the evolving pathogen population. This is captured effectively in the $\boldsymbol{\theta}$ parameters, $\boldsymbol{P}^\star(0) = G(\hat{\boldsymbol{Q}}) = G(\boldsymbol{\theta}/|\boldsymbol{\theta}|)$ (Methods). Much like the production of new naive cells by the thymus and bone marrow throughout life, $\boldsymbol{\theta}$ acts a source term in Eq. 5, steering the repertoire towards $\boldsymbol{P}^\star(0)$. The primordial repertoire can be inhomogeneous, depending on prior expectations about the pathogenic landscape, $\boldsymbol{\theta}/|\boldsymbol{\theta}|$. This bias implies higher levels of pre-existing immunity for pathogenic determinants that occur more frequently over long evolutionary timescales. In the real adaptive immune system, the recombination machinery and initial selection mechanisms that generate immune receptors have large biases [?], which tilt the naive repertoire towards certain regions of the antigenic space. In our interpretation, this bias should be towards pathogens, or rather pathogenic epitopes, that are more prevalent or harmful.

B. Learnability of pathogen distribution implies a sparse pathogenic landscape

The immune system must be prepared to protect us not just from one pathogen but a whole distribution of them. Even restricting recognition to short peptides and accounting for cross-reactivity [25], estimates based on precursor frequencies for common viruses give an effective antigen environment of size $K \sim 10^5$–10^7 [26]. How can the immune system learn anything useful about such a high dimensional distribution from a limited number of pathogenic encounters? Naively, one might expect that the number of samples needed to learn the distribution of pathogens must be larger than of the number of pathogens, i.e., $\lambda t \sim K$, where t is the time over which learning takes place. Although little is known about the receptor-antigen encounter rate λ, this estimate suggests that the pathogenic environment is not easily learnable and therefore memory has limited utility

This apparent paradox can be resolved by the fact that the pathogenic environment may be sparse, meaning that only a small fraction of the possible pathogens are present at any given time. In the neutral theory, this sparsity is controlled by the parameter $\boldsymbol{\theta}$. In the scenario that we are considering, typical pathogen landscapes $\boldsymbol{Q}$ are drawn from the steady state distribution $\rho_s(\boldsymbol{Q})$ of the neutral immigration-drift dynamics, which is a Dirichlet distribution parametrized by $\boldsymbol{\theta}$ (Eq.A1 in SI Text A). When θ_a is small, the distribution is peaked at $Q_a = 0$, meaning that pathogen a is absent a majority of the time. For instance, for uniform $\theta_a \equiv \theta \ll 1$, the effective number of pathogens present at any given time is $K\theta$ (see SI Text C 3). Since the system only needs to learn about the pathogens that are present, the condition for efficient learning should naively be $\lambda t \sim K\theta$, which is much easier to achieve realistically for small θ.

Our theory can be used to quantify the benefit of memory as a function of the different immunological parameters. We compute the optimized cost function $c(t) = \sum_a Q_a(t)c(\tilde{P}_a^\star(t))$, and study how it decreases as a function of age, t, relative to the cost at birth $c_0 = c(t = 0)$, as the organism learns from pathogen

FIG. 2: Advantage of immunological memory depends on sufficient sampling. The mean expected cost of an infection in a changing environment is a function of both the age of the organism t, the timescale τ on which the environment changes, and the sparsity $1/\theta$ of the environment. (A) Relative cost as a function of age for environments with different sparsity (for fixed $\lambda\tau = 10^5$, $K = 2000$). (B) Relative cost as a function of age for environments changing more or less rapidly (for fixed $\theta = 0.02$, $K = 2000$). (C) Collapse of the data by plotting the relative cost gap against the number of samples per effective dimension. Simulations with logarithmically spaced parameters in the ranges λt from 1 to 5000, $\lambda\tau$ from 1 to 1000000, K from 100 to 2000, θ from 0.01 to 0.2. Cost is $c(\tilde{P}_a) = 1/\tilde{P}_a$ and receptors are assumed to be uniquely specific, $f_{a,r} = \delta_{a,r}$.

encounters. This relative cost also depends on the encounter rate λ, the size of the pathogenic space K, the sparsity of pathogenic space θ and timescale of change in the environment τ. Fig. 2A shows that the benefit of memory increases with pathogen sparsity – when θ is small, even a few encounters suffice to seed enough memory to reduce the cost of future infections. The cost saturates with age to a value c_∞, either because memory approaches optimality, or because memory eventually gets discarded and renewed as the environment changes. Fast changing environments lead to an earlier and higher saturation of the cost with age (Fig.2B) since learning and prediction are limited by decorrelation of the environment.

Analytical arguments show that in the limit of few samples the relative cost c/c_0 achievable in a static environment scales as $\lambda t/K\theta$ (SI Text C). In general, we find that the cost is a function of $\lambda t_e/K\theta$, where the effective time t_e is defined via $\lambda t_e = |\boldsymbol{n}(t)| - |\boldsymbol{n}(0)| \approx \sqrt{2\lambda\tau}\tanh(\lambda t/\sqrt{2\lambda\tau})$ with $\boldsymbol{n}(t)$ being the vector of the encounter counts discussed above (see SI Text A 4 for derivation from Eqs. 4-5). Plotted in terms of this variable the relative cost gap as function of time, $(c - c_\infty)/(c_0 - c_\infty)$ collapses onto a single curve for all parameter choices (Fig. 2C). Fig. 2C shows that the cost drops by a factor of ~ 2, when $\lambda t_e/K\theta \sim 1$. Thus, there is a substantial benefit to memory already when the effective number of encounters is comparable to the effective number of pathogens. At young ages (small t) or with slowly changing environments (large τ), $t_e \approx t$ and so this condition is simply $\lambda t \sim K\theta$, i.e. the total number of encounters should be comparable to effective number of pathogens that are present.

C. Memory production in sparse environments should be large and decrease with prior exposure

The theory can be used to make quantitative and testable predictions about the change in the level of protection that should follow a pathogen encounter. Consider an infection cost function that depends as a power law on the coverage, $c(\tilde{P}_a) = 1/\tilde{P}_a^\alpha$, with a *cost exponent* α that sets how much attention the immune system should pay to recognizing rare threats. (Below we will use the shorthand $\alpha = 0$ to indicate logarithmic cost.) Cost functions of this form can be motivated by considering the time to recognition of an exponentially growing antigen population by the immune system [17], or, alternatively, by considering the time delay of the expansion of the precursor cells to some fixed number of effector cells (SI Text D).

In the simplest model for repertoire updates recognition of pathogens leads to proliferation proportionally to the number of specific precursor cells, followed by a homeostatic decrease of the memory pool [14, 31]. Thus the fold-change $\tilde{P}_a(t^+)/\tilde{P}_a(t^-) = $ const where t^-, t^+ are times just before and after the encounter. By contrast, our Bayesian theory predicts that the fold change upon encountering pathogen a should be

$$\frac{\tilde{P}_a^\star(t^+)}{\tilde{P}_a^\star(t^-)} = \left(1 + \kappa/\tilde{P}_a^\star(t^-)^{(1+\alpha)}\right)^{1/(1+\alpha)}, \qquad (7)$$

where κ depends on prior expectations about the antigenic environment and previous pathogen encounters (derivation and details in SI Text B 1). Setting $\alpha = 0$ gives the result for a logarithmic cost function.

To understand this prediction first consider the effect of a primary infection on a naive repertoire, $\theta_a \equiv \theta$,

FIG. 3: Changes in protection levels upon infections for cost functions $c(\tilde{P}_a) = 1/\tilde{P}_a^{\alpha}$ ($\alpha = 0$ indicates logarithmic cost). (A) The prediction of the Bayesian model (Eq. 7 for different α) closely fits experimental data on antibody titers. Data on pre- and post-vaccination antibody titers against stem and head hemaglutinin epitopes following a booster vaccination with inactivated H5N1 in humans from [27]. (B) Optimal fold change of coverage for $n_i(t^-) = \theta$ for all i. The fold change increases with the sparsity of the environment controlled by parameter θ. (C) Absolute change in coverage for primary infection and reinfections (for $\alpha = 1$) normalized such that the change for the primary infection is 1 and neglecting attrition.

$\tilde{P}^{\star}(0) = 1/K$, and $|\boldsymbol{n}(0)| = K\theta$ where the receptors are uniquely specific ($f_{a,r} = \delta_{a,r}$). In this case $\kappa = 1/K^{1+\alpha}\theta$ (see SI Text B1) and Eq. 7 predicts a fold-change of $(1 + 1/\theta)^{1/(1+\alpha)}$. Thus, in a sparse environment our theory predicts a very large memory production upon an antigenic encounter with a naive immune system. Experimentally, memory production typically leads to the proliferation of antigen-specific cells by a factor of 100-1000-fold [16], in line with the prediction from the theory. Such a large increase in protection upon an encounter implies a sparsity parameter $\theta \sim 10^{-6}$–10^{-4} (here taking $\alpha = 1$ for definiteness) (Fig. 3B). Combined with the estimate $K \sim 10^5$–10^7 [26] this suggests that the effective number of pathogens at any given time ranges from $K\theta = 0.1$ to $1,000$.

To test Eq. 7 on immunological data, we fit the Bayesian update model to experiments reporting fold-changes in antigen titers upon booster vaccinations against influenza from [27] (Fig. 3A) using least-squares. Titers correspond to the concentration of antibodies that are specific to the antigen a, and can thus be viewed as an experimental estimate of $\tilde{P}_a$. The optimal Bayesian strategy explains the data, accounting for the larger boosting at small prevaccination titers and showing no increase for large titers, while the proportional model predicts constant boosting for all titers.

A number of further studies have observed a negative correlation between fold-changes and pre-existing immunity for both B cells [28] and T cells [29]. Mechanistic models have been proposed to explain how the population dynamics of expanding lymphocytes might give rise to such a negative correlation [30–32]. Our Bayesian theory provides an adaptive explanation for this effect.

D.   Long-term dynamics of a well-adapting repertoire

Our model makes predictions for the dynamics of growth and attrition of memory over time, with consequences for immunity and for the diversity of the immune repertoire. We quantify the dynamics in terms of a *memory fraction* defined as a sum of the coverage fractions $\tilde{P}_{a_i}$ over all previously encountered pathogens $\{a_i\}$. The memory fraction measures the size of memory relative to size of the whole immune repertoire. Early in life every infection is new and even modest increases in the memory fraction lead to large drops in infection susceptibility (measured by the expected cost of new infections in Fig. 4A). At the same time, the memory fraction increases rapidly (Fig. 4B), but the growth of memory slows as subsequent infections lead to less memory production following the optimal fold-change rule in Eq. 7, and as attrition begins to play a role. The fraction of the repertoire devoted to memory in mid-life is largely determined by how the cost of infections scales with coverage. The observed memory fraction of $\sim 50\%$ at mid-life suggests a cost exponent of $\alpha \approx 0.5$ (Fig. 4B). The diversity of the memory repertoire increases with time at a rate that slows with age (quantified in Fig. 4C by richness, which measures the number of unique specificities, and the Shannon entropy of the repertoire frequency distribution.)

To gain insight into these dynamics of our model we average the stochastic equations over the statistics of pathogen encounters. We show in SI Text B2 that this mean-field approximation yields a differential equation for the population fraction of different clones with two opposing contributions which balance alignment of the immune repertoire with the current pathogenic environment (i.e. memory production) against alignment with the long-term mean environment (i.e. attrition). Interestingly, the mean-field equation broadly coincides with dynamics that were proposed in [17] to self-organize an optimal immune repertoire. The essential difference here is that the time-scale of learning slows down with increasing experience following the rules of optimal sequential update in Eq. 7.

FIG. 4: (A,D) Relative cost, (B,E) memory cell fraction, and (C) memory diversity as a function of age. (A-C) shows long term dynamics for a repertoire following optimal Bayesian update dynamics for three different α. Memory diversity is plotted as richness, i.e. the number of unique memory specificities, as well as the exponential of Shannon entropy S of the memory compartement defined for a probability distribution p_i with $\sum_i p_i = 1$ as $S = -\sum p_i \log p_i$. (D-E) compares the optimal Bayesian dynamics with a constant fold change update by a factor 30, and the same proportional update but with a cap to 10^5 of the total fold expansion for any clone. (F) Comparison of the cost as a function of age for the optimal dynamics (solid lines) and a dynamics without attrition (dotted lines) for two environmental correlation time scales $\tau_c = 2\tau/K\theta$ (see SI Text Eq. **??**). Parameters: encounter rate $\lambda = 40$/year, antigen space dimensionality $K = 10^5$, antigen sparsity $\theta = 2.5 \cdot 10^{-4}$, and for panel A-E $\tau \to \infty$. To reduce fluctuations the statistics were averaged over multiple runs of the dynamics.

We then asked which features of the proposed repertoire dynamics are most relevant to ensure its effectiveness. How important is the negative correlation between fold expansion and prior immune levels, and how important is attrition? Furthermore, if the immune system follows Bayesian dynamics it must have integrated on an evolutionary time scale a prior about composition and evolution of the pathogen environment through the parameters $\boldsymbol{\theta}$ and τ – however, the prior may be inaccurate. How robust is the benefit of memory to imperfections of the host's prior assumptions about pathogen evolution? To answer these questions we compare the long term immune repertoire dynamics using the optimal Bayesian scheme to other simplified schemes. We find that a constant fold expansion dynamics quickly leads to very suboptimal repertoire compositions (Fig. 4D, pink line), since the exponential amplification of cells specific to recurrent threats quickly leads to a very large fraction of the repertoire consisting of memory of those pathogens (Fig. 4E, pink line). This sub-optimality persists even if we assume that some global regulation caps the constant fold expansion such that no individual receptor clone can take over all of the repertoire (Fig. 4D,E grey line). Thus,

negative feedback in T cell expansion to individual antigens is very important to maintain a properly balanced diverse repertoire. In contrast, within a dynamics with a negative correlation, the precise levels of updating do not need to be finely tuned to the environmental statistics: varying the assumed sparsity of the pathogen distribution, which controls fold expansion upon primary infection in the optimal dynamics, leads to a relatively modest deterioration of the convergence speed of the learning process (Fig. S3A) and does not matter asymptotically (Fig. S3B). Attrition does not matter at young age, but can play an important role for long-term adaptation to relatively rapidly changing pathogen distributions (Fig. 4F). However, the attrition time scale need not be finely tuned to get close to optimal dynamics (Fig. S4).

E. Adapting a cross-reactive repertoire

Above, we described adaptation of immune repertoires in terms of changes in the effective coverage $P_a = \sum_r f_{a,r} P_r$, where the cross-reactivity matrix $F = (f_{a,r})$ reflects the ability of each receptor to recognize many

antigens, and also the propensity of each antigen to bind to many receptors [25]. Because of cross-reactivity, each pathogen encounter should result in the expansion of not just one but potentially many receptor clones. Here we ask how the optimal immune response is distributed among clones with different affinities.

Following Perelson and Oster [33], we will represent the interaction of receptors and antigens by embedding both in a high-dimensional metric recognition "shape space", where receptors are points surrounded by recognition balls. Antigens that fall within a ball's radius will be recognized by the corresponding receptor. In this presentation a and r are the coordinates of antigens and receptors respectively and their recognition propensity depends on their distance, $f_{a,r} = f(|a - r|)$.

Earlier sections have already discussed the optimal dynamics of the coverage $\tilde{P}_a$, which is a convolution of the cross-reactivity matrix with the receptor clone distribution P_r. Thus, the optimal dynamics of the clone distribution can be derived by deconvolving the cross-reactivity subject to the constraint that P_r cannot be negative. Carrying out this analysis in SI Text B 3, reveals a general qualitative phenomenon – competitive exclusion between clones expressed in the repertoire and their close neighbors within the cross-reactivity radius (Fig. S5, blue line). This exclusion is not an assumption of the model, but rather stems from the optimal Bayesian theory. Given a receptor clone that covers one region of antigenic shape space, the global likelihood of detecting infections increases by placing other clones to cover other regions. This can be shown analytically when cross-reactivity is limited, memory updates are small in magnitude, and the pathogen distribution is assumed to be uncorrelated (SI Text B 3).

In general, the frequencies of pathogens might be correlated in antigenic space, for example because mutations from a dominant strain give rise to new neighboring strains. An optimally adapting immune system should incorporate such correlations as a prior probability favoring smoothness of the pathogen distribution. Such priors work their way through the optimal belief update scheme that we have described, and weaken the competitive exclusion between clones with overlapping cross-reactivity (Fig. S5, orange line).

In general, when cross-reactivity is wide or the required clone fraction update is large, numerical analysis shows that achieving optimally predictive immunity after a pathogen encounter requires a global reorganization of the entire repertoire (Fig. S6, blue line). There is no plausible mechanism for such a large scale reorganization since it would involve up- and down-regulation even of unspecific clones. However, in SI Text B 3 we show that the optimal update can be well-approximated by changes just to the populations of specific clones with pathogen binding propensities $f_{a,r}$ that exceed a threshold. The optimizing dynamics with this constraint exhibits strong competitive exclusion, where only the highest affinity clones proliferate, while nearby clones with lower affinity are depleted from the repertoire (Fig. S6, orange line). The constrained optimum reaches a cost within 1 percent of the global optimum. Thus, reorganization of pathogen-specific receptor clone populations following an infection, as seen in vertebrates, can suffice to achieve near-optimal predictive adaptation of the immune repertoire.

IV. DISCUSSION

The adaptive immune system has long been viewed as a system for learning the pathogenic environment [11]. We developed a mathematical framework in which this notion can be made precise. We proposed a general scheme for inferring the frequencies of pathogens undergoing an immigration-drift dynamics and showed how such inference might be performed by the population dynamics of lymphocyte clones. We also argued that the antigenic environment must be effectively sparse to be learnable with a realistic rate of pathogen encounters. The optimal repertoire dynamics in sparse antigenic environments naturally produces a number of known properties of the adaptive immune system including a large memory production in naive individuals, a negative correlation of memory production with preexisting immune levels, and a sublinear scaling with age of the fraction of the repertoire taken up by memory of past infections.

Our framework is easily extended to incorporate further aspects of pathogen evolution, e.g. mutational dynamics in antigenic space. Such dynamics will lead to correlations in the pathogen distribution which we showed will influence the structure of the optimal conjugate repertoire. In particular, the optimal response should spread around the currently dominant antigens to also provide protection against potential future mutations. Hypermutations in B cells may play a role in this diversification, in addition to their known function of generating receptors with increased affinity for antigens of current interest. It would also be interesting to extend our framework to other immune defense mechanisms, including innate immunity, where the role of memory has received recent attention [35].

Although our study was motivated by the adaptive immune system, many of our results extend to other statistical inference problems. In effect, we have derived an exactly computable solution to the stochastic filtering problem for a categorical distribution evolving according to a Wright-Fisher diffusion process. Except for special cases such as the well-known Kalman filter, which exactly solves the stochastic filtering problem for continuous Gaussian-distributed latent and measurement variables, stochastic filtering mostly relies on numerical approximations [9]. Our analytical results might therefore be of use in other contexts where changing distributions must be inferred from samples at different time points, e.g., in population genetics. Additionally, we have derived the convergence rate for Bayesian inference of cate-

gorical distributions in high dimensions in the undersampled regime, showing that effectively sparse distributions can be inferred much more quickly. These results add to the growing literature on high-dimensional inference from few samples [36, 37], which has arisen in the context of the big data revolution.

We propose that the adaptive immune system balances integration of new evidence against prior knowledge, while discounting previous observations to account for environmental change. Similar frameworks have been developed for other biological systems. In neuroscience, leaky integration of cues has been proposed as an adaptive mechanism to discount old observations in change-point detection tasks [38?], and close-to-optimal accumulation and discounting of evidence has been reported in a behavioral study of rat decision-making in dynamic environments [40]. Inference from temporally sparse sampling has been considered in the framework of infotaxis, which is relevant for olfactory navigation [41]. In the context of immunity, related ideas about inference and prediction of pathogen dynamics have been used to predict flu strain and cancer neoantigen evolution in silico [42, 43]. Finally, ideas similar to those developed here could be used in ecology or microbiome studies to reconstruct evolutionary or ecological trajectories of population dynamics from incomplete sampling of data at a finite number of time points, e.g., from animal sightings or metagenomics.

V. METHODS

A. Modeling pathogen dynamics by a immigration-drift process

In our model we describe the stochastic dynamics of the pathogenic environment (Fig. 1B) by a Fokker-Planck equation for the conditional probability distribution $\rho(\boldsymbol{Q}, t)$

$$\frac{\partial \rho(\boldsymbol{Q}, t)}{\partial t} = \mathcal{A}\rho(\boldsymbol{Q}, t), \tag{8}$$

where $\mathcal{A}$ is a differential operator acting on ρ that controls the dynamics. For concreteness, we consider a population that changes due to genetic drift and immigration from an external reservoir, which we describe by a Wright-Fisher diffusion equation [44, 46]

$$\tau \frac{\partial \rho(\boldsymbol{Q}, t)}{\partial t} = -\frac{1}{2} \sum_a \frac{\partial}{\partial Q_a} \left[(\theta_a - |\boldsymbol{\theta}| Q_a)\rho(\boldsymbol{Q}, t) \right] \\ + \frac{1}{2} \sum_{a,b} \frac{\partial^2}{\partial Q_a \partial Q_b} \left[Q_a(\delta_{a,b} - Q_b)\rho(\boldsymbol{Q}, t) \right], \tag{9}$$

where τ sets the time scale of dynamics, $\boldsymbol{\theta}$ is a K-dimensional vector of immigration rates, and $\delta_{a,b}$ is the Kronecker delta, which is 1 if $a = b$ and 0 otherwise. Here and in the following we denote the norm of a vector $\boldsymbol{x}$ by $|\boldsymbol{x}| = \sum_i x_i$. To efficiently simulate trajectories according to this dynamics we sample the new distribution of frequencies directly from the transition density of the stochastic process as described in App. A 5. This dynamics retains key features of real pathogen environments. First, at a given point in time the environment contains many different pathogens with different frequencies determined by genetic drift and immigration. Second, the dominant pathogens change over time, such as is the case for many viruses e.g. the flu or HIV.

B. Minimizing the cost of infection

To solve the optimization problem Eq. 3 analytically a set of necessary conditions for optimality, the so-called Karush-Kuhn-Tucker conditions, can be derived. When all receptors are present at a non-zero frequency in the optimal repertoire $P_r^\star > 0$ these conditions imply [17]

$$\left. \frac{\partial \sum_a Q_a c(\tilde{P}_a)}{\partial P_r} \right|_{\boldsymbol{P}^\star} = -\lambda^\star, \tag{10}$$

where $\lambda^\star$ is set by the condition $\sum_r P_r^\star = 1$. If we further simplify the problem by assuming that there is no cross-reactivity between different pathogens and by considering power-law cost functions then this simplifies to the explicit solution

$$\tilde{P}_a^\star = \frac{1}{Z} Q_a^{1/(1+\alpha)}, \tag{11}$$

where Z is a normalization constant. Other cases are discussed in detail in [17] including how to solve the optimization problem numerically using a projected gradient algorithm in the general case.

Acknowledgements. The work was supported by grant ERCStG n. 306312, Simons MMLS grant 400425, and NSF grant PHY-1734030. Work on this project at the Aspen Center for Physics was supported by NSF grant PHY-1607611.

[1] T. J. Perkins and P. S. Swain, Molecular systems biology **5**, 326 (2009).

[2] T. J. Kobayashi, Physical Review Letters **104**, 1 (2010).

[3] E. D. Siggia and M. Vergassola, Proceedings of the Na-

tional Academy of Sciences pp. 3704–3712 (2013).

[4] D. A. Sivak and M. Thomson, PLoS Computational Biology **10** (2014).

[5] M. R. DeWeese and A. M. Zador, Neural Computation **10**, 1179 (1998).

[6] J. M. Beck, W. J. Ma, R. Kiani, T. Hanks, A. K. Churchland, J. Roitman, M. N. Shadlen, P. E. Latham, and A. Pouget, Neuron **60**, 1142 (2008).

[7] S. Deneve, Neural computation **20**, 91 (2008).

[8] B. Wark, A. Fairhall, and F. Rieke, Neuron **61**, 750 (2009).

[9] Z. H. E. Chen, Statistics **182**, 1 (2003).

[10] K. Murphy, P. Travers, and M. Walport, *Janeway's Immunobiology*, vol. 2 (Garland Science, 2001), 7th ed.

[11] J. D. Farmer, N. H. Packard, and A. S. Perelson, Physica D: Nonlinear Phenomena **22**, 187 (1986).

[12] R. Ahmed and D. Gray, Science **272**, 54 (1996).

[13] A. S. Perelson and G. Weisbuch, Reviews of Modern Physics **69**, 1219 (1997).

[14] R. Antia, V. V. Ganusov, and R. Ahmed, Nature Reviews Immunology **5**, 101 (2005).

[15] D. L. Farber, N. a. Yudanin, and N. P. Restifo, Nature Reviews Immunology **14**, 24 (2014).

[16] V. R. Buchholz, T. N. M. Schumacher, and D. H. Busch, Annual review of immunology **34**, 65 (2016).

[17] A. Mayer, V. Balasubramanian, T. Mora, and A. M. Walczak, Proceedings of the National Academy of Sciences **112**, 5950 (2015).

[18] K. L. Y. Christensen, R. C. Holman, C. a. Steiner, J. J. Sejvar, B. J. Stoll, and L. B. Schonberger, Clinical infectious diseases **49**, 1025 (2009).

[19] P. Saule, J. Trauet, V. Dutriez, V. Lekeux, J. P. Dessaint, and M. Labalette, Mechanisms of Ageing and Development **127**, 274 (2006).

[20] W. T. Shearer, H. M. Rosenblatt, R. S. Gelman, R. Oymopito, S. Plaeger, E. R. Stiehm, D. W. Wara, S. D. Douglas, K. Luzuriaga, E. J. McFarland, et al., Journal Of Allergy And Clinical Immunology **112**, 973 (2003).

[21] R. Antia, S. S. Pilyugin, and R. Ahmed, Immunology **95**, 14926 (1998).

[22] I. J. Amanna, N. E. Carlson, and M. K. Slifka, New England Journal of Medicine **357**, 1903 (2007).

[23] F. Sallusto, A. Lanzavecchia, K. Araki, and R. Ahmed, Immunity **33**, 451 (2010).

[24] D. C. Macallan and B. Asquith, Vaccines **5** (2017).

[25] D. Mason, Immunology Today **19**, 395 (1998).

[26] V. I. Zarnitsyna, B. D. Evavold, L. N. Schoettle, J. N. Blattman, and R. Antia, Frontiers in Immunology **4** (2013).

[27] A. Ellebedy, F. Krammer, and A. Rafi, Proc Natl Acad Sci U S A **111**, 13133 (2014).

[28] S. Li, N. L. Sullivan, N. Rouphael, T. Yu, S. Banton, M. S. Maddur, M. Mccausland, C. Chiu, J. Canniff, S. Dubey, et al., Cell **169**, 862 (2017).

[29] J. Quiel, S. Caucheteux, A. Laurence, N. J. Singh, G. Bocharov, S. Z. Ben-Sasson, Z. Grossman, and W. E. Paul, Proceedings of the National Academy of Sciences of the United States of America **108**, 3312 (2011).

[30] G. Bocharov, J. Quiel, T. Luzyanina, H. Alon, E. Chiglintsev, V. Chereshnev, M. Meier-Schellersheim, W. E. Paul, and Z. Grossman, Proceedings of the National Academy of Sciences **108**, 3318 (2011).

[31] R. J. De Boer and A. S. Perelson, The Journal of Immunology **190**, 5454 (2013).

[32] V. I. Zarnitsyna, J. Lavine, A. Ellebedy, R. Ahmed, and R. Antia, PLoS Pathogens **12**, e1005692 (2016).

[33] A. S. Perelson and G. F. Oster, Journal of theoretical biology **81**, 645 (1979).

[34] K. Friston, K. Stephan, B. Li, and J. Daunizeau, Mathematical Problems in Engineering **2010** (2010).

[35] M. G. Netea, L. A. B. Joosten, E. Latz, K. H. G. Mills, G. Natoli, H. G. Stunnenberg, L. A. J. O'Neill, and R. J. Xavier, Science **352**, aaf1098 (2016).

[36] I. M. Johnstone, D. M. Titterington, K. P. Adragni, R. D. Cook, D. L. Banks, L. House, K. Killhoury, D. Barber, M. J. Beal, Z. Ghahramani, et al., Philosophical transactions. Series A, Mathematical, physical, and engineering sciences **367**, 4237 (2009).

[37] M. Advani and S. Ganguli, Physical Review X **6**, 1 (2016).

[38] C. M. Glaze, J. W. Kable, and J. I. Gold, eLife **4**, 1 (2015).

[39] C. M. Glaze, A. L. Filipowicz, J. W. Kable, V. Balasubramanian, and J. I. Gold, Nature Human Behaviour **2**, 213 (2018).

[40] A. Piet, A. El Hady, and C. D. Brody, arXiv preprint arXiv:1710.05945 (2017).

[41] M. Vergassola, E. Villermaux, and B. I. Shraiman, Nature **445**, 406 (2007).

[42] M. Łuksza and M. Lässig, Nature **507**, 57 (2014).

[43] M. Luksza, N. Riaz, V. Makarov, V. P. Balachandran, M. D. Hellmann, A. Solovyov, N. A. Rizvi, T. Merghoub, A. J. Levine, T. A. Chan, et al., Nature **551**, 517 (2017).

[44] A. Etheridge, *Some Mathematical Models from Population Genetics* (Springer-Verlag, 2012), lecture no ed.

[45] A. Gelman, J. B. Carlin, H. S. Stern, and D. B. Rubin, *Bayesian data analysis* (CRC Press, Boca Raton, 2004).

[46] R. C. Griffiths and D. Spanò, arXiv preprint arXiv:1003.4650 (2010).

[47] Y. S. Song and M. Steinrücken, Genetics **190**, 1117 (2012).

[48] R. C. Griffiths, pp. 67–75 (1984).

[49] P. A. Jenkins and D. Spano, arXiv p. 23 (2015).

[50] L. Zdeborová and F. Krzakala, arXiv preprint arXiv:1511.02476 **8732**, 1 (2015).

[51] D. Zehn, S. Y. Lee, and M. J. Bevan, Nature **458**, 211 (2009).

SI Text A: The immigration-drift model of pathogen dynamics

1. Steady-state distribution and the backward equation

The steady-state distribution of the pathogen dynamics (Eq. 9) is a Dirichlet distribution with the parameter vector $\boldsymbol{\theta}$ [44],

$$\rho_s(\boldsymbol{Q}) = \frac{1}{Z(\boldsymbol{\theta})} \prod_i Q_i^{\theta_i - 1} =: \mathcal{D}(\boldsymbol{Q}, \boldsymbol{\theta}). \tag{A1}$$

The normalizing constant is the multivariate Beta function, defined in terms of Gamma functions as $Z(\boldsymbol{\theta}) = \prod_i \Gamma(\theta_i)/\Gamma(|\boldsymbol{\theta}|)$.

To obtain a solution to Eq. 9 we write $\rho(\boldsymbol{Q}, t)$ as the product of the steady-state distribution $\rho_s(\boldsymbol{Q})$ with a time-varying function $f(\boldsymbol{Q}, t)$

$$\rho(\boldsymbol{Q}, t) = \rho_s(\boldsymbol{Q}) f(\boldsymbol{Q}, t). \tag{A2}$$

$f(\boldsymbol{Q}, t)$ can be shown to obey the backward equation

$$\tau \frac{\partial f(\boldsymbol{Q}, t)}{\partial t} = \frac{1}{2} \sum_a \left[(\theta_a - |\boldsymbol{\theta}| Q_a) \frac{\partial}{\partial Q_a} f(\boldsymbol{Q}, t) \right] + \frac{1}{2} \sum_{a,b} \left[Q_a (\delta_{a,b} - Q_b) \frac{\partial^2}{\partial Q_a \partial Q_b} f(\boldsymbol{Q}, t) \right]. \tag{A3}$$

By considering a coalescent dual process a probabilistic expansion can be derived from the backward equation (App. A 2). This probabilistic expansion forms the basis of the approximate count dynamics Eqs. 4,5. It also suggest an efficient method to sample from the transition density used in the simulation of the pathogen dynamics (Sec. A 5). A decomposition of f in Jacobi polynomials, which are the eigenfunctions of Eq. A3, allows us to derive an efficient method to analytically solve the Bayesian prediction and update steps (App. A 3). We use this alternative method to benchmark the quality of the inference achieved by the approximate counting scheme (Fig. S1).

2. Dirichlet mixture expansion of the belief distribution

Let us assume that the belief distribution at some time $B(\boldsymbol{Q}, t)$ is a mixture of Dirichlet distributions

$$B(\boldsymbol{Q}, t) = \sum_{\boldsymbol{m}} c_{\boldsymbol{m}}(t) \mathcal{D}(\boldsymbol{Q}, \boldsymbol{m} + \boldsymbol{\theta}), \tag{A4}$$

with $\boldsymbol{m} = (m_1, m_2, \cdots)$ for integer m_i and mixture coefficients $c_{\boldsymbol{m}} \geq 0, \sum_{\boldsymbol{m}} c_{\boldsymbol{m}} = 1$. In particular this assumption holds true for the prior belief at $t = 0$ which is described by the single component $\boldsymbol{m} = 0$. Both the update and prediction step of the inference procedure can be reduced to a much simpler procedure in terms of the mixture coefficients as derived below. Using the mixture assumption $f(\boldsymbol{Q}, t) = B(\boldsymbol{Q}, t)/\rho_s(\boldsymbol{Q})$ is a mixture $f(\boldsymbol{Q}, t) = \sum_{\boldsymbol{m}} c_{\boldsymbol{m}}(t) f_{\boldsymbol{m}}(\boldsymbol{Q})$ of functions

$$f_{\boldsymbol{m}}(\boldsymbol{Q}) = (|\boldsymbol{\theta}|)_{|\boldsymbol{m}|} \prod_i \frac{Q_i^{m_i}}{(\theta_i)_{m_i}}, \tag{A5}$$

where $(a)_m = a(a+1) \cdots (a+m-1)$. We can interpret the mixture coefficients as a probability distribution and define an average of a quantity x as $\langle x \rangle = \sum_{\boldsymbol{m}} c_{\boldsymbol{m}} x_{\boldsymbol{m}}$. Using this notation the expected frequency is given by

$$\bar{\boldsymbol{Q}} = \langle (\boldsymbol{m} + \boldsymbol{\theta})/(|\boldsymbol{m}| + |\boldsymbol{\theta}|) \rangle. \tag{A6}$$

To perform the update step Eq. 1 we calculate

$$Q_a f_{\boldsymbol{m}}(\boldsymbol{Q}) = \frac{\theta_a + m_a}{|\boldsymbol{\theta}| + |\boldsymbol{m}|} f_{\boldsymbol{m} + \boldsymbol{e}_a}(\boldsymbol{Q}). \tag{A7}$$

Here and in the following we define $\boldsymbol{e}_i$ as the unit basis vector with i-th entry 1 and all other entries 0. It follows that the new belief distribution is still a mixture of Dirichlet distributions with

$$c_{\boldsymbol{m} + \boldsymbol{e}_a}(t^+) = \frac{\theta_a + m_a}{|\boldsymbol{\theta}| + |\boldsymbol{m}|} c_{\boldsymbol{m}}(t^-)/Z, \tag{A8}$$

where $Z = \langle (\theta_a + m_a)/(|\boldsymbol{\theta}| + |\boldsymbol{m}|) \rangle_{c_{\boldsymbol{m}}(t^-)}$ is a normalization constant. To gain intuition about Eq. A8 let us consider the static environment limit $\tau \to \infty$. Applying Eq. A8 sequentially starting from the prior belief shows that the belief distribution remains composed of a single component ($c_{\boldsymbol{m}} = 1$ for $\boldsymbol{m} = \boldsymbol{m}^\star(t)$, $c_{\boldsymbol{m}} = 0$ otherwise) in this limit, where $\boldsymbol{m}^\star(t)$ is the vector of counts of how often the different pathogens have been encountered. This is a classical result in Bayesian statistics [45]. The fact that the belief distribution stays within the Dirichlet family when performing Bayesian updates starting from a Dirichlet prior is known as the conjugacy property of the Dirichlet prior with regards to the categorical likelihood function. The interpretation of $\boldsymbol{m}$ as the number of observations of a particular category motivates thinking about the θ_i as pseudocounts, as according to Eq. A4 they are added to the counts of different categories to pull the estimate of the frequencies closer to the prior expectations.

The prediction step Eq. 2 asks for an application of the backward operator $\mathcal{A}^\dagger$ to $f(\boldsymbol{Q}, t)$. As the backward equation is linear it acts independently on the different components. Applying $\mathcal{A}^\dagger$ to $f_{\boldsymbol{m}}$ yields

$$\mathcal{A}^\dagger f_{\boldsymbol{m}}(\boldsymbol{Q}) = \frac{1}{2}(|\boldsymbol{m}| + |\boldsymbol{\theta}| - 1) \left[\sum_{i=1}^{K} m_i f_{\boldsymbol{m}-\boldsymbol{e}_i}(\boldsymbol{Q}) - |\boldsymbol{m}| f_{\boldsymbol{m}}(q) \right]. \tag{A9}$$

By considering $\mathcal{A}^\dagger$ in Eq. A9 as acting on $\boldsymbol{m}$ we obtain a dual death process description of the Wright-Fisher diffusion [46]. Transitions from $\boldsymbol{m} \to \boldsymbol{m} - \boldsymbol{e}_i$ happen at a rate $\frac{1}{2}(|\boldsymbol{m}| + |\boldsymbol{\theta}| - 1)m_i$. The belief distribution continues to be a mixture of Dirichlet distributions with mixture coefficients that change as

$$\tau \frac{\mathrm{d}c_{\boldsymbol{m}}}{\mathrm{d}t} = \frac{1}{2}(|\boldsymbol{m}| + |\boldsymbol{\theta}|) \left(\sum_i (m_i + 1)c_{\boldsymbol{m}+\boldsymbol{e}_i} \right) - \frac{1}{2}(|\boldsymbol{m}| + |\boldsymbol{\theta}| - 1)|\boldsymbol{m}|c_{\boldsymbol{m}}. \tag{A10}$$

Let us define the mean count vector $\langle \boldsymbol{m} \rangle$. Its dynamical equation is

$$\tau \frac{\mathrm{d}\langle \boldsymbol{m} \rangle}{\mathrm{d}t} = \sum_{\boldsymbol{m}} \boldsymbol{m} \frac{\mathrm{d}c_{\boldsymbol{m}}}{\mathrm{d}t}. \tag{A11}$$

Some algebra leads to the surprisingly simple yet exact equation

$$\tau \frac{\mathrm{d}\langle \boldsymbol{m} \rangle}{\mathrm{d}t} = -\frac{1}{2} \langle (|\boldsymbol{m}| + |\boldsymbol{\theta}| - 1)\boldsymbol{m} \rangle. \tag{A12}$$

The probabilistic expansion allows the derivation of an efficient approximate scheme for inference. For peaked mixture distributions $c_{\boldsymbol{m}}$ we can to a good approximation invert the order of calculating the expectation value and the product in Eq. A12, i.e.

$$\langle (|\boldsymbol{m}| + |\boldsymbol{\theta}| - 1)\boldsymbol{m} \rangle \approx (|\langle \boldsymbol{m} \rangle| + |\boldsymbol{\theta}| - 1)\langle \boldsymbol{m} \rangle. \tag{A13}$$

For peaked mixture distribution the update equation for the mean counts upon encountering pathogen a is approximately

$$\langle \boldsymbol{m}(t^+) \rangle \approx \langle \boldsymbol{m}(t^-) \rangle + \boldsymbol{e}_a, \tag{A14}$$

and the expected frequencies are approximately

$$\bar{\boldsymbol{Q}} \approx \langle \boldsymbol{m} + \boldsymbol{\theta} \rangle / (|\langle \boldsymbol{m} + \boldsymbol{\theta} \rangle|). \tag{A15}$$

Dropping the explicit notation for the average and replacing $\boldsymbol{n} := \boldsymbol{m} + \boldsymbol{\theta}$ we obtain Eqs. 4–6 from the main text.

3. Spectral expansion of the belief distribution

As in App. A 2 we decompose the belief distribution but now we decompose $f(\boldsymbol{Q}, t)$ into the eigenfunctions $g_{\boldsymbol{n}}(\boldsymbol{Q})$ of the backward operator. This approach leads to simpler decoupled equations for the prediction step of the Bayesian inference. To simplify notations we consider the case $K = 2$ but the derivation generalizes to arbitrary K [46]. Note that to describe the dynamics of the frequency of any particular pathogen, we can lump together the frequencies of the other $K - 1$ pathogens for the model we consider. In particular, as the rates of immigration of different types are independent of composition of the population, the dynamics of the chosen pathogen does not depend on distribution of pathogens among the other types. The dynamics of the i-th type for arbitrary K can thus be mapped to $K = 2$ by setting $\theta_1 := \theta_i$ and $\theta_2 := \sum_{j \neq i} \theta_j$.

The steady-state distribution specializes from the Dirichlet distribution to the Beta distribution for $K = 2$

$$\rho_s(q) = q^{\theta_1 - 1}(1 - q)^{\theta_2 - 1}/Z, \tag{A16}$$

where $Z = B(\theta_1, \theta_2) = \Gamma(\theta_1)\Gamma(\theta_2)/\Gamma(\theta_1 + \theta_2)$ is the Beta function. The eigenfunction g_n for an eigenvalue λ_n needs to fulfill the backward equation (Eq. A3), which leads to

$$\frac{1}{2}q(1 - q)\frac{d^2 g_n}{dq^2}(q) + \frac{1}{2}(-\theta_2 q + \theta_1(1 - q))\frac{dg_n}{dq}(q) = -\tau\lambda_n g_n(q), \tag{A17}$$

where $q = Q_1$. Up to a rescaling this equation is a Jacobi differential equation. The solutions of this differential equation are the modified Jacobi Polynomials

$$g_n(q) = P_n^{(\theta_2 - 1, \theta_1 - 1)}(2q - 1), \tag{A18}$$

where $P_n^{(a,b)}(x)$ is the n-th Jacobi polynomial with eigenvalue

$$\lambda_n = \frac{1}{2\tau}n(n + \theta_1 + \theta_2 - 1). \tag{A19}$$

Let us first state a number of properties of these polynomials which we will need later. The polynomials form an orthogonal system with respect to the weight function $q^{\theta_1 - 1}(1 - q)^{\theta_2 - 1}$, i.e.

$$\int_0^1 dq\, g_n(q)g_m(q)q^{\theta_1 - 1}(1 - q)^{\theta_2 - 1} = \delta_{n,m}\Delta_n(\theta_1, \theta_2), \tag{A20}$$

where the normalization coefficients $\Delta_n(\theta_1, \theta_2)$ are given by

$$\Delta_n(\theta_1, \theta_2) = \frac{\Gamma(n + \theta_1)\Gamma(n + \theta_2)}{(2n + \theta_1 + \theta_2 - 1)\Gamma(n + \theta_1 + \theta_2 - 1)\Gamma(n + 1)}. \tag{A21}$$

For $n \geq 1$ the polynomials are related by the recursion formula [47]

$$qg_n(q) = \phi_n^- g_{n-1}(q) + \phi_n^0 g_n(q) + \phi_n^+ g_{n+1}(q), \tag{A22}$$

with coefficients

$$\phi_n^- = \frac{(n + \theta_1 - 1)(n + \theta_2 - 1)}{(2n + \theta_1 + \theta_2 - 1)(2n + \theta_1 + \theta_2 - 2)}, \tag{A23}$$

$$\phi_n^0 = \frac{1}{2} - \frac{\theta_2^2 - \theta_1^2 - 2(\theta_2 - \theta_1)}{2(2n + \theta_1 + \theta_2)(2n + \theta_1 + \theta_2 - 2)}, \tag{A24}$$

$$\phi_n^+ = \frac{(n + 1)(n + \theta_1 + \theta_2 - 1)}{(2n + \theta_1 + \theta_2)(2n + \theta_1 + \theta_2 - 1)}, \tag{A25}$$

while for $n = 0$

$$qg_0(q) = \phi_0^0 g_0(q) + \phi_0^+ g_1(q), \tag{A26}$$

with coefficients

$$\phi_0^0 = \frac{\theta_1}{\theta_1 + \theta_2}, \tag{A27}$$

$$\phi_0^+ = \frac{1}{\theta_1 + \theta_2}. \tag{A28}$$

Now let us write $f(q, t)$ as a linear combination of the eigenfunctions g_n of the backward equation as,

$$f(q, t) = 1 + \sum_{n=1}^{\infty} d_n(t)g_n(q), \tag{A29}$$

with time-varying coefficients $d_n(t)$. The prediction step of the Bayesian inference leads through Eq. A17 to a simple exponential decay of the d_n,

$$\frac{\mathrm{d}d_n(t)}{\mathrm{d}t} = -\lambda_n. \tag{A30}$$

The update step of the Bayesian inference leads through Eq. A22 to

$$d_n(t^+) = \chi_n^- d_{n-1}(t^-) + \chi_n^0 d_n(t^-) + \chi_n^+ d_{n+1}(t^-), \tag{A31}$$

where $\chi_n^-, \chi_n^0, \chi_n^+$ are constants. These constants are normalized versions of the coefficients ϕ that appear in Eq. A22,

$$\chi_n^x = \begin{cases} \phi_n^x / (c_0^0 + d_1(t^-)c_1^-) & \text{if pathogen 1 encountered,} \\ (\delta_{x,0} - \phi_n^x)/(1 - c_0^0 - d_1(t^-)c_1^-) & \text{if pathogen 2 encountered,} \end{cases} \tag{A32}$$

for x in $-, 0, +$, and where $\delta_{x,0} = 1$ for $x = 0$ and $\delta_{x,0} = 0$ otherwise. For efficient numerical computation note that the update step can be performed as a matrix multiplication of the triadiagonal matrix which has c_n^0 along the diagonal, c_{n+1}^- above the diagonal, and c_{n-1}^+ below the diagonal with the vector of coefficients $\boldsymbol{d}(t^-)$ followed by a normalization step. We finally note that the expected frequency is obtained from Eq. A29 as

$$\langle q(t) \rangle = c_0^0 + c_1^- d_1(t). \tag{A33}$$

This implies that the pathogen frequencies change with an autocorrelation timescale

$$\tau_c = \frac{1}{\lambda_1} = \frac{2\tau}{\theta_1 + \theta_2}. \tag{A34}$$

4. Defining an effective timescale for learning in changing environments

The total number of counts $|\boldsymbol{n}|$ follows a piecewise deterministic decay process interspersed by updates at random times. We approximate this stochastic dynamics by the deterministic equation,

$$\frac{\mathrm{d}|\boldsymbol{n}|}{\mathrm{d}t} = -\frac{1}{2\tau}(|\boldsymbol{n}| - 1)(|\boldsymbol{n}| - |\boldsymbol{\theta}|) + \lambda, \tag{A35}$$

where we have replaced the stochastic jumps in the counts do to pathogen encounters by a source-term λ equal to the rate of such jumps. This differential equation is separable and can be solved, which for $|\boldsymbol{n}(0)| = |\boldsymbol{\theta}|$ yields

$$|\boldsymbol{n}(t)| = \frac{|\boldsymbol{\theta}| + 1}{2} + \eta \tanh\left(\frac{t\sqrt{\eta}}{2\tau} + \operatorname{artanh}\left(\frac{|\boldsymbol{\theta}| - 1}{2\sqrt{\eta}}\right)\right), \tag{A36}$$

for

$$\eta = 2\lambda\tau + \frac{(|\boldsymbol{\theta}| - 1)^2}{4} \tag{A37}$$

Considering the limit of long environmental correlation times, $2\lambda\tau \gg \frac{(|\boldsymbol{\theta}| - 1)^2}{4}$, we obtain the simplified expression

$$|\boldsymbol{n}(t)| - |\boldsymbol{\theta}| = \sqrt{2\lambda\tau} \tanh\left(\frac{\lambda t}{\sqrt{2\lambda\tau}}\right). \tag{A38}$$

If we further take the limit $2\lambda\tau \gg (\lambda t)^2$ of no attrition, then we recover the simple linear scaling of the expected number of encounters with time

$$|\boldsymbol{n}(t)| - |\boldsymbol{\theta}| = \lambda t. \tag{A39}$$

The number of remembered encounters $|\boldsymbol{n}(t)| - |\boldsymbol{\theta}|$ relative to the effective number of present pathogens $|\boldsymbol{\theta}|$ controls how much memory improves protection (Fig. 2).

5. Efficient simulation by sampling from the transition density

The transition density of going from $\boldsymbol{x}$ to $\boldsymbol{y}$ in time t for the Wright-Fisher diffusion can be written as [46]

$$f(\boldsymbol{x},\boldsymbol{y},t) = \sum_{|\boldsymbol{l}|=0}^{\infty} q_{|\boldsymbol{l}|}^{|\boldsymbol{\theta}|}(t) \sum_{\{\boldsymbol{l}:|\boldsymbol{l}|fixed\}} \mathcal{M}(\boldsymbol{l},\boldsymbol{x})\mathcal{D}(\boldsymbol{y},\boldsymbol{\theta}+\boldsymbol{l}), \tag{A40}$$

where $\mathcal{M}(\boldsymbol{l},\boldsymbol{x})$ is the multinomial distribution

$$\mathcal{M}(\boldsymbol{l},\boldsymbol{x}) = \binom{|\boldsymbol{l}|}{\boldsymbol{l}} x_1^{l_1} \cdots x_K^{l_K}, \tag{A41}$$

and where $q_{|\boldsymbol{l}|}^{|\boldsymbol{\theta}|}(t)$ are the transition functions of a dual pure death process. This description of the transition density can be derived based on similar arguments to those made in App. A 2 [46]. The death process describes the loss of un-mutated lineages going backward in time through coalescence and mutations. For small times $q_{|\boldsymbol{l}|}^{|\boldsymbol{\theta}|}(t)$ is asymptotically normal with mean $\mu(t)$ and variance $\sigma^2(t)$ [48, 49]

$$\mu(t) = \frac{2\eta}{t}, \tag{A42}$$

$$\sigma^2(t) = \begin{cases} \frac{2\eta}{t}(\eta+\beta)^2 \left(1 + \frac{\eta}{\eta+\beta} - 2\eta\right)\beta^{-2}, & \beta \neq 0, \\ \frac{2}{3t}, & \beta = 0, \end{cases} \tag{A43}$$

$$\text{where } \beta = \frac{1}{2}(|\boldsymbol{\theta}|-1)t, \tag{A44}$$

$$\text{and } \eta = \begin{cases} \frac{e^\beta}{e^\beta-1}, & \beta \neq 0, \\ 1, & \beta = 0. \end{cases} \tag{A45}$$

To sample from the transition density function we thus proceed in three steps: Generate a normally distributed random number $|\boldsymbol{l}|$ according to the asymptotic distribution of $q_{|\boldsymbol{l}|}^{|\boldsymbol{\theta}|}(t)$. Then draw $\boldsymbol{l}$ from $\mathcal{M}(l,\boldsymbol{x})$. Finally, draw $\boldsymbol{y}$ from $\mathcal{D}(\boldsymbol{\theta}+\boldsymbol{l})$.

SI Text B: Induced repertoire dynamics

1. Memory fold change upon a pathogen encounter

.

The inference dynamics induces via the mapping from $\hat{\boldsymbol{Q}}$ to $\boldsymbol{P}^\star$ (Eq. 3) a dynamics of an optimally adapting immune repertoire. To get some intuition we derive more explicitly the form this induced dynamics takes in some limits.

As long as the solution to the optimization problem stays in the interior of the optimization domain the optimal coverage for a cost $c(\tilde{P}_a) = 1/\tilde{P}_a^\alpha$ is [17]

$$\tilde{P}_a = \frac{1}{Z}Q_a^{1/(1+\alpha)}, \text{ with } Z = \sum_a Q_a^{1/(1+\alpha)} \tag{B1}$$

where Z is a normalization factor. (For the logarithmic cost the expression above holds for $\alpha = 0$.) By combining Eqs.4,6 we can write an update equation for the expected frequencies upon encounter of antigen a as

$$\hat{\boldsymbol{Q}}^+ = \frac{\boldsymbol{n}^- + \boldsymbol{e}_a}{|\boldsymbol{n}^+|} = \frac{|\boldsymbol{n}^-|\hat{\boldsymbol{Q}}^- + \boldsymbol{e}_a}{|\boldsymbol{n}^+|}, \tag{B2}$$

where to simplify notations we use $\hat{\boldsymbol{Q}}(t^+) = \hat{\boldsymbol{Q}}^+$, and where $|\boldsymbol{n}^+| = |\boldsymbol{n}^-| + 1$. It follows that the update of the coverages is given by

$$\tilde{P}_b^+ = \frac{[(Q_b^-|\boldsymbol{n}^-| + \delta_{a,b})/(|\boldsymbol{n}^+|)]^{1/(1+\alpha)}}{Z^+} \tag{B3}$$

$$= \frac{[((\tilde{P}_b^-)^{1+\alpha}|\boldsymbol{n}^-| + \delta_{a,b}/(Z^-)^{1+\alpha})/|\boldsymbol{n}^+|]^{1/(1+\alpha)}}{Z^+/Z^-} \tag{B4}$$

The change in coverage for antigen a is

$$\tilde{P}_a^+/\tilde{P}_a^- \approx [1 + \kappa/(\tilde{P}_a^-)^{1+\alpha}]^{1/(1+\alpha)}, \tag{B5}$$

where $\kappa = 1/(|\boldsymbol{n}^+|(Z^-)^{1+\alpha})$ and where we have neglected the change in normalization which is of order $1/K$ relative to the update size. To fit the data set we note that a proportional rescaling of $\tilde{P}_a$ by a factor k can be subsumed within the model by redefining $\kappa \to \kappa k^{1+\alpha}$. Therefore the scaling of $\tilde{P}_a$ to an antibody titer can be subsumed within κ.

For a uniform prior distribution we have $|\boldsymbol{n}| \approx K\theta$ and $Z^- \approx \sum_a 1/K^{1/(1+\alpha)} = K^{\alpha/(1+\alpha)}$, which for $K\theta \gg 1$ leads to $\kappa = 1/(K^{1+\alpha}\theta)$. The fold change upon an encounter of a pathogen starting from a naive repertoire $\tilde{P}_a^- = 1/K$ thus depends as follows on the sparsity of the environment,

$$\tilde{P}_a^+/\tilde{P}_a^- = [1 + 1/\theta]^{1/(1+\alpha)}. \tag{B6}$$

2. Mean-field dynamics

Besides the large changes of the naive repertoire upon a primary infection there are situations in which the inferred distribution is changing in a more continuous manner, e.g. updating in the limit of many previous samples, or the prediction step. We thus now ask how small changes in the expected frequencies of pathogens $\hat{\boldsymbol{Q}}$ change the coverage $\tilde{\boldsymbol{P}}$. We assume that there is no cross-reactivity $f_{r,a} = \delta_{r,a}$, and consider power-law cost functions, where we have optimal receptor frequency distribution $P_r^\star = Q_r^\beta/Z$ with $\beta = 1/(1+\alpha)$. As a preliminary we calculate the Jacobian

$$\frac{\partial P_r^\star}{\partial \hat{Q}_{r'}} = \delta_{r,r'}\frac{1}{Z}\frac{\partial \hat{Q}_r^\beta}{\partial \hat{Q}_r} - \frac{Q_r^\beta}{Z^2}\frac{\partial Z}{\partial \hat{Q}_{r'}}, \tag{B7}$$

$$= \beta P_r^\star \left[\frac{\delta_{r,r'}}{\hat{Q}_r} - \frac{P_{r'}}{\hat{Q}_{r'}} \right], \tag{B8}$$

We can then show that the dynamics in terms of $P_r^\star$ follows,

$$\frac{\mathrm{d}P_r^\star}{\mathrm{d}t} = \sum_{r'} \frac{\partial P_r^\star}{\partial \hat{Q}_{r'}}\frac{\mathrm{d}\hat{Q}_{r'}}{\mathrm{d}t}, \tag{B9}$$

$$= P_r^\star \left[\beta\frac{\mathrm{d}\ln\hat{Q}_r}{\mathrm{d}t} - \sum_{r'} P_{r'}\beta\frac{\mathrm{d}\ln\hat{Q}_{r'}}{\mathrm{d}t} \right], \tag{B10}$$

which is of the form of a replicator equation

$$\frac{\mathrm{d}P_r^\star}{\mathrm{d}t} = P_r^\star \left[f_r - \bar{f} \right] \tag{B11}$$

with "fitness" $f_r = \beta\frac{\mathrm{d}\ln\hat{Q}_r}{\mathrm{d}t}$ and mean fitness $\bar{f} = \sum_{r'} P_{r'}f_{r'}$.

Based on this general result we now analyze the dynamics of the repertoire due to the sequential Bayesian filtering. Equivalently to Eq. B2 the change of inferred distribution $\Delta\hat{\boldsymbol{Q}} = \hat{\boldsymbol{Q}}^+ - \hat{\boldsymbol{Q}}$ upon encountering antigen a is given by

$$\Delta\hat{\boldsymbol{Q}} = \frac{\boldsymbol{e}_a - \hat{\boldsymbol{Q}}}{|\boldsymbol{n}| + 1}, \tag{B12}$$

where $\boldsymbol{e}_a$ is the unit vector with a-th entry one and all other zero. Asymptotically for large $|\boldsymbol{n}|$ every update has a small effect only, and we might consider a mean-field description. In this description we replace $\boldsymbol{e}_a$ by its expectation value $\boldsymbol{Q}$ and define an average rate of change per unit time by multiplying the update size by the frequency λ of pathogen encounters:

$$\frac{\mathrm{d}\hat{\boldsymbol{Q}}}{\mathrm{d}t} = \frac{\lambda}{|\boldsymbol{n}| + 1}(\boldsymbol{Q} - \hat{\boldsymbol{Q}}). \tag{B13}$$

Here $\boldsymbol{Q}$ is the actual distribution of pathogens, and $\hat{\boldsymbol{Q}}$ are the expected frequencies of pathogens based on the immune system's internal belief. For the prediction step we have a dynamics for counts, which we can convert into a dynamics

for the inferred distribution. We have $\hat{Q}_r = n_r/|\boldsymbol{n}|$ and a dynamics on counts given by Eq. 5. From there we obtain

$$\frac{d\hat{Q}_r}{dt} = \frac{1}{|\boldsymbol{n}|}\frac{dn_r}{dt} - \frac{n_r}{|\boldsymbol{n}|^2}\frac{d|\boldsymbol{n}|}{dt}, \tag{B14}$$

$$= -\frac{(|\boldsymbol{n}|-1)|\boldsymbol{\theta}|}{2\tau|\boldsymbol{n}|}(\hat{Q}_r - \hat{Q}_r^0), \tag{B15}$$

where $\hat{Q}_r^0 = \theta_r/|\boldsymbol{\theta}|$ is the prior guess for the distribution. Taken together we have

$$\frac{d\hat{\boldsymbol{Q}}}{dt} = \gamma(t)(\boldsymbol{Q} - \hat{\boldsymbol{Q}}) - \delta(t)(\hat{\boldsymbol{Q}} - \hat{\boldsymbol{Q}}^0). \tag{B16}$$

with the (time-varying) coefficients

$$\gamma(t) = \frac{\lambda}{|\boldsymbol{n}(t)|+1}, \tag{B17}$$

$$\delta(t) = \frac{(|\boldsymbol{n}(t)|-1)|\boldsymbol{\theta}|}{2\tau|\boldsymbol{n}(t)|}. \tag{B18}$$

The fitness in the replicator equation is then

$$f_r = \beta\left(\gamma(t)\frac{Q_r}{\hat{Q}_r} + \delta(t)\frac{\hat{Q}_r^0}{\hat{Q}_r}\right). \tag{B19}$$

The fixed point of the dynamics in a static environment $\delta(t) = 0$ is the optimal repertoire as expected from the asymptotic optimality of Bayesian inference. Replacing $\hat{Q}_r = (ZP_r^\star)^{1+\alpha}$ we then obtain a fitness

$$f_r = \frac{\gamma(t)}{Z^{1+\alpha}}\frac{Q_r}{(P_r^\star)^{1+\alpha}} \tag{B20}$$

which except for the prefactor is equivalent to the population dynamics proposed previously in [17]. That work did not consider the prefactor that leads to a slowing down of the dynamics with time to reflect a tradeoff between new evidence and past experience. The prediction steps relaxes the inferred distribution towards the prior distribution with a speed that for large $|\boldsymbol{n}|$ is proportional to $|\boldsymbol{\theta}|/\tau$.

3. Updating a cross-reactive repertoire

We now consider the repertoire dynamics in the presence of cross-reactivity. In a first order Taylor expansion the change in the repertoire composition upon a pathogen encounter is given by

$$\Delta \boldsymbol{P}^\star \approx \boldsymbol{J_G}(\hat{\boldsymbol{Q}})\Delta\hat{\boldsymbol{Q}}, \tag{B21}$$

where

$$(\boldsymbol{J_G})_{r,r'}(\hat{\boldsymbol{Q}}) = \frac{\partial G_r(\hat{\boldsymbol{Q}})}{\partial Q_r'} \tag{B22}$$

is the Jacobian of the mapping function (Eq. 3).

The mapping between pathogen frequencies and the optimal repertoire takes the form $\boldsymbol{G}(\boldsymbol{Q}) = \boldsymbol{F}^{-1}\tilde{\boldsymbol{P}}^\star(\boldsymbol{Q})$ (if achievable given the constraint that no receptor frequency can be negative). $\tilde{\boldsymbol{P}}^\star(\boldsymbol{Q})$ is a function that depends on the cost function. The Jacobian can thus be calculated using the chain rule as

$$\boldsymbol{J_G}(\hat{\boldsymbol{Q}}) = \boldsymbol{F}^{-1}\boldsymbol{J}_{\tilde{\boldsymbol{P}}^\star}(\hat{\boldsymbol{Q}}). \tag{B23}$$

For the power-law cost function we have $\tilde{P}_a^\star = R\hat{Q}_a^\beta/Z$, where $R = \sum_a f_{r,a}$ is the row sum of F, which assume to be constant. Analogously to the derivation of the Jacobian in the previous section we derive

$$\frac{\partial \tilde{P}_a^\star}{\partial \hat{Q}_{a'}} = \beta\tilde{P}_a^\star\left[\frac{\delta_{a,a'}}{\hat{Q}_a} - \frac{\tilde{P}_{a'}}{R\hat{Q}_{a'}}\right], \tag{B24}$$

from which with some algebra follows

$$\Delta \boldsymbol{P}^\star \approx \left(\frac{R}{\tilde{P}_a^\star}\right)^\alpha \frac{1}{(1+\alpha)(|\boldsymbol{n}|+1)Z^{1+\alpha}}\left(R\boldsymbol{F}^{-1}\boldsymbol{e}_a - \boldsymbol{P}^\star\right). \tag{B25}$$

Here, there is a departure from the dynamics of the number N_r of lymphocytes with receptor r proposed in [17],

$$\Delta N_r = N_r \Delta t \left[A(\tilde{N}_a)f_{r,a} - d\right], \tag{B26}$$

where proliferation is proportional to $f_{r,a}$, instead of $(F^{-1})_{r,a}$.

SI Text C: Inference of high-dimensional categorical distributions from few samples

1. Mean cost versus time

In this Appendix we will derive analytical expressions for the optimized cost as a function of time

$$c(t) = \sum_{a=1}^{K} Q_a(t)c(\tilde{P}_a^\star(t)) \tag{C1}$$

with the following simplifying assumptions: absence of cross-reactivity, $f_{a,r} = \delta_{a,r}$ and $\tilde{P}_a^\star(t) = P_a^\star(t)$; no attrition, $\tau \to +\infty$; and a power-law cost function $c(P) = P^{-\alpha}$. The prior on $\boldsymbol{Q}$ is a homogeneous Dirichlet distribution:

$$P(\boldsymbol{Q}) \propto \prod_{a=1}^{K} Q_a^{\theta-1}. \tag{C2}$$

This problem is equivalent to the Bayesian inference of a distribution drawn from a Dirichlet meta-distribution. Asymptotic convergence properties of Bayesian inference procedures are well-established [45], but the convergence speed of Bayesian estimators of the distribution in the non-asymptotic regime has been much less studied to our knowledge. Analysing the behaviour of $c(t)$ is equivalent to analysing the convergence of the estimated distribution to the true one with increasing number of samples. Here we will establish the relevant scaling for few samples.

We consider the biologically relevant regime of high dimension but effective sparsity of the distribution, $K\theta \gg 1$, $\theta \ll 1$. Our main insight is that for such sparse distribution Bayesian inference is effective when the number of samples is on the order of a few $K\theta$, instead of the potentially much larger K.

The prominent role sparsity plays in allowing for more efficient estimation is reminiscent of compressed sensing [50]. Non-asymptotic results about inference in high-dimensional settings have been explored recently in the context of machine learning [37]. Both connections merit further exploration.

We define the expected cost as $\langle c(t)\rangle$, where the average is taken over both random choices of $\boldsymbol{Q}$, and random realizations of the pathogen encounters, $\boldsymbol{n}$, which are distributed according to:

$$n_a(t) \sim \text{Poisson}(\lambda t Q_a), \tag{C3}$$

where λ is the encounter rate. The number of encounters determine the average belief for $\boldsymbol{Q}$,

$$\hat{Q}_a(t) = \frac{\theta + n_a}{K\theta + \sum_{a=1}^{L} n_a} \approx \frac{\theta + n_a}{K\theta + \lambda t}, \tag{C4}$$

which itself shapes the optimal response and thus the cost through Eq. C1:

$$P_a^\star(t) = \frac{\hat{Q}_a(t)^{1/(1+\alpha)}}{Z}, \tag{C5}$$

where Z is a normalization constant.

Note that the cost can be expressed in terms of a divergence between the best receptor distribution $\boldsymbol{P}^\star$ given full knowledge of $\boldsymbol{Q}$ and the actual receptor distribution $\boldsymbol{P}$. Defining $P_a^\star = Q_a^{1/(1+\alpha)}/\tilde{Z}$ and replacing into Eq. C1, we obtain:

$$c(t) = \tilde{Z}^{1+\alpha} \sum_a (P_a^\star)^{1+\alpha} P_a^{-\alpha} = c_\infty \exp\left[\alpha D_{1+\alpha}(\boldsymbol{P}^\star \| \boldsymbol{P})\right], \tag{C6}$$

where $c_\infty = \tilde{Z}^{1+\alpha}$ is the asymptotic cost for $P = P^\star$ and where $D_\beta(\boldsymbol{P} \| \boldsymbol{Q}) := (\beta-1)^{-1}\ln[\sum_a P_a^\beta Q_a^{1-\beta}]$ is the Rényi divergence of order β, which reduces to the standard Kullback-Leibler divergence for $\beta = 1$, i.e. $\alpha = 0$.

2. Reducing the problem to a single pathogen

By symmetry all terms in the sum of Eq. C1 are the equal on average and we have

$$\langle c(t) \rangle = K \int dq\, \rho(q) q\, \langle c(p) \rangle_n, \tag{C7}$$

where we have introduced the short hand notations $q := Q_a$ and $p := \tilde{P}_a$, and where the average is taken over $n := n_a$. The pathogen frequency q is approximately Gamma-distributed:

$$\rho(q) \approx \frac{(K\theta)^\theta}{\Gamma(\theta)} q^{\theta-1} e^{-K\theta q}. \tag{C8}$$

In general, the expectation value depends through p on all previous encounters with any of the pathogens (i.e. all the other $n_{a'}$, $a' \neq a$). In high dimensions we can approximate this dependence by neglecting the correlation of the normalization factor Z with $\hat{q}$ and using an effective Z. For the power law cost functions we then have

$$p = \hat{q}^{\frac{1}{1+\alpha}}/Z \quad \text{with } Z \approx K\langle \hat{q}^{\frac{1}{1+\alpha}} \rangle. \tag{C9}$$

For logarithmic cost this simplifies to $p = \hat{q}$ and $Z = 1$. From Eq. C9 it follows that

$$\langle c(t) \rangle \approx K Z^\alpha \langle q\hat{q}^{-\frac{\alpha}{1+\alpha}} \rangle = K^{1+\alpha} \langle \hat{q}^{\frac{1}{1+\alpha}} \rangle^\alpha \langle q\hat{q}^{-\frac{\alpha}{1+\alpha}} \rangle. \tag{C10}$$

We have $\hat{q} = (\theta+n)/(K\theta+\lambda t)$, but as the equation is invariant to a linear rescaling of $\hat{q}$ we can replace $\hat{q}$ by simply $\theta + n$.

3. Costs for perfect or no information

Asymptotically the distribution is learned perfectly and we have $\hat{q} = q$. Plugging this into the expressions derived previously we obtain

$$\bar{c}_\infty := \langle c(\infty) \rangle \approx K^{1+\alpha} \langle q^{1/(1+\alpha)} \rangle^{1+\alpha}, \quad \bar{c}_\infty \approx -K\langle q \ln(q) \rangle, \tag{C11}$$

for the power-law and logarithmic cost function respectively. Performing the integrals we obtain

$$\bar{c}_\infty \approx (K\theta)^\alpha \Gamma(1/(1+\alpha))^{1+\alpha}, \quad \bar{c}_\infty \approx \ln(K\theta) + \gamma, \tag{C12}$$

where $\Gamma(z)$ is the Gamma function and γ the Euler-Mascheroni constant. For $\alpha = 1$ this specializes to $\bar{c}_\infty = \pi K\theta$. For the logarithmic cost $\bar{c}_\infty$ is equal to the Shannon entropy of the distribution, which suggests an interpretation of $K\theta$ as the effective number of pathogens that are present.

We can compare these costs for those obtained for a uniform repertoire

$$\bar{c}_0 := \langle c(0) \rangle = K^\alpha, \quad \bar{c}_0 = \ln(K), \tag{C13}$$

to obtain

$$\bar{c}_\infty/\bar{c}_0 \propto \theta^\alpha, \quad \bar{c}_\infty/\bar{c}_0 = (\ln(K\theta) + \gamma)/\ln(K) \tag{C14}$$

for power law and logarithmic cost respectively. As expected, in more sparse environments a larger relative improvement can be obtained by learning the distribution.

4. Scaling in the limit of few samples

In the limit of small sampling, each pathogen has been seen at most once, meaning that n is binary and distributed according to a Bernouilli variable with mean $\lambda t q$. Then we can use the approximation $\langle (\theta+n)^\beta \rangle \approx \theta^\beta(1-\lambda t q) + \lambda t q$ to obtain

$$\langle \hat{q}^{\frac{1}{1+\alpha}} \rangle = \theta^{\frac{1}{1+\alpha}} + (1 - \theta^{\frac{1}{1+\alpha}})\lambda t \langle q \rangle \approx \theta^{\frac{1}{1+\alpha}} + \frac{\lambda t}{K} \tag{C15}$$

$$\langle q\hat{q}^{-\frac{\alpha}{1+\alpha}} \rangle = \theta^{-\frac{\alpha}{1+\alpha}}\langle q \rangle + (1 - \theta^{-\alpha}1 + \alpha)\lambda t \langle q^2 \rangle \approx \frac{\theta^{-\frac{\alpha}{1+\alpha}}}{K}\left[1 - \frac{\lambda t}{K\theta}\right] \tag{C16}$$

Putting things together we obtain

$$\langle c(t) \rangle \approx K^\alpha \left(1 + \frac{\lambda t}{K\theta} \theta^{\frac{\alpha}{1+\alpha}} \right)^\alpha \left(1 - \frac{\lambda t}{K\theta} \right) \approx K^\alpha \left(1 - \left(1 - \alpha\theta^{\frac{\alpha}{1+\alpha}} \right) \frac{\lambda t}{K\theta} \right), \tag{C17}$$

which, except for a correction that vanishes as $\theta \to 0$, scales with $\lambda t/(K\theta)$.

For the logarithmic cost we approximate similarly $\langle \ln(\theta + n) \rangle \approx (1 - \lambda t q)\ln(\theta) + \lambda t q \ln(1 + \theta) \approx (1 - \lambda t q)\ln(\theta)$. We then have

$$\langle c(t) \rangle \approx \ln(K\theta + \lambda t) - K\ln(\theta) \left[\langle q \rangle - \lambda t \langle q^2 \rangle \right]. \tag{C18}$$

Using the formulas for the first and second moments we obtain

$$\langle c(t) \rangle \approx \ln(K\theta + \lambda t) - \ln(\theta) - \frac{\lambda t}{K\theta} \ln(1/\theta) \tag{C19}$$

Approximating further we have

$$\langle c(t) \rangle \approx \ln(K) \left(1 - \left(1 - \frac{\ln(K\theta) - 1}{\ln(K)} \right) \frac{\lambda t}{K\theta} \right). \tag{C20}$$

Again the relative cost depends solely on $\lambda t/(K\theta)$ except for logarithmic corrections that vanish as $K \to \infty$ for fixed $K\theta$.

SI Text D: Infection cost in the expansion-delay regime

We have previously described mechanistic models that give rise to a power-law dependency of infection cost on the coverage [17], in which we have assumed thatthe crucial determinant of infection cost is set by the time delay to recognition of the pathogen by the immune system. Experimental evidence shows that the initial recruitment of a large fraction of all specific lymphocytes often happens rapidly compared to the time it takes for the adaptive immune system to start clearing the infection [51]. We thus might hypothesize that the advantage of higher precursor numbers lies not in shortening the time to detection but in shortening the time to response by a sufficiently large number of effector cells.

To derive the scaling of infection cost with coverage under these conditions, we consider that after an infection at time 0, the number of specific cells grows exponentially with a rate γ, $N(t) = N(0)e^{-\gamma t}$. During the same time the pathogen population grows exponentially as well at a rate γ_p, $P(t) = P(0)e^{-\gamma_p t}$ until a time $t^\star$ at which a threshold level $N^\star$ of specific cells is reached. The expansion-delay time scales as $t^\star = \ln(N^\star/N(0))/\gamma$. If we assume that the cost of an infection is proportional $P(t^\star)$, then the cost scales as a power law with the initial number of specific cells $N(0)^{-\gamma_p/\gamma}$.

SI Text E: Supplementary figures

FIG. S1: Frequency Q of a pathogen over time in a dynamically changing environment along with inferred frequencies $\hat{Q}$. The approximate inference based on the count dynamics (Eqs. 4,5) provides a very good approximation to the exact Bayesian inference implemented as described in App. A 3. Parameter: $K = 500$, $\theta = 0.02$, $\tau = 10$ year, $\lambda = 10$/year.

FIG. S2: Comparison of the dynamic scaling of the relative cost gap for $c(\tilde{P}_a) = \log \tilde{P}_a$ (colored) and $c(\tilde{P}_a) = 1/\tilde{P}_a$ (grey, see Fig. 2). For both cost functions rescaling parameters as $\lambda t_e/K\theta$ collapses the relative cost gaps for different parameters onto a similar master curve.

FIG. S3: Most of the benefit of memory is still achieved if the immune system has a slightly wrong prior about the antigen distribution sparsity θ. (A) Scaling of mean cost of infections with different priors $\tilde{\theta}$ for a correct $\theta = 0.01$ ($K = 1000$). (B) Relative increase in cost by using a wrong prior vs age.

FIG. S4: The timescale of memory attrition does not need to be finely tuned for near-optimal prediction. Relative cost of infection at age 40 years as a function of the relative attrition timescale $\tilde{\tau}/\tau$ for environments with different correlation times $\tau_c = 2\tau/K\theta$. Parameters: $\alpha = 0.5$, K, θ, λ as in Fig. 4.

FIG. S5: While the optimal update of the receptor frequencies for uncorrelated pathogen environments (blue line) shows a depletion of close but suboptimal receptors such an effect might be reversed for correlated pathogen environments (orange line). We use a Gaussian kernel to describe cross-reactivity $f(|a - r|) = e^{|a-r|^2/(2\sigma^2)}$, and a spacing of receptors 0.5σ. Assuming a change in inferred pathogen frequencies upon encountering pathogen a of e_a for the uncorrelated case (unit vector in a direction) and $f_a = e^{(x-a)/(2\sigma_c^2)}$ with $\sigma_c = 2\sigma$ for the correlated case we obtain the optimal update in terms of receptor frequencies by multiplying the inverse of the cross-reactivity matrix F with the change in frequencies.

FIG. S6: For large changes in inferred frequencies the optimal repertoire can change drastically (blue line). A restriction to local changes yields a repertoire providing close to optimal protection, which exhibits a depletion of good, but suboptimal receptors (orange line). To obtain the global (local) optimal repertoire we use a projected gradient algorithm to minimize the cost function over all receptor distributions (over only the disribution of receptors within 2σ of the pathogen in shape space). We use a logarithmic cost function, $|\boldsymbol{\theta}| = 10$, Gaussian kernel $f(|a - r|) = e^{|a-r|^2/(2\sigma^2)}$, and a spacing of receptors 0.5σ.