Modulatory effects of dynamic fMRI-based neurofeedback on emotion regulation networks in adolescent females

CATHARINA ZICH[1,2], NICOLA JOHNSTONE[3], MICHAEL LÜHRS[4,5], STEPHEN LISK[6], SIMONE P W HALLER[1], ANNALISA LIPP[3], JENNIFER Y F LAU[6], & KATHRIN COHEN KADOSH[1,3]

[1]*Department of Experimental Psychology, University of Oxford, Oxford, OX1 3UD, UK*
[2]*Department of Psychology, University of Oldenburg, 26111 Oldenburg, Germany*
[3]*School of Psychology, University of Surrey, Guildford GU2 7XH, UK*
[4]*Department of Cognitive Neuroscience, Maastricht University, Maastricht, 6211 LK, The Netherlands*
[5]*Department of Research and Development, Brain Innovation, Maastricht, 6229 EV, The Netherlands*
[6]*Psychology Department, Institute of Psychiatry, King's College London, London SE5 8AF, UK*

Corresponding author:
Catharina Zich
Nuffield Department of Clinical Neurosciences
University of Oxford
Oxford, UK
Phone: ++44(0) 1865 283 800
Email: catharina.zich@oldenburg.de

1 ABSTRACT

Research has shown that difficulties with emotion regulation abilities in childhood

and adolescence increase the risk for developing symptoms of mental disorders, e.g

anxiety. We investigated whether functional magnetic resonance imaging (fMRI)-

based neurofeedback (NF) can modulate brain networks supporting emotion

regulation abilities in adolescent females.

We performed three studies (total N=63). We first compared different NF

implementations regarding their effectiveness of modulating prefrontal cortex (PFC)-

amygdala functional connectivity (fc). Further we assessed the effects of fc-NF on

neural measures, emotional/metacognitive measures and their associations. Finally,

we probed the mechanism underlying fc-NF by examining concentrations of

inhibitory and excitatory neurotransmitters.

Results showed that NF implementations differentially modulate PFC-amygdala fc.

Using the most effective NF implementation we observed important relationships

between neural and emotional/metacognitive measures, such as practice-related

change in fc was related with change in thought control ability. Further, we found

that the relationship between state anxiety prior to the MRI session and the effect of

fc-NF was moderated by GABA concentrations in the PFC and anterior cingulate

cortex.

To conclude, we were able to show that fc-NF can be used in adolescent females to

shape neural and emotional/metacognitive measures underlying emotion regulation.

We further show that neurotransmitter concentrations moderate fc-NF-effects.

Keywords
Adolescence, Brain-computer-interface, Connectivity, Emotion regulation, Magnetic
Resonance Spectroscopy, Neurofeedback

1. INTRODUCTION

Adolescence is marked by a multitude of neural, emotional/metacognitive and

behavioural changes, such as functional and structural maturation and

improvements in cognitive abilities and socio-emotional behaviour (Blakemore, 2008;

Burnett et al., 2011; Cohen Kadosh et al., 2013; Linscott and van Os, 2013). It has

been suggested that these complex, transformational processes may shape poor

emotion regulation abilities and increase the risk for developing mental disorders

(Haller et al., 2015, 2013; Keshavan et al., 2014; Paus et al., 2008). Anxiety is one of

the most common and impairing mental disorders in adolescence. Youth with anxiety

experience impairing levels of fears and worries that they are unable to regulate,

which can impact friendships, school performance, and mark the beginning of long-

term mental health difficulties (Beddington et al., 2008; Pine et al., 2001; Trentacosta

and Fine, 2010). Gaining a better understanding of the mechanisms underlying the

successful regulation of emotions during this unique developmental period therefore

represents an important step towards devising efficient, and targeted strategies for

early support and intervention.

In the human brain, the regulation of emotions relies on a network of brain regions,

comprising the prefrontal cortex (PFC) and the amygdala, as well as additional

regions in the orbitofrontal, frontal and cingulate cortex (Kohn et al., 2014; Ochsner

and Gross, 2005). A particular focus has been placed on the PFC-amygdala

relationship, a relationship that manifests in both structural and functional

connections (Banks et al., 2007; Davidson et al., 2000; Kim et al., 2003; Kim and

Whalen, 2009; Kohn et al., 2014; Quirk and Beer, 2006). With regard to the regulatory

aspect of the PFC-amygdala connection, the functional connectivity (fc) between the

two regions has been theorized to reflect top-down PFC regulation of amygdala

reactivity (Hare et al., 2008; Hariri et al., 2003; Kim et al., 2003; Pezawas et al., 2005),

with increased PFC responses leading to decreases in amygdala activation, i.e.

negative fc (Kim et al., 2011). Interestingly, this fc pattern only emerges during

adolescence (Dougherty et al., 2015; Gee et al., 2013; Wu et al., 2016), with research

showing that children at the beginning of adolescence (up until around 10 years of

age) exhibit predominantly positive fc, rather than negative fc ((Dumontheil et al.,

2012, 2010; Silvers et al., 2012), also reviewed in (Ahmed et al., 2015)). What is less

clear however is whether we can intervene to actively shape fc patterns in the

developing brain to improve emotional/metacognitive abilities and thereby affect

behaviour.

The current study aimed to address this question by using real-time functional
magnetic resonance imaging (fMRI)-based neurofeedback (NF). NF provides the user
with real-time information of one's own brain activity, and it has already been shown
to be a promising intervention tool to (re-)shape neural activity in a number of
psychiatric and neurological diseases (Linden, 2012; Lubianiker et al., 2019). In the
case of emotion regulation, NF can aid the modulation of networks underlying this
process, such as PFC-amygdala fc. Previous studies suggest that fMRI-based NF
represents a promising approach to train individuals in the self-modulation of brain
regions or networks (Johnston et al., 2010; Koush et al., 2017, 2013; Scharnowski
and Weiskopf, 2015; Weiskopf et al., 2004; Zilverstand et al., 2014), also in the
emotion processing domain (adults: (Johnston et al., 2010; Koush et al., 2017; Paret
et al., 2014; Zotev et al., 2013); children: (Cohen Kadosh et al., 2016)).
Building upon these studies, we conducted a set of three experiments that
investigated whether fMRI-based fc-NF of PFC-amygdala fc can be used to
modulate neural measures (e.g. change in fc) and emotional/metacognitive measures
(i.e. self-reported measures of anxiety and thought control ability) relevant for
emotion regulation in adolescent females. To achieve this overall objective, we first
contrasted three different NF implementations with regard to their effectiveness in
modulating PFC-amygdala fc (**Experiment 1**). The most effective NF implementation
was then used to assess the effects of one session fc-NF on neural measures and
emotional/metacognitive measures in a larger sample (**Experiment 2**). Finally, we
assessed the effects of a longer fc-NF block in order to further enhance the
effectiveness of the fc-NF and explored how neurotransmitter concentrations in
relevant regions influence the observed effects (**Experiment 3**). Regarding the latter,
we used proton magnetic resonance spectroscopy (^{1}H-MRS) to extract the
neurotransmitter profile from two voxels of interest (VOI), the PFC and the anterior
cingulate cortex (ACC). We choose the PFC because of the key role of γ-
aminobutyric acid (GABA)-ergic neurotransmission within the PFC in balancing
amygdala activity (Constantinidis et al., 2002). Further, we choose the ACC because
of its centrality in the reward processing network, especially for explicit processing
of reward, such as NF (Emmert et al., 2016; Sitaram et al., 2016). For both of these
VOIs we quantified GABA and glutamate, the major inhibitory and excitatory
neurotransmitters in the human brain.

2. MATERIALS AND METHODS

2.1. Experiment 1

Participants

18 female adolescent participants (*M* = 14.83 years; *SD* = 0.99 years) were recruited

from local schools in the Oxfordshire/Gloucestershire area. The single-sex approach

allowed us to minimize variation introduced by differences in hormonal development

and puberty (Goddings et al., 2014, 2012). All participants had normal or corrected-

to-normal vision and reported no history of neurological and psychiatric disorders

(determined via self-report). Informed written consent was obtained from the primary

caregiver and informed written assent was obtained from the adolescent.

Participants received an Amazon voucher (£20) for their participation. The study was

approved by the Central Oxfordshire Ethics Committee (MSD-IDREC-C2-2015-023)

and conducted in accordance with the Declaration of Helsinki. This work was

registered as preclinical trial (ClinicalTrials.gov Identifier: NCT02463136).

Self-report questionnaires

Immediately prior to the MRI session, participants completed several self-report

questionnaires, which will be referred to as *emotional/metacognitive measures*.

Specifically, we assessed psychological variables using the Emotion Regulation

Questionnaire (CERQ) (Derryberry and Reed, 2002), the Mood and Feelings

Questionnaire (MFQ) (Angold et al., 1995), Social Anxiety Scale for Adolescents

(SAS-S) (La Greca and Lopez, 1998), the State-Trait Anxiety Inventory (STAI-T, STAI-

S) (Spielberger et al., 1999), the Thought Control Questionnaire (TCQ) (Wells and

Davies, 1994) and the Thought Control Ability Questionnaire (TCAQ) (Luciano et al.,

2005). These are established self-report measures, which are frequently used in

psychological testing. A subset of the questionnaires (i.e. CERQ, MFQ, SATI-S, TCQ,

TCAQ) were repeated after the MRI session in order to assess changes. Participants

also completed a Demographic and Health Questionnaire and the Wechsler

Abbreviated Scale of Intelligence (Wechsler, 2011).

MRI data acquisition

MRI data acquisition was performed on a 3 T Siemens MAGNETOM Prisma MRI

scanner (Siemens AG, Erlangen, Germany) using a standard 32-channel head matrix

coil. First, a high-resolution structural scan was acquired, which was followed by

functional imaging during the localizer task and the NF task (see **Supplementary**

Methods for details on the MRI sequences).

Localizer task

A modified version of the social scenes task (Haller et al., 2016) was used to identify

the NF regions of interest (ROI), as it was expected that this task activates the key

regions involved in cognitive and emotional appraisal and reappraisal. The localizer

task lasted 8.9 min. and comprised 30 trials. Each trial started with a social scene

presented for four volumes (3.73 s). Scenes depicted negative, rejecting social

situations viewed from the perspective of a female protagonist depicted from the

back. Participants were instructed to interpret the scene freely (appraisal). This was

followed by a positively valanced interpretative statement (4 volumes, 3.73 s), after

which the same scene was shown again for four volumes (3.73 s). For the duration

of the second presentation, participants were instructed to reappraise the scene

based on an interpretative statement (reappraisal). Finally, participants were asked

to rate how much they were able to change, i.e. reappraise, their thoughts and

feelings from the first to the second presentation of the scene. Participants indicated

their perceived change on a Likert scale ranging from no change (1) to much change

(4) via button press on each trial. A fixation cross was presented for one volume

between two trials. Stimulus presentation was controlled via BrainStim 1.1.0.1 (open

source   stimulation   software,   Maastricht   University,

http://svengijsen.github.io/BrainStim/). Participants were encouraged to use these

emotional reappraisal strategies during the NF tasks.

Neurofeedback task

The NF task consisted of four identical runs lasting 4.8 min each. Stimulus

presentation was controlled with BrainStim 1.1.0.1. Each run started with a fixation

cross that was displayed for 20 volumes (18.66 s), followed by seven fc-NF mini-

blocks (20 volumes each) and seven no-NF mini-blocks (20 volumes each) that were

presented in alternating order, with the start condition being randomized and

counterbalanced across individuals. During both fc-NF and no-NF mini-blocks a ten-

segment thermometer was presented at the centre of the screen on a dark grey

background. During the no-NF mini-blocks, the 'temperature' of the thermometer

was frozen at the sixth segment (i.e. 6/10) throughout the mini-block (**Fig. 1a**). During

the fc-NF mini-blocks, the thermometer featured a green frame and the 'temperature'

was a direct reflection of PFC-amygdala fc and was updates with every TR (see

section 'Online MRI data analysis' for more details). All participants were asked to

up-regulate the thermometer by controlling their thoughts and feelings and by

revisiting emotional reappraisal strategies as practised in the localizer task (see

Supplementary Methods for more details on the instructions).

We evaluated three different NF implementations with regard to their effectiveness

of modulating PFC-amygdala fc (N = 6 per implementation). The three different NF

implementations only differed in the upper value of the thermometer (the lower value

of the thermometer was set to zero). The upper value of the thermometer was set to

-1 (maximal negative correlation) for the *negative NF implementation*, which

reinforced negative fc; to -0.3[†] for the *weighted negative NF implementation*, which

reinforced negative fc; and to +1 (maximal positive correlation) for the *positive NF*

implementation, which reinforced positive fc. To illustrate this: if the thermometer

changes from 2/10 to 3/10 segments, this corresponds to a more negative correlation

by 0.1 for the negative NF implementation, a more negative correlation by 0.03 for

the weighted negative NF implementation, and a change towards a more positive

correlation by 0.1 for the positive NF implementation.

Online MRI data analysis

To enable real-time fMRI-based NF, MR images were passed from the MRI console

computer to the real-time computer via a direct TCP/IP network link using the Server

Message Block (SMB) network layer. Following completion of the structural scan,

anatomical images were processed using BrainVoyager QX 2.8.2 (Brain Innovation,

Maastricht, The Netherlands). Each 3D volume was corrected for B0 in-

homogeneities (4-cycle bias field estimation), followed by brain extraction from the

skull and separating bone and cerebro-spinal fluid (2-cycle iteration).

Functional images obtained during the localizer and the NF runs were processed in

real-time using Turbo-BrainVoyager 3.2 (Brain Innovation, Maastricht, The

Netherlands). To correct head motion, each volume was realigned to the first volume

of the localizer. Each realigned volume was smoothed with a three-dimensional

Gaussian kernel of 8 mm full-width-half-maximum. After completion of the localizer,

three ROIs (voxel size = 12 mm^3, 6 x 6 x 6) were selected. To this end, GLM t-statistics

of the brain activity during the localizer task, i.e. the sum of the three contrasts:

appraisal > fixation, reappraisal > fixation, reappraisal > appraisal (threshold t = 3),

---

[†] Based on pilot data (N = 5). Cumulative distribution function of PFC-amygdala fc, i.e. partial correlations r_p between the PFC, amygdala and corticospinal tract.

was projected onto the processed structural scan. The local maximum of the t-

statistics within the left dorsolateral and medial PFC constituted the centre of the

PFC ROI. Similarly, the local maximum of the t-statistics within the left amygdala

constituted the centre of the amygdala ROI. A ROI in the left corticospinal tract (CST)

served as control ROI[‡]. Probability maps of the selected ROIs across subjects are

depicted in **Fig. 1c.**

During the NF task, PFC-amygdala fc was calculated in real-time. PFC-amygdala fc

was defined as the partial correlation r_p between PFC and amygdala activity, while

controlling for CST 'activity'. As shown previously, partial correlation analysis can be

used to quantify fc between areas while controlling for noise from a task-unrelated

region (Dawson et al., 2016). Partial correlations were based on a moving window,

which was updated with every incoming volume. The length of the correlation

window was 20 volumes (see **Supplementary Discussion**). Therefore, the ratio of

fc-NF to no-NF volumes in the correlation window changed with every incoming TR

(**Fig. 1b**). To illustrate this: for the fc-NF mini-blocks, the ratio of fc-NF : no-NF

volumes in the correlation window was 1:19 at the first volume of the mini-block, 2:18

at the second volume of the mini-block and 20:0 at the last volume of the mini-block,

and vice versa for no-NF mini-blocks. Calculations were performed using a custom-

made plugin for Turbo-BrainVoyager, which also provided a direct TCP/IP based link

between the real-time analysis software and the stimulus application BrainStim.

PFC-amygdala fc was displayed via thermometer during the fc-NF mini-blocks.

Offline MRI data analysis

We also we analysed the MRI data offline. While online and offline analysis of the

MRI data follow a similar processing pipeline, offline algorithms tend to be more

robust. Offline analysis of the MRI data was performed using SPM12 (FIL, Wellcome

Trust Centre for Neuroimage, UCL, London, UK). Here head motion was corrected

by realigning the functional time series of the localizer or the NF runs to its first

volume. Due to motion artefacts (i.e. motion exceeded 3 mm on any axis) three

individuals (i.e. one individual per group) were excluded from all analyses. Each

individual's structural image was registered to their mean functional image and

segmented, in order to normalize structural and functional images to the Montreal

---

[‡] We note that the current study focused on left-lateralized ROIs as those were reliably activated in all subjects in the localiser task, whereby a lateralization in favour of the left hemisphere is in line with the literature (Ochsner and Gross, 2005).

Neurological Institute (MNI) template. Finally, normalized functional images were

smoothed with a three-dimensional Gaussian kernel of 8 mm full-width-half-

maximum. To increase the consistency between online and offline analysis we used

the same ROIs. Therefore, the ROIs defined in Turbo-BrainVoyager during the MRI

session were transformed from DICOM to NIFTI format, from radiological to

neurological convention and from voxel to mm space. Subsequently, ROIs were

normalized to the MNI template by applying the same transformation matrix used for

the subject-specific normalization of structural and functional images. Time series

were extracted from the three ROIs using MarsBaR 0.44 (Brett et al., 2002) and PFC-

amygdala fc calculated using the same procedure as during real-time NF, i.e. partial

correlation r_p between PFC and amygdala activity, while controlling for CST 'activity'.

Statistical analysis

To identify the most effective NF implementation we evaluated the difference

between no-NF and fc-NF for each NF implementation. The primary measure for this

comparison constitutes PFC-amygdala fc, i.e. partial correlations r_p. Fc was

averaged at each of the 20 volumes (i.e. length of a mini-block) across the seven

mini-blocks within one condition (fc-NF, no-NF), NF runs (1, 2, 3, 4) and individuals

within one NF implementation (negative NF implementation, weighted negative NF

implementation, positive NF implementation). The relationship of fc-NF and fc was

assessed by correlating (Pearson's *r* correlations as implemented in SPSS version

25; SPSS Inc, Chicago, IL, USA) the average fc at each of the 20 volumes with the

number of fc-NF volumes within the correlation window ($N = 40$, i.e. 20 volumes per

mini-block). To statistically compare these Pearson's *r* correlations between the

different NF implementations, Fisher's z-transformation was applied to each

correlation coefficient, resulting in normally distributed values *r'* with standard errors

$s_{r'}$. The null hypotheses ($r'1$-$r'2 = 0$) were tested in R(psych) (Revelle, 2015) using

Student *t* test (Howell, 2011).

The same analysis was conducted to assess the frequency of negative partial

correlations (secondary measure) and the frequency of scenarios describing different

patterns of activation changes (**Supplementary Fig. 5**).

Fig. 1 Schematic representation of the experimental procedures (**Experiment 1** and **2** in green and **Experiment 3** in orange). **(a)** Experimental timelines. To activate the key regions involved in emotion regulation participants performed the social scenes task (localizer). Based on an individual's localizer activity, we then defined three regions of interests (ROIs): left prefrontal cortex (PFC), left amygdala, left corticospinal tract (CST). Each NF run comprised fc-NF mini-blocks (20 volumes in **Experiments 1 and 2**; 40 volumes in **Experiment 3**) and no-NF mini-blocks (20 volumes), presented in alternating order. **(b)** Detailed view of the procedure during the NF task. NF was based on partial correlations r_p between the PFC and amygdala activity, while controlling for CST 'activity', whereby partial correlations were obtained from a moving window comprising 20 subsequent volumes, which was updated with every incoming volume. Thus, for **Experiments 1** and **2**, the ratio of fc-NF : no-NF volumes in the correlation window is 1:19 at the first volume of the mini-block, 2:18 at the second volume of the mini-block and 20:0 at the last volume of the mini-block, and vice versa

for no-NF mini-blocks. Note that due to the extended window length for the fc-NF mini-blocks
in **Experiment 3**, the ratio of fc-NF : no-NF volumes in the correlation window is 20:0 for
twenty volumes. Number of fc-NF volumes in the correlation window (N of fc-NF Volumes in
r_p) is illustrated in form of a grey shape. **(c)** Probabilistic maps of subject-specific ROIs (i.e.
PFC and amygdala) superimposed on the groups' average structural image in MNI space.
The respective centre of the PFC probabilistic maps (Experiment 1 and 2: x=-20, y=50, z=24;
Experiment 3: x=-26, y=47, z=22) are classified as Brodmann area 10 within the left middle
frontal gyrus. The respective centre of the amygdala probabilistic maps (Experiment 1 and 2:
x=-17, y=-1, z=-14; Experiment 3: x=-16, y=-3, z=-15) are classified as left amygdala (Eickhoff
et al., 2005).
11

2.2. Experiment 2

This study comprises data from 30 adolescent females (*M* = 15.20 years; *SD* = 1.10

years; 25 naïve individuals and the 5 individuals tested using the weighted negative

NF implementation in **Experiment 1**). The recruitment procedure was identical to

Experiment 1. The 'Self-report questionnaires', 'MRI data acquisition', 'Localizer

task', 'Neurofeedback task' as well as the 'Online MRI data analysis' and 'Offline MRI

data analysis' were identical to **Experiment 1**. However, all individuals received NF

using the weighted negative NF implementation. Due to motion artefacts (motion

exceeded 3 mm on any axis) datasets from three individuals were excluded from all

analyses.

Statistical analysis

Firstly, we performed the same analysis as in **Experiment 1**.

We extended this by investigating practice-related change in

emotional/metacognitive measures and neural measures. *Practice-related change in*

emotional/metacognitive measures was defined as difference between

emotional/metacognitive measures obtained before and after the MRI session and

assessed using paired samples *t* test (SPSS). Practice-related change in neural

measures, i.e. *practice-related change in fc*, was defined as the slope of the linear

regression of the total fc across runs (**Supplementary Fig. 4a**) and was tested using

a one-sample *t* test (SPSS).

Moreover, we performed correlation analyses between different

emotional/metacognitive and neural measures. Relevant emotional/metacognitive

measures were practice-related change in emotional/metacognitive measures (as

defined in the previous paragraph) and *initial emotional/metacognitive measures*, i.e.

measures obtained before the MRI session. Relevant neural measures were practice-

related change in fc (as defined in the previous paragraph), initial fc and fc-NF-effect.

Initial fc was defined as the average fc of the first two mini-blocks (**Supplementary**

Fig. 4b). *Fc-NF-effect* was defined as the difference between no-NF and fc-NF (**Supplementary Fig. 4c**). Correlation analysis was performed using Bayesian inference (JASP, JASP Team 2019, version 0.8.1.2) with default priors. Results are reported for the whole sample and after outlier removal. Outliers were identified for each correlation separately by bootstrapping the Mahalanobis distance (Schwarzkopf et al., 2012).

Missing values analysis revealed that 2.56% of the self-report questionnaire data were missing. We performed Little's test of Missing Completely at Random (MCAR) (Little, 1988), as implemented in SPSS. MCAR was not significant ($X^2_{(35, N = 27)} = 41.74$, $p = .201$), i.e. there is no evidence to suggest that the data were not MCAR. As such, pairwise deletion was used in the statistical analysis.

2.3. Experiment 3

20 naïve female adolescent participants (*M* age = 15.85 years; *SD* = 1.04 years) were tested using the weighted negative NF implementation as described in **Experiment 1**. The recruitment procedure was identical to **Experiments 1** and **2**.

The 'Self-report questionnaires' were identical to **Experiments 1** and **2**. 'MRI data acquisition' was similar to **Experiments 1** and **2**, the only modification beeing the number of volumes during the localizer and the NF runs (see **Supplementary Methods**). The 'Localizer task' was similar to **Experiments 1** and **2**, however, only half of the trials were performed (i.e. 15 trials, 4.7 min.).

The 'Neurofeedback task' was similar to **Experiments 1 and 2**, with one critical variation: we lengthened the fc-NF mini-blocks to 40 volumes while the no-NF mini-blocks remained 20 volumes. We collected data from 3 identical runs, each comprising five fc-NF mini-blocks and five no-NF mini-blocks. The 'Online MRI data analysis' and 'Offline MRI data analysis' was similar to **Experiments 2** taking into account the change from 20 to 40 volumes per fc-NF mini-block. Moreover, we collected MRS data from two VOIs, the PFC and the ACC. See **Supplementary Methods** for details on MRS data acquisition. Due to motion artefacts (motion exceeded 3 mm on any axis) the dataset from one individual was excluded from all analyses.

MRS analysis

MRS post-processing was performed using the MATLAB (Mathworks, Natick, MA) toolbox MRspa (version 1.5f, https://www.cmrr.umn.edu/downloads/mrspa/).

Motion corrupted spectral averages were removed and frequency and phase drifts

corrected before spectral averaging. Averaged spectra were quantified in LCmodel

(Provencher, 2001, 1993) in reference to a simulated basis set. GABA and glutamate

were quantified in reference to total Creatine (Creatine + Phosphocreatine, t/CrPCr)

for each VOI. Cramér–Rao lower bounds were <20% for GABA (t/CrPr) indicating

reliable estimates extracted from spectra for all but two values in the PFC which were

22% and 27%. Both were retained for analysis. GABA (t/CrPr) concentrations were

similar in each VOI ($t_{(18)}$ = 0.89, p = .386; ACC: M = 5.811%, SD = 1.40%, PFC: M =

5.29%, SD = 2.25%).   Cramér–Rao lower bounds were all <20% for Glutamate

(t/CrPr) indicating reliable estimates extracted from spectra. Glutamate (t/CrPr)

concentrations differed by VOI ($t_{(18)}$ = 4.85, p < .001), with estimates higher in the

ACC (M = 10.57%, SD = 0.76%), than the PFC (M = 9.34%, SD = 0.75%).

Statistical analysis

We performed the same statistical analysis as in **Experiment 1** and **2**. In addition,

we conducted a moderation analysis following mean centering using the SPSS add-

on tool PROCESS macro (version 3.3) (Hayes, 2018). Significant interactions were

followed-up with simple slope analysis.

3. RESULTS

3.1. Experiment 1

The aim of **Experiment 1** was to compare different NF implementations regarding

their effectiveness of modulating PFC-amygdala fc. The primary measure for this

comparison, i.e. PFC-amygdala fc, is illustrated in **Supplementary Fig. 1a**.

The relationship of fc-NF and fc was assessed by correlating the average fc at each

of the 20 volumes with the number of fc-NF volumes within the respective correlation

window (**Supplementary Fig. 1b**). For the weighted negative NF implementation, a

significant negative relationship was found ($r_{(38)}$ = -0.72, p < 0.001), i.e. the more fc-

NF volumes were included in the correlation window, the less positive the resulting

partial correlation. The opposite relationship was found for the negative NF

implementation ($r_{(38)}$ = 0.37, p = 0.02), i.e. the more fc-NF volumes were included in

the correlation window, the more positive the resulting partial correlation. No

significant relationship was observed for the positive NF implementation ($r_{(38)}$ = 0.22,

p = 0.16).

Finally, the relationship between the average fc at each volume and the number of

fc-NF volumes within the correlation window observed for the weighted negative fc-

NF implementation was significantly different from the other two implementations

(both p's < 0.001), which did not differ from each other (p = 0.48). The picture is

similar for the secondary measure, i.e. the frequency of negative partial correlations

(see **Supplementary Results, Supplementary Fig. 2, Supplementary Fig. 3** for

details).

In sum, these findings demonstrate the superiority of the weighted negative NF

implementation in modulating PFC-amygdala fc towards more negative fc[§].

3.2. Experiment 2

The aim of **Experiment 2** was to assess the effects of one session fc-NF (weighted

negative NF implementation) on neural measures, emotional/metacognitive

measures and their associations in a larger sample. The fc-NF-effects observed in

Experiment 1 were replicated, that is, the number of fc-NF volumes within the

correlation window was negatively related with the average fc at each volume (**Fig.**

2a) and positively related with the frequency of negative partial correlations at each

volume (**Fig. 2b**).

[§] Note that the cognitive and psychological measures were comparable between the three groups (see **Supplementary Table 1**).

We further investigated practice-related change on emotional/metacognitive

measures and neural measures for the whole sample. At emotional/metacognitive

level no significant changes between measures obtained before and after the MRI

session were observed (**Supplementary Table 2**). Similarly, at neural level no

significant practice-related change in fc could be observed ($M = .005$, $SD = .035$, p

$> .01$). Thirteen individuals showed a negative slope (i.e. change in the desired

direction, $M = -.023$, $SD = .019$), fourteen individuals exhibited a positive slope ($M =$

$.031$, $SD = .025$). Note that the emotional/metacognitive measures were comparable

between these two subsamples (see **Supplementary Table 3** and **Supplementary**

Table 4).

Correlation analysis

We first investigated whether practice-related change in emotional/metacognitive

measures and practice-related change in fc were related. We found that practice-

related change in fc was negatively related with change in TCAQ (**Fig. 3a** top; $r_{(21)} = -$

$.37$, $B_{10} = 1.03$; after the removal of one outlier: $r_{(20)} = -.58$, $B_{10} = 11.00$). Hence, the

larger the practice-related change towards negative fc, the better the thought control

ability after the MRI session, when compared to the before the MRI session.

Moreover, we were interested whether, neural measures could be predicted by initial

emotional/metacognitive measures, i.e. measures obtained before the MRI session.

Correlation analysis revealed that initial fc was positively related with STAI-T (**Fig. 3b**

top; $r_{(25)} = .36$, $B_{10} = 1.17$; after the removal of two outliers: $r_{(23)} = .59$, $B_{10} = 23.66$) and

negatively related with thought control ability (TCAQ: **Fig. 3c** top; $r_{(21)} = -.36$, $B_{10} =$

1.03; after the removal of two outliers: $r_{(19)} = -.66$, $B_{10} = 35.01$). In other words, initial

fc was more negative in individuals with lower trait anxiety and better ability to control

thoughts before the MRI session. Further, we found that STAI-S and fc-NF-effect

were negatively related (**Fig. 3d** top; $r_{(22)} = -.41$, $B_{10} = 1.67$; after the removal of one

outlier: $r_{(21)} = -.55$, $B_{10} = 7.78$). This suggests that individuals with lower state anxiety

before the MRI session showed a larger difference between fc-NF and no-NF than

individuals with higher state anxiety.

Fig. 2 Difference in average functional connectivity (fc) and frequency of negative partial correlations (N of negative r_p) between fc-NF (red) and no-NF (blue) for **Experiment 2** (green frame) and **Experiment 3** (orange frame). **(a, c)** Top: Means and standard error for average fc, i.e. partial correlations r_p, for fc-NF and no-NF at each volume. Fc is averaged across mini-blocks within one condition (fc-NF, no-NF), NF runs and individuals. Number of fc-NF volumes in the correlation window (N of fc-NF Volumes in r_p) is illustrated in form of a grey shape. Bottom: Relationship between average fc at each volume and the number of fc-NF volumes within the correlation window. **(b, d)** Same as (a, c), but for the frequency of negative partial correlations.

Fig. 3 Associations between the emotional/metacognitive and neural measures for **Experiment 2** (top, green) and **Experiment 3** (bottom, orange). (**a**) Association between change in fc and change in thought control ability (**b,c**) Association between the initial fc and trait anxiety (STAI-T) as well as thought control ability (TCAQ) obtained before the MRI session. (**d**) Association between fc-NF-effect and state anxiety (STAI-S) obtained before the MRI session. Counter lines represent the bootstrapped Mahalanobis distance from the bivariate mean in steps of six squared units. Open circles represent outliers and the solid line is the regression over the data after outlier removal. Direction of solid arrows indicate levels related with higher anxiety. Direction of dashed arrows indicate the desired direction of practice-related change and fc-NF-effect.

3.3. Experiment 3

The aim of **Experiment 3** was three-fold: to assess the effect of longer fc-NF mini-

blocks, to replicate the relationships between the neural and

emotional/metacognitive measures as observed in **Experiment 2**, and to assess how

neurotransmitter concentrations in NF relevant regions influence these relationships.

As in **Experiment 1** and **2**, the number of fc-NF volumes within the correlation

window was negatively related with the average fc at each volume (**Fig. 2c**) and

positively related with the frequency of negative partial correlations (**Fig. 2d**). We

further found that these two measures, i.e. average fc and the frequency of negative

partial correlations, plateaued when the fc-NF was estimated consecutively from

volumes stemming from fc-NF mini-blocks only (i.e. the ratio of fc-NF : no-NF

volumes in the correlation window was 20:0).

Similar to **Experiment 2**, no practice-related change in emotional/metacognitive

measures (**Supplementary Table 5**) or practice-related change in fc could be

observed at group level (M = -.006, SD = .061, p > .01). Ten individuals showed a

negative slope (i.e. change in the desired direction, M = -.057, SD = .022), nine

individuals exhibited a positive slope (M = .052, SD = .026).

Correlation analysis

We first computed the correlation coefficients for the relationships between the

neural and emotional/metacognitive measures that were found to be significant in

Experiment 2. Again, all four correlations were found to be significant. Practice-

related change in fc was negatively related with change in thought control ability

(TCAQ; **Fig. 3a** bottom; $r_{(17)}$ = -.57, B_{10} = 6.04; after the removal of two outliers: $r_{(15)}$ =

-.66, B_{10} = 13.27). Initial fc was positively related with trait anxiety (STAIT T; **Fig. 3b**

bottom; $r_{(17)}$ = .43, B_{10} = 1.32; after the removal of one outlier: $r_{(16)}$ = .56, B_{10} = 4.31)

and negatively related with thought control ability (TCAQ: **Fig. 3c** bottom; $r_{(17)}$ = -.55,

B_{10} = 4.61; after the removal of three outliers: $r_{(14)}$ = -.69, B_{10} = 16.54). The effect of

fc-NF was negatively related to state anxiety (STAI-S; **Fig. 3d** bottom; $r_{(17)}$ = -.69, B_{10}

= 40.49; after the removal of three outliers: $r_{(14)}$ = -.80, B_{10} = 170.20).

In addition, **Fig. 4** shows the correlation matrix for the relevant

emotional/metacognitive measures, neural measures and neurotransmitter

concentrations.

Moderation analysis

We conducted moderation analyses to assess if neurotransmitter concentrations, i.e.

glutamate and GABA, in relevant regions influence the significant relationships found

between neural measures and emotional/metacognitive measures. PFC GABA and

ACC GABA concentrations moderate the relationship between state anxiety before

the MRI session and fc-NF-effect (model 3, $F_{(7,11)} = 3.65$, $p = .028$, $R^2 = .70$, **Fig. 5a**).

We found STAI-S ($b = -.009$ $t_{(11)} = -3.514$, $p = .005$, 95% CI [-.015, -.003]) and PFC

GABA concentrations ($b = .036$ $t_{(11)} = 2.635$, $p = .023$, 95% CI [.006, .066]) to be

significant predictors as well as the interaction term between PFC GABA and ACC

GABA concentrations to be significant ($b = -.037$ $t_{(11)} = -2.394$, $p = .036$, 95% CI [-

.071, -.003], **Table 1**)**.**

To further explore the nature of the interaction we performed a simple slopes

analysis. We found that the relationship between STAI-S and the fc-NF-effect

reached significance at medium ACC GABA and medium PFC GABA concentrations

($b = -.009$ $t_{(11)} = -3.513$, $p = .005$, 95% CI [-.015, -.003]) and is at trend level for

medium ACC GABA and low PFC GABA concentrations ($b = -.022$, $t_{(11)} = -1.894$, p

$= .085$, 95% CI [-.048, .004]), whereas no such effect was found at medium ACC

GABA and high PFC GABA concentrations ($b = .004$ $t_{(11)} = .324$, $p = .752$, 95% CI [-

.023, .031]). No significant effects were observed for low and high ACC GABA (all p's

$> .1$). To test the specificity of this effect we ran the same model using glutamate

only (i.e. PFC glutamate and ACC glutamate concentrations) and a combination of

GABA and glutamate (e.g. PFC GABA and ACC glutamate concentrations). In none

of these cases the overall model was found to be significant (all p's $> .1$).

Fig. 4 Correlation matrix for relevant emotional/metacognitive measures, neural measures
and neurotransmitter concentrations. Bayesian correlation values are shown after outlier
removal using Mahalanobis distance.

1
2 **Fig. 5** Moderation model and results. Moderation of the effect of state anxiety before the
3 MRI session on the fc-NF-effect by neurotransmitter concentrations in NF relevant regions,
4 depicted as a conceptual diagram (**a**) and a statistical diagram (**b**). (**c**) Simple slopes for
5 STAI-S predicting fc-NF-effect for different levels (+1 *SD*, 0, -1 *SD*) of PFC GABA and ACC
6 GABA concentrations.
7
8 **Table 1** Estimating the influence of neurotransmitter concentrations in NF relevant regions
9 on the relationship between state anxiety before the MRI session and fc-NF-effect.

		Coeff	*SE*	*t*	*p*	LLCI	ULCI
Constant	i_Y	-.028	.020	-1.438	.178	-.072	.015
STAI-S (*X*)	b_1	-.009	.003	-3.514	**.005**	-.015	-.003
PFC GABA (*W*)	b_2	.036	.014	2.635	**.023**	.006	.066
XW	b_3	.006	.005	1.121	.286	-.006	.017
ACC GABA (*Z*)	b_4	-.029	.025	-1.185	.261	-.084	.025
XZ	b_5	-.001	.005	-.233	.820	-.011	.009
WZ	b_6	-.037	.015	-2.394	**.036**	-.071	-.003
XWZ	b_7	.002	.008	-.231	.822	-.018	.015

4. DISCUSSION

Here we aimed at comparing different NF implementations with regard to their

effectiveness in modulating PFC-amygdala fc (**Experiment 1**) and at assessing the

effects of the most effective NF implementation on neural measures and

emotional/metacognitive measures, and their associations (**Experiment 2** and **3**). We

further assessed the effect of a longer fc-NF block and whether neurotransmitter

concentrations in NF relevant regions influence fc-NF-effects or practice-related

changes (**Experiment 3**).

Differential effects of NF implementations

In **Experiment 1**, we compared three NF implementations and found that the

weighted negative NF implementation was most effective in achieving a more

negative fc when comparing fc-NF to no-NF. Considering that younger individuals

and anxious individuals of all ages exhibit generally lower levels of negative PFC-

amygdala fc, or even positive PFC-amygdala fc, it is plausible to suggest that a NF

implementation ranging from zero (no correlation) to minus one (maximum negative

correlation) was too unspecific to benefit the individuals (Hattie and Timperley, 2007;

Lotte et al., 2013). In other words, the naturally more rarely occurring negative

connectivity pattern does not receive enough reinforcement to allow the individuals

to learn and improve. This finding holds important implications for the development

of NF interventions, as it suggests that individually tailored NF implementations

should enhance the modulation of task-specific activity patterns even more.

Absence of significant effects at group level

In **Experiment 2** and **3** we did not find significant changes in

emotional/metacognitive measures or in the change in fc at group level. This could

be due to several reasons, such as the large inter-individual differences in neural and

emotional/metacognitive measures, or the relatively short practice time. We note

though that the relationship between the amount of practice and manifestation of

emotional/metacognitive changes is not yet clear, and may depend on various

aspects (e.g. population, task, NF implementation). While Zilverstand and colleagues

(Zilverstand et al., 2015) showed clinically relevant effects with only one NF session,

others argue that multiple sessions may be necessary (Linden et al., 2012; Scheinost

et al., 2013). For one, this shows that there is an urgent need to actively research the

optimal dose of NF practice. Moreover, it may be beneficial to link the NF practice

more closely to cognitive-behavioural approaches, such as cognitive behavioural

therapy.

Relationships between emotional/metacognitive and neural measures

The significant associations between emotional/metacognitive and neural measures

that we observed in **Experiment 2** were replicated in an independent sample in

Experiment 3. The observed correlational results can be subdivided into effects of

change and predictive effects. Correlations representing effects of change

demonstrated that practice-related change in fc is related to change in thought

control abilities. Effects of change show that we can use fc-NF not only to selectively

modulate specific brain network connections, but also that these neural changes

relate to changes in emotional/metacognitive measures, which further supports the

effectiveness of the NF implementation.

Results depicting predictive effects found that lower trait anxiety predicted more

negative fc. This finding is consistent with previous studies (Delli Pizzi et al., 2017;

Kim et al., 2011). Our results further indicate that thought control abilities were

negatively related to the initial fc, i.e. the better the thought control ability, the more

negative the initial fc. This is not surprising given the negative relationship between

trait anxiety and thought control abilities. Together, these results highlight that levels

of anxiety and thought control abilities can be used to predict PFC-amygdala fc

during emotion regulation.

Our results also provide some insights into individual differences in fc-NF-effects.

Specifically, we found that state anxiety prior to the MRI session was negatively

related with the fc-NF-effect, i.e. the lower the state anxiety, the higher the effect of

fc-NF on fc. This highlights the importance of an individual's state prior to the NF.

Future studies aiming at increasing NF success should therefore ensure that

individuals, especially those with high levels of state anxiety, are relaxed and

comfortable prior to the NF session. It is fair to assume that an individual's state not

only plays a pivotal role prior to the NF session, but also during the NF session. One

approach could therefore be to only initiate the next trial if the individual is in a 'good'

state (i.e. relaxed, low state anxiety), which could be assessed either behaviourally

or by means of an identified a biomarker (Meinel et al., 2016).

Moderation through neurotransmitter concentrations

In **Experiment 3**, we extended our protocol to obtain glutamate and GABA
concentrations from the PFC and ACC. We choose the PFC due to the key role of γ-
aminobutyric acid (GABA)-ergic neurotransmission within the PFC in balancing the
amygdala activity (Constantinidis et al., 2002), whereas the ACC was chosen due to
its centrality for explicit processing of reward, such as NF (Emmert et al., 2016;
Sitaram et al., 2016). Our results showed that the relationship between state anxiety
(STAI-S) before the MRI session and fc-NF-effect (**Fig. 3d**) is moderated by GABA
concentrations in the ACC and PFC. Simple slope analysis revealed that the
relationship between STAI-S and the fc-NF-effect reached significance at medium
ACC GABA and medium PFC GABA concentrations and is at trend level for medium
ACC GABA and low PFC GABA concentrations. This demonstrates that medium
ACC GABA concentrations constitutes an optimal level of inhibition. Due to the
central role of the ACC for explicit reward processing (Emmert et al., 2016; Sitaram
et al., 2016), this result suggests that medium inhibition levels in the ACC are ideal
for explicit reward processing, such as NF. Our results further demonstrate that
medium and low PFC GABA concentrations, but not high PFC GABA concentrations,
promote the relationship between STAI-S and fc-NF-effect. This is in agreement with
animal studies demonstrating that higher GABA concentrations in the ventromedial
PFC reduces GABA-ergic inhibition of the amygdala, promoting its hyperactivity
(Akirav and Maroun, 2007; Chefer et al., 2011; Courtin et al., 2014; Gauthier and
Nuss, 2015). Delli Pizzi and colleagues provided supporting evidence for this in
humans, i.e. they found a positive relationship between medial PFC GABA
concentrations and resting state ventromedial PFC-amygdala fc (Delli Pizzi et al.,
2017). This suggests that if high PFC GABA concentrations imply higher amygdala
activity, this could in turn result in a more positive PFC-amygdala fc and
consequently reduce fc-NF-effects. In sum, our results provide first empirical
evidence for the relevance of neurotransmitter concentrations for the effectiveness
of fc-NF.
Given the importance of the optimal neurotransmitter concentrations for fc-NF to be
effective approaches to alter neurotransmitter concentrations prior to NF should be
explored. However, the use of GABA-ergic drugs during childhood and adolescence
is controversial due to the risk of developing dependence and severe adverse effects
(Sidorchuk et al., 2018). Yet other means that alter GABA-ergic activity, such as yoga
(Streeter et al., 2010, 2007), may be adopted. Such studies would also shed further
light on the causal interpretation.

A pressing question is whether the observed moderation effect is specific to the
domain of emotion regulation, i.e. domain specific, or universal. Within our
moderation model ACC GABA concentrations (Z) and fc-NF-effects (Y) constitute NF
specific variables, whereas STAI-T (X) and PFC GABA concentrations (Z) are domain
specific variables. It is thus reasonable to assume that the model is still valid if X and
Z are replaced by corresponding variables in the different domain, such as residual
motor function in the affected limb as X and GABA concertation from the
sensorimotor area of the affected hemisphere as Z for motor control following post-
stroke. It can further be assumed that this effect may not specific to fc-NF only, but
also apply for activity-based NF. Empirical research and computational modelling
should be consulted to answer the question wheatear our results are domain specific
or universal.
13
Implications for translational, clinical application
The results from the current series of experiments have provided strong evidence
that fc-NF is feasible in adolescent females with different anxiety levels. We further
found that practice-related changes in fc is related to changes in thought control
ability. Thus, on a broader scale, our results provide first evidence for a possible
clinical application that aims to shape emerging fc patterns non-invasively in the
developing brain(Cohen Kadosh et al., 2013). This is significant given that one in four
children have increased levels of worry and fear as they enter adolescence and
paediatric anxiety predicts lifelong mental disorders (Keshavan et al., 2014). Aberrant
emotional regulation strategies arising in anxiety mirror fc patterns in younger
individuals where lower levels of negative PFC-amygdala fc or even positive PFC-
amygdala fc are exhibited (Kim et al., 2011).
Further research is now necessary to evaluate the effects of multi-session protocols
in clinical samples using both subjective and objective clinical outcome measures.
28
Limitations
The following limitations merit comment: Despite converging results across the
experiments, multi-centre studies are now necessary to examine the effect of
psychosocial and socioeconomic factors. Furthermore, we acknowledge that we
performed a relatively large number of correlations, whereby we have focussed on
priori hypotheses, i.e. correlations with measures of anxiety and thought control
ability, and, crucially, used Bayesian inference.

While we underline the acquisition of neurotransmitter levels as a strength of this

study, we emphasize that single-voxel MRS has limited spatial resolution. Recent

advances in magnetic resonance spectroscopic imaging (MRSI) have the potential

to overcome the shortcomings related to single-voxel MRS in future (Steel et al.,

2018).

Conclusions

Our results showed that NF implementations differentially modulate PFC-amygdala

fc. Using the most effective NF implementation in a larger sample yielded important

associations between neural measures and emotional/metacognitive measures,

such as practice-related change in fc was related with change in thought control

ability. Further, we found that the relationship between state anxiety and the effect

of fc-NF was moderated by GABA concentrations in the PFC and ACC. Future

studies that investigate the effects of multi-session protocols in clinical samples and

whether the moderation model is domain specific or universal are recommended.

ACKNOWLEDGEMENTS

We thank Dr. Ralf Veit for discussions regarding the offline MRI data analysis, Prof.

Roi Cohen Kadosh for discussions regarding the moderation analysis and Prof. David

Linden for comments on the manuscript.

We thank Verena Koppe for her valuable contribution to data management as well

as Samantha Clarkstone, Kerenza Hood, Rachel McNamara, Rebecca Playle,

Vincent Poile, Elizabeth Randell and Gareth Watson (Centre for Trials Research,

College of Biomedical & Life Sciences, Cardiff University, UK), and all BRAINTRAIN

coordinators and collaborators for helpful and inspiring discussions.

This work has received funding from the European Union's Seventh Framework

Programme for research, technological development and demonstration under grant

agreement no. 602186 and registered as preclinical trial #NCT02463136.

DISCLOSURES

ML was employee of Brain Innovation B.V., Maastricht, The Netherlands. The other

authors disclose no financial or non-financial competing interests or potential

conflicts of interests.

AUTHOR CONTRIBUTIONS

CZ, ML, SPWH, JL and KCK designed experiment 1 and 2. CZ and KCK designed

experiment 3. CZ, NJ, SL and KCK collected the data. NJ analysed the MRS data,

CZ performed all other data analyses. CZ and KCK wrote the manuscript and all of

the other authors edited the manuscript.

REFERENCES

Ahmed, S.P., Bittencourt-Hewitt, A., Sebastian, C.L., 2015. Neurocognitive bases of emotion regulation development in adolescence. Dev. Cogn. Neurosci. 15, 11–25. https://doi.org/10.1016/j.dcn.2015.07.006

Akirav, I., Maroun, M., 2007. The role of the medial prefrontal cortex-amygdala circuit in stress effects on the extinction of fear. Neural Plast. 2007, 30873. https://doi.org/10.1155/2007/30873

Angold, A., Costello, E.J., Messer, S.C., Pickles, A., 1995. Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. Int. J. Methods Psychiatr. Res. 5, 237–249.

Banks, S.J., Eddy, K.T., Angstadt, M., Nathan, P.J., Phan, K.L., 2007. Amygdala–frontal connectivity during emotion regulation. Soc. Cogn. Affect. Neurosci. 2, 303–312. https://doi.org/10.1093/scan/nsm029

Beddington, J., Cooper, C., Field, J., 2008. The mental wealth of nations. Nature 455, 1057–1059.

Blakemore, S.-J., 2008. The social brain in adolescence. Nat. Rev. Neurosci. 9, 267–277. https://doi.org/10.1038/nrn2353

Brett, M., Anton, J.-L., Valabreque, R., Poline, J.-B., 2002. Region of interest analysis using an SPM toolbox. Available on CD-ROM in Neuroimage. Present. 8th Int. Conf. Funct. Mapp. Hum. Brain.

Burnett, S., Sebastian, C., Cohen Kadosh, K., Blakemore, S.-J., 2011. The social brain in adolescence: Evidence from functional magnetic resonance imaging and behavioural studies. Neurosci. Biobehav. Rev. 35, 1654–1664. https://doi.org/10.1016/j.neubiorev.2010.10.011

Chefer, V.I., Wang, R., Shippenberg, T.S., 2011. Basolateral Amygdala-Driven Augmentation of Medial Prefrontal Cortex GABAergic Neurotransmission in Response to Environmental Stimuli Associated with Cocaine Administration. Neuropsychopharmacology 36, 2018–2029. https://doi.org/10.1038/npp.2011.89

Cohen Kadosh, K., Linden, D.E.J., Lau, J.Y.F., 2013. Plasticity during childhood and adolescence: innovative approaches to investigating neurocognitive development. Dev. Sci. 16, 574–83. https://doi.org/10.1111/desc.12054

Cohen Kadosh, K., Luo, Q., de Burca, C., Sokunbi, M.O., Feng, J., Linden, D.E.J., Lau, J.Y.F., 2016. Using real-time fMRI to influence effective connectivity in the developing emotion regulation network. Neuroimage 125, 616–626. https://doi.org/10.1016/j.neuroimage.2015.09.070

Constantinidis, C., Williams, G. V., Goldman-Rakic, P.S., 2002. A role for inhibition in shaping the temporal flow of information in prefrontal cortex. Nat. Neurosci. 5, 175–180. https://doi.org/10.1038/nn799

Courtin, J., Chaudun, F., Rozeske, R.R., Karalis, N., Gonzalez-Campo, C., Wurtz, H., Abdi, A., Baufreton, J., Bienvenu, T.C.M., Herry, C., 2014. Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. Nature 505, 92–96. https://doi.org/10.1038/nature12755

Davidson, R.J., Putnam, K.M., Larson, C.L., 2000. Dysfunction in the neural circuitry of emotion regulation--a possible prelude to violence. Science 289, 591–4.

Dawson, D.A., Lam, J., Lewis, L.B., Carbonell, F., Mendola, J.D., Shmuel, A., 2016. Partial Correlation-Based Retinotopically Organized Resting-State Functional Connectivity Within and Between Areas of the Visual Cortex Reflects More Than Cortical Distance. Brain Connect. 6, 57–75. https://doi.org/10.1089/brain.2014.0331

Delli Pizzi, S., Chiacchiaretta, P., Mantini, D., Bubbico, G., Ferretti, A., Edden, R., Di Giulio, C., Onofrj, M., Bonanni, L., 2017. Functional and neurochemical interactions within the amygdala- medial prefrontal cortex circuit and their relevance to emotional processing. Brain Struct Funct 222, 1267–1279. https://doi.org/10.1007/s00429-016-1276-z.

Derryberry, D., Reed, M.A., 2002. Anxiety-related attentional biases and their regulation by attentional control. J. Abnorm. Psychol. 111, 225–36.

Dougherty, L.R., Blankenship, S.L., Spechler, P.A., Padmala, S., Pessoa, L., 2015. An fMRI Pilot Study of Cognitive Reappraisal in Children: Divergent Effects on Brain and

Behavior. J. Psychopathol. Behav. Assess. 37, 634–644.
https://doi.org/10.1007/s10862-015-9492-z

Dumontheil, I., Gilbert, S.J., Frith, C.D., Burgess, P.W., 2010. Recruitment of lateral rostral prefrontal cortex in spontaneous and task-related thoughts. Q. J. Exp. Psychol. (Hove). 63, 1740–56. https://doi.org/10.1080/17470210903538114

Dumontheil, I., Hillebrandt, H., Apperly, I.A., Blakemore, S.-J., 2012. Developmental Differences in the Control of Action Selection by Social Information. J. Cogn. Neurosci. 24, 2080–2095. https://doi.org/10.1162/jocn_a_00268

Eickhoff, S.B., Stephan, K.E., Mohlberg, H., Grefkes, C., Fink, G.R., Amunts, K., Zilles, K., 2005. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. Neuroimage 25, 1325–1335. https://doi.org/10.1016/j.neuroimage.2004.12.034

Emmert, K., Kopel, R., Sulzer, J., Brühl, A.B., Berman, B.D., Linden, D.E.J., Horovitz, S.G., Breimhorst, M., Caria, A., Frank, S., Johnston, S., Long, Z., Paret, C., Robineau, F., Veit, R., Bartsch, A., Beckmann, C.F., Van De Ville, D., Haller, S., 2016. Meta-analysis of real-time fMRI neurofeedback studies using individual participant data: How is brain regulation mediated? Neuroimage 124, 806–812. https://doi.org/10.1016/j.neuroimage.2015.09.042

Gauthier, I., Nuss, P., 2015. Anxiety disorders and GABA neurotransmission: a disturbance of modulation. Neuropsychiatr. Dis. Treat. 11, 165. https://doi.org/10.2147/NDT.S58841

Gee, D.G., Humphreys, K.L., Flannery, J., Goff, B., Telzer, E.H., Shapiro, M., Hare, T.A., Bookheimer, S.Y., Tottenham, N., 2013. A Developmental Shift from Positive to Negative Connectivity in Human Amygdala-Prefrontal Circuitry. J. Neurosci. 33, 4584–4593. https://doi.org/10.1523/JNEUROSCI.3446-12.2013

Goddings, A.-L., Burnett Heyes, S., Bird, G., Viner, R.M., Blakemore, S.-J., 2012. The relationship between puberty and social emotion processing. Dev. Sci. 15, 801–811. https://doi.org/10.1111/j.1467-7687.2012.01174.x

Goddings, A.-L., Mills, K.L., Clasen, L.S., Giedd, J.N., Viner, R.M., Blakemore, S.-J., 2014. The influence of puberty on subcortical brain development. Neuroimage 88, 242–251. https://doi.org/10.1016/j.neuroimage.2013.09.073

Haller, S.P.W., Cohen Kadosh, K., Lau, J.Y.F., 2013. A developmental angle to understanding the mechanisms of biased cognitions in social anxiety. Front. Hum. Neurosci. 7, 846. https://doi.org/10.3389/fnhum.2013.00846

Haller, S.P.W., Cohen Kadosh, K., Scerif, G., Lau, J.Y.F., 2015. Social anxiety disorder in adolescence: How developmental cognitive neuroscience findings may shape understanding and interventions for psychopathology. Dev. Cogn. Neurosci. 13, 11–20. https://doi.org/10.1016/j.dcn.2015.02.002

Haller, S.P.W., Raeder, S.M., Scerif, G., Cohen Kadosh, K., Lau, J.Y.F., 2016. Measuring online interpretations and attributions of social situations: Links with adolescent social anxiety. J. Behav. Ther. Exp. Psychiatry 50, 250–256. https://doi.org/10.1016/j.jbtep.2015.09.009

Hare, T.A., Tottenham, N., Galvan, A., Voss, H.U., Glover, G.H., Casey, B.J., 2008. Biological substrates of emotional reactivity and regulation in adolescence during an emotional go-nogo task. Biol. Psychiatry 63, 927–34. https://doi.org/10.1016/j.biopsych.2008.03.015

Hariri, A.R., Mattay, V.S., Tessitore, A., Fera, F., Weinberger, D.R., 2003. Neocortical modulation of the amygdala response to fearful stimuli. Biol. Psychiatry 53, 494–501.

Hattie, J., Timperley, H., 2007. The Power of Feedback. Rev. Educ. Res. 77, 81–112.

Hayes, A., 2018. Introduction to Mediation, Moderation, and Conditional Process Analysis - A Regression-Based Approach, Second Edi. ed. Guilford Press, New York.

Howell, D., 2011. Statistical methods for psychology. Cengage Learning, Wadsworth.

Johnston, S.J., Boehm, S.G., Healy, D., Goebel, R., Linden, D.E.J., 2010. Neurofeedback: A promising tool for the self-regulation of emotion networks. Neuroimage 49, 1066–1072. https://doi.org/10.1016/j.neuroimage.2009.07.056

Keshavan, M.S., Giedd, J., Lau, J.Y.F., Lewis, D.A., Paus, T., 2014. Changes in the adolescent brain and the pathophysiology of psychotic disorders. The Lancet Psychiatry 1, 549–558. https://doi.org/10.1016/S2215-0366(14)00081-9

Kim, H., Somerville, L.H., Johnstone, T., Alexander, A.L., Whalen, P.J., 2003. Inverse amygdala and medial prefrontal cortex responses to surprised faces. Neuroreport 14, 2317–22. https://doi.org/10.1097/01.wnr.0000101520.44335.20

Kim, M.J., Loucks, R.A., Palmer, A.L., Brown, A.C., Solomon, K.M., Marchante, A.N., Whalen, P.J., 2011. The structural and functional connectivity of the amygdala: From normal emotion to pathological anxiety. Behav. Brain Res. 223, 403–410. https://doi.org/10.1016/j.bbr.2011.04.025

Kim, M.J., Whalen, P.J., 2009. The structural integrity of an amygdala-prefrontal pathway predicts trait anxiety. J. Neurosci. 29, 11614–8. https://doi.org/10.1523/JNEUROSCI.2335-09.2009

Kohn, N., Eickhoff, S.B., Scheller, M., Laird, A.R., Fox, P.T., Habel, U., 2014. Neural network of cognitive emotion regulation — An ALE meta-analysis and MACM analysis. Neuroimage 87, 345–355. https://doi.org/10.1016/j.neuroimage.2013.11.001

Koush, Y., Meskaldji, D.-E., Pichon, S., Rey, G., Rieger, S.W., Linden, D.E.J., Van De Ville, D., Vuilleumier, P., Scharnowski, F., 2017. Learning Control Over Emotion Networks Through Connectivity-Based Neurofeedback. Cereb. Cortex 27, 1193–1202. https://doi.org/10.1093/cercor/bhv311

Koush, Y., Rosa, M.J., Robineau, F., Heinen, K., W. Rieger, S., Weiskopf, N., Vuilleumier, P., Van De Ville, D., Scharnowski, F., 2013. Connectivity-based neurofeedback: Dynamic causal modeling for real-time fMRI. Neuroimage 81, 422–430. https://doi.org/10.1016/j.neuroimage.2013.05.010

La Greca, A.M., Lopez, N., 1998. Social Anxiety Among Adolescents: Linkages with Peer Relations and Friendships. J. Abnorm. Child Psychol. 26, 83–94. https://doi.org/10.1023/A:1022684520514

Linden, D.E.J., 2012. The Challenges and Promise of Neuroimaging in Psychiatry. Neuron 73, 8–22. https://doi.org/10.1016/j.neuron.2011.12.014

Linden, D.E.J., Habes, I., Johnston, S.J., Linden, S., Tatineni, R., Subramanian, L., Sorger, B., Healy, D., Goebel, R., 2012. Real-time self-regulation of emotion networks in patients with depression. PLoS One 7. https://doi.org/10.1371/journal.pone.0038115

Linscott, R.J., van Os, J., 2013. An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: on the pathway from proneness to persistence to dimensional expression across mental disorders. Psychol. Med. 43, 1133–49. https://doi.org/10.1017/S0033291712001626

Little, R.J.A., 1988. A Test of Missing Completely at Random for Multivariate Data with Missing Values. J. Am. Stat. Assoc. 83, 1198–1202. https://doi.org/10.1080/01621459.1988.10478722

Lotte, F., Larrue, F., Mühl, C., 2013. Flaws in current human training protocols for spontaneous Brain-Computer Interfaces: lessons learned from instructional design. Front. Hum. Neurosci. 7, 568.

Lubianiker, N., Goldway, N., Fruchtman-Steinbok, T., Paret, C., Keynan, J.N., Singer, N., Cohen, A., Kadosh, K.C., Linden, D.E.J., Hendler, T., 2019. Process-based framework for precise neuromodulation. Nat. Hum. Behav. 3, 436–445. https://doi.org/10.1038/s41562-019-0573-y

Luciano, J. V., Algarabel, S., Tomás, J.M., Martínez, J.L., 2005. Development and validation of the thought control ability questionnaire. Pers. Individ. Dif. 38, 997–1008. https://doi.org/10.1016/J.PAID.2004.06.020

Meinel, A., Castaño-Candamil, S., Reis, J., Tangermann, M., 2016. Pre-Trial EEG-Based Single-Trial Motor Performance Prediction to Enhance Neuroergonomics for a Hand Force Task. Front. Hum. Neurosci. 10, 170. https://doi.org/10.3389/fnhum.2016.00170

Ochsner, K., Gross, J., 2005. The cognitive control of emotion. Trends Cogn. Sci. 9, 242–249. https://doi.org/10.1016/j.tics.2005.03.010

Paret, C., Kluetsch, R., Ruf, M., Demirakca, T., Hoesterey, S., Ende, G., Schmahl, C., 2014. Down-regulation of amygdala activation with real-time fMRI neurofeedback in a healthy female sample. Front. Behav. Neurosci. 8, 299. https://doi.org/10.3389/fnbeh.2014.00299

Paus, T., Keshavan, M., Giedd, J.N., 2008. Why do many psychiatric disorders emerge during adolescence? Nat. Rev. Neurosci. 9, 947–957. https://doi.org/10.1038/nrn2513

Pezawas, L., Meyer-Lindenberg, A., Drabant, E.M., Verchinski, B.A., Munoz, K.E.,
Kolachana, B.S., Egan, M.F., Mattay, V.S., Hariri, A.R., Weinberger, D.R., 2005. 5-
HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic
susceptibility mechanism for depression. Nat. Neurosci. 8, 828–834.
https://doi.org/10.1038/nn1463
Pine, D.S., Cohen, P., Brook, J.S., 2001. Emotional Reactivity and Risk for Psychopathology
Among Adolescents. CNS Spectr. 6, 27–35.
https://doi.org/10.1017/S1092852900022860
Provencher, S.W., 2001. Automatic quantitation of localized in vivo 1H spectra with
LCModel. NMR Biomed. 14, 260–264. https://doi.org/10.1002/nbm.698
Provencher, S.W., 1993. Estimation of metabolite concentrations from localizedin vivo
proton NMR spectra. Magn. Reson. Med. 30, 672–679.
https://doi.org/10.1002/mrm.1910300604
Quirk, G.J., Beer, J.S., 2006. Prefrontal involvement in the regulation of emotion:
convergence of rat and human studies. Curr. Opin. Neurobiol. 16, 723–727.
https://doi.org/10.1016/j.conb.2006.07.004
Revelle, W., 2015. psych: Procedures for Personality and Psychological Research.
Scharnowski, F., Weiskopf, N., 2015. Cognitive enhancement through real-time fMRI
neurofeedback. Curr. Opin. Behav. Sci. 4, 122–127.
https://doi.org/10.1016/J.COBEHA.2015.05.001
Scheinost, D., Stoica, T., Saksa, J., Papademetris, X., Constable, R.T., Pittenger, C.,
Hampson, M., 2013. Orbitofrontal cortex neurofeedback produces lasting changes in
contamination anxiety and resting-state connectivity. Transl. Psychiatry 3, e250-6.
https://doi.org/10.1038/tp.2013.24
Schwarzkopf, D.S., De Haas, B., Rees, G., 2012. Better Ways to Improve Standards in
Brain-Behavior Correlation Analysis. Front. Hum. Neurosci. 6, 1–6.
Sidorchuk, A., Isomura, K., Molero, Y., Hellner, C., Lichtenstein, P., Chang, Z., Franck, J.,
Fernández de la Cruz, L., Mataix-Cols, D., 2018. Benzodiazepine prescribing for
children, adolescents, and young adults from 2006 through 2013: A total population
register-linkage study. PLoS Med. 15, e1002635.
https://doi.org/10.1371/journal.pmed.1002635
Silvers, J.A., McRae, K., Gabrieli, J.D.E., Gross, J.J., Remy, K.A., Ochsner, K.N., 2012. Age-
related differences in emotional reactivity, regulation, and rejection sensitivity in
adolescence. Emotion 12, 1235–1247. https://doi.org/10.1037/a0028297
Sitaram, R., Ros, T., Stoeckel, L., Haller, S., Scharnowski, F., Lewis-Peacock, J., Weiskopf,
N., Blefari, M.L., Rana, M., Oblak, E., Birbaumer, N., Sulzer, J., 2016. Closed-loop
brain training: the science of neurofeedback. Nat. Rev. Neurosci. 18, 86–100.
https://doi.org/10.1038/nrn.2016.164
Spielberger, C.., Sydeman, S., Owen, A., Marsh, B., 1999. Measuring anxiety and anger
with the State-Trait Anxiety Inventory (STAI) and the State-Trait Anger Expression
Inventory (STAXI)., in: Maruish, M. (Ed.), The Use of Psychological Testing for
Treatment Planning and Outcomes Assessment. Lawrence Erlbaum Associates
Publishers, Mahwah, NJ, pp. 993–1021.
Steel, A., Chiew, M., Jezzard, P., Voets, N.L., Plaha, P., Thomas, M.A., Stagg, C.J., Emir,
U.E., 2018. Metabolite-cycled density-weighted concentric rings k-space trajectory
(DW-CRT) enables high-resolution 1H magnetic resonance spectroscopic imaging at
3-Tesla. Sci. Rep. 8, 7792. https://doi.org/10.1038/s41598-018-26096-y
Streeter, C.C., Jensen, J.E., Perlmutter, R.M., Cabral, H.J., Tian, H., Terhune, D.B., Ciraulo,
D.A., Renshaw, P.F., 2007. Yoga Asana Sessions Increase Brain GABA Levels: A Pilot
Study. J. Altern. Complement. Med. 13, 419–426.
https://doi.org/10.1089/acm.2007.6338
Streeter, C.C., Whitfield, T.H., Owen, L., Rein, T., Karri, S.K., Yakhkind, A., Perlmutter, R.,
Prescot, A., Renshaw, P.F., Ciraulo, D.A., Jensen, J.E., 2010. Effects of Yoga Versus
Walking on Mood, Anxiety, and Brain GABA Levels: A Randomized Controlled MRS
Study. J. Altern. Complement. Med. 16, 1145–1152.
https://doi.org/10.1089/acm.2010.0007
Trentacosta, C.J., Fine, S.E., 2010. Emotion Knowledge, Social Competence, and Behavior
Problems in Childhood and Adolescence: A Meta-Analytic Review. Soc. Dev. 19, 1–29.

https://doi.org/10.1111/j.1467-9507.2009.00543.x

Wechsler, D., 2011. Wechsler Abbreviated Scale of Intelligence - Second Edition (WASI-II). Harcourt Assessment, Texas.

Weiskopf, N., Mathiak, K., Bock, S.W., Scharnowski, F., Veit, R., Grodd, W., Goebel, R., Birbaumer, N., 2004. Principles of a Brain-Computer Interface (BCI) Based on Real-Time Functional Magnetic Resonance Imaging (fMRI). IEEE Trans. Biomed. Eng. 51, 966–970. https://doi.org/10.1109/TBME.2004.827063

Wells, A., Davies, M.I., 1994. The thought control questionnaire: A measure of individual differences in the control of unwanted thoughts. Behav. Res. Ther. 32, 871–878. https://doi.org/10.1016/0005-7967(94)90168-6

Wu, M., Kujawa, A., Lu, L.H., Fitzgerald, D.A., Klumpp, H., Fitzgerald, K.D., Monk, C.S., Phan, K.L., 2016. Age-related changes in amygdala-frontal connectivity during emotional face processing from childhood into young adulthood. Hum. Brain Mapp. 37, 1684–1695. https://doi.org/10.1002/hbm.23129

Zilverstand, A., Sorger, B., Sarkheil, P., Goebel, R., 2015. fMRI neurofeedback facilitates anxiety regulation in females with spider phobia. Front. Behav. Neurosci. 9, 1–12. https://doi.org/10.3389/fnbeh.2015.00148

Zilverstand, A., Sorger, B., Zimmermann, J., Kaas, A., Goebel, R., 2014. Windowed Correlation: A Suitable Tool for Providing Dynamic fMRI-Based Functional Connectivity Neurofeedback on Task Difficulty. PLoS One 9, e85929. https://doi.org/10.1371/journal.pone.0085929

Zotev, V., Phillips, R., Young, K.D., Drevets, W.C., Bodurka, J., 2013. Prefrontal Control of the Amygdala during Real-Time fMRI Neurofeedback Training of Emotion Regulation. PLoS One 8, e79184. https://doi.org/10.1371/journal.pone.0079184

Supplementary Materials for:
Modulatory effects of dynamic fMRI-based neurofeedback on emotion regulation networks in adolescent females

Catharina Zich, Nicola Johnstone, Michael Lührs, Stephen Lisk, Simone P W Haller, Annalisa Lipp, Jennifer Y F Lau, Kathrin Cohen Kadosh

Supplementary Methods

MRI data acquisition

Before the experimental tasks a high-resolution structural scan was obtained from each subject using a T1-weighted magnetization-prepared rapid-acquisition gradient echo (MPRAGE) sequence (TR = 1900 ms, TE = 3.97 ms, filp angle = 8°, slice thickness 1 mm, in-plane resolution 1 x 1 mm, orientation = sagittal).

Functional data were recorded using 2D multiband gradient echo planer imaging (Todd et al., 2016) (TR = 933 ms, TE = 33.40 ms, flip angle = 64°, slice thickness 2 mm, in-plane resolution 2 x 2 mm, orientation = transversal, 72 slices, multi-band factor = 6).

For **Experiment 1** and **2** the localizer comprised 570 volumes and each of the four NF runs comprised 310 volumes. For **Experiment 3** the localizer comprised 300 volumes and each of the three NF runs comprised 330 volumes. At the beginning of the localizer and each NF run ten additional volumes were acquired, but not analysed, to avoid saturation effects.

For **Experiment 3** we collected MRS data. MRS data were acquired using a semi-LASER sequence (Scheenen et al., 2008) (32 averages, TR = 3500 ms, TE = 28 ms, voxel size = 20 x 20 x 20 mm) using VAPOR (variable power RF pulses with optimized relaxation delays) water suppression (Tkác et al., 1999). Two voxels of interest (VOI) were manually positioned. One VOI was positioned at the same position as the PFC ROI, i.e. local maximum of the GLM t-statistics of the brain activity during the localizer (the sum of the three contrasts: appraisal > fixation, reappraisal > fixation, reappraisal > appraisal) within the left dorsolateral and medial PFC. The other VOI, the anterior cingulate cortex (ACC) VOI, was centred on the interhemispheric fissure on the coronal plane superior to the frontal aspect of the corpus callosum on the sagittal plane.

Neurofeedback task – Instructions

All participants were blind to their NF implementation and were instructed as follows: *"You will see a thermometer with a green rim on the screen. The red bars show how much the regions that are important for emotions are active. Your job is to get these regions as active as possible! So try to get this thermometer up as much as possible. Similar to the task before, try to control your thoughts towards a positive feeling. When the thermometer does not have a green rim, the thermometer is not working. However, even if the thermometer is not working, your task will be the same and we are still measuring how much your brain is active. The two different thermometers will alternate.*

When the thermometer has a green rim, it might need some time to properly estimate the brain activity. This leads to a delay in the thermometer display." Debriefing revealed that, in order to change their thoughts and emotions towards a happier mood, participants thought about certain people, things that happened in the past and things that they would like to happen in the future.

Supplementary Results

NF implementations differentially modulate the frequency of negative PFC-amygdala partial correlations

The observed differences in fc between fc-NF and no-NF as illustrated in Supplementary Fig. 1 are most likely due to changes in the frequency of negative partial correlations (secondary measure, **Supplementary Fig. 2a**).

The relationship of fc-NF and fc was assessed by correlating the frequency of negative partial correlations at each of the 20 volumes with the number of fc-NF volumes within the correlation window (**Supplementary Fig. 2b**). For the negative NF implementation and the positive NF implementation, the frequency of negative partial correlations and the number of fc-NF volumes in the correlation window were related negatively at trend level (negative NF implementation: $r_{(38)} = -.31$, $p = .05$; positive NF implementation: $r_{(38)} = -.32$, $p = .05$). Contrary, for the weighted negative NF implementation a positive relationship was observed ($r_{(38)} = .47$, $p < .01$), indicating that a higher number of fc-NF volumes within the correlation window equalled a higher frequency of negative partial correlation.

The relationship between the frequency of negative partial correlations and the number of fc-NF volumes in the correlation window was significantly different for the weighted negative NF implementation in comparison to the other two NF implementations (both p's < .001), whereby the other two NF implementations did not differ regarding this aspect ($p = .96$).

NF implementations differentially modulate the frequency of correlation scenarios

One limitation of using a fc-NF approach that is based on correlational measures is that we cannot directly infer the direction of change within each region. In other words, a negative correlation exits when variable 1 increases and variable 2 decreases, or when variable 1 decreases and variable 2 increases. However, given the top-down PFC regulation of amygdala reactivity the main goal of PFC-amygdala fc-NF constitutes PFC increase and amygdala decrease.

To understand the nature of the results depicted in Supplementary Fig. 1 and 2, we assigned each negative correlation to one out of four scenarios (see **Supplementary Fig. 5** for details). **Supplementary Fig. 3a** illustrates the percentage of each scenario for fc-NF and no-NF mini-blocks.

As above, we further investigated the relationship between the number of fc-NF volumes within the correlation window and the frequency of scenario 1, i.e. PFC up-regulation & amygdala down-regulation, and of scenario 2, i.e. PFC down-regulation & amygdala up-regulation, at each volume (**Supplementary Fig. 3b**). For the negative NF implementation and the weighted negative NF implementation the number of fc-NF volumes in the correlation window was positively related to the frequency of scenario 1 (negative NF implementation: $r_{(38)} = 0.39$, $p = .01$; weighted negative NF implementation: $r_{(38)} = .70$, $p < .001$) and negatively related to the frequency of scenario 2 (negative NF implementation: $r_{(38)} = -.50$, $p < .01$; weighted negative NF implementation: $r_{(38)} = -.64$, $p < .001$). In contrast, for the positive NF implementation the pattern was reversed, i.e. the number of fc-NF volumes in the correlation window was negatively related to the frequency of scenario 1 ($r_{(38)} = -.44$, $p < .01$) and positively related to the frequency of scenario 2 ($r_{(38)} = .55$, $p < .001$). This suggests that only NF implementations reinforcing negative PFC-amygdala fc support the task-relevant negative fc pattern.

Further, we compared the two negative NF implementations with regard to the strength of the relationship between the number of fc-NF volumes in the correlation window and the frequency of scenario 1. We found a significant difference, indicating that the relationship is strongest for the weighted negative NF implementation.

Supplementary Discussion

Length of the correlation window

We chose a window length of 20 volumes based on a previous study by Zilverstand and colleagues (Zilverstand et al., 2014), who found that the fc measures in a well-controlled, simple motor task were weaker and less reliable for a shorter (12 s) window but not qualitatively different form a longer (26 s) window. Given the higher reliability of the longer correlation windows, we matched the size of our time-windows with the length of the mini-blocks. This allowed us to directly assess the effect of fc-NF as a function of the ratio of fc-NF : no-NF volumes in the correlation window. We found that the PFC-amygdala fc (Supplementary Fig. 1, Fig. 2a,c), frequency of negative partial correlations (Supplementary Fig. 2, Fig. 2b,d), and frequency of scenarios describing different patterns of activation changes (Supplementary Fig. 3) changed as a function of fc-NF volumes within a correlation window. This finding is not only of theoretical interest, but also has important practical implications. Specifically, it suggests that with regards to clinical applications the relation between task window and correlation window should be adapted, so that the length of the task window exceeds the correlation window, in order to further enhance the fc-NF-effect and practice-related changes.

Supplementary Figures

Supplementary Fig. 1 Difference in average functional connectivity (fc) between fc-NF (red) and the no-NF (blue) for each NF implementation ($N = 5$ per NF implementation). **(a)** Means and standard error for the average fc, i.e. partial correlations r_p, at each volume within a mini-block. Fc is averaged across mini-blocks within one condition (fc-NF, no-NF), NF runs (1, 2, 3, 4) and individuals within one NF implementation (negative NF implementation, weighted negative NF implementation, positive NF implementation). Number of fc-NF volumes in the correlation window (N of fc-NF Volumes in r_p) is illustrated in form of a grey triangle. **(b)** Relationship between average fc at each volume and the number of fc-NF volumes within the correlation window.

Supplementary Fig. 2 As Supplementary Fig. 1, but for the frequency of negative partial correlations (N of negative r_p).

Supplementary Fig. 3 Direction of activity change within the negative partial correlations for each NF implementation ($N = 5$ per implementation). **(a)** Percentage of negative partial correlations between PFC and amygdala for scenario 1 (PFC up-regulation & amygdala down-regulation; green), scenario 2 (PFC down-regulation & amygdala up-regulation; black) or scenario 3 and 4 (white). Percentages are averaged across mini-blocks within one condition (fc-NF, no-NF), NF runs (1, 2, 3, 4) and individuals within one NF implementation (negative NF implementation, weighted negative NF implementation, positive NF implementation). The number of fc-NF volumes in the correlation window is illustrated in form of a grey triangle. **(b)** Relationships between the number of fc-NF volumes within the correlation window (N of fc-NF Volumes in r_p) and frequency of scenario 1 (PFC up-regulation & amygdala down-regulation; green) or scenario 2 (PFC down-regulation & amygdala up-regulation; black).

a

$$\text{practice-related change in fc} = \frac{\sum_{i=1}^{4}(\text{Run}_i - \overline{\text{Run}})(\text{fc}_i - \overline{\text{fc}})}{\sum_{i=1}^{4}(\text{Run}_i - \overline{\text{Run}})^2}$$

b

intitial fc

c

$$\text{fc-NF-effect} = \sum_{i=1}^{10}(\text{fc\_NF}_i - \text{no\_NF}_i) + \sum_{j=10}^{20}\text{no\_NF}_j - \text{fc\_NF}_j$$

Supplementary Fig. 4 Illustration of quantification of neural measures. (**a**) Visualization of practice-related change in fc, i.e. slope of the linear regression for the average fc within each NF run. (**b**) Illustration of initial fc, i.e. average fc of the first two mini-blocks. (**c**) Visualization of the fc-NF-effect (hatched) on exemplary data from one subject. Due to the dynamic character of the fc-NF, the fc-NF-effect is composed of two parts. For fc-NF mini-bocks (red) the ratio of fc-NF : no-NF volumes in the correlation window is 1:19 at the first volume of the mini-block, 2:18 at the second volume of the mini-block and 20:0 at the last volume of the mini-block. Consequently, in fc-NF mini-blocks volumes 1 to 10 are dominated by no-NF volumes, and volumes 11 to 20 are dominated by fc-NF volumes. Exactly the opposite is the case for no-NF mini-blocks.

Supplementary Fig. 5 Schematic representation of the procedure used to quantify the direction of activity change within a negative partial correlation. **Centre**: Exemplary negative partial correlation between PFC and amygdala activity accounting for the 'activity' in the CST. The direction of activity change within this correlation was determined for the PFC and the amygdala separately. In detail, both, the PFC and the amygdala, were correlated separately with time while accounting for CST 'activity'. Theoretically four different scenarios are possible: scenario 1 – PFC up-regulation & amygdala down-regulation, scenario 2 – PFC down-regulation & amygdala up-regulation, scenario 3 – PFC & amygdala up-regulation and scenario 4 – PFC & amygdala down-regulation.

Supplementary Tables

Supplementary Table 1 Emotional/metacognitive measures for **Experiment 1**.

Pre	Negative NF implementation (N=5)	Weighted negative NF implementation (N=5)	Positive NF implementation (N=5)
$CERQ_{pre}$	M = 59.50; SD = 9.26	M = 57.40; SD = 7.33	M = 53.65; SD = 10.65
MFQ_{pre}	M = 2.60; SD = 2.07	M = 1.50; SD = 1.91	M = 2.20; SD = 1.71
SAS-S	M= 46.00; SD = 13.41	M = 48.00; SD = 11.22	M = 56.80; SD = 14.62
$STAI-S_{pre}$	M = 31.40; SD = 8.91	M = 23.00; SD = 6.38	M = 36.80; SD = 5.22
STAI-T	M = 35.20; SD = 7.92	M = 35.80; SD = 8.56	M = 38.00; SD = 9.92
TCQ_{pre}	M = 65.75; SD = 4.27	M = 67.20; SD = 9.26	M = 59.60; SD = 7.50
$TCAQ_{pre}$	M = 79.40; SD = 14.62	M = 82.00; SD = 10.42	M = 75.60; SD = 13.63

CERQ = Emotion Regulation Questionnaire; *MFQ* = Mood and Feelings Questionnaire; *SAS-S* = Social Anxiety Scale for Adolescents; *STAI-S, STAI-T* = State-Trait Anxiety Inventory; *TCQ* = Thought Control Questionnaire; *TCAQ* = Cognitive Thought Control Ability Questionnaire

Supplementary Table 2 Emotional/metacognitive measures for **Experiment 2**.

	Pre	Post	Difference
CERQ	N = 27; M = 60.52; SD = 11.09	N = 27; M = 60.93; SD = 11.18	$t_{(26)}$ = -.442; p = .662
MFQ	N = 25; M = 5.44; SD = 4.81	N = 27; M = 5.56; SD = 4.56	$t_{(24)}$ = -.778; p = .444
SAS-S	N = 27; M= 49.15; SD = 15.01		
STAI-S	N = 24; M = 32.79; SD = 8.61	N = 27; M = 31.78; SD = 9.83	$t_{(23)}$ = 1.784; p = .088
STAI-T	N = 27; M = 43.52; SD = 11.49		
TCQ	N = 27; M = 61.04; SD = 8.67	N = 27; M = 60.70; SD = 8.99	$t_{(26)}$ = .251; p = .804
TCAQ	N = 23; M = 78.91; SD = 15.37	N = 27; M = 79.85; SD = 14.77	$t_{(22)}$ = .041; p = .968

CERQ = Emotion Regulation Questionnaire; MFQ = Mood and Feelings Questionnaire; SAS-S = Social Anxiety Scale for Adolescents; STAI-S, STAI-T = State-Trait Anxiety Inventory; TCQ = Thought Control Questionnaire; TCAQ = Cognitive Thought Control Ability Questionnaire

Supplementary Table 3: Emotional/metacognitive measures for a subsample of **Experiment 2** (negative level, i.e. change in the desired direction, *N*=13).

	Pre	Post	Difference
CERQ	N = 13; M = 62.92; SD = 12.43	N = 13; M = 64.15; SD = 12.43	$t_{(12)}$ = -.854; p = .410
MFQ	N = 12; M = 4.92; SD = 5.00	N = 13; M = 4.92; SD = 4.92	$t_{(11)}$ = -.561; p = .586
SAS-S	N = 13; M= 52.15; SD = 13.45		
STAI-S	N = 13; M = 31.83; SD = 9.04	N = 13; M = 30.85; SD = 7.86	$t_{(12)}$ = .616; p = .551
STAI-T	N = 13; M = 42.54; SD = 12.14		
TCQ	N = 13; M = 62.69; SD = 8.60	N = 13; M = 60.77; SD = 10.69	$t_{(12)}$ = 1.402; p = .133
TCAQ	N = 11; M = 81.27; SD = 15.38	N = 13; M = 80.54; SD = 15.01	$t_{(10)}$ = 1.258; p = .237

CERQ = Emotion Regulation Questionnaire; *MFQ* = Mood and Feelings Questionnaire; *SAS-S* = Social Anxiety Scale for Adolescents; *STAI-S, STAI-T* = State-Trait Anxiety Inventory; *TCQ* = Thought Control Questionnaire; *TCAQ* = Cognitive Thought Control Ability Questionnaire

Supplementary Table 4: Emotional/metacognitive measures for a subsample of **Experiment 2** (positive level, i.e. change in the undesired direction *N*=14).

	Pre	Post	Difference
CERQ	N = 14; M = 58.29; SD = 9.60	N = 14; M = 57.93; SD = 9.34	$t_{(13)}$ = .302; p = .768
MFQ	N = 13; M = 5.92; SD = 4.79	N = 14; M = 6.14; SD = 4.29	$t_{(12)}$ = -.519; p = .613
SAS-S	N = 14; M= 46.36; SD = 16.33		
STAI-S	N = 12; M = 33.75; SD = 8.43	N = 14; M = 32.64; SD = 11.59	$t_{(11)}$ = 1.170; p = .115
STAI-T	N = 14; M = 44.43; SD = 11.24		
TCQ	N = 14; M = 59.50; SD = 8.76	N = 14; M = 61.57; SD = 7.38	$t_{(13)}$ = -.965; p = .352
TCAQ	N = 12; M = 76.75; SD = 15.70	N = 14; M = 79.21; SD = 15.08	$t_{(11)}$ = -1.105; p = .293

CERQ = Emotion Regulation Questionnaire; *MFQ* = Mood and Feelings Questionnaire; *SAS-S* = Social Anxiety Scale for Adolescents; *STAI-S, STAI-T* = State-Trait Anxiety Inventory; *TCQ* = Thought Control Questionnaire; *TCAQ* = Cognitive Thought Control Ability Questionnaire

Supplementary Table 5 Emotional/metacognitive measures for **Experiment 3**.

	Pre	Post	Difference
CERQ	N = 19; M = 58.16; SD = 9.00	N = 19; M = 60.84; SD = 11.08	$t_{(18)}$ = -1.326; p = .202
MFQ	N = 19; M = 6.21; SD = 5.72	N = 19; M = 6.42; SD = 5.82	$t_{(18)}$ = -.200; p = .844
SAS-S	N = 19; M= 48.00; SD = 15.37		
STAI-S	N = 19; M = 31.42; SD = 6.01	N = 19; M = 29.00; SD = 5.85	$t_{(18)}$ = 1.490; p = .153
STAI-T	N = 19; M = 42.42; SD = 10.70		
TCQ	N = 19; M = 60.42; SD = 7.52	N = 19; M = 62.74; SD = 7.90	$t_{(18)}$ = -1.180; p = .254
TCAQ	N = 19; M = 74.47; SD = 18.20	N = 19; M = 75.58; SD = 19.27	$t_{(18)}$ = -.045; p = .657

CERQ = Emotion Regulation Questionnaire; MFQ = Mood and Feelings Questionnaire; SAS-S = Social Anxiety Scale for Adolescents; STAI-S, STAI-T = State-Trait Anxiety Inventory; TCQ = Thought Control Questionnaire; TCAQ = Cognitive Thought Control Ability Questionnaire

References

Scheenen, T.W.J., Klomp, D.W.J., Wijnen, J.P., Heerschap, A., 2008. Short echo time1H-MRSI of the human brain at 3T with minimal chemical shift displacement errors using adiabatic refocusing pulses. Magn. Reson. Med. 59, 1–6. https://doi.org/10.1002/mrm.21302

Tkác, I., Starcuk, Z., Choi, I.Y., Gruetter, R., 1999. In vivo 1H NMR spectroscopy of rat brain at 1 ms echo time. Magn. Reson. Med. 41, 649–56.

Todd, N., Moeller, S., Auerbach, E.J., Yacoub, E., Flandin, G., Weiskopf, N., 2016. Evaluation of 2D multiband EPI imaging for high-resolution, whole-brain, task-based fMRI studies at 3T: Sensitivity and slice leakage artifacts. Neuroimage 124, 32–42. https://doi.org/10.1016/j.neuroimage.2015.08.056

Zilverstand, A., Sorger, B., Zimmermann, J., Kaas, A., Goebel, R., 2014. Windowed Correlation: A Suitable Tool for Providing Dynamic fMRI-Based Functional Connectivity Neurofeedback on Task Difficulty. PLoS One 9, e85929. https://doi.org/10.1371/journal.pone.0085929