Gas seepage pockmark microbiomes suggest the presence of sedimentary coal seams in the Öxarfjörður graben of NE-Iceland

Guðný Vala Þorsteinsdóttir[1,2], Anett Blischke[3], M. Auður Sigurbjörnsdóttir[1], Finnbogi Óskarsson[4], Þórarinn Sveinn Arnarson[5†], Kristinn P. Magnússon[1,2,6], and Oddur Vilhelmsson[1,6]

[1]University of Akureyri, Faculty of Natural Resource Sciences, Borgir v. Nordurslod, 600 Akureyri, Iceland.

[2]Icelandic Institute of Natural History, Borgir v. Nordurslod, 600 Akureyri, Iceland

[3]Íslenskar orkurannsóknir / Iceland GeoSurvey (ISOR), Akureyri Branch, Rangarvollum, 600 Akureyri, Iceland

[4]Íslenskar orkurannsóknir / Iceland GeoSurvey (ISOR), Department of Geothermal Engineering, Grensasvegi 9, 108 Reykjavik, Iceland

[5]Orkustofnun / The Icelandic Energy Authority, Grensasvegi 9, 108 Reykjavik, Iceland ([†]Passed away)

[6]Biomedical Center, University of Iceland, Vatnsmyrarvegur 16, 101 Reykjavik, Iceland

Correspondence: Oddur Vilhelmsson, oddurv@unak.is

Abstract

Natural gas seepage pockmarks are found off and onshore in the Öxarfjörður graben, NE Iceland. The bacterial communities of two onshore seepage sites were analysed by amplicon sequencing of 16S rDNA, along with determining the geochemical characteristics, hydrocarbon content and the carbon isotope composition of the sites.

While one site was found to be characterised by biogenic origin of methane gas, with carbon isotope ratio $\delta^{13}C$ [‰] = -63.2, high content of organic matter and complex

hydrocarbons, the other site showed a mixed origin of the methane gas (δ^{13}C [‰] = -26.6)

with geothermal characteristics and lower organic matter content. While both sites

harboured *Proteobacteria* as the most abundant bacterial phyla, the *Deltaproteobacteria*

were more abundant at the geothermal site, and the *Alphaproteobacteria* at the biogenic

site. The *Dehalococcoidia* class of the *Chloroflexi* phylum was abundant at the

geothermal site while the *Anaerolineae* class was more abundant at the biogenic site.

Bacterial strains from the seepage pockmarks were isolated on selective and differential

media targeting bacteria with bioremediation potential. A total of 106 strains were

isolated and characterised, including representatives from the phyla *Proteobacteria*,

Bacterioidetes, *Firmicutes*, and *Actinobacteria*.

Introduction

Natural gas seepage, the emission of gaseous hydrocarbons from the subsurface, has been studied extensively in the context of petroleum exploration because it can be used as an indicator of petroleum generation in subsurface sediments (1–3). Natural methane gas seepage is the result of subsurface generation or accumulation of methane and the methane concentration in the gas varies according to its source (4). At geothermal and hydrothermal sites, methane is generated by thermogenic processes and seeps up to the surface through cracks and pores, whereas, in deep sea sediments the accumulation of methane can result in cold seeps or methane hydrates where no direct input of heat is found. This is often linked to biogenic methane which is a product of microbial processes in various anaerobic environments, like bog lakes and sea sediments (5, 6). In many cases the methane generation is of mixed origin, that is both thermogenic and biogenic. For example, methane that is formed during early coalification processes (coal bed methane) is not only of thermogenic origin but also produced by microbes utilizing the lignite (7). In these environments one would expect to find microbes that participate in methanogenesis and are capable of methane oxidation, respectively.

Where natural methane gas seeps up to the surface, pockmarks can develop, that are a habitat for diverse microorganisms (8) and can be regarded as hotspots for anaerobic oxidation of methane (AOM). AOM is often dependent on archaea and sulphate-reducing bacteria (47), but can in some cases be driven by bacteria through intra-aerobic-denitrification (9), or possibly reductive dehalogenation, as suggested in a recent study on an ice-covered Antarctic lake (10). Microbial communities of hydrocarbon gas seepage environments have been studied around the world, including the Gulf of Mexico (11), Pacific Ocean Margin (12), Cascadia Margin (13) and Barents Sea (14), mainly because of their sulfate-reducing capabilities and AOM.

In Öxarfjörður bay, NE Iceland, natural gas seepage pockmarks are found both on the seafloor and on shore. Öxarfjörður is located along the lithospheric boundaries of the North-American and the Eurasian plates and forms a graben bounded by the Tjörnes Fracture Zone in the west and the eastern rim of the North Iceland Volcanic Zone in the east. Geothermal activity in Öxarfjörður bay is

confined to three major fissure swarms, cross-sectioning the volcanic zone. The area is prevailed by

the river delta of Jökulsá-á-Fjöllum, causing the Öxarfjörður bay to be even more dynamic in nature.

Geological settings in the Öxarfjörður area were studied extensively in the 1990s (15–18), leading to

the discovery that the methane-rich seepage gas likely originates from thermal alteration of lignite and

coal seams from beneath the 1 km thick sediment (18). Taken together, these studies strongly suggest

the presence of sedimentary lignite in the Öxarfjörður graben (19).

Very little geomicrobiological work has thus far been conducted in Iceland, with most

environmental microbiology work being bioprospective in nature, often paying little attention to

community structures or biogeochemical activity. Natural gas seeps such as those found in

Öxarfjörður, have thus far not been investigated from a microbiological standpoint despite their

unique character which makes them ideal for geomicrobiological studies as both sparsely vegetated

geothermal gas seepage pockmarks and colder, more vegetated seeps are found in close proximity to

one another. Each methane seep system is thought to be unique in terms of the composition of

geological and biological features (8), so taking a snapshot of the microbial community at a methane

gas seepage site can provide valuable insight into the dynamics of the system and initiate biological

discoveries.

In this article, we report the first microbial analysis of the natural gas seepage pockmarks in

Öxarfjörður. Hypothesizing that the microbial community in this environment ought to be dominated

by methane oxidizing microbes, we performed microbial community analysis on 16S rRNA gene

amplicon libraries from two sites differing in visible vegetation and complex hydrocarbon content.

Further hypothesizing that these environments would be a source of hydrocarbon-degrading bacteria,

we isolated a collection of microbes on various media and tested them for degradation of naphthalene.

<u>Materials and methods</u>

Sampling and in-field measurements

Samples were collected at Skógalón (site SX, 66°09'N, 16°37'W) on August 21st, 2014, and on

September 11th, 2015, and at Skógakíll (site AEX, 66°10'N, 16°34'W) on August 13th, 2015 (Fig. 1).

At site SX, where the natural gas seepage pockmarks are somewhat difficult to distinguish from

ordinary marsh gas pockmarks, sites were selected where pockmarks were visibly active and appeared

to form straight lines extending NW-SE. Temperature, pH and conductivity were measured *in-situ*

during sampling with hand-held meters. Sediment samples were collected from shallow cores

obtained using a corer constructed from a 3-cm diameter galvanized-iron pipe that was hammered into

the ground using a sledgehammer, and transferred aseptically to sterile, airtight IsoJars (IsoTech

laboratories, Champaign, Illinois), according to manufacturer´s protocol. Surface soil samples were

collected aseptically directly into sterile IsoJars. Water samples were collected aseptically into sterile

glass bottles. Gas samples were collected into evacuated double-port glass bottles by means of an

inverted nylon funnel connected to silicone rubber tubing. All samples for microbial analysis were

immediately put on dry ice where they were kept during transport to laboratory facilities at University

of Akureyri where they were either processed immediately or stored in a freezer at -18°C until

processing. Samples collected, along with *in-situ* measurements and types of sample are listed in

Table 1.

Chemical analysis of geothermal fluids

Dissolved sulphide in the water samples was determined on-site by titration with mercuric acetate

using dithizone in acetone as indicator (48). Major components in the water samples were determined

at the laboratories of Iceland GeoSurvey (ÍSOR) in Reykjavík: Dissolved inorganic carbon was

determined by alkalinity titration (pH 8.2 to 3.8), purging with nitrogen gas and back-titration (pH 3.8

to 8.2) as described previously (48). Silica was analysed by colorimetric determination of a silica-

molybdate complex at 410 nm using a Jenway 6300 spectrophotometer. Total dissolved solids were

determined by gravimetry. Anions were determined by suppressed ion chromatography on a

ThermoScientific ICS-2100 with an AS-20 column. Major metals were analysed by atomic absorption

spectrometry on a Perkin Elmer 1100B spectrometer. The composition of dry gas was also determined

at the ÍSOR laboratories by gas chromatography on a Perkin Elmer Arnel 4019 light gas analyser

equipped with HayeSep and MolSieve columns and three TCDs.

The concentration of trace elements in water samples were determined by ICP methods at the ALS

Laboratories, Luleå, Sweden. Stable water isotopes (^2H and ^{18}O) were determined by mass

spectrometry using a Delta V Advantage IRMS coupled with a Gasbench II at the Institute of Earth

Sciences, University of Iceland.

Headspace gas analysis from sediment samples was performed at Applied Petroleum

Technologies, Kjeller, Norway, using standard techniques. Briefly as follows:

Sample preparation and extraction. Sediment samples were washed in water to remove mud

before extraction using a Soxtec Tecator instrument. Thimbles were pre-extracted in dichloromethane

with 7% (vol/vol) methanol, 10 min boiling and 20 min rinsing. The crushed sample was weighed

accurately in the pre-extracted thimbles and boiled for 1 hour and rinsed for 2 hours in 80 cc of

dichloromethane with 7% (vol/vol) methanol. Copper blades activated in concentrated hydrochloric

acid were added to the extraction cups to cause free sulphur to react with the copper. An aliquot of

10% of the extract was transferred to a pre-weighed bottle and evaporated to dryness. The amount of

extractable organic matter (EOM) was calculated from the weight of this 10% aliquot.

Deasphaltening. Extracts were evaporated almost to dryness before a small amount of

dichloromethane (3 times the amount of EOM) was added. Pentane was added in excess (40 times the

volume of EOM/oil and dichloromethane). The solution was stored for at least 12 hours in a dark

place before the solution was filtered or centrifuged and the weight of the asphaltenes measured.

GC analysis of gas components. Aliquots of the samples were transferred to exetainers. 0.1-1ml

were sampled using a Gerstel MPS2 autosampler and injected into a Agilent 7890 RGA GC equipped

with Molsieve and Poraplot Q columns, a flame ionisation detector (FID) and 2 thermal conductivity

detector (TCD). Hydrocarbons were measured by FID. H_2, CO_2, N_2, and O_2/Ar by TCD.

Carbon isotope analysis of hydrocarbon compounds and CO_2. The carbon isotopic

composition of the hydrocarbon gas components was determined by a GC-C-IRMS system. Aliquots

were sampled with a syringe and analysed on a Trace GC2000, equipped with a Poraplot Q column,

connected to a Delta plus XP IRMS. The components were burnt to CO_2 and water in a 1000 °C

furnace over Cu/Ni/Pt. The water was removed by Nafion membrane separation. Repeated analyses of

standards indicate that the reproducibility of $\delta^{13}C$ values is better than 1 ‰ PDB (2 sigma).

Carbon isotope analysis of low concentration methane using the Precon. The carbon isotopic

composition of methane was determined by a Precon-IRMS system. Aliquots were sampled with a

GCPal autosampler. CO_2, CO and water were removed on chemical traps. Other hydrocarbons than

CH_4 and remaining traces of CO_2 were removed by cryotrapping. The methane was burnt to CO_2 and

water in a 1000 °C furnace over Cu/Ni/Pt. The water was removed by Nafion membrane separation.

The sample preparation system described (Precon) was connected to a Delta plus XP IRMS for $\delta\,^{13}C$

analysis. Repeated analyses of standards indicate that the reproducibility of $\delta^{13}C$ values is better than

1 ‰ PDB (2 sigma).

GC of EOM fraction. A HP7890 A instrument was used. The column was a CP-Sil-5 CB-MS,

length 30 m, i.d. 0.25 mm, film thickness 0.25 μm. C20D42 is used as an internal standard.

Temperature programme: 50°C (1 min), -4 °C/min, -320 °C (25 min).

Bacterial community analysis

Total DNA was extracted from sediment samples in duplicates, using the PowerSoil kit (MoBio

laboratories) following the manufacturer's protocol. The DNA isolated was measured with Qubit

fluorometer (Invitrogen, Carlsbad, CA) to confirm dsDNA in the samples. Paired-end library of the

16S rDNA hypervariable region V3/V4, was sequenced on Illumina MiSeq platform by Macrogen,

Netherlands, four replicates from AEX1 and four from SX1. The data was processed and analysed

using CLC Genomics Workbench 10.1.1 (https://www.qiagen bioinformatics.com/) and the CLC

Microbial Genomics Module 2.5.1, with default parameters. Operational taxonomic units (OTU) were

clustered by reference based OTU clustering using default parameters in the Microbial Genomics

Midule and tree alignment was performed by using the GreenGenes v.15.5 database for 97%

similarity. For statistical analysis only alpha-diversity of samples was performed since the sequencing

data only contained technical replicates, which does not allow analyses of beta-diversity. Differential

abundance analysis (Likelihood Ratio test) was performed to see statistically significant differences in

taxa between sampling sites.

Initial culturing and isolation of bacteria

Samples were serially diluted to 10^{-6} in sterile Butterfield's buffer and all dilutions plated in duplicate

onto Reasoner's agar 2A (Difco) and several selective and differential media including medium 9K for

iron oxidizers (21), Mn medium for manganese oxidizers (990 mL basal agar B [0.42 g NaOAc, 0.1 g

peptone, 0.1 g yeast extract, 15 agar, 990 mL sample water, autoclaved and cooled to 50°C], 10 mL

pre-warmed filter-sterilized 1 M HEPES at pH 7.5, 100 µL filter-sterilized 100 mM $Mn(II)SO_4$), Gui

medium for laccase producers (990 mL basal agar B, 0,01% guiaicol), Hex medium for hexane

degraders (990 mL basal agar B, 1.3 mL filter-sterilized 99 parts hexane/1 part dishwashing

detergent), Naph medium for naphthalene degraders (basal agar B with several crystals of naphthalene

added to the lid of inverted plates and then sprayed with fast blue for degradation indication), and 2,4-

D medium for dichlorophenoxyacetate degraders (basal agar B supplemented with 2 mM 2,4-D).

Three atmospheric incubation conditions were used: an unmodified atmosphere in sealed plate bags, a

propane-enriched aerobic atmosphere in sealed plate bags flushed daily with propane, and an

anaerobic, propane-supplemented atmosphere in anaerobic jars scrubbed of oxygen with a palladium

catalyst (BBL GasPak) and monitored for anaerobicity with a resazurin strip. The jars were injected

with 100 mL propane through a septum. Plates were incubated in the dark at 5, 15, or 22°C until no

new colonies appeared (up to 4 weeks).

Colony morphotypes were examined by visual features, such as colour and form of elevation and

margins. A representative of each morphotype was aseptically restreaked on fresh media and

restreaked up to three times or until considered isolated strain. Stocks of isolates were prepared by

suspending a loopful of growth in 1.0 mL 28% (v/v) glycerol and are stored at -70°C in the University

of Akureyri culture collection.

16S rRNA gene-based identification of cultured strains

For each strain,1 µl of freezer stock was suspended in 25 µl of lysis buffer (1% Triton x-100, 20 mM

Tris, 2 mM EDTA, pH 8,0) and incubated for 10 minutes at 95°C in the thermocyler (MJR PTC-200

thermocycler, MJ Research Inc. Massachusetts, USA). The lysis buffer solution (1 µl), or 1 µl of

extracted DNA (using UltraClean® Microbial DNA Isolation Kit (MoBio Laboratories, Carlsbad,

California, USA)), was used as a DNA template for Polymerase chain reaction (PCR) using Taq

DNA-polymerase to amplify the DNA using the 'universal' bacterial primers 8F (5'-

AGTTTGATCCTGGCTCAG'3) and 1522R: (5'-AAGGAGGTGATCCAGC CGCA-'3).

The PCR reaction was conducted as follows: 95°C for 3 min, followed by 35 cycles of 95°C for 30

sec, 50°C for 30 sec and 68°C for 90 sec, then final extension at 68°C for 7 min. The PCR products

were loaded on 0,8% agarose gel and run at 100 V for approximately 30 minutes to verify the

presence of approximately 1500-bp amplicons.

The PCR products were purified for sequencing using 22 µl of sample in 10 µl ExoSap Mix (mix

for 50 reactions; 1.25 µl Exonuclease I [20 U/µl], 2.5 µl Antartic phosphatase [5 U/µl] (New England

BioLabs Inc.), 496.5 µl distilled H_2O). The products were incubated at 37°C for 30 minutes and

heated to 95°C for 5 minutes.

The purified PCR products were sequenced with BigDye terminator kit on Applied Biosystem

3130XL DNA analyser (Applied Biosystems, Foster City, California, USA) at Macrogen Europe,

Amsterdam, the Netherlands. Two sequencing reactions were run for each strain, using the primers

519F (5'-CAGCAGCCGCGGTAATAC-'3) and 926R (5'-CCGTCAATTCCTTTGAG TTT-'3). The

resulting sequences were trimmed using ABI Sequence Scanner (Applied Biosystems), the forward

sequence and the reverse complement of the reverse sequence aligned and combined, and taxonomic

identities obtained using the EzTaxon server (22).

Results

Water chemical analysis revealed several differences in major components at the two sites (Table

2). Although the pH of the water did not differ significantly as judged by a two-tailed Student's t-test,

electrical conductivity was nearly 12-fold higher at the AEX site, significant at the 99.5% confidence

level. The AEX site water contained more than 420-times as much silica as did the SX site water, and

several other ions, such as sodium, potassium chloride and bromide were also found to be present at

significantly higher levels at the AEX site, underscoring the more geothermal character of the

environment (Table 2).

Headspace gas analysis revealed similar amounts of hydrocarbon gas at the two sites (Table 3).

Although methane content was found (n=3) to be lower at the AEX site than at the SX site, the

difference was not deemed significant by a two-tailed Student's t-test (not shown). Isotope

composition suggests a thermogenic origin of the AEX-site headspace gas, whereas a biogenic origin

is suggested for the SX-site headspace gas (Table 3). Thermogenic origin of the AEX gas was further

supported by a high methane/ethane ratio (17.8). Both the composition of the headspace gas and the

methane isotope composition were similar to those reported by Ólafsson *et al*. for borehole gasses in

the Skógalón area (18). EOM fractions showed difference in hydrocarbon content in terms of lower-

chain hydrocarbons in AEX and higher amounts of higher-chain hydrocarbons at the SX site (Fig. 2).

Bacterial community results.

Four technical replicates from each study site were used to compare the microbial communities in

the gas seepage pockmarks of Skógalón (SX) and Skógakíll (AEX). The DNA extraction yielded on

average 2.8 μg/ml (± 0,3 μg/ml) and the amplicon library generated on average 42.8 ng/μL (± 1,3

ng/μL) of amplicons with the length of 587±5 bp. Over 4 million paired sequences, were analysed and

trimmed to the average of 523,241 reads per sample with the length of 301 bp. A total of 595,137

reads generated the OTU table after filtering out chimeric sequences, where predicted OTUs were in

total 26,786 OTUs (Table 4).

Alpha diversity metrics were measured at the depth of 60,000 sequences per sample. The number

of OTUs had reached plateau at 25,000 sequences, meaning the dataset was sufficient to estimate the

diversity of the bacterial communities in the natural seepage pockmarks. The species richness as

estimated with Chao1 index and Shannon´s diversity index are shown in Table 4.

Taxonomic composition

The focus was set on analysing the most abundant taxa, so after filtering out chloroplast OTUs, the

OTUs with the lowest combined abundance (<=1000 reads) were omitted. A total of 14 bacterial

phyla was observed as the most abundant at AEX and SX sites, divided up to 23 classes and 45

observed genera (Fig. 3). *Proteobacteria* was the most abundant phylum at both AEX and SX sites,

28% and 30%, respectively. At the AEX site, *Proteobacteria* was followed by *Chloroflexi* (22%) and

Aminicenantes (10%) at phylum level. At the SX site, the *Bacteroidetes* had significantly higher

abundance than in the AEX site, with relative abundance of 24%, followed by *Chloroflexi* (13%).

The bacterial community structure differed between sites, but only one class, within the phylum of

Aminicenantes was found by likelihood ratio analysis to be significantly more abundant at the AEX

site than at the SX site. An unnamed order within the *Bacteriodetes* was found to be significantly

more abundant at the SX site compared to the AEX. The family of *Syntrophaceae* within the class of

Deltaproteobacteria was more abundant at the SX site with 2.6-fold higher abundance. Three genera

within the *Clostridia* class had over 3.0-fold higher abundance at AEX than SX site.

Cultured microbiota and isolates.

Plate counts after 7 days at 22°C (Table 5) of samples from site SX indicated the presence of

substantial communities of naphthalene and hexane degraders, particularly under aerobic conditions.

One hundred and eighty-six colonies were restreaked for isolation in pure culture (Table 9 in

supplements).

Putatively facultative chemoautotrophs were surprisingly numerous judging by growth on Mn

media, but the extremely restrictive medium 9K only yielded a few colonies, all from sample OX06.

Spraying colonies with fast blue confirmed the presence of alpha-naphthol in some of the colonies on

Naph-agar, but not all. Strains OX0102 and OX0103 tested positive for naphthalene degradation by

fast blue; strains OX0304 and OX0306 tested positive for 2,4-diphenoxyacetate degradation by fast

blue. One hundred and six strains have been identified by partial 16S rDNA sequencing using the

Sanger method and found to comprise 38 genera in 8 classes (Table 6, Table 10 in supplements).

Discussion

The study sites, AEX and SX, were found to be distinct in terms of geochemistry. The AEX site

contained higher concentrations of silica, very similar to the concentrations of previous studies on

geothermal activity in Öxarfjörður (18) indicating geothermal water coming from the pockmarks. The

sodium chloride concentration was higher than in previous studies which implies a mixture of

seawater with the geothermal water. However, the pockmarks at AEX are located in a river delta and

water samples were taken at the pockmark surface so the intermixture of seawater is not surprising.

The water chemistry at the SX site shows low concentration of silica indicating little or no geothermal

activity and less intermixture of seawater than at AEX. The stable isotope ratio $\delta^{13}C$ of methane also

indicates a biogenic origin of the methane at SX, while at AEX the $\delta^{13}C$ suggests a mixture of

thermogenic and biogenic origin of methane, which can most likely be linked to microbial lignite

utilization at the site as well as the geothermal activity previously described (18). The hydrocarbon

content also shows more complex and higher chain hydrocarbons at SX. This can be related to more

vegetation and organic matter accumulation, in contrast with lower-chain hydrocarbons at the AEX

site with less vegetation (Fig.2). These geochemical factors underline how disparate the two sites are:

the AEX pockmarks containing geothermal groundwater with thermogenic methane generation and

the SX pockmarks the result of biogenic natural gas accumulation. The location of the SX site and the

lining up of the pockmarks can easily suggest thermogenic methane seepage at the site, but our

analysis of the SX-site pockmarks explored in this study is more suggestive of marsh gas seepage,

whereas at the AEX site, a thermogenic origin is more strongly supported. The difference in

hydrocarbon content of the two sites, is most likely to explain the difference in biodiversity presented

with biodiversity indices in Table 4. Howewer, further studies are needed to demonstrate the

correlation between hydrocarbon content and biodiversity.

The seepage pockmarks were found to harbour diverse microbiotas consisting largely of anaerobic

heterotrophs. Given the lack of visible vegetation at the AEX site, available organic matter seems

likely to be restricted to the gas seep itself, to a large extent. This kind of environment thus contains a

microbial community composed largely of facultative chemolithotrophs and oxidizers of methane,

lighter alkanes, and aromatics. The inter-site diversity of both sampling sites was notable. However,

several groups of bacteria were shown to vary in relative abundance between the two sampling sites,

as discussed below.

Hydrocarbon and methyl halide metabolism

The high relative abundance of *Dehalococcoidia* in the microbial consortia at the study sites,

particularly site AEX, is noteworthy and underscores the profound effect that petrochemical seepage

has on the composition of the local microbiota. The class *Dehalococcoidia* contains at the present

time only one validly described order (*Dehalococcoidales*), one family (*Dehalococcoidaceae*), and

three genera (*Dehalococcoides, Dehalobium* and *Dehalogenimonas*), comprising a total of four

species, all of which are capable of anaerobic reductive dehalogenation (23–27).

A large fraction of the dehalococcoidal OTUs in this study were found to be assignable to the

order *Dehalococcoidales* (Table 7), where the majority (9120 of 9579) being further assignable to the

family *Dehalococcoidaceae,* containing the described, dehalorespiring members of this class. In

further support of dehalorespiration being an important process in the seepage pockmark microbiotas,

genera known to contain facultative dehalorespirers, like the betaproteobacterial genus

Dechloromonas (28) ant the deltaproteobacterial genus *Anaeromyxobacter* (29), were significantly

more abundant at the AEX site. Furthermore, cultured bacteria from the Öxfjörður seeps, while not

including *Dehalococcoidia*, do include isolates assigned to genera known to include aerobic

facultative dechlorinators, such as *Dechloromonas* and *Shewanella* (Table 7).

The as-yet unnamed and uncharacterized order GIF9 was highly abundant in the AEX and SX

microbiomes (Table 7) and may consist of bacteria that posess other metabolic pathways than just

organohalide respiration. The order is suggested to be an important bacterial group for the degradation

of organic matter in sediments (49). Thus, a recent metagenomic study indicated that some members

of this group may be homoacetogenic fermenters that possess a complete Wood-Ljungdahl CO_2

reduction pathway (31). It should thus be stressed that the presence of a large contingent of

Dehalococcoidia, as was found to be the case in the present study, need not necessarily be indicative

of dehlaorespiration constituting a major metabolic activity in the environment under study. Indeed,

considerable variation in metabolic characteristics occurs in most well-characterized bacterial classes

and hence it must be considered likely that other, perhaps non-dehalorespiring taxa remain to be

characterized within this class.

Recently, it was suggested, in part because of the notable abundance of *Dehalococcoides*, that in

certain Antarctic lakebeds, anaerobic methane oxidation may be fuelled by reductive dehalogenation

(10). The results of the present study are suggestive of the presence of such an ecosystem in the

methane seeps in the Öxarfjörður graben. Methyl halides are often associated with coal combustion

(32), further suggesting subsurface interaction of geothermal matter with lignite as a source of

chloromethane.

Also among abundant groups observed in the present study, the *Atribacteria* (group 'OP9') are

often found to be predominant in methane-rich anaerobic environments such as marine sediments and

subseafloor "mud volcanoes" (43, 44). Although they have not been directly linked to AOM in these

environments, they have been suggested to mediate AOM in some cold seep environments (10). In

general, the *Atribacteria* are thought to play heterotrophic roles, likely fermentative (43, 45), but a

single-cell genomics studies on representative *Atribacteria* suggests that these organisms may be

indirectly responsible for methane production through the production of acetate or CO_2 (43, 46).

Another highly abundant class within the *Chloroflexi* was the *Anaerolineae* (Table 7), a class

originally described as consisting of strictly anaerobic chemo-organotrophs (33), and frequently

detected in subsurface environments (34–37). However, due to the scarcity of cultured

representatives, the metabolic capabilities of this class have remained elusive. A recent study of seven

single-cell genomes from deep submarine hydrothermal vent sediments indicated the presence of a

Wood-Ljungdahl CO_2 reduction pathway, as well as a number of ABC transporters, and in one case a

putative reductive dehalogenase (38). In the present study, the *Anaerolineae* appear fairly diverse,

with 81% of *Chloroflexi* paired-end reads being assigned to four orders: the *Anaerolineales* and three

putative orders without cultured representation, envOPS12, GCA004, and SHA-20 (Table 7).

***Proteobacteria* and the sulfur cycle**

Members of the *Proteobacteria* phylum observed in the seepage pockmarks mainly consisted of

Deltaproteobacteria and *Alphaproteobacteria* (Table 8). The alphaproteobacterial fraction was fairly

homogeneous, consisting mostly of reads assigned to the order *Rhizobacteriales*, of which 77% could

be assigned to a single genus, *Bosea,* a genus of chemolithoheterotrophs noted for their ability to

oxidize inorganic sulfur compounds (39). The deltaproteobacterial fraction was found to be more

diverse although most of the OTUs could be assigned to either of two orders, the *Syntrophobacterales*

and the *Desulfobacterales* (Table 8), both of which contain mostly, albeit not exclusively, sulfate-

reducing organisms.

The *Syntrophobacterales,* known to be frequently associated with anoxic aquatic environments

(40), are significantly enriched in the SX marshland site as compared to the AEX site, perhaps

reflecting an influx of marshland-associated bacteria into the seepage pockmark environment. Most of

the *Syntrophomonadales* reads (77%) can be assigned to the family *Syntrophaceae*, which contains

both sulfate-reducing and non-sulfate-reducing bacteria (41). Many of the *Syntrophaceae* reads (59%)

could not be confidently assigned to genera, rendering the question of the importance of sulfate

reduction of this taxon in the seepage pockmarks unresolved. However, taken together with the high

abundance of the sulfate-reducing order *Desulfobacterales*, sulfate reduction is likely to be a major

process in the seepage pockmarks, likely supporting AOM consortia. Families within

Desulfobacterales have been reported to actively oxidize short and long chain alkanes and are

suggested to be the key alkane degraders in marine seeps (42). Furthermore, considering the high

abundance of the sulfur-oxidizing *Bosea*, we can surmise that *Proteobacteria* consitute an important

driver of sulfur cycling within the seepage pockmark microbiota.

Bioremediative potential of isolated bacterial strains

Microbial communities containing facultative chemolithotrophs and hydrocarbon oxidizers, could

be valuable for bioremediation of petroleum contamination in basaltic oligotrophic environments (54,

55) such as Icelandic beach environments. Previously it has been demonstrated that the microbial

activity in Icelandic soils is more affected by substrate availability than temperature (50, 51), wich

means the possibility of bioremediative strategies like biostimulation is quite possible. Screening

environmental isolates for possible bioremediative capabilities is therefore meaningful for further

research into potential bioremediation in contaminated seashore environments in Iceland. Two of the

isolated strains showed degradation potential of naphthalene. Strain OX0102 was assigned to the

genus *Brevundimonas* (Table 10 in supplements) and Strain OX0103 was most closely related to

bacteria in the genus *Rahnella* (Table 10 in supplements), but strains from both genera have been

isolated with high hydrocarbonoclastic activity in relation to the degradation of naphthalene (52, 53).

Strains OX0304 and OX0306, both assigned by EzTaxon as *Pseudomonas* species (Table 10 in

supplements), tested positive for 2,4-diphenoxy acetate degradation. Strain OX0627 was assigned as a

member of *Dechloromonas*. Isolates from that genus are capable of anaerobic oxidation of benzene

(30) and could possibly be used for bioremediation, as well as the isolated strains mentioned above.

This substantiates the significance of isolated strains presented in the study for future research in

Iceland, related to hydrocarbon degradation and bioremediation.

Concluding remarks

This study reveals natural gas seeps of biogenic origin in Öxarfjörður in addition to known

geothermal gas seepage pockmarks in the Jökulsá-á-Fjöllum river delta, as well as revealing the first

microbial community analysis of gas seepage in Iceland. Further studies are needed to demonstrate

the connection between the gas origin and the pockmark microbiota, establishing the need for further

geomicrobiological research in Icelandic natural gas seeps. The microbial communities associated

with the pockmarks show higher biodiversity at biogenic gas seepage than in thermogenic gas seepage

pockmarks, presenting diverse microbiota that consists largely of anaerobic heterotrophs. The

abundant taxa in the pockmarks indicate that the microbial community is most likely involved in

hydrocarbon degradation linked to sulfur cycling and AOM, and the abundance of *Dehalococcoidia*

suggests the presence of anaerobic reductive dehalogenation in natural gas seepage pockmarks of

thermogenic origin. Isolated strains from the study showed potential in degrading hydrocarbons, such

as naphthalene, and are pivotal for future studies in enviromental biotechnology in Iceland.

Acknowledgments

This work was funded by Orkustofnun and Orkurannsoknarsjodur Landsvirkjunar. We thank Geir

Hansen & Co. at Applied Petroleum Technology for their contribute, and we thank Helga Helgadóttir

and Silja Rúnarsdóttir for their work related to the research. Last but not least we would like to

dedicate this research to our co-author, Þórarinn Sveinn Arnarson, who pased away during the

submission process.

References

1. Link WK. 1952. Significance of oil and gas seeps in world oil exploration. Am Assoc Pet Geol Bull 36:1505–1540.

2. Jones, V. T., & Drozd RJ. 1983. Predictions of oil or gas potential by near-surface geochemistry. Am Assoc Pet Geol Bull 67:932–952.

3. Abrams MA, Dahdah NF. 2010. Surface sediment gases as indicators of subsurface hydrocarbons - examining the record in laboratory and field studies. Mar Pet Geol 27:273–284.

4. Etiope G, Feyzullayev A, Baciu CL. 2009. Terrestrial methane seeps and mud volcanoes: A global perspective of gas origin. Mar Pet Geol 26:333–344.

5. Zengler K, Richnow HH, Rosselló-Mora R, Michaelis W, Widdel F. 1999. Methane formation from long-chain alkanes by anaerobic microorganisms. Nature 401:266–269.

6. Martini AM, Budai JM, Walter LM, Schoell M. 1996. Microbial generation of economic accumulations of methane within a shallow organic-rich shale. Nature 383:155–158.

7. Ritter D, Vinson D, Barnhart E, Akob DM, Fields MW, Cunningham AB, Orem W, McIntosh JC. 2015. Enhanced microbial coalbed methane generation: A review of research, commercial activity, and remaining challenges. Int J Coal Geol 146:28–41.

8. Zhang CL, Lanoil B. 2004. Geomicrobiology and biogeochemistry of gas hydrates and cold seeps. Chem Geol 205:187–194.

9. Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM, Schreiber F, Dutilh BE, Zedelius J, de Beer D, Gloerich J, Wessels HJCT, van Alen T, Luesken F, Wu ML, van de Pas-Schoonen KT, Op den Camp HJM, Janssen-Megens EM, Francoijs K-J, Stunnenberg H, Weissenbach J, Jetten MSM, Strous M. 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. Nature 464:543–548.

10. Saxton MA, Samarkin VA, Schutte CA, Bowles MW, Madigan MT, Cadieux SB, Pratt LM, Joye SB. 2016. Biogeochemical and 16S rRNA gene sequence evidence supports a novel mode of anaerobic methanotrophy in permanently ice-covered Lake Fryxell, Antarctica.

448    Limnol Oceanogr 61:S119–S130.

449 11. Mills HJ, Martinez RJ, Story S, Sobecky PA, Sobecky A. 2005. Characterization of Microbial

450    Community Structure in Gulf of Mexico Gas Hydrates : Comparative Analysis of DNA- and

451    RNA-Derived Clone Libraries Characterization of Microbial Community Structure in Gulf of

452    Mexico Gas Hydrates : Comparative Analysis of DNA-. Appl Environ Microbiol 71:3235–

453    3247.

454 12. Inagaki F, Nunoura T, Nakagawa S, Teske A, Lever M, Lauer A, Suzuki M, Takai K,

455    Delwiche M, Colwell FS, Nealson KH, Horikoshi K, D'Hondt S, Jørgensen BB. 2006.

456    Biogeographical distribution and diversity of microbes in methane hydrate-bearing deep

457    marine sediments on the Pacific Ocean Margin. Proc Natl Acad Sci U S A 103:2815–20.

458 13. Knittel K, Boetius A, Lemke A, Eilers H, Lochte K, Pfannkuche O, Linke P, Amann R. 2003.

459    Activity, Distribution, and Diversity of Sulfate Reducers and Other Bacteria in Sediments

460    above Gas Hydrate (Cascadia Margin, Oregon). Geomicrobiol J 20:269–294.

461 14. Niemann H, Lösekann T, de Beer D, Elvert M, Nadalig T, Knittel K, Amann R, Sauter EJ,

462    Schlüter M, Klages M, Foucher JP, Boetius A. 2006. Novel microbial communities of the

463    Haakon Mosby mud volcano and their role as a methane sink. Nature 443:854–858.

464 15. Flóvenz ÓG, Gunnarsson K. 1991. Seismic crustal structure in Iceland and surrounding area.

465    Tectonophysics 189:1–17.

466 16. Gunnarsson K. 1998. Sedimentary basins of the N-Iceland shelf. Draft Version for Discussion

467    (April–May 1998). Report OS-98014. Reykjavík.

468 17. Friðleifsson GÓ. 1994. Geothermal gradient and hydrothermal systems off North Iceland.

469    Statement GÓF-94-05. Reykjavík.

470 18. Ólafsson M, Friðleifsson GÓ, Eiríksson J, Sigvaldason H, Ármansson H. 1993. On the origin

471    of organic gas in Öxarfjörður, NE-Iceland. Report No. OS-93015/JHD-05. Reykjavík.

472 19. Richter B, Gunnarsson K. 2010. Overview of hydrocarbon related research in Tjörnes.

473    GagnavefsjaÍs.

474 31. Sæmundsson K, Hjartarson Á, Kaldal I, Sigurgeirsson MA, Kristinsson SG, S V. 2017.

475    Jarðfræðikort af Norðurgosbelti. Nyrðri hluti. 1:100.000. Íslenskar Orkurannsóknir and

Landsvirkjun., Reykjavík.

21.   Silverman MP, Lundgren DG. 1959. Studies on the chemoautotrophic iron bacterium

Ferrobacillus ferrooxidans. J Bacteriol 77:642–647.

22.   Kim O-S, Cho Y-J, Lee K, Yoon S-H, Kim M, Na H, Park S-C, Jeon YS, Lee J-H, Yi H, Won

S, Chun J. 2012. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database

with phylotypes that represent uncultured species. Int J Syst Evol Microbiol 62:716–21.

23.   Löffler F, Yan J, Ritalahti K, Adrian L, Edwards E, Konstantinidis K, Müller J, Fullerton H,

Zinder S SA. 2013. Dehalococcoides mccartyi gen. nov., sp. nov., obligately organohalide-

respiring anaerobic bacteria relevant to halogen cycling and bioremediation, belong to a novel

bacterial class, Dehalococcoidia classis nov., order Dehalococcoidales ord. nov. and famil. Int

J Syst Evol Microbiol 63:625–635.

24.   Bowman KS, Nobre MF, da Costa MS, Rainey F a, Moe WM. 2012. Dehalogenimonas

alkenigignens sp. nov., a chlorinated alkane dehalogenating bacterium isolated from

groundwater. Int J Syst Evol Microbiol.

25.   Moe WM, Yan J, Nobre MF, da Costa MS, Rainey FA. 2009. Dehalogenimonas

lykanthroporepellens gen. nov., sp. nov., a reductively dehalogenating bacterium isolated from

chlorinated solvent-contaminated groundwater. Int J Syst Evol Microbiol 59:2692–2697.

26.   May HD, Sowers KR. 2016. "Dehalobium chlorocoercia" DF-1-from discovery to application,

p. 563–586. *In* Organohalide-Respiring Bacteria.

27.   Nuzzo A, Negroni A, Zanaroli G, Fava F. 2017. Identification of two organohalide-respiring

Dehalococcoidia associated to different dechlorination activities in PCB-impacted marine

sediments. Microb Cell Fact 16:127.

28.   Achenbach LA, Michaelidou U, Bruce RA, Fryman J, Coates JD. 2001. Dechloromonas

agitata gen. nov., sp. nov. and Dechlorosoma suillum gen. nov., sp. nov., two novel

environmentally dominant (per)chlorate-reducing bacteria and their phylogenetic position. Int

J Syst Evol Microbiol 51:527–533.

29.   Sanford RA, Cole JR, Tiedje JM. 2002. Characterization and description of Anaeromyxobacter

dehalogenans gen. nov., sp. nov., an aryl-halorespiring facultative anaerobic myxobacterium.

504   Appl Environ Microbiol 68:893–900.

505 30. Coates JD, Chakraborty R, Lack JG, O'Connor SM, Cole KA, Bender KS, Achenbach LA.

506   2001. Anaerobic benzene oxidation coupled to nitrate reduction in pure culture by two strains

507   of Dechloromonas. Nature 411:1039–1043.

508 31. Hug LA, Castelle CJ, Wrighton KC, Thomas BC, Sharon I, Frischkorn KR, Williams KH,

509   Tringe SG, Banfield JF. 2013. Community genomic analyses constrain the distribution of

510   metabolic traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling.

511   Microbiome 1:22.

512 32. McCulloch A, Aucott ML, Benkovitz CM, Graedel TE, Kleiman G, Midgley PM, Li Y-F.

513   1999. Global emissions of hydrogen chloride and chloromethane from coal combustion,

514   incineration and industrial activities: Reactive Chlorine Emissions Inventory. J Geophys Res

515   104:8391–8403.

516 33. Yamada T, Sekiguchi Y, Hanada S, Imachi H, Ohashi A, Harada H, Kamagata Y. 2006.

517   Anaerolinea thermolimosa sp. nov., Levilinea saccharolytica gen. nov., sp. nov. and Leptolinea

518   tardivitalis gen. nov., sp. nov., novel filamentous anaerobes, and description of the new classes

519   Anaerolineae classis nov. and Caldilineae classis nov. in the . Int J Syst Evol Microbiol

520   56:1331–1340.

521 34. Schippers A, Kock D, Höft C, Köweker G, Siegert M. 2012. Quantification of Microbial

522   Communities in Subsurface Marine Sediments of the Black Sea and off Namibia. Front

523   Microbiol 3:16.

524 35. Galand PE, Potvin M, Casamayor EO, Lovejoy C. 2010. Hydrography shapes bacterial

525   biogeography of the deep Arctic Ocean. ISME J 4:564–576.

526 36. Kato S, Kobayashi C, Kakegawa T, Yamagishi A. 2009. Microbial communities in iron-silica-

527   rich microbial mats at deep-sea hydrothermal fields of the Southern Mariana Trough. Environ

528   Microbiol 11:2094–2111.

529 37. Cleary DFR, Coelho FJRC, Oliveira V, Gomes NCM, Polónia ARM. 2017. Sediment depth

530   and habitat as predictors of the diversity and composition of sediment bacterial communities in

531   an inter-tidal estuarine environment. Mar Ecol 38:e12411.

38.   Fullerton H, Moyer CL. 2016. Comparative Single-Cell Genomics of Chloroflexi from the

Okinawa Trough Deep-Subsurface Biosphere. Appl Environ Microbiol 82:3000–3008.

39.   Das SK. 2015. BoseaBergey's Manual of Systematics of Archaea and Bacteria. John Wiley &

Sons, Ltd.

40.   Kuever J, Rainey FA, Widdel F. 2015. Syntrophobacterales ord. novBergey's Manual of

Systematics of Archaea and Bacteria. John Wiley & Sons, Ltd.

41.   Kuever J, Rainey FA, Widdel F. 2015. Syntrophaceae fam. nov.Bergey's Manual of

Systematics of Archaea and Bacteria. John Wiley & Sons, Ltd.

42.   Kleindienst S, Herbst F-A, Stagars M, von Netzer F, von Bergen M, Seifert J, Peplies J,

Amann R, Musat F, Lueders T, Knittel K. 2014. Diverse sulfate-reducing bacteria of the

Desulfosarcina/Desulfococcus clade are the key alkane degraders at marine seeps. ISME J

8:2029–2044.

43.   Carr SA, Orcutt BN, Mandernack KW, Spear JR. 2015. Abundant Atribacteria in deep marine

sediment from the Adélie Basin, Antarctica. Front Microbiol 6:872.

44.   Hoshino T, Toki T, Ijiri A, Morono Y, Machiyama H, Ashi J, Okamura K, Inagaki F. 2017.

Atribacteria from the Subseafloor Sedimentary Biosphere Disperse to the Hydrosphere

through Submarine Mud Volcanoes. Front Microbiol 8:1135.

45.   Dodsworth JA, Blainey PC, Murugapiran SK, Swingley WD, Ross CA, Tringe SG, Chain

PSG, Scholz MB, Lo C-C, Raymond J, Quake SR, Hedlund BP. 2013. Single-cell and

metagenomic analyses indicate a fermentative and saccharolytic lifestyle for members of the

OP9 lineage. Nat Commun 4:1854.

46.   Nobu MK, Dodsworth JA, Murugapiran SK, Rinke C, Gies EA, Webster G, Schwientek P,

Kille P, Parkes RJ, Sass H, Jørgensen BB, Weightman AJ, Liu W-T, Hallam SJ, Tsiamis G,

Woyke T, Hedlund BP. 2016. Phylogeny and physiology of candidate phylum "Atribacteria"

(OP9/JS1) inferred from cultivation-independent genomics. ISME J 10:273–8

47.   Valentine DL. 2002. Biogeochemistry and microbial ecology of methane oxidation in anoxic

environments: a review. Antonie Van Leeuwenhoek 81:271-282

48.   Arnórsson S, Bjarnason JÖ, Giroud N, Gunnarsson I, Stefánsson A. 2006. Sampling and

analysis of geothermal fluids. Geofluids 6:203-216.

49.   Oni OE, Schmidt F, Miyatake T, Kasten S, Witt M, Hinrichs KU, Friedrich MW. 2015.

Microbial communities and organic matter composition in surface and subsurface sediments of

the Helgoland mud area, North Sea. Front Microbiol 6:1290.

50.   Guicharnaud R, Arnalds O, Paton GI. 2010. Short term changes of microbial processes in

Icelandic soils to increasing temperatures. Biogeosciences 7:671-82.

51.   Lehtinen T, Mikkonen A, Sigfusson B, Ólafsdóttir K, Ragnarsdóttir KV, Guicharnaud R. 2014.

Bioremediation trial on aged PCB-polluted soils—a bench study in Iceland. Environ Sci Pollut

Res 21:1759-68.

52.   Mansur AA, Adetutu EM, Kadali KK, Morrison PD, Nurulita Y, Ball AS. 2014. Assessing the

hydrocarbon degrading potential of indigenous bacteria isolated from crude oil tank bottom

sludge and hydrocarbon-contaminated soil of Azzawiya oil refinery, Libya. Environ Sci Pollut

Res 21:10725-35.

53.   Ma Y, Wang L, Shao Z. 2006. Pseudomonas, the dominant polycyclic aromatic hydrocarbon-

degrading bacteria isolated from Antarctic soils and the role of large plasmids in horizontal

gene transfer. Environ Microbiol 8:455-65.

54.   Markúsdóttir M, Heiðmarsson S, Eyþórsdóttir A, Magnússon KP, Vilhelmsson O. 2013. The

natural and anthropogenic microbiota of Glerá, a sub-arctic river in northeastern Iceland. Int

Biodeterior Biodegradation 84:192-203.

55.   Jóelsson JP, Friðjónsdóttir H, Vilhelmsson O. 2013. Bioprospecting a glacial river in Iceland

for bacterial biopolymer degraders. Cold Reg Sci Technol 96:86-95.

Table 1. Location and description of sampling sites. Temperature and pH were measured in situ *with a hand-held probe.*

Site	Location	Coordinates	Description	pH	T (°C)
AEX-A	Skógakíll	66°10'2.250 N 16°33'59.687 W	Gas seepage sediment	6,4	35,2
AEX-B	Skógakíll	66°10'8.014 N 16°33'56.422 W	Gas seepage water	6,4	53,3
AEX-C	Skógakíll	66°10'12.029 N 16°33'56.998 W	Mud-mire	ND	30,4
AEX-D	Skógakíll	66°10'9.554 N 16°33'57.793 W	Sand-mud	6,4	19,3
SX-A	Skógalón	66°09.222 N 016°37.089 W	Lagoon sediment	8,5	12,1
SX-B	Skógalón	66°09.244 N 016°37.240 W	Puddle water	8,5	13,4

Table 2. Major physicochemical characteristics[1] of water from the two study sites.

	AEX	SX	
pH at 21°C	7.57 ± 0.31	8.64 ± 0.73	(p = 0.19)
Conductivity at 25°C (μScm^{-1})	8945 ± 728	773 ± 392	(p = 0.005)
DIC (as ppm CO_2)	44.8 ± 31.0	94.5 ± 43.1	(p = 0.32)
SiO_2 (ppm)	147.5 ± 2.1	0.35 ± 0.07	(p = 0.0001)
Na (ppm)	1900 ± 56.6	107.4 ± 57.5	(p = 0.001)
K (ppm)	104.9 ± 18.6	4.01 ± 1.96	(p = 0.02)
Mg (ppm)	24.0 ± 15.9	17.8 ± 10.5	(p = 0.69)
Ca (ppm)	311.5 ± 111.0	17.1 ± 9.0	(p = 0.06)
F (ppm)	0.45 ± 0.13	0.21 ± 0.01	(p = 0.11)
Cl (ppm)	3355 ± 276	169.4 ± 105.5	(p = 0.004)
Br (ppm)	12.2 ± 0.8	0.34 ± 0.31	(p = 0.002)
SO_4 (ppm)	215 ± 76.4	5.13 ± 0.60	(p = 0.06)

[1]Dissolved sulphide was determined on-site by titration with mercuric acetate using dithizone as indicator (48). Dissolved inorganic carbon was determined by alkalinity titration and back-titration (48). Silica was analysed by colorimetric determination of a silica-molybdate complex at 410 nm. Total dissolved solids were determined by gravimetry. Anions were determined by suppressed ion chromatography. Major metals were analysed by atomic absorption spectrometry.

Table 3. Headspace gas analysis[1] on sediment samples from seepage pockmarks at the two study sites. The concentrations are shown as parts of Total Hydrocarbon Gas (THCG).

	AEX	SX
N_2 [%total]	39.2 ($\pm$31.5)	81.2 ($\pm$2.62)
O_2 + Ar [%total]	8.46 ($\pm$8.53)	14.10 ($\pm$5.66)
ppm THCG	5.09×10^5 ($\pm 3.88 \times 10^5$)	0.389×10^5 ($\pm 0.296 \times 10^5$)
CO_2 [%THCG]	99.9 ($\pm$0.23)	93.2 ($\pm$1.77)
Methane (CH_4) [%THCG]	46.1×10^{-3} ($\pm 48.3 \times 10^{-3}$)	6.78 ($\pm$1.66)
Ethane (C_2H_6) [%THCG]	2.60×10^{-3} ($\pm 4.50 \times 10^{-3}$)	0 ($\pm$0)
Propane (C_3H_8) [%THCG]	1.03×10^{-3} ($\pm 1.79 \times 10^{-3}$)	0 ($\pm$0)
iso-Pentane (i-C_5H_{12}) [%THCG]	18.7×10^{-3} ($\pm 32.3 \times 10^{-3}$)	8.00×10^{-3} ($\pm 11.3 \times 10^{-3}$)
Pentane (n-C_5H_{12}) [%THCG]	73.7×10^{-3} ($\pm 127 \times 10^{-3}$)	49.5×10^{-3} ($\pm 70.0 \times 10^{-3}$)
Benzene (C_6H_6) [%THCG]	3.80×10^{-3} ($\pm 6.24 \times 10^{-3}$)	7.15×10^{-3} ($\pm 5.45 \times 10^{-3}$)
Methane $\delta^{13}C$ [‰]	-26.6	-63.2
CO_2 $\delta^{13}C$ [‰]	-3.3	nd

[1]Samples were extracted in dichloromethane/methanol, deasphalted in pentane, and analysed by gas chromatography.

Table 4. Number of predicted operational taxanomic units (OTUs) and alpha diversity metrics as calculated at 25,000 sequences from the two study sites.

	Paired seqs	Reads in OTUs	Predict OTUs	Alpha diversity metrics		
				OTUs	Shannon	Chao1
AEX	2.064.234	307.384	14.019	3.461 ± 623	8,3 ± 0,5	3.980 ± 751
SX	2.121.996	287.753	12.767	3.762 ± 333	9,1 ± 0,2	4.203 ± 504

Table 5. Colony-forming units per gram sediment sample after 7 days at 22°C on selective media.

	OX01	OX01	OX03	OX03	OX06	OX06
	Aerobic	Anaerobic	Aerobic	Anaerobic	Aerobic	Anaerobic
R2A	1.4×10^5	2.8×10^4	5.0×10^5	2.4×10^5	1.7×10^7	5.0×10^6
Naph	2.0×10^3	$<1\times10^3$	$<1\times10^3$	$<1\times10^3$	2.5×10^5	1.0×10^5
Hex	7.5×10^4	$<1\times10^3$	2.0×10^5	$<1\times10^3$	$>2.5\times10^5$	2.0×10^5
2,4-D	nd	nd	$<1\times10^3$	$<1\times10^3$	nd	nd
Mn	9.3×10^4	5.0×10^3	2.0×10^5	$<1\times10^3$	$>2.5\times10^5$	7.5×10^4
9K	$<1\times10^3$	$<1\times10^3$	$<1\times10^3$	$<1\times10^3$	1.0×10^3	3.0×10^3
Gui	nd	nd	nd	nd	1.0×10^6	3.0×10^6

Table 6. Bacterial isolates and their taxonomic classification by partial 16S rRNA gene sequencing.

Strains	GenBank accession numbers	EzTaxon classification			
		Class	Order	Family	Genus
SX0206, SX1205	MG575948, MG575949	*Actinobacteria*	*Corynebacteriales*	*Nocardiaceae*	*Rhodococcus*
OX0615	MG575950	´´	*Micrococcales*	*Cellulomonadaceae*	*Oerskovia*
OX0315, OX0319	MG575951, MG575952	´´	´´	*Microbacteriaceae*	*Cryobacterium*
OX0117, OX0308, OX0313	MG575953-MG575955	´´	´´	*Micrococcaceae*	*Arthrobacter*
OX0107	MG575956	´´	´´	´´	*Pseudarthrobacter*
OX0614	MG575957	´´	´´	*Sanguibacteraceae*	*Sanguibacter*
OX0316	MG575958	´´	*Streptomycetales*	*Streptomycetaceae*	*Streptomyces*
OX0104, OX0118	MG575959, MG575960	*Cytophagia*	*Cytophagales*	*Cyclobacteriaceae*	*Algoriphagus*
OX0108, OX0126, OX0312	MG575961-MG575963	´´	´´	´´	*Aquiflexum*
OX0623, OX0125, OX0129, OX0311, OX0122, OX0123, OX0314, OX0127	MG575964-MG575971	*Flavobacteriia*	*Flavobacteriales*	*Flavobacteriaceae*	*Flavobacterium*
OX0625	MG575972	*Sphingobacteria*	*Sphingobacteriales*	*Sphingobacteriaceae*	*Pedobacter*
OX1213, OX2513, OX1505, OX2509, OX1011, OX1604, OX1805, OX1004, OX1006, OX1007, OX1210, OX1212, OX2308, OX2309, OX2506	MG575973 - MG575987	*Bacilli*	*Bacillales*	*Bacillaceae*	*Bacillus*
OX0317	MG575988	´´	´´	´´	*Psychrobacillus*
OX2205, OX0307, OX2310, OX0301, OX0302	MG575989 - MG575993	´´	´´	*Paenibacillaceae*	*Paenibacillus*
OX0310	MG575994	´´	´´	*Planococcaceae*	*Jeotgalibacillus*
OX0626	MG575995	´´	´´	*Staphylococcaceae*	*Staphylococcus*
OX1107, OX1109, OX2515, OX2106, OX2313	MG575996-MG576000	´´	*Lactobacillales*	*incertae sedis*	*Exiguobacterium*
OX0102, OX2008, OX0309	MG576001-MG576003	*Alphaproteobacteria*	*Caulobacterales*	*Caulobacteraceae*	*Brevundimonas*
OX0106, OX1214, OX0119, SX1214	MG576004-MG576007	´´	*Rhizobiales*	*Rhizobiaceae*	*Rhizobium*
OX0632	MG576008	´´	*Rhodobacterales*	*Rhodobacteraceae*	*Cereibacter*
OX1314	MG576009	´´	´´	´´	*Paracoccus*

SX0604	MG576010	´´	*Sphingomonadales*	*Erythrobacteraceae*	*Porphyrobacter*
OX0620	MG576011	´´	´´	*Sphingomonadaceae*	*Sphingobium*
OX1313, OX1403, OX1702, OX1216	MG576012-MG576015	*Betaproteobacteria*	*Burkholderiales*	*Burkholderiaceae*	*Paraburkholderia*
OX0105, OX0120, OX0124	MG576016-MG576018	´´	´´	*Comamonadaceae*	*Acidovorax*
OX0630, OX0321	MG576019, MG576020	´´	´´	´´	*Rhodoferax*
OX0130	MG576021	´´	´´	´´	*Variovorax*
OX0611	MG576022	´´	´´	*incertae sedis*	*Paucibacter*
OX0627	MG576023	´´	*Rhodocyclales*	*Rhodocyclaceae*	*Dechloromonas*
OX0617, OX0622, SX0303, OX0110, OX0112, OX0612	MG576024-MG576029	*Gammaproteobacteria*	*Aeromonadales*	*Aeromonadaceae*	*Aeromonas*
OX0619	MG576030	´´	*Alteromonadales*	*Shewanellaceae*	*Shewanella*
OX1909, OX1208	MG576031-MG576032	´´	*Chromatiales*	*Chromatiaceae*	*Rheinheimera*
OX1012	MG576033	´´	*Enterobacteriales*	*Enterobacteriaceae*	*Escherichia*
OX0103, OX0101	MG576034-MG576035	´´	´´	´´	*Rahnella*
OX0604, OX0606, OX0607	MG576036-MG576038	´´	´´	´´	*Shigella*
OX0601, OX0621	MG576039-MG576040	´´	*Pseudomonadales*	*Moraxellaceae*	*Acinetobacter*
OX1807, OX1808, OX0306, OX0322, OX1110, OX0613, SX0305, SX0307, SX1216, SX0304, OX0631, SX1213, OX0304	MG576041-MG576053	´´	´´	*Pseudomonadaceae*	*Pseudomonas*

Table 7. Fractional abundance of Chloroflexi *classes and orders in the seepage pockmarks microbiomes as determined with amplicon sequencing.*

Class	AEX[1]	SX[1]	Order	AEX[2]	SX[2]
Anaerolineae	**0.42**	**0.63**			
			Anaerolineales	0.31	0.22
			'envOPS12'	**0.21**	**0.43**
			'GCA004'	0.09	0.13
			'SHA-20'	**0.20**	**0.04**
			Others	0.19	0.19
Dehalococcoidia	**0.53**	**0.27**			
			Dehalococcoidales	0.18	0.48
			'GIF9'	0.76	0.32
			'FS117-23B-02'	**0.02**	**0.06**
			Others	0.04	0.14
'Ellin6529'	0.03	0.06			
'S085'	0.02	0.02			
Others	0.00	0.01			

[1]Fraction of *Chloroflexi* paired-end sequences (total of 86,729 paired-end reads). Figures in **bold** indicate a significant difference between the AEX and SX microbiota by a two-tailed two-sample Student's *t*-test (p<0.05).

[2]Fraction of class-level paired-end sequences (total of 44,384 paired-end *Anaerolineae* reads and 36,549 paired-end *Dehalococcoidia* reads). Figures in **bold** indicate a significant difference between the AEX and SX microbiota by a two-tailed two-sample Student's *t*-test (p<0.05).

Table 8. Fractional abundance of Proteobacteria *classes and orders in the seepage pockmarks microbiomes as determined with amplicon sequencing.*

Class	AEX[1]	SX[1]	Order	AEX[2]	SX[2]
Delta-proteobacteria	**0.43**	**0.23**			
			Syntrophobacterales	**0.27**	**0.55**
			Desulfobacterales	**0.26**	**0.07**
			Myxococcales	0.07	0.08
			Desulfarculales	0.06	0.06
			Desulfuromonadales	**0.06**	**0.04**
			'BCP076'	0.14	0.06
			Others	0.15	0.12
Alpha-proteobacteria	**0.48**	**0.63**			
			Rhizobiales	0.88	0.92
			Rhodobacterales	0.06	0.02
			Others	0.06	0.05
Gamma-proteobacteria	**0.11**	**0.06**			
Beta-proteobacteria	0.06	0.07			
Others	0.02	0.01			

[1]Fraction of *Proteobacteria* paired-end sequences (total of 152,153 paired-end reads). Figures in **bold** indicate a significant difference between the AEX and SX microbiota by a two-tailed two-sample Student's *t*-test ($p<0.05$).

[2]Fraction of class-level paired-end sequences (total of 83,764 paired-end *Deltaproteobacteria* reads and 43,823 paired-end *Alphaproteobacteria* reads). Figures in **bold** indicate a significant difference between the AEX and SX microbiota by a two-tailed two-sample Student's *t*-test ($p<0.05$).

Table 9. Description of strains, cultured and isolated on differential and selective media, and conserved in 30% glycerol at -70 °C.

Strain no.	Media	T (°C)	Description	Strain no.	Media	T (°C)	Description
OX0101	Naph	22	White	OX0306	2,4-D	22	Tan
OX0102	Naph	22	White	OX0307	2,4-D	22	Greyish white
OX0103	Naph	22	White	OX0308	2,4-D	22	White
OX0104	HEX	22	Pink	OX0309	2,4-D	22	White
OX0105	HEX	22	Clear-white	OX0310	HEX	22	Tan
OX0106	HEX	22	Yellow, shiny	OX0311	HEX	22	Pink
OX0107	HEX	22	White	OX0312	HEX	22	Pale pink, flat, somewhat swarmy
OX0108	HEX	22	Pink-clear	OX0313	HEX	22	Yellow
OX0109	HEX	22	Pink	OX0314	HEX	22	White
OX0110	HEX	22	White, small	OX0315	HEX	22	Clear-white, swarming
OX0111	HEX	22	Yellow	OX0316	Mn	22	White with a grey centre
OX0112	HEX	22	Clear-white, swarming	OX0317	Mn	22	Buff
OX0113	HEX	22	White	OX0318	Mn	22	Red
OX0116	Mn	22	White with a grey centre	OX0319	R2A	22	Bright yellow
OX0117	Mn	22	bright yellow	OX0320	R2A	22	White, irregular
OX0118	Mn	22	Pink	OX0321	R2A	22	Bright red
OX0119	Mn	22	snowy white	OX0322	R2A	22	Light yellowish brown, irregular
OX0120	Mn	22	Clear-yellow "fried egg"	OX0323	R2A	22	Very slightly buff
OX0121	Mn	22	Yellow, shiny	OX0601	Naph	22	White
OX0122	R2A	22	Bright orange	OX0602	Naph	22	Orange
OX0123	R2A	22	Bright orange	OX0603	Naph	22	White, flat
OX0127	R2A	22	Dark yellow	OX0607	Naph	22	White

Strain no.	Media	T (°C)	Description	Strain no.	Media	T (°C)	Description
OX0128	R2A	22	White w. dark red peak, coherent	OX0608	LAS	22	Snowy white
OX0129	R2A	22	Bright yellow	OX0611	HEX	22	White, shiny/oily
OX0130	R2A	22	Pale yellow	OX0612	HEX	22	White, shiny/oily
OX0620	Gui	22	bright yellow	OX1215	Hex	22°C	Orange
OX0621	Gui	22	White, runny consistency	OX1216	BS4 (R2A)*	15°C	Transparent-white, transparent edge
OX0622	Gui	22	Buff, lobate margins	OX17B03	R2A	22°C	Very transparent
OX0623	Gui	22	bright yellow	OX17B05	R2A	22°C	Light-pink
OX0624	Gui	22	Buff	OX17B06	R2A	22°C	Transparent
OX0625	R2A	22	Bright red	OX1804	R2A	22°C	Red, crawls a lot
OX0626	R2A	22	Pastel yellow	OX1805	R2A	22°C	White-transparent, crawls
OX0627	R2A	22	Yellow "fried egg"	OX1806	Hex	22°C	White, round
OX0628	R2A	22	White	OX1807	Hex	22°C	Light yellow, slimy
OX0629	R2A	22	Clear-white	OX1808	Hex	22°C	Pink
OX0630	R2A	22	Reddish brown	OX1809	BS4 (R2A)*	15°	White-transparent, transparent edge
OX0631	R2A	22	Bright yellow with a clear, flat halo	OX1905	R2A	22°C	Transparent-White, crawls
OX0632	R2A	22	Pink	OX1906	R2A	22°C	White, irregular, crawls
OX0633	R2A	22	Red	OX1907	R2A	22°C	Yellow, lighter edge, crawls
OX1004	R2A	22°C	Red, irregular	OX1908	Hex	22°C	White, small, irregular
OX1005	R2A	22°C	Small light, crawl much	OX1909	Hex	22°C	Transparent, slimy
OX1006	R2A	22°C	Yellow, small, very irregular	OX2005	R2A	22°C	Very light-yellow, crawls, small
OX1007	R2A	22°C	Drab (yellow-ish even), irregular	OX2006	R2A	22°C	Light-drab, small, crawls alot
OX1008	R2A	22°C	Light-orange, crawls	OX2007	R2A	22°C	Orange, crawls a lot, lighter edge

Strain no.	Media	T (°C)	Description	Strain no.	Media	T (°C)	Description
OX1011	Hex	15°C	White	OX2103	R2A	22°C	Light-transparent
OX1012	BS4 (R2A)*	35°C	White-transparent, small	OX2104	R2A	22°C	White-transpaeant, crawls a lot
OX1105	R2A	22°C	Small, transparent, round	OX2105	R2A	22°C	Yellow, crawls
OX1106	R2A	22°C	Small, drab (yellow-ish)	OX2106	Hex	22°C	Orange, round
OX1107	R2A	22°C	Light-orange, lighter irregular edge	OX2107	Hex	22°C	Light-orange, round
OX1211	R2A	22°C	Yellow, very irregular, lighter edge	OX1502	R2A	22°C	Drab, regular, thin edge
OX1212	R2A	22°C	White, round	OX1503	R2A	22°C	Creamy, irregular edge
OX1213	R2A	22°C	Orange, very irregular, lighter edge	OX1504	R2A	22°C	Drab, crawls
OX1214	Hex	22°C	White, small	OX1505	Hex	15°C	Very light orange, small
OX1602	R2A	22°C	White-transparent, thin edge	SX1204	R2A	22°C	Light-tranparent, transparent edge
OX1603	R2A	22°C	Yellow, regular, stuck to the agar	SX1205	R2A	22°C	Light-pink, cramped, crawls a lot
OX1604	Hex	15°C	White, small	SX1206	R2A	22°C	Light-pink, small
OX1702	R2A	22°C	Creamy, transparent, irregular edge	SX1208	Hex	22°C	Light-transparent
OX2208	R2A	22°C	Light-orange, crawls	SX1209	Hex	22°C	Light orange, round
OX2209	Hex	22°C	White, small, crawls	SX1210	Hex	22°C	Light-white, small
OX2210	Hex	22°C	White, small	SX1212	Hex	22°C	White, round
OX2305	R2A	22°C	Light-transparent, crawls	SX1213	Hex	22°C	White, very slimy
OX2307	R2A	22°C	Orange, lighter edge, irregular	SX1214	Hex	22°C	White-transparent, small
OX2308	R2A	22°C	Orange, round	SX1215	Hex	22°C	Light-drab, round
Strain no.	Media	T (°C)	Description	Strain no.	Media	T (°C)	Description

OX2309	R2A	22°C	Red-pink, round, lighter edge	**SX201**	R2A	22°C	White-transparent, crawls a lot
OX2310	R2A	22°C	White w/ zone	**SX202**	R2A	22°C	Light-transparent-yellow, crawls
OX2311	R2A	22°C	Orange, shines, crawls	**SX203**	R2A	22°C	Pink, crawls, slimy
OX2312	R2A	22°C	Light-transparent, crawls	**SX204**	R2A	22°C	Light-pink, big transparent edge
OX2313	Hex	22°C	Orange, slimy, transparent	**SX301**	R2A	22°C	Light-pink, big, slimy
OX2314	BS4 (R2A)*	22°C	White-transparent, transparent edge	**SX504**	R2A	22°C	White, irregular, crawls a lot
OX2404	R2A	22°C	Light-yellow, crawls	**SX601**	R2A	22°C	White, lighter edge, crawls a lot
OX2405	R2A	22°C	Light semi-transparent, crawls	**SX602**	R2A	22°C	White, crawls
OX2406	Hex	22°C	Light-orange, irregular	**SX0505**	Hex	22°C	Pink, round
OX2514	Hex	22°C	Light orange	**SX1201**	R2A	22°C	White, crawls a lot
OX2515	Hex	22°C	Orange	**SX1202**	R2A	22°C	Shiny white, crawls
OX2516	9K (R2A)*	15°	Very light-pink, small, round	**SX0501**	R2A	22°C	Light-pink, lighter edge, slimy
SX0302	R2A	22°C	Light-white, crawls (long branches)	**SX0502**	R2A	22°C	Pink, crawls
SX0303	Hex	22°C	Transparent	**SX0503**	R2A	22°C	Dark-yellow, crawls a lot, very slimy
SX0304	Hex	22°C	White, small , round				

Table 10. Taxanomic assignment of strains identified by 16S rDNA sequencing and their GenBanka ccession numbers.

Strain	EzTaxon assignment	Seq. length	GenBank accession number	%ID	Order	Site[1]
Alphaproteobacteria						
OX0102	*Brevundimonas bullata*	636	MG576001	99.8	Caulobacterales	SX
OX0309	*Brevundimonas bullata*	1394	MG576003	99.9	Caulobacterales	SX
OX2008	*Brevundimonas halotolerans*	1423	MG576002	98.9	Caulobacterales	AEX
SX0604	*Porphyrobacter colymbi*	349	MG576010	98.6	Sphingomonadales	SX
OX0620	*Sphingobium xenophagum*	1451	MG576011	99.4	Sphingomonadales	SX
OX0106	*Rhizobium selenitireducens*	804	MG576004	99.9	Rhizobiales	SX
OX0119	*Rhizobium selenitireducens*	1131	MG576007	99.9	Rhizobiales	SX
OX1214	*Rhizobium sphaerophysae*	872	MG576006	98.1	Rhizobiales	AEX
SX1214	*Rhizobium selenitireducens*	1447	MG576005	97.3	Rhizobiales	SX
OX1314	*Paracoccus homiensis*	721	MG576009	98.1	Rhodobacterales	AEX
OX0632	*Cereibacter changlensis*	1396	MG576008	99.9	Rhodobacterales	SX
Betaproteobacteria						
OX0105	*Acidovorax radicis*	744	MG576016	98.8	Burkholderiales	SX
OX0120	*Acidovorax radicis*	1472	MG576017	99.7	Burkholderiales	SX
OX0124	*Acidovorax radicis*	1380	MG576018	99.5	Burkholderiales	SX
OX0611	*Paucibacter toxinivorans*	1441	MG576022	97.9	Burkholderiales	SX
OX1216	*Paraburkholderia fungorum*	1404	MG576015	99.9	Burkholderiales	AEX
OX1313	*Paraburkholderia fungorum*	1489	MG576012	99.9	Burkholderiales	AEX
OX1403	*Paraburkholderia fungorum*	1492	MG576013	100	Burkholderiales	AEX
OX1702	*Paraburkholderia fungorum*	959	MG576014	100	Burkholderiales	AEX
OX0630	*Rhodoferax fermentans*	1360	MG576019	98.5	Burkholderiales	AEX
OX0321	*Rhodoferax saidenbachensis*	1440	MG576020	99.3	Burkholderiales	AEX
OX0130	*Variovorax*	1465	MG576021	99.4	Burkholderiales	AEX

ginsengisoli

OX0627	*Dechloromonas hortensis*	1501	MG576023	98.8	Rhodocyclales	SX

Gammaproteobacteria

OX0110	*Aeromonas popoffii*	1122	MG576027	99.8	Aeromonadales	SX
OX0112	*Aeromonas popoffii*	1506	MG576028	99.7	Aeromonadales	SX
OX0612	*Aeromonas popoffii*	1451	MG576029	99.7	Aeromonadales	SX
OX0622	*Aeromonas hydrophila*	960	MG576025	100	Aeromonadales	SX
SX0303	*Aeromonas piscicola*	876	MG576026	100	Aeromonadales	SX
OX0617	*Aeromonas cavernicola*	721	MG576024	100	Aeromonadales	SX
OX0631	*Pseudomonas pictorum*	1517	MG576051	99.2	Pseudomonadales	SX
OX1807	*Pseudomonas aestusnigri*	671	MG576041	99.4	Pseudomonadales	AEX
OX1808	*Pseudomonas anguillisepticum*	1536	MG576042	97.9	Pseudomonadales	AEX
OX0306	*Pseudomonas extremaustralis*	1429	MG576043	99.7	Pseudomonadales	AEX
OX0322	*Pseudomonas extremaustralis*	1342	MG576044	99.7	Pseudomonadales	AEX
OX1110	*Pseudomonas guineae*	1502	MG576045	99.8	Pseudomonadales	AEX
OX0613	*Pseudomonas linyingensis*	1450	MG576046	99.9	Pseudomonadales	AEX
SX0305	*Pseudomonas mandelii*	1416	MG576047	99.9	Pseudomonadales	SX
SX0307	*Pseudomonas mandelii*	1503	MG576048	99.9	Pseudomonadales	SX
SX1216	*Pseudomonas mandelii*	1503	MG576049	99.9	Pseudomonadales	SX
SX0304	*Pseudomonas peli*	702	MG576050	99.5	Pseudomonadales	SX
SX1213	*Pseudomonas vancouverensis*	1500	MG576052	99.7	Pseudomonadales	SX
OX0304	*Pseudomonas veronii*	702	MG576053	99.6	Pseudomonadales	SX
OX0601	*Acinetobacter pakistanensis*	1501	MG576039	100	Pseudomonadales	SX
OX0621	*Acinetobacter pakistanensis*	839	MG576040	100	Pseudomonadales	SX
OX0619	*Shewanella putrefaciens*	1511	MG576030	99.5	Alteromonadales	SX
OX0103	*Rahnella aquatilis*	576	MG576034	**94.4**	Enterobacteriales	SX
OX0101	*Rahnella inusitata*	1507	MG576035	99.7	Enterobacteriales	SX
OX0604	*Shigella flexneri*	1505	MG576036	99.7	Enterobacteriales	SX

OX0606	*Shigella flexneri*	1510	MG576037	99.7	Enterobacteriales	SX
OX0607	*Escherichia coli*	858	MG576038	99.5	Enterobacteriales	SX
OX1012	*Escerichia fergusonii*	1504	MG576033	99.7	Enterobacteriales	AEX
OX1208	*Rheinheimera soli*	1515	MG576032	99.0	Chromatiales	AEX
OX1909	*Rheinheimera aestuari*	977	MG576031	99.1	Chromatiales	AEX
Bacilli						
OX0301	*Paenibacillus xylanexedens*	1214	MG575992	98.8	Bacillales	SX
OX0302	*Paenibacillus xylanexedens*	1510	MG575993	99.7	Bacillales	SX
OX2310	*Paenibacillus tundrae*	1497	MG575991	99.3	Bacillales	AEX
OX0307	*Paenibacillus terrae*	1502	MG575990	98.9	Bacillales	SX
OX2205	*Paenibacillus alba*	961	MG575989	99.7	Bacillales	AEX
OX0310	*Jeotgalibacillus campisalis*	1484	MG575994	99.4	Bacillales	SX
OX0626	*Staphylococcus argentus*	1409	MG575995	99.9	Bacillales	SX
OX0317	*Psychrobacillus psychrodurans*	1496	MG575988	99.5	Bacillales	SX
OX1213	*Bacillus aquimaris*	1507	MG575973	99.5	Bacillales	AEX
OX2513	*Bacillus halmapalus*	872	MG575974	98.8	Bacillales	AEX
OX1505	*Bacillus hwajinpoensis*	1262	MG575975	99.0	Bacillales	AEX
OX2509	*Bacillus hwajinpoensis*	1282	MG575976	99.4	Bacillales	AEX
OX1011	*Bacillus oceanisediminis*	937	MG575977	100	Bacillales	AEX
OX1604	*Bacillus oceanisediminis*	363	MG575978	99.2	Bacillales	AEX
OX1805	*Bacillus safensis*	885	MG575979	99.6	Bacillales	AEX
OX1004	*Bacillus vietnamensis*	1434	MG575980	99.9	Bacillales	AEX
OX1006	*Bacillus vietnamensis*	1351	MG575981	99.9	Bacillales	AEX
OX1007	*Bacillus vietnamensis*	1508	MG575982	99.9	Bacillales	AEX
OX1210	*Bacillus vietnamensis*	1394	MG575983	99.8	Bacillales	AEX
OX1212	*Bacillus vietnamensis*	1509	MG575984	99.9	Bacillales	AEX
OX2308	*Bacillus vietnamensis*	1205	MG575985	99.8	Bacillales	AEX

OX2309	*Bacillus vietnamensis*	1059	MG575986	99.7	Bacillales	AEX
OX2506	*Bacillus vietnamensis*	1374	MG575987	99.9	Bacillales	AEX
OX2515	*Exiguobacterium oxidotolerans*	861	MG575998	99.8	Lactobacillales	AEX
OX2106	*Exiguobacterium profundum*	754	MG575999	99.9	Lactobacillales	AEX
OX2313	*Exiguobacterium profundum*	1053	MG576000	99.4	Lactobacillales	AEX
OX1107	*Exiguobacterium profundum*	809	MG575996	99.5	Lactobacillales	AEX
OX1109	*Exiguobacterium profundum*	1463	MG575997	98.2	Lactobacillales	AEX

Flavobacteria

OX0623	*Flavobacterium glaciei*	992	MG575964	99.0	Flavobacteriales	SX
OX0125	*Flavobacterium granuli*	1446	MG575965	97.8	Flavobacteriales	SX
OX0129	*Flavobacterium granuli*	1449	MG575966	97.8	Flavobacteriales	SX
OX0311	*Flavobacterium hydatis*	1506	MG575967	97.6	Flavobacteriales	SX
OX0314	*Flavobacterium succinians*	1496	MG575970	98.4	Flavobacteriales	SX
OX0122	*Flavobacterium succinians*	1451	MG575968	98.5	Flavobacteriales	SX
OX0123	*Flavobacterium succinians*	1402	MG575969	98.4	Flavobacteriales	SX
OX0127	*Flavobacterium terrigena*	1447	MG575971	98.6	Flavobacteriales	SX

Cytophagia

OX0104	*Algoriphagus alkaliphilus*	1481	MG575959	97.9	Cytophagales	SX
OX0118	*Algoriphagus alkaliphilus*	1471	MG575960	97.5	Cytophagales	SX
OX0108	*Aquiflexum balticum*	1060	MG575961	**94.7**	Cytophagales	SX
OX0126	*Aquiflexum balticum*	1459	MG575962	**95.5**	Cytophagales	SX
OX0312	*Aquiflexum balticum*	1459	MG575963	**95.5**	Cytophagales	SX

Sphingobacteria

OX0625	*Pedobacter ruber*	1452	MG575972	98.5	Sphingobacteriales	SX

Actinobacteria

OX0615	*Oerskovia*	1353	MG575950	98.4	Micrococcales	SX

	paurometabola					
OX0315	*Cryobacterium arcticum*	1463	MG575951	99.3	Micrococcales	SX
OX0319	*Cryobacterium arcticum*	932	MG575952	100	Micrococcales	SX
OX0117	*Arthrobacter alpinus*	1471	MG575953	99.0	Micrococcales	SX
OX0308	*Arthrobacter humicola*	1460	MG575954	99.9	Micrococcales	SX
OX0313	*Arthrobacter oryzae*	1494	MG575955	99.3	Micrococcales	SX
OX0107	*Pseudarthrobacter siccitolerans*	1487	MG575956	99.0	Micrococcales	SX
OX0614	*Sanguibacter suarezii*	1243	MG575957	99.1	Micrococcales	SX
OX0316	*Streptomyces clavifer*	1156	MG575958	99.6	Streptomycetales	SX
SX1205	*Rhodococcus globerulus*	1294	MG575949	99.7	Corynebacteriales	SX
SX0206	*Rhodococcus globerulus*	1482	MG575948	99.7	Corynebacteriales	SX

[1] *Site SX is at Skógaeyralón (66.15°N, 16.62°W). Vegetated, wetland area. Some suspected gas seepage, but little or no evidence of geothermal influence. Site AER is at Skógakíll near Ærlækjarsel (66.17°N, 16.57°W). Mostly barren, sandy, estuarine area characterized by geothermal activity.*