1

Bleeding, cramping, and satisfaction among new copper IUD users: A prospective study

7 Jessica N Sanders[1][¶], Daniel E Adkins[2,3,¶,*], Simranvir Kaur[4], Kathryn Storck[1], Lori M Gawron[1],

8 David K Turok[1]

11 [1] University of Utah, Department of Obstetrics and Gynecology, Salt Lake City, Utah, United
12 States of America
13
14 [2] University of Utah, Department of Sociology, Salt Lake City, Utah, United States of America
15
16 [3] University of Utah, Department of Psychiatry, Salt Lake City, Utah, United States of
17 America
18
19 [4] University of Utah, School of Medicine, Salt Lake City Utah, United States of America
20

23 * Corresponding author

24 E-mail: daniel.adkins@soc.utah.edu

26 [¶] These authors contributed equally to this work.

Abstract

Objective: We assess change in bleeding, cramping, and satisfaction among new copper (Cu) IUD users during the first six months of use; and evaluate the impact of bleeding and cramping on method satisfaction.

Methods: We recruited 77 women ages 18-45 with a history of regular menstrual cycles for this prospective longitudinal observational cohort study. Eligible women reported regular menses, had no exposure to hormonal contraception in the last three months, and desired a Cu IUD for contraception. We collected data prospectively for 180 days following IUD insertion. Participants reported bleeding scores using the validated pictorial blood loss assessment chart (PBAC), IUD satisfaction using a monthly five-point Likert scale, and cramping using a seven-level ordinal scale. We used multiple imputation to address nonrandom attrition. Structural equation modeling growth curves for count and ordered outcomes modeled bleeding, cramping, and satisfaction trajectories over time. Multiple imputation analyses corrected for moderate attrition bias.

Results: Bleeding significantly decreased (approximately 25%) over the course of the study from an estimated PBAC=195 at one month post IUD insertion to PBAC=151 at six months (t=-2.38; p<0.05). Additionally, IUD satisfaction improved overtime (t=2.65; p<0.01), increasing from between "Neutral" and "Satisfied" to "Satisfied", over the six month study. Cramping decreased sharply over the six-month study from between bimonthly and weekly, to once or twice a month (t=-4.38; p<0.001). Finally, bleeding, but not cramping, was associated with IUD satisfaction (mean IUD satisfaction across study: t=-2.31; p<0.05; IUD satisfaction at study end: t=-2.81; p<0.01).

51 **Conclusions:** New Cu IUD users reported decreasing bleeding and cramping, and increased

52 IUD satisfaction, over the first six months. Method satisfaction was inversely related to

53 bleeding.

Introduction

The intrauterine device (IUD) is the most common reversible method of

contraception worldwide, with its use increasing in both developed and developing

countries [1-3]. On average, 23% of contracepting women (with a range of <2% to >40%,

depending on country) use either the hormonal or non-hormonal (copper) IUD [4]. At

present, the copper (CuT380A) IUD is the only non-hormonal long-acting reversible

contraceptive device approved by the United States Food and Drug Administration (FDA).

The copper IUD is equally effective to sterilization and is designed for extended use (up to

twelve years) [5]. However some women discontinue use due to undesired side effects. The

most common reasons for discontinuation of a copper IUD during the first-year of use are

symptoms related to pain or cramping and complaints of heavy bleeding [6, 7]. Perceived

changes in vaginal bleeding may impact user preference, acceptability, method satisfaction

and continuation [8].

There is a need for updated research evaluating IUD associated bleeding patterns for

providers to better counsel patients on expectations when initiating a copper IUD. The

primary objective of this study is to evaluate IUD associated bleeding and cramping patterns

over the first six months after copper IUD insertion among individuals with regular menses

who have not been exposed to any hormonal contraception in the past three months. A

secondary objective of this study is to relate bleeding and cramping to IUD satisfaction in

the months following insertion.

Materials and methods

For this prospective cohort study, participants were recruited using convenience

sampling through paper and electronic fliers, social media, and word-of-mouth, as well as by

clinicians seeing patients for contraceptive services. A trained research assistant screened

individuals interested in the copper IUD for contraception over the phone to ensure they

met study criteria. Women, aged 18-45 years, with self-reported normal menses, lasting

between three and seven days, were eligible for participation. Eligible participants needed

to be able to provide bleeding information for three cycles free of any hormonal

contraception prior to enrollment and be willing to track bleeding for an additional 180

days. Individuals who were six or more months postpartum, and not breastfeeding, were

eligible if they had returned to normal menses for three cycles.

Exclusion criteria included use of any hormonal contraceptive methods in the three

months, irregular length of any of the previous three menstrual cycles (less than 21 days, or

longer than 35 days), or irregular duration of vaginal bleeding in the previous three cycles

(less than three days, or longer than seven days) at the time of screening. Other exclusion

criteria included any medical contraindication to the copper T380A IUD per the package

insert [9], desire for pregnancy within six months, and participation in any other clinical

research investigation.

This study aimed to incorporate validated methods to evaluate bleeding profiles with

electronic data capture and popular menstrual tracking mobile applications to examine

changes in bleeding patterns among women initiating the copper IUD [10-12]. The analysis

uses two validated measures: the World Health Organization (WHO) recommendations and

the pictorial blood loss assessment chart (PBAC). The WHO method includes data collection

on bleeding days (defined as days when blood loss requires the use of a menstrual pad or

tampon) and spotting days (defined as bloody vaginal discharge, but when no protection

was needed) [10]. The second tool, the PBAC, is an inexpensive and reliable method of

assessing frequency, duration, and intensity of vaginal bleeding [12]. We calculated

monthly PBAC scores based on the published recommendations: one point for each lightly

stained pad or tampon reported, five points for moderately saturated pads or tampon, and

10 points for a completely saturated tampon or 20 points for completely saturated pad.

Additional points were added for small clots (1pt), large clots (5pt), or flooding (5pt). We

also assessed the frequency of the following hormonal/menstrual symptoms adapted from

the Menstrual Symptoms Questionnaire (MSQ): headache, bloating, breast tenderness,

moodiness/irritability, acne, cramping, weight gain, weight loss, depression, constipation

and, diarrhea [13]. Additionally, participants completed a monthly survey that assessed IUD

use, satisfaction, and potential adverse events. We assessed contraceptive method

satisfaction using a five-point Likert scale (very unsatisfied, unsatisfied, neutral, satisfied,

very satisfied) and cramping frequency was assessed using a seven-level ordinal scale (never

to every day of the month).

We used the Research Electronic Data Capture system (REDcap), a secure online

database for survey creation, data collection and management,[14] to collect survey data

and bleeding diaries. Retrospective data were collected for the 90 days prior to copper IUD

insertion and prospective data was collected for 180 days after insertion using REDCap. We

conducted repeated measures analysis at nine time points: three months before insertion

and monthly for six months after insertion. Participants were encouraged to use electronic

period trackers or daily diaries to track the information that they entered monthly into the

REDCap survey.

We modeled longitudinal growth curve trajectories of bleeding scores, IUD

satisfaction, and cramping (as reported by the MSQ) using structural equation models (SEM)

for count and ordinal outcomes.[15-17] This modeling strategy permits the specification of

fixed and random intercepts and slopes allowing us to formally test whether bleeding

decreases, on average, across the study course (fixed time effect), and whether the rate of

change varies across individuals (random time effect).

As none of the three outcomes are continuous variables (instead, they are a count

variable [i.e., PBAC score assessing bleeding] and two ordinal variables [i.e., IUD satisfaction

Likert scale; cramping MSQ item]), we use models for discrete outcomes. Specifically, we

use count link functions for the bleeding growth curve, and ordinal link function for the

satisfaction and cramping growth curves [17, 18]. Given marked overdispersion in the

bleeding measure (mean=148.4; variance=21242.2), we present negative binomial results in

the main text and verify the robustness using Poisson and standard linear link functions in

the Supporting Information. For the ordinal measures, IUD satisfaction and cramping, we

present the conventional ordered logistic link function results in the main text and verify the

robustness of the specification using ordered probit and linear link functions.

We anticipated a missing data pattern in which women with less favorable outcomes

are more likely to attrite. To avoid biased estimates overstating bleeding and cramping

improvements over the study period we used a well-validated multiple imputation (MI)

method (i.e., multiple imputation with chained equations [MICE}) [19]. We used a

conservative 30 imputations in all MI analyses, which were conducted using the "mi

estimate" command in Stata. Finally, we output subject-level estimates for post-insertion

bleeding and post-insertion cramping from the growth curve models, as well as

retrospectively reported baseline bleeding, and tested their association to IUD method

satisfaction (specified as both mean satisfaction across the study, and satisfaction at study

conclusion).

We informed the power calculations from two prior studies [11, 20]. Belsey found

that women on non–hormonal methods had a mean length of bleeding/spotting lasting five

days out of 30 [21] and Tolley et al. found that each additional day of bleeding increased the

likelihood of discontinuation by about 3% for IUD users [20]. Thus, we estimated that

changes longer than 2 days may be considered a menstrual disruption and could potentially

impact user satisfaction and continuation. Using these assumptions with a standard

deviation of 4 days, a sample size of 64 would allow for 80% power with an alpha of 0.05 to

detect 2-day change in bleeding. Our enrollment goal of 77 accounted for a 20% loss to

follow up. This study protocol was submitted to ClinicalTrials.gov (Identifier NCT02311478).

The University of Utah Institutional Review Board approved this study. Teva Pharmaceutical

Industry, Ltd. funded this investigator initiated study and reviewed the manuscript prior to

submission. All analyses were conducted in Stata 15.1.

Results and discussion

We enrolled 79 women into this study, 78 women had a successful IUD insertion.

One individual immediately withdrew from the study and 5 (8%) were lost to follow-up after

the insertion. Seventy-two women (92%) completed the 6-month follow up: 1 (0.01%)

participant had an expulsion, 8 (11%) elected to have removals, and 63 (88%) participants

were still using their device at 6 months (Supporting Figure S1). The mean age of

participants was 25 years (SD 5.2). Three-quarters of women reported using hormonal

contraception in the past, with a median of nine months (mean 18.7 months; SD 25.3 range

3-120 months) since last use. Most participants reported being single/non-cohabitating

(62%) and had a normal body mass index (18.5 - 24.9kg/m^2) (65%). In the enrollment survey,

half (49%) of women reported 3 to 4 days of bleeding per month and half (51%) reported 5

to 7 days of bleeding per month. Over the course of the study the total accumulated person-

months for key assessments ranged between 406 (88%) for IUD satisfaction and 415 (90%)

for bleeding outcomes. In all MI analyses, there were 462 person months (77 subjects*6

assessments). For additional description of participants see Table 1. The MI distributions for

longitudinal outcomes of bleeding, cramping and IUD satisfaction are depicted in the

dotplots in Figure 1 A-C.

204   **Table 1. Sample summary statistics (N=77)**

Highest level of education	N	%
Completed high school or GED	13	17%
Currently in college	28	36%
Graduated from college	26	34%
Education beyond college	10	13%
Annual household income		
Less than $12,000	24	31%
$12,000-$23,999	15	19%
$24,000-$35,999	16	21%
$36,000-$47,999	5	6%
$48,000-$60,000	9	12%
More than $60,000	8	10%
Insurance status		
No insurance	11	14%
Private insurance	62	82%
Medicaid	1	1%
Don't know	2	3%
Insurance covers birth control		
Yes	29	48%
No	6	10%
Don't know	25	42%
Marital status		
Single, never married	48	62%
Single, living with partner	12	16%
Married	10	13%
Divorced	5	6%
Separated	2	3%
BMI		
Underweight	3	4%
Normal	50	65%
Overweight	14	18%
Obese	10	13%
Ever Used Hormonal BC	57	74%
Ever Used Cu IUD	2	3%
Mean period duration in 3 months prior to IUD insertion		
3 to 4 days	38	49%
5 to 7 days	39	51%
Age (mean and SD)	25.4 ($\bar{x}$)	5.20 (σ)

GED-General Equivalency Diploma; BC-birth control; IUD – intrauterine device

Fig 1. Time series plots summarizing distributions, and predicted mean growth curves, of

(A) Bleeding, (B) IUD Satisfaction, and (C) Cramping over study month. Black diamonds

represent variable means (i.e., Bleeding, IUD satisfaction, and Cramping), by month. Interval

bars represent variable means +/- 0.5 variable SDs, by month. The red lines represent the

model-implied growth curve as predicted by the SEM models presented in Table 2. The

underlying dotplot depicts the distributions of the outcomes, by month. For the first plot,

(A), values are binned, and the Y axis is top coded at ≥400 for the PBAC score to improve

resolution in the middle of the distribution. Abbreviations: PBAC – Pictorial Blood

Assessment Chart; IUD – intrauterine device; MSQ - Menstrual Symptoms Questionnaire

As shown in the second column of Table 2, the fixed time slope coefficient for the

bleeding negative binomial SEM growth curve was significantly negative (b=-0.055, p<0.05),

indicating that bleeding declined over the six-month study period. Comparison of the

complete case analysis (see Supplemental Table 1) to the MI findings of the same outcome

(Supplemental Table 2) indicate that the MI approach captures and corrects a moderate

attrition bias; and contributes robust evidence for significant decrease in bleeding in the six-

month period following IUD insertion. As visualized in Figure 1, Panel A, the significant

random intercept term (p<0.001) indicates significant variation in baseline bleeding

between women in the study, and our growth curve estimates (visualized as a red line)

indicate a ~25% reduction in bleeding across the six-month study period. Cramping also

varied across women at at baseline, and notably decreased from between bimonthly and

weekly to once or twice a month across the six-month study period (b=-1.77; p<0.001;

Figure 1, panel B).

The IUD satisfaction ordered logistic SEM growth curve (Figure 1, panel C) indicated

     Table 2. **Structural equation models** **parameter estimates for growth curves of bleeding**

(PBAC Score), IUD satisfaction (5-point Likert), and Cramping (6-level MSQ item) using

multiple imputations (30 imputations)

Outcome:	Bleeding	IUD Satisfaction	Cramping
Link Function:	Negative Binomial	Ordered logistic	Ordered logistic
Time (study month)	-0.055*	0.154**	-0.280***
	(-2.384)	(2.652)	(-4.378)
Intercept	5.126***		
	(54.695)		
Overdispersion (ln(α))	-0.620***		
	(-8.039)		
Random intercept variance	0.361***	3.458***	4.474***
	(4.210)	(3.839)	(4.203)
Threshold 1		-3.529***	-3.709***
		(-9.504)	(-9.743)
Threshold 2		-2.548***	-2.354***
		(-7.725)	(-6.893)
Threshold 3		-1.087***	1.072***
		(-3.716)	(3.426)
Threshold 4		1.637***	2.242***
		(5.529)	(6.703)
Threshold 5			4.716***
			(9.122)
N_j (Assessments)	462	462	462
N_i (Subjects)	77	77	77

t statistics in parentheses
* p<0.05; ** p<0.01; *** p<0.001
PBAC – Pictorial Blood Assessment Chart; IUD – intrauterine
device; MSQ - Menstrual Symptoms Questionnaire

a significant increase in IUD satisfaction in the six months following placement (b=0.154;

p<0.01). As depicted in Figure 1 Panel B, the model-implied IUD satisfaction growth curve

increases about one third of a Likert category from between "Neutral" and "Satisfied" to

"Satisfied", on average, across the six months following IUD placement. Comparing IUD

satisfaction complete case results (Supplemental Table 3) to MI results (Supplemental Table

4), and cramping complete case results (Supplemental Table 5) to MI results (Supplemental

Table 6), indicates that the for both outcomes, the MI approach captures and corrects a

moderate "unhappy user" attrition bias, but that the fixed time coefficients remain

significant in MI analyses, indicating significant improvement in satisfaction and cramping in

the six following Cu IUD placement. In all growth curve models, repeated assessment error

variances were constrained equal, and random time variances were fixed, based on nested

model likelihood ratio tests [15, 22].

In our final analysis, post-insertion bleeding was significantly negatively associated

with IUD satisfaction across the study (b = -1.052, p<0.05), and at study conclusion (b= -1.77,

p<0.01), controlling for cramping and retrosepctive-report baseline bleeding (Table 3,

Columns 2 and 3, respectively). Cramping was not significantly associated with IUD

satisfaction in either specification, controlling for retrospective baseline bleeding and

prospective post-placement bleeding. The sensitivity analysis shown in Supplemental Table

6 show demonstrate this finding was robust across all tested (i.e., ordered probit and linear)

link functions (p<0.001). Retrospective bleeding, reported by participants at the time of

enrollment, showed mixed evidence of a negative, independent effect on IUD satisfaction

(Table 3: average satisfaction: p=ns; satisfaction at study end: p<0.05).

Main Findings

New copper IUD users experience decreasing bleeding and cramping and increasing

IUD satisfactionacross the first six months of Cu IUD use. Increases in overall method

satisfaction across the study is significantly associated with post-insertion bleeding

decreases (Table 3, Model 2: b=-1.77, p<0.01).

Table 3: IUD Satisfaction predicted by bleeding during study, cramping, and retrospective

report baseline bleeding; using multiple imputations (30 imputations)

Outcome:	IUD Satisfaction (study mean)	IUD Satisfaction (study end)
Link Function:	Linear	Ordered logistic
IUD-induced bleeding	-1.052*	-1.766**
	(-2.306)	(-2.812)
Cramping	-0.033	-0.084
	(-0.334)	(-0.752)
Baseline bleeding	-0.002	-0.003*
	(-1.310)	(-1.963)
Intercept	0.235	
	(0.913)	
Threshold 1		-5.003***
		(-4.770)
Threshold 2		-3.093***
		(-6.047)
Threshold 3		-1.269***
		(-3.741)
Threshold 4		0.200
		(0.648)
N	77	77
R-sq	0.092	
Pseudo R-sq		0.066

t statistics in parentheses
* p<0.05; ** p<0.01; *** p<0.001
 IUD – intrauterine device

Strengths

This study has a number of strengths including an adequately powered, prospective

assessment, and focused analysis with high (>90%) retention rates. Additionally, the use of

validated bleeding measures in combination with electronic period trackers and electronic

data reporting potentially strengthen reporting reliability. Statistical methods employed in

the analysis also provided a robust assessment of the impact of bleeding patterns on

satisfaction. The use of SEM allowed us to avoid pooling of the means, and parse out the

impact of baseline bleeding and IUD induced bleeding. The use multiple imputation allowed

us to address potential attrition bias due to nonrandom missing data among women who

discontinued their device due to dissatisfaction, bleeding, or cramping.

Limitations

A main limitation was the lack of prospective pre-insertion bleeding data. We

attempted to collect retrospective bleeding data leading up to IUD insertion, but felt that

there were too many limitations to use this data for the primary analysis. Therefore, this

manuscript focuses the findings on the prospective reports of decreasing bleeding and

cramping and increasing satisfaction. Future research that incorporates prospective

collection of pre-insertion bleeding using similar analytic approaches would enhance the

findings presented here.

Interpretations

289   Similar to other contraception studies, women report high levels of IUD method

290 satisfaction over time [23]. New copper IUD users experience decreasing bleeding and

291 cramping and increasing IUD satisfaction over the first six months of Cu IUD use. Overall

292 method satisfaction across the study is associated with prospective post-insertion bleeding.

Conclusions

Patient-centered contraceptive counseling helps individuals select methods that are

best for them. These patient-driven decisions are associated with higher patient satisfaction

[24]. Conversely, patients who felt their method selection was influenced by perceived

provider bias towards or against a specific method reported lower satisfaction [25].

Unbiased provider counseling may help women anticipate the likelihood, duration, and

intensity of side effects, provide resources for treatment plans to ameliorate side effects,

and support patient's reproductive goals. Because the most common reason for early IUD

discontinuation is related to the side effects, specifically bleeding and cramping, it is

important for providers to have accurate information about those side effects and the

changes over time. These findings support the practical recommendation that providers give

anticipatory patient counseling about their diminishing bleeding and cramping patterns.

Additionally, researchers and practitioners should continue to seek practices that minimize

irregular bleeding post IUD insertion as higher levels of bleeding is associated with lower

satisfaction with the Cu IUD [26-28]. One option is use of a low-cost, 5-day course of over-

the-counter non-steroidal anti-inflammatory drugs (NSAIDs) to decrease bleeding and

cramping with menses during the first three months when most women who experience

these side effects report them at the highest levels [29, 30]. Although such interventions

may not improve continuation rates [29], they may accelerate the return to baseline

bleeding and cramping [28]. In the absence of highly effective treatment for increased

bleeding in the first several months post copper IUD insertion, medical providers should be

providing women with objective information on the likely changes in bleeding they will

experience. To that end, the findings from this study are helpful and reassuring.

317
318

Acknowledgements

The authors would like to extend our deepest gratitude to Amy Orr, our exceptional

research nurse and study coordinator. Additional thanks to Cindy Berg and the Consortium

for Families and Health Research (CFAHR) and University of Utah. The use of REDCap for

survey and data management was provided by Eunice Kennedy Shriver National Institute of

Child Health and Development grant (8UL1TR000105 (formerly UL1RR025764) NCATS/NIH).

References

1.   d'Arcangues C. Worldwide use of intrauterine devices for contraception.
Contraception. 2007;75(6 Suppl):S2-7. Epub 2007/05/29. doi:
10.1016/j.contraception.2006.12.024. PubMed PMID: 17531612.
2.   Finer LB, Jerman J, Kavanaugh ML. Changes in use of long-acting contraceptive
methods in the United States, 2007-2009. Fertility and sterility. 2012;98(4):893-7. Epub
2012/07/17. doi: 10.1016/j.fertnstert.2012.06.027
S0015-0282(12)00680-2 [pii]. PubMed PMID: 22795639; PubMed Central PMCID:
PMC3462302.
3.   Buhling KJ, Zite NB, Lotke P, Black K, Group IW. Worldwide use of intrauterine
contraception: a review. Contraception. 2014;89(3):162-73. doi:
10.1016/j.contraception.2013.11.011. PubMed PMID: 24369300.
4.   Intrauterine contraception: Devices, candidates, and selection [Internet]. 2017 [cited
3/22/2017].
5.   Long-term reversible contraception. Twelve years of experience with the TCu380A
and TCu220C. Contraception. 1997;56(6):341-52. Epub 1998/03/12. PubMed PMID:
9494767.
6.   Hubacher D, Chen P-L, Park S. Side effects from the copper IUD: do they decrease
over time? Contraception. 2009;79(5):356-62. doi: 10.1016/j.contraception.2008.11.012.
PubMed PMID: PMC2702765.
7.   Grunloh DS, Casner T, Secura GM, Peipert JF, Madden T. Characteristics associated
with discontinuation of long-acting reversible contraception within the first 6 months of use.
Obstetrics and gynecology. 2013;122(6):1214-21. Epub 2013/11/10. doi:
10.1097/01.AOG.0000435452.86108.59. PubMed PMID: 24201685; PubMed Central PMCID:
PMCPMC4051392.
8.   Diedrich JT, Desai S, Zhao Q, Secura G, Madden T, Peipert JF. Association of short-
term bleeding and cramping patterns with long-acting reversible contraceptive method
satisfaction. American journal of obstetrics and gynecology. 2015;212(1):50 e1-8. doi:
10.1016/j.ajog.2014.07.025. PubMed PMID: 25046805; PubMed Central PMCID:
PMCPMC4275360.
9.   Duramed Pharmaceuticals. Prescribing Information: ParaGard T380A, Intrauterine
Copper Contraceptive. Pomona, NY: Duramed Pharmaceuticals; 2006.
10.   Belsey EM, Machin D, d'Arcangues C. The analysis of vaginal bleeding patterns
induced by fertility regulating methods. World Health Organization Special Programme of
Research, Development and Research Training in Human Reproduction. Contraception.
1986;34(3):253-60. Epub 1986/09/01. PubMed PMID: 3539509.
11.   Belsey EM. Vaginal bleeding patterns among women using one natural and eight
hormonal methods of contraception. Contraception. 1988;38(2):181-206. Epub 1988/08/01.
PubMed PMID: 2971505.

12.   Higham JM, O'Brien PM, Shaw RW. Assessment of menstrual blood loss using a
pictorial chart. British journal of obstetrics and gynaecology. 1990;97(8):734-9. Epub
1990/08/01. PubMed PMID: 2400752.
13.   Negriff S, Dorn LD, Hillman JB, Huang B. The measurement of menstrual symptoms:
factor structure of the menstrual symptom questionnaire in adolescent girls. J Health
Psychol. 2009;14(7):899-908. doi: 10.1177/1359105309340995. PubMed PMID: 19786516;
PubMed Central PMCID: PMCPMC4301608.
14.   Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic
data capture (REDCap)--a metadata-driven methodology and workflow process for providing
translational research informatics support. J Biomed Inform. 2009;42(2):377-81. doi:
10.1016/j.jbi.2008.08.010. PubMed PMID: 18929686; PubMed Central PMCID:
PMCPMC2700030.
15.   Bollen KA, Curran PJ. Latent curve models : a structural equation perspective.
Hoboken, N.J.: Wiley-Interscience; 2006.
16.   Bollen K, Kirby J, Curran P, Paxton P, Chen F. Latent Variable Models under
Misspecification: Two Stage Least Squares ({2SLS}) and Maximum Likelihood ({ML})
Estimators. 2005.
17.   Masyn KE, Petras H, Liu W. Growth curve models with categorical outcomes.
Encyclopedia of criminology and criminal justice: Springer; 2014. p. 2013-25.
18.   Mehta PD, Neale MC, Flay BR. Squeezing interval change from ordinal panel data:
latent growth curves with ordinal outcomes. Psychol Methods. 2004;9(3):301-33. Epub
2004/09/10. doi: 10.1037/1082-989x.9.3.301. PubMed PMID: 15355151.
19.   White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues
and guidance for practice. Statistics in medicine. 2011;30(4):377-99.
20.   Tolley E, Loza S, Kafafi L, Cummings S. The impact of menstrual side effects on
contraceptive discontinuation: findings from a longitudinal study in Cairo, Egypt.
International family planning perspectives. 2005;31(1):15-23. Epub 2005/05/13. doi:
10.1363/ifpp.31.15.05. PubMed PMID: 15888405.
21.   Belsey EM. The association between vaginal bleeding patterns and reasons for
discontinuation of contraceptive use. Contraception. 1988;38(2):207-25. Epub 1988/08/01.
PubMed PMID: 2971506.
22.   Bollen KA. Structural Equations with Latent Variables. New York: John Wiley and
Sons; 1989.
23.   Peipert JF, Zhao Q, Allsworth JE, Petrosky E, Madden T, Eisenberg D, et al.
Continuation and satisfaction of reversible contraception. Obstetrics and gynecology.
2011;117(5):1105-13. Epub 2011/04/22. doi: 10.1097/AOG.0b013e31821188ad. PubMed
PMID: 21508749; PubMed Central PMCID: PMC3548669.
24.   Dehlendorf C, Grumbach K, Schmittdiel JA, Steinauer J. Shared decision making in
contraceptive counseling. Contraception. 2017. doi: 10.1016/j.contraception.2016.12.010.
PubMed PMID: 28069491.
25.   Higgins JA, Kramer RD, Ryder KM. Provider Bias in Long-Acting Reversible
Contraception (LARC) Promotion and Removal: Perceptions of Young Adult Women.
American journal of public health. 2016;106(11):1932-7. doi: 10.2105/AJPH.2016.303393.
PubMed PMID: 27631741; PubMed Central PMCID: PMCPMC5055778.
26.   Hatcher R, Trussell J, Nelson A, Cates W, Stewart F, Kowal D. Contraceptive
Technology. 19th ed: Ardent Media, Inc; 2011.

27.   Roy S, Shaw ST, Jr. Role of prostaglandins in IUD-associated uterine bleeding--effect
of a prostaglandin synthetase inhibitor (ibuprofen). Obstetrics and gynecology.
1981;58(1):101-6. PubMed PMID: 7243136.
28.   Makarainen L, Ylikorkala O. Ibuprofen prevents IUCD-induced increases in menstrual
blood loss. British journal of obstetrics and gynaecology. 1986;93(3):285-8. PubMed PMID:
3516203.
29.   Hubacher D, Reyes V, Lillo S, Pierre-Louis B, Zepeda A, Chen PL, et al. Preventing
copper intrauterine device removals due to side effects among first-time users: randomized
trial to study the effect of prophylactic ibuprofen. Human reproduction (Oxford, England).
2006;21(6):1467-72. Epub 2006/02/18. doi: 10.1093/humrep/del029. PubMed PMID:
16484309.
30.   Ylikorkala O. Prostaglandin synthesis inhibitors in menorrhagia, intrauterine
contraceptive device-induced side effects and endometriosis. Pharmacol Toxicol. 1994;75
Suppl 2:86-8. PubMed PMID: 7816792.
425
426

Supporting Information

Supplemental Table 1. Structural equation model parameter estimates for growth curves of bleeding (PBAC Score); Unimputed, complete case analysis

Supplemental Table 2. Structural equation model parameter estimates for growth curves of bleeding (PBAC Score); Multiple imputation, 30 imputations

Supplemental Table 3. Structural equation model parameter estimates for growth curves of IUD Satisfaction (5-level Likert scale); Unimputed, complete case analysis

Supplemental Table 4. Structural equation model parameter estimates for growth curves of IUD Satisfaction (5-level ordinal Likert scale); Multiple imputation, 30 imputations

Supplemental Table 5. Structural equation model parameter estimates for growth curves of cramping (6-level ordinal MSQ item); Unimputed, complete case analysis

Supplemental Table 6. Structural equation model parameter estimates for growth curves of cramping (6-level ordinal MSQ item); Multiple imputation, 30 imputations

Supplemental Table 7. IUD Satisfaction predicted by bleeding during study and cramping; Multiple imputation, 30 imputations