Endothelial Protein C Receptor Could Contribute to Experimental Malaria-
Associated Acute Respiratory Distress Syndrome

Luana dos Santos Ortolan[1,4], Michelle Klein Sercundes[2], Gabriel Candido Moura[2],
Thatyane de Castro Quirino[2], Daniela Debone[2], Douglas de Sousa Costa[3], Oscar
Murillo[3], Claudio Romero Farias Marinho[3], Sabrina Epiphanio[1,2]

[1] Department of Immunology, Biomedical Science Institute, University of São Paulo
(USP), 1730 Professor Lineu Prestes avenue, 05508-900, São Paulo, Brazil.

[2] Department of Clinical and Toxicological Analyses, Faculty of Pharmaceutical
Science, University of São Paulo (USP), 580 Professor Lineu Prestes avenue, 05508-
000, São Paulo, Brazil São Paulo, Brazil.

[3] Department of Parasitology, Biomedical Science Institute, University of São Paulo
(USP), 1374 Professor Lineu Prestes avenue, 05508-000, São Paulo, Brazil São Paulo,
Brazil.

[4] Present address: Department of Dermatology, Carver College of Medicine, University
of Iowa, 500 Newton Road, 2110 Medical Laboratories, 52242, Iowa City, IA, United
States.

Correspondence should be addressed to Sabrina Epiphanio; sabrinae@usp.br

Abstract

The severity of *Plasmodium falciparum* malaria is associated with parasite
cytoadherence, but there is limited knowledge about the effect of parasite cytoadherence
in malaria-associated acute respiratory distress syndrome (ARDS). Our objective was to
evaluate the cytoadherence of infected red blood cells (iRBCs) in a murine model of
ARDS and to appraise a potential function of endothelial protein C receptor (EPCR) in
ARDS pathogenesis. DBA/2 mice infected with *P. berghei* ANKA were classified as
ARDS- or hyperparasitemia (HP)-developing mice according to respiratory parameters
and parasitemia. Lungs, blood and bronchoalveolar lavage were collected for gene
expression or protein analyses. Primary cultures of microvascular lung endothelial cells
from DBA/2 mice were analyzed for iRBC interactions. Lungs from ARDS-developing
mice showed evidence of iRBC accumulation along with an increase in EPCR and TNF
concentrations. Furthermore, TNF increased iRBC adherence *in vitro*. Dexamethasone-
treated infected mice showed low levels of TNF and EPCR mRNA expression and,
finally, decreased vascular permeability, thus protecting mice from ARDS. In
conclusion, we identified that increased iRBC cytoadherence in the lungs underlies
malaria-associated ARDS in DBA/2-infected mice and that inflammation increased
cytoadherence capacity, suggesting a participation of EPCR and a conceivable target for
drug development.

Keywords: malaria; endothelial cells; cytoadherence; endothelial protein C receptor

INTRODUCTION

Malaria infection by *Plasmodium falciparum* is responsible for the largest number of severe and fatal diseases in the tropics [1,2]. The main complications of *P. falciparum* infection include cerebral malaria, pulmonary complications, acute renal failure, severe anemia, bleeding and placental malaria [3]. An important aspect of the pathogenesis of severe malaria results from the ability of infected red blood cells (iRBCs) to adhere to the microvasculature. This interaction between iRBCs and the endothelium can cause blocking of blood flow and/or a local inflammatory response [3–5]. Furthermore, these adhesions promote the disappearance of asexual forms of the parasite in the peripheral circulation, thus preventing them from being destroyed in the spleen [3,5,6]. Pulmonary complications caused by severe malaria include acute respiratory distress syndrome (ARDS), which has been associated with not only severe malaria but also different diseases [7–9]. Although malaria-associated ARDS often causes a high mortality rate, not so much research has been performed. Murine models have been used to study malaria-associated ARDS [10,11], and DBA/2 mice infected with *P. berghei* ANKA (PbA) develop ARDS and die between the 7th to 12th days post-infection (dpi) with pleural effusion, edema and inflammatory infiltration in the lungs but without signals of cerebral malaria. In contrast, DBA/2 mice that died after the 13th dpi exhibited pale lungs, no pleural effusion and high levels of parasitemia. The cause of death was attributed to hyperparasitemia (HP) and consequent anemia. In addition, we established predictive criteria to distinguish which mice would die from ARDS on the 7th dpi using respiratory and parasitemia data [12]. Using this model, we found that on average 50% of the mice died from ARDS and 50% died from HP. We found that recruitment of neutrophils and vascular endothelial growth factor (VEGF) is essential to the pathogenesis of malaria-associated ARDS [13,14] and that the induction of heme oxygenase-1 (HO-1) has a protective effect against the development of ARDS in mice [15]. Epiphanio *et al* showed sFLT1 (soluble form of VEGF receptor, known to neutralize excess VEGF in circulation) - expressing adenoviruses treated mice protected against ARDS which was correlated with significant decrease in VEGF levels in circulation, even as carbon monoxide administration by inhalation suppresses the onset of this syndrome [13]. In addition, it was demonstrated that hemin (inducer of HO-1) treated mice had also VEGF levels reduced and were protected against ARDS, defending the alveolar capillary barrier in vitro [15]. Although VEGF is important in PbA-infection of DBA/2 mice, this is not a universal effect, since neutralization of VEGF receptor-2 did not decrease ARDS pathology in *P. berghei* NK65-infected C57BL/6 mice, another well-defined and accepted model of ARDS in malaria [16].
Additionally, it has already been demonstrated that PbA-iRBCs adhere to MVECS (microvascular lung endothelial cells from CBA/Ca mice) and that TNF-stimulated cells express more ICAM-1 and VCAM than control cells [17]. In the *P. chabaudi* model, infected mice in the absence of ICAM-1 showed less anemia and weight loss, reduced parasite accumulation in both the spleen and liver and higher peripheral blood parasitemia during acute stage malaria, which presented the possible role of ICAM-1 in adhesion and pathogenesis of *P. chabaudi* [18]. Endothelial receptors have been studied to understand iRBC adherence in the microvasculature of different organs in severe malaria, and some of these receptors are well established such as ICAM-1 in cerebral malaria and CSA in placental malaria [19]. However, both the mechanism and consequences of PbA-iRBC adhesion in DBA/2 lung endothelial cells remain unknown. In 2013, endothelial protein C receptor (EPCR) was shown to be a new receptor for *Plasmodium falciparum* erythrocyte protein 1 (PfEMP1) in *P. falciparum*-induced severe malaria [20], which may result in a new branch of research on severe malaria. As its primary function, EPCR binds with activated protein C (APC) and cleaves protease-

activated receptor 1 (PAR-1) in a specific Ras-related C3 botulinum toxin substrate 1
(RAC1) pathway, which inhibits the activation of nuclear factor-κB and provides barrier
protection [21]. The EPCR facilitates the activation of protein C (PC) by the thrombin-
thrombomodulin complex, promoting cytoprotective effects in vessels and tissue
protection in the brain, lungs, kidneys, and liver [22]. Additionally, APC provides
neuroprotective effects such as anti-inflammatory and anti-apoptotic effects and
protection of the blood-brain barrier, kidneys, and lungs and thus may be directly
relevant to the complications associated with severe malaria [23].
Some authors proposed that in malaria, *P. falciparum*-infected erythrocytes prompt a
decrease in EPCR levels, and, consequently, APC production is disabled, resulting in
enhanced coagulation and inducing proinflammatory factors and endothelial
dysfunction via PAR-1 [24]. Indeed, *in vitro* studies show that purified amino-terminal
cysteine-rich interdomain region (CIDRα1), especially that from domain cassette 8
(DC8), member of the group containing PfEMP1, interferes with protein C binding to
EPCR, resulting in acquired functional PC systemic deficiency [20,23,25]. However, it
has unexpectedly been shown that iRBCs expressing DC13 and the HB3var03 or
IT4var07 variants of PfEMP1 do not bind to the EPCR of brain endothelial cells *in*
vitro. On the other hand, it has been shown that the DC8 variant IT4var19 may bind to
the EPCR, but this interaction was inhibited when human serum or plasma was added to
the assay [25]. Therefore, the disagreement concerning PfEMP1-EPCR interactions
indicates the need for further studies to understand the outcome of severe malaria.
Notwithstanding, there is no evidence to date of *P. berghei* ANKA binding to EPCR,
and the effects of EPCR on ARDS pathogenesis must be elucidated. Our study,
therefore, was developed in a murine model that mimics various types of human ARDS
[12] and in primary culture of microvascular lung endothelial cells from DBA/2 mice
(PMLEC) not only to clarify the adhesion of infected erythrocytes to murine lung
microvascular endothelial cells but also to understand the relevant aspects of EPCR
modulation by the immune response, which can bring important contributions to
understanding malaria-associated ARDS.
119

MATERIALS AND METHODS

Experimental outline

DBA/2 mice were infected with 10^6 infected red blood cells (iRBC) with *Plasmodium*
berghei ANKA (PbA) and classified as ARDS-developing or HP-developing mice
before death according to previously described criteria [12]. Briefly, we used two
groups of infected mice: the survival group (infected control) and the euthanized group,
in which the mice were euthanized on the 7th day postinfection (10–12 mice per group).
By using respiratory patterns (enhanced pause and respiratory frequency) and the degree
of parasitemia as predictive criteria, we established cut-off values using receiver
operating characteristic (ROC) curves for these parameters measured on the 7th dpi
based on data from mice whose cause of death was known (survival group). In the
survival group, for mice showing pleural effusion or red and congested lungs at
necropsy, the cause of death was designated as ARDS. For mice without pleural
effusion that died after 13 days postinfection with pale lungs and high levels of
parasitemia, the cause of death was designated as hyperparasitemia (HP). Afterward, we
retrospectively diagnosed the euthanized mice as suffering from ARDS or HP by
comparing their respiratory patterns and parasitemia measured on the 7th dpi with the

cut-off values from the survival group at the end of each experiment (20th dpi) [14,15].
All lung tissue, blood, and bronchoalveolar lavage (BAL) samples were collected from
mice on the 7th day postinfection (dpi) always after perfusion of the right ventricle with
20 mL of 1× PBS until the lungs remained clear.

Mice, parasites and euthanasia

Male DBA/2 mice between 6-10 weeks old (purchased from the Department of
Parasitology, University of São Paulo, Brazil) were infected with 1×10^6 *P. berghei*
ANKA (clone 1.49L) iRBCs kindly provided by the laboratory of Dr. Maria Mota from
the Institute of Molecular Medicine (IMM) in Portugal. Parasitemia and mortality were
monitored daily. Parasitemia was determined by Giemsa staining and expressed as the
percentage of infected red blood cells. The euthanasia of mice was performed using
ketamine (150 mg/kg)/xylazine (15 mg/kg).

Determination of Respiratory Pattern

150   Respiratory patterns (respiratory frequency [RF], tidal volume [TV] and enhanced pause
151   [Penh]) were monitored on the 7th dpi by an unrestrained whole-body plethysmography
152   chamber (WBP, Buxco Electronics, USA) for 10 minutes (basal level) according to
153   previously described methods [12].

Histopathological analyses and hemozoin count

Lung tissue fragments were fixed with 10% buffered formalin for 24 hours and kept in
70% ethanol until being embedded in paraffin, and 4–5 μm sections were stained with
hematoxylin-eosin (H&E). To determine the hemozoin (Hz)-containing area, lungs from
euthanized mice (ARDS-developing or HP-developing mice according to the predictive
model on the 7th day postinfection [12]) were stained with H&E, and 10 images were
captured from each tissue sample with polarized light (400× magnification) using a
Zeiss color camera (CAM Axio HRc) connected to a Zeiss light microscope (Axio
Imager.M2). The corresponding percentage of Hz in each image was identified using
the program ImageJ, in which the areas containing Hz were distinguished by brightness
adjustment. The area containing Hz was normalized by the total area in each image.

Quantification of gene expression

Quantitative RT-PCR was performed for the relative quantification of gene expression
from the lungs of noninfected and infected mice. RNA extraction was performed
according to the "Animal Cell I" protocol from a RNeasy Mini kit (Qiagen, USA).
cDNA synthesis was performed with a 1 μg RNA sample using the First Strand cDNA
Synthesis kit RT-PCR (Roche, USA) according to the manufacturer's instructions.
Finally, the PCR sample to test gene expression was prepared with SYBR Green PCR
Master Mix (Applied Biosystems, USA), and the relative quantification $2^{(-\Delta\Delta CT)}$ method
was used as described before (51). The qRT-PCR reactions were performed in the ABI
7500 Fast instrument (Applied Biosystems, USA) using the following oligonucleotides:
PbA 18S (forward 5'-agcattaaataaagcgaatacatccttac-3'; reverse 5'-
ggagattggttttgacgtttatgtg-3'); HPRT (forward 5'-aagcttgctggtgaaaagga-3'; reverse 5'-
ttgcgctcatcttaggcttt-3'); ICAM-1 (forward 5'-cgaaggtggttcttctgagc-3'; reverse 5'-
gtctgctgagscccctcttg-3'); VCAM (forward 5'-agtccgttctgaccatggag-3'; reverse 5'-

tgtctggagccaaacacttg-3'); EPCR (forward 5'-gacgaagtttctgccgctac-3'; reverse 5'-
ctggaggatggtgacgtttt-3').

Parasite localization by bioluminescence

DBA/2 mice were infected with luciferase-expressing *P. berghei* parasite (parasite line
354cl4) kindly provided by the laboratory of Dr. Maria Motta from the Institute of
Molecular Medicine (IMM) in Portugal. On the 7^{th} day postinfection, 150 µl of luciferin
(2.25 mg) (VivoGlo ™ Luciferin, In Vivo Grid, Catalog #: P1041, Promega) was
injected intraperitoneally, allowing the emission of parasite luminescence. Then, the
localization of the parasite was analyzed with an IVIS Spectrum system (PerkinElmer).
For this purpose, mice were sedated with isoflurane for pictures (approximately 6
minutes after the injection of luciferin). Later, they were euthanized and perfused in the
right ventricle with 20 ml of $1\times$ PBS. After perfusion, a new image of the mice was
captured, and the organs were collected and placed in sterile Petri dishes to observe the
bioluminescence of each tissue.

Cytokines quantification

194 The quantities of TNF, IL-6, and IL-33 in serum and culture supernatants were
195 determined by Mouse ELISA Ready-SET-Go! ® commercial kits from eBioscience (San
196 Diego, CA, USA) specific for TNF (ref 88-7324-88), IL-6 (ref 88-7064-88) and IL-33
197 (ref 88-7333-88) according to the manufacturer's protocols.

Isolation of the primary microvascular lung endothelial cells

Primary microvascular lung endothelial cells (PMLECs) were obtained from naive
DBA/2 mice as described before (15,52). Briefly, after euthanasia, the body of the
animal was disinfected with iodine/alcohol. Then, the mice had all of their blood
removed by cutting the carotid artery. In a laminar flow chamber, the lung tissue was
cut into fragments of approximately 1 mm^2 and distributed among 6-well polystyrene
plates in low glucose DMEM (Invitrogen)-supplemented culture medium [20% heat
inactivated FBS, 40 µg/ml gentamicin (Invitrogen-15710064) and 1:100
antibiotic/antimycotic (Gibco-25200)] at 37°C and 5% CO2. After 72 hours, the tissue
fragments were removed, and 50% of the medium was replaced. After 7 days of
incubation, the cells were removed with trypsin 0.25% EDTA (Gibco) and replaced in a
75 cm^2 culture flask. The trypsinization procedure was repeated every 5 to 7 days.
Finally, the cells were cultured for 15 to 20 days (3^{rd} and 5^{th} passage) until being used in
the trials. The purity of isolated DBA/2-PMLECs was characterized by
immunofluorescence with lectin from *Ulex europaeus* (Sigma-Aldrich, USA– l9006)
and the anti-VWF (Santa Cruz Biotechnology, USA – sc14014), anti-CD31 (Abcam,
UK – ab28364), anti-ACE (Abcam, UK – ab85955), anti-CD62E (Abcam, UK -
ab18981), anti-eNOS (Abcam, UK – ab87750) and anti-VE-cadherin (Abcam, UK –
ab205336) antibodies. All of the cells were stained with all of the markers, indicating
high purity (Figure S1).

***Plasmodium* synchronization and enrichment of parasitized erythrocytes**

To obtain mature forms of *P. berghei* ANKA, iRBCs were synchronized as described
previously (53). Briefly, iRBCs were collected from infected mice exhibiting 10 to 20%
parasitemia through cardiac puncture and transferred to RPMI 1640 culture medium

(Gibco, Thermo Fisher Scientific, USA) supplemented with 25% fetal bovine serum
(FBS). The iRBCs were subsequently maintained *in vitro* at 37°C for 14 hours in an
atmosphere containing 5% CO_2, 85% N_2, and 10% O_2. The parasitized erythrocytes
were then enriched using a magnetic separation column (Miltenyi Biotec, USA) to
generate cell populations consisting of approximately 95% iRBCs, as assessed by thick
blood smears. Lysate containing *P. berghei* was obtained from iRBCs subjected to
several freeze-thaw cycles.

Peritoneal Macrophages

Peritoneal macrophages (Mφ) were collected from noninfected DBA/2 mice euthanized
with halothane, the skin was then removed and 5 ml of 1× sterile cold PBS (at 4°C) was
inserted into the peritoneum. The liquid was then removed, aspirated with a 24-gauge
needle, and centrifuged at 1000 rpm for 5 minutes at 4°C. The supernatant was
discarded, and the cells were resuspended in supplemented DMEM medium [20% heat-
inactivated FBS, 40 μg/ml gentamicin (Invitrogen-15710064) and 1:100 antibiotic and
antimycotic (Gibco-25200)]. Cells were counted in a Neubauer chamber and arranged in
Transwell culture plates pretreated with 0.2% gelatin in 1× PBS (gelatin from bovine
skin, G9391, Sigma-Aldrich).

Transwell assay with endothelial cells and peritoneal macrophages

Sterile 13 mm diameter coverslips (Knittel) were placed in 24-well culture plates. The
coverslips were treated with 0.2% gelatin (gelatin from bovine skin - Sigma-Aldrich)
diluted in 1× PBS. PMLECs were seeded in the culture plate (7×10^4 cells/well) in 400
μl of complete DMEM medium, and 1×10^5 peritoneal macrophages (Mφ) were seeded
in the Transwell membrane (6.5 mm membrane diameter, 0.4 μm membrane pore,
Corning, Costar 3470) without contacting PMLECs on the bottom of the 24-well plate.
They only shared the supernatant through the Transwell membrane. After 24 hours, 2.5
$\times 10^6$ mature iRBCs or RBCs were added over the Mφ and coincubated for 24 hours.
After this time, the supernatant between the Mφ and endothelial cells was collected for
the quantification of cytokines by ELISA. Subsequently, endothelial cells were
coincubated with 1.75×10^6 PbA-iRBCs for the adhesion assay.

***Plasmodium berghei* ANKA adhesion assay under static conditions**

DBA/2-PMLECs were plated in a Lab-Tek chamber slide system made of Permanox
containing 8 wells (3×10^4 cells/well) (Thermo Fisher Scientific - 177 455). Afterwards,
TNF (50 ng/ml) was added to the cells and incubated for 24, 48 or 72 hours. After this
stimulation, synchronized PbA-iRBCs were added for 1 hour to the culture at a ratio of
25 iRBCs/PMLEC at 37°C and 5% CO_2. After the incubation period, the culture
medium and the removable chamber were removed, and the slides were dipped in
preheated DMEM medium without FBS to eliminate unbound PbA-iRBCs. The slides
were then fixed with methanol, stained with Giemsa and observed under an optical
microscope immersion objective (1000×). The counting pattern used was the number of
adhered PbA-iRBCs for every 100 DBA/2-PMLECs.

Parasite adhesion assay in flowing conditions

PMLECs between the 3rd and 5th passages were seeded (8×10^4/well) in 2-well
Permanox chamber slides from Lab-tek (Thermo Scientific, Nunc). The cells were

incubated at 37°C and 5% CO_2 for 24 hours. After this time, TNF was added for 24
hours or not added (for the control group). The stimulus was removed, and mature PbA-
iRBCs (25 per cell) in DMEM supplemented with 20% FBS were added and allowed to
interact with the endothelial cells for 1 hour at 37°C in 5% CO_2. Subsequently, the wells
were detached from the polystyrene slides, and the slides were added to a flow system
composed of a chamber (cell adhesion flow chamber, Immunetics) that kept the slide
adhered through the formation of a vacuum (coupled by a vacuum pump), a syringe
pump (Insight Inc.), an inverted microscope (Zeiss Vert. A1) connected to a camera
(Axio Cam ERc 5s, Zeiss) and a computer with an image capture system (Zen 2011
program, AxioVision Rel 4.8.2 SP2). The chamber was initially filled with DMEM
medium, and then an initial photo with the parasites adhered to the PMLEC-DBA/2
cells was taken. Then, a medium continuous stream was run through the chamber and
maintained at a flow rate of 2 ml/hour with a pump syringe (Insight Inc.). Five images,
taken every 3 minutes, of each well were captured while maintaining the same field as
the initial image to determine the erythrocyte binding efficiency. Image processing and
analysis were performed using ImageJ (version 1.46 r). The images were opened
individually in ImageJ and analyzed through the "cell counter" plugin, and all PbA-
iRBC images were selected on each image. After counting all the images, the total
number of PbA-iRBCs that remained in the final image (taken after 15 minutes) was
recorded. The experiment was repeated with two independent cultures, each with two
technical replicates. Image processing and analysis were performed in ImageJ.

Treatment with dexamethasone

287 A total of 200 µl (80 mg/kg) of dexamethasone (8 mg/ml Decadronal, Aché, Brazil) was
288 taken directly from the bottle and injected intraperitoneally (ip) on days 5 and 6
289 postinfection. The dexamethasone dose was chosen according to the therapeutic dose
290 observed by Van den Steen in 2010 [29]. Control mice received 200 µl of $1\times$ PBS ip on
291 the same days.

Lung permeability and edema quantification

To investigate lung permeability, noninfected or PbA-infected mice (dexamethasone-
treated or not) on the 7[th] dpi were injected intravenously with 0.2 mL of 1% Evans Blue
(Sigma-Aldrich). The mice were euthanized 45 minutes later, and the lungs were
weighed immediately and placed in 2 mL of formamide (Merck) for 48 hours at 37°C
[12]. The absorbance of the formamide was then measured at 620 nm and 740 nm. The
amount of Evans Blue staining per gram of lung tissue was calculated from a standard
curve.

Measure of primary microvascular lung endothelial cell permeability

The increased lung vascular permeability was analyzed in DBA/2-PMLECs plated on
permeable membrane inserts with 0.4 µM pores (Transwell Corning) pretreated with
gelatin 0.2% in $1\times$ PBS (gelatin from bovine skin, G9391, Sigma-Aldrich), coupled in
24-well polystyrene plates at a concentration of 2.2×10^4 cells per insert and maintained
in DMEM culture at 37°C as previously described (15). After 96 hours, when the cells
reached confluency, PbA lysate was applied for 1 hour after incubation with
dexamethasone (500 ng/ml for 24 h) or solely with 20% FBS-supplemented DMEM
culture medium. The culture medium was subsequently replaced by Hank's balanced
salt solution, and in the upper compartment of each insert in contact with the cells, 200

μl of Evans Blue was incubated at a 2 mg/mL concentration at 37°C. After 30 minutes,
the liquid from the lower compartment was collected and analyzed in a
spectrophotometer at a wavelength of 650 nm (NanoDrop 2000, Thermo Scientific).
Finally, the concentration of Evans Blue was determined from a standard curve (0.2
mg/mL to 0.0031 mg/mL) as previously described [30].

Actin microfilament identification by immunofluorescent and morphometric
analysis of the opening of interendothelial junctions

To analyze the area of opening of interendothelial junctions (OIJ), PMLECs were plated
in 24-well plates (7×10^4 cells/well), adhered to gelatin on glass coverslips, and
maintained at 37°C and 5% CO_2. The cells were coincubated with either iRBCs or
RBCs for 1 hour after incubation with dexamethasone (24 hours) or DMEM culture
medium supplemented with 20% FBS, and incubations were done in triplicate.
Subsequently, the cells were fixed with 3.7% formaldehyde, permeabilized with acetone
at -20°C, and blocked with bovine serum albumin solution (1% BSA). Actin was
marked with Texas Red-phalloidin (T7471, Life Technologies) for 20 minutes. The cell
nuclei were marked with Hoechst stain (H33342, Life Technologies). Each slide with
fully confluent cells was chosen randomly, and ten to twenty pictures were taken and
scanned in a "zig-zag" pattern from top to bottom. The images were acquired with an
Axio Imager M2 microscope (Zeiss) using Axiocam HRc (Zeiss) and Axio Vision
software, version 4.9.1.0. The total OIJ area was measured in each picture using ImageJ
software (version 1.52a - NIH, USA). To calculate the OIJ, the "measure" tool and the
"wand (tracing)" tool in ImageJ were used. Each image was analyzed individually, and
all of the OIJs were circled with the "wand (tracing)" tool, from which the area of each
space was calculated. Analysis of the differences between the openings of the
interendothelial junctions following different stimuli was performed using GraphPad
Prism 8® software.

TNF blockage

TNF was blocked in DBA-PMLECs using mouse TNF neutralizing (D2H4) rabbit mAb
#11969 (Cell Signaling®) at a concentration of 1 µg/ml for 24 hours in PMLECs.

Soluble EPCR quantification

Soluble endothelial protein C receptor (sEPCR) was measured with an ELISA kit
(Elabscience®, E-EL-M1073) according to the manufacturer's instructions.

EPCR Western Blot

Fresh frozen mouse lung tissues collected on the 7th dpi were sonicated and
homogenized at 4°C using Radio-Immunoprecipitation Assay (RIPA) buffer composed
of 50 mM Tris-HCl pH 8.0, 150 mM NaCl, 0.5% sodium deoxycholate, 0.2% sodium
dodecyl sulfate (SDS), 1 mM sodium orthovanadate, 1 mM NaF, and a protease
inhibitor tablet. The total protein concentration was determined using a SpectraMax®
Plus 384 (Molecular Devices) spectrophotometer with a 562 nm filter using the
Pierce™ BCA Protein Assay kit (Thermo Scientific) according to the manufacturer's
instructions. Then, 30 µg of protein from each sample was electrophoretically separated
in a 10% polyacrylamide mini-gel (Bio-Rad) by SDS-PAGE at 100 V for 2 hours in a
Mini-PROTEAN® Tetra cell apparatus (Bio-Rad). The Kaleidoscope commercial

standard Precision Plus Protein Standard (Bio-Rad) was used. Subsequently, proteins
separated on the gel were transferred to PVDF (polyvinylidene difluoride, 0.2 μm, Bio-
Rad) membranes in transfer buffer (0.1 M Tris, 20% methanol and MilliQ water)
overnight at 4°C in a wet transfer apparatus at 70 mA constant voltage (Bio-Rad).
Detection of the chemiluminescence immunochemical reaction was performed on a
ChemiDoc ™ XRS + Molecular Imager® (Bio-Rad) transilluminator with the ImageLab
™ program (Bio-Rad). Blocking of nonspecific sites was carried out after the transfer in
Tris-buffered saline and Tween-20 buffer (0.05 M Tris, 0.1 M NaCl, pH 7.3, and 0.1%
Tween-20, Synth) containing 3% BSA for 2 hours while stirring at room temperature
(RT). The membrane was then incubated with the primary anti-EPCR antibody
(#151403, Abcam) at 1:1,000 while stirring overnight at 4°C. After washing in TBS-T,
the membrane was incubated with the peroxidase-conjugated rabbit anti-rabbit IgG
(HRP) secondary antibody (#AP307P, Millipore) at 1:6,000 in TBS-T for 1 hour (RT).
Bands corresponding to EPCR and the housekeeping protein β-actin (mouse IgG
#NB600-501, Novus, at 1:40,000) were detected by the chemiluminescence method
(Clarity Western ECL Bio-Rad) and finally densitometrically measured by ImageJ 1.6.0
software.

Bronchoalveolar lavage and VEGF quantification

371 On the 7[th] dpi, infected mice (untreated or treated with dexamethasone) were
372 anesthetized, and having bronchoalveolar lavage (BAL) collected. For that, the trachea
373 was exposed and cannulated, and the lungs were washed once with 1.0 ml of 1× PBS.
374 An ELISA kit (R&D Systems, USA) was used to quantify VEGF levels in BAL and in
375 culture supernatant according to the manufacturer's instructions.

Transfection of EPCR by siRNA

To transfect EPCR by interference RNA assay, Lipofectamine® RNAiMAX reagent
(Invitrogen by Life Technologies) and 3 different predesigned and validated oligo
silencers (Ambion by Life Technologies) were used according to the manufacturer's
protocol. Additionally, a Select Negative Control silencer [(C-) (Ambion 4390843)] and
a select GAPDH positive control silencer [(C+) (Ambion 4390849) were employed as
control reactions.
siRNA1: forward: 5' CAACCGGACUCGGUAUGAATT 3';
reverse: 5' UUCAUACCGAGUCCGGUUGta3'
siRNA2: forward: 5' ACGCAAAACAUGAAAGGGATT 3';
reverse: 5' UCCCUUUCAUGUUUUGCGUGG 3'
siRNA3: forward: 5' CGCCCUUUGUAACUCCGAUTT 3';
reverse: 5' AUCGGAGUUACAAAGGGCGCA3'

Immunohistochemistry of ICAM-1 and VCAM in lungs

To perform immunohistochemistry of ICAM-1 and VCAM, slides containing the
paraffin tissue sections were placed in an incubator at 60°C for 20 minutes to melt the
paraffin. Then, they were incubated in xylene twice for 15 min at 60°C and then in
absolute ethanol, 95% alcohol, 70% alcohol, distilled water, and finally 1× PBS, pH 7.2
to 7.4. For antigen retrieval, the slides were incubated in sodium citrate buffer, pH 6, for
45 minutes at 95°C. Endogenous peroxidase was blocked with 3% hydrogen peroxide
for 15 minutes twice at room temperature while protected from light. The tissue was
probed with rabbit polyclonal to ICAM-1 (1:400) antibody (Abcam, ab124759) and

rabbit monoclonal to VCAM (1:600) antibody (Abcam ab134047) overnight at 4°C, and
then the REVEAL mouse/rabbit kit (Spring, Code SPD-015) was used in accordance
with the manufacturer's instructions. The quantification of ICAM-1 and VCAM in lung
tissue was performed by calculating the marked area normalized to the total tissue
section. The calculation was performed in ImageJ (version 1.50b) software using the
IHC toolbox plugin [31].

Statistical Analysis

The statistical analyses were performed using GraphPad Prism$^{®}$ 5.0 software for
analysis and graphing. The data were analyzed for normality by the Kolmogorov-
Smirnov test or the Shapiro-Wilk normality test and for variance with the Bartlett test.
Nonparametric variables for two groups were compared by using the Mann-Whitney
test. For analysis of three groups, we used the Kruskal-Wallis test followed by the
Dunn's post hoc test. To compare parametrical variables for two groups, a t-test was
employed, and for three or more groups, one-way ANOVA followed by the Bonferroni
posttest was used. For the survival curves, log-rank and Gehan-Breslow Wilcoxon tests
were applied. The differences between the groups were considered significant when
$p \leq 0.05$ (5%). To establish a cut-off from the data, ROC curves were generated by using
the results of the control group obtained in MedCalc version 8.2.1.0.

Ethics Statement

All experiments were performed in accordance with the ethical guidelines for
experiments with mice, and the protocols were approved by the Animal Health
Committee of the Biomedical Sciences Institute of the University of São Paulo (CEUA
nº 24, page 16, book 03). The guidelines for animal use and care were based on the
standards established by the National Council for Control of Animal Experimentation
(CONCEA: Conselho Nacional de Controle de Experimentação Animal) and Brazilian
Federal Law nº 11.794.

RESULTS

ARDS-developing mice show higher levels of pulmonary parasite than HP-developing mice

In our previous study, it was shown that a proportion of DBA/2 mice infected with
PbA-iRBC developed singular characteristics of ARDS and died between the 7th and
12th dpi [12–15]. To investigate whether adherence is essential in the evolution of this
phenotype, we observed that luciferase-expressing *P. berghei* parasites were distributed
in the peripheral blood and tissues of DBA/2 mice. However, when they were perfused
with 1× PBS, the bioluminescence (luciferase/luciferin) signal remained concentrated in
the spleen, lungs and slightly in the liver especially in ARDS-developing mice (Figure 1
a-c), corroborating our published data where we showed more bioluminescence signal
in ARDS mice compared to HP [14]. We demonstrated that ARDS-developing mice
showed higher levels of 18S subunit PbA rRNA expression (Figure 1d) and higher
hemozoin concentrations in the lungs compared to those of HP-developing mice on the
7th dpi (Figure 1 e-g). In addition, we analyzed histological lung sections of mice that
died with ARDS and found several iRBCs in close contact with endothelial cells (Figure

1h). These results taken together indicated that ARDS-developing mice accumulated a
considerable amount of iRBCs in the lungs, suggesting their essential participation in
ARDS pathogenesis.

TNF increases *P. berghei* adherence in the ARDS experimental model

ARDS-developing mice on the 7^{th} dpi displayed higher TNF levels in serum compared
to those of HP-developing mice (Figure 2a), suggesting that this inflammatory cytokine
may be critical to ARDS development. We further investigated whether iRBCs could
contribute to TNF release by endothelial cells. First, we demonstrated that primary
microvascular lung endothelial cells (PMLECs) stimulated with iRBCs can directly
contribute to TNF production (Figure 2b). Then, we examined the influence of TNF on
the adhesion of iRBCs to PMLECs using static and flow conditions. TNF-stimulated
cells increased the capacity of iRBCs to adhere to PMLECs after 24 and 72 hours in the
static assay (Figure 2 c-e). In the flow adherence assay, which mimics physiologic
conditions, the iRBCs had more cytoadherence with TNF than without TNF stimulation
(Figure 2 f-h). Additionally, peritoneal macrophages (Mφ), collected from noninfected
DBA/2 mice, were seeded in Transwell membranes, and then PMLECs were seeded on
the bottom of a 24-well plate with no contact with the Mφ being stimulated or not by
red blood cells (RBCs) or iRBCs (Figure 2i-k). Mφ stimulated by iRBCs produced
more TNF than Mφ without contact with iRBCs (Figure 2j). Additionally, PMLECs that
were in indirect contact with Mφ stimulated with iRBCs demonstrated more adherence
to iRBCs than PMLECs with no previous contact with Mφ (Figure 2k). Finally, to
evaluate the contribution of TNF to the adhesion of iRBCs to PMLECs, the cells were
blocked with TNF antibody, and as a result, the blockage reduced the capacity of iRBCs
to adhere to PMLECs (Figure 2l). All these data show that ARDS-developing mice
express more TNF, which contributes to iRBC adherence in endothelial cells than in
HP-developing mice.

EPCR could contribute to *P. berghei* cytoadherence

Looking for adhesion molecules that could be involved in PbA-iRBC cytoadherence in
ARDS pathogenesis, we observed that ARDS-developing mice showed higher levels of
VCAM and ICAM-1 in lungs than HP-developing or noninfected mice (NI) as analyzed
by immunohistochemistry (Figure S2 a-c). Additionally, recombinant TNF upregulated
the mRNA expression of ICAM-1 and VCAM in PMLECs (Figure S2 d-i). However,
the most interesting finding was the upregulation of mRNA EPCR expression in the
lungs of ARDS-developing mice compared to NI mice (28.48-fold increase) and HP-
developing mice (13.16-fold increase) (Figure 3a). There was also an increase in soluble
EPCR (sEPCR) (Figure 3b) in serum and EPCR protein in lungs (Figure 3c) in ARDS-
developing mice when compared to HP-developing mice. We further investigated the
influence of TNF on the regulation of EPCR expression in PMLECs. TNF-stimulated
cells showed an upregulation of EPCR expression at 48 and 72 hours, but this was not
seen with PbA-iRBC stimulation (Figure 3 d-f). Moreover, EPCR knockdown with
siRNA transfection (siEPCR) in PMLECs reduced EPCR expression compared to that
in nontransfected PMLECs (Figure 3g). In addition, in siEPCR cells, iRBC adhesion
was reduced compared to TNF-stimulated and nontransfected cells (Figure 3h).

Dexamethasone reduces VEGF, TNF and EPCR, protecting mouse lungs and PMLECs from increased vascular permeability

Once TNF seemed to be important to induce PbA-iRBC adherence and consequently ARDS development, DBA/2 mice were infected with PbA-iRBC and treated with dexamethasone (80 mg/kg) on days 5 and 6 post-infection to investigate whether it could protect them from lung injury.

Dexamethasone treatment decreased TNF concentrations in the plasma (Figure 4a) apart from IL-6 and IL-33 concentrations in treated mice compared to those of nontreated mice (Figure S3 a and b). Dexamethasone also downregulated EPCR (Figure 4b) and VCAM (Figure S2 c) expression but not ICAM expression (Figure S2 d) in the lungs of infected mice. The soluble EPCR concentrations in the serum and bronchoalveolar lavage (BAL) are shown (Figure 4c and d). The concentration of VEGF, an essential factor to increase vascular permeability and ARDS development, also decreased after dexamethasone treatment in the BAL of infected mice on the 7th dpi (Figure 4e).

Consequently, we analyzed lung vascular permeability, and dexamethasone-treated infected mice were protected compared to nontreated mice (Figure 4f-h). To clarify the mechanism that reduced lung permeability, the openings of interendothelial junctions in PMLECs were evaluated through actin analysis. The PMLECs stimulated with iRBCs increased the spaces between the interendothelial junctions and led to disruption and reorganization in the actin filaments, which were protected by treatment with dexamethasone (Figure 4i and j). iRBC-stimulated PMLECs also demonstrated more permeability for Evans Blue dye in a Transwell assay compared to dexamethasone-treated PMLECs (Figure 4k). Additionally, dexamethasone-treated PMLECs showed a reduction in VEGF concentration compared to cells untreated with dexamethasone (Figure 4l). With everything considered, these data suggest that dexamethasone is capable of reducing inflammation and decreasing EPCR levels and vascular permeability in PMLECs and the lungs of DBA/2 mice.

Anti-inflammatory drug protects mice from ARDS but not from malaria infection

Dexamethasone-treated mice improved their respiratory parameters [respiratory frequency, tidal volume and enhanced pause (Penh)] compared to infected nontreated mice on the 7th dpi (Figure 5 a-c). On the other hand, parasitemia levels increased in the peripheral blood of dexamethasone-treated mice on the 7th dpi (Figure 5d), and these mice succumbed between 8 and 20 dpi (Figure 5e) with characteristics of HP without edema and lower inflammatory infiltration according to the histopathological analyses (Figure 5 f-g) and count of iRBC data (Figure 5d).

Discussion

In the present study, we describe the cytoadherence of PbA-iRBCs in microvascular lung endothelial cells in ARDS-developing mice, as well as potential contribution of the EPCR in this mechanism and, consequently, in ARDS pathogenesis. An important advantage of this study is that our animal model reproduces the physiopathological aspects found in human ARDS patients [12,13,32].

We have previously shown that PbA-infected DBA/2 mice can develop ARDS (between
7 to 12 days after infection) or develop HP later. ARDS-developing mice show, from
7^{th} dpi, interstitial and lung edema, higher vascular permeability, lung opacification,
worse breathing capacity unlike HP- developing mice [12]. ARDS-developing mice also
produce more neutrophils-attracting chemokines, myeloperoxidase and reactive oxygen
species resulting in increase of neutrophils in lungs and BAL [14]. In this work, we
investigated how PbA-iRBC interaction with endothelial cells could mediate some of
those changes in the DBA/2 model, leading to one of the two different phenotypes.
It is well known that mature forms of *P. falciparum*-iRBCs can adhere to the
microvasculature of different organs, which is an important feature in the pathogenesis
of severe malaria in humans [5,33]. However, there are only a few studies focusing on
cytoadherence in ARDS [11,29,34,35], and the mechanism of this is not entirely known.
Previously, we found that ARDS-developing mice showed a higher level of parasites in
their lungs than HP-developing mice, which may be important to induce malaria [14].
Our work reinforces that PbAluc-iRBC accumulated in the lungs of ARDS-developing
mice, which could contribute to pathogenesis. Additionally, ARDS-developing mice
showed a larger hemozoin (Hz) area in the lungs than HP-developing mice on the 7^{th}
dpi, which is consistent with the onset of ARDS development as patients from Thailand
with ARDS development presented a significant amount of pulmonary Hz [36]. Some
previous works suggest that Hz, which is released from the food vacuole into circulation
during erythrocyte lysis, is rapidly taken up by circulating monocytes and tissue
macrophages, inducing the production of pro-inflammatory mediators [37,38]. In
addition, Hz linked by *Plasmodium* DNA is a ligand for TLR 9, which activates innate
immune responses *in vivo* and *in vitro*, resulting in the production of cytokines and
chemokines and the upregulation of costimulatory molecules [39,40] and
proinflammatory mediators such as IL-1β via NLP3 inflammasome complex activation
[38].
It is known that iRBCs adhere to endothelial cells *in vivo* and *in vitro*. Recently, PbA-
iRBCs have been demonstrated to adhere to MVECs (microvascular lung endothelial
cells from CBA/Ca mice) *in vitro*, and TNF-stimulated cells expressed more ICAM-1
and VCAM [17]. Other studies showed that endothelial cells stimulated with TNF
showed an increase in ICAM-1 expression [34,41], corroborating our results *in vivo* and
in vitro that recently demonstrated that TNF and ICAM-1 markers were increased in the
lungs of patients who had developed the syndrome [36]. The inhibition of *P. falciparum*
iRBC adherence in TNF-activated HUVECs incubated with either anti-ELAN-1 mAB
BBII or anti-VCAM mAB 4B9 [42] was not observed, suggesting that other molecules
could contribute to the adhesion of iRBCs. It is interesting to highlight that the parasites
by themselves (PbA-iRBC or PbA lysate) cannot upregulate ICAM-1, VCAM or EPCR
expression in DBA/2-PMLECs. In fact, PbA-iRBCs adhere more to TNF-stimulated
DBA/2-PMLECs than nonstimulated cells. We previously observed the close contact
between PbA-iRBCs and endothelial cell membranes in the lungs of ARDS-developing
mice [32]. Nevertheless, the study reported here suggests the mechanism of PbA-iRBC
adhesion using DBA/2 mice that had developed ARDS. We found that ARDS-
developing mice have approximately 28.5 times more EPCR expression than uninfected
mice; they showed more EPCR protein in their lungs than HP-developing mice, and
TNF-stimulated PMLECs displayed an upregulation in EPCR expression compared to
that of nonstimulated cells (NS) or cells stimulated with iRBCs, indicating that EPCR
can contribute to ARDS progression. Similarly, a study indicated that children in Papua
New Guinea with severe malaria had higher antibody levels against EPCR binding to
CIDRα1 domains compared to uncomplicated malaria and this difference was only

found in older children [43]. However, in a recent study, the expression of EPCR
decreased in the lungs of *P. falciparum*-infected patients who developed ARDS when
compared to those who did not develop ARDS as analyzed by immunohistochemistry.
The authors suggest that the change in EPCR together with thrombomodulin and in
association with the deposition of hemozoin in the lungs plays an important role in the
pathogenesis of ARDS [36]. Additionally, reduced sEPCR serum levels were detected
in children with severe malaria, and these levels were higher in cerebral liquid from
children with cerebral malaria. In spite of this, sEPCR levels were not related to
mortality or neurologic manifestations at discharge or 6-month follow-up [44].
Despite the fact that *P. berghei* ANKA does not express PfEMP1, it can cause severe
malaria. PfEMP-homologies proteins are not displayed from *P. berghei* and exported
proteins from this parasite species, responsible for iRBC cytoadherence, are not
unrevealed yet. On the other hand, *P. berghei*-iRBC have shown close contact with lung
endothelial cells in ARDS-developing mice [32]. Nevertheless, a recent malaria
proteomic and genetic study shows that *P. berghei* blood stages export diversified
collection of proteins [45]. One molecule that have been previously reported to mediate
adhesion of iRBC in endothelial cells is the schizont membrane-associated
cytoadherence protein (SMAC), that is exported into the cytoplasm of the host
erythrocyte. Mutants lacking expression of SMAC show a strongly reduced CD36-
mediated iRBC sequestration [46]. Unfortunately, we have not found any correlation
between CD36 mRNA levels and ARDS development in our model. Passini *et al*, 2013
identified erythrocyte membrane associated proteins, but related in sequestration [45].
Therefore, further studies should be developed to recognize this *P. berghei*-proteins,
especially with potential to connected EPCR adhesion.
With the hypothesis that inflammation is essential to expose adherence molecules and
consequently to ARDS pathogenesis, we treated DBA/2 mice with 80 mg/kg of
dexamethasone. The 80 mg/kg dose is the therapeutic dose for ARDS treatment in mice
observed by Van den Steen in 2010, because lower doses did not protect mice from
ARDS and edema [29]. In contrast, dexamethasone was administered to patients
suffering from ARDS (different etiologies) at lower doses: 5 mg/day for 4 days in
ARDS patients with bacterial pneumonia [47] and 10 mg for 6 hours in ARDS patients
with acute leucocyte leukemia [48]. In two well-conducted studies, dexamethasone
failed to improve the fatality rate of cerebral malaria [49,50]. Unfortunately, the use of
corticosteroids in acute respiratory distress syndrome (ARDS) due to malaria in humans
has not been well explored [51]. In our experiments, we observed that dexamethasone-
treated mice were protected from ARDS and showed the downregulation in EPCR
expression, which indicates that EPCR may be necessary for iRBC adherence and the
progression of disease. In other models, using endothelial cells (HUVECs),
dexamethasone also had a direct effect on the down-regulation of the EPCR expression
[52].
EPCR bind to *Plasmodium* was suggested to block the interaction of EPCR with protein
C. [53]. On the other hand, APC bind with EPCR was required to inhibit NETosis [54].
We have previously shown that *P. berghei* and *P.falciparum*-iRBC induced NET
formation and its importance in ARDS development [14].
However, there are specific domains of the PfEMP1 subfamily that do not bind to
EPCR while others do [20], although the presence of serum inhibits this binding during
in vitro assays [25]. Nevertheless, recently, a new study showed that using human brain
microvascular endothelial cells and 10% human serum did not affect the binding
between *P. falciparum*-infected red blood cells and EPCR. In addition, parasites
isolated from patients with cerebral malaria displayed higher binding capacity of

cytoadherence under flow conditions, compared with isolated from uncomplicated malaria patients [55]. Parasite var transcripts encoding EPCR-binding domains, in association with high load of parasite and low platelet levels, are hardy indicators of cerebral malaria. Patients with cerebral edema showed an increased transcript of parasite PfEMP1 DC8 and EPCR-binding domains. In addition 62B1-1-CIDRa1.7 affected EPCR-APC interaction and the authors proposed a harmful role for EPCR-binding domains in these patients [56]. Therefore, the exact role of EPCR is controversial and needs to be better understood. In addition, dexamethasone-treated mice are less susceptible to increased vascular permeability. Likewise, iRBC-stimulated PMLECs treated with dexamethasone showed less permeability than untreated cells. *P. berghei* NK65-infected C57BL/6 mice treated with the same dose of dexamethasone also showed an increase in peripheral parasitemia and ARDS protection [29] but not an effect on pulmonary vascular permeability.

Based on our results, treatment with dexamethasone, a potent anti-inflammatory drug, decreased the serum concentrations of IL-33, IL-6, and TNF in PbA-infected mice. In malaria, high level of TNF was associated with severe malaria, especially in children. TNF can participate in parasite killing and, consequently, parasitemia reduction. On the other hand, increased TNF levels were observed in children with high-density parasitemia and when parasite clearance occurs, TNF decays. It is described, in a recent review, that TNF polymorphisms could control parasite development in early stages of infection, but can enlarge the probability of patients to develop severe malaria syndromes [57].

TNF is an inflammatory mediator strongly implicated in the development of ARDS [58] signaling through two receptors, p55 and p75, that play differential roles in pulmonary edema formation during ARDS. It was shown by a novel domain antibody (dAb™) that p55 attenuated ventilator-induced lung injury [59] and lung injury and edema formation in models of ARDS induced by acid aspiration [58]. Recently, the selective blockage of TNFR1 (GSK1995057 antibody) inhibited cytokine and neutrophil adhesion molecule expression in activated HMVEC-L monolayers *in vitro* and attenuated inflammation and signs of lung injury in primates [60]. Moreover, treatment with this antibody attenuated pulmonary neutrophilia, inflammatory cytokine release and signs of endothelial injury in BAL and serum samples in healthy humans challenged with a low dose of inhaled endotoxin [60]. However, the precise mechanisms by which TNF is involved in severe malaria remains unknown [57].

IL-33 is a tissue-derived nuclear cytokine from the IL-1 family, and it is highly expressed during homeostasis and inflammation in endothelial, epithelial and fibroblast-like cells, working as an alarm signal released upon cell injury or tissue damage to alert immune cells to express the ST2 receptor (IL-1RL1) [61]. Bronchial IL-33 expression is significantly increased in severe malaria patients with pulmonary edema [62]. Moreover, in PbA-induced experimental cerebral malaria (ECM), IL-33 expression is increased in the brain, and ST2-deficient mice were resistant to PbA-induced neuropathology [63,64]. On the other hand, PbA-infected C57BL/6 mice treated with recombinant IL-33 presented no signs of neurological pathology associated with CM and had reduced production of pro-inflammatory cytokines and chemokines [65].

Circulating IL-6 levels are elevated in nearly all infectious, traumatic, and inflammatory states, including ARDS. Elevated levels of IL-6 are found in the BAL and plasma of patients with ARDS and those at risk [66,67].

Finally, we proposed the essential mechanisms of protection for dexamethasone in experimental malaria (Figure 6), suggesting that PbA-iRBCs induce TNF release by endothelial cells and by Mφ, and then TNF upregulates EPCR expression in PMLECs,

increasing iRBC adhesion through the EPCR pathway. The adhesion of iRBCs in pulmonary endothelial cells leads to an increase in gap formation in the interendothelial junction, raising vascular permeability and consequently the formation of edema. Activated alveolar macrophages also produce TNF, which contributes to the activation of endothelial cells and possibly the recruitment of neutrophils [14] and alveolar damage. The drastic reduction in inflammation with dexamethasone treatment may be responsible for the increase in parasitemia and mortality. These data suggest that TNF is essential to increase the expression of EPCR and consequently increase adhesion of iRBCs in malaria-associated ARDS.

Conclusions

Our data together suggest that *P. berghei* infection induces TNF production by inflammatory and endothelial cells, leading to the expression of adhesion molecules such as ICAM-1, VCAM and, especially, EPCR. These results allow us to infer that those factors could contribute to PbA-iRBC cytoadhesion and thus suggest the participation of these mechanisms in the pathogenesis of malaria-associated ARDS. Knowing that inhibition of cytoadhesion mechanisms may improve the prognosis of infection [68], we emphasize that the processes described above may be important in the treatment of ARDS due to *Plasmodium* infection, acting as adjunctive therapy in association with the use of antimalarials, not only making the treatment more efficient but also reducing the morbidity and mortality of these patients.

Conflicts of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding Statement

This work was supported by the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo: FAPESP, Brazil), the Coordination of Improvement of HigherLevel Personnel; (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior: CAPES) and the National Council for Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e Tecnológico: CNPq, Brazil). LSO (CAPES and FAPESP 2013/20718-3), MKS (CAPES/ PNPD), GCM (FAPESP 2017/00077-4), TCQ (CNPq 131431/2017-0), DD (CNPq 162730/2014-4), DSC (CNPq 133890/2016-3), OM (FAPESP 2013/00981-1), CRFM (FAPESP 2016/07030-3), SE (FAPESP 2014/20451-0 and 2017/05782-8), and CNPQ (455863/2014-8) received fellowship grants.

Acknowledgments

The authors thank Patricia Mendonça da Silva Amorim, Erika Paula Machado Peixoto, and Bernardo Paulo Albe for their technical assistance.

REFERENCES

1. Craig AG, Khairul MFM, Patil PR. Cytoadherence and severe malaria. Malays J
Med Sci [Internet]. 2012 Apr [cited 2013 Mar 19];19(2):5–18. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3431742&tool=pmce
ntrez&rendertype=abstract
2. Beeson JG, Brown G V. Pathogenesis of Plasmodium falciparum malaria: the
roles of parasite adhesion and antigenic variation. Cell Mol Life Sci [Internet].
2002 Feb;59(2):258–71. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11915943
3. Miller LH, Baruch DI, Marsh K, Doumbo OK. The pathogenic basis of malaria.
Nature [Internet]. 2002 Feb 7;415(6872):673–9. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11832955
4. Greenwood BM, Fidock DA, Kyle DE, Kappe SHI, Alonso PL, Collins FH, et al.
Malaria: progress, perils, and prospects for eradication. J Clin Invest [Internet].
2008 Apr [cited 2013 Feb 27];118(4):1266–76. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2276780&tool=pmce
ntrez&rendertype=abstract
5. Pouvelle B, Buffet P a, Lépolard C, Scherf a, Gysin J. Cytoadhesion of
Plasmodium falciparum ring-stage-infected erythrocytes. Nat Med [Internet].
2000 Nov;6(11):1264–8. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11062539
6. Rénia L, Wu Howland S, Claser C, Charlotte Gruner A, Suwanarusk R, Hui Teo
T, et al. Cerebral malaria: mysteries at the blood-brain barrier. Virulence
[Internet]. 2012 [cited 2013 Mar 8];3(2):193–201. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3396698&tool=pmce
ntrez&rendertype=abstract
7. Grommes J, Soehnlein O. Contribution of neutrophils to acute lung injury. Mol
Med [Internet]. 2011 [cited 2011 Jul 20];17(3–4):293–307. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3060975&tool=pmce
ntrez&rendertype=abstract
8. Bhatia M, Zemans RL, Jeyaseelan S. Role of chemokines in the pathogenesis of
acute lung injury. Am J Respir Cell Mol Biol [Internet]. 2012 May [cited 2013
Mar 7];46(5):566–72. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/22323365
9. Matute-Bello G, Martin TR. Science review: apoptosis in acute lung injury. Crit
Care [Internet]. 2003 Oct [cited 2011 Jul 29];7(5):355–8. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=270707&tool=pmcen
trez&rendertype=abstract
10. Van den Steen PE, Deroost K, Deckers J, Van Herck E, Struyf S, Opdenakker G.
Pathogenesis of malaria-associated acute respiratory distress syndrome. Trends
Parasitol [Internet]. 2013 Jun 3 [cited 2013 Jun 10];29(7):346–58. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/23742967
11. Lovegrove FE, Gharib S a, Peña-Castillo L, Patel SN, Ruzinski JT, Hughes TR,
et al. Parasite burden and CD36-mediated sequestration are determinants of acute
lung injury in an experimental malaria model. PLoS Pathog [Internet]. 2008 May
[cited 2012 Mar 18];4(5):e1000068. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2364663&tool=pmce
ntrez&rendertype=abstract
12. Ortolan LS, Sercundes MK, Barboza R, Debone D, Murillo O, Hagen SCF, et al.
Predictive criteria to study the pathogenesis of malaria-associated ALI/ARDS in
mice. Mediators Inflamm [Internet]. 2014 Jan [cited 2014 Dec 16];2014:872464.

Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4167651&tool=pmce
ntrez&rendertype=abstract

13.   Epiphanio S, Campos MG, Pamplona A, Carapau D, Pena AC, Ataíde R, et al.
VEGF promotes malaria-associated acute lung injury in mice. PLoS Pathog
[Internet]. 2010 May [cited 2011 Jul 12];6(5):e1000916. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2873913&tool=pmce
ntrez&rendertype=abstract

14.   Sercundes MK, Ortolan LS, Debone D, Soeiro-Pereira P V., Gomes E, Aitken
EH, et al. Targeting neutrophils to Prevent Malaria-Associated Acute Lung
Injury/Acute respiratory Distress Syndrome in Mice. PLOS Pathog [Internet].
2016;12(12):e1006054. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/27926944%0Ahttp://dx.plos.org/10.1371/j
ournal.ppat.1006054

15.   Pereira MLM, Ortolan LS, Sercundes MK, Debone D, Murillo O, Lima FA, et al.
Association of Heme Oxygenase 1 with Lung Protection in Malaria-Associated
ALI/ARDS. Mediators Inflamm. 2016;2016.

16.   Pham T-T, Verheijen M, Vandermosten L, Deroost K, Knoops S, Van den Eynde
K, et al. Pathogenic CD8+ T Cells Cause Increased Levels of VEGF-A in
Experimental Malaria-Associated Acute Respiratory Distress Syndrome, but
Therapeutic VEGFR Inhibition Is Not Effective. Front Cell Infect Microbiol.
2017;7(September):1–10.

17.   El-Assaad F, Wheway J, Mitchell AJ, Lou J, Hunt NH, Combes V, et al.
Cytoadherence of Plasmodium berghei-infected red blood cells to murine brain
and lung microvascular endothelial cells in vitro. Infect Immun [Internet]. 2013
Aug 12 [cited 2013 Aug 20];81(11):3984–91. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23940206

18.   Cunningham DA, Lin J, Brugat T, Jarra W, Tumwine I, Kushinga G, et al.
ICAM-1 is a key receptor mediating cytoadherence and pathology in the
Plasmodium chabaudi malaria model. Malar J [Internet]. 2017;16(1):185.
Available from:
http://malariajournal.biomedcentral.com/articles/10.1186/s12936-017-1834-8

19.   Aird WC, Mosnier LO, Fairhurst RM. Plasmodium falciparum picks (on) EPCR.
Blood. 2014;123(2):163–7.

20.   Turner L, Lavstsen T, Berger SS, Wang CW, Petersen JE V, Avril M, et al.
Severe malaria is associated with parasite binding to endothelial protein C
receptor. Nature [Internet]. 2013;498(7455):502–5. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3870021&tool=pmce
ntrez&rendertype=abstract

21.   Rao LVM, Esmon CT, Pendurthi UR. Review Article Endothelial cell protein C
receptor⬜: a multiliganded and multifunctional receptor. Blood.
2014;124(10):1553–63.

22.   Shabani E, Opoka RO, Bangirana P, Park GS, Vercellotti GM, Guan W, et al.
The endothelial protein C receptor rs867186-GG genotype is associated with
increased soluble EPCR and could mediate protection against severe malaria. Nat
Publ Gr. 2016;

23.   Mosnier LO, Lavstsen T. The role of EPCR in the pathogenesis of severe
malaria. Thromb Res [Internet]. 2016;141:S46–9. Available from:
http://dx.doi.org/10.1016/S0049-3848(16)30364-4

24.   Moxon CA, Wassmer SC, Milner DA, Chisala N V., Taylor TE, Seydel KB, et

al. Loss of endothelial protein C receptors links coagulation and inflammation to
parasite sequestration in cerebral malaria in African children. Blood.
2013;122(5):842–51.

25. Azasi Y, Lindergard G, Ghumra A, Mu J, Miller LH, Rowe JA. Infected
erythrocytes expressing DC13 PfEMP1 differ from recombinant proteins in
EPCR-binding function. Proc Natl Acad Sci [Internet]. 2018;201712879.
Available from: http://www.pnas.org/lookup/doi/10.1073/pnas.1712879115

26. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-
time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods [Internet].
2001 Dec [cited 2011 Jul 15];25(4):402–8. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11846609

27. Chen SF, Fei X, Li SH. A new simple method for isolation of microvascular
endothelial cells avoiding both chemical and mechanical injuries. Microvasc Res
[Internet]. 1995 Jul [cited 2013 May 14];50(1):119–28. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/7476573

28. Trager W, Jensen JB. Human malaria parasites in continuous culture. Science
(80-). 1976 Aug;193(4254):673–5.

29. Van den Steen PE, Geurts N, Deroost K, Van Aelst I, Verhenne S, Heremans H,
et al. Immunopathology and dexamethasone therapy in a new model for malaria-
associated acute respiratory distress syndrome. Am J Respir Crit Care Med
[Internet]. 2010 May 1 [cited 2011 Nov 1];181(9):957–68. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/20093644

30. Thumwood CM, Hunt NH, Clark IA, Cowden WB. Breakdown of the blood-
brain barrier in murine cerebral malaria. Parasitology [Internet]. 1988 Jun
6;96(3):579–89. Available from:
https://www.cambridge.org/core/product/identifier/S0031182000080203/type/jou
rnal_article

31. Shu J, Qiu G, Mohammad Ilyas. Immunohistochemistry (IHC) Image Analysis
Tollbox [Internet]. 2015. Available from: http://rsb.info.nih.gov/ij/plugins/ihc-
toolbox/index.html

32. Aitken EH, Negri EM, Barboza R, Lima MRI, Alvarez JM, Marinho CRF, et al.
Ultrastructure of the lung in a murine model of malaria-associated acute lung
injury/acute respiratory distress syndrome. Malar J [Internet]. 2014;13:230.
Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4062769&tool=pmce
ntrez&rendertype=abstract

33. Pino P, Vouldoukis I, Kolb JP, Mahmoudi N, Desportes-Livage I, Bricaire F, et
al. Plasmodium falciparum--infected erythrocyte adhesion induces caspase
activation and apoptosis in human endothelial cells. J Infect Dis [Internet]. 2003
Apr 15;187(8):1283–90. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/12696008

34. De las Salas B, Segura C, Pabón A, Lopes SCP, Costa FTM, Blair S. Adherence
to human lung microvascular endothelial cells (HMVEC-L) of Plasmodium vivax
isolates from Colombia. Malar J [Internet]. 2013 Jan [cited 2014 Dec 17];12:347.
Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3850517&tool=pmce
ntrez&rendertype=abstract

35. Lagasse HAD, Anidi IU, Craig JM, Limjunyawong N, Poupore AK, Mitzner W,
et al. Recruited monocytes modulate malaria-induced lung injury through CD36-
mediated clearance of sequestered infected erythrocytes. J Leukoc Biol [Internet].

2016 May 1;99(5):659–71. Available from:
http://www.jleukbio.org/cgi/doi/10.1189/jlb.4HI0315-130RRR

36. Maknitikul S, Luplertlop N, Grau GER, Ampawong S. Dysregulation of
pulmonary endothelial protein C receptor and thrombomodulin in severe
falciparum malaria-associated ARDS relevant to hemozoin. PLoS One [Internet].
2017;12(7):e0181674. Available from:
http://dx.plos.org/10.1371/journal.pone.0181674

37. Jaramillo M, Godbout M, Olivier M. Hemozoin induces macrophage chemokine
expression through oxidative stress-dependent and -independent mechanisms. J
Immunol [Internet]. 2005 Jan 1;174(1):475–84. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/15611273

38. Shio MT, Kassa FA, Bellemare MJ, Olivier M. Innate inflammatory response to
the malarial pigment hemozoin. Microbes Infect. 2010;12(12–13):889–99.

39. Coban C, Ishii KJ, Kawai T, Hemmi H, Sato S, Uematsu S, et al. Toll-like
receptor 9 mediates innate immune activation by the malaria pigment hemozoin.
J Exp Med [Internet]. 2005 Jan 3 [cited 2013 Mar 6];201(1):19–25. Available
from: http://www.jem.org/lookup/doi/10.1084/jem.20041836

40. Parroche P, Lauw FN, Goutagny N, Latz E, Monks BG, Visintin A, et al. Malaria
hemozoin is immunologically inert but radically enhances innate responses by
presenting malaria DNA to Toll-like receptor 9. Proc Natl Acad Sci U S A
[Internet]. 2007 Feb 6;104(6):1919–24. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1794278&tool=pmce
ntrez&rendertype=abstract

41. El-Assaad F, Wheway J, Mitchell AJ, Lou J, Hunt NH, Combes V, et al.
Cytoadherence of plasmodium berghei-infected red blood cells to murine brain
and lung microvascular endothelial cells in vitro. Infect Immun [Internet]. 2013
Aug 12 [cited 2013 Aug 20];81(11):3984–91. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23940206

42. Ockenhouse CF, Tegoshi T, Maeno Y, Benjamin C, Ho M, Kan KE, et al.
Human vascular endothelial cell adhesion receptors for Plasmodium falciparum-
infected erythrocytes: roles for endothelial leukocyte adhesion molecule 1 and
vascular cell adhesion molecule 1. J Exp Med [Internet]. 1992 Oct
1;176(4):1183–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/1383378

43. Rambhatla JS, Turner L, Manning L, Laman M, Davis TME, Beeson JG, et al.
Acquisition of antibodies against endothelial protein c receptor-binding domains
of plasmodium falciparum erythrocyte membrane protein 1 in children with
severe malaria. J Infect Dis. 2019;219(5):808–18.

44. Shabani E, Opoka RO, Bangirana P, Park GS, Vercellotti GM, Guan W, et al.
The endothelial protein C receptor rs867186-GG genotype is associated with
increased soluble EPCR and could mediate protection against severe malaria. Sci
Rep [Internet]. 2016;6(April):27084. Available from:
http://dx.doi.org/10.1038/srep27084

45. Pasini EM, Braks JA, Fonager J, Klop O, Aime E, Spaccapelo R, et al.
Proteomic and Genetic Analyses Demonstrate that Plasmodium berghei Blood
Stages Export a Large and Diverse Repertoire of Proteins . Mol Cell Proteomics.
2013;12(2):426–48.

46. Fonager J, Pasini EM, Braks J a M, Klop O, Ramesar J, Remarque EJ, et al.
Reduced CD36-dependent tissue sequestration of Plasmodium-infected
erythrocytes is detrimental to malaria parasite growth in vivo. J Exp Med
[Internet]. 2012 Jan 16 [cited 2012 Mar 16];209(1):93–107. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260870&tool=pmce
ntrez&rendertype=abstract

47. Meijvis SCA, Hardeman H, Remmelts HHF, Heijligenberg R, Rijkers GT, Van
Velzen-Blad H, et al. Dexamethasone and length of hospital stay in patients with
community-acquired pneumonia: A randomised, double-blind, placebo-
controlled trial. Lancet [Internet]. 2011;377(9782):2023–30. Available from:
http://dx.doi.org/10.1016/S0140-6736(11)60607-7

48. Azoulay É, Canet E, Raffoux E, Lengliné E, Lemiale V, Vincent F, et al.
Dexamethasone in patients with acute lung injury from acute monocytic
leukaemia. Eur Respir J. 2012;39(3):648–53.

49. Warrell DA, Looareesuwan S, Warrell MJ, Kasemsarn P, Intaraprasert R,
Bunnag D, et al. Dexamethasone Proves Deleterious in Cerebral Malaria. N Engl
J Med [Internet]. 1982 Feb 11;306(6):313–9. Available from:
http://www.nejm.org/doi/abs/10.1056/NEJM198202113060601

50. Hoffman SL, Rustama D, Punjabi NH, Surampaet B, Sanjaya B, Dimpudus AJ,
et al. High-dose dexamethasone in quinine-treated patients with cerebral malaria:
a double-blind, placebo-controlled trial. J Infect Dis [Internet]. 1988
Aug;158(2):325–31. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/3042874

51. Pasvol G. The treatment of complicated and severe malaria. Br Med Bull.
2005;75–76(1):29–47.

52. Gao Y, Ma G, Liu S, Teng Y, Wang Y, Su Y. Thalidomide and multiple
myeloma serum synergistically induce a hemostatic imbalance in endothelial
cells in vitro. Thromb Res [Internet]. 2015;135(6):1154–9. Available from:
http://dx.doi.org/10.1016/j.thromres.2015.03.019

53. Sampath S, Brazier AJ, Avril M, Bernabeu M, Vigdorovich V, Mascarenhas A,
et al. Plasmodium falciparum adhesion domains linked to severe malaria differ in
blockade of endothelial protein C receptor. Cell Microbiol. 2015;17(12):1868–
82.

54. Healy LD, Puy C, Fernández JA, Mitrugno A, Keshari RS, Taku NA, et al.
Activated protein C inhibits neutrophil extracellular trap formation in vitro and
activation in vivo. J Biol Chem. 2017;292(21):8616–29.

55. Storm J, Jespersen JS, Seydel KB, Szestak T, Mbewe M, Chisala N V, et al.
Cerebral malaria is associated with differential cytoadherence to brain endothelial
cells. EMBO Mol Med. 2019;11(2):e9164.

56. Kessler A, Dankwa S, Bernabeu M, Harawa V, Danziger SA, Duffy F, et al.
Linking EPCR-Binding PfEMP1 to Brain Swelling in Pediatric Cerebral Malaria.
Cell Host Microbe [Internet]. 2017;22(5):601-614.e5. Available from:
https://doi.org/10.1016/j.chom.2017.09.009

57. Penha-Gonçalves C. Genetics of Malaria Inflammatory Responses: A
Pathogenesis Perspective. Front Immunol. 2019;10(July):1–15.

58. Wilson MR, Wakabayashi K, Bertok S, Oakley CM, Patel B V., O'Dea KP, et al.
Inhibition of TNF receptor p55 by a domain antibody attenuates the initial phase
of acid-induced lung injury in mice. Front Immunol. 2017;8(FEB):1–12.

59. Wilson MR, Goddard ME, O'Dea KP, Choudhury S, Takata M. Differential roles
of p55 and p75 tumor necrosis factor receptors on stretch-induced pulmonary
edema in mice. Am J Physiol Cell Mol Physiol [Internet]. 2007;293(1):L60–8.
Available from: http://www.physiology.org/doi/10.1152/ajplung.00284.2006

60. Proudfoot A, Bayliffe A, O'Kane CM, Wright T, Serone A, Bareille PJ, et al.
Novel anti-tumour necrosis factor receptor-1 (TNFR1) domain antibody prevents

pulmonary inflammation in experimental acute lung injury. Thorax [Internet].
2018;1:thoraxjnl-2017-210305. Available from:
http://thorax.bmj.com/lookup/doi/10.1136/thoraxjnl-2017-210305
61.   Cayrol C, Girard JP. Interleukin-33 (IL-33): A nuclear cytokine from the IL-1
family. Immunol Rev. 2018;281(1):154–68.
62.   Ampawong S, Chaisri U, Viriyavejakul P, Prapansilp P, Grau GE, Turner GDH,
et al. A potential role for interleukin-33 and γ-epithelium sodium channel in the
pathogenesis of human malaria associated lung injury. Malar J. 2015;14(1):1–15.
63.   Palomo J, Reverchon F, Piotet J, Besnard AG, Couturier-Maillard A, Maillet I, et
al. Critical role of IL-33 receptor ST2 in experimental cerebral malaria
development. Eur J Immunol. 2015;45(5):1354–65.
64.   Reverchon F, Mortaud S, Sivoyon M, Maillet I, Laugeray A, Palomo J, et al. IL-
33 receptor ST2 regulates the cognitive impairments associated with
experimental cerebral malaria. PLoS Pathog. 2017;13(4):1–25.
65.   Besnard AG, Guabiraba R, Niedbala W, Palomo J, Reverchon F, Shaw TN, et al.
IL-33-Mediated Protection against Experimental Cerebral Malaria Is Linked to
Induction of Type 2 Innate Lymphoid Cells, M2 Macrophages and Regulatory T
Cells. PLoS Pathog. 2015;11(2):1–21.
66.   Park WY, Goodman RB, Steinberg KP, Ruzinski JT, Ii FR, Park DR, et al.
Cytokine Balance in the Lungs of Patients with Acute Respiratory Distress
Syndrome. 2001;(17):1896–903.
67.   Takala A, Jousela I, Takkunen O, Kautiainen H, Jansson S-E, Orpana A, et al. A
prospective study of inflammation markers in patients at risk of indirect acute
lung injury. Shock [Internet]. 2002;17(4):252–7. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11954822
68.   Lin JW, Sodenkamp J, Cunningham D, Deroost K, Tshitenge TC, McLaughlin S,
et al. Signatures of malaria-associated pathology revealed by high-resolution
whole-blood transcriptomics in a rodent model of malaria. Sci Rep [Internet].
2017;7(September 2016):1–13. Available from:
http://dx.doi.org/10.1038/srep41722

1000

1001 **Figure Captions**

1002

Figure 1: ARDS-developing mice have higher levels of pulmonary parasites than
HP-developing mice. (a and b) The distribution of infected red blood cells (iRBCs) was
analyzed in DBA/2 mice infected with luciferase-expressing *P. berghei* parasite (PbA-
luciferase) on the 7[th] day postinfection (dpi) and analyzed *in vivo* by IVIS imaging after
luciferin. (c) The organs were individually analyzed after perfusion and results were
plotted with black circles (mice with pleural effusion) and with gray squares (mice
without pleural effusion); one-way ANOVA (**p<0.01; ***p<0.001). (d) Perfused
lungs from ARDS- and HP-developing mice were collected on the 7[th] dpi and analyzed
by rRNA gene expression of *P. berghei* ANKA (18S subunit) using the $2^{-\Delta\Delta CT}$ method.
Representative image of a perfused lung collected from an ARDS-developing mouse
under (e) normal and polarized lights (f) showing the hemozoin pigments, on the 7[th] dpi.
(g) Quantification of the hemozoin area analyzed by ImageJ (magnification: 400×, scale
bar: 20 μm). Bars represent the average ± SD (*p<0.05; **p< 0.01). (d) Unpaired t-test
and (g) Mann-Whitney test. (h) Lung histological analyses showing iRBCs in close
contact with endothelial cells (inserts) in an infected mouse that died by ARDS
(magnification: 630X, scale bar: 10 μm). ARDS: acute respiratory distress syndrome;
HP: hyperparasitemia; Lu: lung; Sp: spleen; Ki: kidney; Li: liver; Br: brain; He: heart.

Figure 2: TNF increases *Plasmodium berghei* adherence in the ARDS experimental model. (a) Serum from ARDS- and HP-developing mice was collected on the 7th day postinfection (dpi) and submitted for TNF analysis by ELISA. (b) Primary microvascular lung endothelial cells (PMLECs) from naïve DBA/2 mice were stimulated with infected red blood cells (iRBCs) and TNF was measured in the supernatant. (c-e) PMLECs were also stimulated with TNF (50 ng/ml) and the adhesion of iRBCs to PMLECs under static conditions and (f-h) under flow conditions (after 24 hours of TNF-stimulation) were analysed. (i-k) iRBC adhered in PMLEC in a static condition assay after coincubation with peritoneal macrophages (Mφ), seeded in transwell membranes and coincubated with iRBCs or red blood cells (RBCs) for 24 hours, and then (j) TNF release was measured in the supernatant. (k) PMLECs (that had contact with the Mφ supernatant) were incubated with iRBCs having their adhesion analyzed. (l) iRBC adherence was assessed in TNF-stimulated cells and TNF-neutralized antibodies plus TNF. Bars represent the average ± SEM, Graphics a: one-way ANOVA from two grouped experiments; (b-h) Mann-Whitney test representative of (b and h) two independent experiments and (c-e and l) two grouped experiments; (j and k) Kruskal-Wallis test representative of two independent experiments; Bars represent the average ± SD (*p<0.05; **p<0.01; ***p<0.001). ARDS: acute respiratory distress syndrome; HP: hyperparasitemia; Arrows in f, g and i indicates iRBC adhered in PMLEC.

Figure 3: EPCR could contribute to *P. berghei* cytoadherence. Lungs and serum from ARDS- and HP-developing mice were analyzed by (a) EPCR mRNA expression, (b) soluble EPCR (sEPCR) in serum and (c) EPCR protein in lungs, on the 7th day postinfection. (d-f) Primary microvascular endothelial cells (PMLECs) were stimulated with either infected red blood cells (iRBCs) or TNF (50 µg/ml) for (d) 24, (e) 48 and (f) 72 hours to analyze EPCR mRNA expression. (g) PMLECs were subjected to EPCR knockdown with siRNA, and (h) iRBC adherence in PMLECs was evaluated. (a, b and d) Unpaired t-test representative of (a) three grouped, (b) two independent and (d) two grouped experiments; (c) Mann-Whitney test representative of two independent experiments; (e-h) One-way ANOVA from two grouped experiments. Bars represent the average ± SD (*p<0.05; **p<0.01; ***p<0.001; ****p<0.0001; n=4-7 replicates/group). ARDS: acute respiratory distress syndrome; HP: hyperparasitemia; NS: nonstimulated cells; NT: nontransfected cells. Red dashed lines: noninfected mice.

Figure 4: Dexamethasone reduces VEGF, TNF and EPCR, protecting mouse lungs and PMLECs from increased vascular permeability. (a-h) *Plasmodium berghei* ANKA-infected DBA/2 mice were treated with dexamethasone and their serum and perfused lungs were analyzed, on the 7th day postinfection. (a) The TNF concentration in the serum, (b) EPCR mRNA expression in the lungs, (c) sEPCR concentration in both sera and (d) bronchoalveolar lavage (BAL) and (e) VEGF concentration in BAL were evaluated. (f-g) Evans Blue (EB) accumulation was measured in the lungs and (h) weight lungs was analyzed after perfusion. (i-j) Primary microvascular lung endothelial cells (PMLECs) were treated, or not, with dexamethasone and, subsequently, coincubated with either infected red blood cells (iRBCs) or red blood cells (RBCs). (i) PMLECs were stained for actin microfilaments (gray), and nuclei were stained with Hoechst stain (blue), and (j) openings in the interendothelial junctions (OIJ) (indicated by yellow arrows) were compared among all groups. (k) The permeability of dexamethasone-treated PMLECs was related using Evans blue concentration in a Transwell assay. (l) VEGF levels were measured in these groups. Infected+Dexa: *P.*

berghei ANKA-infected and dexamethasone-treated mice. Graphics (a, c, d, g and h) Mann-Whitney test representative of (a, d, g and h) two independent and (c) two grouped experiments. (b, e and k) Unpaired t-test from (b) two grouped, (e) from two independent and (k) three grouped experiments; (i) Kruskall-Wallis test from three grouped experiments; (l) One-way ANOVA representative of two independent experiments. Bars represent the average ± SD (*p<0.05; **p<0.01; ****p<0.0001). Red dashed lines: noninfected mice.

Figure 5: Dexamethasone protects mice from ARDS but not from malaria infection. *Plasmodium berghei* ANKA-infected DBA/2 mice were treated with dexamethasone and compared to infected-untreated mice, on the 7th day post-infection for respiratory parameters: (a) respiratory frequency, (b) tidal volume and (c) enhanced pause (Penh) analyzed in whole body plethysmography chambers. (d) Parasitemia (%) in the peripheral blood and (e) the survival curves were compared between groups. Representative image of histological analysis of lungs from (f) an infected mouse and (g) an Infected+Dexa mouse (Scale bar: 50 μm). Infected+Dexa: *P. berghei* ANKA-infected and dexamethasone-treated mice. Graphics (a-d) Unpaired t test, representative data from two independent experiments. Bars represent the average ± SD. (*p<0.05; ** p<0.01). Red dashed lines: noninfected mice.

Figure 6: Schematic representation of ARDS development involving adhesion of iRBCs to EPCR. *Plasmodium berghei* ANKA-infected red blood cells (iRBCs) induce TNF release by endothelial and inflammatory cells. TNF upregulates EPCR expression in lung endothelial cells, increasing iRBCs adhesion in these cells through EPCR pathway. The adhesion of iRBCs to lung endothelial cells leads to an increase in gap formation in the interendothelial junction, an increase in vascular permeability with infiltration of inflammatory cells and red blood cells and edema formation. Activated alveolar macrophages also produce TNF, which contributes to the activation of endothelial cells, recruitment of neutrophils and alveolar damage. Treatment with dexamethasone decreases inflammatory cytokine release, including TNF release and, consequently, downregulates EPCR expression in lung endothelial cells; dexamethasone also decreases VEFG released by endothelial cells, protecting mice from gap formations, vascular permeability, inflammatory cell infiltration and alveolar damage.

a Before perfusion

b After perfusion

He   Br
Lu
Sp
Li   Ki

He   Br
Lu
Li   Sp
Ki

c

Luciferase (ROI)

4×10^8

3×10^8

2×10^8

1×10^8

Lu   Sp   Ki   Li   Br   He

● With pleural effusion   ■ Without pleural effusion

d

P. berghei ANKA rRNA expression (18s)
Fold increased relative to hprt

*

ARDS   HP

e

f

g

Hemozoin area in lungs (%) normalized by total area

1.0

0.8

0.6

0.4

0.2

0.0

-0.2

**

ARDS   HP

h